Persistence and Free Chlorine Disinfection of Human Coronaviruses and Their Surrogates

in Water

Mengyang Zhang,[1] Michelle Wei Leong,[2] William A. Mitch,[1] Catherine A. Blish,[2] Alexandria

Boehm[1*]

[1] Department of Civil and Environmental Engineering, School of Engineering and Doerr School

of Sustainability, Stanford University, Stanford, CA 94305, United States

[2] Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, United

States

* Corresponding Author:

Alexandria Boehm, Phone: 650 724-9128 Email: aboehm@stanford.edu

Abstract

The COVID-19 pandemic illustrates the importance of understanding the behavior and

control of human pathogenic viruses in the environment. Exposure via water (drinking, bathing,

and recreation) is a known route of transmission of viruses to humans, but the literature is

relatively void of studies on the persistence of many viruses, especially coronaviruses, in water

and their susceptibility to chlorine disinfection. To fill that knowledge gap, we evaluated the

persistence and free chlorine disinfection of human coronavirus OC43 (HCoV-OC43) and its

surrogates, murine hepatitis virus (MHV) and porcine transmissible gastroenteritis virus

(TGEV), in drinking water and laboratory buffer using cell culture methods. The decay rate

constants of human coronavirus and its surrogates in water varied depending on virus and water

matrix. In drinking water prior to disinfectant addition, MHV showed the largest decay rate

constant (2.25 day^{-1}) followed by HCoV-OC43 (0.99 day^{-1}) and TGEV (0.65 day^{-1}); while in

phosphate buffer, HCoV-OC43 (0.51 day^{-1}) had a larger decay rate constant than MHV (0.28

day^{-1}) and TGEV (0.24 day^{-1}). Upon free chlorine disinfection, the inactivation rates of

coronaviruses were independent of free chlorine concentration and not affected by water matrix,

though they still varied between viruses. TGEV showed the highest susceptibility to free chlorine

disinfection with the inactivation rate constant of 113.50 mg^{-1} min^{-1} L, followed by MHV (81.33

mg^{-1} min^{-1} L) and HCoV-OC43 (59.42 mg^{-1} min^{-1} L).

Importance: This study addresses an important knowledge gap on enveloped virus persistence

and disinfection in water. Results have immediate practical applications for shaping evidence-

based water policies, particularly in the development of disinfection strategies for pathogenic

virus control.

Keywords: Human coronavirus | persistence | disinfection | drinking water

Introduction

Individuals can acquire viral infections via exposure to contaminated water, through
drinking, bathing, and recreating. Globally, at least 2 billion people use a drinking water source
contaminated with feces; exposure to resultant microbial contamination can cause diarrhoeal
diseases, acute respiratory diseases and various neglected tropical diseases.[1] Waterborne viral
pathogens are classified with moderate to high health significance by the World Health
Organization (WHO) because they can have prolonged persistence in water supplies, moderate
resistance to chlorine disinfection, and high relative infectivity.[2]

Human coronaviruses, which are single-stranded positive-sense RNA viruses from the family
of *Coronaviridae*, have been circulating in human populations for decades. The first human
coronavirus was isolated in 1965, while three highly pathogenic human coronaviruses emerged
in the last 20 years: severe acute respiratory syndrome coronavirus (SARS-CoV-1), Middle East
respiratory syndrome coronavirus (MERS-CoV), and SARS-CoV-2.[3,4] The other four human
coronaviruses (HCoVs) HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-HKU1 are
seasonal or endemic human coronaviruses, typically result in mild to moderate upper-respiratory
tract illness, are responsible for 15-30% of common colds in adults, and represent the second
most prevalent cause of the common cold.[5–7] Notably, though HCoVs typically induce mild and
self-limiting symptoms in immunocompetent adults, they could provoke lower respiratory tract
inflammation and result in fatal outcomes among vulnerable populations such as infants, the
elderly, pregnant women, and other immunocompromised individuals.[7–9] HCoV-OC43 has also
shown neuroinvasive potential, supporting observations of persistent replication in primary
neuronal cell cultures and confirmed HCoV-OC43 neuroinvasive disease cases in humans.[10–13]

Regarding the global epidemiology of HCoVs, they exhibit seasonal patterns with peak activity

typically occurring between December and March (Northern hemisphere) annually.[6,14] The

predominant HCoV species can vary by region, year, or age group, with HCoV-OC43 being the

most commonly detected and peaking annually.[6,14,15]

During the COVID-19 pandemic, a number of studies were completed that investigated the

environmental occurrence, persistence, and disinfection of HCoVs, including SARS-CoV-2, but

these studies primarily focused on surfaces of fomite and food or in the air/aerosol droplets,[16–18]

as well as wastewater surveillance[19–21]. HCoVs have demonstrated the ability to survive in

aerosols for hours to days, with persistence influenced by factors such as temperature and

humidity.[16,18] HCoVs also exhibit high persistence on fomites, with examples like HCoV-229E

and SARS-CoV-2 surviving for 3-5 days on stainless steel surfaces.[16] However, there are very

limited studies on the persistence of HCoVs in water, particularly drinking water, and their

susceptibility to chlorine disinfection.[22–24]

A few studies imply HCoVs could be persistent in water but susceptible to free chlorine

disinfection in water but detailed evaluations are lacking. HCoV-229E could survive in

dechlorinated tap water for ~ 8 days (time to 99% inactivation, T99, at 23°C) and decayed

significantly slower at a lower temperature (i.e., T_{99} >100 days at 4°C).[25] Bivins et al.[26] reported

a T_{99} of ~ 3.9 days for SARS-CoV-2 at 20°C in tap water had not been dechlorinated. Several

other studies on the persistence of HCoVs and SARS-CoV-2 in water matrices were conducted

but targeting the viral RNA which does not reflect the virus infectivity and usually shows higher

persistence compared with intact, infectious viruses.[22,24] As a common surrogate for enveloped

virus, *Pseudomonas phage* Phi6 showed great free chlorine susceptibility at 23°C in phosphate-

buffered saline (PBS) at pH 7.4 with an inactivation rate constant of 276.0 ± 30 mg^{-1} min^{-1} L.[27]

However, there has been no human coronavirus inactivation rate constant with exposure to free

chlorine reported so far. Thus, there is a significant research gap regarding HCoVs persistence in

drinking water and their susceptibility to free chlorine disinfection.

This study fills an important research gap on human coronavirus survival in water and their

potential for chlorine disinfection. In particular, we demonstrate the persistence and free chlorine

disinfection of infectious human coronaviruses and their surrogates in drinking water and

laboratory buffer. HCoV-OC43 was selected as a representative human coronavirus because of

its high prevalence in humans.[6,14,15] We also selected murine hepatitis virus (MHV) and

transmissible gastroenteritis virus (TGEV), which are mammalian coronaviruses infecting mice

and pigs, respectively, and are common alternative viruses used in HCoV-related research.[28,29]

The decay rate constants of coronaviruses in water and their free chlorine disinfection kinetics

were measured through cell culture-based assays. The results of this study will be of immediate

use to inform water policies for pandemic preparedness and also highlight the necessity of

selecting appropriate virus surrogates and water matrices in research on human pathogenic virus

persistence and disinfection in water.

Experimental Methods

Overview. The persistence and free chlorine disinfection of human coronaviruses and their

surrogates in drinking water were investigated by measuring their decay rate constants and free

chlorine disinfection kinetics in drinking water. Factors including different viruses (HCoV-

OC43, MHV, TGEV), water matrix (drinking water, phosphate buffer), and free chlorine

concentration (mean value $\pm$ standard deviation: 1.31 ± 0.06 mg L^{-1}, 2.53 ± 0.08 mg L^{-1}) were

investigated and statistically analyzed to identify the significant predictors for coronavirus decay

and free chlorine disinfection kinetics in water. The concentration of infectious coronaviruses

was quantified using a cell culture-based endpoint dilution assay known as the $TCID_{50}$ method.

Cells and virus propagation. Delayed brain tumor (DBT) cells were kindly provided by Dr.

Krista Wigginton (University of Michigan, Ann Arbor, MI); swine testicular cells (ST) (CRL-

1746, ATCC) and Vero E6 cells (CRL-1586, ATCC) were purchased from ATCC. All cell lines

were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal

bovine serum (FBS) and 1% Penicillin-Streptomycin (P/S) and incubated at 37°C/5% CO_2 until

confluence. HCoV-OC43 and MHV were kindly provided by Dr. Jeffrey S. Glenn (Stanford

University, Stanford, CA) and Dr. Krista Wigginton, respectively. For virus propagation, HCoV-

OC43, MHV, and TGEV (VR-1995, ATCC) were inoculated to confluent Vero E6, DBT, and

ST cells respectively, which were all maintained in serum free DMEM, at a multiplicity of

infection (MOI) of 0.001-0.01. Infected cells were incubated at 37°C/5% CO_2 until significant

cytopathic effect (CPE) was observed, which usually took 6, 1, and 2 days for HCoV-OC43,

MHV, and TGEV respectively. Virus suspension was collected and stored at -20 °C until further

virus concentration and purification (see **Virus purification**).

Virus purification. The presence of cell culture components and culture medium in virus

stocks during the virus propagation process could cause significant bias in virus inactivation

studies, by, for example, introducing artificial chlorine demand to experimental water matrices

in virus seeding steps.[30] In particular, DMEM is enriched with amino acids and vitamins (**Table**

S1), which react with free chlorine rapidly[30]. To minimize the introduction of constituents with

significant chlorine demand, all coronavirus stocks were purified through an ultracentrifugation-

Amicon filtration process. First, crude coronavirus suspension was centrifuged at 4,000 ×g for 10

minutes. Then, the supernatant was filtered through 0.22 µm-pore size membranes to remove cell

debris, and 120 mL of the filtrate was processed by ultracentrifugation at 24,000 rpm for 2.5

hours with a 30% of sucrose cushion. The coronavirus pellets were rinsed with phosphate buffer

(10 μM, pH = 7.1) once and then resuspended in 1 mL autoclaved phosphate buffer. The 1 mL

coronavirus suspension was then washed twice using AmiconUltra (Merck Millipore) with 10

mL phosphate buffer. The result was approximately 150-200 μL of purified coronavirus stock.

The purified coronavirus stocks were stored at -80 °C for up to two weeks until used in

persistence and free chlorine disinfection experiments.

Persistence experiments. We obtained water samples from the Vineyard Water Treatment

Plant in Sacramento, California, USA, which produces potable water for the surrounding

communities. Water samples were collected right after the sedimentation process and before

chlorine addition. Water quality parameters are provided in **Table S2**. The water samples were

filtered through 0.7 μm pore size glass fiber membranes (Whatman) to mimic the filtration

process in the Vineyard Water Treatment Plant and then stored at -80 °C for up to a year until

use. This water is subsequently referred to as "drinking water".

290 μL of purified HCoV-OC43, MHV, and TGEV stocks were spiked into either 2,610 μL

of drinking water or autoclaved phosphate buffer at a ratio of 1:10 separately to achieve final

coronavirus concentrations of ~ 10^5-10^6 TCID$_{50}$ mL^{-1}; incubated at room temperature (22°C) for

up to 30 days. During the incubation period, reactors were kept on a sample mixer

(HulaMixer^{TM}) and rotated vertically at 1 rpm. 60 μL of subsamples were collected on days 0, 1,

2, 3, 5, 7, 11, 17, and 30, and tested immediately for the TCID$_{50}$ to quantify the concentration of

infectious coronaviruses (see below **TCID$_{50}$ for virus titration**). Two reactors were used for

each coronavirus (experimental duplicates) and two subsamples were collected at each time point

from each reactor (technical duplicates), resulting in four replicates for each time point.

The decay rate constants (k_{decay}) were estimated by fitting a first-order decay model to the

corresponding persistence data from four replicates, while excluding the data under the lower

limit of detection (LoD):

$$ln\left(\frac{C}{C_0}\right) = \beta_0 - k_{decay}t$$

where C is the average infectious concentration of each coronavirus at each time point in each

water matrix (TCID$_{50}$ mL^{-1}), C_0 is the average infectious concentration of each coronavirus at

time 0 in each water matrix (TCID$_{50}$ mL^{-1}), β_0 is the intercept, k_{decay} is the first-order decay rate

constant for each condition (day^{-1}), and t is the incubation time (day). k_{decay} and its standard error

were determined using the function "lm" in R (version 4.2.2) and Rstudio (Version

2022.12.0+353), and the goodness of fit of the first-order decay model was assessed through

examination of the coefficient of determination (R^2).

Free chlorine disinfection. Previous work has indicated that enveloped viruses are

particularly susceptible to free chlorine exposure[27], and our own pilot experiments suggested that

coronaviruses are as well (data not shown). We therefore created a modified lab-scale continuous

quench-flow system (**Figure S1**) to investigate the free chlorine disinfection of coronaviruses.

This system allows one to measure reaction kinetics for reactions that are too fast to observe in

batch reactors.[31] In brief, free chlorine working solution and coronavirus working solution were

pumped through a six-channel syringe pump (Masterflex®), for which the flow rate was set as 30

mL h^{-1}, and then mixed in a PEEK micro static mixing tee (IDEX Health & Science). The

reacting mixture flowed through a sample loop of pre-defined length and then was quenched by a

four-fold molar excess of ascorbic acid (A61-100, Fisher Scientific) relative to the initial free

chlorine. The ascorbic acid was dissolved directly in DMEM to minimize the requirement of

dilution of coronaviruses at the end of the disinfection experiments. Reaction times were

controlled by using different sample loop lengths. As the flow rate of reacting mixture flowing

through the sample loop is fixed (i.e., 60 mL h^{-1}), the contact time of the free chlorine

disinfection process is only dependent on the length of the sample loop, and can be calculated

by:

$$t = \frac{60 \times (L \times S - V_{dead})}{2F}$$

where t is the contact time (min), 60 represents 60 min h^{-1}, L is the length of the sample loop

(cm), S is the surface area of the sample loop (1/16" OD · 0.010" ID tubing, 0.0005 cm^2), V_{dead}

is the dead volume in the micro static mixing tee (0.00095 cm^3), and F is the flow rate of the

syringe pump (30 mL h^{-1}).

The free chlorine working solution was prepared by diluting a NaOCl stock solution (Sigma-

Aldrich) in 10 mM autoclaved phosphate buffer at pH of 7.1, where HOCl and OCl$^-$ are at a ratio

of 72:28 in molar concentrations. Phosphate buffer was selected instead of phosphate buffer

saline since high levels of chloride can affect speciation of free chlorine and thus the reaction

kinetics of some organic compounds.[32] The HCoV-OC43, MHV, and TGEV working solutions

were prepared by diluting purified HCoV-OC43, MHV, and TGEV stocks at least 1:100 either in

drinking water or phosphate buffer to minimize chlorine demand introduced by DMEM residual.

The pH of the mixture of drinking water and free chlorine working solution at 1:1 ratio was 7.1.

Therefore, the pH was consistent between the chlorine disinfection experiments conducted using

drinking water or phosphate buffer. Two free chlorine concentrations of approximately 1.30 mg

L^{-1} (referred to as "low concentration" subsequently) and 2.50 mg L^{-1} (referred to as "high"

concentration" subsequently) were used to investigate the impact of free chlorine concentration

on coronavirus chlorine disinfection kinetics in water. The concentration of free chlorine

working solution was prepared as 2.60 mg L^{-1} and 5.00 mg L^{-1} as Cl$_2$ to obtain the targeted final

free chlorine concentration of 1.30 mg L^{-1} and 2.50 mg L^{-1} as Cl_2 in the reacting mixture after

mixing with the coronavirus working solution. The free chlorine concentration was measured by

a modified *N,N*-diethyl-*p*-phenylenediamine colorimetric method (see **Supplemental Material**

for details).[33] The chlorine demand of coronavirus working solutions and the actual free chlorine

concentration in the reacting mixture were measured before every free chlorine disinfection

experiment. The chlorine demand of drinking water was negligible after exposure to the highest

chlorine concentration and contact time (**Table S3**). The samples after free chlorine disinfection

experiments were immediately tested using the $TCID_{50}$ assay for the measurement of infectious

virus concentration (see below for **$TCID_{50}$ for virus titration**). For time 0 samples, coronavirus

working solutions were mixed with pre-quenched free chlorine working solutions. Three

independent experiments were conducted at each time point across all conditions shown in **Table**

S4.

The inactivation rate constants ($k_{inactivate}$) were estimated by fitting a pseudo-first-order model

to the corresponding free chlorine disinfection data from 3 replicates while excluding the data

under the lower limit of detection (LoD), since the chlorine concentration was constant:

$$ln\left(\frac{C}{C_0}\right) = \beta_1 - k_{inactivate} \times C_{Cl} \times t$$

where C is the average infectious concentration of each coronavirus at each time point ($TCID_{50}$

mL^{-1}), C_0 is the average infectious concentration of each coronavirus at time 0 ($TCID_{50}$ mL^{-1}), β_1

is the intercept, $k_{inactivate}$ is the pseudo-first-order inactivation rate contestant for each condition

(mg^{-1} min^{-1} L), C_{Cl} is the measured initial free chlorine concentration in each experiment (mg L^{-}

^{1}, **Table S4**), and t is the contact time (min, **Table S4**). $k_{inactivation}$ and its standard error were

determined using the function "lm" in Rstudio for data of $ln(C/C_0)$ versus CT ($C_{Cl} \times t$), and the

fitness of the pseudo-first-order decay model was assessed by determining the coefficient of

determination (R^2).

$TCID_{50}$ for virus titration. A endpoint dilution assay of $TCID_{50}$ was used to measure

coronavirus infectivity.[34] In brief, HCoV-OC43, MHV, and TGEV samples were serially diluted

10-fold (i.e., 10^0-10^{-7}) with DMEM supplemented with 1% P/S and 100 μL of the diluted

samples was inoculated to confluent Vero E6, DBT, and ST cells respectively, which were all

maintained in 96-well plates and rinsed with serum free DMEM once. Five replicate wells were

applied for each dilution. After incubation at 37°C/5% CO_2 for 7, 2, or 4 days for HCoV-OC43,

MHV, and TGEV, respectively, when significant cytopathic effect was observed, supernatant

was removed and the cells were stained with 1% crystal violet (HT901, Sigma-Aldrich, diluted

with ethanol) for 10 min[35]. After the crystal violet was rinsed away using tap water, the number

of positive wells that were transparent or had large clearings in the cell lawn were counted and

the infectious concentration of coronaviruses was calculated as $TCID_{50}$ mL^{-1} as per Spearman–

Kärber method[36]. The LoD was defined as the concentration where only one out of five wells

was scored as positive at the lowest dilution, corresponding to 5.0 $TCID_{50}$ mL^{-1} in this study.

Statistical Analysis. Analysis of covariance (ANCOVA) was applied to test whether two

decay or inactivation constants were significantly different using GraphPad Prism 9 software.

For the persistence results, a total of 9 pairs of comparison was conducted via ANCOVA

(HCoV-OC43/MHV, HCoV-OC43/TGEV, and MHV/TGEV in drinking water or phosphate

buffer; drinking water/phosphate buffer condition for each virus); for the disinfection results, a

total of 21 pairs of comparison was conducted via ANCOVA (drinking water/phosphate buffer

condition with low/high concentration of free chlorine for each virus; HCoV-OC43/MHV,

HCoV-OC43/TGEV, and MHV/TGEV for merged condition). The null hypotheses of no

difference between two decay or inactivation rate constants were rejected if p values were <0.05.

All data from this study (time series of measured viral concentrations, calculated k_{decay} and

$k_{inactivation}$ values, and statistical analysis results) are available in the Stanford Digital Repository

DOI: https://purl.stanford.edu/tm847rm8309.

Results

Persistence of coronaviruses in water.

Purified HCoV-OC43, MHV, and TGEV stocks were spiked into either drinking water prior

to disinfectant addition or phosphate buffer and incubated for 30 days at room temperature to

observe their decay rates. As shown in **Figure 1** and **Table S5**, the times required for a 99%

reduction in infectivity (T_{99}) of MHV and TGEV in drinking water were notably shorter, with

MHV taking 2.0 days and TGEV taking 7.1 days, whereas a significantly longer duration of 16.5

days for MHV and 19.4 days for TGEV was necessary for achieving the same level of reduction

when they were in phosphate buffer. For HCoV-OC43, the T_{99} values were 4.6 days in drinking

water and 9.0 days in phosphate buffer.

The decay rate constants k_{decay} were estimated by fitting a first-order model (**Figure 2**).

HCoV-OC43, MHV, and TGEV all exhibited significantly higher decay rate constants in

drinking water than in phosphate buffer, i.e., around 1.9-8.1 fold higher (p<0.05, **Tables S5** and

S6). In drinking water, MHV demonstrated the highest decay rate constant with a k_{decay} of

2.25±0.09 day^{-1}, while in phosphate buffer, HCoV-OC43 had the largest decay rate constant of

k_{decay} of 0.51±0.10 day^{-1} (**Table S5**). Notably, TGEV has the smallest decay rate constants in

both drinking water and phosphate buffer, i.e., 0.65±0.06 day^{-1} and 0.24±0.02 day^{-1}, respectively.

Decay rate constants for MHV and TGEV in phosphate buffer were not significantly different

(0.28 ± 0.03 day^{-1} and 0.24 ± 0.02 day^{-1} respectively, p=0.5724).

Free chlorine disinfection kinetics of coronaviruses.

The susceptibility of coronaviruses to disinfection by free chlorine was evaluated with initial

free chlorine concentrations of 1.31 ± 0.06 and 2.53 ± 0.08 mg L^{-1}, which fall in the range of 0.5-4

mg L^{-1} recommended by EPA for both effective disinfection and safety in drinking water.[2] As

shown in **Figure 3**, HCoV-OC43 was inactivated by ~ 2.1 log$_{10}$ after exposure to free chlorine

with a CT value of 0.08 mg min L^{-1} (i.e., free chlorine concentration $\cdot$ contact time: 1.30 mg L^{-1}

of free chlorine $\cdot$ 3.5 s or 2.50 mg L^{-1} of free chlorine $\cdot$ 1.8 s) in drinking water and phosphate

buffer. MHV and TGEV demonstrated greater inactivation compared to HCoV-OC43 when

exposed to an equivalent CT value of 0.08 mg min L^{-1}, exhibiting reductions in virus titer by $\sim$

2.8 log$_{10}$ and 4.0 log$_{10}$, respectively.

Although inactivation was faster at higher free chlorine concentrations, second order

inactivation rate constants ($k_{inactivate}$; mg^{-1} min^{-1} L estimated from the pseudo-first-order model

were not significantly different either with different initial free chlorine concentrations or for

different water matrices (p>0.05, **Figure 4**, **Table S7**). Thus, the inactivation rate constants for

each virus were estimated through the consolidation of data obtained from all disinfection

experiments for each virus, including exposure to low and high concentrations of free chlorine in

both drinking water and phosphate buffer, followed by fitting the data to a pseudo-first-order

model (**Table S8**). The inactivation rates of coronaviruses by free chlorine varied significantly

between viruses (p<0.05, **Table S9**). TGEV had the highest inactivation rate constant of 113.50

mg^{-1} min^{-1} L followed by MHV (81.33 mg^{-1} min^{-1} L) and HCoV-OC43 (59.42 mg^{-1} min^{-1} L).

Discussion

This study demonstrated that HCoV-OC43 and surrogate viruses MHV and TGEV are

persistent in water, and that the decay varied between water matrices and viruses. Prolonged

survival of MHV, TGEV, and SARS-CoV in laboratory buffers or water has been observed in

previous studies,[37,38] and the decay rate constants reported here are within the range of those

previously reported for viruses in water[23]. Previous work on virus persistence in water suggests

that enzymatic activity, predation, particles, and the presence of solvents, detergents, and organic

matter can accelerate virus inactivation;[25,37,39] these were all absent or present in low

concentrations in the simplified waters studied herein. The different decay rate constants of

HCoV, MHV, and TGEV in water suggests that intrinsic properties of the viruses affect their

stability in water. A meta-analysis of decay rate constants of enveloped viruses in water found

that virus type significantly affected viral persistence.[40] For example, they reported that Influenza

A subtype H11N6 had larger decay rate constants than other influenza A subtypes; enveloped

viruses from the *Togaviridae* family decayed slower than influenza A viruses and some viruses

from the Rhabdoviridae and *Coronaviridae* families.[40] As more data are collected on virus

persistence in water, it is our hope that greater insights will be developed on how virus

characteristics affect their decay.

HCoV-OC43, MHV, and TGEV all showed susceptibility to free chlorine disinfection

(Figure 3). We tested both biphasic and pseudo-first-order models for the inactivation, and the

pseudo-first-order model fits better than the biphasic model (see **Supplemental Material** for

model equation) as illustrated by lower Akaike Information Criteria (AIC) values for all free

chlorine disinfection conditions (**Table S10**). The pseudo-first-order inactivation rate constants

(using the product of free chlorine concentration and contact time (CT) as the variable) of

coronaviruses were independent of free chlorine concentration and not affected by water matrix,

though they still varied between viruses. Lipids containing saturated fatty acids are widely

recognized for their high resistance to oxidation by free chlorine,[41] particularly in contrast to

proteins and genomes.[27] However, other research has demonstrated that lipids containing

unsaturated fatty acids react rapidly with free chlorine.[42] Previous work suggests that genome

damage might be a major mechanism for free chlorine inactivation of viruses like hepatitis A

virus and bacteriophage MS2,[43,44] with adenosine and cytosine nucleobases being more chlorine-

reactive[45]. However, the three viruses studied herein (HCoV-OC43, MHV, and TGEV) have

similar proportions of chlorine-reactive bases in their genomes (71-74% as total chlorine-reactive

bases (C+A+G), **Table S11**), which indicates that their differential susceptibility to free chlorine

might not solely be driven by their genome composition. Chlorine-induced protein damage is

another significant contributor to virus inactivation by free chlorine.[27,43] Specifically, the

inactivation of *Pseudomonas phage* Phi6, a common surrogate for enveloped viruses, by free

chlorine was reportedly driven by the oxidation of nucleocapsid proteins and proteins in the

polymerase complex; free chlorine was shown to readily penetrate the envelope membrane.[27]

Methionine (Met) and cysteine (Cys) are two of the most chlorine-reactive amino acid residues

in viral proteins, with second-order reaction rate constants of $3.8 \cdot 10^7 \ M^{-1} \ s^{-1}$ and $3.0 \cdot 10^7 \ M^{-1} \ s^{-1}$,

respectively.[46] Notably, Phi6 membrane proteins predominantly lack these amino acids. In

contrast, the presence of Met and Cys is pervasive among proteins in HCoV-OC43, MHV, and

TGEV (**Table S12**). In the present study, the relationship between the absolute counts of Met

and Cys in viral proteins and virus inactivation rate constants by free chlorine varied depending

on the protein's location. For example, a negative correlation (correlation coefficient r=-0.74,

p<0.001) was observed between Met counts in spike proteins and virus inactivation rate

constants, while a positive correlation (r=0.83, p<0.001) was found between Cys counts in

membrane proteins and virus inactivation rate constants (**Table S13**). The varied correlations

between Met and Cys counts in viral proteins and virus inactivation rate constants by free

chlorine suggests the protein composition might not be the only factor driving the protein

degradation by free chlorine. Both Ye et al[27] and Sigstam et al[43] emphasized the significant role

of solvent-accessible surface area of individual amino acids in the degradation of proteins or

peptides, i.e., slower degradation of Met- and Cys- containing peptides was observed when they

were surrounded by other amino acid residues. Remarkably, despite the striking structural

similarities among HCoV-OC43, MHV, and TGEV, it is noteworthy that only HCoV-OC43

exhibits the expression of the Hemagglutinin-esterase protein HE,[47] which is located on the

enveloped membrane and might potentially play a role in the inactivation of HCoV-OC43 due to

the presence of Met and Cys. Nevertheless, the oxidation kinetics of individual proteins or

peptides in HCoV-OC43, MHV, and TGEV remain unclear, calling for further investigation.

It is increasingly recognized that various viruses, even those closely related or possessing

similar structural features, can display markedly divergent responses when exposed to

environmental water or disinfectants.[43,48] However, the underlying mechanisms driving these

distinct viral responses remain elusive, necessitating further research efforts. For example,

comprehensive analyses such as proteomic and lipidomic assessments, as well as investigations

into the degradation kinetics of individual virus peptides, could be systematically conducted on

viruses after exposure to environmental stressors or disinfectants. Moreover, advanced genetic

approaches, like sequencing, may provide insight into resultant genome damage, i.e., by

identifying and quantifying damage in specific regions of the genome.

Experimental artifacts can also significantly impact results and potentially lead to

misinterpretations in studies of virus disinfection. For instance, the impurities present in virus

stocks used during experiments can introduce additional oxidant demand, complicating virus

disinfection investigations.[30] The presence of chloride in commonly used laboratory buffers,

such as PBS buffer, may enhance the degradation of viral genomes when exposed to free

chlorine, as it can form more effective halogenating agents than HOCl (e.g., Cl_2).[49] Moreover,

given that the choice of host cell line is indispensable to virus culture and titration, the observed

kinetics of enterovirus inactivation by free chlorine have been noted to be host cell-dependent.[50]

Discrepancies in inactivation kinetics of rotavirus during free chlorine disinfection were also

observed in experiments using mice model versus cell culture model.[41] These considerations

underscore the importance of meticulous experimental design in future virus research to ensure

accurate and meaningful results.

Environmental Implication

This study addresses a significant knowledge gap by investigating the behavior of HCoVs in

water, particularly focusing on their persistence and disinfection potential. While extensive

research has been conducted on non-enveloped viruses in water,[51] limited information exists

regarding enveloped viruses like coronaviruses[23,40]. With the ongoing emergence of new

enveloped viruses with pandemic potential, understanding how enveloped viruses like HCoVs

behave in water can inform the development of water treatment and disinfection methods to

mitigate the spread of infectious diseases, particularly when waterborne transmission is a

concern. Advanced research and adaptive management strategies are critical to stay ahead of

emerging threats.

Our research on the disinfection of HCoVs by free chlorine in water has immediate practical

applications for shaping evidence-based water policies, particularly in the development of

disinfection strategies for pathogenic virus control. With a CT value of 0.08 mg min L^{-1} of free

chlorine, which can achieve at least a 2 $\log_{10}$ reduction in virus infectivity, our findings align

with the concept of low resistance of pathogenic viruses to free chlorine (CT<1 mg min L^{-1} for

99% inactivation) as defined by WHO criteria[2]or typical free chlorine residual concentrations (~

0.5-1.5 mg/L), 2 $\log_{10}$ inactivation would be achieved in <20 seconds. Note that slower

inactivation kinetics are expected for chloramines than for free chlorine disinfection. However,

utilities employing chloramines for maintenance of a distribution system residual frequently

incorporate free chlorine contact times prior to ammonia addition to form chloramines for

distribution. Even where only short free chlorine contact times are employed (e.g., 3 min), our

results suggest the potential for significant viral inactivation. While many utilities have switched

from free chlorine. This information is pertinent not only to the safety of potable water but also

for safeguarding recreational water sources like swimming pools, as both are integral to public

health. The EPA recommends a range of 0.5-4 mg L^{-1} of free chlorine for both effective

disinfection and safety in drinking water,[2] while the CDC advocates for maintaining a minimum

free chlorine concentration of 1 mg L^{-1} in swimming pools, and at least 3 mg L^{-1} in hot tubs and

spas,[52] underscoring the practical significance of our research in promoting public health and

safety.

Furthermore, this research underscores the critical importance of meticulous selection of

appropriate virus surrogates and environmentally relevant matrices, when assessing the

persistence of human pathogenic viruses in the environment and their resistance to disinfectants.

The significant variations in both persistence and resistance to free chlorine observed among

HCoV-OC43, MHV, and TGEV herein highlight that even closely related or structurally similar

viruses can exhibit markedly distinct behavior when exposed to environmental conditions and

disinfectants. The conventional practice of using a universal surrogate for multiple viruses within

the same family may no longer be applicable, a viewpoint supported by other researchers as

well[43,48]. For instance, Aquino de Carvalho et al.[48] advocate for the necessity of working with the

virus of interest in environmental persistence studies, after their evaluation of Phi6 as a surrogate

for enveloped viruses. Many previous studies on viruses in the environment have traditionally

been carried out using laboratory buffers or artificial media. However, it is important to

recognize that real environmental matrices are considerably more complex, encompassing not

only a great variety of particles, solvents, detergents, and organic materials but also microbial

communities, which all might impact the virus infectivity, persistence, and disinfection in the

environment. This emphasizes the need for a more comprehensive approach to studying viruses

in their true environmental contexts.

Acknowledgment

This work was supported by the Stanford Woods Institute for the Environment.

References
(1) *State of the World's Drinking Water: An Urgent Call to Action to Accelerate Progress on*
Ensuring Safe Drinking Water for All; Global report ISBN 978-92-4-006081-4; WHO,
UNICEF, World Bank, 2022; p 114.
https://www.who.int/publications/i/item/9789240060807 (accessed 2023-07-04).
(2) *Guidelines for Drinking-Water Quality: Fourth Edition Incorporating the First and Second*
Addenda; Global report ISBN 978-92-4-004506-4; WHO, 2022; p 614.
https://www.who.int/publications-detail-redirect/9789240045064 (accessed 2023-07-05).
(3) Mulabbi, E. N.; Tweyongyere, R.; Byarugaba, D. K. The History of the Emergence and
Transmission of Human Coronaviruses. *Onderstepoort J Vet Res* **2021**, *88* (1), 1872.
https://doi.org/10.4102/ojvr.v88i1.1872.
(4) Hamre, D.; Procknow, J. J. A New Virus Isolated from the Human Respiratory Tract.
Proceedings of the Society for Experimental Biology and Medicine **1966**, *121* (1), 190–193.
https://doi.org/10.3181/00379727-121-30734.
(5) Liu, D. X.; Liang, J. Q.; Fung, T. S. Human Coronavirus-229E, -OC43, -NL63, and -HKU1
(Coronaviridae). *Encyclopedia of Virology* **2021**, 428–440. https://doi.org/10.1016/B978-0-
12-809633-8.21501-X.
(6) Killerby, M. E.; Biggs, H. M.; Haynes, A.; Dahl, R. M.; Mustaquim, D.; Gerber, S. I.;
Watson, J. T. Human Coronavirus Circulation in the United States 2014–2017. *Journal of*
Clinical Virology **2018**, *101*, 52–56. https://doi.org/10.1016/j.jcv.2018.01.019.

(7) Su, S.; Wong, G.; Shi, W.; Liu, J.; Lai, A. C. K.; Zhou, J.; Liu, W.; Bi, Y.; Gao, G. F.
Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends in*
Microbiology **2016**, *24* (6), 490–502. https://doi.org/10.1016/j.tim.2016.03.003.
(8) Konca, C.; Korukluoglu, G.; Tekin, M.; Almis, H.; Bucak, I. H.; Uygun, H.; Altas, A. B.;
Bayrakdar, F. The First Infant Death Associated With Human Coronavirus NL63 Infection.
The Pediatric Infectious Disease Journal **2017**, *36* (2), 231.
https://doi.org/10.1097/INF.0000000000001390.
(9) Riski, H.; Hovi, T. Coronavirus Infections of Man Associated with Diseases Other than the
Common Cold. *Journal of Medical Virology* **1980**, *6* (3), 259–265.
https://doi.org/10.1002/jmv.1890060309.
(10) Kasereka, M. C.; Hawkes, M. T. Neuroinvasive Potential of Human Coronavirus OC43:
Case Report of Fatal Encephalitis in an Immunocompromised Host. *J Neurovirol* **2021**, *27*
(2), 340–344. https://doi.org/10.1007/s13365-020-00926-0.
(11) Arbour, N.; Côté, G.; Lachance, C.; Tardieu, M.; Cashman, N. R.; Talbot, P. J. Acute and
Persistent Infection of Human Neural Cell Lines by Human Coronavirus OC43. *Journal of*
Virology **1999**, *73* (4), 3338–3350. https://doi.org/10.1128/jvi.73.4.3338-3350.1999.
(12) Turgay, C.; Emine, T.; Ozlem, K.; Muhammet, S. P.; Haydar, A. T. A Rare Cause of Acute
Flaccid Paralysis: Human Coronaviruses. *J Pediatr Neurosci* **2015**, *10* (3), 280–281.
https://doi.org/10.4103/1817-1745.165716.
(13) Morfopoulou, S.; Brown, J. R.; Davies, E. G.; Anderson, G.; Virasami, A.; Qasim, W.;
Chong, W. K.; Hubank, M.; Plagnol, V.; Desforges, M.; Jacques, T. S.; Talbot, P. J.;
Breuer, J. Human Coronavirus OC43 Associated with Fatal Encephalitis. *New England*
Journal of Medicine **2016**, *375* (5), 497–498. https://doi.org/10.1056/NEJMc1509458.
(14) Nichols, G. L.; Gillingham, E. L.; Macintyre, H. L.; Vardoulakis, S.; Hajat, S.; Sarran, C.
E.; Amankwaah, D.; Phalkey, R. Coronavirus Seasonality, Respiratory Infections and
Weather. *BMC Infectious Diseases* **2021**, *21* (1), 1101. https://doi.org/10.1186/s12879-021-
06785-2.
(15) Talbot, H. K.; Shepherd, B. E.; Crowe, J. E. J.; Griffin, M. R.; Edwards, K. M.; Podsiad, A.
B.; Tollefson, S. J.; Wright, P. F.; Williams, J. V. The Pediatric Burden of Human
Coronaviruses Evaluated for Twenty Years. *The Pediatric Infectious Disease Journal* **2009**,
28 (8), 682. https://doi.org/10.1097/INF.0b013e31819d0d27.
(16) Aboubakr, H. A.; Sharafeldin, T. A.; Goyal, S. M. Stability of SARS-CoV-2 and Other
Coronaviruses in the Environment and on Common Touch Surfaces and the Influence of
Climatic Conditions: A Review. *Transboundary and Emerging Diseases* **2021**, *68* (2), 296–
312. https://doi.org/10.1111/tbed.13707.
(17) Carraturo, F.; Del Giudice, C.; Morelli, M.; Cerullo, V.; Libralato, G.; Galdiero, E.; Guida,
M. Persistence of SARS-CoV-2 in the Environment and COVID-19 Transmission Risk
from Environmental Matrices and Surfaces. *Environmental Pollution* **2020**, *265*, 115010.
https://doi.org/10.1016/j.envpol.2020.115010.
(18) Geng, Y.; Wang, Y. Stability and Transmissibility of SARS-CoV-2 in the Environment.
Journal of Medical Virology **2023**, *95* (1), e28103. https://doi.org/10.1002/jmv.28103.
(19) Parida, V. K.; Saidulu, D.; Bhatnagar, A.; Gupta, A. K.; Afzal, M. S. A Critical Assessment
of SARS-CoV-2 in Aqueous Environment: Existence, Detection, Survival, Wastewater-
Based Surveillance, Inactivation Methods, and Effective Management of COVID-19.
Chemosphere **2023**, *327*, 138503. https://doi.org/10.1016/j.chemosphere.2023.138503.
(20) Graham, K. E.; Loeb, S. K.; Wolfe, M. K.; Catoe, D.; Sinnott-Armstrong, N.; Kim, S.;

Yamahara, K. M.; Sassoubre, L. M.; Mendoza Grijalva, L. M.; Roldan-Hernandez, L.;
Langenfeld, K.; Wigginton, K. R.; Boehm, A. B. SARS-CoV-2 RNA in Wastewater Settled
Solids Is Associated with COVID-19 Cases in a Large Urban Sewershed. *Environ. Sci.*
Technol. **2021**, *55* (1), 488–498. https://doi.org/10.1021/acs.est.0c06191.
(21) Ahmed, W.; Angel, N.; Edson, J.; Bibby, K.; Bivins, A.; O'Brien, J. W.; Choi, P. M.;
Kitajima, M.; Simpson, S. L.; Li, J.; Tscharke, B.; Verhagen, R.; Smith, W. J. M.; Zaugg,
J.; Dierens, L.; Hugenholtz, P.; Thomas, K. V.; Mueller, J. F. First Confirmed Detection of
SARS-CoV-2 in Untreated Wastewater in Australia: A Proof of Concept for the
Wastewater Surveillance of COVID-19 in the Community. *Science of The Total*
Environment **2020**, *728*, 138764. https://doi.org/10.1016/j.scitotenv.2020.138764.
(22) Paul, D.; Kolar, P.; Hall, S. G. A Review of the Impact of Environmental Factors on the
Fate and Transport of Coronaviruses in Aqueous Environments. *npj Clean Water* **2021**, *4*
(1), 1–13. https://doi.org/10.1038/s41545-020-00096-w.
(23) Silverman, A. I.; Boehm, A. B. Systematic Review and Meta-Analysis of the Persistence
and Disinfection of Human Coronaviruses and Their Viral Surrogates in Water and
Wastewater. *Environmental Science & Technology Letters* **2020**, *7* (8), 544–553.
https://doi.org/10.1021/acs.estlett.0c00313.
(24) Mahlknecht, J. Presence and Persistence of SARS-CoV-2 in Aquatic Environments: A
Mini-Review. *Curr Opin Environ Sci Health* **2022**, *29*, 100385.
https://doi.org/10.1016/j.coesh.2022.100385.
(25) Gundy, P. M.; Gerba, C. P.; Pepper, I. L. Survival of Coronaviruses in Water and
Wastewater. *Food Environ Virol* **2009**, *1* (1), 10–14. https://doi.org/10.1007/s12560-008-
9001-6.
(26) Bivins, A.; Greaves, J.; Fischer, R.; Yinda, K. C.; Ahmed, W.; Kitajima, M.; Munster, V. J.;
Bibby, K. Persistence of SARS-CoV-2 in Water and Wastewater. *Environ. Sci. Technol.*
Lett. **2020**, *7* (12), 937–942. https://doi.org/10.1021/acs.estlett.0c00730.
(27) Ye, Y.; Chang, P. H.; Hartert, J.; Wigginton, K. R. Reactivity of Enveloped Virus Genome,
Proteins, and Lipids with Free Chlorine and UV254. *Environ. Sci. Technol.* **2018**, *52* (14),
7698–7708. https://doi.org/10.1021/acs.est.8b00824.
(28) Körner, R. W.; Majjouti, M.; Alcazar, M. A. A.; Mahabir, E. Of Mice and Men: The
Coronavirus MHV and Mouse Models as a Translational Approach to Understand SARS-
CoV-2. *Viruses* **2020**, *12* (8), 880. https://doi.org/10.3390/v12080880.
(29) Cantero, M.; Carlero, D.; Chichón, F. J.; Martín-Benito, J.; De Pablo, P. J. Monitoring
SARS-CoV-2 Surrogate TGEV Individual Virions Structure Survival under Harsh
Physicochemical Environments. *Cells* **2022**, *11* (11), 1759.
https://doi.org/10.3390/cells11111759.
(30) Dunkin, N.; Weng, S.; Jacangelo, J. G.; Schwab, K. J. Minimizing Bias in Virally Seeded
Water Treatment Studies: Evaluation of Optimal Bacteriophage and Mammalian Virus
Preparation Methodologies. *Food Environ Virol* **2017**, *9* (4), 473–486.
https://doi.org/10.1007/s12560-017-9307-3.
(31) Buffle, M.-O.; Schumacher, J.; Salhi, E.; Jekel, M.; von Gunten, U. Measurement of the
Initial Phase of Ozone Decomposition in Water and Wastewater by Means of a Continuous
Quench-Flow System: Application to Disinfection and Pharmaceutical Oxidation. *Water*
Research **2006**, *40* (9), 1884–1894. https://doi.org/10.1016/j.watres.2006.02.026.
(32) Sivey, J. D.; McCullough, C. E.; Roberts, A. L. Chlorine Monoxide (Cl2O) and Molecular
Chlorine (Cl2) as Active Chlorinating Agents in Reaction of Dimethenamid with Aqueous

Free Chlorine. *Environ. Sci. Technol.* **2010**, *44* (9), 3357–3362.
https://doi.org/10.1021/es9038903.

(33) Association, A. P. H.; Association, A. W. W.; Federation, W. P. C. *Standard Methods for*
the Examination of Water and Wastewater. 17th Edition.; American Public Health
Association, 1989.

(34) Leibowitz, J.; Kaufman, G.; Liu, P. Coronaviruses: Propagation, Quantification, Storage,
and Construction of Recombinant Mouse Hepatitis Virus. *Current Protocols in*
Microbiology **2011**, *21* (1), 15E.1.1-15E.1.46.
https://doi.org/10.1002/9780471729259.mc15e01s21.

(35) Ghosh, S.; Kumar, M.; Santiana, M.; Mishra, A.; Zhang, M.; Labayo, H.; Chibly, A. M.;
Nakamura, H.; Tanaka, T.; Henderson, W.; Lewis, E.; Voss, O.; Su, Y.; Belkaid, Y.;
Chiorini, J. A.; Hoffman, M. P.; Altan-Bonnet, N. Enteric Viruses Replicate in Salivary
Glands and Infect through Saliva. *Nature* **2022**, *607* (7918), 345–350.
https://doi.org/10.1038/s41586-022-04895-8.

(36) Hierholzer, J. C.; Killington, R. A. Virus Isolation and Quantitation. *Virology Methods*
Manual **1996**, 25–46. https://doi.org/10.1016/B978-012465330-6/50003-8.

(37) Casanova, L.; Rutala, W. A.; Weber, D. J.; Sobsey, M. D. Survival of Surrogate
Coronaviruses in Water. *Water Research* **2009**, *43* (7), 1893–1898.
https://doi.org/10.1016/j.watres.2009.02.002.

(38) Wang, X.-W.; Li, J.-S.; Jin, M.; Zhen, B.; Kong, Q.-X.; Song, N.; Xiao, W.-J.; Yin, J.; Wei,
W.; Wang, G.-J.; Si, B.; Guo, B.-Z.; Liu, C.; Ou, G.-R.; Wang, M.-N.; Fang, T.-Y.; Chao,
F.-H.; Li, J.-W. Study on the Resistance of Severe Acute Respiratory Syndrome-Associated
Coronavirus. *J Virol Methods* **2005**, *126* (1), 171–177.
https://doi.org/10.1016/j.jviromet.2005.02.005.

(39) Ye, Y.; Ellenberg, R. M.; Graham, K. E.; Wigginton, K. R. Survivability, Partitioning, and
Recovery of Enveloped Viruses in Untreated Municipal Wastewater. *Environ. Sci. Technol.*
2016, *50* (10), 5077–5085. https://doi.org/10.1021/acs.est.6b00876.

(40) Silverman, A. I.; Boehm, A. B. Systematic Review and Meta-Analysis of the Persistence of
Enveloped Viruses in Environmental Waters and Wastewater in the Absence of
Disinfectants. *Environ. Sci. Technol.* **2021**. https://doi.org/10.1021/acs.est.1c03977.

(41) Zhang, M.; Ghosh, S.; Li, M.; Altan-Bonnet, N.; Shuai, D. Vesicle-Cloaked Rotavirus
Clusters Are Environmentally Persistent and Resistant to Free Chlorine Disinfection.
Environ. Sci. Technol. **2022**, *56* (12), 8475–8484. https://doi.org/10.1021/acs.est.2c00732.

(42) Simpson, A. M.-A.; Suh, M.-J.; Plewa, M. J.; Mitch, W. A. Formation of Oleic Acid
Chlorohydrins in Vegetables during Postharvest Chlorine Disinfection. *Environ. Sci.*
Technol. **2022**, *56* (2), 1233–1243. https://doi.org/10.1021/acs.est.1c04362.

(43) Sigstam, T.; Gannon, G.; Cascella, M.; Pecson, B. M.; Wigginton, K. R.; Kohn, T. Subtle
Differences in Virus Composition Affect Disinfection Kinetics and Mechanisms. *Applied*
and Environmental Microbiology **2013**, *79* (11), 3455–3467.
https://doi.org/10.1128/AEM.00663-13.

(44) Li, J. W.; Xin, Z. T.; Wang, X. W.; Zheng, J. L.; Chao, F. H. Mechanisms of Inactivation of
Hepatitis A Virus by Chlorine. *Appl Environ Microbiol* **2002**, *68* (10), 4951–4955.
https://doi.org/10.1128/AEM.68.10.4951-4955.2002.

(45) Dennis, W. H.; Olivieri, V. P.; Krusé, C. W. The Reaction of Nucleotides with Aqueous
Hypochlorous Acid. *Water Research* **1979**, *13* (4), 357–362. https://doi.org/10.1016/0043-
1354(79)90023-X.

(46) Pattison, D. I.; Davies, M. J. Absolute Rate Constants for the Reaction of Hypochlorous
568   Acid with Protein Side Chains and Peptide Bonds. *Chem. Res. Toxicol.* **2001**, *14* (10),
569   1453–1464. https://doi.org/10.1021/tx0155451.
570   (47) Bakkers, M. J. G.; Lang, Y.; Feitsma, L. J.; Hulswit, R. J. G.; de Poot, S. A. H.; van Vliet,
571   A. L. W.; Margine, I.; de Groot-Mijnes, J. D. F.; van Kuppeveld, F. J. M.; Langereis, M.
572   A.; Huizinga, E. G.; de Groot, R. J. Betacoronavirus Adaptation to Humans Involved
573   Progressive Loss of Hemagglutinin-Esterase Lectin Activity. *Cell Host Microbe* **2017**, *21*
574   (3), 356–366. https://doi.org/10.1016/j.chom.2017.02.008.
575   (48) Aquino de Carvalho, N.; Stachler, E. N.; Cimabue, N.; Bibby, K. Evaluation of Phi6
576   Persistence and Suitability as an Enveloped Virus Surrogate. *Environ. Sci. Technol.* **2017**,
577   *51* (15), 8692–8700. https://doi.org/10.1021/acs.est.7b01296.
578   (49) Qiao, Z.; Ye, Y.; Szczuka, A.; Harrison, K. R.; Dodd, M. C.; Wigginton, K. R. Reactivity
579   of Viral Nucleic Acids with Chlorine and the Impact of Virus Encapsidation. *Environ. Sci.*
580   *Technol.* **2022**, *56* (1), 218–227. https://doi.org/10.1021/acs.est.1c04239.
581   (50) Torii, S.; David, S. C.; Larivé, O.; Cariti, F.; Kohn, T. Observed Kinetics of Enterovirus
582   Inactivation by Free Chlorine Are Host Cell-Dependent. *Environ. Sci. Technol.* **2023**.
583   https://doi.org/10.1021/acs.est.2c07048.
584   (51) Boehm, A. B.; Silverman, A. I.; Schriewer, A.; Goodwin, K. Systematic Review and Meta-
585   Analysis of Decay Rates of Waterborne Mammalian Viruses and Coliphages in Surface
586   Waters. *Water Research* **2019**, *164*, 114898. https://doi.org/10.1016/j.watres.2019.114898.
587   (52) *Water Treatment and Testing | Healthy Swimming | Healthy Water | CDC*.
588   https://www.cdc.gov/healthywater/swimming/residential/disinfection-testing.html (accessed
589   2023-09-05).
590

594   Or?

Figure 1. Decay of coronaviruses in drinking water and phosphate buffer. Error bar represents
standard deviation. n=4 for each data point.

600
601 Figure 2. First-order decay rate constants of coronaviruses in drinking water and phosphate
602 buffer. Error bar represents standard error.
603

604
605
606　Figure 3. Inactivation of coronaviruses by free chlorine in drinking water and phosphate buffer.
607　Water Matrix represents the virus working solution matrices in free chlorine disinfection
608　experiments; High (2.53±0.08 mg L^{-1}) or low (1.31±0.06 mg L^{-1}) Chlorine Concentration
609　represents the initial free chlorine concentration used in free chlorine disinfection experiments.
610　Error bar represents standard deviation. n=3 for each data point.

611
612
613 Figure 4. Pseudo-first-order inactivation rate constants of coronaviruses by free chlorine
614 disinfection. Drinking Water or Phosphate Buffer represents the virus working solution matrices
615 in free chlorine disinfection experiments; High and low represents the initial free chlorine
616 concentration of 2.53 ± 0.08 mg L^{-1} and 1.31 ± 0.06 mg L^{-1} used in free chlorine disinfection
617 experiments, representatively. Error bar represents standard error.
618
619