The coral pathogen *Vibrio coralliilyticus* utilizes the compatible solute and signaling molecule *myo*-inositol as a sole carbon source

Katherine E. Boas Lichty and E. Fidelma Boyd

Department of Biological Sciences, University of Delaware

Abstract. Marine bacteria experience fluctuations in osmolarity that they must adapt to,

and most bacteria respond to high osmolarity by accumulating compatible solutes. The

osmotic stress response and compatible solutes used by the marine pathogen *Vibrio*

coralliilyticus are unknown. In this study, we showed that to alleviate osmotic stress *V.*

coralliilyticus biosynthesized the compatible solute glycine betaine (GB) and transported

choline, GB, ectoine, dimethylglycine, and dimethylsulfoniopropionate, for

osmoprotection, but not *myo*-inositol. *Myo*-inositol is a stress protectant and a signaling

molecule that is biosynthesized by algae. Bioinformatics analysis identified two

contiguous *myo*-inositol metabolic gene clusters in all *V. coralliilyticus* genomes. Growth

pattern analysis demonstrated that *V. coralliilyticus* utilized *myo*-inositol as a sole carbon

source, with a short 3 h lag time, and the inositol 2-dehydrogenase (*iolG)* gene was

required for this growth. Genetic analysis identified both an MFS-type (IolT1) and ABC-

type (IolXYZ) transporter in the metabolic gene clusters that transported *myo*-inositol.

Further genomics and phylogenetic analyses showed that the *myo*-inositol metabolic

gene clusters were widespread among *Vibrionaceae* (*Vibrio, Photobacterium, Enterovibrio,*

and *Grimontia* species) and acquired several times within the group. IolG was most

closely related to IolG from *Aeromonas, Pectobacterium (*formerly *Erwinia),* and *Dickeya*

species. These species contained the *iol* genes in a single metabolic gene cluster with an

iolXYZ transporter. In *Aeromonas*, the cluster was always found within mobile genetic

elements, but present only in a few species, predominantly in strains of *A. hydrophila*

and *A. dhakensis*.

Importance. Bacterial pathogens such as *V. coralliilyticus* encounter competition for

nutrient sources and have evolved metabolic strategies to adapt to nutrient

opportunities. Emerging studies show that *myo*-inositol is exchanged in the coral-algae

symbiosis, likely involved in signaling, but is also a compatible solute in algae. Our

findings that the coral pathogens *V. coralliilyticus* and *V. mediterranei* contained *myo*-

inosistol metabolism gene clusters suggest this could be an important adaptation. The

consumption of *myo*-inositol by these coral pathogens could contribute to a breakdown

of the coral-algae symbiosis by inhibition of algal survival or algal expulsion leading to

coral bleaching. Phylogenetic analysis showed that a *myo*-inositol metabolism gene

cluster is ancestral to *Pectobacterium* and *Dickeya*, but was acquired recently by

horizontal gene transfer in hypervirulent strains of *Aeromonas*.

Introduction

Vibrio coralliilyticus is a Gram-negative rod-shaped marine bacterium, a member of the family Vibrionaceae, and important pathogen of coral and larval oysters. This species was first characterized as the etiological agent for coral bleaching of Pocillopora damicornis, and has since been identified as a pathogen in numerous other species of coral (1-16). Recent studies also have shown that V. coralliilyticus and a related species V. tubiashii cause high mortalities in oyster larvae of both Pacific oyster (Crassostrea gigas) and Eastern oyster (Crassostrea virginica) in the United States (6, 17-22).

Under homeostatic conditions, the relationship between V. coralliilyticus and P. damicornis is commensal (23-25). However, with increased temperatures the abundance of V. coralliilyticus expands, and this in addition to other factors can turn the relationship pathogenic (26-34). One route of pathogenesis is the expulsion of the corals' endosymbiotic dinoflagellate algae (genus Symbiodinium) that occurs with increased V. coralliilyticus abundance (2, 12, 26, 27, 35, 36). Dinoflagellate algae cells are given protection and live intracellularly within coral cells, while providing their coral host with metabolites as a reliable food source (37-42). Corals thrive because of this mutualistic symbiosis with dinoflagellate algae and central to maintaining this relationship is metabolic exchange (43-48).

One metabolite exchanged between dinoflagellate algae and coral is myo-inositol (38, 39, 49). While the function of myo-inositol is not yet established in the coral-algae symbiosis, this compound is known to be biosynthesized by algae (43, 50, 51) and hypothesized to be transported to coral (39, 52). Myo-inositol is a six-carbon polyol compound and the most abundant form of inositol in the environment (53-55). In response to increased salinity; algae, plants, and marine invertebrates use myo-inositol as a compatible solute for osmotic stress protection to maintain cellular turgor pressure

(56-58). Additionally, in algae and plants, *myo*-inositol has been implicated as an

important signaling molecule (49, 59-61).

Studies have shown that soil bacteria and human bacterial pathogens can utilize

myo-inositol as an energy source (62, 63). Genetics and enzyme functionality studies for

bacterial *myo*-inositol catabolism genes was first established and investigated in Gram-

positive *Bacillus subtilis* (64-67) (**Fig. 1**). *Myo*-inositol is degraded by inositol 2-

dehydrogenase (IolG) into 2-keto-*myo*-inositol, which is then converted by myo-inosose

dehydratase (IolE) to 3,5/4-Trihydroxycylohexane-1,2-dione. This is subsequently

hydrolyzed into 5-deoxy-D-gluconate by the acylhydrolase (IolD). This product is then

isomerized by 5-deoxy-glucuronate isomerase (IolB) and phosphorylated by 5-dehydro-

2-deoxygluconokinase (IolC). Then 2-deoxy-5-keto-D-gluconate-6-P is converted to

dihydroxyacetone (DHA) and malonate semialdehyde by an aldolase (IolJ). Malonate

semialdehyde is converted to acetyl-CoA by the CoA-acylating methylmalonate-

semialdehyde dehydrogenase (IolA). The product DHA can enter glycolysis and acetyl-

CoA can enter the TCA cycle (65-68). The transport of *myo*-inositol into *B. subtilis* for

catabolism is performed by two major facilitator superfamily (MFS)-type transporters,

named IolT and IolF (65). Several Gram-negative bacteria have been shown to use *myo*-

inositol as a metabolite with a similar pathway, species in the genera *Klebsiella*,

Caulobacter, Rhizobium, Sinorhizobium, Pseudomonas, Yersinia, Corynebacterium, Salmonella,

and *Serratia* (69-78). For example, the human pathogen *Salmonella enterica* serovar

Typhymurium utilized *myo*-inositol as a sole carbon source, but growth on *myo*-inositol

was characterized by a long lag phase of > 45 h and a bistable phenotype (63, 71, 72, 79,

80). This species was found to use two MFS family transporters, named IolT1 and IolT2

for *myo*-inositol uptake (72). In the Alpha-proteobacteria *Caulobacter vibrioides* (formerly

crescentus) catabolism of *myo*-inositol was demonstrated and uptake was shown by an

ABC-type transporter (70, 81). *Myo*-inositol utilization has not been previously

examined in marine bacteria.

Here, we established the osmotic stress response of *V. coralliilyticus* at different

temperatures and showed a salinity growth range between 1% NaCl to 4% NaCl with

optimal growth in 2% NaCl at 28°C. Bioinformatics analysis of the genomes of *V.*

coralliilyticus identified the genes required for the biosynthesis of glycine betaine (GB)

and homologs of six compatible solute transporters. Our work determined that *V.*

coralliilyticus can uptake a range of compatible solutes for osmotic protection and

showed that GB can be biosynthesized from choline. Our analysis also identified a large

genomic region containing two distinct metabolic gene clusters with homologs of genes

for the regulation, transport, and catabolism of *myo*-inositol. Growth pattern analysis

demonstrated that *V. coralliilyticus* isolates used *myo*-inositol as a sole carbon source. We

also provided experimental evidence that these metabolic regions are required for

transport and catabolism of *myo*-inositol. Comparative genomics and phylogenetic

analyses demonstrated that the two gene clusters were acquired independently among

closely related species of the family *Vibrionaceae*. IolG from *Vibrio* species was most

closely related to IolG from *Aeromonas*, *Dickeya*, and *Pectobacterium* species. Within each

of these genera, the *iol* genes were present in a single cluster and were highly related to

one another except for the *Aeromonas* species. In this group the cluster was divergent

among species and present on a mobile genetic element indicative of recent horizontal

transfer into this genus.

Results

Vibrio coralliilyticus **has a relatively narrow salinity range compared to other** *Vibrio*

species. Previous work has shown that several *Vibrio* species can grow in 1% to 7%

NaCl without the addition of compatible solutes (82-87). To establish the ability of *V.*

coralliilyticus to thrive in a range of NaCl concentrations, we examined growth patterns

in M9 minimal media supplemented with glucose (M9G) and 1% to 5% NaCl at 21°C,

28°C 34°C and 37°C for 24 h (**Fig. 2**). Under all the conditions tested, *V. coralliilyticus*

grew optimally in M9G 2% NaCl at 28°C (**Fig. 2A**). At this temperature, the final OD

was 0.81 in 2% NaCl whereas in 1% NaCl, the final OD reached was 0.47. When the

salinity was increased to 3% NaCl the maximum growth rate and area under the curve

(0.66 h^{-1}, 15.3) were lower than what was observed at 2% NaCl (0.76 h^{-1}, 16.5). Growth of

V. coralliilyticus in 4% NaCl showed 2 h lag time and lower maximum growth rate (0.40

h^{-1}) whereas no growth occurred in 5% NaCl (**Fig. 2A**). Next, we tested whether

temperature affected the salinity growth range by examining growth at 21°C, 34°C and

37°C (**Fig. 2B-D**). Two observations were made from these analyses; first incubation at

21°C negatively affected growth in 4% NaCl resulting in a significantly longer lag phase

than at any other temperature (**Fig. 2B**). Secondly, incubation at 37°C impacted the

ability to grow in low salinity; at this temperature there was no growth in 1% NaCl (**Fig.**

2D). Overall, growth in 1% NaCl at 28°C, 34°C and 37°C resulted in a final OD lower

than what was observed at the higher NaCl concentrations up to 4% NaCl (**Fig. 2**).

***V. coralliilyticus* biosynthesizes glycine betaine (GB) and uptakes numerous**

compatible solutes for osmotic protection. Analysis of *V. coralliilyticus* genomes

identified a gene cluster involved in the biosynthesis of GB from the precursor choline

(*betIBA*) and six putative compatible solute transporters (**Fig. 3A**). ^{1}H-NMR spectra

analysis of ethanol extracts for *V. coralliilyticus* grown in M9G 4% NaCl supplemented

with 1 mM choline confirmed glycine betaine biosynthesis (**Fig. 3B**). To establish the

range of compatible solutes taken up by *V. coralliilyticus* for osmotic protection, cells

were grown in M9G 5% NaCl supplemented with one of ten different compatible

solutes. The growth pattern in each compatible solute was monitored for 24 h at 28°C.

Seven of the ten compatible solutes tested restored growth (**Table 1**). GB,

dimethylsulfoniopropionate (DMSP), ectoine, choline, and dimethylglycine (DMG)

rescued growth to the greatest extent reaching a final OD ranging from 0.45 to 0.69

(**Table 1**). The compatible solutes γ-amino-butyric acid (GABA) and creatine had longer

lag phase times and reached a final OD of 0.23 and 0.31, respectively (**Table 1**).

Trimethylamine-N-oxide, *myo*-inositol (MI), and glycerol were unable to rescue growth

of *V. coralliilyticus* in M9G 5% NaCl indicating they are not used for osmotic protection

in this species (**Table 1**).

***Myo*-inositol metabolic gene clusters are present in all strains of the coral pathogens**

***V. coralliilyticus* and *V. mediterranei*.** Bioinformatics analysis identified putative *myo*-

inositol metabolic gene clusters that were present in all *V. coralliilyticus* genomes

examined. The region contained two distinct gene clusters for the regulation, transport,

and catabolism of *myo*-inositol (**Fig. 4**), named region 1 and region 2. Gene annotation

and naming were established based upon amino acid similarity to *Salmonella enterica*.

The gene clusters and gene order were highly conserved among *Vibrio* species. No

synteny was found between *myo*-inositol gene clusters in *Vibrio* species and *S. enterica*,

B. subtilis, or *C. vibrioides* (**Fig.4**). Functionally similar genes present in region 1 and

region 2 in *V. coralliilyticus* shared low sequence similarity (27% to 50%) and lacked

gene synteny.

Region 1 contained fifteen genes, which included a predicted operon composed

of ten genes *iolEDOCBJRIBG* encoding putative enzymes responsible for *myo*-inositol

conversion to dihydroxyacetone and malonate semialdehyde (**Fig.4**). Region 1 also

contained two MFS transporters, a LysR family transcriptional regulator, a

glycosyltransferase and a polysaccharide biosynthesis protein. A single copy of the *iolG*

gene (inositol 2-dehydrogenase) was present in the metabolic gene clusters, this enzyme

is required for the first step in the pathway (**Fig. 1**). IolG shares 36.1% amino acid

identity to IolG2 from *S. enterica* (**Fig.4**). A copy of *iolE* and *iolD* are present within

region 1 and encode enzymes involved in the second and third steps of the catabolic

pathway. Two copies of IolE and IolD are present, a copy in each region. Region 1

copies are named IolE2 and IolD2 as they share only 38% amino acid identity to the

enzymes from *S. enterica* (**Fig.4**). A putative 5-dehydro-2-deoxyglucokinase is encoded

by *iolC* is also found in region 1 and shares 33% amino acid identify with IolC from *S.*

enterica. The next step in the pathway is completed by an oxidoreductase encoded by

iolB; two copies of this gene were found in region 1 (IolB1 and IolB2), sharing 63.5% and

31% amino acid identify to IolB from *S. enterica* (**Fig. 4**). Region 1 also contained a

hypothetical Gfo/Idh/MocA family oxidoreductase that we named *iolO*.

Region 2 contained a predicted operon *iolCDEA*, which overall shared greater

amino acid identity ranging from 67.4% to 43.7% with proteins from *S. enterica* (**Fig.4**).

In region 2, an *iolA* homolog shared 65.2% amino acid identity to IolA from *S. enterica*

and encoded a CoA-acylating methylmalonate-semialdehyde dehydrogenase involved

in the conversion of malonate semialdehyde to acetyl-CoA. There is no homolog for

IolA in region 1. The *iolCDE* genes were named *iolC1D1E1* as they share the closest

homology to proteins in *S. enterica*. Divergently transcribed from this operon is the

putative operon *iolXYZR* that encodes an ABC-type transporter IolXYZ and an RpiR

type regulator IolR (**Fig.4**). *S. enterica* does not possess an ABC-type transporter that

uptakes *myo*-inositol therefore this transporter was compared to another Gram-negative

species known to transport *myo*-inositol for catabolism. The substrate binding protein

IolX from *V. coralliilyticus* shares 31% amino acid identity with IbpA a known

transporter of *myo*-inositol in *Caulobacter vibrioides* (81).

A similar gene arrangement in region 1 was present in the genomes of the coral

pathogen *V. mediterranei* (formerly *V. shilonii*) and related species *V. maritimus* and *V.*

barjaei (**Fig.4**). However, in these species region 1 was contiguous with region 2, and this

cluster contained the operon *iolCA* divergently transcribed from *iolXYZR* (**Fig.4**).

200 *V. coralliilyticus* **can utilize *myo*-inositol as a sole carbon source.** To demonstrate *myo*-

201 inositol catabolism by *V. coralliilyticus*, strain CN52H-1 was grown in M9 2% NaCl

202 media supplemented with 20 mM Glucose or 20mM *myo*-inositol as carbon sources at

203 28°C for 30 h. *V. coralliilyticus* growth in M9 2% NaCl with 20mM *myo*-inositol reached a

204 final OD of 0.29 with a lag time of 3 h (**Fig. 5A**). We also examined *V. harveyi* growth, a

205 species that contains region 1, but not region 2, and found that this species grew in

206 glucose but did not grow in *myo*-inositol as a sole carbon source (**Fig. 5B**). This

207 suggested that region 2 is essential for *myo*-inositol catabolism. To demonstrate that

208 region 1 is also required for *myo*-inositol catabolism, an in-frame deletion of *V.*

209 *coralliilyticus iolG* (HRJ43_RS06535) was generated and the Δ*iolG* strain showed no

210 growth in M9 2% NaCl media supplemented with 20 mM *myo*-inositol (**Fig. 5C**).

211 Overall, the data showed that *V. coralliilyticus* has a functioning *myo*-inositol pathway

212 and both region 1 and region 2 are required for catabolism in *Vibrio* species.

213 **Ligand modeling predicts *V. coralliilyticus* IolT1 and IolXYZ are *myo*-inositol**

214 **transporters**. Three transporters, two MFS-type transporters named IolT1 and IolT2

215 within region 1 and one ABC-type transporter IolXYZ within region 2 were identified.

216 Ligand modeling was performed to predict which of these are likely responsible for

217 *myo*-inositol transport by comparison to known transporters of *myo*-inositol. The MFS

218 transporters from *B. subtilis* IolT (WP_003234027.1) and *S. enterica* IolT1 (NP_463279.1)

219 were selected as these previously showed transport of *myo*-inositol (65). *V. coralliilyticus*

220 IolT1 (WP_099607352.1) had 48.8% amino acid identity to that of *S. enterica* IolT1

221 whereas *V. coralliilyticus* IolT2 (WP_172853823.1) shared only 27% similarity. Docking of

222 *myo*-inositol with *B. subtilis* IolT and *S. enterica* IolT1 predicted a mode of interaction

223 with seven and five putative binding residues, respectively and favorable binding free

224 energy changes (ΔG) of -7.0 kcal/mol and -6.1 kcal/mol. *Myo*-inositol docking with *V.*

225 *coralliilyticus* IolT1 predicted a favorable ΔG of -6.2 kcal/mol and four putative binding

residues. Alignment of *B. subtilis* IolT, *S. enterica* IolT1, *V. coralliilyticus* IolT1 and IolT2 was performed and predicted transmembrane domains are shown (**Fig. 6A**). MFS-type transporters are known to coordinate substrate binding with transmembrane (TM) domains TM7 and TM10 (88). Predicted binding residues found in the three models are shown with blue (*V. coralliilyticus* IolT1), green (*S. enterica* IolT1), and orange (*B. subtilis* IolT) arrows. (**Fig. 6B**). The two glutamine (Q273, Q274) and asparagine (N279) in TM7 are found in all three modeled sequences. In TM10 three of the four sequences also contained a conserved tryptophan (W377) residue. All three models predicted a residue that aligns at Y368. Both *V. coralliilyticus* IolT1 and *S. enterica* IolT1 contain a conserved tyrosine (Y) residue here, but *B. subtilis* IolT contained glutamine (Q) (**Fig. 6A**). Taken together this similarity in the binding pockets of these transporters suggested that *V. coralliilyticus* IolT1 is likely able to transport *myo*-inositol.

The ABC-type transporter substrate binding protein IbpA from *C. vibrioides* had previously been shown to bind *myo*-inositol (81) and IolX (WP_227495657.1) from *V. coralliilyticus* had 87% coverage, 31% amino acid identity to IbpA. Ligand docking analysis of *myo*-inositol with IbpA and IolX predicted favorable modes of action with ΔG of -7.4 kcal/mol and -5.3 kcal/mol respectively. Putative binding residues were identified for each substrate binding protein and are presented in the amino acid alignment (**Fig. 6B**). The green arrows indicate IbpA predicted binding residues and blue arrows indicate IolX predicted binding residues. Five aligned residues were predicted to be important in both docking models: 63 aspartic acid (D)/ glutamine (Q), 141 aspartic acid (D)/ asparagine (N), 187 glutamine (Q)/ serine (S), 269 aspartic acid (D), and 289 glutamine (Q) (**Fig. 6B**). Based upon the favorable docking model and select conserved residues in the alignment with a known *myo*-inositol transporter, IolXYZ is predicted to be capable of *myo*-inositol transport.

The MFS transporter IolT1 and ABC-type transporter IolXYZ are responsible for *myo*-inositol uptake for catabolism. To evaluate the ability of IolT1 and IolXYZ to transport *myo*-inositol, in-frame deletion mutants were generated in the *iolT1* (HRJ43_RS06585) and *iolX (HRJ43_RS06670)* genes. The Δ*iolT1* mutant grew in *myo*-inositol as the sole carbon source with a lag time of 6 h compared to wild type with a lag time of 3 h and decreased maximum growth rate (Δ*iolT1;* 0.11 h^{-1}, wild type; 0.29 h^{-1}) (**Fig. 7**). Additionally, Δ*iolT1* reached a lower final OD (0.12) as compared to wild type (0.29) (**Fig. 7**). This data indicated that IolT1 is utilized for the transport of *myo*-inositol, but another transporter also uptakes this compound. Similarly, the growth pattern analysis of the Δ*iolX* mutant showed growth in *myo*-inositol with a final OD of 0.17 compared to an OD of 0.29 in wild type (**Fig. 7**). *V. coralliilyticus* Δ*iolX* displayed a maximum growth rate of 0.10 h^{-1} as compared to 0.29 h^{-1} observed for wild type (**Fig. 7**). These data suggested IolXYZ also transported *myo*-inositol. Next, we constructed a double mutant Δ*iolT1*/Δ*iolX* to determine whether any other transporters could uptake *myo*-inositol. The double mutant failed to grow in M9 supplemented with *myo*-inositol as a sole carbon source and demonstrated that *V. coralliilyticus* has two transporters for *myo*-inositol uptake for catabolism (**Fig. 7**).

Phylogenetic analysis of IolG from region 1 and IolA from region 2 among *Vibrionaceae* reveals different evolutionary histories. The low sequence similarity between comparable proteins in each of the metabolic gene clusters and lack of synteny suggested different evolutionary histories of these regions. To examine this further, a phylogeny based on the IolG protein from region 1 was constructed among members of the family *Vibrionaceae* (**Fig. 8**). In addition, a phylogeny based on the housekeeping protein RpoB was also constructed among the same set of species to represent their ancestral relationships (**Fig. 9**). BLAST analysis was performed using IolG (WP_038162996.1), the inositol-2-dehydrogenase from region 1, as a seed to identify

protein sequences in the NCBI genome database. Sequences with >95% query coverage

and >60% amino acid similarity were downloaded and aligned using CLUSTALW (89).

The genome context of these IolG proteins within each species was determined and all

IolG proteins showed the presence of additional *myo*-inositol metabolism genes.

Phylogenetic trees were constructed, and a representative tree is presented in **Figure**

8A. IolG was present in *Vibrio* (red), *Photobacterium* (blue), *Enterovibrio* (orange), and

Grimontia (brown) species. Among *Vibrio* species, the IolG proteins are found dispersed

within three major branches on the tree. The *V. coralliilyticus* IolG clustered divergently

from members of the Mediterranei clade (**Fig. 8A**). Comparative genomics showed that

in the Mediterranei group region 1 was contiguous with region 2 whereas in *V.*

coralliilyticus the regions were separated by between 6 kb to 28 kb depending on the

strain. Branching distantly from these species were a highly similar cluster of IolG

proteins indicated by short branch lengths with species in the Harveyi clade. Within

this tight cluster was IolG from *V. vulnificus*, a species which is not a member of the

Harveyi clade. These data indicated that IolG and by extension region 1 were acquired

by horizontal gene transfer in *V. vulnificus* from a *Vibrio* species, region 2 was absent

among these species (**Fig. 8B**). The second major clustering of IolG from *Vibrio* species

contained long-branch lengths and relatively low bootstrap values. In this group, IolG

from *V. kanaloae, V. toranzoniae*, and *V. rumoiensis* clustered together. Branched distantly

from this group were IolG from two *Photobacterium* species that clustered divergently

from *V. anguillarum, V. ponticus*, and *V. metschnikovii* (**Fig. 8**). The *myo*-inositol gene

clusters in these species shared a similar gene order comprised of *iolDCEA* contiguous

with *iolB, iolR, iolG*, and *iolXYZ*, which differed significantly from the gene order

present in all other *Vibrio* species (**Fig. 8B**). The third cluster of IolG proteins from *Vibrio*

species contained IolG from four members of the Splendidus clade, which branched

divergently from IolG in nine *Photobacterium* species (**Fig. 8**). These all contained a

similar region 1 as the Harveyi clade, but region 2 showed different gene contents (**Fig.**

8B). The IolG proteins from five *Grimontia* species clustered closely together with short branch lengths and grouped with IolG from *Enterovibrio calviensis and E. norvegicus* (**Fig. 8**). IolG was also present in four additional *Enterovibrio* species, branched separately from the other two *Enterovibrio* species. Lastly, we observed *V. penaeicida* and *V. nigripulchritudo* clustering divergently from the *Enterovibrio* and *Grimontia* groups.

The IolA protein, a CoA-acylating methylmalonate-semialdehyde dehydrogenase, was selected as the representative of region 2 and used as a seed for BLAST analysis. Eight *Vibrio* species that contained IolG did not have a copy of IolA or any of the duplicate proteins present in region 2. These included species from the Harveyi clade, *V. harveyi, V. campbellii, V. owensii,* and *V. jasicida,* as well as *V. astriarenae, V. agarivorans, V. ponticus,* and *V. vulnificus.* In the IolA tree, the bootstrap values for internal branch nodes for most *Vibrio* and *Photobacterium* species was low and indicated that the relationships of IolA from these species is not clear (**Fig. 9**). In contrast, the bootstrap values for IolA from *Grimontia* species were robust with short branch lengths and indicated that they are highly related (**Fig. 9**). Branched divergently from these were IolA from *Enterovibrio* species with greater than 85% confidence in many external nodes and IolA from *V. penaeicida* and *V. nigripulchritudo* also clustered distantly within this group (**Fig. 9**). The position of IolA from *Photobacterium* species within the tree could not be determined as the internal bootstrap values were < 40%. In most *Vibrio* species, we observed divergent branches with long-branch lengths, which represented low IolA similarity to each other and a polyphyletic origin for IolA. The IolA protein from *Photobacterium chitinilyticum*, but not *P. atrarenae,* similar to the IolG phylogeny, branched distantly with IolA from *Vibrio* species.

The differences between the IolG and IolA trees and the comparative genomic analysis showed that the two regions were acquired separately likely via horizontal transfer, each with a different origin and ancestral history. Overall, the data suggested a

polyphyletic origin for the *myo*-inositol metabolic gene clusters and that a similar region

1 was acquired by *Vibrio* and *Photobacterium* species. Whereas a separate unrelated

region 1 was acquired by *Grimontia* and *Enterovibrio* species and a handful of *Vibrio*

species. The data also showed that when a region was present in a species, it was

present in all representatives of that species in the NCBI genome database. In addition,

there were no signatures of recent horizontal transfer such as the presence of IS

elements, transposases, integrases, phage, or plasmid elements within 10 genes of the

metabolic gene clusters.

To examine *myo*-inositol evolutionary history further, an IolG tree was

constructed from species outside of the *Vibrionaceae* family (**Fig. 10**). Within this tree,

IolG from *Vibrio* species clustered most closely with members of the genera *Aeromonas*,

Pectobacterium (formerly Erwinia species), and *Dickeya* with each genus tightly clustered

within distinct clades (**Fig. 10**). Genomic analysis of *Pectobacterium (formerly Erwinia),*

and *Dickeya* showed that the *myo*-inositol metabolic genes were present in a single

cluster; in *Pectobacterium* the operon structure was *iolADGXYZOCEB* with the ABC-type

transporter. In *Dickeya* species, the region was similar to what was seen in

Pectobacterium with the exception of three genes inserted between *iolD* and *iolG*. For

both these genera, IolG showed short branch lengths indicating these proteins are

highly related to one another within each genus. In six *Aeromonas* species, the

distribution and divergence of the region and proteins suggested the *iol* cluster was

acquired via horizontal gene transfer. The *iol* metabolism genes were present as a single

cluster, but the cluster differed significantly between species. In a large group of *A.*

hydrophila strains, the region contained the *iolXYZ* transporter genes and IolG clustered

divergently with *Pectobacterium* and *Dickeya* species. An *iol* cluster containing a Sodium

Solute Symporter (SSS) type transporter was present in all other *Aeromonas* species

examined and in these species was only present in a few to one isolate of each species.

In *A. hydrophila* strains, the cluster was inserted adjacent to a tRNA-Leu locus and

clustered with integrase and phage genes, markers of horizontal gene transfer. In *A.*

dhakensis strains, also an important pathogen of fish, insertion occurred at the tRNA-Ser

locus, a 27 gene region with homology to phage genes and a restriction modification

system was present followed by the *iol* gene cluster. In *A. media, A. allosaccharophila,* and

A. finlandersis, the *iol* gene cluster was also present adjacent to a tRNA-Ser locus but had

different flanking genes and was only present in one strain of these species.

Discussion

Our examination of *V. coralliilyticus* response to changes in osmotic stress under

different growth temperatures demonstrated the importance of NaCl for optimal

growth and growth at high temperatures. We have shown that *V. coralliilyticus* can

grow in 1% - 4% NaCl but grows optimal in 2% NaCl at 28°C and has an absolute

requirement for NaCl for growth at 37°C. Compared to other *Vibrio* species that grow in

up to 7% NaCl without the addition of compatible solutes, *V. coralliilyticus* has a

relatively low salinity tolerance of 4% NaCl. A major difference between *V. coralliilyticus*

and these *Vibrio* species is the absence of the ectoine biosynthesis cluster *ectABC*, which

is required for biosynthesis of ectoine *de novo*. However, similar to other *Vibrio* species,

V. coralliilyticus could uptake a wide range of compatible solutes from their surrounding

environment such choline, GB, DMSP, DMG, and ectoine for osmotic protection. *Myo-*

inositol is a compatible solute more commonly associated with a few hyperthermophile

bacteria as well as many eukaryotic species such as algae and plants (90-93). Therefore,

it came as no surprise that *V. coralliilyticus* did not use *myo*-inositol for osmotic

protection. However, our bioinformatics analysis did identify a genomic region on

chromosome 2 that contained *myo*-inositol metabolic gene clusters present in all *V.*

coralliilyticus isolates. We showed *V. coralliilyticus* can utilize *myo*-inositol as a sole

carbon source with a 3 h lag phase. This is significant since previous studies on the

Gram-negative human pathogens such as *Salmonella enterica* have shown a lag time of over 40 hours (69, 71, 72, 75). The Gram-negative soil bacteria *Caulobacter vibrioides* (formerly *crescentus*) (70, 81), *Rhizobium leguminosarum* (formerly *trifolii*) (76), and *Sinorhizobium meliloti* (94) were also shown to have a long lag times of about 14 h when grown on *myo*-inositol as a sole carbon source (81). A possible reason for this efficient use of *myo*-inositol in *V. coralliilyticus* is the presence of two transporters for uptake into the cell, an MFS-type IolT and an ABC-type family transporter IolXYZ. As shown in the deletion mutant analysis of these systems, both have somewhat similar *myo*-inositol uptake affinities. Alternatively, the regulation of the catabolism genes may be different in these species.

Examination of the evolutionary history of the proteins IolG and IolA among *Vibrionaceae* demonstrated that the two regions were acquired separately in many species indicating convergent evolution of this phenotypic trait. The closest relative to IolG outside of the Vibrionales were members of the Enterobacterales (*Dickeya*, *Pectobacterium*) and Aeromonadales (*Aeromonas*). In all fifteen *Pectobacterium* species, the genes are present within a highly homologous single cluster indicating it was acquired once within the genus (**Fig. 10**). In the ten *Dickeya* species examined a similar gene cluster as found in *Pectobacterium* was present with the addition of 3 genes inserted between *iolD* and *iolG* (**Fig. 10**). Two different *myo*-inositol clusters were present in *Aeromonas* species. In the hypervirulent *A. hydrophila* species associated with epidemic disease in fish aquaculture farms in China and the USA, the single *iol* cluster contained the IolXYZ ABC-type transporter and was inserted at a tRNA-Leu locus. This insertion was within a region that contained an integrase and phage genes on one flank and an IS3 element on the opposite flank. Previous studies have showed that hypervirulent *A. hydrophila* strains all contained an *iol* cluster and could utilize *myo*-inositol as a sole carbon source. This is an important trait differentiating these strains from other *A.*

hydrophila isolates (95-97). In *A. hydrophila* strains WCX23 and 23-C-23, the *iol* metabolic

gene cluster was inserted next to a tRNA-His locus, contained a sodium solute

symporter (SSS) and was flanked on both side by genes encoding a type six secretion

system (T6SS). Overall, the data suggest that in *Aeromonas,* the *iol* gene cluster is likely a

recent addition acquired on a mobile genetic element. This contrasts with what we

observed in *Vibrio* species.

Heightened ocean temperatures have resulted in increased *Vibrio* species abundance

worldwide (4, 29, 98-101). The increased abundance of *V. coralliilyticus* during ocean

warming has also been documented and believed to be a major contributing factor to

coral bleaching(2). We hypothesize that this increased abundance could contribute to a

breakdown in the coral-algae symbiosis. Perhaps *V. coralliilyticus* consumption of *myo-*

inositol is depleting the availability of *myo*-inositol as a stress protectant for algae,

which could then lead to a decline in algal abundance. In addition, *myo*-inositol is used

as a building block in various signal transduction pathways, and the ability of *V.*

coralliilyticus to catabolize this sugar poses a means for signaling interference. Both

potentially contributing to increased *myo*-inositol consumption and an inhibition of

algal survival eventually leading to coral bleaching. Alternatively, the consumption of

myo-inositol by *V. coralliilyticus* may be a fitness factor that gives it a competitive

advantage over members of the coral microbiome.

Materials and Methods

Bacterial strains, media, and culture conditions. All strains and plasmids used in this

study are listed in **Table 2**. *Vibrio coralliilyticus* strain CN52H-1 (7) was used in this

study. Unless stated otherwise, *V. coralliilyticus* was grown at 28°C in either lysogeny

broth (LB) (Fisher Scientific, Fair Lawn, NJ) with 2% (wt/vol) NaCl (LB 2%NaCl) or M9

minimal media (47.8 mM Na_2HPO_4, 22 mM KH_2PO_4, 18.7 mM NH_4Cl, 8.6 mM NaCl;

Sigma Aldrich) supplemented with 2 mM $MgSO_4$, 0.1 mM $CaCl_2$, 2% (wt/vol) NaCl and

20 mM glucose as the sole carbon source (M9G 2% NaCl). *Escherichia coli* strains were

grown in LB supplemented with 1% (wt/vol) NaCl (LB1%) at 37°C. The strain *E. coli*

β2155 *λpir* is a diaminopimelic acid (DAP) auxotroph and was grown supplemented

with 0.3 mM DAP (102). The *V. harveyi* strain used for analysis was strain 393 (103).

Chloramphenicol (Cm) was added to media at a concentration of 12.5 μg/mL when

specified.

Osmotic Stress Growth Analysis. *Vibrio coralliilyticus* was grown overnight in M9G 1%

NaCl at 28°C with aeration. A 2% inoculation of the overnight culture into fresh M9G

1% NaCl was grown to OD ~ 0.50 at 28°C with aeration. Exponential cultures were

inoculated 1:40 into 200 μL fresh M9G media containing 1% to 6% NaCl. When using

compatible solutes glycine betaine (GB), dimethylsulfoniopropionate (DMSP), ectoine,

choline, dimethylglycine (DMG), trimethylamine N-oxide (TMAO), gamma-

aminobutyric acid (GABA), creatine, *myo*-inositol (MI), and glycerol were used at a final

concentration of 1 mM in M9G 5% NaCl. Plates were incubated at 28°C with

intermittent shaking for 1 min every hour. Optical densities were taken at 595 nm every

hour for a total of 24 h using a Tecan Sunrise microplate reader and Magellan plate

reader software (Tecan Systems Inc., San Jose, CA).

Deletion Mutant Construction. In-frame nonpolar deletion mutants of *iolG*

(HRJ43_RS06535), *iolT1* (HRJ43_RS06585), and *iolX* (HRJ43_RS06670) were constructed

using SOE PCR and allelic exchange as previously described (104). Truncations of the

genes of interest, *iolG* (18-bp of the 984-bp gene), *iolT1* (30-bp of the 1452-bp gene), and

iolX (24-bp of the 1008-bp) were generated using the primers listed in **Table 3**. The

truncated gene products were ligated with the suicide vector pDS132 using the Gibson

assembly protocol and then transformed into *E. coli* Dh5α *λ*pir (105, 106). The plasmids

pDSΔ*iolG*, pDSΔ*iolT1*, and pDSΔ*iolX* were then purified and transformed into *E. coli*

β2155 *λpir* (DAP auxotroph). Then conjugation followed by homologous recombination

into the *V. coralliilyticus* CN52H-1 genome was completed. To select for the single

crossover of the plasmids into the genome, plating onto chloramphenicol (Cm) was

used and colonies were screened via PCR for the truncated allele. To induce a double-

crossover event, the single-cross strains were grown overnight in the absence of Cm.

The cultures were then spread on 10% sucrose plates, and healthy colonies were

screened for gene truncation, and plasmid excision. Colonies that still contained the

plasmid would appear soupy on the 10% sucrose plates because of the presence of *sacB*

selectable marker. In-frame deletions were confirmed with sequencing.

Metabolic Growth Analysis. Strains were grown overnight in M9G 2% NaCl at 28°C

with aeration. Overnight cultures were washed twice with PBS and then inoculated 1:40

into 200 μL fresh M9 media supplemented with 2% NaCl, and specified carbon source

(glucose (20 mM) or *myo*-inositol (20 mM)). Plates were incubated at 28°C with

intermittent shaking for 1 min during every hour. Optical densities were taken at 595

nm every hour for a total of 30 h using a Tecan Sunrise microplate reader and Magellan

plate reader software (Tecan Systems Inc., San Jose, CA).

Preparation of cellular extracts and proton nuclear magnetic resonance (^{1}H-NMR).

Wild type *V. coralliilyticus* was grown overnight at 28°C in M9G 4% NaCl supplemented

with 1 mM choline. Stationary phase cells were then pelleted and washed twice with

PBS. Three freeze-thaw cycles were performed with the cell pellets to increase lysis, and

the cells were then resuspended in 750 μL ethanol. Debris was pelleted by

centrifugation, and the ethanol solution was transferred to a clean tube and evaporated

under vacuum. Then deuterium oxide (D_2O) was used to resuspend the pellet, and

insoluble material was removed by centrifugation. The solution was then transferred to

a 5-mm NMR tube for analysis on a Bruker AVANCE 600NMR spectrometer at a proton

frequency of 600.13 MHz with a sweep of 12,376 Hz and a relaxation delay of 5 s.

Sixteen scans were co-added for each spectrum.

Phylogenetic analysis. Phylogenetic analysis was conducted using the program MEGA

X (107). The evolutionary history of IolG, IolA and RpoB was inferred using the

Neighbor-Joining method (108). The evolutionary distances were computed using the

JTT matrix-based method and are in the units of the number of amino acid substitutions

per site (109). The rate variation among sites was modeled with a gamma distribution

(shape parameter = 5). All ambiguous positions were removed for each sequence pair

(pairwise deletion option).

Ligand Docking Modeling. The transporter amino acid sequences were downloaded

from NCBI and aligned using ClustalW(89): *B. subtilis* IolT (WP_003234027.1), *S. enterica*

IolT1 (NP_463279.1), *V. coralliilyticus* IolT1 (WP_099607352.1), *V. coralliilyticus* IolT2

(WP_172853823.1), *C. crescentus* IbpA (WP_010918744.1), and *V. coralliilyticus* IolX

(WP_227495657.1). Transporter alignments were displayed and annotated using ESPript

(http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi) (110). Obtained predicted protein

structures from AlphaFold (111). The Ligand model for *myo*-inositol (MI) was obtained

from Chemical Entities of Biological Interest (CheBI) EMBL (112). Polar hydrogens were

added to the ligand structures. The docking experiments were performed using

SeamDock which was set to use docking parameters for the Vina software (Spacing: 1.0,

Mode: 2, Energy Range: 5, Exhaustiveness: 8) (113).

Figure Legends

Fig. 1. Pathway of m*yo*-inositol catabolism. The pathway presented is based upon enzymological studies in *B. subtilis* and genome comparison. The proteins and intermediate compounds in the pathway are displayed.

Fig. 2. *Vibrio coralliilyticus* NaCl tolerance range. Growth curve analysis of *V. coralliilyticus* CN52H-1 in minimal media with glucose (M9G) containing 1% to 5% NaCl. Growth was measured every hour for 24 h at 28°C. Mean and standard deviation of two biological replicates are shown.

Fig. 3. *V. coralliilyticus* compatible solute biosynthesis and uptake systems. A. Predicted genes for glycine betaine biosynthesis and compatible solute transporters in *V. coralliilyticus* **B. Glycine betaine biosynthesis from choline.** H-NMR spectra of *V. coralliilyticus* CN52H-1 cellular extract grown in minimal media (M9G) with 4% NaCl with the addition of 1mM choline. The spectra peaks corresponding to glycine betaine are labeled (2 and 3) and the chemical structure is provided.

Fig. 4. High variability in chromosomal organization of *iol* genes among bacteria. Schematic of the gene order of myo-inositol transporter, catabolism and regulatory genes among previously characterized species and *Vibrio*. Arrows represent open reading frames and direction of arrow represents direct of transcription. Arrows of similar color represent similar proteins/genes.

Fig. 5. *V. coralliilyticus* utilized *myo*-inositol as a sole carbon source. Growth curve analysis in minimal media (M9) containing either 20 mM Glucose (black squares) or 20mM *myo*-inositol (blue triangles). **A.** *V. coralliilyticus* CN52H-1. **B.** *V. harveyi* 393 **C.** *V. coralliilyticus* Δ*iolG.* Growth was measured every hour for 30 h at 28°C. Mean and standard deviation of two biological replicates are shown.

Fig. 6. Favorable *myo*-inositol ligand docking predicted for *V. coralliilyticus* transporters. Results of *myo*-inositol docking study showing putative residues mediating the interactions with the ligand. Residues that are predicted to interact with *myo*-inositol are shown with an up arrow. **A.** MFS-type transporters; *V. coralliilyticus* IolT1 (WP_099607352.1, blue arrows), *V. coralliilyticus* IolT2 (WP_172853823.1), *B. subtilis* IolT (WP_003234027.1, orange arrows), *S. enterica* IolT1 (NP_463279.1, green arrows). Both *B. subtilis* IolT and *S. enterica* IolT1 take up *myo*-inositol. **B.** ABC-type transporter substrate binding proteins; *V. coralliilyticus* IolX (WP_227495657.1, blue arrows) and *C. vibrioides* IbpA (WP_010918744.1, green arrows). *C. vibrioides* has been shown to uptake *myo*-inositol using this ABC-type transporter.

Fig. 7. Identification of *myo*-inositol transporters in *V. coralliilyticus*. Growth analysis of wild type, Δ*iolT1*, Δ*iolX*, and Δ*iolT1*/ Δ*iolX* in minimal media (M9G) with 20 mM *myo*-inositol as a sole carbon source. Growth was measured every hour for 30 h at 28°C. Mean and standard deviation of two biological replicates are shown.

Fig. 8. Phylogenetic distribution of IolG and Iol cluster organization among *Vibrionaceae*. A. IolG evolutionary history was inferred using the Neighbor-Joining method (108) with bootstrap value of 1000 (114). The optimal tree with the sum of branch length = 4.11930205 is shown. The tree is drawn to scale, and evolutionary distances were computed using the JTT matrix-based method (109) with units of the number of amino acid substitutions per site. Shown are 53 amino acid sequences, ambiguous positions were removed for each sequence pair (pairwise deletion option) with a total of 332 positions. Evolutionary analyses were conducted in MEGA X (107). Adjacent to the tree are gene order diagrams representative of the designated species. **B.** Iol cluster organization among *Vibrionaceae* species. Arrows represent open rreading frame and direction of transcription.

Fig. 9. Phylogenetic distribution of IolA and the housekeeping protein RpoB. A. IolA

evolutionary history was inferred using the Neighbor-Joining method (108) with

bootstrap value of 1000 (114). The optimal tree with the sum of branch length =

2.37747408 is shown. The tree is drawn to scale, and evolutionary distances were

computed using the JTT matrix-based method (109) with units of the number of amino

acid substitutions per site. Shown are 43 amino acid sequences, ambiguous positions

were removed for each sequence pair (pairwise deletion option) with a total of 510

positions. Evolutionary analyses were conducted in MEGA X (107). **B**. Housekeeping

protein RpoB phylogeny based on the same set of species that contained IolG

constructed using the Neighbor-Joining method.

Fig. 10. IolG from *Vibrio* clustered with members of *Aeromonas*, *Pectobacterium*, and

Dickeya. Evolutionary analyses were conducted in MEGA X (107). IolG phylogeny was

inferred using the Neighbor-Joining method (108) and the optimal tree with the sum of

branch length = 5.23592824 is shown. The percentage of replicate trees in which the

associated taxa clustered together in the bootstrap test (1000 replicates) are shown next

to the branches and distances were computed using the JTT matrix-based method and

are in the units of the number of amino acid substitutions per site (114). The rate

variation among sites was modeled with a gamma distribution (shape parameter = 5).

This analysis involved 81 amino acid sequences. All ambiguous positions were

removed for each sequence pair (pairwise deletion option). There were a total of 333

positions in the final dataset.

Table 1. Growth curve analysis of *V. coralliilyticus* in minimal media containing glucose

(M9G) and 5% NaCl supplemented with 1mM of GB (glycine betaine), DMSP

(dimethylsulfoniopropionate), Ectoine, Choline, MI (*myo*-inositol), DMG

(dimethylglycine), trimethylamine N-oxide (TMAO), gamma-aminobutyric acid

(GABA), Creatine, or Glycerol. Growth was measured every hour for 24 h at 28°C.

Maximum growth rate, Area under the curve, and final OD presented here for two

biological replicates.

M9G + 5% NaCl	Max. Growth Rate (h⁻¹)	Area Under Curve	Final OD
No CS	0	0	0
GB	0.56	6.1	0.55
DMSP	0.55	6.3	0.67
Ectoine	0.48	3.9	0.69
Choline	0.48	1.9	0.45
***myo*-inositol**	0	0	0
DMG	0.53	4.6	0.52
TMAO	0	0	0
GABA	0.31	0.9	0.23
Creatine	0.38	1.2	0.31
Glycerol	0	0	0

590   **Table 2. Strains and plasmids used in this study**

Strain or plasmid	Genotype or description	Reference(s) or source
Vibrio coralliilyticus		
CN52H-1	Environmental isolate from reef	Ushijima et al., 2020
Δ*iolG*	Deletion of HRJ43_RS06535 from CN52H-1	This study
Δ*iolT1*	Deletion of HRJ43_RS06585 from CN52H-1	This study
Δ*iolX*	Deletion of HRJ43_RS06670 from CN52H-1	This study
Δ*iolT1*/Δ*iolX*	*iolX* deletion in Δ*iolT2* mutant background	This study
Vibrio harveyi		
393	Isolated from barramundi in Australia	Pederson et al., 1998
Escherichia coli		
DH5α *λpir*		ThermoFisher Scientific
B2155 *λpir*	Δ*dapA*::erm pir for bacterial conjugation	(Dehio & Meyer, 1997)
Plasmids		
pDS132	Suicide plasmid; CmR; *sacB*, R6Kg ori	(Philippe et al., 2004)
pDS*iolG*	pDS132 harboring truncated *iolG* allele	This study
pDS*iolT1*	pDS132 harboring truncated *iolT1* allele	This study
pDS*iolX*	pDS132 harboring truncated *iolX* allele	This study

596   **Table 3. Primers used in this study**

Primer Name	Sequence (5′ – 3′)	Length (bp)
Mutant Primers		
AB fwd_*iolG*	accgcatgcgatatcgagctGTGGTTGAGGTGTACACC	549
AB rev_*iolG*	cttatagtgaATTAAACATTCTGTACTCCTTACAG	
CD fwd_*iolG*	aatgtttaatTCACTATAAGAAATTCAACCGGAAG	549
CD rev_*iolG*	gtggaattcccgggagagctTCCGGAAATCGTCCGCTATTTC	
flank fwd_*iolG*	acatcggtcagcgtatgtct	2369
flank rev_*iolG*	aaccgcgactacctttacca	
AB fwd_*iolT1*	accgcatgcgatatcgagctATGCTTTTCATCTCTCGTTC	549
AB rev_*iolT1*	gtacgtgcgcATGACTCGATGACATGTTTC	
CD fwd_*iolT1*	atcgagtcatGCGCACGTACAATAACATTG	549
CD rev_*iolT1*	gtggaattcccgggagagctCGATACCCCTGAAATGAGG	
flank fwd_*iolT1*	actttgttcccagctccgta	2809
flank rev_*iolT1*	gcgaacaaacaatggtgctg	
AB fwd_*iolX*	accgcatgcgatatcgagctCAACTATCGCCGAATAAGGTTATTTG	606
AB rev_*iolX*	aacctttttgTAATGATGGCATCATTGCC	
CD fwd_*iolX*	gccatcattaCAAAAAGGTTAATGGAATAACAGC	465
CD rev_*iolX*	gtggaattcccgggagagctATCGCGATACATCTTATCGTG	
flank fwd_*iolX*	cgatgacccaccgagatact	2248
flank rev_*iolX*	caccgtcagttcagtcatcg	

References

1.   Santos Ede O, Alves N, Jr., Dias GM, Mazotto AM, Vermelho A, Vora GJ, Wilson

B, Beltran VH, Bourne DG, Le Roux F, Thompson FL. 2011. Genomic and

proteomic analyses of the coral pathogen Vibrio coralliilyticus reveal a diverse

virulence repertoire. Isme j 5:1471-83.

2.   Ben-Haim Y, Zicherman-Keren M, Rosenberg E. 2003. Temperature-regulated

bleaching and lysis of the coral Pocillopora damicornis by the novel pathogen

Vibrio coralliilyticus. Appl Environ Microbiol 69:4236-42.

3.   Kooperman N, Ben-Dov E, Kramarsky-Winter E, Barak Z, Kushmaro A. 2007.

Coral mucus-associated bacterial communities from natural and aquarium

environments. FEMS Microbiol Lett 276:106-13.

4.   Kimes NE, Grim CJ, Johnson WR, Hasan NA, Tall BD, Kothary MH, Kiss H,

Munk AC, Tapia R, Green L, Detter C, Bruce DC, Brettin TS, Colwell RR, Morris

PJ. 2012. Temperature regulation of virulence factors in the pathogen Vibrio

coralliilyticus. Isme j 6:835-46.

5.   Godwin S, Bent E, Borneman J, Pereg L. 2012. The role of coral-associated

bacterial communities in Australian Subtropical White Syndrome of Turbinaria

mesenterina. PLoS One 7:e44243.

6.   Ushijima B, Saw JH, Videau P, Häse CC. 2022. Comparison of Vibrio

coralliilyticus virulence in Pacific oyster larvae and corals. Microbiology

(Reading) 168(4).

7.   Ushijima B, Meyer JL, Thompson S, Pitts K, Marusich MF, Tittl J, Weatherup E,

Reu J, Wetzell R, Aeby GS, Häse CC, Paul VJ. 2020. Disease Diagnostics and

Potential Coinfections by Vibrio coralliilyticus During an Ongoing Coral Disease

Outbreak in Florida. Front Microbiol 11:569354.

8.   Vidal-Dupiol J, Ladrière O, Meistertzheim AL, Fouré L, Adjeroud M, Mitta G.

2011. Physiological responses of the scleractinian coral Pocillopora damicornis to

bacterial stress from Vibrio coralliilyticus. J Exp Biol 214:1533-45.

9.   Ben-Haim Y, Rosenberg E. 2002. A novel Vibrio sp. pathogen of the coral

Pocillopora damicornis. Marine Biology 141:47-55.

10.   Rosenberg E, Ben-Haim Y. 2002. Microbial diseases of corals and global

warming. Environ Microbiol 4:318-26.

11.   Ben-Haim Y, Thompson FL, Thompson CC, Cnockaert MC, Hoste B, Swings J,

Rosenberg E. 2003. Vibrio coralliilyticus sp. nov., a temperature-dependent

pathogen of the coral Pocillopora damicornis. Int J Syst Evol Microbiol 53:309-

315.

12.   Bourne DG, Munn CB. 2005. Diversity of bacteria associated with the coral

Pocillopora damicornis from the Great Barrier Reef. Environ Microbiol 7:1162-74.

13.   Sussman M, Willis BL, Victor S, Bourne DG. 2008. Coral pathogens identified for

639   White Syndrome (WS) epizootics in the Indo-Pacific. PLoS One 3:e2393.

640   14.   Munn CB. 2015. The Role of Vibrios in Diseases of Corals. Microbiol Spectr 3(4).

641   15.   Ushijima B, Videau P, Burger AH, Shore-Maggio A, Runyon CM, Sudek M, Aeby

642   GS, Callahan SM. 2014. Vibrio coralliilyticus strain OCN008 is an etiological

643   agent of acute Montipora white syndrome. Appl Environ Microbiol 80:2102-9.

644   16.   Ushijima B, Videau P, Poscablo D, Stengel JW, Beurmann S, Burger AH, Aeby

645   GS, Callahan SM. 2016. Mutation of the toxR or mshA genes from Vibrio

646   coralliilyticus strain OCN014 reduces infection of the coral Acropora cytherea.

647   Environ Microbiol 18:4055-4067.

648   17.   Richards GP, Watson MA, Needleman DS, Church KM, Häse CC. 2015.

649   Mortalities of Eastern and Pacific oyster Larvae caused by the pathogens Vibrio

650   coralliilyticus and Vibrio tubiashii. Appl Environ Microbiol 81:292-7.

651   18.   Ushijima B, Richards GP, Watson MA, Schubiger CB, Häse CC. 2018. Factors

652   affecting infection of corals and larval oysters by Vibrio coralliilyticus. PLoS One

653   13:e0199475.

654   19.   Richards GP, Watson MA, Madison D, Soffer N, Needleman DS, Soroka DS,

655   Uknalis J, Baranzoni GM, Church KM, Polson SW, Elston R, Langdon C,

656   Sulakvelidze A. 2021. Bacteriophages against Vibrio coralliilyticus and Vibrio

tubiashii: Isolation, Characterization, and Remediation of Larval Oyster

Mortalities. Appl Environ Microbiol 87(10):e00008-21.

20. Kim HJ, Giri SS, Kim SG, Kim SW, Kwon J, Lee SB, Park SC. 2020. Isolation and

Characterization of Two Bacteriophages and Their Preventive Effects against

Pathogenic Vibrio coralliilyticus Causing Mortality of Pacific Oyster (Crassostrea

gigas) Larvae. Microorganisms 8(6):926.

21. Kim HJ, Jun JW, Giri SS, Chi C, Yun S, Kim SG, Kim SW, Han SJ, Kwon J, Oh

WT, Lee SB, Kim JH, Park SC. 2020. Identification and Genome Analysis of

Vibrio coralliilyticus Causing Mortality of Pacific Oyster (Crassostrea gigas)

Larvae. Pathogens 9(3):206.

22. Richards GP, Kingham BF, Shevchenko O, Watson MA, Needleman DS. 2018.

Complete Genome Sequence of Vibrio coralliilyticus RE22, a Marine Bacterium

Pathogenic toward Larval Shellfish. Microbiol Resour Announc 7(17):e01332-18.

23. Arboleda M, Reichardt W. 2009. Epizoic communities of prokaryotes on healthy

and diseased scleractinian corals in Lingayen Gulf, Philippines. Microb Ecol

57:117-28.

24. Koenig JE, Bourne DG, Curtis B, Dlutek M, Stokes HW, Doolittle WF, Boucher Y.

2011. Coral-mucus-associated Vibrio integrons in the Great Barrier Reef: genomic

hotspots for environmental adaptation. Isme j 5:962-72.

25.   Kvennefors EC, Sampayo E, Ridgway T, Barnes AC, Hoegh-Guldberg O. 2010.

677   Bacterial communities of two ubiquitous Great Barrier Reef corals reveals both

678   site- and species-specificity of common bacterial associates. PLoS One 5:e10401.

679   26.   Gibbin E, Gavish A, Krueger T, Kramarsky-Winter E, Shapiro O, Guiet R, Jensen

680   L, Vardi A, Meibom A. 2019. Vibrio coralliilyticus infection triggers a

681   behavioural response and perturbs nutritional exchange and tissue integrity in a

682   symbiotic coral. Isme j 13:989-1003.

683   27.   Vega Thurber R, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA,

684   Angly F, Dinsdale E, Kelly L, Rohwer F. 2009. Metagenomic analysis of stressed

685   coral holobionts. Environ Microbiol 11:2148-63.

686   28.   Tout J, Siboni N, Messer LF, Garren M, Stocker R, Webster NS, Ralph PJ,

687   Seymour JR. 2015. Increased seawater temperature increases the abundance and

688   alters the structure of natural Vibrio populations associated with the coral

689   Pocillopora damicornis. Front Microbiol 6:432.

690   29.   Vezzulli L, Previati M, Pruzzo C, Marchese A, Bourne DG, Cerrano C. 2010.

691   Vibrio infections triggering mass mortality events in a warming Mediterranean

692   Sea. Environ Microbiol 12:2007-19.

693   30.   Garren M, Son K, Raina JB, Rusconi R, Menolascina F, Shapiro OH, Tout J,

694   Bourne DG, Seymour JR, Stocker R. 2014. A bacterial pathogen uses

dimethylsulfoniopropionate as a cue to target heat-stressed corals. Isme j 8:999-

1007.

31. Vidal-Dupiol J, Dheilly NM, Rondon R, Grunau C, Cosseau C, Smith KM, Freitag

M, Adjeroud M, Mitta G. 2014. Thermal stress triggers broad Pocillopora

damicornis transcriptomic remodeling, while Vibrio coralliilyticus infection

induces a more targeted immuno-suppression response. PLoS One 9:e107672.

32. Garren M, Son K, Tout J, Seymour JR, Stocker R. 2016. Temperature-induced

behavioral switches in a bacterial coral pathogen. Isme j 10:1363-72.

33. Wu Y, Zhou Z, Wang J, Luo J, Wang L, Zhang Y. 2019. Temperature regulates the

recognition activities of a galectin to pathogen and symbiont in the scleractinian

coral Pocillopora damicornis. Dev Comp Immunol 96:103-110.

34. Zhou Z, Zhao S, Tang J, Liu Z, Wu Y, Wang Y, Lin S. 2019. Altered Immune

Landscape and Disrupted Coral-Symbiodinium Symbiosis in the Scleractinian

Coral Pocillopora damicornis by Vibrio coralliilyticus Challenge. Front Physiol

10:366.

35. Rosenberg E, Kushmaro A, Kramarsky-Winter E, Banin E, Yossi L. 2009. The role

of microorganisms in coral bleaching. Isme j 3:139-46.

36. Roth MS. 2014. The engine of the reef: photobiology of the coral-algal symbiosis.

Front Microbiol 5:422.

37. Kopp C, Domart-Coulon I, Escrig S, Humbel BM, Hignette M, Meibom A. 2015.

Subcellular investigation of photosynthesis-driven carbon assimilation in the

symbiotic reef coral Pocillopora damicornis. mBio 6(1):e02299-14.

38. Sproles AE, Kirk NL, Kitchen SA, Oakley CA, Grossman AR, Weis VM, Davy SK.

2018. Phylogenetic characterization of transporter proteins in the cnidarian-

dinoflagellate symbiosis. Mol Phylogenet Evol 120:307-320.

39. Matthews JL, Oakley CA, Lutz A, Hillyer KE, Roessner U, Grossman AR, Weis

VM, Davy SK. 2018. Partner switching and metabolic flux in a model cnidarian-

dinoflagellate symbiosis. Proc Biol Sci 285(1892):20182336.

40. LaJeunesse TC, Thornhill DJ, Cox EF, Stanton FG, Fitt WK, Schmidt GW. 2004.

High diversity and host specificity observed among symbiotic dinoflagellates in

reef coral communities from Hawaii. Coral Reefs 23:596-603.

41. Tanaka Y, Miyajima T, Koike I, Hayashibara T, Ogawa H. 2006. Translocation

and conservation of organic nitrogen within the coral-zooxanthella symbiotic

system of Acropora pulchra, as demonstrated by dual isotope-labeling

techniques. Journal of Experimental Marine Biology and Ecology 336:110-119.

42. Rädecker N, Pogoreutz C, Voolstra CR, Wiedenmann J, Wild C. 2015. Nitrogen

cycling in corals: the key to understanding holobiont functioning? Trends

Microbiol 23:490-7.

43.   Rosic N, Ling EY, Chan CK, Lee HC, Kaniewska P, Edwards D, Dove S, Hoegh-

734   Guldberg O. 2015. Unfolding the secrets of coral-algal symbiosis. Isme j 9:844-56.

735   44.   Cui G, Liew YJ, Konciute MK, Zhan Y, Hung SH, Thistle J, Gastoldi L, Schmidt-

736   Roach S, Dekker J, Aranda M. 2022. Nutritional control regulates symbiont

737   proliferation and life history in coral-dinoflagellate symbiosis. BMC Biol 20:103.

738   45.   Rands ML, Loughman BC, Douglas AE. 1993. The Symbiotic Interface in an Alga-

739   Invertebrate Symbiosis. Proceedings: Biological Sciences 253:161-165.

740   46.   Muscatine L, Porter JW. 1977. Reef Corals: Mutualistic Symbioses Adapted to

741   Nutrient-Poor Environments. BioScience 27:454-460.

742   47.   Aranda M, Li Y, Liew YJ, Baumgarten S, Simakov O, Wilson MC, Piel J, Ashoor

743   H, Bougouffa S, Bajic VB, Ryu T, Ravasi T, Bayer T, Micklem G, Kim H, Bhak J,

744   LaJeunesse TC, Voolstra CR. 2016. Genomes of coral dinoflagellate symbionts

745   highlight evolutionary adaptations conducive to a symbiotic lifestyle. Sci Rep

746   6:39734.

747   48.   LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR,

748   Santos SR. 2018. Systematic Revision of Symbiodiniaceae Highlights the

749   Antiquity and Diversity of Coral Endosymbionts. Curr Biol 28:2570-2580.e6.

750   49.   Ashley IA, Kitchen SA, Gorman LM, Grossman AR, Oakley CA, Suggett DJ, Weis

751   VM, Rosset SL, Davy SK. 2022. Genomic conservation and putative downstream

functionality of the phosphatidylinositol signalling pathway in the cnidarian-

dinoflagellate symbiosis. Front Microbiol 13:1094255.

50.   Lohia A, Hait NC, Majumder AL. 1999. L-myo-Inositol 1-phosphate synthase

from Entamoeba histolytica. Mol Biochem Parasitol 98:67-79.

51.   Nikitashina V, Stettin D, Pohnert G. 2022. Metabolic adaptation of diatoms to

hypersalinity. Phytochemistry 201:113267.

52.   Lehnert EM, Mouchka ME, Burriesci MS, Gallo ND, Schwarz JA, Pringle JR. 2014.

Extensive differences in gene expression between symbiotic and aposymbiotic

cnidarians. G3 (Bethesda) 4:277-95.

53.   Michell RH. 2008. Inositol derivatives: evolution and functions. Nat Rev Mol Cell

Biol 9:151-61.

54.   Majumder AL, Chatterjee A, Ghosh Dastidar K, Majee M. 2003. Diversification

and evolution of L-myo-inositol 1-phosphate synthase. FEBS Lett 553:3-10.

55.   Turner BL, Papházy MJ, Haygarth PM, McKelvie ID. 2002. Inositol phosphates in

the environment. Philos Trans R Soc Lond B Biol Sci 357:449-69.

56.   Al-Mushhin AAM, Qari SH, Fakhr MA, Alnusairi GSH, Alnusaire TS, AA AL,

Latef A, Ali OM, Khan AA, Soliman MH. 2021. Exogenous Myo-Inositol

Alleviates Salt Stress by Enhancing Antioxidants and Membrane Stability via the

Upregulation of Stress Responsive Genes in Chenopodium quinoa L. Plants

(Basel) 10(11):2416.

57.   Das-Chatterjee A, Goswami L, Maitra S, Dastidar KG, Ray S, Majumder AL.

2006. Introgression of a novel salt-tolerant L-myo-inositol 1-phosphate synthase

from Porteresia coarctata (Roxb.) Tateoka (PcINO1) confers salt tolerance to

evolutionary diverse organisms. FEBS Lett 580:3980-8.

58.   Nelson DE, Koukoumanos M, Bohnert HJ. 1999. Myo-inositol-dependent sodium

uptake in ice plant. Plant Physiol 119:165-72.

59.   Blatt MR, Thiel G, Trentham DR. 1990. Reversible inactivation of K+ channels of

Vicia stomatal guard cells following the photolysis of caged inositol 1,4,5-

trisphosphate. Nature 346:766-9.

60.   Thiel G, MacRobbie EA, Hanke DE. 1990. Raising the intracellular level of

inositol 1,4,5-trisphosphate changes plasma membrane ion transport in

characean algae. Embo j 9:1737-41.

61.   Gillaspy GE. 2011. The cellular language of myo-inositol signaling. New Phytol

192:823-839.

62.   Weber M, Fuchs TM. 2022. Metabolism in the Niche: a Large-Scale Genome-

Based Survey Reveals Inositol Utilization To Be Widespread among Soil,

Commensal, and Pathogenic Bacteria. Microbiol Spectr 10:e0201322.

63.   Felsl A, Brokatzky D, Kröger C, Heermann R, Fuchs TM. 2023. Hierarchic

regulation of a metabolic pathway: H-NS, CRP, and SsrB control myo-inositol

utilization by Salmonella enterica. Microbiol Spectr:e0272423.

64. Yoshida KI, Aoyama D, Ishio I, Shibayama T, Fujita Y. 1997. Organization and

transcription of the myo-inositol operon, iol, of Bacillus subtilis. J Bacteriol

179:4591-8.

65. Yoshida K, Yamamoto Y, Omae K, Yamamoto M, Fujita Y. 2002. Identification of

two myo-inositol transporter genes of Bacillus subtilis. J Bacteriol 184:983-91.

66. Yoshida K, Yamaguchi M, Morinaga T, Kinehara M, Ikeuchi M, Ashida H, Fujita

Y. 2008. myo-Inositol catabolism in Bacillus subtilis. J Biol Chem 283:10415-24.

67. Yoshida KI, Sanbongi A, Murakami A, Suzuki H, Takenaka S, Takami H. 2012.

Three inositol dehydrogenases involved in utilization and interconversion of

inositol stereoisomers in a thermophile, Geobacillus kaustophilus HTA426.

Microbiology (Reading) 158:1942-1952.

68. Yoshida KI, Shibayama T, Aoyama D, Fujita Y. 1999. Interaction of a repressor

and its binding sites for regulation of the Bacillus subtilis iol divergon. J Mol Biol

285:917-29.

69. Berman T, Magasanik B. 1966. The pathway of myo-inositol degradation in

Aerobacter aerogenes. Ring scission. J Biol Chem 241:807-13.

70. Boutte CC, Srinivasan BS, Flannick JA, Novak AF, Martens AT, Batzoglou S,

Viollier PH, Crosson S. 2008. Genetic and computational identification of a

conserved bacterial metabolic module. PLoS Genet 4:e1000310.

71.   Kröger C, Fuchs TM. 2009. Characterization of the myo-inositol utilization island

812   of Salmonella enterica serovar Typhimurium. J Bacteriol 191:545-54.

813   72.   Kröger C, Stolz J, Fuchs TM. 2010. myo-Inositol transport by Salmonella enterica

814   serovar Typhimurium. Microbiology (Reading) 156:128-138.

815   73.   Fry J, Wood M, Poole PS. 2001. Investigation of myo-inositol catabolism in

816   Rhizobium leguminosarum bv. viciae and its effect on nodulation

817   competitiveness. Mol Plant Microbe Interact 14:1016-25.

818   74.   Gauchat-Feiss D, Frey J, Belet M, Deshusses J. 1985. Cloning of genes involved in

819   myo-inositol transport in a Pseudomonas sp. J Bacteriol 162:324-7.

820   75.   Legakis NJ, Papavassiliou JT, Xilinas ME. 1976. Inositol as a selective substrate

821   for the growth of Klebsiellae and Serratiae. Zentralbl Bakteriol Orig A 235:453-8.

822   76.   Primrose SB, Ronson CW. 1980. Polyol metabolism by Rhizobium trifolii. J

823   Bacteriol 141:1109-14.

824   77.   Reber G, Belet M, Deshusses J. 1980. Energetics of myo-inositol transport in

825   Pseudomonas putida. Experientia 36:528-9.

826   78.   Krings E, Krumbach K, Bathe B, Kelle R, Wendisch VF, Sahm H, Eggeling L.

827   2006. Characterization of myo-inositol utilization by Corynebacterium

828   glutamicum: the stimulon, identification of transporters, and influence on L-

829   lysine formation. J Bacteriol 188:8054-61.

79.   Hellinckx J, Fuchs TM. 2017. Hysteresis in myo-inositol utilization by Salmonella

Typhimurium. Microbiologyopen 6(2):e00431.

80.   Müller J, Spriewald S, Stecher B, Stadler E, Fuchs TM. 2019. Evolutionary

Stability of Salmonella Competition with the Gut Microbiota: How the

Environment Fosters Heterogeneity in Exploitative and Interference

Competition. J Mol Biol 431:4732-4748.

81.   Herrou J, Crosson S. 2013. myo-inositol and D-ribose ligand discrimination in an

ABC periplasmic binding protein. J Bacteriol 195:2379-88.

82.   Naughton LM, Blumerman SL, Carlberg M, Boyd EF. 2009. Osmoadaptation

among Vibrio species and unique genomic features and physiological responses

of Vibrio parahaemolyticus. Appl Environ Microbiol 75:2802-10.

83.   Ongagna-Yhombi SY, Boyd EF. 2013. Biosynthesis of the osmoprotectant ectoine,

but not glycine betaine, is critical for survival of osmotically stressed Vibrio

parahaemolyticus cells. Appl Environ Microbiol 79:5038-49.

84.   Whitaker WB, Parent MA, Naughton LM, Richards GP, Blumerman SL, Boyd EF.

2010. Modulation of responses of Vibrio parahaemolyticus O3:K6 to pH and

temperature stresses by growth at different salt concentrations. Appl Environ

Microbiol 76:4720-9.

85.   Pflughoeft KJ, Kierek K, Watnick PI. 2003. Role of ectoine in Vibrio cholerae

osmoadaptation. Appl Environ Microbiol 69:5919-27.

86.   Kapfhammer D, Karatan E, Pflughoeft KJ, Watnick PI. 2005. Role for glycine

betaine transport in Vibrio cholerae osmoadaptation and biofilm formation

within microbial communities. Appl Environ Microbiol 71:3840-7.

87.   Rao NV, Shashidhar R, Bandekar JR. 2013. Comparative analysis of induction of

osmotic-stress-dependent genes in Vibrio vulnificus exposed to hyper- and

hypo-osmotic stress. Can J Microbiol 59:333-8.

88.   Drew D, North RA, Nagarathinam K, Tanabe M. 2021. Structures and General

Transport Mechanisms by the Major Facilitator Superfamily (MFS). Chem Rev

121:5289-5335.

89.   Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the

sensitivity of progressive multiple sequence alignment through sequence

weighting, position-specific gap penalties and weight matrix choice. Nucleic

Acids Res 22:4673-80.

90.   Martins LO, Carreto LS, Da Costa MS, Santos H. 1996. New compatible solutes

related to Di-myo-inositol-phosphate in members of the order Thermotogales. J

Bacteriol 178:5644-51.

91.   Empadinhas N, Mendes V, Simões C, Santos MS, Mingote A, Lamosa P, Santos

H, Costa MS. 2007. Organic solutes in Rubrobacter xylanophilus: the first

example of di-myo-inositol-phosphate in a thermophile. Extremophiles 11:667-

73.

92.   Lamosa P, Rodrigues MV, Gonçalves LG, Carr J, Ventura R, Maycock C, Raven

871   ND, Santos H. 2013. Organic solutes in the deepest phylogenetic branches of the

872   Bacteria: identification of α(1-6)glucosyl-α(1-2)glucosylglycerate in

873   Persephonella marina. Extremophiles 17:137-46.

874   93.   Rodrigues MV, Borges N, Almeida CP, Lamosa P, Santos H. 2009. A unique beta-

875   1,2-mannosyltransferase of Thermotoga maritima that uses di-myo-inositol

876   phosphate as the mannosyl acceptor. J Bacteriol 191:6105-15.

877   94.   Galbraith MP, Feng SF, Borneman J, Triplett EW, de Bruijn FJ, Rossbachl S. 1998.

878   A functional myo-inositol catabolism pathway is essential for rhizopine

879   utilization by Sinorhizobium meliloti. Microbiology (Reading) 144 (Pt 10):2915-

880   2924.

881   95.   Xu T, Rasmussen-Ivey CR, Moen FS, Fernández-Bravo A, Lamy B, Beaz-Hidalgo

882   R, Khan CD, Castro Escarpulli G, Yasin ISM, Figueras MJ, Azzam-Sayuti M,

883   Karim MM, Alam KMM, Le TTT, Thao NHP, Addo S, Duodu S, Ali S, Latif T,

884   Mey S, Somony T, Liles MR. 2023. A Global Survey of Hypervirulent Aeromonas

885   hydrophila (vAh) Identified vAh Strains in the Lower Mekong River Basin and

886   Diverse Opportunistic Pathogens from Farmed Fish and Other Environmental

887   Sources. Microbiol Spectr 11:e0370522.

888   96.   Hossain MJ, Waldbieser GC, Sun D, Capps NK, Hemstreet WB, Carlisle K,

889   Griffin MJ, Khoo L, Goodwin AE, Sonstegard TS, Schroeder S, Hayden K,

Newton JC, Terhune JS, Liles MR. 2013. Implication of lateral genetic transfer in

the emergence of Aeromonas hydrophila isolates of epidemic outbreaks in

channel catfish. PLoS One 8:e80943.

97. Rasmussen-Ivey CR, Hossain MJ, Odom SE, Terhune JS, Hemstreet WG,

Shoemaker CA, Zhang D, Xu DH, Griffin MJ, Liu YJ, Figueras MJ, Santos SR,

Newton JC, Liles MR. 2016. Classification of a Hypervirulent Aeromonas

hydrophila Pathotype Responsible for Epidemic Outbreaks in Warm-Water

Fishes. Front Microbiol 7:1615.

98. Boroujerdi AF, Vizcaino MI, Meyers A, Pollock EC, Huynh SL, Schock TB, Morris

PJ, Bearden DW. 2009. NMR-based microbial metabolomics and the temperature-

dependent coral pathogen Vibrio coralliilyticus. Environ Sci Technol 43:7658-64.

99. Rubio-Portillo E, Gago JF, Martínez-García M, Vezzulli L, Rosselló-Móra R,

Antón J, Ramos-Esplá AA. 2018. Vibrio communities in scleractinian corals differ

according to health status and geographic location in the Mediterranean Sea. Syst

Appl Microbiol 41:131-138.

100. Vezzulli L, Brettar I, Pezzati E, Reid PC, Colwell RR, Höfle MG, Pruzzo C. 2012.

Long-term effects of ocean warming on the prokaryotic community: evidence

from the vibrios. Isme j 6:21-30.

101. Vezzulli L, Colwell RR, Pruzzo C. 2013. Ocean warming and spread of

pathogenic vibrios in the aquatic environment. Microb Ecol 65:817-25.

102.   Dehio C, Meyer M. 1997. Maintenance of broad-host-range incompatibility group

911   P and group Q plasmids and transposition of Tn5 in Bartonella henselae

912   following conjugal plasmid transfer from Escherichia coli. J Bacteriol 179:538-40.

913   103.   Pederson K, Verdonck L, Austin B, Austin D, Blanch AR, Grimont PAD, Jofre J,

914   Koblavi S, Larsen JL, Tianinen T, Vigneulle M, Swings J. 1998. Taxonomic

915   evidence that Vibrio carchariae Grimes et al. 1985 is a junior synonym of Vibrio

916   harveyi (Johnson and Shunk 1936) Baumann et al. 1981. International Journal of

917   Systematic and Evolutionary Microbiology 48:748-749.

918   104.   Gregory GJ, Morreale DP, Carpenter MR, Kalburge SS, Boyd EF. 2019. Quorum

919   Sensing Regulators AphA and OpaR Control Expression of the Operon

920   Responsible for Biosynthesis of the Compatible Solute Ectoine. Appl Environ

921   Microbiol 85(22):e01543-19.

922   105.   Gibson DG. 2011. Enzymatic assembly of overlapping DNA fragments. Methods

923   Enzymol 498:349-61.

924   106.   Philippe N, Alcaraz JP, Coursange E, Geiselmann J, Schneider D. 2004.

925   Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria.

926   Plasmid 51:246-55.

927   107.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

928   Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol

929   35:1547-1549.

930   108.   Saitou N, Nei M. 1987. The neighbor-joining method: a new method for

931   reconstructing phylogenetic trees. Mol Biol Evol 4:406-25.

932   109.   Jones DT, Taylor WR, Thornton JM. 1992. The rapid generation of mutation data

933   matrices from protein sequences. Comput Appl Biosci 8:275-82.

934   110.   Robert X, Gouet P. 2014. Deciphering key features in protein structures with the

935   new ENDscript server. Nucleic Acids Res 42:W320-4.

936   111.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,

937   Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C,

938   Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J,

939   Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M,

940   Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,

941   Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with

942   AlphaFold. Nature 596:583-589.

943   112.   Hastings J, Owen G, Dekker A, Ennis M, Kale N, Muthukrishnan V, Turner S,

944   Swainston N, Mendes P, Steinbeck C. 2016. ChEBI in 2016: Improved services

945   and an expanding collection of metabolites. Nucleic Acids Res 44:D1214-9.

946   113.   Murail S, de Vries SJ, Rey J, Moroy G, Tufféry P. 2021. SeamDock: An Interactive

947   and Collaborative Online Docking Resource to Assist Small Compound

948   Molecular Docking. Front Mol Biosci 8:716466.

949   114.   Felsenstein J. 1985. CONFIDENCE LIMITS ON PHYLOGENIES: AN

950   APPROACH USING THE BOOTSTRAP. Evolution 39:783-791.

Fig. 1

Fig. 2

A. Glycine betaine synthesis and transporter genes

B. Glycine betaine biosynthesis from choline

M9G4% NaCl + 1mM Choline

Fig. 3

Fig. 4

Fig. 5. *V. coralliilyticus* can utilize *myo*-inositol as a sole carbon source. Growth curve analysis in minimal media (M9) containing either 20 mM Glucose (black squares) or 20mM *myo*-inositol (blue triangles). **A.** *V. coralliilyticus* CN52H-1 **B.** *V. harveyi* 393 **C.** *V. coralliilyticus ΔiolG.* Growth was measured every hour for 30 h at 28°C. Mean and standard deviation of two biological replicates are shown.

Fig. 5

A. Predicted *myo*-inositol ligand docking in IolT1 (MFS-type transporter)

B. Predicted *myo*-inositol ligand docking in IolX (ABC-type transporter)

Fig. 6

Fig. 7. Growth pattern analysis of transporter deletion mutants. Growth analysis of wild type, Δ*iolT1*, Δ*iolX*, and Δ*iolT1*/ Δ*iolX*. In minimal media (M9G) with 20 mM *myo*-inositol as a sole carbon source. Growth was measured every hour for 30 h at 28°C. Mean and standard deviation of two biological replicates are shown.

Fig. 7

A. IolG phylogeny

B. myo-inositol metabolism gene clusters

Fig. 8

A. IolA phylogeny

B. Housekeeping RpoB phylogeny

Fig. 9

Fig. 10