Title: Testosterone-Induced Metabolic Changes in Seminal Vesicle Epithelial cells Alter

Plasma Components to Enhance Sperm Fertility

Authors: Takahiro Yamanaka[1], Zimo Xiao[1], Natsumi Tsujita[1], Mahmoud Awad[1,2], Takashi

Umehara[1], Masayuki Shimada[1,3*]

Affiliations:

[1] Graduate School of Integrated Sciences for Life, Hiroshima University; Higashihiroshima,

Hiroshima, Japan

[2] Department of Histology, Faculty of Veterinary Medicine, South Valley University; Qena,

Egypt

[3] Graduate School of Innovation and Practice for Smart Society, Hiroshima University;

Higashihiroshima, Hiroshima, Japan

*Corresponding author. Masayuki Shimada, Ph.D. E-mail: mashimad@hiroshima-u.ac.jp,

Tel/Fax: (+81)-824-24-7899

One Sentence Summary: Testosterone induces ACLY expression in seminal vesicle, a key

factor of forming seminal plasma to acquire *in vivo* fertilization ability of sperm.

Abstract:

Male infertility depends on both sperm and seminal plasma and is induced by aging. In this

study, male infertility was examined with seminal plasma and its synthesis mechanism. The

factors ensuring *in vivo* fertilization potential was secreted from seminal vesicle where the

factors were synthesized in an androgen-dependent manner. Androgen increased glucose uptake

and glycolytic capacity in seminal vesicles, which caused activation of oleic acid synthesis

rather than mitochondrial ATP synthesis. ACLY was identified as a key player in this metabolic

mechanism for producing oleic acid that was incorporated into the sperm and enhanced

fertilization potential *in vivo*. In conclusion, an important role of testosterone-induced metabolic

pathways in the seminal vesicle was to ensure the synthesis of oleic acid, which is essential for

sperm fertilization *in vivo*. These results provide new perspectives for the development of

biochemical markers of semen to assess male fertility and for artificial insemination techniques.

Main Text:

INTRODUCTION

Infertility is defined by the World Health Organization (WHO) as the failure to conceive within 12 months despite contraceptive-free sexual intercourse. Its prevalence has increased over the decades and now accounts for 17.5% of couples worldwide *(1)*. The male partner is thought to be solely or in combination responsible for 50% of infertility cases, and the male factor in the etiology of infertility is highly recognized. The male fertility depends largely on the quality of semen composed of sperm and seminal plasma and decreases with aging.

Spermatogenesis, the process by which mature sperm are generated, takes place partially in the testis, where the characteristic sperm morphology of the head, middle, and tail is formed. Within testis however, sperm are functionally immature and cannot penetrate an oocyte until they undergo a maturation process that occurs during their passage through the epididymis *(2–4)*. Sperm recovered from the epididymis can fertilize a mature oocyte *in vitro* but cannot maintain sustained liner motility required for migration from the uterus to the oviduct *(5, 6)*. Thus, when epididymal sperm are artificially inseminated to the cervix or uterus, fertilization rates are low *(7–9)*. However, when epididymal sperm are mixed with seminal plasma at the time of ejaculation or when sperm are exposed to seminal plasma *in vitro*, even for only a brief period of time, they acquire *in vivo* fertilization potential and are capable of passing through the female reproductive tract and fertilizing ovulated oocyte *in vivo (10)*. Thus, sperm formation occurs in the testis, they mature in the epididymis, but only acquire the *in vivo* fertilization ability after exposure to seminal plasma.

Seminal plasma is mainly produced in the seminal vesicles (approximately 65-75%), with approximately 20-30% produced by the prostate, and small amounts secreted by the bulbourethral and urethral glands *(11)*. Seminal plasma or TGFβ, a component of seminal plasma, increases antigen-specific Treg cells in uterus of mice and humans which induce immune tolerance, resulting in pregnancy *(12, 13)*. Furthermore, SVS2 that is secreted from the seminal vesicle is also known to be required for sperm to escape attack by the female immune system in uterus *(14)*. In addition, seminal plasma is known to contain various biochemical components that support sperm metabolism, including lipids and organic acids. Specifically, Rosecrans et al. *(15)* found significant differences in the levels of most biochemical components

in seminal plasma compared to blood, with specific factors like fructose being rarely detected in

blood but abundant in seminal plasma. Our previous studies *(16–18)* documented that seminal

plasma also contains creatine, and fatty acids, that are taken up by sperm within minutes. These

substances play crucial roles in regulating sperm metabolic pathways and enhancing sperm

motility. Collectively these observations indicate that seminal plasma not only alters the female

immune system to protect sperm but also directly influences sperm metabolic pathways,

facilitating the transformation of sperm capable of *in vivo* fertilization.

Some specific factors that male accessory glands contribute to seminal plasma and impact

male fertility have been elucidated. For example, surgical removal of seminal vesicles in male

mice and rats was associated with infertility *(14, 19, 20)*. The observation that fructose *(21)* and

citric acid *(22)* concentrations in seminal plasma of control mice and rats are higher than in

castrated animals indicates that specific functions of the accessory glands might be affected by

testicular-derived testosterone that is known to activate the intracellular androgen receptor

(AR;NR3C4), required for gene regulation of transcription. AR-deficient mice have been found

to be infertile *(23)*, and long-term administration of androgen receptor antagonists, like

flutamide, abolished male fertility *(24, 25)*. AR-deficient mice and transcriptome analyses have

identified specific targets of AR *(26, 27)*. In both mice and human, androgen levels decrease

with aging, and their concentrations have been reported to cause a decrease in male fertility,

especially ejaculated sperm motility. Therefore, the reduction of androgen concentrations may

affect the metabolic mechanisms of the accessory reproductive organs and alter the composition

of seminal plasma with aging. However, the metabolic mechanisms that impact sperm functions

are governed by a complex of enzymatic reactions and therefore, the intricate interplay between

testosterone and seminal plasma synthesis in the accessory reproductive organs requires further

investigation.

In this study, we first performed various bioassays to identify which accessory glands are

essential for sperm fertility. Subsequently, comparative *in vivo* and *in vitro* analyses were

performed using a hypoandrogenic model. In addition, biochemical and molecular techniques

such as RNA-seq, flux analysis, mass spectrometry, and knockdown experiments were

employed. These analyses reveal the pathways by which seminal vesicles synthesize seminal

plasma, ensure fertilization ability, and provide new therapeutic and preventive strategies for

male infertility.

Seminal vesicle and prostate-derived fluids induce functional sperm motility

A bioassay using epididymal sperm that had not been exposed to seminal plasma was done to

clarify the effect of seminal plasma on sperm functions. For this bioassay, fluids were prepared

from seminal vesicle secretion and homogenized prostate tissues, and then the fluid was mixed

with the basic medium, Human Tubal Fluid (HTF) medium in various proportions (Fig.1A-D).

The addition of seminal vesicle secretion alone to the basic medium (HTF medium)

significantly increased the rate of sperm linear motility (LIN) compared to the HTF medium

alone (Fig.1D). The straight line velocity (VSL) of those sperm was also significantly increased

(Fig.1C). This significant increase in straightening of sperm by the seminal vesicle-derived

secretion was further significantly positively affected by the addition of prostate extract 1:9 or

2:8 to the seminal vesicle secretion (Fig.1C,D). However, no significant improvement in

straightness was observed with the mixture of prostate extract alone (Fig.1C,D). These results

clearly indicate that sperm exposure to seminal plasma factors significantly improves sperm

linear motility and that this positive effect is mainly due to the seminal vesicle secretory factors.

The functions of male accessory reproductive organs are regulated by androgens. Therefore,

to elucidate whether the seminal vesicle factors, which enhance sperm linear motility, are

synthesized in an androgen-dependent manner, bioassays were performed using seminal vesicle

secretions collected from control mice or mice treated for 7 consecutive days with flutamide, an

androgen receptor antagonist, and sperm collected from the epididymis of mice not treated with

flutamide (Fig.1E). The results showed that LIN and VSL were significantly reduced in the

seminal vesicle secretions from flutamide-treated mice (Flutamide) compared to the control

mice (Ctrl) (Fig.1F,G). The reduction of sperm linier motility speed (VSL) was also

significantly lower in the sperm treated with seminal vesicle secretions from over 12 months old

mice that showed low circular testosterone level as compared with the sperm treated with

seminal vesicle secretions from young adult mice (3 months old mice) (Supplemental Fig.1A,B).

Furthermore, JC-1 staining to measure mitochondrial membrane potential, which is important

for sperm linear motility, showed significantly lower mitochondrial membrane potential activity

in sperm treated with the secretions of seminal vesicles from flutamide-treated mice (Fig.1H).

These results indicated that androgen-dependent changes in seminal vesicle functions alter the

levels/activity of seminal vesicle secretion factors, which regulate sperm linear motility. The

changes of seminal vesicle functions were also induced with aging.

Testosterone-androgen receptor activity suppresses the proliferation of seminal vesicle

epithelial cells

To understand the difference between seminal vesicles that secrete factors for enhancing sperm

linear motility and those that are impaired in this function, we performed histological analysis of

seminal vesicles recovered from mice treated and non-treated with flutamide. In control mice, the

seminal vesicle epithelial cells were in a single layer, and androgen receptors were localized in the

nuclei of seminal vesicle epithelial cells (Fig.2A). However, in mice treated with continuous

administration of flutamide, seminal vesicle epithelial cells were multilayered, and the strong

accumulation of androgen receptors in nucleus was not observed (Fig.2A). To investigate the

cause of this multilayering of cells, we performed Ki67 staining, a marker of cell proliferation, and

apoptosis detection by TUNEL staining (Fig.2B). The results showed that the percentage of

Ki67-positive cells increased significantly after treatment with flutamide, but the percentage of

TUNEL-positive cells did not change, suggesting that the testosterone-androgen receptor pathway

is responsible for the inhibition of proliferation of seminal vesicle epithelial cells (Fig.2C). These

changes of abnormal morphology and high proliferation activity were also observed in seminal

vesicle of more than 12 months old mice that showed the low level of testosterone in serum

(Supplemental Figure 1C-E). Therefore, seminal vesicle epithelial cells were collected and

cultured in primary culture as shown in Fig. 2D, and the changes in cell number were counted and

cell cycle was analyzed by flow cytometry. When cultured without testosterone (c), the number of

seminal vesicle epithelial cells increased in a day-dependent manner. However, the addition of

testosterone significantly decreased the number of seminal vesicle epithelial cells during the 8-day

culture period in a concentration-dependent manner (Fig.2E). Flow cytometric analysis of the cell

cycle on day 7 showed that the G0/G1 phase was significantly increased and the G2/M1 phase

was significantly decreased by the addition of testosterone (Fig. 2F,G). These results indicate that

seminal vesicle epithelial cells are inhibited in cell division by the testosterone-androgen receptor

pathway.

Testosterone alters metabolic gene expression in seminal vesicle epithelial cells

To better understand the roles of testosterone in suppressing proliferation of seminal vesicle

epithelial cells, RNA sequencing was done using the epithelial cells cultured with or without

testosterone. DEG (Differential Expressed Genes) analysis revealed that a total of 23,459 genes

were identified, including 997 upregulated genes and 3463 downregulated genes in seminal

vesicle epithelial cells by testosterone relative to control (Fig.3A). Genes largely upregulated

by testosterone included those already reported in seminal vesicles *(14, 19, 28)*, such as the

Seminal Vesicle Secretion (*Svs*) family (*Svs1~6*), *Pate4*, *Spinkl*, *Ceacam10*, and *Sva* (Fig.3B,C).

Therefore, the primary culture of seminal vesicle epithelial cells used in this experiment was

considered to maintain the characteristics of seminal vesicle epithelial cells *in vivo*. Many

genes were downregulated by testosterone, and those with a high rate of variation included

collagen synthase and *Mmp2*, which are involved in extracellular matrix formation (Fig.3C).

However, KEGG analysis, rather than individual genes, was used to predict functions altered

by testosterone, and it was found that the expression of genes involved in the metabolic

pathway was the most highly affected by testosterone (Fig.3D). Because there is a close

relationship between cell proliferation and metabolic activity, and between metabolic

substrates in seminal plasma and sperm motility, we hypothesized that testosterone induced

changes in seminal vesicle epithelial cell metabolism alter the components of seminal plasma

to mediate the enhancement of sperm motility.

Testosterone sifts metabolic activity of seminal vesicle epithelial cells

In the presence of testosterone, the addition of glucose significantly activated glycolysis as

compared with that in epithelial cells cultured without testosterone (Fig.4A,B). However, there

was no difference in the glycolytic capacity, indicating that testosterone may increase glucose

metabolism but does not to enhance the expression of a group of enzymes involved in the

glycolytic pathway. To determine the metabolic fate of glycolytic end products, seminal vesicle

epithelial cells were cultured with and without testosterone and then concentrations of pyruvate

and lactate were measured. The concentration of pyruvate was significantly decreased by the

addition of testosterone; however, the concentration of lactate was not significantly changed

(Fig.4D,E). Therefore, we analyzed mitochondrial metabolism using a flux analyzer, predicting that

more glucose is metabolized, pyruvate is synthesized in response to testosterone, and is further

metabolized in the mitochondria. However, measuring mitochondrial respiratory metabolism with

a Flux analyzer, the basal respiratory rate (the activity of the electron transport chain) was

significantly decreased by testosterone (Fig.4F-I). ATP production was also significantly

suppressed exposure to testosterone. Furthermore, the maximum respiration and spare

respiration capacity were also significantly decreased in response to testosterone treatment,

suggesting that testosterone does not educe transient enzyme activity in mitochondria, but to

rather weakens the metabolic pathway of the mitochondrial TCA cycle and/or the electron

transport chain due to the changes in gene expression patterns in seminal vesicle epithelial cells.

Testosterone causes changes in the expression of genes encoding enzymes involved in

glucose catabolism and anabolism

The expression of genes involved in glucose catabolism and anabolism in the *in vitro* culture

system of seminal vesicle epithelial cells was analyzed by qPCR (Fig.5). The gene expression of

glucose transporter (*Glut*) and enzymes involved in the glycolytic pathway were not

significantly affected by testosterone, consistent with the results of metabolic flux analysis,

which showed that the glycolytic pathway was not enhanced despite the activation of the

glycolysis. Furthermore, the expression of genes encoding enzymes involved in the TCA cycle

in mitochondria was also not significantly affected by testosterone, but the expression of

enzymes involved in the electron transport chain was significantly decreased. Interestingly, the

expression of *Acly*, that encodes an enzyme that reconverts citric acid formed in the TCA cycle

to Acetyl CoA and induces cholesterol synthesis and fatty acid synthesis, was significantly

increased by testosterone. Although *Hmgcr*, which encodes the rate-limiting enzyme involved in

cholesterol synthesis, was unchanged, the expression of *Acc*, which encodes the rate-limiting

enzyme for fatty acid synthesis, was significantly upregulated by testosterone. These

characteristic testosterone-induced changes in gene expression observed in cultured cells were

also detected by ACLY immunostaining of *in vivo* seminal vesicle samples (Supplemental

Figure 2). Specifically, the expression of genes encoding a group of enzymes of the electron

transport chain was significantly upregulated by continuous flutamide administration *in vivo*,

whereas the gene expression of *Acly* and *Acc* was significantly decreased by flutamide

administration. Immunostaining results showed that ACLY was detected specifically in seminal

vesicle epithelial cells, and their staining appeared to be reduced by flutamide treatment. The

lower level of ACLY was also observed in seminal vesicle epithelial cells of over 12 months old

mice (Supplemental Figure 2B).

Mechanism of testosterone-induced enhancement of glucose uptake and thereby fatty acid

synthesis

The effect of testosterone on glucose uptake in seminal vesicle epithelial cells was analyzed

using fluorescence-labeled glucose. As expected from the results of flux metabolic analyses,

testosterone treatment significantly increased glucose uptake (Supplemental Figure 3A).

Because the expression of each *Glut* family gene encoding glucose transporters was not altered

by testosterone (Fig.5), we focused on GLUT4, whose cellular localization is regulated, and

analyzed the effects of Indinavir, a functional inhibitor of GLUT4 *(29, 30)*. Indinavir

significantly suppressed glucose uptake and significantly decreased fatty acid synthesis (the

amount of fatty acids accumulated in the cell), which was increased by testosterone

(Supplemental Figure 3A,B). Furthermore, immunostaining analysis of the localization of

GLUT4 in seminal vesicles revealed that GLUT4 was localized outside the nucleus of seminal

vesicle epithelial cells (Supplemental Figure 4), suggesting that GLUT4 was partially localized

at the plasma membrane and contributed to glucose uptake, whereas in seminal vesicles from

mice treated with flutamide continuously, no strong signal was detected at the presumed

location of GLUT4 near the plasma membrane of seminal vesicle epithelial cells

(Supplemental Figure 4). The same characteristics were observed in aged mice.

Qualitative and quantitative analysis of the fatty acids synthesis was performed by GC-MS.

In seminal vesicles, C16:0, C18:0 and C18:1 were detected, in which C18:1 was decreased to

an undetectible level in seminal vesicles of mice treated with the flutamide (Supplemental

Figure 5A). In the cultured seminal vesicle epithelial cells, C18:1, oleic acid, was significantly

increased by the addition of testosterone as observed in seminal vesicle (Supplemental Figure

3C). The expression of *Elovl6* encoding ELOVL6, which desaturates C18:0 saturated fatty

acids to C18:1, was significantly up-regulated by testosterone (Supplemental Figure 3D). The

testosterone-induced change was also detected in *in vivo* samples (seminal vesicle), where the

expression of *Evovl6* was decreased by the flutamide treatment (Supplemental Figure 5B).

Oleic acid (18:1) is taken up by sperm and promotes their linear movement

We observed that seminal vesicles secrete testosterone-dependent factors that induce sperm

to move straight and that seminal vesicle epithelial cells synthesize and secrete oleic acid in a

testosterone-dependent manner. In order to examine whether oleic acid is a factor that induces

sperm linear motility, we examined whether or not oleic acid is taken up by sperm and then

changes of sperm motility pattern. When epididymal sperm was incubated with fluorescently

labeled oleic acid, the fluorescent signal was selectively detected in the midpiece of the sperm

after quenching of extracellular fluorescence (Fig.6A). The fluorescence intensity of sperm

detected by flow cytometry significantly increased in a dose-dependent manner (Fig.6B, C). In

oleic acid-treated sperm, VSL was significantly increased in a dose-dependent manner in the

range of 0 to 50 nM (Fig.6D). LIN values were highest when 10 nM oleic acid was added

(Fig.6E). To quantify metabolic changes in mitochondrial activity, a flux analyzer was used to

monitor real-time changes in mitochondrial β-oxidation. A significant increase in OCR was

induced by the injection of 10 nM oleic acid. Etomoxir, a β-oxidation inhibitor, significantly

decreased the OCR that was induced by oleic acid (Fig.6F, Supplemental Figure 6). In situ ATP

detection by BioTracker ATP-Red revealed that the percentage of sperm with high ATP

production was significantly increased under the presence of oleic acid (Fig.6G).

To examine whether the addition of oleic acid to low quality of seminal plasma (collected

from flutamide-treated mice) restores *in vivo* the fertilization potential of epididymal sperm,

sperm treated with seminal plasma collected from control or flutamide-treated mice was used

for artificial insemination. The *in vivo* fertilization rate was about 70% when sperm was treated

with the seminal vesicle secretions from control mice (SV); however, the fertilization rate was

significantly lower in the group that sperm was treated with the seminal vesicle secretions from

flutamide-treated mice (Flu) (SV: 65.5 ± 3.8%, Flu: 39.4 ± 7.8%). Although not significant,

adding oleic acid to the seminal secretions of flutamide-treated mice (Flu+OA) increased the

fertilization rate to 53.1 ± 1.3% (Fig.6H).

ACLY expression is upregulated by testosterone and is essential for the metabolic shift of

seminal vesicle epithelial cells that mediates sperm linear motility

ACLY is up-regulated in seminal epithelial cells as a factor that enables the utilization of

citric acid from the TCA cycle for fatty acid synthesis outside the mitochondria. Therefore, we

performed siRNA knockdown experiments of ACLY to clarify the impact of ACLY on the

cellular metabolic shift. The addition of testosterone significantly increased the amount of

ACLY protein in seminal epithelial cells (Fig.7A,B). Viral infection with siRNAs encoding

Acly-targeted siRNAs markedly decreased the amount of ACLY (Fig.7C). This knockdown of

ACLY not only increased the oxygen consumption of seminal epithelial cells in the control and

testosterone-treated group, but also significantly increased the ATP production related oxygen

consumption (Fig.7D,E,F). Thus, ACLY expression is associated with mitochondrial metabolic

pathway activity in seminal epithelial cells. Furthermore, the knockdown of ACLY also

significantly suppressed testosterone-induced fatty acid synthesis in seminal epithelial cells

(Fig.7G). Supernatants from epithelial cells of the testosterone-added group induced sperm to

move linearly (Fig.7H). However, the knockdown of ACLY significantly reduced the effects of

the culture supernatant on sperm linear motility (Fig.7H).

This characteristic metabolic mechanism induced by testosterone is also observed in

human seminal epithelial cells

To determine whether these findings in mice could be applied to humans, the commercially

available human seminal vesicle epithelial cells were used for following studies (Fig.8A). The

growth curves showed that cell proliferation was inhibited by 100 ng/mL of testosterone

compared to the control; a pattern similar to that observed in murine seminal vesicle epithelial

cells (Fig.8B). The characteristics of metabolic activity of human seminal epithelial cells were

also similar to those of mice, with testosterone significantly enhancing the glycolytic pathway

(Fig.8C), but significantly reducing the capacity for oxygen consumption in mitochondria

(Fig.8D). Moreover, fluorescent-glucose uptake was increased by testosterone and decreased

by the inhibitor of GLUT4 function (Fig.8E). Additionally, while testosterone also significantly

increased fatty acid synthesis, this increase was completely abolished by the GLUT4 inhibitor

(Fig.8F).

DISCUSSION

Seminal plasma has a variety of functions, including sperm transport, nutritional support,

interactions with the female reproductive tract, and especially the regulation of its relationship

with the female immune system. Mammalian seminal plasma is secreted mainly from the

seminal vesicle and is known to be rich in cytokines, prostaglandins, and SVS family members

(31). For this reason, recent studies have focused on the immunomodulatory function of

seminal plasma in the female reproductive tract. On the other hand, fructose and lipids in

seminal plasma are the major energy sources for sperm *(32)*. However, although metabolomic

analyses of seminal plasma have shown that there are metabolic substrates present at higher

concentrations in seminal plasma than in blood *(15, 21, 22)*, how these substrates are

synthesized and specifically how they alter sperm motility has not been determined. Moreover,

though the motility of ejaculated spermatozoa is known to decrease with aging, few reports

have revealed age-related changes in the seminal vesicle glands, which are responsible for

seminal plasma synthesis.

In this study, we focused on the reproductive organs that synthesize seminal vesicle

secretion as target organs for testosterone, and found that factors formed by

testosterone-dependent metabolic changes in seminal vesicles activate sperm mitochondria and

enhance their linear motility. Testosterone is known to activate intracellular ARs and alter

intracellular metabolic pathways in skeletal muscle *(33)*, prostate cells *(34)* and

cardiomyocytes *(30)*. ARs are known to directly regulate the expression of glycolytic and lipid

metabolic genes *(30, 35, 36)*. In the present study, the nuclear localization of AR in seminal

vesicle epithelial cells (i.e., functioning as a transcription factor) and the activation of the

glycolytic system by testosterone were similar to those in the above cell types. Interestingly,

however, in the seminal vesicle epithelial cells, expression of *Glut* family members was

unchanged but GLUT4 localization was altered in a testosterone-dependent manner to promote

glucose uptake. Such gene expression-independent effects of testosterone on glucose

metabolism have also been reported in skeletal muscle, liver, and adipose tissue, where

testosterone leads to the phosphorylation of GLUT4 and enhances its translocation to the

plasma membrane *(37, 38)*. This is consistent with the results of our metabolic flux analyses,

that show testosterone increases the rate of glycolysis in seminal vesicle epithelial cells by

increasing glucose uptake rather than by increasing glycolytic capacity. We further confirmed

that GLUT4 is the transporter that mediates this glucose uptake using indinavir, a

GLUT4-specific inhibitor.

In seminal epithelial cells, ATP production was low and cell proliferation was suppressed

despite testosterone-induced activation of the glycolytic system. We hypothesized that this

could be explained by the testosterone-dependent regulation of genes involved in

mitochondrial metabolism, including the up-regulation of ACLY, a strategic enzyme that links

the glycolytic system with lipid metabolism. Manipulation of ACLY expression in mouse

myoblasts and embryonic stem cells has been shown to alter the TCA cycle into a "non-normal

cycle" *(39)*. One advantage of the non-normal TCA cycle is its ability to retain carbon instead

of burning it and to regenerate the cytosolic NAD+ needed to maintain glycolysis *(39, 40)*.

This suggests that ACLY activated by testosterone may inhibit proliferation of seminal

epithelial cells by skipping some steps of the mitochondrial TCA cycle and minimizing ATP

production.

Testosterone has been reported to cause hypertrophy of cardiac myocytes *(41, 42)*. Cell

hypertrophy is partly due to the accumulation of fatty acids in the cytoplasm, which causes

significant damage to cellular functions *(43, 44)*. On the other hand, the normal cell

hypertrophy, especially in myofibers, involves the activation of signaling molecules such as

Myc, hypoxia-inducible factor, and Pi3k-Akt-mTor, which regulate metabolic reprogramming

in cancer by increasing glucose uptake in the presence of oxygen *(45–47)*. In the present study,

induction of ACLY expression by testosterone not only decreased cycling in the TCA cycle, but

also promoted glucose assimilation, which played an important role in fatty acid synthesis.

Thus, in tissues where fatty acid synthesis is promoted, either β-oxidation-dependent ATP

production in mitochondria or the secretion of fatty acids occurs to maintain cellular function

(48–50); providing evidence to explain why the presence of fatty acids in the cell has no

adverse effect. In accessory reproductive glands, such as the prostate and seminal vesicles, the

glycolytic system is activated in a testosterone-dependent, but ATP production is not enhanced
and cell proliferation is not induced and the metabolites are used as a substrate for sperm
motility. This may be because in the prostate, pyruvate synthesized by the glycolytic system in
response to testosterone is converted to acetyl CoA, which is then converted to citric acid in
the TCA cycle, citric acid is released as the main component of seminal plasma *(34)*.

The testosterone-dependent growth of cancerous prostate epithelial cells may be due to
genetic mutations, such as elevated expression of the gene encoding aconitase, which converts
citric acid to isocitrate, which blocks the TCA cycle and inhibits citric acid secretion *(51)*. On
the other hand, in the analysis of seminal epithelial cells, testosterone treatment induced the
expression of *Acc*, which encodes the rate-limiting enzyme for fatty acid synthesis, and *Elovl6*,
which desaturates to oleic acid, significantly increasing total saturated fatty acid and oleic acid
content. Therefore, testosterone promotes glucose conversion into fatty acids, especially oleic
acid, in seminal epithelial cells. Although there is a commonality in the activation of the
glycolytic system testosterone target tissues, the different expression patterns of genes involved
in metabolism in mitochondria may exert their specific functions in each tissue. In addition to
genes involved in fatty acid synthesis such as *Fasn* and *Elovl6*, *Acly* is also encoded in a region
in close proximity in human chromosome 17 and mouse chromosome 11, suggesting that in
seminal vesicle epithelial cells, this region is targeted to promote gene expression. The
co-regulation of these genes in seminal vesicle epithelial cells leads to the synthesis of seminal
plasma components that ensure sperm motility.

Decreased testosterone synthesis in the testes is a phenomenon observed with aging, and
associated with decreased spermatogenesis*(52)*. However, the probability of obtaining sperm
without ICSI or IVF is extremely low, despite the presence of sufficient numbers of sperm with
normal-morphology *(53–55)*. This indicates that testosterone plays a major role not only in
spermatogenesis but also in functional maturation of sperm. In this study, metabolic
abnormalities induced by inhibition of androgen receptor function caused abnormal
proliferation of seminal vesicle epithelial cells in aged mice. In other words, the findings of the
seminal vesicle epithelial cell culture experiments and flutamide treatment may mimic the
changes that occur with aging. Oleic acid synthesis by testosterone-dependent metabolic
pathways was regulated in seminal vesicle epithelial cells and the oleic acid stimulated sperm

motility and enhanced *in vivo* fertilization success. Thus, results of these studies document that

the quality and presence of key metabolic products in seminal plasma influence male fertility.

Specifically, the results open the possibility of translational studies to understand human male

infertility, especially in cases of failure to conceive by natural mating or artificial insemination.

Because seminal plasma contains components specific to certain male reproductive organs,

differences in protein composition may reflect pathological processes in specific organs *(56)*.

However, although proteomic and metabolomic analyses of seminal plasma have been

performed *(28, 56)*, the functional changes in seminal plasma, especially the effects on *in vivo*

fertilization potential, have not been well understood. Differences in fatty acid concentrations,

including oleic acid, in seminal plasma may provide a predictive marker for the fertilization

potential of human and domestic animal semen. Furthermore, the addition of oleic acid to

low-quality semen improved fertilization rates associated with artificial insemination,

providing a strategy for improving artificial insemination. In our previous study, adding fatty

acids to the thawing medium of frozen bull sperm improved their linear motility and viability

(18). The similarities between mouse and human seminal vesicle epithelial cells demonstrated

in this study have strong potential for clinical application of these observations.

Overall, the results of the studies described herein strongly support the hypothesis that

testosterone-dependent glucose to oleic acid conversion processes are required for normal

seminal plasma functions in the seminal vesicle and normal reproductive performance in males.

This study advances our understanding of the role of testosterone in glucose metabolism and

fatty acid synthesis and lays the groundwork for future research and potential therapeutic

interventions in reproductive biology and fertility.

MATERIALS AND METHODS

Study design

The objective of this study was to elucidate the mechanisms of sperm plasma synthesis that

enhance sperm fertility. Testosterone-dependent changes in the seminal plasma function were

examined by comparative analysis of adult mice and flutamide-treated mice. The bioassays of

extracts from male accessory glands by epididymal sperm motility analysis were used to

examine the seminal plasma components important for sperm fertilization. We also isolated the

mouse seminal vesicle epithelial cultured cells, and investigated the synthesis mechanism of the

seminal plasma components using RNA-seq, FluxAnalyzer, and shRNA knockdown. Finally,

we performed validation using the human epithelial cells.

Chemicals and Animals

Unless otherwise noted, chemicals used in this study were purchased from Sigma-Aldrich (St.

Louis, MO, USA) or Nacalai tesque (Osaka, Japan). Pregnant mare serum gonadotropin

(PMSG) and human chorionic gonadotropin (hCG) were purchased from Asuka Seiyaku (Tokyo,

Japan). All procedures during animal experiments were reviewed and approved by the Animal

Care and Use Committee of Hiroshima University (Hiroshima, Japan; C21-9-3) and conducted

according to regulations. Specific pathogen-free C57BL/6NCrl male mice or Crl: CD1 (ICR)

female mice were purchased from Jackson Laboratory Japan (Kanagawa, Japan) and housed in

an environmentally controlled room with a 12-hour light/dark cycle, a temperature of $23 \pm 3°C$,

and free access to laboratory food (MF; Oriental Yeast Co., Ltd., Tokyo, Japan) and tap water. 8-

to 16-week-old male or female mice were used for the experiments. Flutamide-treated mice

were injected subcutaneously for 7 days with 50 μg/g (body weight) of flutamide dissolved in

corn oil. Vehicle-treated mice received 100 μL of corn oil subcutaneously for 7 days. The aged

males were 12 to 16 months old.

Bioassay using pseudo-seminal plasma and epididymal sperm

Prostate (mixed anterior, dorsal, lateral, and ventral) and seminal vesicles were collected from

adult C57BL/6NCrl mice. Seminal vesicle secretions were obtained by pressing the seminal

vesicles immediately after euthanasia. Prostatic extracts were obtained by homogenizing the

prostate immediately after euthanasia. Seminal vesicle secretions and prostate extracts collected

from one male mouse were dissolved in 500 μL and 200 μL of mHTF medium without BSA

(57), respectively, then centrifuged at $2000 \times g$ for 5 min at room temperature, and the

supernatant was collected, and protein levels were determined. The seminal vesicle secretions

from flutamide-treated mice were also collected similarly.

Cauda epididymis sperm recovered from male mouse were dispersed in 500 μL of mHTF

without BSA drops and centrifuged at $300 \times g$, 3 min to wash out epididymis-derived factors.

The epididymis sperm was mixed with seminal vesicle secretion and/or prostate extract as

shown in Fig.1A. In flutamide administration experiments, the final protein concentration of

seminal vesicle secretion was adjusted to 10 mg/mL. The samples were incubated at 37°C for

60 min in a humidified atmosphere of 5% CO_2 in air. Sperm motility patterns were then

examined using computer-assisted sperm analysis (CASA; HT CASA-Ceros II, Hamilton

Thorne, Beverly, MA, USA).

Serum testosterone levels

The serum was separated from the clot by centrifugation at $1,500 \times g$ for 15 min.

Testosterone levels in the serum samples were determined by a rodent testosterone ELISA test

kit (ERKR7016, Endocrine Technologies, Inc., Newark, CA, USA) according to the

manufacturer's manual.

Mitochondrial activity

Mitochondrial activity of sperm was measured using a MitoPT® JC-1 Assay Kit (911,

Immuno Chemistry Technologies, LLC, Bloomington, MN, USA) according to our previous

study *(17)*. Briefly, sperm were incubated with 200 µL of working solution containing

5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolyl carbocyanine iodide (JC-1) dye at 37 °C

for 30 min in the dark. The sperm suspension was centrifuged and washed twice with the base

medium. After washing, the sperm pellet was resuspended in the base medium and analyzed

with an Attune® NxT Acoustic Focusing Cytometer (Thermo Fisher Scientific Inc., Waltham,

MA, USA) using a 488 nm laser and a filter with a bandwidth of 574/26 nm. The intensity of

the average value was analyzed as the mean fluorescence intensity (MFI) of JC-1 orange

aggregates. A total of 100,000 sperm events were analyzed. The sperm population is shown in

Supplementary Figure 8.

Immunohistochemistry and immunofluorescence

Seminal vesicles were fixed overnight in 4% (w/v) paraformaldehyde/PBS, dehydrated in

70% (v/v) ethanol, and embedded in paraffin. Paraffin-embedded fixed sections (4 µm) were

deparaffinized and hydrated with xylene and ethanol, respectively. Some slides were stained

with hematoxylin and eosin (HE). For antigen retrieval, sections were placed in citrate buffer

(pH 6.0), microwaved until boiling and allowed to stand at the temperature of pre-boil (95°C)

for 15 minutes. Sections were incubated with 3% (v/v) hydrogen peroxide in methanol to block

endogenous peroxidase and then incubated with Normal Goat Serum Blocking Solution for

rabbit antibody (S-1000, Vector Laboratories, Newark, CA, USA) to block nonspecific sites.

The sections were then incubated overnight with primary rabbit antibodies: anti-AR antibody

(1:100; ab133273; Abcam, Cambridge, UK), anti-Ki67 antibody (1:1000; ab15580; Abcam) or

anti-ACLY antibody (1:200; ab40793; Abcam). Positive signals were visualized by DAB

(3,3′-Diaminobenzidine, 040-27001, Fujifilm Wako, Osaka, Japan) using the VECTASTAINE

LifeABC kit (PK-6101, Vector Laboratories) according to the manufacturer's protocol.

Apoptotic cells (including fragmented DNA) were detected using the In Situ Cell Death

Detection Kit POD (11684817910, Roche Ltd., Basel, Switzerland) according to manufacturer's

protocol. Nuclei were visualized with hematoxylin (Sakura Finetek, Tokyo, Japan). Ki-67 and

apoptotic positive cells were quantified using ImageJ software.

For immunofluorescence, after blocking nonspecific sites, the sections were incubated with a

primary antibody: anti-GLUT4 antibody (1:200; ab33780; Abcam). After washing with 0.3%

(v/v) Triton X-100 in PBS (-), Cy3-labeled goat anti-rabbit IgG (1:200, C-2306, Sigma) and

DAPI (VECTESHIELD Mounting Medium with DAPI, H-1200, Vector Laboratories) were

used to visualize the antigen and nuclei. Digital images were taken using an APX100 Digital

Imaging System (EVIDENT, Tokyo, Japan).

Isolation and culture of seminal vesicle epithelial cells

Seminal vesicles were removed from euthanized mice, cut open vertically to expose the

seminal vesicle cavity, and incubated with Hanks' balanced salt solution (HBSS; $CaCl_2$ • $2H_2O$

0.185 g/L, $MgSO_4$ 0.098 g/L, KCl 0.400 g/L, KH_2PO_4 0.060 g/L, NaHCO3 0.350 g/L, NaCl

8.000 g/L, Na_2HPO_4 0.048 g/L, D-Glucose 1.0 g/L, dissolved in ultrapure water). Tissues were

digested with 10 mL of HBSS containing 25 mg/mL pancreatin and 0.25% (v/v) trypsin EDTA

for 1 hour at 4°C, 45 minutes at room temperature, and 15 minutes at 37°C, and epithelial cells

were obtained. These were passed through a 70 μm nylon filter (352350, Corning Inc., Corning,

NY, USA) and seeded into collagen-coated 12-well plates (4815-010, IWAKI, Shizuoka, Japan).

Cells were cultured in DMEM/F-12, 10% fetal bovine serum (FBS; 10438-018, Thermo Fisher

Scientific Inc.), 100 U/mL penicillin, 100 μg/mL streptomycin. The culture condition was 37°C,

5% CO_2, 95% air, and humidity.

Human seminal vesicle epithelial cells (4460, ScienCell Research Laboratories, Carlsbad, CA,

USA) were purchased, seeded in 2 μg/cm^2 poly-L-lysine-coated plates according to

manufacturer's protocol, and cultured in epithelial cell medium (4101, ScienCell Research

Laboratories). The cells were seeded at 10000 cells/cm^2 on collagen-coated plates the day

before 100 ng/mL testosterone treatment. Testosterone was dissolved in ethanol, and the final

concentration of ethanol was adjusted to 0.1% (v/v) when used for cultured cells. The culture

medium was replaced every 48 hours.

Cell proliferation curve

The seminal vesicle epithelial cells were seeded at 10000 cells/cm^2 and then were cultured in

various concentrations of testosterone. The cells were washed with HBSS (without

$CaCl_2$-2H_2O/$MgSO_4$) at the time of cell counting and collected using 0.25% (v/v) trypsin. The

recovered cells were suspended in DMEM/F-12 medium, and cell numbers were determined

using an Automated Cell Counter TC20TM (BIO-RAD Laboratories, Hercules, CA, USA) to

calculate the number of cells contained per well (4 cm^2).

Cell cycle analysis

The recovered cells as described above were fixed in 2-5 ml of cold (4°C) 70% methanol for

at least 30 minutes on ice. After centrifugation at 500g for 5 minutes at 4°C to obtain pellets, the

cell pellet was suspended in 50 μM propidium iodide and 20 × diluted RNase (19101, QIAGEN

Inc., Venlo, Netherlands) and then were incubated at 37°C for 30 minutes. After twice washing

with 1 ml of PBS at 300 g for 3 min at 4°C, the cells were analyzed by flow cytometry. The cell

population is shown in Supplementary Figure 9. A total of 20,000 events were analyzed.

RNA-sequencing

Total RNA was extracted from cells using the RNeasy Mini Kit (74106, QIAGEN) according

to the manufacturer's protocol. Three independent samples were assayed to evaluate the

reproducibility of the experimental procedures. The quality of total RNA (1 μg) submitted for

sequencing was checked using a Bioanalyzer (Agilent Technologies Inc., Santa Clara, CA,

USA), and all RNA integrity numbers were >7.0. The libraries for RNA sequencing were

sequenced with 2×150-bp paired-end reads on a NovaSeq6000 (Illumina Inc., San Diego, CA,

USA). The obtained sequencing data were analyzed using RaNA-seq *(58)*, an open

bioinformatics tool for rapid RNA-seq data analysis. FASTQ files were preprocessed in the

pipeline using the Fastp tool, and expression was quantified using Salmon. DEseq2 was used to

identify differentially expressed genes (DEGs) with adjusted p-values less than 0.05 between

the testosterone-treated groups in transcripts per million (TPM) between control and

testosterone-treated cells. The pathways were identified from the KEGG database on RaNA-seq.

Extracellular flux analysis

Oxygen consumption rate (OCR) and extracellular oxidation rate (ECAR) were measured

using an extracellular flux analyzer (XF HS Mini; Agilent Technologies). Eight-well plates for

the XF HS Mini were coated with fibronectin (5 μg/mL, F1141, Sigma-Aldrich) dissolved in

phosphate-buffered saline (PBS), incubated overnight, and then washed with Roswell Park

Memorial Laboratory medium for XF (Roswell Park Memorial Institute medium; RPMI,

103576-100, Agilent Technologies; containing 1% FBS) for XF before analysis. Mouse seminal

vesicle epithelial cells or human seminal vesicle epithelial cells treated with 100 ng/mL

testosterone for 7 days were suspended in RPMI medium at a concentration of 10,000 cells/180

μL, or mouse seminal vesicle epithelial cells infected with shRNA were suspended at a

concentration of 2000 cells/180 μL in each well. After centrifugation at $300 \times g$ for 3 minutes,

the plates were placed in an incubator at 37°C and 100% air and used for analysis within 1 hour.

For glucose metabolism assessment, cells were sequentially treated as indicated with

D-Glucose (10 mM; 103577-100, Agilent Technologies), Oligomycin (1 μM; O4876,

Sigma-Aldrich), and 2-Deoxy-Glucose (50 mM; D8375, Sigma-Aldrich).

For mitochondrial respiration measurement, cells were sequentially treated as indicated with

Oligomycin (1 μM), carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP; 5 μM,

SML2959, Sigma-Aldrich), and Rotenone (1 μM; R8875, Sigma-Aldrich)/Antimycin A (1 μM;

A8674, Sigma-Aldrich). Cellular metabolism readouts, such as glycolysis, glycolytic capacity,

glycolytic reserve, basal OCR, Maximum respiration, ATP Production, Spare respiratory

capacity, were determined using the Agilent Seahorse Wave software.

For mouse sperm analysis, NaHCO3-free HTF medium was prepared according to Balbach et

al. *(59)*. Plates were coated with concanavalin A (0.5 mg/mL, Fujifilm Wako Chemicals)

overnight the day before assay. Each well was seeded with 180 μL of NaHCO3- Free HTF

medium containing 3,000,000 sperm cells. Sperm were injected with oleic acid (final

concentration: 10 nM) with or without etomoxir (final concentration: 40 nM, E1905,

Sigma-Aldrich).

Measurement of pyruvate and lactic acid

For measurements, mouse or human seminal vesicle epithelial cells were treated with 100

ng/ml testosterone for 6 days. The culture supernatant was collected after 24 hr of additional

culture. Pyruvate consumption was measured using the Pyruvate Assay kit (MET-5029, Cell

Biolabs, Inc., San Diego, CA, USA) according to the manufacturer's protocol. The fluorescence

intensity was measured using a microplate reader (ex. 570 nm/em. 615 nm). Lactate released

into the medium was measured using the Lactate Assay kit (L256, DOJINDO

LABOLATORIES Co., Ltd., Kumamoto, Japan) according to the manufacturer's protocol. The

absorbance was measured using a microplate reader (wave length=450nm).

qPCR

The cDNA was synthesized from 40 ng of total RNA using oligo(dT)$_{15}$ primers (3805, Takara

Bio Inc., Shiga, Japan) and an Avian myeloblastosis virus (AMV) reverse transcriptase from

Promega (M5101, Madison, WI, USA). The cDNA and primers were added to Power SYBR

Green PCR Master Mix (4367659, Applied Biosystems, Foster City, CA, USA) in a total

reaction volume of 15 μl. Conditions were set to the following parameters: 10 min at 95°C, 15

sec at 95°C, and 1 min at 60°C. Cycles were repeated 45 times. Specific primer pairs were

selected and analyzed, as shown in Table S1. Expression was first normalized to housekeeping

gene *Rpl19*, and fold change was calculated relative to the mean of the control samples.

Glucose uptake test of seminal vesicle epithelial cells and lipid determination of culture supernatants

Mouse or human seminal vesicle epithelial cells treated with or without 100 ng/mL testosterone for 6 days were used for this experiment. Cells were treated with or without the GLUT4 inhibitor indinavir (100 nM) and 100 ng/mL testosterone for an additional 24 hours. Collected 15,000 cells were subjected to Glucose Uptake Assay Kit-Green (UP02; DOJINDO), and fluorescence intensity was measured using a microplate reader (ex. 485 nm/em. 535 nm).

Total lipid content in cultured medium was quantified using a lipid quantification kit (STA-617 Fluorometric, Cell Biolabs, Inc.) according to the manufacturer's protocol. The fluorescence intensity was measured using a microplate reader (ex. 485 nm/em. 535 nm).

Gas chromatography-mass spectrometry analysis

Fatty acid measurements in culture supernatant and seminal vesicle secretions were done according to our previous work *(18)*. Fatty acids in the samples were extracted and methylated using a fatty acid methylation kit (06482-04, Nacalai Tesque) and then purified using a fatty acid methyl ester purification kit (06483-94, Nacalai Tesque) according to the manufacturer's protocol. The methylated sample (3.0 mL) was dried and dissolved in 50 μL of elution solution. Samples were subjected to GC-MS.

Fatty acids were determined on an Agilent 7890A GC system coupled with a JMS-T 100 GCv mass detector at the Natural Science Research Center, Hiroshima University (N-BARD). Sample (1.0 μL aliquot) was injected to SP$^{®□}$-2560 capillary GC column (100 m × 0.25 mm i.d. × 0.2 μm film thickness; 24056, Sigma-Aldrich). For measurements, helium was used as the carrier gas at a constant flow rate of 2.0 mL/min. The GC oven temperature was gradually increased from 100°C to 300°C at a rate of 10°C/min and held at 300°C for 20 minutes. Ionization was performed using electron ionization with an electron energy of 70 eV. Mass spectra were obtained in scan mode (mass scan range 29-800 m/z). NIST MS Search v.2.0 was used to detect and identify of fatty acids.

Quantification of sperm oleic acid uptake by flow cytometry

To detect the fluorescence intensity of each sperm sample using flow cytometry, sperm

collected from the epididymis were incubated with 1, 10 or 100 μM of fluorescently labeled

oleic acid (F-OA-01, NANOCS, New York, NY, USA). After washing, fluorescence intensities

were analyzed by flow cytometry using a 530/30 nm bandwidth filter and measured as mean

intensity. A total of 100,000 sperm events were analyzed. The localization of fluorescence in

sperm was determined using an APX100 Digital Imaging System.

ACLY knock down experiment using shRNA

The U6 shRNA lentiviral vector of the *Acly* gene (mAcly[shRNA#1]; a virus packaging

service, VectorBuilder, Chicago, IL, USA) was used for knockdown experiments. The ACLY

sequence   targeted   by   the   shRNA   vector   used   in   this   study   was

5′-GCAGCAAAGATGTTCAGTAAA-3′. Lentivirally transduced cells were prepared according

to manufacture protocol. Immediately after isolation, the mouse seminal vesicle epithelial cells

were cultured overnight in a medium containing 5 μg/mL polybrene (VectorBuilder) and

lentiviral particles expressing an shRNA targeting ACLY or a scramble (Multiplicity of

Infection of 10). After an additional overnight incubation in a fresh culture medium, the cells

were incubated for 5 days in a medium containing 1 μg/ml puromycin (19752-64, Nacalai

tesque). 100 ng/mL of testosterone was added to all procedures for the testosterone-treated

group. Infection efficiency, calculated based on the number of GFP-positive cells using an

APX100 Digital Imaging System, ranged between 70% and 90%.

Western blot analysis

Protein samples of mouse seminal vesicle epithelial cells and seminal vesicle tissue were

prepared by homogenization in cell lysis buffer (04719964001, cOmplete™ Lysis-M

EDTA-free, Roche) and diluted with an equal volume of sample buffer solution containing

2-ME (2 ×) for SDS-PAGE (30566-22, Nacalai tesque). The extracts (10 μg protein) were

separated by SDS-polyacrylamide gel (10%) electrophoresis and transferred to polyvinylidene

fluoride membranes (10600069, Cytiva, Tokyo, Japan). The membranes were then treated with

Tris-buffered saline and Tween 20 (TBST, 20 mM Tris, pH 7.5, 150 mM NaCl, 0.1% Tween 20)

containing 5% (w/v) Non-Fat Dry milk (MORINAGA MILK INDUSTRY Co., Ltd., Tokyo,

Japan) for the nonspecific reaction was blocked. The membranes were incubated overnight at

4°C with primary antibodies: anti-ACLY antibody (1:10,000; ab40793; Abcam) or

anti-α/β-Tubulin antibody (1:1,000; 2148S; Cell Signaling). The next day, membranes were

washed with TBST and incubated with HRP-labeled antibody specific for rabbit IgG (1:4000;

7074S; Cell Signaling). The bands were visualized by using Enhanched Chemiluminescence

detection systems (RPN2232, Cytiva) and ChemiDoc^{TM} MP Imaging System (BIO-RAD

Laboratories). The bands were analyzed by using the software ImageJ (National Institutes of

Health; Bethesda, MD, USA).

ATP production in sperm

After 1 hour of incubation in the presence or absence of oleic acid (10 nM), ATP in sperm

was visualized by BioTracker ATP-Red (SCT045, Sigma-Aldrich). The sperm were incubated

with 10 nM of working solution containing BioTracker ATP-Red dye at 37 °C for 60 min in the

dark. After washing, the sperm pellet was resuspended in the base medium and analyzed with

the flow cytometry using a 488 nm laser and a filter with a bandwidth of 574/26 nm. A total of

100,000 sperm events were analyzed. The sperm population is shown in Supplementary Figure

6.

Artificial insemination

To examine the involvement of seminal vesicle secretions in fertilization, sperm from

C57BL/6N mice were injected directly into the uterine bodies of superovulated CD1 mice by

syringe, as previously described with minor modifications *(14, 19)*. CD1 female mice were

injected intraperitoneally with 8 IU eCG to stimulate follicle development and used as

recipients for artificial insemination 48 hours later. Artificial seminal fluid was prepared in HTF

medium to a final concentration of 10 mg/mL of collected control or frutamide-derived seminal

vesicle secretions. As rescue experiment, 10 nM of oleic acid was added to the

frutamide-derived seminal secretions. Cauda epididymis sperm was suspended with above

artificial seminal fluid (1×10^6 sperm/mL), and a total of 40 µL (20 µL each per uterus) was

injected through the cervix of the female mice. After artificial insemination, 7.5 IU of hCG was

injected to stimulate ovulation. Subsequently, a mixture of vaseline and paraffin with a ratio of

10:1 was injected to form an artificial plug around the cervix. After 36~48 hours of sperm

injection, the putative eggs were retrieved from the oviducts. The number of two-cell embryos

and mature unfertilized eggs were counted and used to estimate the fertilization success rate.

Statistical analysis

Quantitative data were presented as means ± SEM. All statistical details of the experiments

are given in the figure legends. R [(version 4.3.1 (2023-06-16)] was used for statistical

analysis .

Data availability

Nucleotide sequence data reported are available in the DDBJ Sequenced Read Archive under

the accession numbers DRA017090. Additional information needed to reanalyze the data

reported in this paper is available from the principal contact upon request.

List of Supplementary Materials

Supplemental Figure 1 to 9

Table S1

Supplementary Data

Acknowledgments:

The authors would like to thank JoAnne S. Richards for carefully proofreading the manuscript

and useful comments.

Funding:

This work was supported in part by JSPS KAKENHI Grant Number 23H00361 (to MS).

Author contributions:

Conceptualization: TY, TU, MS

Methodology: TY, ZX, NT, MA, TU

Investigation: TY, ZX, NT, MA, TU

Visualization: TY, MS

Funding acquisition: MS

Supervision: MS

Writing – original draft: TY

Writing – review & editing: TY, TU, MS

Competing interests:

MS holds stocks in and receives a salary from Hiroshima Cryopreservation Service Co. as a

director. In addition, MS has received royalties and grants from Hiroshima Cryopreservation

Service Co. M.S. has received consulting fees from Rohto Pharmaceutical Co., Ltd.. TU holds

stocks in and receives a salary from Lullabio Inc. as a director and has received honoraria from

Rohto Pharmaceutical Co., Ltd. The other authors declare they have no competing interests.

References

1. World Health Organization., Infertility prevalence estimates: 1990–2021. *Global report* , 1–

98 (2023).

2. S. Elbashir, Y. Magdi, A. Rashed, R. Henkel, A. Agarwal, Epididymal contribution to male

infertility: An overlooked problem. *Andrologia* **53**, e13721 (2021).

3. M. J. Cosentino, A. T. K. Cockett, Review Article: Structure and Function of the Epididymis

Transport of Spermatozoa. *Urol Res* **14**, 229–240 (1986).

4. B. Howarth, Fertilizing ability of cock spermatozoa from the testis epididymis and vas

deferens following intramagnal insemination. *Biol Reprod* **28**, 586–590 (1983).

5. S. Elbashir, Y. Magdi, A. Rashed, R. Henkel, A. Agarwal, Epididymal contribution to male

infertility: An overlooked problem. *Andrologia* **53**, e13721 (2021).

6. T. G. Cooper, Sperm maturation in the epididymis: A new look at an old problem. *Asian J*

Androl **9**, 533–539 (2007).

7. P. Gaddum-Rosse, Some observations on sperm transport through the uterotubal junction of

the rat. *Am J Anat* **160**, 333–341 (1981).

8. R. Shalgi, T. T. Smith, R. Yanagimachi2, A Quantitative Comparison of the Passage of

Capacitated and Uncapacitated Hamster Spermatozoa through the Uterotubal Junction. *Biol*

Reprod **46**, 419–424 (1992).

9. H. Li, P. Hung, S. S. Suarez, Ejaculated Mouse Sperm Enter Cumulus-Oocyte Complexes

More Efficiently In Vitro than Epididymal Sperm. (2015), doi:10.1371/journal.pone.0127753.

10. T. Noda, M. Ikawa, Physiological function of seminal vesicle secretions on male fecundity.

Reprod Med Biol **18**, 241–246 (2019).

11. P. Verze, T. Cai, S. Lorenzetti, The role of the prostate in male fertility, health and disease.

Nat Rev Urol **13**, 379–386 (2016).

12. E. Balandya, W. Wieland-Alter, K. Sanders, T. Lahey, Human Seminal Plasma Fosters

CD4+ Regulatory T-cell Phenotype and Transforming Growth Factor-β1 Expression. *American*

Journal of Reproductive Immunology **68**, 322–330 (2012).

13. T. Shima, K. Inada, A. Nakashima, A. Ushijima, M. Ito, O. Yoshino, S. Saito, Paternal

antigen-specific proliferating regulatory T cells are increased in uterine-draining lymph nodes

just before implantation and in pregnant uterus just after implantation by seminal

plasma-priming in allogeneic mouse pregnancy. *J Reprod Immunol* **108**, 72–82 (2015).

14. N. Kawano, N. Araki, K. Yoshida, T. Hibino, N. Ohnami, M. Makino, S. Kanai, H. Hasuwa,

M. Yoshida, K. Miyado, A. Umezawa, Seminal vesicle protein SVS2 is required for sperm

survival in the uterus. *Proc Natl Acad Sci U S A* **111**, 4145–4150 (2014).

15. R. R. ROSECRANS, R. S. JEYENDRAN, M. PEREZ-PELAEZ, W. P. KENNEDY,

Comparison of Biochemical Parameters of Human Blood Serum and Seminal Plasma.

Andrologia **19**, 625–628 (1987).

16. T. Umehara, T. Kawai, M. Goto, J. A. S. Richards, M. Shimada, Creatine enhances the

duration of sperm capacitation: a novel factor for improving in vitro fertilization with small

numbers of sperm. *Human Reproduction* **33**, 1117–1129 (2018).

17. Z. Zhu, T. Kawai, T. Umehara, S. A. M. Hoque, W. Zeng, M. Shimada, Negative effects of

ROS generated during linear sperm motility on gene expression and ATP generation in boar

sperm mitochondria. *Free Radic Biol Med* **141**, 159–171 (2019).

18. M. M. Islam, T. Umehara, N. Tsujita, M. Shimada, Saturated fatty acids accelerate linear

motility through mitochondrial ATP production in bull sperm. *Reprod Med Biol* **20**, 289–298

(2021).

19. T. Noda, Y. Fujihara, T. Matsumura, S. Oura, S. Kobayashi, M. Ikawa, Seminal vesicle secretory protein 7, PATE4, is not required for sperm function but for copulatory plug formation to ensure fecundity. *Biol Reprod* **100**, 1035–1045 (2019).

20. K. Queen, C. B. Dhabuwala, C. G. Pierrepoint, The effect of the removal of the various accessory sex glands on the fertility of male rats. *J Reprod Fertil* **62**, 423–426 (1981).

21. Mann T, Studies on the metabolism of semen; fructose as a normal constituent of seminal plasma; site of formation and function of fructose in semen. *Biochem J* **40**, 481–491 (1946).

22. Humphrey GF, Mann T, Studies on the metabolism of semen; citric acid in semen. *Biochem J* **44**, 97–105 (1949).

23. C. Chang, S. O. Lee, R. S. Wang, S. Yeh, T. M. Chang, Androgen receptor (AR) physiological roles in male and female reproductive systems: lessons learned from AR-knockout mice lacking AR in selective cells. *Biol Reprod* **89**, 1–16 (2013).

24. C. Fix, C. Jordan, P. Cano, W. H. Walker, Testosterone activates mitogen-activated protein kinase and the cAMP response element binding protein transcription factor in Sertoli cells. *Proc Natl Acad Sci U S A* **101**, 10919–10924 (2004).

25. K. Nagaosa, A. Kishimoto, R. Kizu, A. Nakagawa, A. Shiratsuchi, Y. Nakanishi, Perturbation of spermatogenesis by androgen antagonists directly injected into seminiferous tubules of live mice. *Reproduction* **133**, 21–27 (2007).

26. L. B. Smith, W. H. Walker, The regulation of spermatogenesis by androgens. *Semin Cell Dev Biol* **30**, 2–13 (2014).

27. P. S. Cooke, W. H. Walker, Male fertility in mice requires classical and nonclassical androgen signaling. *Cell Rep* **36**, 109557 (2021).

28. S. P. Smyth, B. Nixon, A. L. Anderson, H. C. Murray, J. H. Martin, L. A. MacDougall, S. A. Robertson, D. A. Skerrett-Byrne, J. E. Schjenken, Elucidation of the protein composition of mouse seminal vesicle fluid. *Proteomics* **22**, e2100227 (2022).

29. H. Murata, P. W. Hruz, M. Mueckler, Indinavir inhibits the glucose transporter isoform Glut4 at physiologic concentrations. *AIDS* **16**, 859–863 (2002).

30. C. Wilson, A. Contreras-Ferrat, N. Venegas, C. Osorio-Fuentealba, M. Pávez, K. Montoya, J. Durán, R. Maass, S. Lavandero, M. Estrada, Testosterone increases GLUT4-dependent glucose

uptake in cardiomyocytes. *J Cell Physiol* **228**, 2399–2407 (2013).

31. G. F. Gonzales, Function of seminal vesicles and their role on male fertility. *Asian J Androl*

3, 251–258 (2001).

32. A. Lenzi, M. Picardo, L. Gandini, F. Dondero, Lipids of the sperm plasma membrane: from

polyunsaturated fatty acids considered as markers of sperm function to possible scavenger

therapy. *Hum Reprod Update* **2**, 246–256 (1996).

33. M. T. Haren, A. M. Siddiqui, H. J. Armbrecht, R. T. Kevorkian, M. J. Kim, M. J. Haas, A.

Mazza, V. B. Kumar, M. Green, W. A. Banks, J. E. Morley, Testosterone modulates gene

expression pathways regulating nutrient accumulation, glucose metabolism and protein turnover

in mouse skeletal muscle. *Int J Androl* **34**, 55–68 (2011).

34. L. C. Costello, R. B. Franklin, Testosterone and prolactin regulation of metabolic genes and

citrate metabolism of prostate epithelial cells. *Horm Metab Res* **34**, 417–424 (2002).

35. K. Gonthier, R. T. K. Poluri, É. Audet-Walsh, Functional genomic studies reveal the

androgen receptor as a master regulator of cellular energy metabolism in prostate cancer. *J*

Steroid Biochem Mol Biol **191**, 105367 (2019).

36. M. F. Troncoso, M. Pavez, C. Wilson, D. Lagos, J. Duran, S. Ramos, G. Barrientos, P. Silva,

P. Llanos, C. Basualto-Alarcón, B. D. Westenbrink, S. Lavandero, M. Estrada, Testosterone

activates glucose metabolism through AMPK and androgen signaling in cardiomyocyte

hypertrophy. *Biol Res* **54**, 1–16 (2021).

37. Chen X, Li X, Huang HY, Li X, Lin JF, Effects of testosterone on insulin receptor

substrate-1 and glucose transporter 4 expression in cells sensitive to insulin. *Zhonghua Yi Xue*

Za Zhi **86**, 1474–1477 (2006).

38. K. Sato, M. Iemitsu, K. Aizawa, R. Ajisaka, Testosterone and DHEA activate the glucose

metabolism-related signaling pathway in skeletal muscle. *Am J Physiol Endocrinol Metab* **294**,

E961–E968 (2008).

39. P. K. Arnold, B. T. Jackson, K. I. Paras, J. S. Brunner, M. L. Hart, O. J. Newsom, S. P.

Alibeckoff, J. Endress, E. Drill, L. B. Sullivan, L. W. S. Finley, A non-canonical tricarboxylic

acid cycle underlies cellular identity. *Nature* **603**, 477 (2022).

40. A. Luengo, Z. Li, D. Y. Gui, L. B. Sullivan, M. Zagorulya, B. T. Do, R. Ferreira, A. Naamati,

A. Ali, C. A. Lewis, C. J. Thomas, S. Spranger, N. J. Matheson, M. G. Vander Heiden, Increased

demand for NAD+ relative to ATP drives aerobic glycolysis. *Mol Cell* **81**, 691–707 (2021).

41. B. B. Yeap, Testosterone and cardiovascular disease risk. *Curr Opin Endocrinol Diabetes*
Obes **22**, 193–202 (2015).

42. J. D. Marsh, M. H. Lehmann, R. H. Ritchie, J. K. Gwathmey, G. E. Green, R. J. Schiebinger,
Androgen receptors mediate hypertrophy in cardiac myocytes. *Circulation* **98**, 256–261 (1998).

43. G. F. Grabner, H. Xie, M. Schweiger, R. Zechner, Lipolysis: cellular mechanisms for lipid
mobilization from fat stores. *Nat Metab* **3**, 1445–1465 (2021).

44. B. Mittendorfer, Origins of metabolic complications in obesity: adipose tissue and free fatty
acid trafficking. *Curr Opin Clin Nutr Metab Care* **14**, 535–541 (2011).

45. H. Wackerhage, I. J. Vechetti, P. Baumert, S. Gehlert, L. Becker, R. T. Jaspers, M. H. de
Angelis, Does a Hypertrophying Muscle Fibre Reprogramme its Metabolism Similar to a
Cancer Cell? *Sports Med* **52**, 2569–2578 (2022).

46. R. J. De Berardinis, N. S. Chandel, Fundamentals of cancer metabolism. *Sci Adv* **2**,
e1600200 (2016).

47. G. L. Semenza, Hypoxia-inducible factors in physiology and medicine. *Cell* **148**, 399–408
(2012).

48. B. Morant-Ferrando, D. Jimenez-Blasco, P. Alonso-Batan, J. Agulla, R. Lapresa, D.
Garcia-Rodriguez, S. Yunta-Sanchez, I. Lopez-Fabuel, E. Fernandez, P. Carmeliet, A. Almeida,
M. Garcia-Macia, J. P. Bolaños, Fatty acid oxidation organizes mitochondrial supercomplexes
to sustain astrocytic ROS and cognition. *Nature Metabolism 2023 5:8* **5**, 1290–1302 (2023).

49. M. Duta-Mare, V. Sachdev, C. Leopold, D. Kolb, N. Vujic, M. Korbelius, D. C. Hofer, W.
Xia, K. Huber, M. Auer, B. Gottschalk, C. Magnes, W. F. Graier, A. Prokesch, B. Radovic, J. G.
Bogner-Strauss, D. Kratky, Lysosomal acid lipase regulates fatty acid channeling in brown
adipose tissue to maintain thermogenesis. *Biochim Biophys Acta Mol Cell Biol Lipids* **1863**,
467–478 (2018).

50. S. C. C. Huang, B. Everts, Y. Ivanova, D. O'Sullivan, M. Nascimento, A. M. Smith, W.
Beatty, L. Love-Gregory, W. Y. Lam, C. M. O'Neill, C. Yan, H. Du, N. A. Abumrad, J. F. Urban,
M. N. Artyomov, E. L. Pearce, E. J. Pearce, Cell-intrinsic lysosomal lipolysis is essential for
alternative activation of macrophages. *Nat Immunol* **15**, 846–855 (2014).

51. K. H. Tsui, T. H. Feng, Y. F. Lin, P. L. Chang, H. H. Juang, p53 downregulates the gene

expression of mitochondrial aconitase in human prostate carcinoma cells. *Prostate* **71**, 62–70

(2011).

52. S. Almeida, L. Rato, M. Sousa, M. G. Alves, P. F. Oliveira, Fertility and Sperm Quality in

the Aging Male. *Curr Pharm Des* **23**, 4429–4437 (2017).

53. T. Murata, Y. Ishitsuka, K. Karouji, H. Kaneda, H. Toki, Y. Nakai, S. Makino, R. Fukumura,

H. Kotaki, S. Wakana, T. Noda, Y. Gondo, β-cateninC429S mice exhibit sterility consequent to

spatiotemporally sustained Wnt signalling in the internal genitalia. *Scientific Reports 2014 4:1* **4**,

1–7 (2014).

54. B. A. Stone, A. Alex, L. B. Werlin, R. P. Marrs, Age thresholds for changes in semen

parameters in men. *Fertil Steril* **100**, 952–958 (2013).

55. J. L. Frattarelli, K. A. Miller, B. T. Miller, K. Elkind-Hirsch, R. T. Scott, Male age

negatively impacts embryo development and reproductive outcome in donor oocyte assisted

reproductive technology cycles. *Fertil Steril* **90**, 97–103 (2008).

56. A. P. Drabovich, P. Saraon, K. Jarvi, E. P. Diamandis, Seminal plasma as a diagnostic fluid

for male reproductive system disorders. *Nat Rev Urol* **11**, 278–288 (2014).

57. T. Umehara, N. Tsujita, Z. Zhu, M. Ikedo, M. Shimada, A simple sperm-sexing method that

activates TLR7/8 on X sperm for the efficient production of sexed mouse or cattle embryos. *Nat*

Protoc **15**, 2645–2667 (2020).

58. C. Prieto, D. Barrios, RaNA-Seq: Interactive RNA-Seq analysis from FASTQ files to

functional analysis. *Bioinformatics* **36**, 1955–1956 (2019).

59. M. Balbach, J. Buck, L. R. Levin, Using an Extracellular Flux Analyzer to Measure

Changes in Glycolysis and Oxidative Phosphorylation during Mouse Sperm Capacitation. *J Vis*

Exp **2020**, e60815 (2020).

894 **Figures and Figure legends**

895 **Fig.1**

Figure 1. Effects of seminal vesicle fluid and prostate fluids on sperm motility (A) To determine the effect of seminal plasma on sperm motility, we prepared "pseudo" seminal plasma from the mixture of seminal vesicle secretions or prostate extracts. (B-D) Tested various concentrations of these fluids on epididymal sperm: (B) Curvi-Linear Velocity; VCL, (C) Straight-Line Velocity; VSL and (D) Linearity; LIN is the ratio of VSL to VCL of sperm that was incubated with the mixture of seminal vesicle secretions and/or prostate extracts. (E) Experimental design to evaluate the effect of seminal secretions collected from male mice treated with or without flutamide (50 mg/kg subcutaneously for 7 days) on sperm. (F-H) Performed bioassay using seminal vesicle fluid after treatment with vehicle (Ctrl) or flutamide: (F) LIN, (G) VSL and (H) the activity of mitochondria were checked by JC-1 kit. Data are mean ± SEM. n=3-5 independent replicates. Significance was tested in comparison to HTF or Ctrl (vehicle-treated animals) using Student's t-test.

910 **Fig.2**

911
912

Figure 2. The testosterone-androgen receptor pathway inhibits the cell proliferation of
seminal vesicle gland epithelial cells. (A) The localization of the androgen receptor (AR;
NR3C4) in seminal vesicle of control (Ctrl) mice and those treated with flutamide (Fult). Scale
bar=50 μm for left panel and 20 μm for right panel in each group. (B,C) Cell proliferation and
apoptosis in seminal vesicle of control (Ctrl) mice and those treated with flutamide (Fult): (B)
Representative images of Ctrl and Flut staining for Ki67 (top) and TUNEL (bottom). Scale
bar=20 μm. (C) Percentage of Ki67 positive cells (top) and TUNEL positive (bottom) in Ctrl
and Flut-treated seminal vesicle sections. (D) Experimental design to evaluate the effects of
testosterone on the proliferation of seminal vesicle epithelial cells *in vitro*. The epithelial cells
were culutured with or without testosterone for 8 days. (E) Growth curves of seminal vesicle
epithelial cells that cultured with 0, 1, 10 and 100 ng/mL of testosterone. (F,G) Cell cycle status
was determined by flow cytometry. Data are mean ± SEM. n=3-5 mice or independent
replicates. Each replicate experiments with 3-6 wells containing pooled cells from 3-5 mice.
Significance was tested in comparison to Ctrl using Student's t-test. The cell proliferation data
were compared to Ctrl by Dunnett's test.

Fig.3

Figure 3. Testosterone changes the expression of genes involved in metabolic pathway in
seminal vesicle epithelial cells. (A) Volcano plot of differentially expressed genes. RNA
sequencing was performed using RNA extracted from the seminal vesicle epithelial cells
cultured with or without 100 ng/mL testosterone. Genes with a significant expression change are
highlighted as red dots. It found 4460 significant genes for a cutoff of P<0.05. (B) MA plot of
differential expression results. The seminal vesicle specific genes are highlighted as red dots
rather than significant genes. (C) Top 10 genes upregulated or downregulated by testosterone,
respectively, for altered gene expression. (D) Conducted KEGG analysis to identify differences
in pathway enrichment by testosterone, identifying the most variability in metabolic pathway
genes. It shows the number of differential expressed genes annotated to Gene Ontology shows
(x axis). p-value which measure the statistical significance of a possible functional enrichment
for each term.

Fig.4

944
Figure 4. Testosterone regulates glucose metabolism and mitochondrial ATP production in
seminal vesicle epithelial cells. (A-C) Glycolytic pathway analysis by an extracellular flux
analyzer in seminal vesicle gland epithelial cells cultured with 100 ng/mL testosterone (Testo)
or vehicle (Ctrl) for 7 days: (A) Extracellular acidification rate (ECAR) kinetics of seminal
vesicle epithelial cells using an extracellular flux analyzer. (B) Glycolysis and (C) Glycolytic
capacity. (D) Pyruvate concentration in the medium where seminal vesicle epithelial cells were
cultured with or without testosterone for 24 hours. (E) Lactate concentration in the medium
where seminal vesicle epithelial cells were cultured with or without testosterone for 24 hours.
(F-J) Mitochondrial respiration measurement by an extracellular flux analyzer: (F) Oxygen

consumption rate (OCR) kinetics of seminal vesicle epithelial cells with or without 100 ng/mL

testosterone. (G) Basal OCR, (H) Maximum respiration, (I) Spare respiratory capacity and (J)

ATP production. Data are mean ± SEM of n = 3 replicate experiments with 3-6 wells containing

pooled cells from 3-5 mice or medium. Student's t-test was used to compare Ctrl and Testo.

Figure 5. Effect of testosterone on gene expression of enzymes involved in the glucose

metabolic pathway in seminal vesicle epithelial cells. To elucidate the effects of testosterone

on gene expression of enzymes involved in glucose catabolism and anabolism in seminal

vesicle gland epithelial cells. ctrl: 7 days of culture with vehicle. testo: 7 days of culture with

100 ng/mL testosterone. Data are mean ± SEM of n = 3 replicate experiments with 3-6 wells

containing pooled cells from 3-5 mice. Student's t-test was used to compare Ctrl and Testo.

Fig.6

Figure 6. Oleic acid is incorporated to sperm, which enhances linear motility and

mitochondrial activity. (A) Epididymal sperm were incubated with fluorescently labeled oleic

acid (OA) and the fluorescence intensity after quenching was observed by a fluorescence

microscope. (B) The fluorescence intensity of fluorescence-conjugated OA in sperm was

detected by flow cytometry. (C) Average fluorescence intensity of sperm at different

concentrations of fluorescence-conjugated OA. (D,E) OA improved sperm motility parameters.

Sperm collected from mouse epididymis was incubated with OA for 60 min: (D) Straight-Line

Velocity; VSL and (E) Linearity; LIN (F) Increase in the Oxygen consumption rate (OCR) after

treatment with 10 nM OA or 10 nM OA + 40 nM Etomoxir (Eto). (G) The percentage of

ATP-Red positive sperm after 1 hour incubation with or without 10 nM OA, measured using

BioTracker ATP-Red staining and flow cytometry. (H) Artificial insemination using sperm

treated with "pseudo" seminal plasma from control (SV), flutamide-treated (Flu) mice and

supplement of OA (Flu+OA) followed by an evaluation of the fertilization rates. Data are mean

$\pm$ SEM. n=3-6 independent replicates. Student's t-test was used for comparison between the two

groups.

Fig.7

Figure 7. Testosterone-regulated ACLY induces metabolic shifts in seminal vesicle

989 **epithelial cells.** (A) Western blot images of ACLY and α/β-tubulin in three sets of seminal

990 vesicle epithelial cells cultured with 100 ng/mL testosterone (Testo) or in vehicle (Ctrl) for 7

991 days. (B) Quantitative analysis of ACLY relative to α-tubulin obtained from Western blot. (C)

992 siRNA knockdown experiments of ACLY in seminal vesicle epithelial cells. ACLY protein

levels in scrambled shRNA or ACLY shRNA-transfected seminal vesicle epithelial cells cultured with or without 100 ng/mL testosterone were determined by Western blot. (D-F) Changes in oxygen consumption of seminal epithelial cells were analyzed by a flux analyzer: (D) Oxygen consumption rate (OCR) kinetics of seminal vesicle epithelial cells transfected with scrambled shRNA or ACLY shRNA. (E) Basal OCR and (F) Maximum respiration. (G) The impact of ACLY knockdown on testosterone-induced fatty acid synthesis was measured. (H) The effect of culture supernatants of testosterone-treated cells on sperm motility was evaluated, especially after ACLY knockdown. Curvi-Linear Velocity; VCL, Straight-Line Velocity; VSL, Linearity; LIN. Data are mean ± SEM. n=3-6 independent replicates. Repeated experiments were performed with cells recovered from 3 wells containing pooled cells from 3 to 5 mice. Student's t-test was used for comparison between the two groups.

 Fig.8

1006

Figure 8. Testosterone regulates the metabolic activity of human seminal epithelial cells.

(A) Representative image of human seminal vesicle epithelial cells. (B) Growth curves of

seminal vesicle epithelial cells with 100 ng/mL testosterone. (C) Oxygen consumption rate (OCR) kinetics and (D) Extracellular acidification rate (ECAR) kinetics of human seminal vesicle epithelial cells after 6 days of culture with or without 100 ng/mL testosterone using a flux analyzer. (E) Effects of testosterone on fluorescent glucose uptake and inhibitors of GLUT4 function. (F) Effects of testosterone and GLUT4 inhibitors on fatty acid synthesis. Data are mean ± SEM. n=3 independent replicates. Student's t-test was used for comparison between the two groups.