Title: Testosterone-Induced Metabolic Changes in Seminal Vesicle Epithelial cells Alter

Plasma Components to Enhance Sperm Fertility

Authors: Takahiro Yamanaka[1], Zimo Xiao[1], Natsumi Tsujita[1], Mahmoud Awad[1,2], Takashi

Umehara[1], Masayuki Shimada[1,3*]

Affiliations:

[1] Graduate School of Integrated Sciences for Life, Hiroshima University; Higashihiroshima,

Hiroshima, Japan

[2] Department of Histology, Faculty of Veterinary Medicine, South Valley University; Qena, Egypt

[3] Graduate School of Innovation and Practice for Smart Society, Hiroshima University;

Higashihiroshima, Hiroshima, Japan

*Corresponding author. Masayuki Shimada, Ph.D. E-mail: mashimad@hiroshima-u.ac.jp,

Tel/Fax: (+81)-824-24-7899

One Sentence Summary: Testosterone induces ACLY expression in seminal vesicle, a key factor

of forming seminal plasma to acquire *in vivo* fertilization ability of sperm.

Abstract:

Male infertility depends on both sperm and seminal plasma and is induced by aging. In this

study, male infertility was examined with seminal plasma and its synthesis mechanism. The

factors ensuring *in vivo* fertilization potential was secreted from seminal vesicle where the factors

were synthesized in an androgen-dependent manner. Androgen increased glucose uptake and

glycolytic capacity in seminal vesicles, which caused activation of oleic acid synthesis rather than

mitochondrial ATP synthesis. ACLY was identified as a key player in this metabolic mechanism

for producing oleic acid that was incorporated into the sperm and enhanced fertilization potential

in vivo. In conclusion, an important role of testosterone-induced metabolic pathways in the

seminal vesicle was to ensure the synthesis of oleic acid, which is essential for sperm fertilization

in vivo. These results provide new perspectives for the development of biochemical markers of

semen to assess male fertility and for artificial insemination techniques.

Main Text:

INTRODUCTION

Infertility is defined by the World Health Organization (WHO) as the failure to conceive within 12 months despite contraceptive-free sexual intercourse. Its prevalence has increased over the decades and now accounts for 17.5% of couples worldwide *(1)*. The male partner is thought to be solely or in combination responsible for 50% of infertility cases, and the male factor in the etiology of infertility is highly recognized. The male fertility depends largely on the quality of semen composed of sperm and seminal plasma and decreases with aging.

Spermatogenesis, the process by which mature sperm are generated, takes place partially in the testis, where the characteristic sperm morphology of the head, middle, and tail is formed. Within testis however, sperm are functionally immature and cannot penetrate an oocyte until they undergo a maturation process that occurs during their passage through the epididymis *(2–4)*. Sperm recovered from the epididymis can fertilize a mature oocyte *in vitro* but cannot maintain sustained liner motility required for migration from the uterus to the oviduct *(5, 6)*. Thus, when epididymal sperm are artificially inseminated to the cervix or uterus, fertilization rates are low *(7–9)*. However, when epididymal sperm are mixed with seminal plasma at the time of ejaculation or when sperm are exposed to seminal plasma *in vitro*, even for only a brief period of time, they acquire *in vivo* fertilization potential and are capable of passing through the female reproductive tract and fertilizing ovulated oocyte *in vivo (10)*. Thus, sperm formation occurs in the testis, they mature in the epididymis, but only acquire the *in vivo* fertilization ability after exposure to seminal plasma.

Seminal plasma is mainly produced in the seminal vesicles (approximately 65-75%), with approximately 20-30% produced by the prostate, and small amounts secreted by the bulbourethral and urethral glands *(11)*. Seminal plasma or TGFβ, a component of seminal plasma, increases antigen-specific Treg cells in uterus of mice and humans which induce immune tolerance, resulting in pregnancy *(12, 13)*. Furthermore, SVS2 that is secreted from the seminal vesicle is also known to be required for sperm to escape attack by the female immune system in uterus *(14)*. In addition, seminal plasma is known to contain various biochemical components that support sperm metabolism, including lipids and organic acids. Specifically, Rosecrans et al. *(15)* found significant differences in the levels of most biochemical components in seminal plasma compared

to blood, with specific factors like fructose being rarely detected in blood but abundant in seminal

plasma. Our previous studies *(16–18)* documented that seminal plasma also contains creatine, and

fatty acids, that are taken up by sperm within minutes. These substances play crucial roles in

regulating sperm metabolic pathways and enhancing sperm motility. Collectively these

observations indicate that seminal plasma not only alters the female immune system to protect

sperm but also directly influences sperm metabolic pathways, facilitating the transformation of

sperm capable of *in vivo* fertilization.

Some specific factors that male accessory glands contribute to seminal plasma and impact male

fertility have been elucidated. For example, surgical removal of seminal vesicles in male mice and

rats was associated with infertility *(14, 19, 20)*. The observation that fructose *(21)* and citric acid

(22) concentrations in seminal plasma of control mice and rats are higher than in castrated

animals indicates that specific functions of the accessory glands might be affected by

testicular-derived testosterone that is known to activate the intracellular androgen receptor

(AR;NR3C4), required for gene regulation of transcription. AR-deficient mice have been found

to be infertile *(23)*, and long-term administration of androgen receptor antagonists, like flutamide,

abolished male fertility *(24, 25)*. AR-deficient mice and transcriptome analyses have identified

specific targets of AR *(26, 27)*. In both mice and human, androgen levels decrease with aging, and

their concentrations have been reported to cause a decrease in male fertility, especially ejaculated

sperm motility. Therefore, the reduction of androgen concentrations may affect the metabolic

mechanisms of the accessory reproductive organs and alter the composition of seminal plasma

with aging. However, the metabolic mechanisms that impact sperm functions are governed by a

complex of enzymatic reactions and therefore, the intricate interplay between testosterone and

seminal plasma synthesis in the accessory reproductive organs requires further investigation.

In this study, we first performed various bioassays to identify which accessory glands are

essential for sperm fertility. Subsequently, comparative *in vivo* and *in vitro* analyses were

performed using a hypoandrogenic model. In addition, biochemical and molecular techniques

such as RNA-seq, flux analysis, mass spectrometry, and knockdown experiments were employed.

These analyses reveal the pathways by which seminal vesicles synthesize seminal plasma, ensure

fertilization ability, and provide new therapeutic and preventive strategies for male infertility.

Seminal vesicle and prostate-derived fluids induce functional sperm motility

A bioassay using epididymal sperm that had not been exposed to seminal plasma was done to clarify the effect of seminal plasma on sperm functions. For this bioassay, fluids were prepared from seminal vesicle secretion and homogenized prostate tissues, and then the fluid was mixed with the basic medium, Human Tubal Fluid (HTF) medium in various proportions (Fig.1A-D). The addition of seminal vesicle secretion alone to the basic medium (HTF medium) significantly increased the rate of sperm linear motility (LIN) compared to the HTF medium alone (Fig.1D). The straight line velocity (VSL) of those sperm was also significantly increased (Fig.1C). This significant increase in straightening of sperm by the seminal vesicle-derived secretion was further significantly positively affected by the addition of prostate extract 1:9 or 2:8 to the seminal vesicle secretion (Fig.1C,D). However, no significant improvement in straightness was observed with the mixture of prostate extract alone (Fig.1C,D). These results clearly indicate that sperm exposure to seminal plasma factors significantly improves sperm linear motility and that this positive effect is mainly due to the seminal vesicle secretory factors.

The functions of male accessory reproductive organs are regulated by androgens. Therefore, to elucidate whether the seminal vesicle factors, which enhance sperm linear motility, are synthesized in an androgen-dependent manner, bioassays were performed using seminal vesicle secretions collected from control mice or mice treated for 7 consecutive days with flutamide, an androgen receptor antagonist, and sperm collected from the epididymis of mice not treated with flutamide (Fig.1E). The results showed that LIN and VSL were significantly reduced in the seminal vesicle secretions from flutamide-treated mice (Flutamide) compared to the control mice (Ctrl) (Fig.1F,G). The reduction of sperm linier motility speed (VSL) was also significantly lower in the sperm treated with seminal vesicle secretions from over 12 months old mice that showed low circular testosterone level as compared with the sperm treated with seminal vesicle secretions from young adult mice (3 months old mice) (Supplemental Fig.1A,B). Furthermore, JC-1 staining to measure mitochondrial membrane potential, which is important for sperm linear motility, showed significantly lower mitochondrial membrane potential activity in sperm treated with the secretions of seminal vesicles from flutamide-treated mice (Fig.1H). These results indicated that androgen-dependent changes in seminal vesicle functions alter the levels/activity of seminal vesicle secretion factors, which regulate sperm linear motility. The changes of seminal vesicle

functions were also induced with aging.

Testosterone-androgen receptor activity suppresses the proliferation of seminal vesicle

epithelial cells

To understand the difference between seminal vesicles that secrete factors for enhancing sperm

linear motility and those that are impaired in this function, we performed histological analysis of

seminal vesicles recovered from mice treated and non-treated with flutamide. In control mice, the

seminal vesicle epithelial cells were in a single layer, and androgen receptors were localized in the

nuclei of seminal vesicle epithelial cells (Fig.2A). However, in mice treated with continuous

administration of flutamide, seminal vesicle epithelial cells were multilayered, and the strong

accumulation of androgen receptors in nucleus was not observed (Fig.2A). To investigate the cause

of this multilayering of cells, we performed Ki67 staining, a marker of cell proliferation, and

apoptosis detection by TUNEL staining (Fig.2B). The results showed that the percentage of

Ki67-positive cells increased significantly after treatment with flutamide, but the percentage of

TUNEL-positive cells did not change, suggesting that the testosterone-androgen receptor pathway

is responsible for the inhibition of proliferation of seminal vesicle epithelial cells (Fig.2C). These

changes of abnormal morphology and high proliferation activity were also observed in seminal

vesicle of more than 12 months old mice that showed the low level of testosterone in serum

(Supplemental Figure 1C-E). Therefore, seminal vesicle epithelial cells were collected and cultured

in primary culture as shown in Fig. 2D, and the changes in cell number were counted and cell cycle

was analyzed by flow cytometry. When cultured without testosterone (c), the number of seminal

vesicle epithelial cells increased in a day-dependent manner. However, the addition of testosterone

significantly decreased the number of seminal vesicle epithelial cells during the 8-day culture

period in a concentration-dependent manner (Fig.2E). Flow cytometric analysis of the cell cycle on

day 7 showed that the G0/G1 phase was significantly increased and the G2/M1 phase was

significantly decreased by the addition of testosterone (Fig. 2F,G). These results indicate that

seminal vesicle epithelial cells are inhibited in cell division by the testosterone-androgen receptor

pathway.

Testosterone alters metabolic gene expression in seminal vesicle epithelial cells

To better understand the roles of testosterone in suppressing proliferation of seminal vesicle epithelial cells, RNA sequencing was done using the epithelial cells cultured with or without testosterone. DEG (Differential Expressed Genes) analysis revealed that a total of 23,459 genes were identified, including 997 upregulated genes and 3463 downregulated genes in seminal vesicle epithelial cells by testosterone relative to control (Fig.3A). Genes largely upregulated by testosterone included those already reported in seminal vesicles *(14, 19, 28)*, such as the Seminal Vesicle Secretion (*Svs*) family (*Svs1~6*), *Pate4*, *Spinkl*, *Ceacam10*, and *Sva* (Fig.3B,C). Therefore, the primary culture of seminal vesicle epithelial cells used in this experiment was considered to maintain the characteristics of seminal vesicle epithelial cells *in vivo*. Many genes were downregulated by testosterone, and those with a high rate of variation included collagen synthase and *Mmp2*, which are involved in extracellular matrix formation (Fig.3C). However, KEGG analysis, rather than individual genes, was used to predict functions altered by testosterone, and it was found that the expression of genes involved in the metabolic pathway was the most highly affected by testosterone (Fig.3D). Because there is a close relationship between cell proliferation and metabolic activity, and between metabolic substrates in seminal plasma and sperm motility, we hypothesized that testosterone induced changes in seminal vesicle epithelial cell metabolism alter the components of seminal plasma to mediate the enhancement of sperm motility.

Testosterone sifts metabolic activity of seminal vesicle epithelial cells

In the presence of testosterone, the addition of glucose significantly activated glycolysis as compared with that in epithelial cells cultured without testosterone (Fig.4A,B). However, there was no difference in the glycolytic capacity, indicating that testosterone may increase glucose metabolism but does not to enhance the expression of a group of enzymes involved in the glycolytic pathway. To determine the metabolic fate of glycolytic end products, seminal vesicle epithelial cells were cultured with and without testosterone and then concentrations of pyruvate and lactate were measured. The concentration of pyruvate was significantly decreased by the addition of testosterone; however, the concentration of lactate was not significantly changed (Fig.4D,E). Therefore, we analyzed mitochondrial metabolism using a flux analyzer, predicting that

more glucose is metabolized, pyruvate is synthesized in response to testosterone, and is further

metabolized in the mitochondria. However, measuring mitochondrial respiratory metabolism with

a Flux analyzer, the basal respiratory rate (the activity of the electron transport chain) was

significantly decreased by testosterone (Fig.4F-I). ATP production was also significantly

suppressed exposure to testosterone. Furthermore, the maximum respiration and spare respiration

capacity were also significantly decreased in response to testosterone treatment, suggesting that

testosterone does not educe transient enzyme activity in mitochondria, but to rather weakens the

metabolic pathway of the mitochondrial TCA cycle and/or the electron transport chain due to the

changes in gene expression patterns in seminal vesicle epithelial cells.

Testosterone causes changes in the expression of genes encoding enzymes involved in

glucose catabolism and anabolism

The expression of genes involved in glucose catabolism and anabolism in the *in vitro* culture

system of seminal vesicle epithelial cells was analyzed by qPCR (Fig.5). The gene expression of

glucose transporter (*Glut*) and enzymes involved in the glycolytic pathway were not significantly

affected by testosterone, consistent with the results of metabolic flux analysis, which showed that

the glycolytic pathway was not enhanced despite the activation of the glycolysis. Furthermore,

the expression of genes encoding enzymes involved in the TCA cycle in mitochondria was also

not significantly affected by testosterone, but the expression of enzymes involved in the electron

transport chain was significantly decreased. Interestingly, the expression of *Acly*, that encodes an

enzyme that reconverts citric acid formed in the TCA cycle to Acetyl CoA and induces cholesterol

synthesis and fatty acid synthesis, was significantly increased by testosterone. Although *Hmgcr*,

which encodes the rate-limiting enzyme involved in cholesterol synthesis, was unchanged, the

expression of *Acc*, which encodes the rate-limiting enzyme for fatty acid synthesis, was

significantly upregulated by testosterone. These characteristic testosterone-induced changes in

gene expression observed in cultured cells were also detected by ACLY immunostaining of *in*

vivo seminal vesicle samples (Supplemental Figure 2). Specifically, the expression of genes

encoding a group of enzymes of the electron transport chain was significantly upregulated by

continuous flutamide administration *in vivo*, whereas the gene expression of *Acly* and *Acc* was

significantly decreased by flutamide administration. Immunostaining results showed that ACLY

was detected specifically in seminal vesicle epithelial cells, and their staining appeared to be reduced by flutamide treatment. The lower level of ACLY was also observed in seminal vesicle epithelial cells of over 12 months old mice (Supplemental Figure 2B).

Mechanism of testosterone-induced enhancement of glucose uptake and thereby fatty acid synthesis

The effect of testosterone on glucose uptake in seminal vesicle epithelial cells was analyzed using fluorescence-labeled glucose. As expected from the results of flux metabolic analyses, testosterone treatment significantly increased glucose uptake (Supplemental Figure 3A). Because the expression of each *Glut* family gene encoding glucose transporters was not altered by testosterone (Fig.5), we focused on GLUT4, whose cellular localization is regulated, and analyzed the effects of Indinavir, a functional inhibitor of GLUT4 *(29, 30)*. Indinavir significantly suppressed glucose uptake and significantly decreased fatty acid synthesis (the amount of fatty acids accumulated in the cell), which was increased by testosterone (Supplemental Figure 3A,B). Furthermore, immunostaining analysis of the localization of GLUT4 in seminal vesicles revealed that GLUT4 was localized outside the nucleus of seminal vesicle epithelial cells (Supplemental Figure 4), suggesting that GLUT4 was partially localized at the plasma membrane and contributed to glucose uptake, whereas in seminal vesicles from mice treated with flutamide continuously, no strong signal was detected at the presumed location of GLUT4 near the plasma membrane of seminal vesicle epithelial cells (Supplemental Figure 4). The same characteristics were observed in aged mice.

Qualitative and quantitative analysis of the fatty acids synthesis was performed by GC-MS. In seminal vesicles, C16:0, C18:0 and C18:1 were detected, in which C18:1 was decreased to an undetectible level in seminal vesicles of mice treated with the flutamide (Supplemental Figure 5A). In the cultured seminal vesicle epithelial cells, C18:1, oleic acid, was significantly increased by the addition of testosterone as observed in seminal vesicle (Supplemental Figure 3C). The expression of *Elovl6* encoding ELOVL6, which desaturates C18:0 saturated fatty acids to C18:1, was significantly up-regulated by testosterone (Supplemental Figure 3D). The testosterone-induced change was also detected in *in vivo* samples (seminal vesicle), where the expression of *Evovl6* was decreased by the flutamide treatment (Supplemental Figure 5B).

Oleic acid (18:1) is taken up by sperm and promotes their linear movement

We observed that seminal vesicles secrete testosterone-dependent factors that induce sperm to move straight and that seminal vesicle epithelial cells synthesize and secrete oleic acid in a testosterone-dependent manner. In order to examine whether oleic acid is a factor that induces sperm linear motility, we examined whether or not oleic acid is taken up by sperm and then changes of sperm motility pattern. When epididymal sperm was incubated with fluorescently labeled oleic acid, the fluorescent signal was selectively detected in the midpiece of the sperm after quenching of extracellular fluorescence (Fig.6A). The fluorescence intensity of sperm detected by flow cytometry significantly increased in a dose-dependent manner (Fig.6B, C). In oleic acid-treated sperm, VSL was significantly increased in a dose-dependent manner in the range of 0 to 50 nM (Fig.6D). LIN values were highest when 10 nM oleic acid was added (Fig.6E). To quantify metabolic changes in mitochondrial activity, a flux analyzer was used to monitor real-time changes in mitochondrial β-oxidation. A significant increase in OCR was induced by the injection of 10 nM oleic acid. Etomoxir, a β-oxidation inhibitor, significantly decreased the OCR that was induced by oleic acid (Fig.6F, Supplemental Figure 6). In situ ATP detection by BioTracker ATP-Red revealed that the percentage of sperm with high ATP production was significantly increased under the presence of oleic acid (Fig.6G).

To examine whether the addition of oleic acid to low quality of seminal plasma (collected from flutamide-treated mice) restores *in vivo* the fertilization potential of epididymal sperm, sperm treated with seminal plasma collected from control or flutamide-treated mice was used for artificial insemination. The *in vivo* fertilization rate was about 70% when sperm was treated with the seminal vesicle secretions from control mice (SV); however, the fertilization rate was significantly lower in the group that sperm was treated with the seminal vesicle secretions from flutamide-treated mice (Flu) (SV: $65.5 \pm 3.8\%$, Flu: $39.4 \pm 7.8\%$). Although not significant, adding oleic acid to the seminal secretions of flutamide-treated mice (Flu+OA) increased the fertilization rate to $53.1 \pm 1.3\%$ (Fig.6H).

ACLY expression is upregulated by testosterone and is essential for the metabolic shift of seminal vesicle epithelial cells that mediates sperm linear motility

ACLY is up-regulated in seminal epithelial cells as a factor that enables the utilization of citric

acid from the TCA cycle for fatty acid synthesis outside the mitochondria. Therefore, we

performed siRNA knockdown experiments of ACLY to clarify the impact of ACLY on the

cellular metabolic shift. The addition of testosterone significantly increased the amount of

ACLY protein in seminal epithelial cells (Fig.7A,B). Viral infection with siRNAs encoding

Acly-targeted siRNAs markedly decreased the amount of ACLY (Fig.7C). This knockdown of

ACLY not only increased the oxygen consumption of seminal epithelial cells in the control and

testosterone-treated group, but also significantly increased the ATP production related oxygen

consumption (Fig.7D,E,F). Thus, ACLY expression is associated with mitochondrial metabolic

pathway activity in seminal epithelial cells. Furthermore, the knockdown of ACLY also

significantly suppressed testosterone-induced fatty acid synthesis in seminal epithelial cells

(Fig.7G). Supernatants from epithelial cells of the testosterone-added group induced sperm to

move linearly (Fig.7H). However, the knockdown of ACLY significantly reduced the effects of

the culture supernatant on sperm linear motility (Fig.7H).

This characteristic metabolic mechanism induced by testosterone is also observed in

human seminal epithelial cells

To determine whether these findings in mice could be applied to humans, the commercially

available human seminal vesicle epithelial cells were used for following studies (Fig.8A). The

growth curves showed that cell proliferation was inhibited by 100 ng/mL of testosterone

compared to the control; a pattern similar to that observed in murine seminal vesicle epithelial

cells (Fig.8B). The characteristics of metabolic activity of human seminal epithelial cells were

also similar to those of mice, with testosterone significantly enhancing the glycolytic pathway

(Fig.8C), but significantly reducing the capacity for oxygen consumption in mitochondria

(Fig.8D). Moreover, fluorescent-glucose uptake was increased by testosterone and decreased by

the inhibitor of GLUT4 function (Fig.8E). Additionally, while testosterone also significantly

increased fatty acid synthesis, this increase was completely abolished by the GLUT4 inhibitor

(Fig.8F).

DISCUSSION

Seminal plasma has a variety of functions, including sperm transport, nutritional support,

interactions with the female reproductive tract, and especially the regulation of its relationship

with the female immune system. Mammalian seminal plasma is secreted mainly from the

seminal vesicle and is known to be rich in cytokines, prostaglandins, and SVS family members

(31). For this reason, recent studies have focused on the immunomodulatory function of seminal

plasma in the female reproductive tract. On the other hand, fructose and lipids in seminal plasma

are the major energy sources for sperm *(32)*. However, although metabolomic analyses of

seminal plasma have shown that there are metabolic substrates present at higher concentrations

in seminal plasma than in blood *(15, 21, 22)*, how these substrates are synthesized and

specifically how they alter sperm motility has not been determined. Moreover, though the

motility of ejaculated spermatozoa is known to decrease with aging, few reports have revealed

age-related changes in the seminal vesicle glands, which are responsible for seminal plasma

synthesis.

In this study, we focused on the reproductive organs that synthesize seminal vesicle secretion

as target organs for testosterone, and found that factors formed by testosterone-dependent

metabolic changes in seminal vesicles activate sperm mitochondria and enhance their linear

motility. Testosterone is known to activate intracellular ARs and alter intracellular metabolic

pathways in skeletal muscle *(33)*, prostate cells *(34)* and cardiomyocytes *(30)*. ARs are known to

directly regulate the expression of glycolytic and lipid metabolic genes *(30, 35, 36)*. In the

present study, the nuclear localization of AR in seminal vesicle epithelial cells (i.e., functioning

as a transcription factor) and the activation of the glycolytic system by testosterone were similar

to those in the above cell types. Interestingly, however, in the seminal vesicle epithelial cells,

expression of *Glut* family members was unchanged but GLUT4 localization was altered in a

testosterone-dependent manner to promote glucose uptake. Such gene expression-independent

effects of testosterone on glucose metabolism have also been reported in skeletal muscle, liver,

and adipose tissue, where testosterone leads to the phosphorylation of GLUT4 and enhances its

translocation to the plasma membrane *(37, 38)*. This is consistent with the results of our

metabolic flux analyses, that show testosterone increases the rate of glycolysis in seminal vesicle

epithelial cells by increasing glucose uptake rather than by increasing glycolytic capacity. We

further confirmed that GLUT4 is the transporter that mediates this glucose uptake using
indinavir, a GLUT4-specific inhibitor.

In seminal epithelial cells, ATP production was low and cell proliferation was suppressed
despite testosterone-induced activation of the glycolytic system. We hypothesized that this could
be explained by the testosterone-dependent regulation of genes involved in mitochondrial
metabolism, including the up-regulation of ACLY, a strategic enzyme that links the glycolytic
system with lipid metabolism. Manipulation of ACLY expression in mouse myoblasts and
embryonic stem cells has been shown to alter the TCA cycle into a "non-normal cycle" *(39)*. One
advantage of the non-normal TCA cycle is its ability to retain carbon instead of burning it and to
regenerate the cytosolic NAD+ needed to maintain glycolysis *(39, 40)*. This suggests that ACLY
activated by testosterone may inhibit proliferation of seminal epithelial cells by skipping some
steps of the mitochondrial TCA cycle and minimizing ATP production.

Testosterone has been reported to cause hypertrophy of cardiac myocytes *(41, 42)*. Cell
hypertrophy is partly due to the accumulation of fatty acids in the cytoplasm, which causes
significant damage to cellular functions *(43, 44)*. On the other hand, the normal cell hypertrophy,
especially in myofibers, involves the activation of signaling molecules such as Myc,
hypoxia-inducible factor, and Pi3k-Akt-mTor, which regulate metabolic reprogramming in
cancer by increasing glucose uptake in the presence of oxygen *(45–47)*. In the present study,
induction of ACLY expression by testosterone not only decreased cycling in the TCA cycle, but
also promoted glucose assimilation, which played an important role in fatty acid synthesis. Thus,
in tissues where fatty acid synthesis is promoted, either β-oxidation-dependent ATP production
in mitochondria or the secretion of fatty acids occurs to maintain cellular function *(48–50)*;
providing evidence to explain why the presence of fatty acids in the cell has no adverse effect. In
accessory reproductive glands, such as the prostate and seminal vesicles, the glycolytic system is
activated in a testosterone-dependent, but ATP production is not enhanced and cell proliferation
is not induced and the metabolites are used as a substrate for sperm motility. This may be
because in the prostate, pyruvate synthesized by the glycolytic system in response to testosterone
is converted to acetyl CoA, which is then converted to citric acid in the TCA cycle, citric acid is
released as the main component of seminal plasma *(34)*.

The testosterone-dependent growth of cancerous prostate epithelial cells may be due to
genetic mutations, such as elevated expression of the gene encoding aconitase, which converts
citric acid to isocitrate, which blocks the TCA cycle and inhibits citric acid secretion *(51)*. On the
other hand, in the analysis of seminal epithelial cells, testosterone treatment induced the
expression of *Acc*, which encodes the rate-limiting enzyme for fatty acid synthesis, and *Elovl6*,
which desaturates to oleic acid, significantly increasing total saturated fatty acid and oleic acid
content. Therefore, testosterone promotes glucose conversion into fatty acids, especially oleic
acid, in seminal epithelial cells. Although there is a commonality in the activation of the
glycolytic system testosterone target tissues, the different expression patterns of genes involved
in metabolism in mitochondria may exert their specific functions in each tissue. In addition to
genes involved in fatty acid synthesis such as *Fasn* and *Elovl6*, *Acly* is also encoded in a region
in close proximity in human chromosome 17 and mouse chromosome 11, suggesting that in
seminal vesicle epithelial cells, this region is targeted to promote gene expression. The
co-regulation of these genes in seminal vesicle epithelial cells leads to the synthesis of seminal
plasma components that ensure sperm motility.

Decreased testosterone synthesis in the testes is a phenomenon observed with aging, and
associated with decreased spermatogenesis*(52)*. However, the probability of obtaining sperm
without ICSI or IVF is extremely low, despite the presence of sufficient numbers of sperm with
normal-morphology *(53–55)*. This indicates that testosterone plays a major role not only in
spermatogenesis but also in functional maturation of sperm. In this study, metabolic
abnormalities induced by inhibition of androgen receptor function caused abnormal proliferation
of seminal vesicle epithelial cells in aged mice. In other words, the findings of the seminal
vesicle epithelial cell culture experiments and flutamide treatment may mimic the changes that
occur with aging. Oleic acid synthesis by testosterone-dependent metabolic pathways was
regulated in seminal vesicle epithelial cells and the oleic acid stimulated sperm motility and
enhanced *in vivo* fertilization success. Thus, results of these studies document that the quality
and presence of key metabolic products in seminal plasma influence male fertility. Specifically,
the results open the possibility of translational studies to understand human male infertility,
especially in cases of failure to conceive by natural mating or artificial insemination. Because
seminal plasma contains components specific to certain male reproductive organs, differences in

protein composition may reflect pathological processes in specific organs *(56)*. However,

although proteomic and metabolomic analyses of seminal plasma have been performed *(28, 56)*,

the functional changes in seminal plasma, especially the effects on *in vivo* fertilization potential,

have not been well understood. Differences in fatty acid concentrations, including oleic acid, in

seminal plasma may provide a predictive marker for the fertilization potential of human and

domestic animal semen. Furthermore, the addition of oleic acid to low-quality semen improved

fertilization rates associated with artificial insemination, providing a strategy for improving

artificial insemination. In our previous study, adding fatty acids to the thawing medium of frozen

bull sperm improved their linear motility and viability *(18)*. The similarities between mouse and

human seminal vesicle epithelial cells demonstrated in this study have strong potential for

clinical application of these observations.

Overall, the results of the studies described herein strongly support the hypothesis that

testosterone-dependent glucose to oleic acid conversion processes are required for normal

seminal plasma functions in the seminal vesicle and normal reproductive performance in males.

This study advances our understanding of the role of testosterone in glucose metabolism and

fatty acid synthesis and lays the groundwork for future research and potential therapeutic

interventions in reproductive biology and fertility.

MATERIALS AND METHODS

Study design

The objective of this study was to elucidate the mechanisms of sperm plasma synthesis that

enhance sperm fertility. Testosterone-dependent changes in the seminal plasma function were

examined by comparative analysis of adult mice and flutamide-treated mice. The bioassays of

extracts from male accessory glands by epididymal sperm motility analysis were used to examine

the seminal plasma components important for sperm fertilization. We also isolated the mouse

seminal vesicle epithelial cultured cells, and investigated the synthesis mechanism of the seminal

plasma components using RNA-seq, FluxAnalyzer, and shRNA knockdown. Finally, we

performed validation using the human epithelial cells.

Chemicals and Animals

Unless otherwise noted, chemicals used in this study were purchased from Sigma-Aldrich (St.

Louis, MO, USA) or Nacalai tesque (Osaka, Japan). Pregnant mare serum gonadotropin (PMSG)

and human chorionic gonadotropin (hCG) were purchased from Asuka Seiyaku (Tokyo, Japan).

All procedures during animal experiments were reviewed and approved by the Animal Care and

Use Committee of Hiroshima University (Hiroshima, Japan; C21-9-3) and conducted according

to regulations. Specific pathogen-free C57BL/6NCrl male mice or Crl: CD1 (ICR) female mice

were purchased from Jackson Laboratory Japan (Kanagawa, Japan) and housed in an

environmentally controlled room with a 12-hour light/dark cycle, a temperature of $23 \pm 3°C$, and

free access to laboratory food (MF; Oriental Yeast Co., Ltd., Tokyo, Japan) and tap water. 8- to

16-week-old male or female mice were used for the experiments. Flutamide-treated mice were

injected subcutaneously for 7 days with 50 µg/g (body weight) of flutamide dissolved in corn oil.

Vehicle-treated mice received 100 µL of corn oil subcutaneously for 7 days. The aged males were

12 to 16 months old.

Bioassay using pseudo-seminal plasma and epididymal sperm

Prostate (mixed anterior, dorsal, lateral, and ventral) and seminal vesicles were collected from

adult C57BL/6NCrl mice. Seminal vesicle secretions were obtained by pressing the seminal

vesicles immediately after euthanasia. Prostatic extracts were obtained by homogenizing the

prostate immediately after euthanasia. Seminal vesicle secretions and prostate extracts collected

from one male mouse were dissolved in 500 µL and 200 µL of mHTF medium without BSA *(57)*,

respectively, then centrifuged at $2000 \times g$ for 5 min at room temperature, and the supernatant was

collected, and protein levels were determined. The seminal vesicle secretions from

flutamide-treated mice were also collected similarly.

Cauda epididymis sperm recovered from male mouse were dispersed in 500 µL of mHTF

without BSA drops and centrifuged at $300 \times g$, 3 min to wash out epididymis-derived factors. The

epididymis sperm was mixed with seminal vesicle secretion and/or prostate extract as shown in

Fig.1A. In flutamide administration experiments, the final protein concentration of seminal

vesicle secretion was adjusted to 10 mg/mL. The samples were incubated at 37°C for 60 min in a

humidified atmosphere of 5% CO_2 in air. Sperm motility patterns were then examined using

computer-assisted sperm analysis (CASA; HT CASA-Ceros II, Hamilton Thorne, Beverly, MA,

USA).

Serum testosterone levels

The serum was separated from the clot by centrifugation at $1,500 \times g$ for 15 min. Testosterone

levels in the serum samples were determined by a rodent testosterone ELISA test kit (ERKR7016,

Endocrine Technologies, Inc., Newark, CA, USA) according to the manufacturer's manual.

Mitochondrial activity

Mitochondrial activity of sperm was measured using a MitoPT® JC-1 Assay Kit (911, Immuno

Chemistry Technologies, LLC, Bloomington, MN, USA) according to our previous study *(17)*.

Briefly, sperm were incubated with 200 µL of working solution containing

5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolyl carbocyanine iodide (JC-1) dye at 37 °C

for 30 min in the dark. The sperm suspension was centrifuged and washed twice with the base

medium. After washing, the sperm pellet was resuspended in the base medium and analyzed with

an Attune® NxT Acoustic Focusing Cytometer (Thermo Fisher Scientific Inc., Waltham, MA,

USA) using a 488 nm laser and a filter with a bandwidth of 574/26 nm. The intensity of the

average value was analyzed as the mean fluorescence intensity (MFI) of JC-1 orange aggregates.

A total of 100,000 sperm events were analyzed. The sperm population is shown in Supplementary

Figure 8.

Immunohistochemistry and immunofluorescence

Seminal vesicles were fixed overnight in 4% (w/v) paraformaldehyde/PBS, dehydrated in 70%

(v/v) ethanol, and embedded in paraffin. Paraffin-embedded fixed sections (4 µm) were

deparaffinized and hydrated with xylene and ethanol, respectively. Some slides were stained with

hematoxylin and eosin (HE). For antigen retrieval, sections were placed in citrate buffer (pH 6.0),

microwaved until boiling and allowed to stand at the temperature of pre-boil (95°C) for 15

minutes. Sections were incubated with 3% (v/v) hydrogen peroxide in methanol to block

endogenous peroxidase and then incubated with Normal Goat Serum Blocking Solution for rabbit

antibody (S-1000, Vector Laboratories, Newark, CA, USA) to block nonspecific sites. The

sections were then incubated overnight with primary rabbit antibodies: anti-AR antibody (1:100;

ab133273; Abcam, Cambridge, UK), anti-Ki67 antibody (1:1000; ab15580; Abcam) or

anti-ACLY antibody (1:200; ab40793; Abcam). Positive signals were visualized by DAB

(3,3′-Diaminobenzidine, 040-27001, Fujifilm Wako, Osaka, Japan) using the VECTASTAINE

LifeABC kit (PK-6101, Vector Laboratories) according to the manufacturer's protocol. Apoptotic

cells (including fragmented DNA) were detected using the In Situ Cell Death Detection Kit POD

(11684817910, Roche Ltd., Basel, Switzerland) according to manufacturer's protocol. Nuclei

were visualized with hematoxylin (Sakura Finetek, Tokyo, Japan). Ki-67 and apoptotic positive

cells were quantified using ImageJ software.

For immunofluorescence, after blocking nonspecific sites, the sections were incubated with a

primary antibody: anti-GLUT4 antibody (1:200; ab33780; Abcam). After washing with 0.3%

(v/v) Triton X-100 in PBS (-), Cy3-labeled goat anti-rabbit IgG (1:200, C-2306, Sigma) and

DAPI (VECTESHIELD Mounting Medium with DAPI, H-1200, Vector Laboratories) were used

to visualize the antigen and nuclei. Digital images were taken using an APX100 Digital Imaging

System (EVIDENT, Tokyo, Japan).

Isolation and culture of seminal vesicle epithelial cells

Seminal vesicles were removed from euthanized mice, cut open vertically to expose the

seminal vesicle cavity, and incubated with Hanks' balanced salt solution (HBSS; $CaCl_2 \cdot 2H_2O$

0.185 g/L, $MgSO_4$ 0.098 g/L, KCl 0.400 g/L, KH_2PO_4 0.060 g/L, NaHCO3 0.350 g/L, NaCl 8.000

g/L, Na_2HPO_4 0.048 g/L, D-Glucose 1.0 g/L, dissolved in ultrapure water). Tissues were digested

with 10 mL of HBSS containing 25 mg/mL pancreatin and 0.25% (v/v) trypsin EDTA for 1 hour

at 4°C, 45 minutes at room temperature, and 15 minutes at 37°C, and epithelial cells were

obtained. These were passed through a 70 µm nylon filter (352350, Corning Inc., Corning, NY,

USA) and seeded into collagen-coated 12-well plates (4815-010, IWAKI, Shizuoka, Japan). Cells

were cultured in DMEM/F-12, 10% fetal bovine serum (FBS; 10438-018, Thermo Fisher

Scientific Inc.), 100 U/mL penicillin, 100 µg/mL streptomycin. The culture condition was 37°C,

5% CO_2, 95% air, and humidity.

Human seminal vesicle epithelial cells (4460, ScienCell Research Laboratories, Carlsbad, CA,

USA) were purchased, seeded in 2 $\mu g/cm^2$ poly-L-lysine-coated plates according to

manufacturer's protocol, and cultured in epithelial cell medium (4101, ScienCell Research

Laboratories). The cells were seeded at 10000 cells/cm^2 on collagen-coated plates the day before

100 ng/mL testosterone treatment. Testosterone was dissolved in ethanol, and the final

concentration of ethanol was adjusted to 0.1% (v/v) when used for cultured cells. The culture

medium was replaced every 48 hours.

Cell proliferation curve

The seminal vesicle epithelial cells were seeded at 10000 cells/cm^2 and then were cultured in

various concentrations of testosterone. The cells were washed with HBSS (without

CaCl$_2$-2H$_2$O/MgSO$_4$) at the time of cell counting and collected using 0.25% (v/v) trypsin. The

recovered cells were suspended in DMEM/F-12 medium, and cell numbers were determined

using an Automated Cell Counter TC20TM (BIO-RAD Laboratories, Hercules, CA, USA) to

calculate the number of cells contained per well (4 cm^2).

Cell cycle analysis

The recovered cells as described above were fixed in 2-5 ml of cold (4°C) 70% methanol for at

least 30 minutes on ice. After centrifugation at 500g for 5 minutes at 4°C to obtain pellets, the cell

pellet was suspended in 50 µM propidium iodide and 20 × diluted RNase (19101, QIAGEN Inc.,

Venlo, Netherlands) and then were incubated at 37°C for 30 minutes. After twice washing with 1

ml of PBS at 300 g for 3 min at 4°C, the cells were analyzed by flow cytometry. The cell

population is shown in Supplementary Figure 9. A total of 20,000 events were analyzed.

RNA-sequencing

Total RNA was extracted from cells using the RNeasy Mini Kit (74106, QIAGEN) according

to the manufacturer's protocol. Three independent samples were assayed to evaluate the

reproducibility of the experimental procedures. The quality of total RNA (1 µg) submitted for

sequencing was checked using a Bioanalyzer (Agilent Technologies Inc., Santa Clara, CA, USA),

and all RNA integrity numbers were >7.0. The libraries for RNA sequencing were sequenced with

2 × 150-bp paired-end reads on a NovaSeq6000 (Illumina Inc., San Diego, CA, USA). The

obtained sequencing data were analyzed using RaNA-seq *(58)*, an open bioinformatics tool for

rapid RNA-seq data analysis. FASTQ files were preprocessed in the pipeline using the Fastp tool,

and expression was quantified using Salmon. DEseq2 was used to identify differentially

expressed genes (DEGs) with adjusted p-values less than 0.05 between the testosterone-treated

groups in transcripts per million (TPM) between control and testosterone-treated cells. The

pathways were identified from the KEGG database on RaNA-seq.

Extracellular flux analysis

Oxygen consumption rate (OCR) and extracellular oxidation rate (ECAR) were measured

using an extracellular flux analyzer (XF HS Mini; Agilent Technologies). Eight-well plates for

the XF HS Mini were coated with fibronectin (5 µg/mL, F1141, Sigma-Aldrich) dissolved in

phosphate-buffered saline (PBS), incubated overnight, and then washed with Roswell Park

Memorial Laboratory medium for XF (Roswell Park Memorial Institute medium; RPMI,

103576-100, Agilent Technologies; containing 1% FBS) for XF before analysis. Mouse seminal

vesicle epithelial cells or human seminal vesicle epithelial cells treated with 100 ng/mL

testosterone for 7 days were suspended in RPMI medium at a concentration of 10,000 cells/180

µL, or mouse seminal vesicle epithelial cells infected with shRNA were suspended at a

concentration of 2000 cells/180 µL in each well. After centrifugation at $300 \times g$ for 3 minutes, the

plates were placed in an incubator at 37°C and 100% air and used for analysis within 1 hour.

For glucose metabolism assessment, cells were sequentially treated as indicated with

D-Glucose (10 mM; 103577-100, Agilent Technologies), Oligomycin (1 µM; O4876,

Sigma-Aldrich), and 2-Deoxy-Glucose (50 mM; D8375, Sigma-Aldrich).

For mitochondrial respiration measurement, cells were sequentially treated as indicated with

Oligomycin (1 µM), carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP; 5 µM,

SML2959, Sigma-Aldrich), and Rotenone (1 µM; R8875, Sigma-Aldrich)/Antimycin A (1 µM;

A8674, Sigma-Aldrich). Cellular metabolism readouts, such as glycolysis, glycolytic capacity,

glycolytic reserve, basal OCR, Maximum respiration, ATP Production, Spare respiratory capacity,

were determined using the Agilent Seahorse Wave software.

For mouse sperm analysis, NaHCO3-free HTF medium was prepared according to Balbach et

al. *(59)*. Plates were coated with concanavalin A (0.5 mg/mL, Fujifilm Wako Chemicals)

overnight the day before assay. Each well was seeded with 180 µL of NaHCO3- Free HTF

medium containing 3,000,000 sperm cells. Sperm were injected with oleic acid (final

concentration: 10 nM) with or without etomoxir (final concentration: 40 nM, E1905,

Sigma-Aldrich).

Measurement of pyruvate and lactic acid

For measurements, mouse or human seminal vesicle epithelial cells were treated with 100

ng/ml testosterone for 6 days. The culture supernatant was collected after 24 hr of additional

culture. Pyruvate consumption was measured using the Pyruvate Assay kit (MET-5029, Cell

Biolabs, Inc., San Diego, CA, USA) according to the manufacturer's protocol. The fluorescence

intensity was measured using a microplate reader (ex. 570 nm/em. 615 nm). Lactate released into

the medium was measured using the Lactate Assay kit (L256, DOJINDO LABOLATORIES Co.,

Ltd., Kumamoto, Japan) according to the manufacturer's protocol. The absorbance was measured

using a microplate reader (wave length=450nm).

qPCR

The cDNA was synthesized from 40 ng of total RNA using oligo(dT)$_{15}$ primers (3805, Takara

Bio Inc., Shiga, Japan) and an Avian myeloblastosis virus (AMV) reverse transcriptase from

Promega (M5101, Madison, WI, USA). The cDNA and primers were added to Power SYBR

Green PCR Master Mix (4367659, Applied Biosystems, Foster City, CA, USA) in a total reaction

volume of 15 μl. Conditions were set to the following parameters: 10 min at 95°C, 15 sec at 95°C,

and 1 min at 60°C. Cycles were repeated 45 times. Specific primer pairs were selected and

analyzed, as shown in Table S1. Expression was first normalized to housekeeping gene *Rpl19*,

and fold change was calculated relative to the mean of the control samples.

Glucose uptake test of seminal vesicle epithelial cells and lipid determination of culture supernatants

Mouse or human seminal vesicle epithelial cells treated with or without 100 ng/mL testosterone

for 6 days were used for this experiment. Cells were treated with or without the GLUT4 inhibitor

indinavir (100 nM) and 100 ng/mL testosterone for an additional 24 hours. Collected 15,000 cells

were subjected to Glucose Uptake Assay Kit-Green (UP02; DOJINDO), and fluorescence
intensity was measured using a microplate reader (ex. 485 nm/em. 535 nm).

Total lipid content in cultured medium was quantified using a lipid quantification kit (STA-617
Fluorometric, Cell Biolabs, Inc.) according to the manufacturer's protocol. The fluorescence
intensity was measured using a microplate reader (ex. 485 nm/em. 535 nm).

Gas chromatography-mass spectrometry analysis

Fatty acid measurements in culture supernatant and seminal vesicle secretions were done
according to our previous work *(18)*. Fatty acids in the samples were extracted and methylated
using a fatty acid methylation kit (06482-04, Nacalai Tesque) and then purified using a fatty acid
methyl ester purification kit (06483-94, Nacalai Tesque) according to the manufacturer's protocol.
The methylated sample (3.0 mL) was dried and dissolved in 50 µL of elution solution. Samples
were subjected to GC-MS.

Fatty acids were determined on an Agilent 7890A GC system coupled with a JMS-T 100 GCv
mass detector at the Natural Science Research Center, Hiroshima University (N-BARD). Sample
(1.0 µL aliquot) was injected to SP$^{®\square}$-2560 capillary GC column (100 m × 0.25 mm i.d. × 0.2 µm
film thickness; 24056, Sigma-Aldrich). For measurements, helium was used as the carrier gas at a
constant flow rate of 2.0 mL/min. The GC oven temperature was gradually increased from 100°C
to 300°C at a rate of 10°C/min and held at 300°C for 20 minutes. Ionization was performed using
electron ionization with an electron energy of 70 eV. Mass spectra were obtained in scan mode
(mass scan range 29-800 m/z). NIST MS Search v.2.0 was used to detect and identify of fatty
acids.

Quantification of sperm oleic acid uptake by flow cytometry

To detect the fluorescence intensity of each sperm sample using flow cytometry, sperm
collected from the epididymis were incubated with 1, 10 or 100 µM of fluorescently labeled oleic
acid (F-OA-01, NANOCS, New York, NY, USA). After washing, fluorescence intensities were
analyzed by flow cytometry using a 530/30 nm bandwidth filter and measured as mean intensity.
A total of 100,000 sperm events were analyzed. The localization of fluorescence in sperm was
determined using an APX100 Digital Imaging System.

ACLY knock down experiment using shRNA

    The U6 shRNA lentiviral vector of the *Acly* gene (mAcly[shRNA#1]; a virus packaging service,
VectorBuilder, Chicago, IL, USA) was used for knockdown experiments. The ACLY sequence
targeted by the shRNA vector used in this study was 5′-GCAGCAAAGATGTTCAGTAAA-3′.
Lentivirally transduced cells were prepared according to manufacture protocol. Immediately after
isolation, the mouse seminal vesicle epithelial cells were cultured overnight in a medium
containing 5 µg/mL polybrene (VectorBuilder) and lentiviral particles expressing an shRNA
targeting ACLY or a scramble (Multiplicity of Infection of 10). After an additional overnight
incubation in a fresh culture medium, the cells were incubated for 5 days in a medium containing
1 µg/ml puromycin (19752-64, Nacalai tesque). 100 ng/mL of testosterone was added to all
procedures for the testosterone-treated group. Infection efficiency, calculated based on the
number of GFP-positive cells using an APX100 Digital Imaging System, ranged between 70%
and 90%.

Western blot analysis

    Protein samples of mouse seminal vesicle epithelial cells and seminal vesicle tissue were
prepared by homogenization in cell lysis buffer (04719964001, cOmplete™ Lysis-M EDTA-free,
Roche) and diluted with an equal volume of sample buffer solution containing 2-ME (2 ×) for
SDS-PAGE (30566-22, Nacalai tesque). The extracts (10 µg protein) were separated by
SDS-polyacrylamide gel (10%) electrophoresis and transferred to polyvinylidene fluoride
membranes (10600069, Cytiva, Tokyo, Japan). The membranes were then treated with
Tris-buffered saline and Tween 20 (TBST, 20 mM Tris, pH 7.5, 150 mM NaCl, 0.1% Tween 20)
containing 5% (w/v) Non-Fat Dry milk (MORINAGA MILK INDUSTRY Co., Ltd., Tokyo,
Japan) for the nonspecific reaction was blocked. The membranes were incubated overnight at 4°C
with primary antibodies: anti-ACLY antibody (1:10,000; ab40793; Abcam) or anti-α/β-Tubulin
antibody (1:1,000; 2148S; Cell Signaling). The next day, membranes were washed with TBST
and incubated with HRP-labeled antibody specific for rabbit IgG (1:4000; 7074S; Cell Signaling).
The bands were visualized by using Enhanched Chemiluminescence detection systems

655 (RPN2232, Cytiva) and ChemiDoc^{TM} MP Imaging System (BIO-RAD Laboratories). The bands

656 were analyzed by using the software ImageJ (National Institutes of Health; Bethesda, MD, USA).

ATP production in sperm

After 1 hour of incubation in the presence or absence of oleic acid (10 nM), ATP in sperm was

visualized by BioTracker ATP-Red (SCT045, Sigma-Aldrich). The sperm were incubated with 10

nM of working solution containing BioTracker ATP-Red dye at 37 °C for 60 min in the dark.

After washing, the sperm pellet was resuspended in the base medium and analyzed with the flow

cytometry using a 488 nm laser and a filter with a bandwidth of 574/26 nm. A total of 100,000

sperm events were analyzed. The sperm population is shown in Supplementary Figure 6.

Artificial insemination

To examine the involvement of seminal vesicle secretions in fertilization, sperm from

C57BL/6N mice were injected directly into the uterine bodies of superovulated CD1 mice by

syringe, as previously described with minor modifications *(14, 19)*. CD1 female mice were

injected intraperitoneally with 8 IU eCG to stimulate follicle development and used as recipients

for artificial insemination 48 hours later. Artificial seminal fluid was prepared in HTF medium to

a final concentration of 10 mg/mL of collected control or frutamide-derived seminal vesicle

secretions. As rescue experiment, 10 nM of oleic acid was added to the frutamide-derived seminal

secretions. Cauda epididymis sperm was suspended with above artificial seminal fluid (1×10^6

sperm/mL), and a total of 40 µL (20 µL each per uterus) was injected through the cervix of the

female mice. After artificial insemination, 7.5 IU of hCG was injected to stimulate ovulation.

Subsequently, a mixture of vaseline and paraffin with a ratio of 10:1 was injected to form an

artificial plug around the cervix. After 36~48 hours of sperm injection, the putative eggs were

retrieved from the oviducts. The number of two-cell embryos and mature unfertilized eggs were

counted and used to estimate the fertilization success rate.

Statistical analysis

683   Quantitative data were presented as means ± SEM. All statistical details of the experiments are

684 given in the figure legends. R [(version 4.3.1 (2023-06-16)] was used for statistical analysis .

Data availability

Nucleotide sequence data reported are available in the DDBJ Sequenced Read Archive under

the accession numbers DRA017090. Additional information needed to reanalyze the data

reported in this paper is available from the principal contact upon request.

List of Supplementary Materials

Supplemental Figure 1 to 9

Table S1

Supplementary Data

Acknowledgments:

The authors would like to thank JoAnne S. Richards for carefully proofreading the manuscript

and useful comments.

Funding:

This work was supported in part by JSPS KAKENHI Grant Number 23H00361 (to MS).

Author contributions:

Conceptualization: TY, TU, MS

Methodology: TY, ZX, NT, MA, TU

Investigation: TY, ZX, NT, MA, TU

Visualization: TY, MS

Funding acquisition: MS

Supervision: MS

Writing – original draft: TY

Writing – review & editing: TY, TU, MS

Competing interests:

MS holds stocks in and receives a salary from Hiroshima Cryopreservation Service Co. as a director. In addition, MS has received royalties and grants from Hiroshima Cryopreservation Service Co. M.S. has received consulting fees from Rohto Pharmaceutical Co., Ltd.. TU holds stocks in and receives a salary from Lullabio Inc. as a director and has received honoraria from Rohto Pharmaceutical Co., Ltd. The other authors declare they have no competing interests.

References

1. World Health Organization., Infertility prevalence estimates: 1990–2021. *Global report* , 1–98 (2023).

2. S. Elbashir, Y. Magdi, A. Rashed, R. Henkel, A. Agarwal, Epididymal contribution to male infertility: An overlooked problem. *Andrologia* **53**, e13721 (2021).

3. M. J. Cosentino, A. T. K. Cockett, Review Article: Structure and Function of the Epididymis Transport of Spermatozoa. *Urol Res* **14**, 229–240 (1986).

4. B. Howarth, Fertilizing ability of cock spermatozoa from the testis epididymis and vas deferens following intramagnal insemination. *Biol Reprod* **28**, 586–590 (1983).

5. S. Elbashir, Y. Magdi, A. Rashed, R. Henkel, A. Agarwal, Epididymal contribution to male infertility: An overlooked problem. *Andrologia* **53**, e13721 (2021).

6. T. G. Cooper, Sperm maturation in the epididymis: A new look at an old problem. *Asian J Androl* **9**, 533–539 (2007).

7. P. Gaddum-Rosse, Some observations on sperm transport through the uterotubal junction of the rat. *Am J Anat* **160**, 333–341 (1981).

8. R. Shalgi, T. T. Smith, R. Yanagimachi2, A Quantitative Comparison of the Passage of Capacitated and Uncapacitated Hamster Spermatozoa through the Uterotubal Junction. *Biol Reprod* **46**, 419–424 (1992).

9. H. Li, P. Hung, S. S. Suarez, Ejaculated Mouse Sperm Enter Cumulus-Oocyte Complexes More Efficiently In Vitro than Epididymal Sperm. (2015), doi:10.1371/journal.pone.0127753.

10. T. Noda, M. Ikawa, Physiological function of seminal vesicle secretions on male fecundity. *Reprod Med Biol* **18**, 241–246 (2019).

11. P. Verze, T. Cai, S. Lorenzetti, The role of the prostate in male fertility, health and disease. *Nat*

Rev Urol **13**, 379–386 (2016).

12. E. Balandya, W. Wieland-Alter, K. Sanders, T. Lahey, Human Seminal Plasma Fosters CD4+

Regulatory T-cell Phenotype and Transforming Growth Factor-β1 Expression. *American Journal*

of Reproductive Immunology **68**, 322–330 (2012).

13. T. Shima, K. Inada, A. Nakashima, A. Ushijima, M. Ito, O. Yoshino, S. Saito, Paternal

antigen-specific proliferating regulatory T cells are increased in uterine-draining lymph nodes

just before implantation and in pregnant uterus just after implantation by seminal plasma-priming

in allogeneic mouse pregnancy. *J Reprod Immunol* **108**, 72–82 (2015).

14. N. Kawano, N. Araki, K. Yoshida, T. Hibino, N. Ohnami, M. Makino, S. Kanai, H. Hasuwa,

M. Yoshida, K. Miyado, A. Umezawa, Seminal vesicle protein SVS2 is required for sperm

survival in the uterus. *Proc Natl Acad Sci U S A* **111**, 4145–4150 (2014).

15. R. R. ROSECRANS, R. S. JEYENDRAN, M. PEREZ-PELAEZ, W. P. KENNEDY,

Comparison of Biochemical Parameters of Human Blood Serum and Seminal Plasma.

Andrologia **19**, 625–628 (1987).

16. T. Umehara, T. Kawai, M. Goto, J. A. S. Richards, M. Shimada, Creatine enhances the

duration of sperm capacitation: a novel factor for improving in vitro fertilization with small

numbers of sperm. *Human Reproduction* **33**, 1117–1129 (2018).

17. Z. Zhu, T. Kawai, T. Umehara, S. A. M. Hoque, W. Zeng, M. Shimada, Negative effects of

ROS generated during linear sperm motility on gene expression and ATP generation in boar

sperm mitochondria. *Free Radic Biol Med* **141**, 159–171 (2019).

18. M. M. Islam, T. Umehara, N. Tsujita, M. Shimada, Saturated fatty acids accelerate linear

motility through mitochondrial ATP production in bull sperm. *Reprod Med Biol* **20**, 289–298

(2021).

19. T. Noda, Y. Fujihara, T. Matsumura, S. Oura, S. Kobayashi, M. Ikawa, Seminal vesicle

secretory protein 7, PATE4, is not required for sperm function but for copulatory plug formation

to ensure fecundity. *Biol Reprod* **100**, 1035–1045 (2019).

20. K. Queen, C. B. Dhabuwala, C. G. Pierrepoint, The effect of the removal of the various

accessory sex glands on the fertility of male rats. *J Reprod Fertil* **62**, 423–426 (1981).

21. Mann T, Studies on the metabolism of semen; fructose as a normal constituent of seminal

plasma; site of formation and function of fructose in semen. *Biochem J* **40**, 481–491 (1946).

22. Humphrey GF, Mann T, Studies on the metabolism of semen; citric acid in semen. *Biochem J* **44**, 97–105 (1949).

23. C. Chang, S. O. Lee, R. S. Wang, S. Yeh, T. M. Chang, Androgen receptor (AR) physiological roles in male and female reproductive systems: lessons learned from AR-knockout mice lacking AR in selective cells. *Biol Reprod* **89**, 1–16 (2013).

24. C. Fix, C. Jordan, P. Cano, W. H. Walker, Testosterone activates mitogen-activated protein kinase and the cAMP response element binding protein transcription factor in Sertoli cells. *Proc Natl Acad Sci U S A* **101**, 10919–10924 (2004).

25. K. Nagaosa, A. Kishimoto, R. Kizu, A. Nakagawa, A. Shiratsuchi, Y. Nakanishi, Perturbation of spermatogenesis by androgen antagonists directly injected into seminiferous tubules of live mice. *Reproduction* **133**, 21–27 (2007).

26. L. B. Smith, W. H. Walker, The regulation of spermatogenesis by androgens. *Semin Cell Dev Biol* **30**, 2–13 (2014).

27. P. S. Cooke, W. H. Walker, Male fertility in mice requires classical and nonclassical androgen signaling. *Cell Rep* **36**, 109557 (2021).

28. S. P. Smyth, B. Nixon, A. L. Anderson, H. C. Murray, J. H. Martin, L. A. MacDougall, S. A. Robertson, D. A. Skerrett-Byrne, J. E. Schjenken, Elucidation of the protein composition of mouse seminal vesicle fluid. *Proteomics* **22**, e2100227 (2022).

29. H. Murata, P. W. Hruz, M. Mueckler, Indinavir inhibits the glucose transporter isoform Glut4 at physiologic concentrations. *AIDS* **16**, 859–863 (2002).

30. C. Wilson, A. Contreras-Ferrat, N. Venegas, C. Osorio-Fuentealba, M. Pávez, K. Montoya, J. Durán, R. Maass, S. Lavandero, M. Estrada, Testosterone increases GLUT4-dependent glucose uptake in cardiomyocytes. *J Cell Physiol* **228**, 2399–2407 (2013).

31. G. F. Gonzales, Function of seminal vesicles and their role on male fertility. *Asian J Androl* **3**, 251–258 (2001).

32. A. Lenzi, M. Picardo, L. Gandini, F. Dondero, Lipids of the sperm plasma membrane: from polyunsaturated fatty acids considered as markers of sperm function to possible scavenger therapy. *Hum Reprod Update* **2**, 246–256 (1996).

33. M. T. Haren, A. M. Siddiqui, H. J. Armbrecht, R. T. Kevorkian, M. J. Kim, M. J. Haas, A.

Mazza, V. B. Kumar, M. Green, W. A. Banks, J. E. Morley, Testosterone modulates gene

expression pathways regulating nutrient accumulation, glucose metabolism and protein turnover

in mouse skeletal muscle. *Int J Androl* **34**, 55–68 (2011).

34. L. C. Costello, R. B. Franklin, Testosterone and prolactin regulation of metabolic genes and

citrate metabolism of prostate epithelial cells. *Horm Metab Res* **34**, 417–424 (2002).

35. K. Gonthier, R. T. K. Poluri, É. Audet-Walsh, Functional genomic studies reveal the androgen

receptor as a master regulator of cellular energy metabolism in prostate cancer. *J Steroid Biochem*

Mol Biol **191**, 105367 (2019).

36. M. F. Troncoso, M. Pavez, C. Wilson, D. Lagos, J. Duran, S. Ramos, G. Barrientos, P. Silva, P.

Llanos, C. Basualto-Alarcón, B. D. Westenbrink, S. Lavandero, M. Estrada, Testosterone

activates glucose metabolism through AMPK and androgen signaling in cardiomyocyte

hypertrophy. *Biol Res* **54**, 1–16 (2021).

37. Chen X, Li X, Huang HY, Li X, Lin JF, Effects of testosterone on insulin receptor substrate-1

and glucose transporter 4 expression in cells sensitive to insulin. *Zhonghua Yi Xue Za Zhi* **86**,

1474–1477 (2006).

38. K. Sato, M. Iemitsu, K. Aizawa, R. Ajisaka, Testosterone and DHEA activate the glucose

metabolism-related signaling pathway in skeletal muscle. *Am J Physiol Endocrinol Metab* **294**,

E961–E968 (2008).

39. P. K. Arnold, B. T. Jackson, K. I. Paras, J. S. Brunner, M. L. Hart, O. J. Newsom, S. P.

Alibeckoff, J. Endress, E. Drill, L. B. Sullivan, L. W. S. Finley, A non-canonical tricarboxylic acid

cycle underlies cellular identity. *Nature* **603**, 477 (2022).

40. A. Luengo, Z. Li, D. Y. Gui, L. B. Sullivan, M. Zagorulya, B. T. Do, R. Ferreira, A. Naamati,

A. Ali, C. A. Lewis, C. J. Thomas, S. Spranger, N. J. Matheson, M. G. Vander Heiden, Increased

demand for NAD+ relative to ATP drives aerobic glycolysis. *Mol Cell* **81**, 691–707 (2021).

41. B. B. Yeap, Testosterone and cardiovascular disease risk. *Curr Opin Endocrinol Diabetes*

Obes **22**, 193–202 (2015).

42. J. D. Marsh, M. H. Lehmann, R. H. Ritchie, J. K. Gwathmey, G. E. Green, R. J. Schiebinger,

Androgen receptors mediate hypertrophy in cardiac myocytes. *Circulation* **98**, 256–261 (1998).

43. G. F. Grabner, H. Xie, M. Schweiger, R. Zechner, Lipolysis: cellular mechanisms for lipid

mobilization from fat stores. *Nat Metab* **3**, 1445–1465 (2021).

44. B. Mittendorfer, Origins of metabolic complications in obesity: adipose tissue and free fatty

acid trafficking. *Curr Opin Clin Nutr Metab Care* **14**, 535–541 (2011).

45. H. Wackerhage, I. J. Vechetti, P. Baumert, S. Gehlert, L. Becker, R. T. Jaspers, M. H. de

Angelis, Does a Hypertrophying Muscle Fibre Reprogramme its Metabolism Similar to a Cancer

Cell? *Sports Med* **52**, 2569–2578 (2022).

46. R. J. De Berardinis, N. S. Chandel, Fundamentals of cancer metabolism. *Sci Adv* **2**, e1600200

(2016).

47. G. L. Semenza, Hypoxia-inducible factors in physiology and medicine. *Cell* **148**, 399–408

(2012).

48. B. Morant-Ferrando, D. Jimenez-Blasco, P. Alonso-Batan, J. Agulla, R. Lapresa, D.

Garcia-Rodriguez, S. Yunta-Sanchez, I. Lopez-Fabuel, E. Fernandez, P. Carmeliet, A. Almeida,

M. Garcia-Macia, J. P. Bolaños, Fatty acid oxidation organizes mitochondrial supercomplexes to

sustain astrocytic ROS and cognition. *Nature Metabolism 2023 5:8* **5**, 1290–1302 (2023).

49. M. Duta-Mare, V. Sachdev, C. Leopold, D. Kolb, N. Vujic, M. Korbelius, D. C. Hofer, W. Xia,

K. Huber, M. Auer, B. Gottschalk, C. Magnes, W. F. Graier, A. Prokesch, B. Radovic, J. G.

Bogner-Strauss, D. Kratky, Lysosomal acid lipase regulates fatty acid channeling in brown

adipose tissue to maintain thermogenesis. *Biochim Biophys Acta Mol Cell Biol Lipids* **1863**,

467–478 (2018).

50. S. C. C. Huang, B. Everts, Y. Ivanova, D. O'Sullivan, M. Nascimento, A. M. Smith, W. Beatty,

L. Love-Gregory, W. Y. Lam, C. M. O'Neill, C. Yan, H. Du, N. A. Abumrad, J. F. Urban, M. N.

Artyomov, E. L. Pearce, E. J. Pearce, Cell-intrinsic lysosomal lipolysis is essential for alternative

activation of macrophages. *Nat Immunol* **15**, 846–855 (2014).

51. K. H. Tsui, T. H. Feng, Y. F. Lin, P. L. Chang, H. H. Juang, p53 downregulates the gene

expression of mitochondrial aconitase in human prostate carcinoma cells. *Prostate* **71**, 62–70

(2011).

52. S. Almeida, L. Rato, M. Sousa, M. G. Alves, P. F. Oliveira, Fertility and Sperm Quality in the

Aging Male. *Curr Pharm Des* **23**, 4429–4437 (2017).

53. T. Murata, Y. Ishitsuka, K. Karouji, H. Kaneda, H. Toki, Y. Nakai, S. Makino, R. Fukumura, H.

Kotaki, S. Wakana, T. Noda, Y. Gondo, β-cateninC429S mice exhibit sterility consequent to

spatiotemporally sustained Wnt signalling in the internal genitalia. *Scientific Reports 2014 4:1* **4**,

1–7 (2014).

54. B. A. Stone, A. Alex, L. B. Werlin, R. P. Marrs, Age thresholds for changes in semen

parameters in men. *Fertil Steril* **100**, 952–958 (2013).

55. J. L. Frattarelli, K. A. Miller, B. T. Miller, K. Elkind-Hirsch, R. T. Scott, Male age negatively

impacts embryo development and reproductive outcome in donor oocyte assisted reproductive

technology cycles. *Fertil Steril* **90**, 97–103 (2008).

56. A. P. Drabovich, P. Saraon, K. Jarvi, E. P. Diamandis, Seminal plasma as a diagnostic fluid for

male reproductive system disorders. *Nat Rev Urol* **11**, 278–288 (2014).

57. T. Umehara, N. Tsujita, Z. Zhu, M. Ikedo, M. Shimada, A simple sperm-sexing method that

activates TLR7/8 on X sperm for the efficient production of sexed mouse or cattle embryos. *Nat*

Protoc **15**, 2645–2667 (2020).

58. C. Prieto, D. Barrios, RaNA-Seq: Interactive RNA-Seq analysis from FASTQ files to

functional analysis. *Bioinformatics* **36**, 1955–1956 (2019).

59. M. Balbach, J. Buck, L. R. Levin, Using an Extracellular Flux Analyzer to Measure Changes

in Glycolysis and Oxidative Phosphorylation during Mouse Sperm Capacitation. *J Vis Exp* **2020**,

e60815 (2020).

Figures and Figure legends

Figure 1. Effects of seminal vesicle fluid and prostate fluids on sperm motility (A) To

determine the effect of seminal plasma on sperm motility, we prepared "pseudo" seminal plasma

from the mixture of seminal vesicle secretions or prostate extracts. (B-D) Tested various

concentrations of these fluids on epididymal sperm: (B) Curvi-Linear Velocity; VCL, (C)

Straight-Line Velocity; VSL and (D) Linearity; LIN is the ratio of VSL to VCL of sperm that was

incubated with the mixture of seminal vesicle secretions and/or prostate extracts. (E)

Experimental design to evaluate the effect of seminal secretions collected from male mice treated

with or without flutamide (50 mg/kg subcutaneously for 7 days) on sperm. (F-H) Performed

bioassay using seminal vesicle fluid after treatment with vehicle (Ctrl) or flutamide: (F) LIN, (G)

VSL and (H) the activity of mitochondria were checked by JC-1 kit. Data are mean ± SEM. n=3-5

independent replicates. Significance was tested in comparison to HTF or Ctrl (vehicle-treated

animals) using Student's t-test.

Figure 2. The testosterone-androgen receptor pathway inhibits the cell proliferation of

seminal vesicle gland epithelial cells. (A) The localization of the androgen receptor (AR;

NR3C4) in seminal vesicle of control (Ctrl) mice and those treated with flutamide (Fult). Scale

bar=50 µm for left panel and 20 µm for right panel in each group. (B,C) Cell proliferation and

apoptosis in seminal vesicle of control (Ctrl) mice and those treated with flutamide (Fult): (B)

Representative images of Ctrl and Flut staining for Ki67 (top) and TUNEL (bottom). Scale

bar=20 µm. (C) Percentage of Ki67 positive cells (top) and TUNEL positive (bottom) in Ctrl and

Flut-treated seminal vesicle sections. (D) Experimental design to evaluate the effects of

testosterone on the proliferation of seminal vesicle epithelial cells *in vitro*. The epithelial cells

were culutured with or without testosterone for 8 days. (E) Growth curves of seminal vesicle

epithelial cells that cultured with 0, 1, 10 and 100 ng/mL of testosterone. (F,G) Cell cycle status

was determined by flow cytometry. Data are mean ± SEM. n=3-5 mice or independent replicates.

Each replicate experiments with 3-6 wells containing pooled cells from 3-5 mice. Significance

was tested in comparison to Ctrl using Student's t-test. The cell proliferation data were compared

to Ctrl by Dunnett's test.

Figure 3. Testosterone changes the expression of genes involved in metabolic pathway in seminal vesicle epithelial cells. (A) Volcano plot of differentially expressed genes. RNA sequencing was performed using RNA extracted from the seminal vesicle epithelial cells cultured with or without 100 ng/mL testosterone. Genes with a significant expression change are highlighted as red dots. It found 4460 significant genes for a cutoff of P<0.05. (B) MA plot of differential expression results. The seminal vesicle specific genes are highlighted as red dots rather than significant genes. (C) Top 10 genes upregulated or downregulated by testosterone, respectively, for altered gene expression. (D) Conducted KEGG analysis to identify differences in pathway enrichment by testosterone, identifying the most variability in metabolic pathway genes. It shows the number of differential expressed genes annotated to Gene Ontology shows (x axis). p-value which measure the statistical significance of a possible functional enrichment for each term.

Figure 4. Testosterone regulates glucose metabolism and mitochondrial ATP production in seminal vesicle epithelial cells. (A-C) Glycolytic pathway analysis by an extracellular flux analyzer in seminal vesicle gland epithelial cells cultured with 100 ng/mL testosterone (Testo) or vehicle (Ctrl) for 7 days: (A) Extracellular acidification rate (ECAR) kinetics of seminal vesicle epithelial cells using an extracellular flux analyzer. (B) Glycolysis and (C) Glycolytic capacity. (D) Pyruvate concentration in the medium where seminal vesicle epithelial cells were cultured with or without testosterone for 24 hours. (E) Lactate concentration in the medium where seminal vesicle epithelial cells were cultured with or without testosterone for 24 hours. (F-J) Mitochondrial respiration measurement by an extracellular flux analyzer: (F) Oxygen consumption rate (OCR) kinetics of seminal vesicle epithelial cells with or without 100 ng/mL testosterone. (G) Basal OCR, (H) Maximum respiration, (I) Spare respiratory capacity and (J) ATP production. Data are mean ± SEM of n = 3 replicate experiments with 3-6 wells containing pooled cells from 3-5 mice or medium. Student's t-test was used to compare Ctrl and Testo.

Figure 5. Effect of testosterone on gene expression of enzymes involved in the glucose metabolic pathway in seminal vesicle epithelial cells. To elucidate the effects of testosterone on

gene expression of enzymes involved in glucose catabolism and anabolism in seminal vesicle

gland epithelial cells. ctrl: 7 days of culture with vehicle. testo: 7 days of culture with 100 ng/mL

testosterone. Data are mean ± SEM of n = 3 replicate experiments with 3-6 wells containing

pooled cells from 3-5 mice. Student's t-test was used to compare Ctrl and Testo.

Figure 6. Oleic acid is incorporated to sperm, which enhances linear motility and

mitochondrial activity. (A) Epididymal sperm were incubated with fluorescently labeled oleic

acid (OA) and the fluorescence intensity after quenching was observed by a fluorescence

microscope. (B) The fluorescence intensity of fluorescence-conjugated OA in sperm was detected

by flow cytometry. (C) Average fluorescence intensity of sperm at different concentrations of

fluorescence-conjugated OA. (D,E) OA improved sperm motility parameters. Sperm collected

from mouse epididymis was incubated with OA for 60 min: (D) Straight-Line Velocity; VSL and

(E) Linearity; LIN (F) Increase in the Oxygen consumption rate (OCR) after treatment with 10

nM OA or 10 nM OA + 40 nM Etomoxir (Eto). (G) The percentage of ATP-Red positive sperm

after 1 hour incubation with or without 10 nM OA, measured using BioTracker ATP-Red staining

and flow cytometry. (H) Artificial insemination using sperm treated with "pseudo" seminal

plasma from control (SV), flutamide-treated (Flu) mice and supplement of OA (Flu+OA)

followed by an evaluation of the fertilization rates. Data are mean ± SEM. n=3-6 independent

replicates. Student's t-test was used for comparison between the two groups.

Figure 7. Testosterone-regulated ACLY induces metabolic shifts in seminal vesicle epithelial

cells. (A) Western blot images of ACLY and α/β-tubulin in three sets of seminal vesicle epithelial

cells cultured with 100 ng/mL testosterone (Testo) or in vehicle (Ctrl) for 7 days. (B) Quantitative

analysis of ACLY relative to α-tubulin obtained from Western blot. (C) siRNA knockdown

experiments of ACLY in seminal vesicle epithelial cells. ACLY protein levels in scrambled

shRNA or ACLY shRNA-transfected seminal vesicle epithelial cells cultured with or without 100

ng/mL testosterone were determined by Western blot. (D-F) Changes in oxygen consumption of

seminal epithelial cells were analyzed by a flux analyzer: (D) Oxygen consumption rate (OCR)

kinetics of seminal vesicle epithelial cells transfected with scrambled shRNA or ACLY shRNA.

(E) Basal OCR and (F) Maximum respiration. (G) The impact of ACLY knockdown on

testosterone-induced fatty acid synthesis was measured. (H) The effect of culture supernatants of

testosterone-treated cells on sperm motility was evaluated, especially after ACLY knockdown.

Curvi-Linear Velocity; VCL, Straight-Line Velocity; VSL, Linearity; LIN. Data are mean ± SEM.

n=3-6 independent replicates. Repeated experiments were performed with cells recovered from 3

wells containing pooled cells from 3 to 5 mice. Student's t-test was used for comparison between

the two groups.

Figure 8. Testosterone regulates the metabolic activity of human seminal epithelial cells. (A)

Representative image of human seminal vesicle epithelial cells. (B) Growth curves of seminal

vesicle epithelial cells with 100 ng/mL testosterone. (C) Oxygen consumption rate (OCR)

kinetics and (D) Extracellular acidification rate (ECAR) kinetics of human seminal vesicle

epithelial cells after 6 days of culture with or without 100 ng/mL testosterone using a flux

analyzer. (E) Effects of testosterone on fluorescent glucose uptake and inhibitors of GLUT4

function. (F) Effects of testosterone and GLUT4 inhibitors on fatty acid synthesis. Data are mean

± SEM. n=3 independent replicates. Student's t-test was used for comparison between the two

groups.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

ACLY

α/β-Tubulin

Ctrl   Testo

B

p=0.0084

Relative intensity

Ctrl   Testo

C

ACLY

α/β-Tubulin

Testosterone	-	+	-	+
ACLY shRNA	-	-	+	+
Scramble shRNA	+	+	-	-

D

Ctrl+Scramble
Testo+Scramble
Ctrl+shACLY
Testo+shACLY

Oligomycin   FCCP   Rot/Ant

OCR (pmol/min)

E Basal OCR

OCR (pmol/min)

p=0.0011   p=0.0031

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

F ATP production

OCR (pmol/min)

p<0.001   p<0.001

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

G Lipid accumulation

Δmg/dL (10000 cells)

p=0.031

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

H

VCL (μm/min)

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

VSL (μm/min)

p=0.068   p=0.055

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

LIN (%)

p=0.0018   p=0.0073

Testosterone	-	-	+	+
ACLY shRNA	-	+	-	+
Scramble	+	-	+	-

Figure 7

Figure 8