Main Manuscript for

Structural insights into peptidoglycan hydrolysis by the FtsEX system in *Escherichia coli*
during cell division

Jianwei Li[1, #], Yutong He[1, #], Xin Xu[1, #], Martin Alcorlo[2], Jian Shi[3], David I. Roper[4], Juan A. Hermoso[2],

Lok-To Sham[5], and Min Luo[1, 3, *]

[1] Department of Biological Sciences, Faculty of Science, National University of Singapore

[2] Department of Crystallography and Structural Biology, Instituto de Química-Física "Blas Cabrera",

CSIC, Madrid, Spain

[3] Center for Bioimaging Sciences, Department of Biological Sciences, National University of

Singapore

[4] School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK

[5] Infectious Diseases Translational Research Programme and Department of Microbiology and

Immunology, Yong Loo Lin School of Medicine, National University of Singapore

[#] These authors contributed equally.

* Corresponding author. Email : dbslmin@nus.edu.sg

Author Contributions: Jianwei Li and Yutong He performed biochemical reconstitution, functional

characterization, and cryo-EM studies; Xin Xu performed kinetics assays; Martin Alcorlo, David

Roper, Juan A. Hermoso and Lok-To Sham helped with data analysis; Jian Shi helped with EM

data collection; Min Luo designed and supervised research, analyzed data, and wrote the paper

with help from all authors.

Competing Interest Statement: Authors declare no competing interests**.**

Classification: Biological Sciences; Microbiology

Keywords: cell division, peptidoglycan, antimicrobial, cryo-EM

This PDF file includes:

Main Text

Figures 1 to 6

Abstract

Bacterial cell division relies on precise peptidoglycan (PG) remodelling, a process orchestrated by

the FtsEX complex. Comprised of FtsE and FtsX, this complex collaborates with EnvC, a

periplasmic lytic enzyme activator, to regulate septal PG hydrolysis by amidases like AmiB. While

recent structural investigations, particularly of *Pseudomonas aeruginosa* FtsEX (*Pae*FtsEX), have

shed light on complex interactions and proposed activation mechanisms, the structural intricacies

governing PG degradation by the FtsEX complex and EnvC in *Escherichia coli* cytokinesis remain

unexplored. In this study, we present a comprehensive biochemical and structural analysis of *E.*

coli FtsEX complexes, unveiling a key role for ATP in complex stabilization that extends across

bacterial species. Upon EnvC binding, ATPase activity markedly increases. High-resolution

structures of *Eco*FtsEX, both in the presence and absence of EnvC, reveal a symmetrical

conformation of *Eco*FtsEX capable of accommodating the inherent asymmetry of EnvC, mediated

by flexible loops within the periplasmic domain. Our negative-staining imaging showcases an

elongated *Eco*FtsEX/EnvC/AmiB complex reminiscent of the *Pae*FtsEX system. These findings

collectively provide intricate insights into the regulation of PG cleavage by FtsEX in *E. coli* - a pivotal

model system used in pilot genetic studies, suggesting a conserved mechanism for precise

hydrolase activation in bacteria.

Introduction

In most bacteria, peptidoglycan (PG) is remodelled during cytokinesis in a highly coordinated

fashion. This coordination ensures the daughter cells are separated without causing lethal lesions

of PG (1-3). At least 10 essential proteins are involved in cell division in the model organism

Escherichia coli, many of which are organized by the cytoskeletal protein FtsZ (4, 5). FtsZ recruits

FtsA and FtsEX to form the early divisome. After a short pause, the early divisome recruits other

division proteins such as FtsK, FtsQLB, FtsWI, and FtsN to the septum (4, 6). The arrival of FtsN

somehow triggers cell constriction, likely through activating the FtsQLB complex and FtsA (4, 7-9).

Cell separation is then achieved through the amidases AmiA, AmiB, and AmiC. To control their PG

hydrolytic activity, these amidases are activated by coiled-coil domain proteins EnvC (for AmiA and

AmiB) and NlpD (for AmiC) (1, 3, 10, 11). If these systems are simultaneously inactivated, the

daughter cells will be unable to separate and form long chains (12).

PG remodelling is linked to cytokinesis by an ATP-binding cassette (ABC) transporter-like protein

complex FtsEX. Rather than transporting a substrate, the FtsEX complex transduces

conformational changes that span the cytoplasmic membrane (13, 14). Driven by ATP hydrolysis

of FtsE, these changes are required for stimulating EnvC/AmiA/AmiB activities (15, 16) and

initiating cell constriction via FtsA (15, 17-19). The FtsEX complex also recruits other cell division

proteins to the divisome (17, 19). Thus, deletions of *ftsEX* result in non-viable cells unless they are
grown in permissive conditions, such as at a low temperature or in media with high osmolarity (20,
21). Alternatively, the essentiality of *ftsEX* can be bypassed by gain-of-function mutations that
strengthen the interaction between FtsN and FtsA (17, 20).
72
Several partial structures of the FtsEX complex have been reported, some are with their interacting
partners (22-25). Very recently, progress has been made in elucidating the full-length structure of
the FtsEX complex from *Pseudomonas aeruginosa* (*Pae*FtsEX), both in the presence and absence
of critical regulators like ATP, EnvC, and AmiB (26). These structures offer insight into the interfaces
between FtsE, FtsX, and their binding partners. In addition, they suggest a mechanism in which
the rearrangement of regulatory helices of EnvC and AmiB activates peptidoglycan hydrolysis at
the septum (26). Yet, the majority of work regarding FtsEX was done in *Escherichia coli* (15-21),
where an abundance of genetic studies has laid a significant foundation for our understanding of
PG degradation activation in bacterial cell division. However, the intricate molecular basis
orchestrating this process remains elusive, underscoring the need for a direct structural and
biochemical investigation within this extensively studied model system. Specifically, given the
relatively low sequence similarity between the FtsEX complexes from these two species, it is
unclear whether the structural information of *Pae*FtsEX can reconcile the findings from genetic
studies. Furthermore, previous genetic studies have proposed a direct interaction between FtsEX
and components of the constriction ring, such as FtsA and FtsZ (18, 27), to ensure the accurate
localization of FtsEX at the division site. However, there is currently a lack of supporting
biochemical and structural evidence in this regard.
90
In this study, we present a comprehensive analysis of the structure and function of the *Escherichia*
coli FtsEX complex (*Eco*FtsEX) in conjunction with EnvC, focusing on its regulation of AmiB
activation for PG cleavage. Our investigation uncovers a previously unrecognized function of ATP
in stabilizing the FtsEX complex, a role that appears to be conserved across different bacterial
species. Notably, we observe a substantial increase in the ATPase activity of *Eco*FtsEX upon
binding with EnvC. High-resolution structures of *Eco*FtsEX alone and with EnvC were obtained at
3.9 and 3.4 Å, respectively, offering insights into the EnvC recognition mechanism. Interestingly,
*Eco*FtsEX maintains a symmetrical conformation even when complexed with the asymmetrical
EnvC. Comparative analyses identify the periplasmic domain's flexible loops as key to EnvC
interaction, revealing structural adaptability. Furthermore, our negative-staining imaging
experiments unveil an elongated conformation of the *Eco*FtsEX/EnvC/AmiB complex for PG
cleavage, reminiscent of the FtsEX system from *Pseudomonas aeruginosa*. Together, our results
revealed a detailed regulation of PG cleavage by FtsEX in the key model system of *E. coli*,
suggesting a conserved mechanism for hydrolase activation in bacteria.

Results

ATP Binding as a Universal Stabilizing Factor for the FtsEX Complex

In our efforts to reconstitute the full-length *Eco*FtsEX complex in vitro, we explored various strategies involving changing the expression constructs and extraction conditions. During the purification process, we observed that the protein remained stable in the presence of 10% glycerol. However, almost all protein precipitated out after the removal of glycerol either through size-exclusion chromatography (SEC) or during the peptidisc reconstitution (**Figure 1A**). This indicated a low level of stability for the complex. To investigate whether the stability of the FtsEX complex might be dependent on its cognate partner, EnvC, we conducted co-expression experiments by introducing EnvC along with FtsE and FtsX (**Figure 1A**). Despite the presence of EnvC, FtsEX still exhibited a tendency to precipitate.

Given the persistent challenges in achieving a stable FtsEX complex, we redirected our focus towards exploring the potential role of the substrate ligand, ATP, in complex stabilization. ATP, well-known for its role as a biological energy source, has also been associated with the stabilization of proteins, as reported in recent studies (28, 29). To investigate this hypothesis, we conducted a series of experiments to assess the impact of ATP on the FtsEX or the FtsEX/EnvC complex (**Figure 1A**). Remarkably, the introduction of ATP resulted in the formation of a robust *Eco*FtsEX complex, both in its standalone form and in the presence of EnvC. This complex remained tightly associated throughout the SEC process (**Figure supplement 1A**). Further validation through negative-staining electron microscopy confirmed that the purified *Eco*FtsEX complex maintained stability, with clearly discernible features corresponding to an intact ABC transporter containing two FtsE and two FtsX subunits (**Figure 1B**). This compelling evidence suggests that ATP, acting as the substrate of FtsEX, also plays a pivotal role in stabilizing this crucial complex formation.

Interestingly, ATP analogues, including ATPγS, AMP-PNP, and ADP, also exhibited similar stabilizing effects (**Figure 1C**). As they are not hydrolysable for FtsEX, it indicates that ATP binding, rather than hydrolysis, is sufficient for complex stabilization. This was corroborated by the introduction of a FtsE mutant (E163Q), inspired by related ABC transporters (26), which is incapable of ATP hydrolysis. When this mutant was assessed for complex formation with FtsX in the presence of ATP, it indeed formed a stable complex (**Figure 1C**). These findings suggest ATP binding alone is responsible for stabilizing the complex formation.

To expand the scope of our investigation, we explored whether ATP could similarly stabilize FtsEX complexes from other bacterial species, which had also presented challenges in in-vitro reconstitution. We expressed and purified FtsEX from various species, including *Streptococcus*

pneumoniae, Pseudomonas aeruginosa, and *Bacillus subtilis,* in the presence and absence of ATP.
Remarkably, regardless of the bacterial species, ATP consistently enhanced the stability of the
complexes, as demonstrated by either increased complex yield in SEC elution profiles or with a
more accurate stoichiometry between FtsE and FtsX as shown from the SDS-page gel (**Figure**
supplement 1B). Collectively, our findings underscore the pivotal role of ATP in stabilizing FtsEX
complexes. Notably, the ability of other nonhydrolizable ATP analogues to replicate this effect
suggests that ATP binding is a crucial step required to facilitate the coupling of FtsX and FtsE for
the formation of a functionally stable complex.
150
Biochemical Characterization of FtsEX and Its Complexes with EnvC in *E. coli*
The His-tagged full-length FtsE and FtsX proteins from *E. coli* (str. K-12, MG1655) were
overexpressed in *E. coli* BL21(DE3) and subsequently purified using a combination of affinity and
size-exclusion chromatography (SEC) (**Figure supplement 1A**). For subsequent functional and
structural investigations, FtsEX was solubilized in DDM and reconstituted into peptidiscs (see
Methods). Given the proven stabilization role, 0.5 mM ATP was maintained throughout the entire
purification process of *Eco*FtsEX.
158
Assays assessing ATPase activity demonstrated that, in contrast to previously reported FtsEX
complexes from *Pseudomonas aeruginosa*, the *Eco*FtsEX complex exhibited obvious basal
ATPase activity (**Figure 1D**). To confirm this, we introduced the FtsE mutant (E163Q) as a control,
which was not able to hydrolyse ATP. As anticipated, ATPase activity in the mutant was markedly
diminished (**Figure 1D**), while low ATPase activity was still detected, potentially attributable to co-
purified proteins.
165
To understand the role of ATP binding/hydrolysis on EnvC recruitment, a protein pull-down assay
was conducted. As ATP was present in the purified protein, a wash step with buffer in the presence
of Mg^{2+} and 10% glycerol was performed to remove the extra ATP for the FtsEX complex that bind
to the resin. Next, bound FtsEX and EnvC were mixed in the presence of ATP or its non-
hydrolysable analogue, ATPγS (**Figure 1E**). The results revealed that EnvC stably associated with
FtsEX, with its binding seemingly unaffected by ATP hydrolysis. This was evidenced by the
comparable amounts of EnvC bound in the presence of 3 mM ATPγS and 3 mM ATP. To confirm
this, FtsEX^{E163Q} mutant was introduced. Results, similar to those from the wild type, showed that
the mutant still demonstrated similar EnvC binding levels. This confirms that ATP binding is
sufficient to initiate EnvC recruitment.
176
Additionally, the effects of EnvC binding on the ATPase activity of *Eco*FtsEX were probed (**Figure**
1D). An enhanced activity was observed upon the addition of EnvC. This EnvC-mediated increase

in ATPase activity was further validated through ATPase assays involving varying ATP

concentrations for both FtsEX and the FtsEX/EnvC complex. Both sets of data fit well with the

Michaelis-Menten equation (**Figure 1F**). Notably, in the presence of EnvC, the V_{max} (92.7±5.1

nmol/mg/min) was approximately 2.2-fold higher than that observed for FtsEX (42.7±2.1

nmol/mg/min) alone. However, no large differences were observed regarding the K_m (0.11±0.02

mM for FtsEX/EnvC; 0.13±0.02 mM for FtsEX). Together, these findings underscore that in the

presence of ATP, FtsEX and EnvC form a robust complex; while *Eco*FtsEX inherently exhibits

ATPase activity, the presence of EnvC stimulates this activity.

CryoEM structure of *Eco*FtsEX reveals a symmetrical PLD arrangement

To elucidate the full-length structure of *Eco*FtsEX, we subjected the WT *Eco*FtsEX sample,

incubated with 3 mM ATPγS, to single-particle CryoEM analysis and achieved a resolution of 3.7 Å

(**Figure supplement 2**). This map was of sufficient resolution for detailed model-building for the

Transmembrane domains (TMD), the nucleotide-binding domains (NBD), and a large portion of the

Periplasmic domain (PLD); in addition, density for two bound nucleotides was clearly captured as

well (**Figure supplement 3A**).

*Eco*FtsEX, akin to most Type VII ABC transporters, possesses one coupling helice (CH) and a C

terminal tail (C-tail) (**Figure 2A**). Cryo-EM structure reveals the assembly of the full *Eco*FtsEX

system (**Figure 2B**). This comprises two cytosolic FtsE NBD domains and two FtsX, with TMD

domains embedded in the membrane and PLD extending into the periplasmic space. Both TMD

and NBD regions in the *Eco*FtsEX structure exhibit canonical folds, similar to other characterized

Type VII ABC transporters such as MacB, LolCDE, and *Pae*FtsEX. The overall conformation of

*Eco*FtsEX is quite similar to that of *Pae*FtsEX (**Figure supplement 3B**). Notably, the N-terminal

region of FtsX (Residues 53-70) forms an elbow helix (EH) oriented parallel to the membrane

(**Figure 2B**). Like its counterpart in *Pae*FtsEX, the extreme N-terminal region of FtsX (Residues 1-

52) was not resolved, suggesting its highly disordered nature.

The entire PLD domain comprises a smaller loop and a larger domain. For ease of reference, only

the larger domain is termed the PLD (**Figure 2A**). The smaller loop encompasses 14 residues

between TM3 and TM4. The principal PLD domain, spanning residues between TM1 and TM2

(106-208), displays a canonical topology, with a Porter domain and an X-lobe (14, 22). Compared

to the TMD and NBD regions, PLD domains exhibit higher flexibility, evident from their diminished

EM density at a contour level of 0.22 in Chimera. Lowering the contour to 0.1 in Chimera reveals

clear density, enabling the visualization of PLD's secondary structures (**Figure supplement 2F**),

with an exception for the X-lobe segment with residues 161-168 (**Figure 2C**). Structurally, the

individual PLD domains are superimposable (**Figure 2C)**, and also align closely with those in

*Pae*FtsEX (**Figure supplement 4**), despite their amino acid sequence identities with *E. coli* being

only 29%.

The PLD domain of *Eco*FtsEX structure offers a snapshot with well-defined secondary structures,

highlighting a 2-fold symmetrical conformation (**Figure 2D)**. The X-lobes of both PLDs are oriented

mutually inwards with sideways blocked and appear to facilitate EnvC binding strictly from the top

of the FtsEX complex, mirroring findings in the *Pae*FtsEX system.

CryoEM structure of *Eco*FtsEX/EnvC Complex elucidates EnvC binding and ATP-Dependent

LytM Release

To elucidate the EnvC recruitment by FtsEX in *E. coli*, purified FtsEX[E163Q] mutant was reconstituted

in peptidiscs and incubated with EnvC in the presence of 2 mM ATP for cryo-EM analysis (**Figure**

supplement 5). After 3D reconstruction, a major conformation stands out with resolution

determined at 3.4 Å (**Figure supplement 5F**). The quality of the EM density allowed modelling of

nearly all sidechains of FtsEX, the two bound nucleotides, and a portion of EnvC's coiled-coil

domain (CCD) (**Figure 3A & Figure supplement 6**). The density of EnvC's remaining regions,

including part of the CCD domain, the LytM domain and the regulatory helix, was absent,

suggesting significant flexibility. The global configuration of the FtsEX/EnvC complex is similar to

the previously described *Pae*FtsEX system (**Figure supplement 7**), where EnvC inserts from

above between the two PLD domains along the FtsEX central axis.

To dissect the state of the bound EnvC, we compared our cryoEM FtsEX/EnvC complex with the

crystal structure of the PLD domain in complex with EnvC (PLD/EnvC) from *E. coli* (PDB code:

6TPI) *(22)*, where EnvC was in an inactivated state. In the PLD/EnvC structure, a distinct

asymmetrical conformation is evident: one PLD is positioned about 5 Å higher than the other one,

accommodating the bound asymmetric EnvC with its LytM domain locked (**Figure supplement 8A,**

left). However, in our full-length ATP-bound FtsEX/EnvC structure, the two PLDs are highly

symmetrical (**Figure supplement 8A, right**). Besides this, the bound EnvC has also undergone

conformational shifts (**Figure supplement 8B**). In contrast to the PLD/EnvC crystal structure,

where the two CCD helices exhibit huge asymmetry (**Figure supplement 8B, left**), our structure

reveals a more symmetrical arrangement induced by simultaneous compression from both PLD

domains (**Figure supplement 8B, right**). Consequently, the distance between the terminal residue

of the α1 helix (Q127) and the initial residue of the α2 helix (Q143) within the CCD domain is

reduced from 24 Å to a mere 10 Å, while residing at a relatively similar height from the PLD domain.

This compression of the CCD helices may drive the release of the LytM domain, supported by the

lack of LytM density in our EM map. Together, we hypothesize that in the presence of ATP, the

LytM domain is released, facilitating AmiB recruitment.

Detailed EnvC interactions with FtsEX

The ATP-bound *Eco*FtsEX/EnvC structure also revealed detailed interactions with EnvC. This interaction mainly involves two interfaces and covers an approximate surface area of 2290 Å^2, predominantly consisting of hydrophobic contacts (**Figure 3B**).

The primary interface, also resolved in the PLD/EnvC crystal structure (22), features residues Y114, F152, W155, F158, L162, L165, and L170 of FtsX associating with residues L117, L121, L148, Y151, F152, Y154, L155, and I163 of EnvC (**Figure 3C**). Conversely, the secondary interface encompasses the Lip of CCD from EnvC, and the small loop of FtsX. Interacting residues at this interface comprise I133, L135, I136, L137, E141 from EnvC and W210, L214, F309 from FtsX. Although hydrophobic interactions dominate, several electrostatic interactions exist, notably the salt bridges (R126$_{EnvC}$-D207$_{FtsX}$, E141$_{EnvC}$-R213$_{FtsX}$) (**Figure 3D**). These interactions have largely been validated by previous studies (22), mutations like Y114E, F152E, and L165A in FtsX's periplasmic domain notably affected EnvC's binding efficacy, affirming their role in recruiting EnvC.

Recognition of Asymmetrical EnvC by Symmetrical FtsEX: Unveiling Key Loops for Conformational Adaptivity

Intriguingly, our analysis of the FtsEX/EnvC structure showed that even when bound to the asymmetrical EnvC, FtsEX—including its two PLD domains interacting directly with EnvC—maintains a symmetrical conformation (**Figure 4A**). Compared to the *Eco*FtsEX structure without EnvC, which is also symmetrical, the only noticeable change is the outward shift of the two PLD hooks by about 12 Å (**Figure 4A**), a result of their interaction with the CCD domain of EnvC (**Figure 4B**). This observation poses an interesting question: How does the inherently symmetrical FtsEX cater to an asymmetrical binding partner? To explore this, we closely analysed the FtsX-EnvC binding interface.

Upon detailed examination, we noticed that the motifs in FtsEX directly interacting with EnvC are primarily loops (**Figure 4C**). The main loop engaging with EnvC is X-lobe (**Figure 4D**), covering residues G157-L171. This segment, originally disordered in the FtsEX alone structure, becomes crucial for engaging with EnvC, involving a binding interface of about 1204 Å^2. The other interacting regions are loops spanning residues Y104-Q109 (**Figure 4E**) and D208-L214 (**Figure 4F**), and a segment of the small loop in the PLD domain (Q307-T311) (**Figure 4G**). Combined, these loops (eight in total from the two PLD domains) are responsible for the majority of the interactions with EnvC. This inherent adaptability in the loops provides FtsEX the versatility needed for binding the asymmetrical EnvC. Notably, the interactions contributed by these loops are largely hydrophobic (**Figure 4D-G**). Thus, our data suggest that the flexible loops, along with the hydrophobic

interactions they facilitate, equip the symmetrical FtsEX to effectively engage with asymmetrical

partners.

Reconstitution and Elucidating the Elongated Conformation of *E. coli* FtsEX/EnvC/AmiB

Complex for PG cleavage

The absence of the LytM density, in comparison to the crystal structure of PLD/EnvC (22), indicates

that our ATP-bound FtsEX/EnvC sample likely has the LytM domain released, primed for hydrolase

interaction. To further validate this, we conducted pull-down assays (**Figure 5A**). When we

incubated AmiB with either the FtsEX/EnvC complex or the mutant FtsEX^{E163Q}/EnvC in the

presence of ATP or its nonhydrolyzable analogue, ATPγS, both the wild type and mutant

FtsEX/EnvC complexes effectively pulled down AmiB. Notably, ATP and ATPγS showed similar

levels of AmiB binding, confirming that ATP-bound FtsEX/EnvC is prepped for AmiB recruitment.

Additionally, we investigated the influence of AmiB binding on FtsEX's ATPase activity (**Figure 5B**).

There was no obvious effect upon the addition of AmiB to the FtsEX/EnvC complex, suggesting

that AmiB does not regulate FtsEX function.

Next, we sought to check if the bound AmiB is activated for PG cleavage. FtsEX/EnvC complex

was first mixed with a threefold molar excess of AmiB in the presence of 2 mM ATP, and the mixture

was applied to size-exclusion chromatography (SEC) (**Figure 5C**). An early eluting fraction was

observed, and subsequent SDS-PAGE confirmed the co-existence of FtsE, FtsX, EnvC, and AmiB

in this fraction. The reconstituted PG-cleaving complex of *Eco*FtsEX/EnvC/AmiB, was further

analysed using negative staining (NS) electron microscopy (**Figure 5D**). Many of the observed

particles exhibited long tails in the image, a feature suggestive of AmiB binding. A 2D classification

analysis presented an elongated structure, mirroring previous observations in *P. aeruginosa* (26).

The electron densities on the periplasmic side displayed a distinct rod-like form in alignment with

the FtsEX complex's central axis, extending about 290 Å into the periplasm, which we attribute to

the bound EnvC and AmiB. This conformation likely represents an activated state prepared for PG

cleavage. However, due to the *Eco*FtsEX complex's instability during the PG hydrolysis assay, we

couldn't directly check PG cleavage activity. The attempt to obtain a CryoEM structure of the

complex was also unsuccessful, possibly because of its inherent flexibility, preventing the

reconstruction of a clear EM map. Nonetheless, it has been well established in *P. aeruginosa*,

where the active FtsEX/EnvC/AmiB complex also assumes an extended form (**Figure 5E**), we

propose that our observed complex with a similar conformation is also in an activated state, primed

for PG degradation.

Discussion

A structural and functional description is here provided for the complete *E. coli* lytic machinery
regulated by the FtsEX system. We observed an antiparallel arrangement of PLDs in the
periplasmic space in the *Eco*FtsEX complex. Incorporation of ATP, but not hydrolysis, is capable
of facilitating EnvC binding. The ternary complex increases its ATPase activity, compared with that
of the binary complex. Interestingly, upon ATP binding, the flexibility of loops in the PLD domain
allows a symmetric interaction with the EnvC coiled-coil domain that is compressed to allow the
release of its LytM domain. While resolution for the complete FtsEX/EnvC/AmiB complex limited
detailed descriptions, our cryo-EM data points to a very similar disposition of the lytic complex to
that reported for *P. aeruginosa* (26) with a fully expanded EnvC coiled-coil that would place the
amidase at ~300 Å over the inner membrane.

In the context of recruiting an asymmetric partner by the inherently symmetrical FtsEX complex,
our study has unveiled a dynamic and intriguing mechanism. At the centre of this mechanism lies
the highly flexible PLD domain (composed of the porter domain and the loops connecting the
different TM of FtsX), which demonstrates exceptional adaptability due to the presence of multiple
flexible loops, including the x-lobe. These flexible loops within the PLD domain function as dynamic
connectors, enabling the FtsEX complex to finely tune its conformation and engage with
asymmetric partners in an exceptionally adaptable manner. This inherent plasticity empowers the
FtsEX complex to align itself with the specific structural requirements of its binding partners, thereby
facilitating their effective interaction and subsequent regulatory processes.

Moreover, these flexible elements serve a dual role by providing the necessary adaptability to
bridge the gap between the inherent symmetry of the FtsEX complex and the asymmetry present
in its interacting partners. This becomes particularly intriguing in the context of bacterial cell
division, where the same FtsEX complex must interact with a variety of asymmetric binding
partners. These binding partners exhibit diverse structural features and distinct binding interfaces,
necessitating a mechanism that allows the inherently symmetrical FtsEX complex to engage with
them with precision and effectiveness.

Our study has uncovered a previously unrecognized role of ATP in the stabilization of the FtsEX
complex, a role that extends beyond its established function as a substrate. This discovery not only
expands our understanding of the functional versatility of ATP but also unravels a crucial aspect of
how FtsEX complexes are regulated. The stabilizing influence of ATP is particularly intriguing, as
evidenced by our experiments across diverse bacterial species, suggesting that ATP-driven
stabilization might represent a shared feature among FtsEX complexes in various bacteria.
Furthermore, it is noteworthy that both the FtsEX systems from *E. coli* and the recently
characterized system from *P. aeruginosa* exhibit a relatively low basal ATPase activity. However,

this activity is significantly enhanced in the presence of their respective cognate binding partner,
EnvC. Specifically, for *E. coli*, the ATPase activity increases by a factor of 2.2, whereas for *P.*
aeruginosa, the increase is even more pronounced, with a greater than 20-fold enhancement
observed. This additional regulatory mechanism ensures that, under normal physiological
conditions, minimal energy is expended. It is only during the cell division process, when
components like EnvC are involved, that the ATPase activity of FtsEX becomes fully activated.

Interestingly, our cryoEM map of EcoFtsEX, expressed from its native host bacteria *E. coli*,
revealed an additional density located on the bottom side of the complex adjacent to FtsE (**Figure**
supplement 2, and **Figure supplement 9A**). This suggests the co-purification of endogenously
binding proteins with the complex. To identify the proteins, we conducted mass spectrometry on
the purified FtsEX sample. Our results showed that in addition to FtsE and FtsX, FtsA was present
in significant abundance, followed by FtsZ (**Figure supplement 9B**). This observation is consistent
with previous genetic and phenotypic observations on the *E. coli* divisome activation (17, 27). We
also noted during the course of these studies that the interaction of *E.coli* FtsA with FtsEX protein
complexes from other species is indicative of the conserved and fundamental nature of divisome
protein interactions. For example, the overexpression of *S. pneumoniae* FtsEX in *E. coli* also
indicated an association with *E.coli* FtsA and could be isolated using styrene-maleic acid isolation
and affinity purification as shown in **Figure supplement 9C** (30, 31). If this holds true, it would
provide a mechanism for the temporal regulation of PG cleavage at the cell division site, as FtsA
and FtsZ are the key components of the constriction ring at the septum site (15, 18, 27, 32-36).
However, the identity of this density and the association of FtsEX with other Fts components for
divisome formation warrant further investigation to fully elucidate the role of FtsEX in bacterial cell
division.

In combination with previous studies on the inactivated EnvC (22) and AmiB activation by
*Pae*FtsEX (26), we propose a working model that may be conserved across Gram-negative FtsEX
systems (**Figure 6**). Firstly, physiological ATP stabilizes the formation of a functional FtsEX
regulator with low basal ATP hydrolysis activity. Secondly, the inactivated EnvC approaches the
periplasmic space via the relatively flexible PLD domain of FtsEX, which adopts a highly
asymmetric conformation, as shown in the inactivated EnvC structure (22). Thirdly, upon full
recognition by FtsEX, considering the presence of ATP in cell and a symmetrical conformation
defined within the context of a full-length FtsEX complex, recruited EnvC is compelled to adopt a
relatively symmetrical conformation, involving squeezing of the coiled-coil domain to facilitate the
release of the LytM domain; With EnvC engagement, the ATPase activity of FtsEX is fully activated.
Finally, the released LytM recruits amidases through direct interaction, consequently removing the
blocking helix from the catalytic site and activating PG cleavage. Remarkably, expansion of the

coiled-coil domain in EnvC would place the amidase domain ~300 Å over the inner membrane of
the Gram-negative bacteria (as here seen in *E. coli*, and also in *P. aeruginosa* (26)), compatible
with the location of the septal PG (26) and avoiding damages into the non-septal cell wall.
Throughout this process, ATP binding appears to be sufficient, with ATP hydrolysis potentially
serving as a driving force for the release of EnvC and amidases from FtsEX, allowing them to be
recycled to a normal ABC transporter. This transporter by itself may play additional roles, such as
antibiotic resistance, as suggested by previous studies (22). For the temporal regulation of PG
cleavage, the potential interaction between FtsEX and FtsA/FtsZ, as indicated in this study and
previous genetic research (17, 27), could bridge the PG-cleaving complex with septum constriction
ring, facilitating precise cell division at the septum site. This model provides a potential conserved
mechanism for PG cleavage activation by FtsEX and illuminates its temporal regulation. Although
further validation is required, our results offer valuable biochemical and structural insights that
complement and elucidate decades of genetic research in *E. coli* and other Gram-negative bacteria.
413

Materials and Methods

Cloning and Mutagenesis
The primers and templates utilized in this study are listed in **Table supplement 1**. To generate the
PCR amplicon containing FtsEXs in genomic DNA of *Escherichia coli* Str *K-12 MG1655*,
Streptococcus pneumoniae str. D39, *Bacillus_subtilis_subsp_subtilis_ATCC_23857*, and
Pseudomonas aeruginosa ATCC 47085D. ClonExpress II One Step Cloning Kit (Vazyme) was
used, followed by purification. After purification through gel electrophoresis, the resulting product
was ligated to pTD68 vector (PMID: 20300061) digested with the Nde I and Hind III enzymes to
produce pTD68 [PT7::his6-FtsEX]. The construction of pET28-sumo (PT7:his6-sumo-
*Eco*EnvCΔ34), (PT7:his6-sumo-*Eco*AmiBΔ22), involved ligating pET28-sumo digested with BamH
I and Xho I to a PCR amplicon synthesized by EnvC-F, EnvC-R, AmiB-F, AmiB-R which were also
digested with the same enzymes. The plasmids utilized in this study are listed in **Table supplement
2.** Mutations to *Eco*FtsE was generated utilizing the QuikChange kit (Agilent Technologies), using
pTD68 and pET28-sumo as templates and the primers listed in **Table supplement 1**. The
mutations were then verified by DNA sequencing.
429

Protein expression, purification of FtsEX complex and their mutants
FtsEX and its mutants are expressed in *E. coli* strain BL21(DE3) pLysS (Invitrogen) using lysogeny
broth (LB) supplemented with 50 µg/ml ampicillin and 10 µg/ml chloramphenicol. This is incubated
at 37°C until an optical density at 600 nm (OD_{600}) of 0.8 is achieved. Following this, induction with
0.5 mM isopropyl-β-D-thiogalactoside (IPTG) takes place at 20°C for 16 hours. The cells are
subsequently harvested through centrifugation using a Beckman JLA9-1000 rotor at 5,180 xg and
20°C for 15 minutes. Harvested cells are then stored at -80°C.

For membrane extraction, the frozen cell pellets undergo thawing at 4°C and are resuspended in

buffer A, consisting of 50 mM Tris-HCl pH 8.0, 250 mM NaCl, 10% glycerol, and 2 mM ATP/MgCl$_2$.

Lysis of the mixture was achieved with a high-pressure homogenizer (AH-100D, ATS). Subsequent

centrifugation was performed at 27,000 xg for an hour at 4°C. Following this, solubilization of the

membrane takes place in buffer B, containing 50 mM Tris-HCl, 250 mM NaCl, 1.3% n-dodecyl β-

D-maltoside (DDM), pH 8.0, 10% glycerol, and 2 mM ATP/MgCl$_2$, and agitated for 2 hours at 4°C.

Post a 30-minute centrifugation at 100,000 xg, the solubilized membrane fraction gets incubated

with TALON SuperFlow resin (Takara 635502) for 1 hours at 4°C and was later washed using buffer

B supplemented with 20 mM imidazole. The present detergent is substituted by a peptide

(GenScript) to form peptidiscs, employing the "on-beads" method (37, 38) with a final peptide

concentration of 1mg/ml. The FtsEX complex, reconstituted in the peptidiscs, is washed using

buffer B inclusive of 20 mM imidazole and eluted with buffer B rich in 200 mM imidazole. The

purified protein is directed through a gel filtration chromatography column (Superose 6 increase

10/300) and is eluted with buffer C, which includes 25 mM Tris-HCl, 150 mM NaCl, pH 8.0, and 2

mM ATP/MgCl$_2$. The fractions bearing the FtsEX complex are combined together, concentrated to

3 mg/ml by concentrator (Amicon® Ulyra-15 100kDa), and preserved at -80°C for further protein

interaction, pull-down, and ATP hydrolysis assay.

Protein expression, purification of EnvC, AmiB

For the expression of EnvC, AmiB and their respective variants, the *E. coli* strain BL21(DE3) pLysS

was employed. These cells were cultured in LB medium with 50 µg/ml kanamycin at 37°C. Once

the OD$_{600}$ reached a value of 0.6, induction was facilitated by the addition of IPTG to achieve a final

concentration of 0.5 mM, and this was maintained for 16 hours at 20°C. Following the induction,

cells were harvested through centrifugation using a JLA9-1000 rotor in a Beckman centrifuge at

5,000 rpm and 20°C for a duration of 15 minutes. The resulting pellets were stored at -80°C. For

further processing, these frozen pellets were brought to 4°C for thawing, using buffer C (50 mM

Tris-HCl pH 8.0, 150 mM NaCl, and 10% glycerol) and then subjected to lysis via a high-pressure

homogenizer (AH-100D, ATS). Subsequent centrifugation was performed at 27,000 xg for an hour

at 4°C. The resultant supernatant was incubated with TALON resin, followed by a wash using buffer

C complemented with 20 mM imidazole. Post-washing, the His-tagged SUMO protease Ulp1 was

introduced to the resin-buffer at a ratio of 0.1 mg Ulp1 for every liter of culture, facilitating the

cleavage of the SUMO tag over 2 hours at ambient temperature. The processed samples were

isolated, concentrated, and then passed through the Superose 6 increase 10/300 column

previously balanced in buffer D which includes 25 mM Tris-HCl, 150 mM NaCl, pH 8.0. Target

protein was confirmed using SDS-PAGE, further concentrated up to 3 mg/ml, and stored at -80°C.

Analyzing the Impact of ATP on the FtsEX Complex

Expression and purification of FtsEXs which from *Escherichia coli (Eco)*, *Streptococcus*
pneumoniae (Sp), *Bacillus_subtilis (Bs)*, and *Pseudomonas aeruginosa (Pae)* followed the
procedure above. For the ATP-free FstEXs purification, the whole process excludes ATP. For the
samples of FtsEXs with ATP or ATP analogs, 2 mM ATP or ATP analogs present in the purification
process. The processed samples of *Sp*FtsEX, *Bs*FtsEX, *Pae*FtsEX were concentrated, and then
passed through the Superose 6 increase 10/300 column previously balanced in buffer which
includes 25 mM Tris-HCl, 150 mM NaCl, pH 8.0 with or without 2 mM ATP. The eluted proteins
profiles were confirmed using SDS-PAGE gel. For the *Eco*FtsEX/EnvC samples purification add
EnvC with final concentration 1mg/ml after FtsEX reconstituted in the peptidiscs. The eluted ATP-
free *Eco*FtsEX sample was performed at 12,000 xg for 15 min at 4°C due to persistent precipitate
appearing in the eluted sample. Confirming the supernatant and precipitate using SDS-PAGE gel.

Protein interaction assay

To explore the interaction between FtsEX and EnvC, we employed affinity chromatography with
TALON resin. Specifically, 15 μg of FtsEX or its ATPase-defective variants were introduced to the
resin pre-equilibrated at room temperature, utilizing a Micro Gravity Column, for a 15-minute
duration. After washing away unbound FtsEX with 100 μl of buffer E (50 mM Tris-HCl pH 8.0, 150
mM NaCl, 10% glycerol, and 2 mM MgCl$_2$) supplemented with 5 mM imidazole (to get rid of the
extra ATP present in the protein), we incorporated 5 μg of EnvC diluted in 60 μl of the
aforementioned buffer in the presence and absence of ATP or ATP analogs. This mixture was
incubated for an additional 10 minutes. Subsequently, the beads were rinsed with buffer E
supplemented with 5 mM imidazole in the presence and absence of ATP or ATP analogs. To elute
the bound proteins and possible interaction partners, we utilized 20 μl of buffer E with a
concentration of 300 mM imidazole, either with or without 2 mM ATP or ATP analogs. Eluted
proteins were then analyzed using SDS-PAGE.

To explore the interactions between FtsEX/EnvC and AmiB, 20 μg of FtsEX/EnvC or its ATPase-
deficient variants, FtsE^{E163Q}X/EnvC, were added to the TALON resin. This was done either in the
presence or absence of 2 mM ATP or ATP analogs and incubated at room temperature for 15
minutes. Any free FtsEX/EnvC or FtsE^{E163Q}X/EnvC was removed by washing the resin with 100 μl
of buffer E containing 5 mM imidazole. Following this, 10 μg of AmiB were introduced to the resin
with or without 2 mM ATP or ATP analogs. After a 10-minute incubation, a triple wash was executed
using 100 μl of buffer E supplemented with 5 mM imidazole, either with or without 2 mM ATP or
ATP analogs . Finally, bound proteins were eluted using 20 μl of buffer E supplemented with 300
mM imidazole, either with or without 2 mM ATP or ATP analogs, and the elution profile was
assessed via SDS-PAGE.

Reconstitution of FtsEX complexes

For the reconstitution of the FtsEX-EnvC complex, an excess of EnvC, three times the amount of FtsEX, was introduced to the resin-bound FtsEX-peptidiscs during the purification process. This mixture was allowed to interact at room temperature for 15 minutes. Subsequent to this incubation, any free EnvC was removed by washing with buffer A containing 20 mM imidazole. The FtsEX-EnvC complex was subsequently eluted from the resin using buffer A supplemented with 200 mM imidazole. This eluted protein was applied to additional purification through gel filtration on a Superose 6 increase 10/300 column, equilibrated in buffer B. Fractions containing the FtsEX-EnvC complex were assessed using SDS-PAGE, combined, concentrated to a concentration of 3 mg/ml, and stored at -80°C.

For the reconstitution of the AmiB-EnvC-FtsEX supercomplex, AmiB was mixed with the FtsEX-EnvC complex in peptidiscs, at a 3:1 ratio. Following a 15-minute incubation, the supercomplex was further purified via gel filtration. Fractions encompassing the FtsEX-EnvC-AmiB complex were collected, concentrated to 3 mg/ml, and subsequently stored at -80°C.

ATP hydrolysis assay

To assess the ATPase activity of FtsEX and its mutants or complxes, we employed the malachite green phosphate assay (39). We introduced 2 µg of FtsEX with or without 0.7 µg of EnvC or 0.7 µg AmiB of following 0.8 µg of EnvC into a 20 µl reaction buffer comprising 25 mM Tris-HCl (pH 8.0), 150 mM NaCl, and 2 mM ATP/MgCl$_2$. For the elucidation of the Michaelis-Menten constants, varying concentrations of ATP/MgCl$_2$, ranging from 0.05 to 5 mM, were utilized to initiate the enzymatic reaction. After a 10-minute incubation at 37°C, 80 µl of the Malachite Green-Ammonium Molybdate (MG-AG) solution was added and the resultant mixture was quickly vortexed to deactivate the enzyme. A brief 2-minute incubation was followed by the addition of 10 µl of a 34% (w/v) sodium citrate solution, with a subsequent vortexing step. Absorbance was then recorded at 650 nm after a 5-minute interval. Quantification of released inorganic phosphate (Pi) was achieved using a calibration curve derived from known Pi concentrations in a K$_2$HPO$_4$ solution. Data processing and curve fitting to the Michaelis-Menten equation were conducted utilizing Microsoft Excel in conjunction with GraphPad Prism 9.

Negative staining imaging of FtsE^{E163Q}X-EnvC-AmiB complex

For negative staining of the FtsE^{E163Q}X-EnvC-AmiB sample, it was diluted to a concentration of 0.01 mg/ml. A 5 µl aliquot of the sample was deposited onto a 300 mesh copper grid coated with carbon film (EMS). After allowing the sample to interact with the grid for 60 seconds, any surplus was carefully blotted away using filter paper. The grid was then washed with 5 µl of Uranyless EM

Stain (EMS). Subsequent to the washing step, the grid was stained by applying another 5 µl of the

Uranyless EM Stain for 120 seconds. Excess stain was again removed with filter paper. The

prepared grid was left to air-dry under ambient conditions for approximately 15 minutes. Imaging

of the sample was executed with a Tecnai 12 microscope (FEI), operating at 120kV with a LaB6

source, and magnification set to 52, 000 x.

Cryo-Electron microscopy sample preparation and data acquisition

To prepare the sample, 3.5 µl of the protein sample at a concentration of 3 mg/ml was applied to

Quantifoil holey carbon grids (R1.2/1.3, 400 mesh) that were glow-discharged. To obtain the ATPγS

-bound complex, the sample was incubated at 37°C for 8 minutes with 3mM ATPγS-2mM Mg^{2+}

(FtsEX) or 2 mM ATP-Mg^{2+} (FtsE^{E163Q}X/EnvC) before freezing. The grids were blotted for 3.5-4

seconds with 100% relative humidity and then plunge-frozen in liquid ethane that was cooled by

liquid nitrogen using a Vitrobot System (Gatan).

Cryo-electron microscopy data was collected at liquid nitrogen temperature using a Titan Krios

electron microscope (Thermo Fisher Scientific), equipped with a K3 Summit direct electron detector

(Gatan) and GIF Quantum energy filter. All cryo-EM movies were recorded in counting mode with

SerialEM4 (40) with a slit width of 20 eV from the energy filter. Movies were acquired at nominal

magnifications of 81k x, which corresponded to a calibrated pixel size of 1.06 Å on the specimen

level. The total exposure time for each movie was 5 seconds, resulting in a total dose range of 35-

39 electrons per Å^2, fractionated into 34 frames. Additional electron microscopy data collection

parameters can be found in **Table supplement 3**.

Cryo-Electron microscope image processing.

For electron microscopy (EM) data processing, CryoSPARC (41) was utilized. Dose-fractionated

movies, captured with a K3 Summit direct electron detector, was first applied to motion correction

using MotionCor2 (42). A sum of all frames from each movie was generated applying a dose-

weighting scheme. This sum was utilized for all subsequent image-processing steps, except for

defocus determination. For that, CTFFIND4 (43) was used on the non-dose-weighted summed

images from all movie frames. The initial particle selection was achieved using the blob picker,

which was subsequently refined by the template picker. The steps of 2D and 3D classification, as

well as 3D refinement, were carried out using the "2D Classification", "Ab-initio Reconstruction",

and "Heterogeneous Refinement" functions. For further refinement, both the "Homogeneous

Refinement" and "Non-Uniform Refinement" functions were employed. The overall resolutions was

determined using the gold-standard Fourier shell correlation (FSC) threshold of 0.143. Lastly, the

local resolution was assessed via the "Local Resolution Estimation" tool.

Model building and refinement for FtsEX and its complex with EnvC

The cryo-EM maps of the ATPγS-bound FtsEX and ATP-bound FtsE^{E163Q}X/EnvC complexes were resolved at resolutions of 3.7 Å and 3.4 Å, respectively, with sufficient quality to enable accurate model building for the TMD domain, FtsE, and a portion of the PLD region (see **Figure supplement 3** and **Figure supplement 6** for details), leaving only the N-terminal region of FtsX, including residues 1-52, not modeled.

Initial models for FtsE and FtsX were generated using the AlphaFold program (41). Subsequently, the FtsEX models were rigid-body fitted into the cryo-EM map density using UCSF Chimera software (44). These initial models were then refined manually within Coot (45). Restraints for ATPγS were generated using the phenix_elbow program, employing its isomeric SMILES string files obtained from the PDB Chemical Component Dictionary via Ligand Expo. Further refinement was performed using Phenix (46), including 'minimization_global' and 'Real-space refinement' protocols. Multiple rounds of manual adjustments in Coot, in conjunction with real-space refinement in Phenix, were carried out to obtain the final model. Model validation was performed using Ramachandran plots, MolProbity scores, and Clash scores in Phenix (46) (details provided in **Table supplement 3**).

In the case of the FtsE^{E163Q}X/EnvC/ATP complex structure, the initial models for EnvC was derived from AlphaFold (47) and our resolved FtsEX structure was used as initial model for FtsEX part. These initial models were fitted into the cryo-EM map of the FtsE^{E163Q}X/EnvC/ATP complex using UCSF Chimera (44). Subsequent manual adjustments were performed within Coot (45), and the Glu163 was replaced with Gln. Similar to the ATPγS-containing complex, restraints for ATP were generated using the phenix_elbow program. The model was refined using phenix software (46), incorporating 'minimization_global' and 'Real-space refinement' procedures. The final model was refined through multiple iterations of manual adjustments in Coot (45), alongside real-space refinement in Phenix. Model quality assessment included Ramachandran plots, MolProbity scores, and Clash scores, with detailed results provided in **Table supplement 3 (46)**.

Acknowledgments

We appreciate the contributions of Yibei Xiao and Zhaoxing Li from China Pharmaceutical University, whose expertise in Mass Spectrometry greatly enhanced our research. Furthermore, we thank all Luo lab members for their constructive comments throughout the course of this study.

Funding: This work was supported by the National University of Singapore Start-up grant to M. L., the Ministry of Education Tier 2 Grant to M.L. and L.-T. S. (MOE-T2EP30222-0015, MOE-T2EP30220-0012, and T2EP30123-0017), the National Medical Research Council grant to L.-T. S.

(A-8001581-00-00), and the National Research Foundation grant to M.L. and L.-T. S. (NRF-
CRP22-2019-0001 and NRFF11-2019-0005). The work in Spain is supported by grants PID2020-
115331GB-100 funded by MCIN/AEI/10.13039/501100011033 and CRSII5_198737/1 by the Swiss
National Science Foundation to J.A.H.

Data and materials availability: Two three-dimensional cryo-EM density maps of FtsEX and its
complexes with EnvC have been deposited in the Electron Microscopy Data Bank under accession
codes: EMD-37324 (ATPγS-bound FtsEX); EMD-37325(ATP-bound FtsE^{E163Q}X with EnvC). Two
atomic models have been deposited in the Protein Data Bank under accession codes 8W6I (ATPγS
bound FtsEX), and 8W6J (ATP-bound FtsE^{E163Q}X/EnvC). All data presented in this study are
provided within the main text and supplementary materials, or readily accessible upon request.

References

1.   T. Do, J. E. Page, S. Walker, Uncovering the activities, biological roles, and regulation of
bacterial cell wall hydrolases and tailoring enzymes. *J Biol Chem* **295**, 3347-3361 (2020).
2.   A. Typas, M. Banzhaf, C. A. Gross, W. Vollmer, From the regulation of peptidoglycan
synthesis to bacterial growth and morphology. *Nat Rev Microbiol* **10**, 123-136 (2011).
3.   P. D. A. Rohs, T. G. Bernhardt, Growth and Division of the Peptidoglycan Matrix. *Annu Rev*
Microbiol **75**, 315-336 (2021).
4.   S. Du, J. Lutkenhaus, Assembly and activation of the Escherichia coli divisome. *Mol*
Microbiol **105**, 177-187 (2017).
5.   S. Du, J. Lutkenhaus, At the Heart of Bacterial Cytokinesis: The Z Ring. *Trends Microbiol*
27, 781-791 (2019).
6.   J. Lutkenhaus, S. Du, E. coli Cell Cycle Machinery. *Subcell Biochem* **84**, 27-65 (2017).
7.   Z. Lyu *et al.*, FtsN maintains active septal cell wall synthesis by forming a processive
complex with the septum-specific peptidoglycan synthases in E. coli. *Nat Commun* **13**,
5751 (2022).
8.   T. A. Cameron, W. Margolin, Insights into the assembly and regulation of the bacterial
divisome. *Nat Rev Microbiol* 10.1038/s41579-023-00942-x (2023).
9.   M. J. Tsang, T. G. Bernhardt, Guiding divisome assembly and controlling its activity. *Curr*
Opin Microbiol **24**, 60-65 (2015).
10.   M. J. Tsang, A. A. Yakhnina, T. G. Bernhardt, NlpD links cell wall remodeling and outer
membrane invagination during cytokinesis in Escherichia coli. *PLoS Genet* **13**, e1006888
(2017).
11.   T. Uehara, T. G. Bernhardt, More than just lysins: peptidoglycan hydrolases tailor the cell
wall. *Curr Opin Microbiol* **14**, 698-703 (2011).
12.   T. Uehara, K. R. Parzych, T. Dinh, T. G. Bernhardt, Daughter cell separation is controlled
by cytokinetic ring-activated cell wall hydrolysis. *EMBO J* **29**, 1412-1422 (2010).
13.   M. Alcorlo, S. Martinez-Caballero, R. Molina, J. A. Hermoso, Regulation of Lytic
Machineries by the FtsEX Complex in the Bacterial Divisome. *Subcell Biochem* **99**, 285-315
(2022).
14.   S. Pichoff, S. Du, J. Lutkenhaus, Roles of FtsEX in cell division. *Res Microbiol* **170**, 374-380
(2019).

15.   S. Du, S. Pichoff, J. Lutkenhaus, Roles of ATP Hydrolysis by FtsEX and Interaction with FtsA
in Regulation of Septal Peptidoglycan Synthesis and Hydrolysis. *mBio* **11** (2020).
16.   D. C. Yang *et al.*, An ATP-binding cassette transporter-like complex governs cell-wall
hydrolysis at the bacterial cytokinetic ring. *Proc Natl Acad Sci U S A* **108**, E1052-1060
(2011).
17.   S. Du, S. Pichoff, J. Lutkenhaus, FtsEX acts on FtsA to regulate divisome assembly and
activity. *Proc Natl Acad Sci U S A* **113**, E5052-5061 (2016).
18.   S. Du, W. Henke, S. Pichoff, J. Lutkenhaus, How FtsEX localizes to the Z ring and interacts
with FtsA to regulate cell division. *Mol Microbiol* **112**, 881-895 (2019).
19.   S. J. Arends, R. J. Kustusch, D. S. Weiss, ATP-binding site lesions in FtsE impair cell division.
J Bacteriol **191**, 3772-3784 (2009).
20.   M. Reddy, Role of FtsEX in cell division of Escherichia coli: viability of ftsEX mutants is
dependent on functional SufI or high osmotic strength. *J Bacteriol* **189**, 98-108 (2007).
21.   K. L. Schmidt *et al.*, A predicted ABC transporter, FtsEX, is needed for cell division in
Escherichia coli. *J Bacteriol* **186**, 785-793 (2004).
22.   J. Cook *et al.*, Insights into bacterial cell division from a structure of EnvC bound to the
FtsX periplasmic domain. *Proc Natl Acad Sci U S A* **117**, 28355-28365 (2020).
23.   D. Mavrici *et al.*, Mycobacterium tuberculosis FtsX extracellular domain activates the
peptidoglycan hydrolase, RipC. *Proc Natl Acad Sci U S A* **111**, 8037-8042 (2014).
24.   B. E. Rued *et al.*, Structure of the Large Extracellular Loop of FtsX and Its Interaction with
the Essential Peptidoglycan Hydrolase PcsB in Streptococcus pneumoniae. *mBio* **10**
(2019).
25.   Y. Fu, K. E. Bruce, B. Rued, M. E. Winkler, D. P. Giedroc, 1H, 13C, 15N resonance
assignments of the extracellular loop 1 domain (ECL1) of Streptococcus pneumoniae D39
FtsX, an essential cell division protein. *Biomol NMR Assign* **10**, 89-92 (2016).
26.   X. Xu *et al.*, Mechanistic insights into the regulation of cell wall hydrolysis by FtsEX and
EnvC at the bacterial division site. *Proc Natl Acad Sci U S A* **120**, e2301897120 (2023).
27.   B. D. Corbin, Y. Wang, T. K. Beuria, W. Margolin, Interaction between cell division proteins
FtsE and FtsZ. *J Bacteriol* **189**, 3026-3035 (2007).
28.   X. Ou *et al.*, ATP Can Efficiently Stabilize Protein through a Unique Mechanism. *JACS Au* **1**,
1766-1777 (2021).
29.   S. Sridharan *et al.*, Proteome-wide solubility and thermal stability profiling reveals distinct
regulatory roles for ATP. *Nat Commun* **10**, 1155 (2019).
30.   S. C. Lee *et al.*, Nano-encapsulated Escherichia coli Divisome Anchor ZipA, and in Complex
with FtsZ. *Sci Rep* **9**, 18712 (2019).
31.   S. C. Lee *et al.*, A method for detergent-free isolation of membrane proteins in their local
lipid environment. *Nat Protoc* **11**, 1149-1162 (2016).
32.   E. Godino, C. Danelon, Gene-Directed FtsZ Ring Assembly Generates Constricted
Liposomes with Stable Membrane Necks. *Adv Biol (Weinh)* **7**, e2200172 (2023).
33.   C. Ortiz, P. Natale, L. Cueto, M. Vicente, The keepers of the ring: regulators of FtsZ
assembly. *FEMS Microbiol Rev* **40**, 57-67 (2016).
34.   P. Szwedziak, Q. Wang, T. A. Bharat, M. Tsim, J. Lowe, Architecture of the ring formed by
the tubulin homologue FtsZ in bacterial cell division. *Elife* **3**, e04601 (2014).
35.   S. Rueda, M. Vicente, J. Mingorance, Concentration and assembly of the division ring
proteins FtsZ, FtsA, and ZipA during the Escherichia coli cell cycle. *J Bacteriol* **185**, 3344-
3351 (2003).
36.   H. P. Erickson, M. Osawa, FtsZ Constriction Force - Curved Protofilaments Bending
Membranes. *Subcell Biochem* **84**, 139-160 (2017).

37. M. L. Carlson *et al.*, The Peptidisc, a simple method for stabilizing membrane proteins in
detergent-free solution. *Elife* **7** (2018).
38. G. Angiulli *et al.*, New approach for membrane protein reconstitution into peptidiscs and
basis for their adaptability to different proteins. *Elife* **9** (2020).
39. P. A. Lanzetta, L. J. Alvarez, P. S. Reinach, O. A. Candia, An improved assay for nanomole
amounts of inorganic phosphate. *Anal Biochem* **100**, 95-97 (1979).
40. D. N. Mastronarde, Automated electron microscope tomography using robust prediction
of specimen movements. *J Struct Biol* **152**, 36-51 (2005).
41. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
42. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).
43. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216-221 (2015).
44. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612 (2004).
45. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr
D Biol Crystallogr* **60**, 2126-2132 (2004).
46. P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
47. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).

Figures

Figure 1. Biochemical reconstitution of *Eco*FtsEX and *Eco*FtsEX/EnvC complexes. (A)
Evaluation of ATP's impact on FtsEX and FtsEX/EnvC complexes. Purified FtsEX and FtsEX/EnvC
complexes undergo precipitation in the absence of ATP but form stable complexes when ATP is
present during purification. S: Supernatant, P: Precipitate. **(B)** Representative negative staining

image and 2D average of FtsEX samples purified in the presence of ATP, with FtsEX highlighted

in green. The FtsE subunit, Transmembrane Domain (TMD), and Periplasmic Domain (PLD) of

FtsX are clearly visible. **(C)** Assessment of ATP and ATP analogs' influence on the stability of FtsEX

complex formation. ATP and ATP analogs contribute to the stabilization of the FtsEX complex. S:

Supernatant, P: Precipitate. (**D**) Assessment of ATPase activity in FtsEX and FtsEX mutants, with

and without EnvC. Notably, the FtsE^{E163Q}X mutants exhibit significantly reduced ATPase activity.

Data represent an average of three replicates, with error bars indicating mean ± SD. Statistical

analysis performed using a two-tailed unpaired t-test; ***P < 0.0005. (**E**)**.** Pull-down study illustrating

EnvC binding to FtsEX or its ATPase mutant in the presence or absence of ATP or ATP analogs.

The initial FtsEX or FtsE^{E163Q}X added to the resin contains 2 mM ATP, the ATP was subsequently

removed by the wash buffer. (**F**) ATPase activity comparison between FtsEX and the FtsEX/EnvC

complex in peptidiscs. Data represent an average of three replicates, with error bars indicating

mean ± SD.

Figure 2. Cryo-EM study of *Eco*FtsEX complex in peptidiscs. (**A**) Topological diagram of FtsEX,

highlighting the α helices represented as cylinders and β sheets depicted as arrows. Color scheme:

FtsE (magenta and dark green), FtsX (cornflower blue), Porter of PLD$_{FtsX}$ (yellow), and X-lobe of

PLD$_{FtsX}$ (green). CH: coupling helix, EH: elbow helix. (**B**) Left and middle panels depict the side and

front views of the density map of FtsEX, with ATPγS displayed in the front view at 90%

transparency. Ribbon representation of WT FtsEX in the presence of ATPγS with EM density. Color

scheme: FtsE (magenta and dark green), FtsX (cornflower blue and rosy brown), Porter of PCD_{FtsX}

(yellow), and X-lobe of PLD_{FtsX} (green). EH: elbow helix. (**C**) Left panel presents a rainbow

representation of the PLD domain of FtsX, transitioning from blue (N-terminal) to red (C-terminal).

Right panel shows a front view overlay of two PLD domains. (**D**) Surface representations of the

PLD domains in a top-view configuration, with the Porter of the PLD colored in yellow and the X-

lobe in green. In *Eco*FtsEX, the Porter and X-lobe are positioned facing each other.

Figure 3. Cryo-EM structure of the EnvC-bound *Eco*FtsEX complex. (**A**) Domain arrangements
and the overall structure of FtsEX/EnvC. Left and middle panels display side and front views of the
cryo-EM density maps for FtsEX-EnvC complex in the presence of ATP. The right panel presents
a ribbon representation of the structural model overlaid with the EM density. In the front-view, FtsE
is depicted at 90% transparency to reveal the ATP-binding site. Color scheme of EnvC: α1 (plum),
lip (purple), α2 (red), RH1, RH2, and LytM domains (grey). (**B**) EnvC binding to the PLD domain of
FtsEX is depicted with ribbon representations. Color scheme: FtsX (cornflower blue), FtsE
(magenta/dark green), and EnvC (red). (**C**) **&** (**D**). Top-down views provide a detailed depiction of
the interaction between EnvC and FtsEX.

Figure 4. Identification of key loops for EnvC binding in the *Eco*FtsEX/EnvC Complex. (**A**)
Overlay of *Eco*FtsEX and *Eco*FtsEX/EnvC complexes, with a focus on the conformational changes
in the PLD domain induced by EnvC binding. The overall overlay of *Eco*FtsEX and *Eco*FtsEX/EnvC
is depicted in ribbon form. Color scheme: FtsE (magenta and dark green), FtsX (cornflower blue
and rosy brown), Porter of PLD$_{FtsX}$ (yellow), and X-lobe of PLD$_{FtsX}$ (green). The TMD and FtsE
regions are set at 60% transparency. (**B**) Depiction of the conformational changes in the PLD
domains upon EnvC binding. (**C**) Side and front views illustrating the key loops involved in EnvC
binding within the *Eco*FtsEX/EnvC complex. Key loops of FtsX, highlighted in blue, are indicated
by arrows and labeled. (**D**) Side-view displaying the binding details between EnvC and flexible
loops located in the X-lobe (G157-L171). (**E**)**,** (**F**)**, and** (**G**) Top-down views demonstrating the
binding details of EnvC and flexible loops positioned between TM1 and PLD (Y105-Q109) (**E**), PLD
and TM2 (D208-L214) (**F**), and TM3 and TM4 (Q307-T311) (**G**).

Figure 5. Biochemical reconstitution and negative-staining EM study of the AmiB-Bound
_Eco_FtsEX/EnvC Complex. (**A**) Pull-down study illustrating the interaction between AmiB and
FtsEX-EnvC or its ATPase mutant, in the presence or absence of ATP or its analogs. The initial
FtsEX or FtsE^{E163Q}X added to the resin contains 2 mM ATP, the ATP was subsequently removed
by the wash buffer. (**B**) Evaluation of ATPase activity in FtsEX, FtsEX/EnvC, and FtsEX/EnvC/AmiB
complexes. Notably, the binding of AmiB does not significantly impact the ATPase activity of
FtsEX/EnvC. Data represent an average of three replicates, with error bars indicating mean ± SD.
Statistical analysis performed using a two-tailed unpaired t-test; ***$P < 0.0005$. (**C**) SEC profile and
SDS-PAGE gel of purified FtsEX/EnvC/AmiB complex reconstituted in peptidiscs. (**D**) Negative-
staining micrograph and 2D average image of _Eco_FtsEX/EnvC/AmiB. (**E**) Cryo-EM density map of
_Pae_FtsEX/EnvC/AmiB.

Figure 6. A working model illustrating FtsEX complex formation, recruitment of EnvC and

AmiB, and the temporal regulation of PG cleavage activation in Gram-negative bacterial

systems. Initially, the presence of physiological ATP stabilizes a fully functional FtsEX complex.

Subsequently, the asymmetrical and inactive EnvC approaches the relatively flexible PLD domain

of FtsEX. Upon complete recognition by FtsEX, the CCD domain of recruited EnvC undergoes a

conformational change to a symmetrical state, involving compression of the coiled-coil domain,

facilitating the release of the LytM domain. Interaction with EnvC activates FtsEX's ATPase activity.

The liberated LytM recruits amidases through direct interaction with the catalytic domain's blocking

helices, leading to the removal of the obstructing helix from the catalytic site and activation of PG

cleavage. ATP hydrolysis likely triggers the release of EnvC and AmiB, allowing FtsEX to return to

its normal ABC transporter state with a low basal ATPase activity. The potential interaction between

FtsEX and the constriction ring component FtsA/FtsZ precisely locates the PG-cleaving complex

at the septum for temporal regulation of PG cleavage.

819