Visualization of DNA Methylation in Cell Cycle Genes during iPSC Cardiac Differentiation

Ning Li[1, 2], Ba Thong Nguyen[3,4], Zhenhe Zhang[1], Yiqiang Zhang[1, 3, 4*]

From [1] Division of Cardiology, Department of Medicine, University of Washington, Seattle, WA 98109, USA; [2] State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China; [3]Department of Anatomy, Biochemistry and Physiology, and [4]Center for Cardiovascular Research, John A. Burns School of Medicine, University of Hawaii at Manoa, Honolulu, HI 96813, USA.

* Correspondence should be addressed to: Yiqiang Zhang, PhD (yiqiang.zhang@hawaii.edu).

ABSTRACT

Epigenetic DNA methylation is an essential mechanism controlling gene expression and cellular

function. Endpoint analyses with conventional assays have generated significant insights into

static states of DNA methylation, but were short of visualizing the dynamics of epigenetic

regulation. Here, we adopted a genomic DNA methylation reporter (GMR) system to track the

changes of DNA methylation during cardiac differentiation. The promoter region of *Cdk1*

(Cyclin-dependent kinase 1) or *Sox2* (SRY-Box Transcription Factor 2) gene was cloned

upstream of the small nuclear ribonucleoprotein polypeptide N (Snrpn) minimal promoter

followed by a fluorescent reporter gene. Mouse induced pluripotent stem cells (iPSCs) carrying

Sox2 GMR rapidly lost fluorescent reporter signal upon the induction of differentiation. Cdk1

GMR reporter signal was strong in undifferentiated iPSCs, and gradually decreased during

directed cardiomyocyte (CM) differentiation. The present study demonstrated the dynamic DNA

methylation along the course of cell cycle withdrawal during CM differentiation. The GMR

reporter system can be instrumental to monitor real-time epigenetic DNA modification at single-

cell resolution.

KEYWORDS: Epigenetics; DNA Methylation; Cardiac Differentiation; Cell Cycle;

Genomics Methylation Reporter; Pyrosequencing; single-cell imaging; flow cytometry

INTRODUCTION

Epigenetic DNA methylation is highly associated with gene expressions in development and cellular functions such as cell cycle progression and differentiation. Methylation status of gDNA can be determined by a number of available methods, such as gene-specific analysis using Pyrosequencing and PCR quantification after bisulfite conversion, or digestion-based assay followed by PCR or qPCR, and whole-genome sequencing on similarly processed samples.[1] The mounting data in DNA methylation studies have generated significant insights into static states of DNA methylation but were short of depicting epigenetic perturbations in a real-time manner. A live-cell gDNA methylation reporter system is essential in visualizing dynamic genomic methylation states, such as in cardiac differentiation of induced pluripotent stem cells (iPSCs).

Mouse iPSCs have been widely used to study lineage differentiation, which is often associated with a perturbation in the cell cycle activity, but the exact temporal dynamics remained unclear for various lineages. Cardiomyocytes (CMs) withdraw from activity cell cycling shortly after birth,[2] but dynamic epigenetic DNA methylation during cardiac differentiation remained to be visualized.

The small nuclear ribonucleoprotein polypeptide N (*Snrpn*) gene is a maternally imprinted gene that is sensitive to DNA methylation of adjacent genomic regions.[3] *Snrpn* mini promoter region contains the conserved elements between human and mouse, with the endogenous imprinted differentially-methylated region (DMR). The *Snrpn* mini promoter can be used as a methylation sensor for real-time reporting of the DNA methylation state of adjacent sequences such as CpG regions of a particular gene.[4] In this study, we adopted this Snrpn mini-promoter-based

Genomics DNA methylation reporter (GMR) to track DNA methylation states of stemness and

cell cycle genes during CM differentiation from mouse iPSCs.

MATERIALS AND METHODS

Mouse iPSCs cell culture – Mouse iPSCs were maintained in ESC medium with 2iL [MEK

inhibitor (MEKi), GSK3 inhibitor (CHIR) and leukemia inhibitory factor (LIF)]. Stem Cell

Medium was made of DMEM (Gibco; # 11995065) supplemented with 20% FBS (Fetal Bovine

Serum; Gibco; #10082147), 1% penicillin/streptomycin (Gibco; #15070063), 1% NEAA (Non-

Essential Amino Acid; Gibco; #11140050) and 0.1 mM beta-mercaptoethanol (Sigma-Aldrich).

The medium was filtered through a Millipore Stericup 0.22 mm filtration system, and then added

with 2iL [10^3 Units/mL LIF (Millipore; #ESG1107), 1.0 μM MEKi (Stemgent; #PD0325901)

and 1.5 μM CHIR (Selleck Chemicals; #CHIR-99021)] before use in mouse iPSCs. All medium

was prepared under sterile conditions and cells were cultured in a incubator at 37°C temperature

and 5% CO_2.

Construction of cell cycle GMR, and generation of reporter iPSC cell lines – Sox2se-CPG-

TV-dTomatto and Gapdh-CpG-TV-GFP plasmids (Addgene plasmid # 70155, and 70148) were

a gift from Dr. Rudolf Jaenisch.[4] The promoter of mouse *Cdk1* gene that includes 29 CpG was

cloned from wildtype C56/Bl6j mouse gDNA, using a primer pair: Cdk1-pro-F (5'-

ACG*GGTACC*AGCAATAGCCAGGTGGTGGTG-3') and Cdk1-pro-R (5'-

GAC*CAATTG*CATAAGGGATCCCGGACTTAG-3'). Then the PCR amplicon of *Cdk1* CpG

was digested by KpnI and MfeI (New England Biolabs) to remove the overhangs. The resultant

fragment was ligated to the Snrpn-dTomato backbone from Sox2se-CPG-TV, to generate the

Cdk1-Snrpn-dTomato plasmid, which was then amplified in MAX Efficiency DH5α Competent

Cells (Invitrogen).

To generate *Sox2* and *Cdk1* GMR reporter cell lines, we transfected Sox2se-CPG-TV-dTomato

and Cdk1-Snrpn-dTomato plasmids into mouse iPSCs using Lipofectamine 2000 Transfection

Reagent (Invitrogen) according to the provider's protocol. Twenty-four hours following

transfection, cells were treated with puromycin (0.75 µg/ml) for 3 days. Then the puromycin-

resistant cells carrying GMR genes were harvested and sorted with a BD Aria II cell sorter for

single-cell tdTomato expression. Cells were cultured in 96-well plates containing irradiated

mouse embryonic fibroblasts (iMEFs) feeder layer. Single colonies were passaged and used in

subsequent analyses.

Directed cardiomyocyte differentiation from mouse iPSCs – For differentiation assay, we

adopted Yashiro's method with some modifications.[5] Mouse Sox2 and Cdk1 GMR iPSC lines

were maintained in Stem Cell Medium with 2iL for three days, then harvested by trypsin

dissociation, and suspended in 2iL Stem Cell Medium medium, followed by incubation in a

incubator (5% CO_2, 37 °C) for 30 minutes to remove iMEFs. Cells were washed twice with

DPBS (Dulbecco's phosphate buffered saline, Gibco) before medium replacement. The iPSCs

were re-seeded into gelatin-coated 24-well plates at a density of 120,000 cells/well in 2iL ESC

medium. After 8 hours, medium was changed to Serum-free (SF) differentiation medium

containing the followings: 75% Iscove's modified Dulbecco's medium (IMDM, Gibco), 25%

Ham's F12 Nutrient Mix (Gibco), supplemented with 0.5% N-2 supplement (Gibco), 0.5% B-27

supplement (Gibco), 0.05% bovine serum albumin (BSA, Fisher scientific), 1% Pen-Strep

(Gibco), 2 mM L-glutamine (Gibco), 0.5 mM L-ascorbic acid (Sigma), and 0.45 mM 1-

Thioglycerol (Sigma); and cells were cultured for 2 days. It was counted as Day 0 when medium

was changed to SF medium with 8 ng/mL recombinant Activin A (R&D Systems), 5 ng/mL

recombinant human VEGF 165 Protein (R&D Systems) and 0.25 ng/mL recombinant human

BMP-4 Protein (R&D Systems).Cells were cultured for 28 hours; and then the medium was

changed to StemPro-34 Serum-free Medium (SFM, Gibco) supplemented with 2 mM L-

glutamine, 0.5 mM L-ascorbic acid (Sigma), 5 ng/ml recombinant human VEGF 165 protein

(R&D Systems), 10 ng/ml human bFGF (Invitrogen), and 50 ng/ml recombinant human FGF-10

protein, and cells were cultured for another 3.5 days. Subsequently, cells were harvested, and

passaged into gelatin-coated 24-well plates in 500k cells/well density in StemPro-34 SF

supplemented medium. After 2 days, RPMI 1640 medium (with L-glutamine, Gibco) with B-27

supplement (minus vitamin A, Gibco) was used for further cell culture of the differentiated

myocytes.

Flow cytometry – Mouse iPSCs and differentiated cells in tissue cultures were dissociated by

incubation with Trypsin-EDTA (0.25%, Gibco) and then fixed with 4% paraformaldehyde and

0.2% saponin in PBS (Gibco). The fixed cells could be stored in 10% DMSO in FBS at -80°C

for future use. Mouse iPSCs were stained with Oct4-AF488 (Millipore MAB4419A4) to verify

the stemness gene. The cells harvested on different days of cardiac differentiation were stained

with cTnT antibody (Troponin T, mouse monoclonal antibody, Thermo Fisher Scientific, MS-

295-P, 1:200) and then with donkey anti-mouse IgG (H+L) secondary antibody (Alexa Fluor

647; Thermo Fisher; A31571; 1:1000). Immuno stained cells were analyzed on an FACScan

Canto RUO flow cytometer (BD) and the result data were analyzed with FlowJo software

(FlowJo; Ashland, OR).[6-8]

Statistics – After the derivation of GMR reporter clones, experiments were carried out with at

least biologically replicated, unless otherwise specified. Group data are expressed as Mean±

SEM unless otherwise specified. Difference between groups was assessed by t test, with a P

value <0.05 consider significant different.

RESULTS

Mouse iPSC with GMR for Sox2 and Cdk1 – To test the utility of Snrpn mini promoter's

DNA methylation sensing function, we generated mouse iPSC lines carrying Sox2-GMR

construct that composites of Sox2 super enhancer CpG island region fused with the minimal

promoter of Snrpn, followed by dTomato reporter. iPSCs transfected with Sox2-GMR were

subjected to puromycin-resistant selection. Cells with high expression of dTomato reporter

signals were collected by fluorescence-activated cell sorting (FACS), and single clones were

expanded, among which clone C7 was used for subsequent experiments (**Figure 1**). Similar

approaches were used to generate mouse Cdk1-GMR iPSC lines, among which clone B9 was

used for subsequent analyses. The expression of stemness protein such as Oct4 or Nano, Sox2 in

GMR lines remained at comparable to wildtype mouse iPSC as assessed by immunostaining and

flow cytometry. Fluorescent reporter signals in both Sox2 and Cdk1 GMR lines were readily

detected with epifluorescent microscopy imaging.

Diminished reporter signals in cells differentiated from Cdk1 and Sox2 GMR iPSCs – Cell

differentiation is accompanied with epigenetic reprogramming that sets the landscape for

transcriptional activity for lineage specification, cell identity and functions. As a proof of

concept to the utility of GMR, we examined the reporter signal and cardiac gene expression in

cells differentiated from the GMR iPSCs lines. By day 2.5 of directed cardiac differentiation,

dTomato reporter signals in both Sox2 GMR and Cdk1 GMR cells reduced remarkably (**Figure**

2A). By day 6.5 of differentiation, dTomato signal was minimally detected in ~2.08% in Sox2-

GMR cells, consistent with the loss of stemness in differentiated cells assessed by flow

cytometry on the gene expression of Sox2 and Oct4. dTomato signal in Cdk1 GMR cells also

reduced significantly in differentiated cells (6.5 days) compared to iPSCs (64.3%±7.1%

compared to 95.5%±1.5% (n=3, p<0.01). By 3 weeks, there is barely detectably dTomato signal

from Cdk1 GMR-derived cells that were mostly myocytes expressing cTnT as detected by

immunostaining and flow cytometry (**Figure 2B, C**).

Correlations between GMR signals and gene expression – To determine the correlation

between GMR signal in live cells and gene expression during cardiac differentiation, we

performed RT-qPCR on cells differentiated from Cdk1 GMR iPSCs. After 4.5 days of cardiac

differentiation, *Tnnt2* was expressed and remarkably increased in myocytes of later stage (13.5

days) as compared to earlier stages. On the other hand, the transcript *Cdk1* increased in 4.5 days

differentiated cells compared to Cdk1 GMR iPSC, then tapered: *Cdk1* in differentiated cells at

day 13.5 was reduced to only 25% level of that in undifferentiated iPSC (**Figure 3**).

DISCUSSION

Epigenetic DNA and histone modifications, and subsequently the changes in chromatin

accessibility, are the critical mechanisms controlling gene expression, thereby affecting cellular

phenotype and function, such as differentiation and proliferation.[6,9-15] Several approaches can be

used to reveal the status of DNA methylation; for example, fluorescent proteins (FPs) in

conjugation with methyl-CpG-binding domain (MBD) or Synthetic-molecule/protein hybrid

probes,[16] modular fluorescence complementation-based epigenetic biosensors that combine

engineered DNA-binding proteins with domains recognizing defined epigenetic marks, both

fused to non-fluorescent fragments of a fluorescent protein.[17] The live-cell reporter system

leveraging the mini-promoter of imprinted *Snrpn* gene to visualize the CpG island regions of a

target gene represents unique and highly specific system with detail temporal and spatial

resolutions critical in the study of DNA methylation dynamics.[4]

In this study, we confirmed the fidelity of CpG-*Snrpn* based GMR approach in mouse iPSC and

differentiated myocytes. It is important to note that methylation status of genes that have no or

short CpG regions may not be reflective in this GMR system. C2c12 cell lines carrying *Gapdh*

GMR (GFP) reporter and Sox2SE GMR (tdTomato) were also used to verify the expected GMR

signals. In addition, there was no reporter signals in cells transfected with pre-methylated Sox2

or Gapdh GMR gene constructs (Data not shown). Live cell imaging data showed that the

positive GMR reporter signal reflects well the transcript expression of corresponding gene.

Differentiated cells, like cardiomyocytes, have slower proliferation and the cell cycle can be

arrested shortly after birth.[2] Our data show that during early differentiation (4.5 days), there was

an increased expression of *Cdk1* then reduced significantly in more mature cardiomyocytes (e.g.,

13.5 days). It may be plausible that there was a fast proliferative phase right after transition to

cardiac mersoderm lineage, leading to the higher *Cdk1* transcript expression, and the changes in

transcription activity may be delayed behind the GMR reporter signal for that short period.[18] It

is worth noting that Cdk1 itself can affect histone epigenetic landscape as shown in mouse

embryonic stem cells.[19] Combination with other approaches will likely further delineate the

epigenetic-transcriptional regulations in cell differentiation and cell cycle regulation. A caveat of

using this GMR is that it is a "negative-positive" system in which the reporter signal represents

hypomethylation rather than hypermethylation.

In summary, the *Snrpn* mini-promoter-based GMR system,[4] such as for *Cdk1* and other cell cycle

genes, can be instrumental to report dynamic DNA methylation states during cell differentiation

and cell cycling.

Declaration of Competing Interest

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Y.Z. conceived the study. Y.Z. and N.L. designed and performed experiments, collected and

analyzed data. B.T.N. contributed to gene expression analysis. Z.Z. conducted experiments on

cell culture. N.L. and Y.Z. wrote the manuscript. All co-authors provided their edits to the

manuscript.

Acknowledgments

The authors thank Yun-Yu Wu and Lovina Abdi for technical supports, and Dr. W. Robb

MacLellan for assistantce in discussing the experiments. The work was in part supported by

research grants from American Heart Association (18IPA34110210 to Y.Z.) and the National

Institute of Health (1R56HL131898 to Y.Z.) and University of Washington Institute for Stem

Cells and Regenerative Medicine (Innovation Pilot Awards to Y.Z.).

REFERENCES:

1. Ammerpohl O, Martin-Subero JI, Richter J, Vater I, Siebert R. Hunting for the 5th base: Techniques for analyzing DNA methylation. *Biochim Biophys Acta*. 2009;1790:847-862. doi: 10.1016/j.bbagen.2009.02.001

2. Zhang Y, Mignone J, MacLellan WR. Cardiac Regeneration and Stem Cells. *Physiol Rev*. 2015;95:1189-1204.

3. Shemer R, Birger Y, Riggs AD, Razin A. Structure of the imprinted mouse Snrpn gene and establishment of its parental-specific methylation pattern. *ProcNatlAcadSciUSA*. 1997;94:10267-10272.

4. Stelzer Y, Shivalila CS, Soldner F, Markoulaki S, Jaenisch R. Tracing dynamic changes of DNA methylation at single-cell resolution. *Cell*. 2015;163:218-229.

5. Kokkinopoulos I, Ishida H, Saba R, Coppen S, Suzuki K, Yashiro K. Cardiomyocyte differentiation from mouse embryonic stem cells using a simple and defined protocol. *Dev Dyn*. 2016;245:157-165. doi: 10.1002/dvdy.24366

6. Zhang Y, Gago-Lopez N, Li N, Zhang Z, Alver N, Liu Y, Martinson AM, Mehri A, MacLellan WR. Single-cell imaging and transcriptomic analyses of endogenous cardiomyocyte dedifferentiation and cycling. *Cell Discov*. 2019;5:30.

7. Gago-Lopez N, Awaji O, Zhang Y, Ko C, Nsair A, Liem D, Stempien-Otero A, MacLellan WR. THY-1 Receptor Expression Differentiates Cardiosphere-Derived Cells with Divergent Cardiogenic Differentiation Potential. *Stem Cell Reports*. 2014;2:576-591.

8.   Zhang Y, Matsushita N, Eigler T, Marban E. Targeted microRNA interference promotes

postnatal cardiac cell cycle re-entry. *JRegenerative Medicine*. 2013;2:doi:10.4172/2325-

9620.1000108.

9.   Radak M, Fallahi H. The Epigenetic Regulation of Quiescent in Stem Cells. *Glob Med*

Genet. 2023;10:339-344. doi: 10.1055/s-0043-1777072

10.   Chen X, Chakravarty T, Zhang Y, Li X, Zhong JF, Wang C. Single-cell transcriptome

and epigenomic reprogramming of cardiomyocyte-derived cardiac progenitor cells.

SciData. 2016;3:160079.

11.   Zhang Y, Zhong JF, Qiu H, Robb MW, Marban E, Wang C. Epigenomic Reprogramming

of Adult Cardiomyocyte-Derived Cardiac Progenitor Cells. *SciRep*. 2015;5:17686.

12.   Vallaster M, Vallaster CD, Wu SM. Epigenetic mechanisms in cardiac development and

disease. *Acta BiochimBiophysSin(Shanghai)*. 2012;44:92-102.

13.   McKinsey TA, Vondriska TM, Wang Y. Epigenomic regulation of heart failure:

integrating histone marks, long noncoding RNAs, and chromatin architecture. *F1000Res*.

2018;7.

14.   Klemm SL, Shipony Z, Greenleaf WJ. Chromatin accessibility and the regulatory

epigenome. *Nat Rev Genet*. 2019;20:207-220. doi: 10.1038/s41576-018-0089-8

15.   Ailsa M, Elisa De F, Alice F, Christine SF, Sarah JR, Kathryn M, Joe B, Barts Pancreas

Tissue B, Emma MW, Noel GM, et al. Developmentally dynamic changes in DNA

methylation in the human pancreas. *bioRxiv*. 2023:2023.2010.2019.563104. doi:

10.1101/2023.10.19.563104

16.   Kumar N, Hori Y, Kikuchi K. Live-Cell Imaging of DNA Methylation Based on

Synthetic-Molecule/Protein Hybrid Probe. *Chem Rec*. 2018;18:1672-1680. doi:

10.1002/tcr.201800039

17.   Lungu C, Pinter S, Broche J, Rathert P, Jeltsch A. Modular fluorescence

complementation sensors for live cell detection of epigenetic signals at endogenous

genomic sites. *Nature Communications*. 2017;8:649. doi: 10.1038/s41467-017-00457-z

18.   Soonpaa MH, Field LJ. Survey of studies examining mammalian cardiomyocyte DNA

synthesis. *Circ Res*. 1998;83:15-26. doi: 10.1161/01.res.83.1.15

19.   Michowski W, Chick JM, Chu C, Kolodziejczyk A, Wang Y, Suski JM, Abraham B,

Anders L, Day D, Dunkl LM, et al. Cdk1 Controls Global Epigenetic Landscape in

Embryonic Stem Cells. *Mol Cell*. 2020;78:459-476.e413. doi:

10.1016/j.molcel.2020.03.010

Figures and Figure Legends

Figure 1.

Figure 1. Generation of mouse GMR iPSC lines. A, Gene construction of Snrpn mini-
promoter based genomic DNA methylation reporter system. **B,** Microscopy images of wildtype
iPSC (control) and GMR iPSC lines for carrying GMR for Sox2 super enhancer or Cdk1
promoter's CpG region.

Figure 2.

Figure 2. Dynamic reporter signal during cardiac differentiation of GMR iPSCs. A,

Microscopy images of Sox2-GMR and Cdk1-GMR iPSCs (day 0) and differentiated cells. **B,**

Flow cytometry dot plots showing the signal of GMR reporter dTomato and cardiomyocyte

marker cTnT (cardiac troponin T). **C**, Cell population quantified by the expression of GMR

signal dTomato of cardiac myocyte marker cTnT.

Figure 3.

Figure 3. Expression of Cdk1 and Tnnt2 (cTnT) gene during cardiac differentiation from

the Cdk1 GMR iPSCs. Relative quantification of mRNA as normalized to *Gapdh*. *** and ###,

$p<0.05$ vs 0d; n=3.