Dynamic Visualization of DNA Methylation in Cell Cycle Genes during iPSC Cardiac

Differentiation

Ning Li[1,2], Ba Thong Nguyen[3,4]*, Zhenhe Zhang[1]*, W. Robb MacLellan,[1] Yiqiang Zhang[1,3,4]¶

[1] Division of Cardiology, Department of Medicine, University of Washington, Seattle, WA

98109, USA

[2] State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant

Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China

[3] Department of Anatomy, Biochemistry and Physiology, John A. Burns School of Medicine,

University of Hawaii at Manoa, Honolulu, HI 96813, USA.

[4] Center for Cardiovascular Research, John A. Burns School of Medicine, University of Hawaii

at Manoa, Honolulu, HI 96813, USA.

* Equal contribution.

¶ Author for Correspondence: Yiqiang Zhang, PhD (yiqiang.zhang@hawaii.edu).

ABSTRACT

Background: Epigenetic DNA methylation is an essential mechanism controlling gene

expression and cellular function. Existing analyses with conventional assays have generated

significant insights into static states of DNA methylation, but were unable to visualize the

dynamics of epigenetic regulation. **Aim:** We utilized a genomic DNA methylation reporter

(GMR) system to track changes in DNA methylation during cardiac differentiation. **Methods**

and Results: The promoter region of *Cdk1* (Cyclin-dependent kinase 1) or *Sox2* (SRY-Box

Transcription Factor 2) gene was cloned upstream of the small nuclear ribonucleoprotein

polypeptide N (Snrpn) minimal promoter followed by a fluorescent reporter gene. Mouse

induced pluripotent stem cells (iPSCs) carrying Sox2 GMR rapidly lost fluorescent reporter

signal upon the induction of differentiation. Cdk1 GMR reporter signal was strong in

undifferentiated iPSCs, and gradually decreased during directed cardiomyocyte (CM)

differentiation. RT-qPCR and pyrosequencing demonstrated that the reduction of *Sox2* and *Cdk1*

was regulated by hypermethylation of their CpG regions during cardiac differentiation. The

present study demonstrated the dynamic DNA methylation along the course of cell cycle

withdrawal during CM differentiation. **Conclusion:** The GMR reporter system can be a useful

tool to monitor real-time epigenetic DNA modification at single-cell resolution.

KEYWORDS: cardiac differentiation; cell cycle; DNA methylation; epigenetics; GMR

(genomic DNA methylation reporter); pyrosequencing; iPSC (induced pluripotent stem cell);

single-cell imaging

INTRODUCTION

Epigenetic DNA methylation is highly associated with gene expression in development and cellular functions such as cell cycle progression and differentiation. Methylation status of gDNA can be assessed by a number of available methods, such as gene-specific analysis using Pyrosequencing and PCR quantification after bisulfite conversion, or digestion-based assay followed by PCR or qPCR, or whole-genome sequencing on similarly processed samples.[1] DNA methylation studies have generated significant insights into static states of DNA methylation but could not provide data on epigenetic perturbations in a real-time manner. A live-cell gDNA methylation reporter system is essential in visualizing dynamic genomic methylation states, such as in cardiac differentiation of induced pluripotent stem cells (iPSCs).

Mouse iPSCs have been widely used to study lineage differentiation, which is often associated with a perturbation in the cell cycle activity, but the exact temporal dynamics remained unclear for various lineages. Cardiomyocytes (CMs) withdraw from activity cell cycling shortly after birth,[2] but dynamic epigenetic DNA methylation during cardiac differentiation remained to be visualized.

The small nuclear ribonucleoprotein polypeptide N (*Snrpn*) gene is a maternally imprinted gene that is sensitive to DNA methylation of adjacent genomic regions.[3] *Snrpn* mini promoter region contains the conserved elements between human and mouse, with the endogenous imprinted differentially-methylated region (DMR). The *Snrpn* mini promoter can be used as a methylation sensor for real-time reporting of the DNA methylation state of adjacent sequences, such as CpG regions of a particular gene.[4] In this study, we adopted this Snrpn mini-promoter-based Genomics DNA Methylation Reporter (GMR) to track DNA methylation states of stemness gene

(Sox2; SRY-Box Transcription Factor 2) and cell cycle gene (Cdk1; cyclin-dependent kinase 1)

during CM differentiation from mouse iPSCs. Sox2 is a transcription factor that regulates the

pluripotency of stem cells. Its promoter region has a high density of CpG island (186 CpG sites;

1530 C+G counts), while the super enhancer region has a low density of CpG sequence. The

enhancer region is important for Sox2 mRNA and protein expression and the pluripotency of

embryonic stem cells.[5] In addition, developmental cardiomyocytes have a proliferative phase

followed by cell cycle withdrawal and terminal differentiation, which is in part dependent on

Cdk1. There are 29 CpGs in Cdk1's promoter region. Our study showed dynamic signals of

DNA methylation reporter in Cdk1 and Sox2 during mouse iPSC cardiac differentiation.

MATERIALS AND METHODS

Mouse iPSC Cell Culture

Mouse iPSCs were maintained in ESC medium with 2iL [MEK inhibitor (MEKi), GSK3

inhibitor (CHIR), and leukemia inhibitory factor (LIF)]. Stem Cell Medium was made of DMEM

(Gibco; # 11995065) supplemented with 20% FBS (Fetal Bovine Serum; Gibco; #10082147),

1% penicillin/streptomycin (Gibco; #15070063), 1% NEAA (Non-Essential Amino Acid; Gibco;

#11140050) and 0.1 mM beta-mercaptoethanol (Sigma-Aldrich). The medium was filtered

through a Millipore SteriCup 0.22 mm filtration system, and then added with 2iL [10^3 Units/mL

LIF (Millipore; #ESG1107), 1.0 µM MEKi (Stemgent; #PD0325901) and 1.5 µM CHIR (Selleck

Chemicals; #CHIR-99021)] before used in mouse iPSCs. All medium was prepared under sterile

conditions, and the cells were cultured in an incubator at 37°C and 5% CO_2.

Construction of Cell Cycle GMR, and Generation of Reporter iPSC Cell Lines

Sox2se-CPG-TV-dTomatto and Gapdh-CpG-TV-GFP plasmids (Addgene plasmid # 70155, and

70148) were a gift from Dr. Rudolf Jaenisch.[4] The promoter of mouse *Cdk1* gene that includes

29 CpG was cloned from wildtype C56/Bl6j mouse gDNA, using a primer pair: Cdk1-pro-F (5'-

ACG*GGTACC*AGCAATAGCCAGGTGGTGGTG-3') and Cdk1-pro-R (5'-

GAC*CAATTG*CATAAGGGATCCCGGACTTAG-3'). Then the PCR amplicon of *Cdk1* CpG

was digested by KpnI and MfeI (New England Biolabs) to remove the overhangs. The resultant

fragment was ligated to the Snrpn-dTomato backbone from Sox2se-CPG-TV, to generate the

Cdk1-Snrpn-dTomato plasmid, which was then amplified in MAX Efficiency DH5α Competent

Cells (Invitrogen).

To generate *Sox2* and *Cdk1* GMR reporter cell lines, we transfected Sox2se-CPG-TV-dTomato

and Cdk1-Snrpn-dTomato plasmids into mouse iPSCs using Lipofectamine 2000 Transfection

Reagent (Invitrogen) according to the provider's protocol. Twenty-four hours following

transfection, cells were treated with puromycin (0.75 µg/ml) for 3 days. Then the puromycin-

resistant cells carrying GMR genes were harvested and sorted with a BD Aria II cell sorter based

on single-cell tdTomato expression. Cells were cultured in 96-well plates containing irradiated

mouse embryonic fibroblasts (iMEFs) feeder layer. Single colonies were passaged and used in

subsequent analyses.

Directed Cardiomyocyte Differentiation from Mouse iPSCs

For cardiac differentiation assay, we adopted Yashiro's method with some modifications.[6] Mouse

Sox2 and Cdk1 GMR iPSC lines were maintained in Stem Cell Medium with 2iL for three days,

then harvested by trypsin dissociation and suspended in 2iL Stem Cell Medium, followed by

incubation in an incubator (5% CO_2, 37 °C) for 30 minutes to remove iMEFs. Cells were washed

twice with DPBS (Dulbecco's phosphate buffered saline, Gibco) before medium replacement.

The iPSCs were re-seeded into gelatin-coated 24-well plates at a density of 120,000 cells/well in

2iL ESC medium. After 8 hours, medium was changed to Serum-free (SF) differentiation

medium containing the followings: 75% Iscove's modified Dulbecco's medium (IMDM, Gibco),

25% Ham's F12 Nutrient Mix (Gibco), supplemented with 0.5% N-2 supplement (Gibco), 0.5%

B-27 supplement (Gibco), 0.05% bovine serum albumin (BSA, Fisher Scientific), 1% Pen-Strep

(Gibco), 2 mM L-glutamine (Gibco), 0.5 mM L-ascorbic acid (Sigma), and 0.45 mM 1-

Thioglycerol (Sigma); and cells were cultured for 2 days. It was counted as Day 0 when the

medium was changed to SF medium with 8 ng/mL recombinant Activin A (R&D Systems), 5

ng/mL recombinant human VEGF 165 Protein (R&D Systems) and 0.25 ng/mL recombinant

human BMP-4 Protein (R&D Systems). Cells were cultured for 28 hours; and then the medium

was changed to StemPro-34 Serum-free Medium (SFM; Gibco) supplemented with 2 mM L-

glutamine, 0.5 mM L-ascorbic acid (Sigma), 5 ng/ml recombinant human VEGF 165 protein

(R&D Systems), 10 ng/ml human bFGF (Invitrogen), and 50 ng/ml recombinant human FGF-10

protein, and cells were cultured for another 3.5 days. Subsequently, cells were harvested and

passaged into gelatin-coated 24-well plates in 500k cells/well density in StemPro-34 SF

supplemented medium. After 2 days, RPMI 1640 medium (with L-glutamine; Gibco) with B-27

Supplement (minus vitamin A; Gibco) was used for further cell culture of the differentiated

myocytes.

Flow Cytometry

Mouse iPSCs and differentiated cells in tissue cultures were dissociated by incubation with

Trypsin-EDTA (0.25%; Gibco) and then fixed with 4% paraformaldehyde and 0.2% saponin in

PBS (Gibco). The fixed cells could be stored in 10% DMSO in FBS at -80°C for future use.

Mouse iPSCs were stained with Oct4-AF488 (Millipore; # MAB4419A4) to verify the stemness

gene. The cells harvested on different days of cardiac differentiation were stained with mouse

anti-cardiac Troponin T (cTnT) antibody (Thermo Fisher Scientific; #MS-295-P; 1:200) and then

with Alexa Fluor 647-conjugated donkey anti-mouse IgG (H+L) secondary antibody (Thermo

Fisher; A31571; 1:1000). Immunostained cells were analyzed on a FACScan Canto RUO flow

cytometer (BD) and the data were analyzed with FlowJo software (FlowJo; Ashland, OR).[7-9]

RT-qPCR

To evaluate gene expression levels during iPSC differentiation, total RNA in frozen cells was

isolated using a RNeasy mini kit (Qiagen). The Maxima H-minus cDNA synthesis kit (Thermo

Fisher Scientific) with dT18 primer and random hexamer primer option for reverse transcription

was used to synthesize first strain cDNA from total RNA. SYBR Select Real-Time PCR Master

Mix (Thermo Fisher Scientific) was used in qPCR reactions that were performed on a

QuantStudio 12K Flex Real-Time PCR System (Applied Biosystems). Raw data was collected,

processed, and exported using the accompanying QuantStudio 12K Flex software suite. Ct values

were normalized to that of Gapdh, and a comparative $2^{-\Delta\Delta Ct}$ method was used to evaluate relative

gene expression in differentiated cells compared to iPSCs. Relative Expression Software Tool

(REST; version 2.0; Qiagen) was used to assess the changes in gene expression. The primer

sequences for genes of interest are the followings: Gapdh (5'-

CAACTCCCACTCTTCCACCTTC-3' and 5'-TTGCTGTAGCCGTATTCATTGTC-3'); Cdk1

(5'-GGCGAGTTCTTCACAGAGACTTG-3' and 5'-CCCTATACTCCAGATGTCAACCGG-

3'); and Tnnt2 (5'-AGAGGACACCAAACCCAAGC-3' and 5'-

CGACGCTTTTCGATCCTGTC-3').

Bisulfite Conversion of gDNA, and DNA Methylation (CpG) Analysis by Pyrosequencing

Genomic DNA from flash-frozen cells was isolated using a Blood and Tissue DNeasy kit (Qiagen). About 500 ng gDNA was used for CT bisulfite conversion with an EZ DNA Methylation-Gold kit (Zymo Research; #D5005), following the manufacturer's recommended protocol.

DNA methylation was measured by pyrosequencing on PCR products from bisulfite-treated gDNA. PyroMark Assay Design (version 2.0.1; Qiagen) was used for PCR primer pairs and pyrosequencing primer design against CpG regions around the super enhancer of Sox2 and the promoter of Cdk1. The sequence of primers are followings: for Sox2 gene: forward primer: 5' TGATTATAGGGAYGTGGGAGAATTTT 3', reverse primer: Biotin-5' AAAAAACCCTCAAACCTAATAAAATATCTA 3', pyrosequencing primer: 5' GGGAGAATTTTTTTTTGGAG 3'); and for Cdk1: forward primer: 5' TGGAAGGAAAATAGAGTTTAAGAGTTAG 3', reverse primer: Biotin-5' ACTCACRCCATACCTCTCTATCCC 3', pyrosequencing primer: 5' AGTTTTGATTGGTTTTTTTGA 3'). After primer quality verification, targeted CpG regions were amplified with a ZymoTaq Premix kit (Zymo; #E2003). Subsequently, PCR amplicons were sequenced with the pyroseq primers using a PyroMark Q24 DNA sequencer (Qiagen). The accompanied PyroMark Q24 software (version 2.0.8; Qiagen) was used to assess the CpG methylation level (percentage of the relative light unit (RLU) of the C/(C+T) peaks (C, methylated cytosine; T, unmethylated cytosine). CpG sites with reliable CpG methylation measurement were included in the calculation for averaged DNA methylation level for the sequenced CpG regions.

Statistics

After the derivation of GMR reporter clones, experiments were carried out with at least

biological replications, unless otherwise specified. Group data are expressed as Mean± SEM

unless otherwise specified. Difference between groups was assessed by t-test, with a P value

<0.05 considered a significant difference. ANOVA analysis were performed using GraphPad

Prism (version 10.2; GraphPad Software).

RESULTS

Mouse iPSC with GMR for Sox2 and Cdk1 – To test the utility of Snrpn mini promoter's

DNA methylation sensing function, we generated mouse iPSC lines stably carrying the Sox2-

GMR construct that is comprised of the Sox2 super enhancer CpG island region fused with the

Snrpn minimal promoter, followed by dTomato reporter (**Figure 1A**). iPSCs transfected with

Sox2-GMR were subjected to puromycin-resistant selection. Cells with high expression of

dTomato reporter signals were collected by fluorescence-activated cell sorting (FACS), and

single clones were expanded, among which clone C7 was used for subsequent experiments

(**Figure 1B**). Similar approaches were used to generate mouse Cdk1-GMR iPSC lines, which

stably carry Cdk1 promoter's CpG region, Snrpn minimal promoter and dTomato reporter.

Among the stable Cdk1-GMR iPSC lines, clone B9 was used for subsequent analyses. As

assessed by immunostaining and flow cytometry, the expression of stemness proteins such as

Oct4, Nanog, or Sox2 in GMR lines remained at levels comparable to wild-type mouse iPSC.

Fluorescent reporter signals in both Sox2 and Cdk1 GMR iPSCs were readily detected with

epifluorescent microscopy imaging.

Diminished reporter signals in cells differentiated from Cdk1 and Sox2 GMR iPSCs – Cell

differentiation is accompanied by epigenetic reprogramming that sets the landscape for

transcriptional activity for lineage specification, cell identity and cell-specific functions. As a

proof of concept for the utility of GMR, we examined the reporter signal and cardiac gene

expression in cells differentiated from the GMR iPSCs lines. By day 2.5 of directed cardiac

differentiation, dTomato reporter signals in both Sox2 GMR and Cdk1 GMR cells decreased

substantially (**Figure 2A**). By day 6.5 of differentiation, dTomato signal was minimally detected

in ~2.08% in Sox2 GMR cells, consistent with the loss of stemness in differentiated cells

assessed by flow cytometry on the gene expression of Sox2 and Oct4. dTomato signal in Cdk1

GMR cells also reduced significantly in differentiated cells (6.5 days) compared to iPSCs

($64.3\% \pm 7.1\%$ compared to $95.5\% \pm 1.5\%$; n=3; p<0.01). By 3 weeks, there was barely detectably

dTomato signal from Sox2 GMR or Cdk1 GMR-derived cells that were mostly contracting

myocytes (**Supplemental Figure and Videos**) expressing cTnT as detected by immunostaining

and flow cytometry (**Figure 2B, C**).

Correlations between GMR signals, gene expression, and DNA methylation – To further

verify that the GMR system can be used to visualize epigenetic regulation of genes by DNA

methylation, we examined the correlations between GMR reporter signal, transcript expression,

and CpG methylation of stemness and cell cycle genes. We performed RT-qPCR on cells

differentiated from Cdk1 GMR iPSCs. *Tnnt2* was expressed and increased in myocytes at 13.5

days of differentiation as compared to earlier timepoints. On the other hand, the transcript *Cdk1*

increased in 4.5 day differentiated cells compared to Cdk1 GMR iPSC, then decreased: *Cdk1* in

differentiated CM at day 13.5 was reduced to only 25% level of that in undifferentiated iPSCs or

CMs at differentiation data 6.5 (**Figure 3A**). Pyrosequencing showed significantly upregulated

CpG DNA methylation levels (hypermethylation) for stemness gene Sox2 and cell cycle gene

Cdk1 in the Cdk1-GMR reporter cells subject to cardiac differentiation. Sox2 and Cdk1 DNA

methylation was at 10.1% and 2.1% in undifferentiated iPSCs, and became 51.8% and 10.6%,

respectively, after induction of cardiac differentiation (at 4.5 day); and reached to 62.4% and

12.3% at 2 weeks of cardiac differentiation (**Figure 3B**). These results demonstrated DNA

methylation during stem cell differentiation, reflective of the reporter profiles of GMR system

and the accompanying gene expression.

DISCUSSION

Epigenetic DNA and histone modifications, and subsequently the changes in chromatin

accessibility, are the critical mechanisms controlling gene expression, thereby affecting cellular

phenotype and function, such as differentiation and proliferation.[7,10-16] Several approaches can be

used to reveal the status of DNA methylation; for example, fluorescent proteins in conjugation

with methyl-CpG-binding domain (MBD) or Synthetic-molecule/protein hybrid probes,[17]

modular fluorescence complementation-based epigenetic biosensors that combine engineered

DNA-binding proteins with domains recognizing defined epigenetic marks, both fused to non-

fluorescent fragments of a fluorescent protein.[18] The live-cell reporter system leveraging the

mini-promoter of imprinted *Snrpn* gene to visualize the CpG island regions of a target gene

represents a unique and highly specific system with detailed temporal and spatial resolutions

critical in the study of DNA methylation dynamics.[4]

In this study, we confirmed the fidelity of the CpG-Snrpn-based GMR approach in mouse iPSC

and differentiated myocytes. It is important to note that the methylation status of genes that have

too short no CpG regions may not be reflective in this GMR system. C2c12 cell lines carrying

Gapdh GMR (GFP) reporter and Sox2SE GMR (tdTomato) were also used to verify the expected

GMR signals. In addition, there was no reporter signal in cells transfected with pre-methylated

Sox2 or Gapdh GMR gene constructs (Data not shown). Live cell imaging data showed that the

positive GMR reporter signal reflects well the transcript expression of the corresponding gene.

Differentiated cells, like cardiomyocytes, have slower proliferation, and the cell cycle can be

arrested shortly after birth.[2] Our data show that during early differentiation (4.5 days), there was

an increased expression of *Cdk1,* then reduced significantly in more mature cardiomyocytes

(e.g., 13.5 days). It may be plausible that there was a fast proliferative phase right after the

transition to cardiac mesoderm lineage, leading to the higher *Cdk1* transcript expression, and the

changes in transcription activity may be delayed behind the GMR reporter signal for that short

period.[19] A caveat of using this GMR is that it is a "negative-positive" system in which the

reporter signal represents hypomethylation rather than hypermethylation. Nonetheless, the GMR

system is still informative as the change in reporter profile is consistent with epigenetic

mechanism in which an increase in DNA methylation leads to lower transcript expression, as

verified in our integrated transcriptional-epigenetic analysis (Figure 3). Lastly, we recognized

that while cell cycle gene *Cdk1* is reduced, accompanied by significantly upregulation of CpG

methylation in the promoter, the percentage methylation is much lower than Sox2. This could in

part due to the fewer CpG sites in mouse Cdk1 gene compared to Sox2 super enhancer and

promoter. Indeed, when a gene has too short CpG region, like mouse Tnnt2, its sequence will not

elicit well GMR-reporter dynamics as confirmed in our separated experiments (data not shown).

We also postulate that some genes conserved for various stage of cell functions in different cell

types have multiple regulatory mechanisms to fine tune gene expression, such as by microRNA,[9]

and epigenetic histone modifications.[20,21] It is worth noting that Cdk1 itself can affect histone

epigenetic landscape as shown in mouse embryonic stem cells.[22] The GMR reporter system can

be used with both live cell reporters to decipher the intricate cellular processes such as lineage

specification and tracking (e.g., cardiomyocyte) and phenotype (e.g., myocyte maturity or cell

cycle/proliferation activities) and functions (e.g., myocyte electrical and contractile activities).

In summary, the *Snrpn* mini-promoter-based GMR system, such as for *Cdk1* and other cell cycle

genes, can be instrumental in reporting dynamic DNA methylation states during cell

differentiation and cell cycling. Combining the new GMR system with other visualization

technologies will likely further delineate the epigenetic-transcriptional regulations of various

cellular phenotypes and functions.

Declaration of Competing Interest

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

Y.Z. conceived and conceptualized the study. Y.Z. and N.L. designed and performed experiments, collected, and analyzed data. B.T.N. contributed to gene expression analysis. Z.Z. conducted experiments on cell culture. W.R.M. helped analyze the results and discussion. N.L. and Y.Z. wrote the manuscript. All co-authors provided their edits to the manuscript.

Acknowledgments

The authors thank Yun-Yu Wu and Lovina Abdi for technical support and Dr. Karolina Peplowska for assistance in DNA methylation pyrosequencing experiments. N.L. was in part supported by the CSC Scholarship from China's Ministry of Education. The work was in part supported by research grants from the American Heart Association (18IPA34110210 to Y.Z.) the National Institute of Health (1R56HL131898, U54MD007601, P20GM113134, 5P20GM125526 to Y.Z.) and the University of Washington Institute for Stem Cells and Regenerative Medicine (Innovation Pilot Awards to Y.Z.).

REFERENCES:

1. Ammerpohl O, Martin-Subero JI, Richter J, Vater I, Siebert R. Hunting for the 5th base: Techniques for analyzing DNA methylation. *Biochim Biophys Acta*. 2009;1790:847-862. doi: 10.1016/j.bbagen.2009.02.001

2. Zhang Y, Mignone J, MacLellan WR. Cardiac Regeneration and Stem Cells. *Physiol Rev*. 2015;95:1189-1204.

3. Shemer R, Birger Y, Riggs AD, Razin A. Structure of the imprinted mouse Snrpn gene and establishment of its parental-specific methylation pattern. *Proc Natl Acad Sci USA*. 1997;94:10267-10272.

4. Stelzer Y, Shivalila CS, Soldner F, Markoulaki S, Jaenisch R. Tracing dynamic changes of DNA methylation at single-cell resolution. *Cell*. 2015;163:218-229.

5. Zhou HY, Katsman Y, Dhaliwal NK, Davidson S, Macpherson NN, Sakthidevi M, Collura F, Mitchell JA. A Sox2 distal enhancer cluster regulates embryonic stem cell differentiation potential. *Genes Dev*. 2014;28:2699-2711. doi: 10.1101/gad.248526.114

6. Kokkinopoulos I, Ishida H, Saba R, Coppen S, Suzuki K, Yashiro K. Cardiomyocyte differentiation from mouse embryonic stem cells using a simple and defined protocol. *Dev Dyn*. 2016;245:157-165. doi: 10.1002/dvdy.24366

7. Zhang Y, Gago-Lopez N, Li N, Zhang Z, Alver N, Liu Y, Martinson AM, Mehri A, MacLellan WR. Single-cell imaging and transcriptomic analyses of endogenous cardiomyocyte dedifferentiation and cycling. *Cell Discov*. 2019;5:30.

8. Gago-Lopez N, Awaji O, Zhang Y, Ko C, Nsair A, Liem D, Stempien-Otero A, MacLellan WR. THY-1 Receptor Expression Differentiates Cardiosphere-Derived Cells

with Divergent Cardiogenic Differentiation Potential. *Stem Cell Reports*. 2014;2:576-

591.

9.   Zhang Y, Matsushita N, Eigler T, Marban E. Targeted microRNA interference promotes

postnatal cardiac cell cycle re-entry. *JRegenerative Medicine*. 2013;2:doi:10.4172/2325-

9620.1000108.

10. Radak M, Fallahi H. The Epigenetic Regulation of Quiescent in Stem Cells. *Glob Med*

Genet. 2023;10:339-344. doi: 10.1055/s-0043-1777072

11. Chen X, Chakravarty T, Zhang Y, Li X, Zhong JF, Wang C. Single-cell transcriptome

and epigenomic reprogramming of cardiomyocyte-derived cardiac progenitor cells. *Sci*

Data. 2016;3:160079.

12. Zhang Y, Zhong JF, Qiu H, Robb MW, Marban E, Wang C. Epigenomic Reprogramming

of Adult Cardiomyocyte-Derived Cardiac Progenitor Cells. *SciRep*. 2015;5:17686.

13. Vallaster M, Vallaster CD, Wu SM. Epigenetic mechanisms in cardiac development and

disease. *Acta Biochim Biophys Sin (Shanghai)*. 2012;44:92-102.

14. McKinsey TA, Vondriska TM, Wang Y. Epigenomic regulation of heart failure:

integrating histone marks, long noncoding RNAs, and chromatin architecture. *F1000Res*.

2018;7.

15. Klemm SL, Shipony Z, Greenleaf WJ. Chromatin accessibility and the regulatory

epigenome. *Nat Rev Genet*. 2019;20:207-220. doi: 10.1038/s41576-018-0089-8

16. Ailsa M, Elisa De F, Alice F, Christine SF, Sarah JR, Kathryn M, Joe B, Barts Pancreas

Tissue Bank, Emma MW, Noel GM, et al. Developmentally dynamic changes in DNA

methylation in the human pancreas. *bioRxiv*. 2023:2023.2010.2019.563104. doi:

10.1101/2023.10.19.563104

17.   Kumar N, Hori Y, Kikuchi K. Live-Cell Imaging of DNA Methylation Based on

329   Synthetic-Molecule/Protein Hybrid Probe. *Chem Rec*. 2018;18:1672-1680. doi:

330   10.1002/tcr.201800039

331   18.   Lungu C, Pinter S, Broche J, Rathert P, Jeltsch A. Modular fluorescence

332   complementation sensors for live cell detection of epigenetic signals at endogenous

333   genomic sites. *Nature Communications*. 2017;8:649. doi: 10.1038/s41467-017-00457-z

334   19.   Soonpaa MH, Field LJ. Survey of studies examining mammalian cardiomyocyte DNA

335   synthesis. *Circ Res*. 1998;83:15-26. doi: 10.1161/01.res.83.1.15

336   20.   El-Nachef D, Oyama K, Wu YY, Freeman M, Zhang Y, MacLellan WR. Repressive

337   histone methylation regulates cardiac myocyte cell cycle exit. *J Mol Cell Cardiol*.

338   2018;121:1-12.

339   21.   Zhang Z, Freeman M, Zhang Y, El-Nachef D, Davenport G, Williams A, MacLellan WR.

340   Hippo signaling and histone methylation control cardiomyocyte cell cycle re-entry

341   through distinct transcriptional pathways. *PLoS One*. 2023;18:e0281610.

342   22.   Michowski W, Chick JM, Chu C, Kolodziejczyk A, Wang Y, Suski JM, Abraham B,

343   Anders L, Day D, Dunkl LM, et al. Cdk1 Controls Global Epigenetic Landscape in

344   Embryonic Stem Cells. *Mol Cell*. 2020;78:459-476.e413. doi:

345   10.1016/j.molcel.2020.03.010

Dynamic Visualization of DNA Methylation in Cell Cycle Genes during iPSC Cardiac

Differentiation

Ning Li[1,2], Ba Thong Nguyen[3,4*], Zhenhe Zhang[1*], W. Robb MacLellan,[1] Yiqiang Zhang[1,3,4¶]

[1] Division of Cardiology, Department of Medicine, University of Washington, Seattle, WA

98109, USA

[2] State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant

Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China

[3] Department of Anatomy, Biochemistry and Physiology, John A. Burns School of Medicine,

University of Hawaii at Manoa, Honolulu, HI 96813, USA.

[4] Center for Cardiovascular Research, John A. Burns School of Medicine, University of Hawaii

at Manoa, Honolulu, HI 96813, USA.

* Equal contribution.

¶ Author for Correspondence: Yiqiang Zhang, PhD (yiqiang.zhang@hawaii.edu).

Figures (3)

Figures and Figure Legends

Figure 1.

Figure 1. Generation of mouse GMR iPSC lines. A, Gene construction of Snrpn mini-
promoter based genomic DNA methylation reporter system. **B,** Microscopy images of wildtype
iPSC (control) and GMR iPSC lines for carrying GMR for Sox2 super enhancer or Cdk1
promoter's CpG regions.

Figure 2.

Figure 2. Dynamic reporter signal during cardiac differentiation of GMR iPSCs. A,

Microscopy images of Sox2-GMR and Cdk1-GMR iPSCs (day 0) and differentiated cells. **B,**

Flow cytometry dot plots showing the signal of GMR reporter dTomato and cardiomyocyte

marker cTnT (cardiac troponin T). **C**, Cell population quantified by the expression of GMR

signal dTomato of cardiac myocyte marker cTnT.

Figure 3.

Figure 3. Gene expression and DNA methylation during cardiac differentiation from Cdk1

GMR iPSCs. A, Relative quantification of *Cdk1* and *Tnnt2* in undifferentiated iPSCs and

cardiac differentiation day 4.5, 6.5, and 13.5; as normalized to *Gapdh.* *** and ###, p<0.05 vs 0d

For *Cdk1* and *Tnnt2*, respectively; n=3. **B**, CpG DNA methylation in Sox2 and Cdk1 CpG

regions in Cdk1 GMR iPSCs (0d) or cells from directed cardiac differentiation for 4.5, 6.5, and

13.5 days. Data points are % methylation of CpG sites assessed with pyrosequencing assays.

p<0.001 (2-way ANOVA).