Transgenic αβ TCR tonic signaling is leukemogenic while strong stimulation is

leukemia-suppressive

Telmo A. Catarino,[1,2,3] Ivette Pacheco-Leyva,[1,2] João L. Pereira,[1,2] Marina Baessa,[1,2]

Nuno R. dos Santos[1,2,4]

[1]i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto,

Portugal.

[2]Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP),

Porto, Portugal.

[3]Graduate Program in Areas of Basic and Applied Biology, Instituto de Ciências

Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal

[4]Centre for Biomedical Research (CBMR), University of Algarve, Faro, Portugal.

Running title: Dual function for αβ TCR in leukemogenesis

Text word count: 4291

Number of figures: 4

Number of references: 42

Abstract

The pre-T cell receptor (TCR) and TCR complexes are frequently expressed in T-cell acute lymphoblastic leukemia (T-ALL), an aggressive T cell precursor malignancy. Although mutations in TCR components are infrequent in T-ALL, earlier research indicated that transgenic αβ TCR expression in mouse T cell precursors promoted T-ALL development. However, we recently found that stimulation of TCR signaling in T-ALL induced leukemic cell apoptosis and suppressed leukemia. Our aim was to elucidate if a given αβ TCR complex has a dual role in leukemogenesis depending on the nature of the stimulus. We demonstrate that transgenic expression of the Marilyn αβ TCR, specific for the H-Y male antigen presented by major histocompatibility complex class II, triggers T-ALL development exclusively in female mice. This T-ALL exhibited *Notch1* mutations, *Cdkn2a* copy number loss, immature immunophenotype and infiltrated both lymphoid and non-lymphoid organs. Furthermore, leukemic cells expressed surface CD5, a marker of tonic TCR signaling. T-ALL efficiently developed in *Rag2*-deficient Marilyn transgenic females, indicating that Rag2-mediated recombination is not implicated in this T-ALL model. Remarkably, exposure of Marilyn female T-ALL to male antigen in recipient mice resulted in T-ALL apoptosis and prolonged mouse survival. These findings underscore that the same αβ TCR complex has a dual role in T-ALL in that its tonic stimulation is leukemogenic, while strong stimulation is leukemia-suppressive.

Introduction

The pre-T cell receptor (TCR) and αβ TCR are critical protein complexes for T cell development. Pre-TCR expression in CD4⁻CD8⁻ double-negative (DN) thymocytes induces their survival, proliferation and maturation through the β-selection checkpoint,[1,2] whereas surface expression of the αβ TCR in CD4⁺CD8⁺ double-positive (DP), thymocytes drives thymic positive and negative selection to establish major histocompatibility complex (MHC) restriction and eliminate autoreactive thymocytes.[3] The pre-TCR, TCR and CD3 proteins are frequently expressed in T-cell acute lymphoblastic leukemia (T-ALL).[4] Although T-ALL is genetically a very heterogeneous disease, with mutations detected in multiple genes, a number of genetic alterations have been recurrently reported for T-ALL, including *CDKN2A* loss of function, *NOTCH1*-activating mutations, and mutations in JAK/STAT signaling pathway genes (e.g. *IL7R*, *JAK3*, *JAK1*, and *STAT5B*).[5] Mutations in components of the TCR signaling machinery are rare,[5] but the Lck kinase, a key member of the pre-TCR and TCR signaling pathways, has been shown to be activated in T-ALL and put forward as a potential therapeutic target.[6–9] These observations hint that pre-TCR signaling can promote human T cell leukemogenesis. In this line, genetic inactivation of the pre-TCR complex (e.g. CD3ε, pTα or Rag1/2 deficiency) was shown to delay T-ALL development in several mouse models.[10–14] In contrast, the impact and function of TCR signaling in T-ALL development and progression remains controversial. While transgenic TCR expression was shown to promote mouse T-ALL development in cooperation with STAT5 overexpression or deficiency of p53 or Tpl2 proteins,[15–17] expression of endogenous αβ TCR or MHC-dependent antigen presentation were not required for leukemogenesis in the ETV6::JAK2 or TAL/LMO transgenic mouse models.[13,14] Indicating that TCR could rather have an antagonistic action in T-ALL, T-ALL onset in a *Pten*-deficient mouse model was accelerated by TCR genetic inactivation and delayed by expression of transgenic TCR.[18] In this line, it was also reported that strong agonist TCR stimulation of mouse ETV6::JAK2-driven T-ALL or patient T-ALL was anti-leukemic and induced a transcriptional program similar to that involved in thymocyte negative selection.[19] Overall, these observations suggest that, depending on the oncogenic drivers of leukemia and degree of TCR stimulation, the TCR signaling complex might play either a leukemia-promoting or a leukemia-suppressive role.

Here, our aim was to investigate the role of basal versus strong TCR signaling in T-ALL leukemogenesis. In normal T cells, TCR recognition of self-peptides presented by MHC (self-pMHC) induces tonic signaling, while agonist peptides induce strong TCR signaling. Using the Marilyn transgenic TCR mouse strain, specific for the H-Y male antigen, we show that tonic TCR signaling led to *Notch1*-mutated T-ALL exclusively in

78   females. MHC-dependent agonist antigenic stimulation prevented spontaneous T-ALL

79   development in males, induced apoptosis of female leukemic cells and hampered their

80   leukemogenic potential. Our results support the notion that αβ TCR tonic signaling

81   promotes T cell precursor malignant transformation and that strong signaling suppresses

82   leukemia.

Results

Transgenic expression of an αβ TCR specific for the H-Y male antigen drives leukemia development in female mice

To study the effect of TCR expression in leukemia development, we resorted to Marilyn transgenic mice, which express an MHC class II-restricted transgenic TCR (TCR-Vα1.1/Vβ6) specific for the H-Y male antigen (HY-TCR henceforth).[20] In this model, male thymocytes expressing the HY-TCR are exposed to the cognate antigen (DBY peptide), through MHC class II presentation, and are eliminated by apoptosis. In contrast, HY-TCR-expressing female thymocytes undergo positive selection and differentiate as mature CD4$^+$ T cells. To verify if the transgenic Marilyn HY-TCR promoted leukemogenesis, we bred transgenic mice expressing the ETV6::JAK2 fusion protein in lymphoid cells with Marilyn mice. As expected,[13] all ETV6::JAK2 transgenic mice developed T-ALL expressing CD8, CD25, CD24, CD3ε and TCRβ (not shown). Remarkably, co-expression of the Marilyn HY-TCR with ETV6::JAK2 accelerated spontaneous T-ALL development in females (median survival of 10.5 weeks versus 14 weeks; Figure 1A). This suggests that HY-TCR tonic signaling, which is induced by TCR interaction with self-peptides presented by MHC, cooperates with the ETV6::JAK2 transgene in driving leukemogenesis.

To determine whether HY-TCR tonic signaling alone could induce T cell malignant transformation, we monitored a cohort of both Marilyn females and males until 18 months of age. Remarkably, while most females developed fatal hematological disease (median survival of 58 weeks), none of the males did (Figure 1B). Macroscopic analyses showed that Marilyn females developed enlarged lymphoid (thymus, spleen and lymph nodes (LNs)) and non-lymphoid organs (mainly liver and kidneys) (Figure 1C; Supplementary Figure 1), together with high number of blood lymphoblasts (Figure 1D). Histological analyses revealed massive lymphoid cell infiltration of liver, kidneys, lungs and bone marrow (Supplementary Figure 2). These data indicate that tonic (in females) but not strong (in males) transgenic HY-TCR signaling is sufficient to induce leukemia and lymphoma development.

Transgenic HY-TCR expression induced T-cell disease with variable immunophenotype

Macroscopic analysis revealed two distinct groups of diseased Marilyn females: a subset with thymic lymphoma (TL; 9 cases) and a subset characterized by splenomegaly and lymphadenopathy without thymic lymphoma (non-TL; 7 cases; Figure 1E). TL females developed leukemia slightly faster and presented higher total lymphoid organ disease burden than non-TL females, suggestive of a more aggressive disease

course (Supplementary Figures 3A,B). For a better characterization of the hematological disease developing in different Marilyn females, we performed immunophenotyping by flow cytometry. To define the T cell maturation status, we probed expression of CD24, which is downregulated upon thymocyte maturation, CD5, which is upregulated after thymocyte positive selection, and CD44, which is expressed in memory T cells.[21,22] Within the TL subgroup, five mice (nos. 2, 6, 22, 33 and 40) presented thymic T cells with immature phenotype, characterized by surface expression of CD90.2, TCR-Vβ6, and CD5, low levels of CD4 and CD8, high CD24 levels, and CD44 negativity (Figure 2 and Supplementary Table 1). Such immunophenotype was recapitulated in leukemic cells disseminated to the spleen, LNs, and bone marrow (Supplementary Figure 4 and not shown). Two other TL cases (nos. 9 and 29) presented a more mature T cell immunophenotype, characterized by CD4$^+$CD8$^-$TCR-Vβ6$^+$CD5$^+$ leukemic cells with CD44 expression (Figure 2). These cells were also present in spleen and lymph nodes (Supplementary Figure 4). Finally, and rather unexpectedly, two TL cases (nos. 3 and 34) presented a large proportion of B220-positive B cells, concomitantly with TCR-Vβ6$^+$CD5$^+$CD44$^+$ CD4 single-positive (SP) and CD8SP T cells (Figure 2). Of note, TL cases with immature immunophenotype, but not those with mature phenotype, could reinitiate disease in recipient syngeneic mice (Supplementary Table 1), suggesting that only the former were malignantly transformed.

Non-TL Marilyn females exhibited a mixture of B220$^+$ B cells and CD90.2$^+$TCR-Vβ6$^+$CD5$^+$CD44$^+$ CD4SP cells in lymphoid organs (Figure 2 and Supplementary Figure 4 and Table 1). Analysis of CD69 expression, a marker of TCR stimulation,[21] showed that TCR-Vβ6$^+$ cells from mixed B and T lymphomas were CD69-positive, indicating they experienced recent TCR stimulation (Supplementary Figure 5). This suggests that mixed B and T disease found in Marilyn female mice was driven by continuous stimulation of the HY-TCR transgene expressed in mature CD4SP or CD8SP T cells. Immature-type leukemic cells did not express CD69, suggesting that their proliferation was likely driven by oncogenic mechanisms rather than TCR stimulation.

We conclude that while one third (5 of 16) of Marilyn TCR transgenic females developed a transplantable immature disease, akin to T-ALL, marked by pronounced thymic enlargement, the remaining diseased mice developed a non-transplantable and heterogeneous mature T and B nonmalignant lymphoproliferation affecting the thymus, spleen, lymph nodes and other organs, with statistically significant longer delay (median survival of 40 versus 60 weeks; Supplementary Figure 6).

Marilyn T-ALL presents *Notch1* PEST domain mutations and genetic copy number alterations

To determine whether Marilyn T-ALL was associated with secondary genetic alterations, we first sequenced the *Notch1* exon 34 (PEST domain), where most murine T-ALL *Notch1* mutations have been found.[23,24] Four out of 5 Marilyn T-ALL cases analyzed carried exon 34 frameshift mutations (Supplementary Figure 7 and Table 1), while none were detected in cases of mixed T and B cell disease (Supplementary Table 1). Additionally, by performing low-coverage whole-genome sequencing, we found that T-ALL samples had near-diploid genomes, with only a few copy number alterations: a chromosome 15 trisomy (case no. 6), a chromosome 10 trisomy (case no. 33) (Supplementary Figure 8A), and loss of the *Cdkn2a* locus (chromosome 4) in three cases (nos. 2, 22 and 40) (Supplementary Figure 8B). In contrast, no consistent copy number alterations were found in 2 cases of mixed T and B cell disease (Supplementary Figure 8A). Deletions of the *Tcra/d*, *Tcrb* and *Tcrg* loci were also found in Marilyn T-ALL (Supplementary Figure 8C), indicating that these loci underwent genetic rearrangements and that leukemia arose from post-β-selection thymocytes with rearranged TCR loci. As expected from their T lineage origin, T-ALL cases did not exhibit consistent copy number alterations in the immunoglobulin *Ighm* or *Igk* loci (Supplementary Figure 8C). The detection of clonal copy number genetic alterations in Marilyn T-ALL bulk populations further underscores their malignant nature. Given the high frequency of *NOTCH1*-activating mutations and *CDKN2A* inactivation in human T-ALL,[5] our findings show that the Marilyn transgenic HY-TCR T-ALL recapitulates key cellular and molecular features of the human disease.

HY-TCR-induced T-ALL is not dependent on *Rag2* recombinase expression

Since several diseased Marilyn females developed B cell lymphoproliferation, we generated Marilyn mice with *Rag2* constitutive knockout to block B cell development, while maintaining HY-TCR-dependent T-cell development. This approach simultaneously assessed whether Rag-mediated gene recombination plays a role in HY-TCR-induced T cell leukemogenesis. Marilyn HY-TCR cohorts with (Marilyn;*Rag2*^+/-^) or without Rag2 protein (Marilyn;*Rag2*^-/-^) were generated in another animal facility (i3S/Porto). Surprisingly, most Marilyn;*Rag2*^+/-^ female mice (78%) did not develop disease, with the exception of two mice with splenomegaly at nearly 18 months of age, which was composed of B and myeloid cells (Figure 3A and not shown). Although half of Marilyn;*Rag2*^-/-^ males developed hematological splenic disease after 1 year of age (4 cases of myeloid proliferation and 1 case of *Notch1*-mutated TCR-Vβ6⁺ CD4⁻CD8⁻ splenic lymphoma), nearly all Marilyn;*Rag2*^-/-^ females developed lymphoma with much

earlier onset than males (median survival of 46 and 70 weeks, respectively; Figure 3A).
All diseased Marilyn;*Rag2*[-/-] females presented thymic lymphoma, of variable size,
frequently accompanied by enlarged spleen, LNs, liver and kidneys (Figure 3B;
Supplementary Figure 9A). Leukemias from Marilyn;*Rag2*[-/-] females were mostly
composed of CD4[+/low]CD8[+/low] immature T cells expressing TCR-Vβ6, high levels of
CD24, CD44 negativity and variable expression of CD25 and CD5 (Figure 3C, panel i,
and Supplementary Table 2). A similar immunophenotype was detected across different
affected organs, suggesting that diseased cells disseminated from the thymus to other
organs (Figure 3C, panel ii). The pathological and immunophenotypic features, together
with the observed 60% frequency of *Notch1* exon 34 frameshift mutations and ability to
reinitiate disease in syngeneic recipients (Supplementary Table 2), indicate that
Marilyn;*Rag2*[-/-] females developed T-ALL. To assess the degree of leukemic cell
infiltration in different Marilyn;*Rag2*[-/-] organs, we performed histological analysis.
Lymphoblasts were conspicuous in the thymus and spleen, and infiltrated with variable
extent, the liver, kidneys, lungs and bone marrow (Supplementary Figure 9B).
Interestingly, immunohistochemical analyses of thymus histological sections showed
that while Marilyn T-ALL cells were highly proliferative, with conspicuous Ki67 staining,
they maintained a high level of apoptosis, as gauged by cleaved caspase 3 staining
(Figure 3D). In sum, these data confirm that Marilyn HY-TCR tonic signaling drives T-
ALL and demonstrated that Rag recombinase-mediated DNA rearrangements are not
implicated.
213

Hematological disease in HY-TCR transgenic females is associated with microbial
status
Marilyn females in the UAlg/Faro animal facility had higher incidence of
hematological disease than the Marilyn;*Rag2*[+/-] females in the i3S/Porto facility. Since
opportunistic microorganisms were detected in the UAlg but not the i3S facility (see
Materials and methods), we posit that an environmental input present only at the UAlg
facility increased susceptibility for lymphoid disease. Considering that bacteria are
frequently present in tumors, and can be detected by immunohistochemical methods,[25]
we probed lymphomas obtained in both facilities for the presence of bacterial-derived
lipopolysaccharide (LPS). Indeed, Marilyn lymphomas derived from the UAlg facility
displayed more conspicuous LPS immunostaining than Marilyn;*Rag2*[-/-] lymphomas
derived from the i3S facility (Supplementary Figure 10A). Of note, the pattern of LPS
staining in Marilyn splenic lymphomas overlapped with B220[+] B cell areas
(Supplementary Figure 10B). These data suggest that higher bacterial exposure could
have promoted hematological diseases in Marilyn females.

The transgenic HY-TCR is functional in Marilyn T-ALL cells and its stimulation delays leukemogenesis

Given that antigenic stimulation prevented spontaneous T-ALL development in
male HY-TCR mice, we set out to determine the response of female HY-TCR leukemic
cells to TCR stimulation. Marilyn T-ALL cells readily increased CD69 and CD5
expression and enlarged in size when stimulated *in vitro* with plate-bound CD3 antibody,
which directly activates the TCR signaling complex, or phorbol 12-myristate-13-acetate
(PMA) and ionomycin, triggering downstream activation of protein kinase C and
intracellular calcium pathways (Supplementary Figure 11). Previous research
demonstrated that agonist TCR stimulation can induce T-ALL apoptosis *in vitro*.[19]
However, Marilyn leukemic cells underwent extensive apoptosis *in vitro*, impeding
determination of whether TCR stimulation induced cell death. To assess the impact of
TCR stimulation *in vivo*, Marilyn T-ALL cells were injected in syngeneic male and female
mice and disease progression followed by peripheral blood collection. Marilyn T-ALL
cells were found in the blood of females, but not males, as early as 30 days post-injection
(Figure 4A). Most female recipients rapidly developed fatal T-ALL, while male recipients
often failed to develop leukemia and therefore survived longer (Figure 4B). In addition,
females infused with Marilyn T-ALL presented significantly larger spleen and kidneys
(Figure 4C), and more extensive BM leukemia infiltration (Figure 4D), than male
recipients euthanized simultaneously. Furthermore, male spleens showed higher levels
of cleaved caspase 3 immunostaining than female spleens (Figure 4E), indicating that
delayed leukemogenesis in males was caused by apoptosis induction.

To address whether the leukemia-delaying effect of agonist-induced TCR
stimulation in Marilyn T-ALL cells was dependent on MHC class II presentation, Marilyn
T-ALL cells were infused in immunodeficient syngeneic and allogenic males and
females. In contrast to the delayed T-ALL development in syngeneic males, as compared
to female counterparts (Figure 4F, left panel), no significant difference between
allogeneic males and females was observed (Figure 4F, right panel). These findings
indicate that the T-ALL-suppressive effect of the male antigen was dependent on
restricted MHC class II molecules. To confirm that delayed Marilyn leukemogenesis in
male mice was due to male antigen presentation, we next treated syngeneic
immunodeficient females infused with Marilyn T-ALL with the DBY cognate peptide.
Indeed, DBY administration significantly delayed T-ALL development in females (Figure
4F). Overall, these data indicate that strong agonistic HY-TCR signaling suppresses
leukemogenesis.

Discussion

In this report, we addressed the functional role of TCR signaling intensity in

mouse T-ALL. We show that transgenic Marilyn HY-TCR tonic signaling not only had a

synergistic effect with the ETV6::JAK2 fusion kinase, but also was sufficient to induce T-

ALL on its own. Furthermore, our data show that the TCR expressed in Marilyn T-ALL

cells responded to agonist stimuli, inducing apoptosis and leading to leukemogenesis

suppression. Thus, these results support the notion that tonic TCR signaling can be

leukemogenic, and strong signaling emanating from the same TCR is anti-leukemic.

Tonic signaling is generated by relatively weak TCR interactions with self-pMHC

ligands and is important for controlling survival of naive T cells and allow their

responsiveness to foreign antigens.[26] Thymocytes expressing transgenic TCRs against

foreign antigens (such as the male antigen for females) are subjected to self-pMHC-

mediated tonic signaling during their maturation, namely at the positive selection stage.

The notion that TCR signaling can be leukemogenic has been earlier described, and

transgenic TCR mouse strains other than the Marilyn strain, were shown to lead to

spontaneous T-ALL development with variable latency and penetrance.[27–30] It is thus

likely that tonic signaling originates pro-survival or pro-proliferative genetic programs that

can cooperate with stochastic oncogenic genetic alterations, such as *Notch1*-activating

mutations and *Cdkn2a* deletions. *Cdkn2a* inactivation in particular was reported to

facilitate the malignant transformation of immature T cells,[31,32] so it is possible that loss

of *Cdkn2a* function may cooperate with TCR tonic signaling in DN thymocytes.

The finding that tonic TCR signaling can promote malignant transformation finds

parallels in other systems. In fact, tonic B cell receptor signaling, defined as antigen-

independent cell-autonomous signaling, has been shown to drive B cell

lymphomagenesis [33–35] and to be therapeutically targetable.[36] By the same token, tonic

TCR signaling could be a targetable mechanism in human T-ALL, as suggested by the

frequent activation of Lck kinase (a mediator of TCR signaling) in T-ALL and by the

therapeutic potential of Lck inhibitors.[6–9] Lck kinase activation in T-ALL was attributed to

constitutive pre-TCR signaling found in a subset of patients.[6] Since TCR and CD3

proteins are also frequently expressed in T-ALL,[4] further studies are warranted to

determine the role of TCR signaling in human T-ALL pathogenesis.

Although Rag recombinase-mediated mutagenic events can contribute to

leukemia development,[37] that was not the case for Marilyn T-ALL. In fact, T-ALL

developed with higher penetrance and much faster in Marilyn;*Rag2*[-/-] females than in

Marilyn;*Rag2*[+/-] females generated in the same animal facility. Since Rag2 deficiency

arrests thymocyte development at the DN3 stage, these results suggest that ectopic TCR

expression at the immature DN3 stage in the absence of more mature thymocytes has

increased leukemogenic potential, in line with the finding that early expression of mature

TCR in DN thymocytes promotes T-ALL development.[30]

Rag2-sufficient, Marilyn mouse cohorts housed in different animal facilities led to

unexpected different rates of T-ALL incidence. Indeed, Marilyn T-ALL frequency was

much higher in females housed at the UAlg/Faro facility than females housed at the

i3S/Porto facility. Since several microorganisms (*Helicobacter* spp., *Pasteurella*

pneumotropica and murine norovirus) were detected in the UAlg facility, but not in the

i3S facility, increased exposure to pathogenic microbiota could have promoted a

microenvironment favorable for transgenic TCR-induced leukemogenesis. This notion is

supported by the finding of higher levels of LPS staining in lymphomas from UAlg facility

mice than in lymphomas from i3S facility mice. Early research revealed that B-ALL in

Pax5 heterozygous mice was initiated only upon exposure to common pathogens.[38]

Future studies should, therefore, determine whether the presence of microorganims

promotes TCR-induced leukemogenesis and whether TCR interactions with MHC

molecules presenting non-cognate foreign peptides generate pro-leukemogenic basal

TCR signaling.

Although T-ALL in Marilyn females developed in the presence of self-pMHC, we

found that MHC-dependent TCR signaling was not required for its maintenance. Indeed,

secondary leukemia development was not affected by lack of MHC-bound self-peptides

in recipient allogenic mice. These results contrast with the finding that continuous self-

pMHC-mediated TCR signaling was necessary for lymphoma cell growth in a TCR-

expressing reprogrammed T cell model.[39] Of note, expression of a transgenic TCR

recognizing a peptide from the murine survivin protein was also shown to result in

increased TCR signaling and to induce T-ALL.[29] In this case, the specific (self) antigen,

survivin, was expressed constitutively, which contrasts with the absence of the HY-TCR

cognate peptide (DBY) in Marilyn females. Our data together with previous reports

demonstrate that the TCR signaling complex promotes T-ALL, and that this ability very

much depends on the TCR affinity to antigen.

Finally, we showed that Marilyn T-ALL cells had a functional TCR and that *in vivo*

TCR stimulation of transplanted female Marilyn T-ALL cells was associated with

apoptosis and impaired Marilyn T-ALL leukemogenesis. These observations corroborate

previous reports showing that mouse or human T-ALL underwent apoptosis upon

activation of the TCR/CD3 signaling complex.[19,28,40] These findings suggest that, similarly

to thymocytes undergoing thymic selection, TCR activation with low-affinity or high-

affinity antigens will stimulate differentially the TCR signaling pathway and result in cell

survival or death.[41] Future research should determine the TCR signaling regulators that

dictate if a particular antigenic stimulus promotes or antagonizes T-ALL.

Methods

Mice

HY-TCR Marilyn transgenic mice (B6.Cg-Tg(TcraH-Y,TcrbH-Y)1Pas), on a
CD45.1 background, were obtained from Jocelyne Demengeot (IGC, Oeiras). Marilyn
mice were bred with EµSRα-ETV6::JAK2 (B6.Cg-Tg(Emu-ETV6/JAK2)71Ghy)
transgenic mice[13], or *Rag2* (B6.129S6-Rag2^{tm1Fwa}) knockout mice. The ETV6::JAK2
transgene was kept in hemizygosity. *Foxn1*$^{nu/nu}$ (Nude) C57BL/6 and CBA mice were
bred at the i3S facility. Mice were maintained at the UAlg/Faro and i3S/Porto barrier
animal facilities, under 12 h light/dark cycles and with food and water *ad libitum*. HY-TCR
Marilyn mice and ETV6::JAK2;Marilyn double transgenic were bred at the UALg animal
facility, while the Marilyn;*Rag2*$^{+/-}$ and Marily;*Rag2*$^{-/-}$ were bred at the i3S animal facilty.
Microorganism screening detected opportunistic pathogens (*Helicobacter* spp.,
Pasteurella pneumotropica and murine norovirus) in the UAlg, but not i3S, experimental
rooms. All experimental procedures were approved by the i3S and CBMR/UAlg ethics
committees and Portuguese authorities (*Direção-Geral de Agricultura e Veterinária*) and
followed recommendations from the European Commission (Directive 2010/63/UE) and
the local Portuguese authorities (*Decreto-Lei* n°113/2013). Both female and male mice
were used for all experiments. Mice were monitored for signs of disease (e.g. dyspnea,
lethargy, enlarged lymph nodes and enlarged abdomen) and killed by CO_2 inhalation
when reaching predefined experimental endpoints. Mice of different genotypes from the
same litter were kept together in the same cages and monitoring for signs of disease
was done blindly. Adult mice that were euthanized without leukemia were censored in
Kaplan-Meier survival curves.

Mice genotyping

The following primers were used for genotyping. HY-TCR Marilyn transgene: 5'-
CGAGAGGAACCTGGGAGCTGT-3' and 5'-TGCTGTCTGTACCACCAGAAATAC-3';
Rag2: 5'-TGTCCCTGCAGATGGTAACA-3', 5'-CCTTTGTATGAGCAAGTAGC-3', 5'-
CTATTCGGCTATGACTGGG-3' and 5'-AAGGCGATAGAAGGCGATG-3'; *Cdkn2a*: 5'-
GTGATCCCTCTACTTTTTCTTCTGACTT-3', 5'-CGGAACGCAAATATCGCAC-3' and
5'-GAGACTAGTGAGACGTGCTACTTCCA-3'.

***In vivo* experiments**

For mouse leukemia transplantation assays, 0.5-2 x 10^6 leukemic cells collected
from diseased female Marilyn mice were intravenously injected in the tail vein of recipient
8-12-week-old C57BL/6, C57BL/6 Nude or CBA Nude mice of the indicated sex, and
regularly monitored through peripheral blood detection of leukemic cells

(CD45.1$^+$CD45.2$^+$TCR-Vβ6$^+$). Female *Rag2*$^{-/-}$ mice infused with Marilyn T-ALL cells were

treated intraperitoneally once a week for two weeks with 0.01 mg of DBY peptide

(NAGFNSNRANSSRSS; cat. no. AS-61046; Eurogentec).

Flow cytometry

Single-cell suspensions were prepared from lymphoid organs using cell strainers,

washed with FACS buffer (phosphate-buffered saline (PBS) with 3% fetal bovine serum

(FBS) and 10 mM NaN$_3$), and incubated for 30-45 min with fluorochrome-labeled

antibodies in FACS buffer. The following Biolegend antibodies were used: CD25-

fluoresceine isothiocyanate (FITC) (clone PC61), TCRβ-phycoerithrin (PE) (clone H57-

597), CD44-PE/Cyanine5 (clone IM7), CD90.2-PE-Cyanine7 (clone 30-H12), CD24-

allophycocyanin (APC) (clone M1/69), CD4-APC/Cyanine7 (clone GK1.5), CD8α-Pacific

Blue (clone 53-6.7), Gr1-PE (clone RB6-8C5), CD127-PE/Cyanine5 (clone A7R34),

B220-APC/Cyanine7 (clone RA3-6B2), CD24-Pacific Blue (clone M1/69), CD90.2-FITC

(clone 30-H12), TCR-Vb6-PE (clone RR4-7), TCR-Vb5.1, 5.2-PE (clone MR9-4), CD5-

APC (clone 53-7.3), CD69-FITC (clone H1.2F3), CD62L-APC (clone MEL-14), CD45.1-

APC (clone A20) and CD45.2-Peridinin Chlorophyll Protein (PerCP)-Cyanine5.5 (clone

104). Immunostained cells were washed twice with FACS buffer and incubated in PBS

with 10 mM NaN$_3$. Cell viability was determined using the Zombie Aqua Fixable Viability

Kit (Invitrogen). Samples were acquired using BD FACS Calibur, CANTO II or Accuri C6

and analyzed using FlowJo software.

***Ex-vivo* T-ALL cell culture**

Marilyn T-ALL cells were cultured ex-vivo for 16 h in RPMI medium supplemented

with 10% FBS, 1% penicillin-streptomycin and 50 μM of 2-mercaptoethanol, all from

Gibco. For *ex vivo* stimulation of Marilyn T-ALL cells, plates were coated with 10 μg/ml

anti-CD3 (145-2C11, Biolegend) for 2 h at 37°C, and then plates were washed with PBS.

Ten ng/ml of PMA and 250 ng/ml Ionomycin (both from Sigma-Aldrich) treatments were

performed for 18 hours.

***Notch1* mutation detection**

Genomic DNA from mouse leukemia samples was isolated using the GeneJET

Genomic DNA purification kit (Thermo Fisher Scientific), following the manufacturer's

instructions. DNA was used for PCR amplification of two segments of *Notch1* exon 34:

primer pair 5'-GCTCCCTCATGTACCTCCTG-3' and 5'-TAGTGGCCCCATCATGCTAT-

3', generating a predicted amplicon of 904 bp, and primer pair 5'-

ATAGCATGATGGGGCCACTA-3' and 5'-CTTCACCCTGACCAGGAAAA-3', generating

a predicted amplicon of 893 bp. PCR products were Sanger sequenced using the same

primers at i3S Genomics (Porto, Portugal) or CCMAR *Serviços de Biologia Molecular*

(Faro, Portugal).

Low-coverage whole genome sequencing

Genomic DNA was used for copy number analysis performed as described

previously[42] using a HiSeq 4000 (Illumina) sequencer in a single-read 50-cycle run mode.

For copy number analysis, sequencing data was aligned with the mouse genome

(mm10) as reference and further analyzed using QDNAseq (RRID:SCR_003174)

package (version 1.12) in R software. A bin size of 15 kb was used.

Histology and immunohistochemistry

For histological and immunohistochemistry analysis, formalin-fixed paraffin-

embedded tissues were sectioned with 4 μm thickness. Hematoxylin and eosin staining

was performed using standard procedures. For immunohistochemistry, heat-mediated

antigen retrieval was performed for 35-40 min with citrate-based antigen retrieving

solution (cat. no. H-3300, Vector Laboratories), for cleaved-caspase 3, Ki67 and LPS

immunodetection, or 10 μM EDTA, pH 8.0, and 0.05% Tween 20 in PBS, for B220 or

CD3 immunodetection. Endogenous peroxidase was inactivated with 3% H_2O_2 in

methanol, and nonspecific antibody binding was blocked using Ultravision Protein-block

(Thermo Fisher Scientific). Sections were incubated overnight with rat anti-B220 (1:200,

clone RA3-6B2, Biolegend), anti-CD3 (1:200, ab5690, Abcam), anti-cleaved-caspase 3

(1:200, 9661, Cell Signaling), Ki67 (1:500, ab15580, Abcam) or LPS (1:100, WN1222-5,

Hycult Biotech) at 4°C. For B220 detection, slides were incubated sequentially with

avidin/biotin blocking system (Biolegend), goat anti-rabbit secondary antibody

(Biovision), and streptavidin-horseradish peroxidase (HRP) (Enzo Biochem). For CD3,

Ki67 and LPS detection, HRP-conjugated rabbit/mouse secondary antibody (Agilent)

was used. 3,3′-Diaminobenzidine chromogen (Agilent) was used as detection reagent.

For quantification of percentage of cleaved caspase 3-positive cells, five images were

taken from different areas of each tissue section.

Statistics

Statistical analysis was performed with GraphPad Prism 6.0 software

(RRID:SCR_002798). Log-rank test was used to compare survival of different groups.

Unpaired student's t-test was used for comparisons between two groups. Welch's

correction was used when variances were different. Mann-Whitney test was used to

compare white blood cell (WBC) count. Detection of CD45.1$^+$ T-ALL cells in the
peripheral blood of males and females was compared using two-way ANOVA. Sample
numbers are indicated in figure legends. *P*<0.05 was considered statistically significant.
453

454 **Data Sharing Statement**

455   The data that support the findings of this study are available from the
456 corresponding author, upon reasonable request. Raw unaligned sequencing reads
457 (fastq-format) that support the findings of this study have been deposited in the SRA.
458

Acknowledgments

We thank Jocelyne Demengeot (IGC, Oeiras) and Nuno L. Alves (i3S, Porto) for
providing Marylin transgenic mice. We thank members of i3S Intercellular
Communication and Cancer group for fruitful discussions about this work. The authors
acknowledge the support the Animal Facility, Translational Cytometry and Histology,
Electron Microscopy (member of the Portuguese Platform of Bioimaging; PPBI-POCI-
01-0145-FEDER-022122) and Genomics i3S Scientific Platforms. This work was
supported by a fellowship (FAZ Ciencia prize) from Fundação AstraZeneca (Lisbon,
Portugal). This work was supported by European Regional Development Fund (ERDF),
through COMPETE 2020 - Operational Program for Competitiveness and
Internationalisation, Portugal2020, and *Fundação para a Ciência e a Tecnologia* (FCT;
POCI-01-0145-FEDER-007274; PTDC/MED-ONC/32592/2017), and by ERDF, through
the Norte Portugal Regional Program (NORTE2020), Portugal2020 (NORTE-01-0145-
FEDER-000029). T.A.C. was recipient of FCT (PD/BD/114129/2015) and *Liga*
Portuguesa Contra o Cancro – Núcleo Regional do Norte fellowships. J.L.P.
(SFRH/BD/147979/2019) and M.B. (2021.06024.BD) were recipients of FCT fellowships.
475

476 **Contribution:** T.A.C. designed, performed and analyzed experiments, made the figures
477 and wrote the paper; I.P.-L., J.L.P., and M.B. performed experiments; N.R.S. designed
478 the study, performed experiments, made the figures and wrote the paper.
479

480 **Conflict-of-interest disclosures:** No competing financial interests to declare.
481

Correspondence: Nuno R. dos Santos, i3S - Instituto de Investigação e Inovação em
Saúde, Universidade do Porto, Rua Alfredo Allen, 208, 4200-135 Porto, Portugal; e-mail:
nunos@i3s.up.pt

References

1.   Shah, D. K. & Zúñiga-Pflücker, J. C. An Overview of the Intrathymic Intricacies of
T Cell Development. *The Journal of Immunology* **192**, 4017–4023 (2014).

2.   Liu, X. *et al.* Notch-induced endoplasmic reticulum-associated degradation
governs mouse thymocyte β-selection. *Elife* **10**, (2021).

3.   Klein, L., Kyewski, B., Allen, P. M. & Hogquist, K. A. Positive and negative
selection of the T cell repertoire: what thymocytes see (and don't see). *Nat Rev*
Immunol **14**, 377–391 (2014).

4.   Asnafi, V. *et al.* Analysis of TCR, pT alpha, and RAG-1 in T-acute lymphoblastic
leukemias improves understanding of early human T-lymphoid lineage
commitment. *Blood* **101**, 2693–2703 (2003).

5.   Girardi, T., Vicente, C., Cools, J. & De Keersmaecker, K. The genetics and
molecular biology of T-ALL. *Blood* **129**, 1113–1123 (2017).

6.   Gocho, Y. *et al.* Network-based systems pharmacology reveals heterogeneity in
LCK and BCL2 signaling and therapeutic sensitivity of T-cell acute lymphoblastic
leukemia. *Nat Cancer* **2**, 284–299 (2021).

7.   Laukkanen, S. *et al.* Therapeutic targeting of LCK tyrosine kinase and mTOR
signaling in T-cell acute lymphoblastic leukemia. *Blood* **140**, 1891–1906 (2022).

8.   Serafin, V. *et al.* Glucocorticoid resistance is reverted by LCK inhibition in pediatric
T-cell acute lymphoblastic leukemia. *Blood* **130**, 2750–2761 (2017).

9.   Buffière, A. *et al.* Saracatinib impairs maintenance of human T-ALL by targeting
the LCK tyrosine kinase in cells displaying high level of lipid rafts. *Leukemia* **32**,
2062–2065 (2018).

10. Campese, A. F. *et al.* Notch1-dependent lymphomagenesis is assisted by but
does not essentially require pre-TCR signaling. *Blood* **108**, 305–310 (2006).

11. Bellavia, D. *et al.* Combined expression of pTα and Notch3 in T cell leukemia
identifies the requirement of preTCR for leukemogenesis. *Proceedings of the*
National Academy of Sciences **99**, 3788–3793 (2002).

12. Smith, D. P., Bath, M. L., Harris, A. W. & Cory, S. T-cell lymphomas mask slower
developing B-lymphoid and myeloid tumours in transgenic mice with broad
haemopoietic expression of MYC. *Oncogene* **24**, 3544–3553 (2005).

13. dos Santos, N. R. *et al.* Pre-TCR expression cooperates with TEL-JAK2 to
transform immature thymocytes and induce T-cell leukemia. *Blood* **109**, 3972–
3981 (2007).

14. Tremblay, M. *et al.* Modeling T-cell acute lymphoblastic leukemia induced by the
SCL and LMO1 oncogenes. *Genes Dev* **24**, 1093–1105 (2010).

15. Liao, M.-J. *et al.* No Requirement for V(D)J Recombination in p53-Deficient
Thymic Lymphoma. *Mol Cell Biol* **18**, 3495–3501 (1998).

16.   Kelly, J. A. *et al.* Stat5 Synergizes with T Cell Receptor/Antigen Stimulation in the
525   Development of Lymphoblastic Lymphoma. *Journal of Experimental Medicine*
526   **198**, 79–89 (2003).
527   17.   Tsatsanis, C. *et al.* Tpl2 and ERK transduce antiproliferative T cell receptor signals
528   and inhibit transformation of chronically stimulated T cells. *Proceedings of the*
529   *National Academy of Sciences* **105**, 2987–2992 (2008).
530   18.   Gon, S. *et al.* Fit αβ T-cell receptor suppresses leukemogenesis of Pten-deficient
531   thymocytes. *Haematologica* **103**, 999–1007 (2018).
532   19.   Trinquand, A. *et al.* Triggering the TCR Developmental Checkpoint Activates a
533   Therapeutically Targetable Tumor Suppressive Pathway in T-cell Leukemia.
534   *Cancer Discov* **6**, 972–85 (2016).
535   20.   Lantz, O., Grandjean, I., Matzinger, P. & di Santo, J. P. γ chain required for naïve
536   CD4+ T cell survival but not for antigen proliferation. *Nat Immunol* **1**, 54–58 (2000).
537   21.   Bendelac, A., Matzinger, P., Seder, R. A., Paul, W. E. & Schwartz, R. H. Activation
538   events during thymic selection. *J Exp Med* **175**, 731–42 (1992).
539   22.   Azzam, H. S. *et al.* CD5 expression is developmentally regulated by T cell receptor
540   (TCR) signals and TCR avidity. *J Exp Med* **188**, 2301–11 (1998).
541   23.   O'Neil, J. *et al.* Activating Notch1 mutations in mouse models of T-ALL. *Blood* **107**,
542   781–785 (2006).
543   24.   Lin, Y.-W., Nichols, R. A., Letterio, J. J. & Aplan, P. D. Notch1 mutations are
544   important for leukemic transformation in murine models of precursor-T
545   leukemia/lymphoma. *Blood* **107**, 2540–2543 (2006).
546   25.   Nejman, D. *et al.* The human tumor microbiome is composed of tumor type–
547   specific intracellular bacteria. *Science (1979)* **368**, 973–980 (2020).
548   26.   Cho, J.-H. & Sprent, J. TCR tuning of T cell subsets. *Immunol Rev* **283**, 129–137
549   (2018).
550   27.   Brabb, T., Rubicz, R., Mannikko, V. & Goverman, J. Separately expressed T cell
551   receptor α and β chain transgenes exert opposite effects on T cell differentiation
552   and neoplastic transformation. *Eur J Immunol* **27**, 3039–3048 (1997).
553   28.   Strzadala, L., Miazek, A., Matuszyk, J. & Kisielow, P. Role of thymic selection in
554   the development of thymic lymphomas in TCR transgenic mice. *Int Immunol* **9**,
555   127–38 (1997).
556   29.   Cui, Y. *et al.* Thymic expression of a T-cell receptor targeting a tumor-associated
557   antigen coexpressed in the thymus induces T-ALL. *Blood* **125**, 2958–2967 (2015).
558   30.   Laffey, K. G. *et al.* Early expression of mature αβ TCR in CD4 − CD8 − T cell
559   progenitors enables MHC to drive development of T-ALL bearing NOTCH
560   mutations. *Proceedings of the National Academy of Sciences* **119**, (2022).

561   31.   Volanakis, E. J., Williams, R. T. & Sherr, C. J. Stage-specific Arf tumor
562   suppression in Notch1-induced T-cell acute lymphoblastic leukemia. *Blood* **114**,
563   4451–4459 (2009).

564   32.   Catarino, T. A. *et al.* Cdkn2a inactivation promotes malignant transformation of
565   mouse immature thymocytes before the β-selection checkpoint. *Exp Hematol* **116**,
566   30–36 (2022).

567   33.   Minden, M. D. *et al.* Chronic lymphocytic leukaemia is driven by antigen-
568   independent cell-autonomous signalling. *Nature* **489**, 309–312 (2012).

569   34.   Schmitz, R. *et al.* Burkitt lymphoma pathogenesis and therapeutic targets from
570   structural and functional genomics. *Nature* **490**, 116–120 (2012).

571   35.   Havranek, O. *et al.* Tonic B-cell receptor signaling in diffuse large B-cell
572   lymphoma. *Blood* **130**, 995–1006 (2017).

573   36.   Szydlowski, M. *et al.* FOXO1 activation is an effector of SYK and AKT inhibition in
574   tonic BCR signal-dependent diffuse large B-cell lymphomas. *Blood* **127**, 739–748
575   (2016).

576   37.   Bianchi, J. J., Murigneux, V., Bedora-Faure, M., Lescale, C. & Deriano, L.
577   Breakage-Fusion-Bridge Events Trigger Complex Genome Rearrangements
578   and Amplifications in Developmentally Arrested T Cell Lymphomas. *Cell Rep* **27**,
579   2847-2858.e4 (2019).

580   38.   Martín-Lorenzo, A. *et al.* Infection Exposure Is a Causal Factor in B-cell Precursor
581   Acute Lymphoblastic Leukemia as a Result of Pax5 -Inherited Susceptibility.
582   *Cancer Discov* **5**, 1328–1343 (2015).

583   39.   Serwold, T. *et al.* T-cell receptor-driven lymphomagenesis in mice derived from a
584   reprogrammed T cell. *Proceedings of the National Academy of Sciences* **107**,
585   18939–18943 (2010).

586   40.   Tran Quang, C. *et al.* Preclinical efficacy of humanized, non-FcγR-binding anti-
587   CD3 antibodies in T-cell acute lymphoblastic leukemia. *Blood* **136**, 1298–1302
588   (2020).

589   41.   dos Santos, N. R., Ghysdael, J. & Tran Quang, C. The TCR/CD3 complex in
590   leukemogenesis and as a therapeutic target in T-cell acute lymphoblastic
591   leukemia. *Adv Biol Regul* **74**, 100638 (2019).

592   42.   Scheinin, I. *et al.* DNA copy number analysis of fresh and formalin-fixed
593   specimens by shallow whole-genome sequencing with identification and exclusion
594   of problematic regions in the genome assembly. *Genome Res* **24**, 2022–32
595   (2014).

 Figure legends

Figure 1. HY-TCR transgene expression drives leukemia development. (A,B)
Kaplan-Meier leukemia-free survival curves of ETV6::JAK2 (male and female together)
and female ETV6::JAK2;Marilyn transgenic mice (A) and male (M) and female (F)
Marilyn transgenic mice (B). Log-rank test *P* values are shown in graphs. (C) Organ
weights of Marilyn females with leukemia/lymphoma compared to those of 7 month-old
wild-type (WT) females (n=3), or 10-week-old WT males for lymph nodes (n=4). For
Marilyn females n=16, except n=14 for kidneys. (D) White blood cell (WBC) counts of
Marilyn (n=15) and WT (n=3) females. Lower panels show blood smears of
representative WT and diseased Marilyn females. *P* values in (C) were obtained from
unpaired *t* tests with Welch's correction, and in (D) from Mann-Whitney test. Mean and
standard error of the mean (SEM) are shown. (E) Histogram representation of tumor
burden in lymphoid organs (thymus, spleen, and lymph nodes (LNs)) of Marilyn females
with or without thymic lymphoma. Female no. 12 had a mediastinal lymphoid tumor, while
no. 26 had lymphoid tumors on the duodenum wall. Each column represents a mouse.
Wild-type (WT) female mice were used as control.

Figure 2. Immunophenotypic analysis of HY-TCR leukemic cells. Flow cytometry
analysis of thymus-derived leukemic cells from representative Marilyn mice presenting
immature T cell phenotype, based on CD44 negativity and high CD24 expression (nos.
22, 33 and 40), mature T cell phenotype, based on CD44 positivity and CD24
intermediate/low expression (nos. 9 and 29), and mixed B and T cell phenotype, based
on B220 and CD90.2 expression (nos. 25 and 34). In CD44 and CD24 histogram plots,
unstained cells were used as negative control (grey shades). Healthy HY-TCR female
total thymocyte immunostaining is shown as control. TL, thymic lymphoma.

Figure 3. Transgenic HY-TCR expression in *Rag2*$^{-/-}$ mice drives development of T-
ALL with thymic lymphomas. (A) Kaplan-Meier leukemia-free survival curve for the
indicated genotypes and number of mice (n). Log-rank test *P* values are indicated. (B)
Histogram representation of lymphoid organ weight (thymus, spleen and lymph nodes
(LNs)) of Marilyn females with leukemia (left graph) and healthy (right graph). Each
column represents a mouse. (C) Flow cytometry immunostaining of cells from the thymus
(i) and indicated organs (ii) of two representative Marilyn;*Rag2*$^{-/-}$ leukemic mice with the
indicated surface markers. (D) Immunohistochemical Ki67 and cleaved-caspase 3

staining of representative thymic lymphoma sections from Marilyn;*Rag2*$^{-/-}$ and

Marilyn;*Rag2*$^{+/+}$ diseased mice. Black scale bars: 500 μm; green scale bar: 200 μm.

Figure 4. HY-TCR T-ALL antigenic stimulation *in vivo* hampers leukemia

development. (A) Percentage of CD45.1$^+$ T-ALL cells detected by flow cytometry in the

peripheral blood (PB) of male and female C57BL/6 mice infused with female HY-TCR T-

ALL cells. *P* value was obtained with two-way ANOVA. (B) Kaplan-Meier leukemia-free

survival curve for the same mice as in (A), which is representative of two experiments

performed with different primary leukemias. *P* value obtained with log-rank test. (C)

Spleens and kidneys (top panels) and respective organ weights (bottom panels) for male

(n=5) and female (n=6) C57BL/6 mice 21 days after infusion with HY-TCR T-ALL cells.

(D) Percentage of CD45.1$^+$CD90.2$^+$ HY-TCR T-ALL cells present in the bone marrow of

males and females shown in (C). *P* values in (C) and (D) were obtained from unpaired *t*

tests with Welch's correction. (E) Representative sections and quantification of anti-

cleaved caspase 3 immunohistochemistry for spleens from females and males shown in

(C,D; n=3). *P* value from unpaired *t* test. (C-E) Graphs show mean and SEM. (F) Kaplan-

Meier leukemia-free survival curves for syngeneic C57BL/6 nude or allogeneic CBA

nude male and female mice injected with female Marilyn T-ALL cells. Number of mice

and *P* values from log-rank test are indicated. (G) Kaplan-Meier leukemia-free survival

curves for syngeneic *Rag2*$^{-/-}$ females infused on day 0 with female Marilyn T-ALL cells

and treated once a week with DBY peptide (0.01 mg; n=5) or Control (n=4). Gray shade

represents treatment duration of 2 weeks.

Figure 1

Figure 2

Figure 3

Figure 4