Cell-autonomous and non-cell-autonomous effects of arginase-II on cardiac aging

Duilio M. Potenza[1]*, Xin Cheng[1], Guillaume Ajalbert[1], Andrea Brenna[1], Marie-Noelle Giraud[2], Aurelien Frobert[2]; Stephane Cook[2]; Kirsten D. Mertz[3], Zhihong Yang[1]*, Xiu-Fen Ming[1]*

[1]Laboratory of Cardiovascular and Aging Research and [2]Cardiology, Department of Endocrinology, Metabolism, and Cardiovascular System, Faculty of Science and Medicine, University of Fribourg, Switzerland; [3]Institute for Pathology, Cantonal Hospital, Baselland, Liestal, Switzerland

*Corresponding authors

Running title: Arginase-II promotes cardiac aging

Email addresses of authors :
Duilio M. Potenza: duilio.potenza@unifr.ch
Xiu-Fen Ming: xiu-fen.ming@unifr.ch
Zhihong Yang: zhihong.yang@unifr.ch

Statements relating to relevant ethics: Experimental work with animals was approved by the Ethical Committee of the Veterinary Office of Fribourg Switzerland (2020-01-FR) and performed in compliance with guidelines on animal experimentation at our institution. The study was conducted according to the principles expressed in the Declaration of Helsinki. Experiments with human heart biopsies were approved by the Ethics Committee of Northwestern and Central Switzerland (Project-ID 2021-01445).

Abstract

Aging is a predominant risk factor for heart disease. Aging heart reveals low-grade chronic inflammation, cell apoptosis, cardiac fibrosis, and increased vulnerability to ischemic injury. The underlying molecular mechanisms responsible for the cardiac aging phenotype and its susceptibility to injury are far from being fully understood. Although previous literature reports a role of the mitochondrial enzyme arginase-II (Arg-II) in development of heart failure, contradictory results are reported and no systematic analysis of cellular expression and localization of Arg-II in the heart has been performed. Whether and how Arg-II participates in cardiac aging are still unknown. In this study, we demonstrate, to our surprise, that Arg-II is not expressed in cardiomyocytes from aged mice and human patients, but upregulated in non-myocytes of the aging heart, including macrophages, fibroblasts, endothelial cells. Mice with genetic deficiency of *arg-ii* (*arg-ii$^{-/-}$*) are protected from age-associated cardiac inflammation, myocyte apoptosis, interstitial and perivascular fibrosis, endothelial-mesenchymal transition (EndMT), and susceptibility to ischemic injury. Further experiments show that Arg-II mediates IL-1β release from macrophages of old mice, contributing to the above-described cardiac aging phenotype. In addition, Arg-II enhances mitochondrial reactive oxygen species (mtROS) and activates cardiac fibroblasts that is inhibited by inhibition of mtROS. Thus, our study demonstrates a non-cell-autonomous effect of Arg-II on cardiomyocytes, fibroblasts, and endothelial cells mediated by IL-1β from aging macrophages as well as a cell-autonomous effect of Arg-II through mtROS in fibroblasts contributing to cardiac aging phenotype.

Introduction

Aging is a predominant risk factor for cardiovascular disease (CVD) (***Moturi et al., 2022***). Cardiac aging is accompanied by low-grade chronic inflammation termed "inflammaging" (***Cevenini et al., 2013***), which is linked to gradual development of cardiac fibrosis and heart failure, whereby macrophages, cardiac fibroblasts, and endothelial cells play critical roles in age-associated cardiac remodeling and dysfunction (***Abdellatif et al., 2023; Meschiari et al., 2017***). Studies in experimental animal models and in humans provide evidence demonstrating that aging heart is more vulnerable to stressors such as ischemia/reperfusion injury and myocardial infarction and exhibits less efficient reparative capability after injury as compared to the heart of young individuals (***Bujak et al., 2008; Mariani et al., 2000***). Even in the heart of apparently healthy individuals of old age, chronic inflammation, cardiomyocyte senescence, cell apoptosis, interstitial/perivascular tissue fibrosis, endothelial dysfunction and endothelial-mesenchymal transition (EndMT), and cardiac dysfunction either with preserved or reduced ejection fraction rate are observed (***Abdellatif et al., 2023; Ruiz-Meana et al., 2020***). Despite the intensive investigation in the past, the underlying causative cellular and molecular mechanisms responsible for the cardiac aging phenotype and the susceptibility of aging heart to injurious stressors are not fully elucidated, yet. Moreover, most preclinical studies preferentially use relatively juvenile animal models which hardly recapitulate clinical scenario. It is therefore critical to investigate mechanisms of cardiac aging phenotype with a naturally advanced aging animal model.

Research in the past years suggests a wide range of functions of the enzyme arginase, including the cytosolic isoenzyme arginase-I (Arg-I) and the mitochondrial isoenzyme arginase-II (Arg-II) in organismal aging as well as age-related organ structural remodeling and functional changes (***Caldwell et al., 2015; Li et al., 2022***). These functions of arginase have been implicated in cardiac and vascular aging (***Xiong, Yepuri, Montani, et al., 2017; Yepuri et al., 2012***) and cardiac ischemia/reperfusion injury in rodents (***Jung et al., 2010***). Arg-I, originally found in hepatocytes, plays a crucial role in urea cycle in the liver to remove ammonia from the blood (***Sin et al., 2015***), whereas Arg-II is inducible in many cell types under pathological conditions, contributing to chronic tissue inflammation and remodeling (***Moretto et al., 2019***). Both enzymes are able to catalyze the divalent cation-dependent hydrolysis of L-arginine (***Ming et al., 2013***). Studies in the literature including our own, demonstrate that Arg-II is involved in aging process, and promotes organ inflammation and fibrosis and inhibition of arginase or genetic ablation of *arg-ii* slows down aging process and protects against age-associated organ degeneration such as lung (***Zhu et al., 2023***), pancreas (***Xiong, Yepuri, Necetin, et al., 2017***) and heart (***Xiong, Yepuri, Montani, et al., 2017***). However, controversial results on expression of arginase and its isoforms at the cellular

levels in the heart are reported. While some studies linked Arg-I to the cardiac ischemia/reperfusion

injury (***Jung et al., 2010***), others suggest a role of Arg-II in this context (***Heusch et al., 2010***). Even

protective functions of Arg-II are suggested in some studies, based on correlation of Arg-II levels

with cardiac injury or systemic administration of Arg-II in rat models (***Huang et al., 2018; Lu et***

al., 2012). Importantly, cellular localization of arginase and its isoenzyme in the heart, i.e.,

expression of arginase in cardiomyocytes and non-cardiomyocytes have not been systematically

analyzed and confirmed. Finally, whether and how Arg-II participates in cardiac aging and whether

it is through cell-autonomous and/or paracrine effects are not known.

Therefore, our current study is aimed to investigate which cell types in the heart express

arginase and what enzymatic isoforms. How arginase and this isoenzyme in these cells influence

cardiac aging process and enhance susceptibility of the aging heart to ischemia/reperfusion injury.

                                          Results

1. Increased *arg-ii* expression in aging heart

In heart from young (3-4 months in age) and old (22-24 months in age) mice, an age-

associated increase in *arg-ii* expression in both *wt* male and female mice are observed (**Fig. 1A**).

This age-associated increase in *arg-ii* expression is more pronounced in females than in males

(**Fig. 1A**). In contrast to *arg-ii*, *arg-i* mRNA expression is very low in the heart (with very high

Ct values, data not shown), and no age-dependent changes of *arg-i* gene expression are observed

in *wt* and in *arg-ii*$^{-/-}$ mice of either gender (**Suppl. Fig. 1A** and **Suppl. Fig 1B**), suggesting a

specific enhancement of Arg-II in aging heart. Further experiments are thus focused on Arg-II

and female mice. We made effort to analyze Arg-II protein levels by immunoblotting in the heart

tissues. However, immunoblotting is not sensitive enough to detect Arg-II protein in the heart

tissue of either young or old animals. Therefore, further experiments are performed to detect

Arg-II proteins at the cellular level with confocal microscopic immunofluorescence staining as

described in later sections.

2. *Arg-ii* ablation reduces cardiac aging phenotype

Next we investigated the impact of *arg-ii* ablation on cardiac phenotype. Masson's staining

reveals an enhanced peri-vascular and interstitial fibrosis in *wt* old mice, which is mitigated in

age-matched *arg-ii*$^{-/-}$ animals (**Fig. 1B, 1C**). The age-associated increase in cardiac fibrosis and

its inhibition by *arg-ii*$^{-/-}$ are further confirmed by quantitative measurements of collagen content

as assayed by determination of hydroxyproline levels in the heart tissues (**Fig. 1D**). The results

demonstrate that *arg-ii*$^{-/-}$ prevents age-associated fibrosis in the heart. Furthermore, an increased

density of fibroblasts as characterized by positive staining of PDGF-Rα is observed in *wt* old

mice, but not in age-matched *arg-ii*$^{-/-}$ animals (**Fig. 1E** and **Fig. 1F**).

Moreover, a significant increase in macrophage marker *f4/80* (**Fig. 2A**) and elevated gene

expression of numerous pro-inflammatory cytokines such as *mcp1*, *il-1β* and *tnf-α* are observed

in aging heart of *wt* mice (**Fig. 2B to Fig. 2D**). This age-associated increase in the inflammatory

markers and cytokine gene expression (except *tnf-α*) are overall significantly reduced or

prevented in *arg-ii*$^{-/-}$ mice (**Fig. 2A** to **Fig. 2D**). Furthermore, an age-associated increase in total

numbers of apoptotic cells as demonstrated by TUNEL staining is observed in the heart of *wt*

mice, which is reduced in age-matched *arg-ii*$^{-/-}$ animals (**Fig. 2E** and **Fig. 2F**). Further

experiments with wheat germ agglutinin (WGA)-Alexa Fluor 488 (used to define cell boarders)

(**Fig. 2G**) show a higher percentage of apoptotic cardiomyocytes (CM) than non-cardiomyocytes

(NCM) in *wt* mice (**Fig. 2H**). This increased percentage of cardiomyocyte apoptosis is also

prevented in age-matched *arg-ii⁻/⁻* animals (**Fig. 2H**). The percentage of apoptotic non-
cardiomyocytes in aged *arg-ii⁻/⁻* mice tended to be reduced as compared to *wt* mice, but it does
not reach statistical significance (**Fig. 2H**).

In addition, heart from old *wt* mice reveals elevated levels of SNAIL (the master regulator of
End-MT process) and vimentin (mesenchymal marker), which is prevented in *arg-ii⁻/⁻* mice (**Fig.**
3A to **Fig. 3C**). No changes in CD31 (endothelial marker) levels are observed among the four
groups (**Fig. 3A** and **Fig. 3D**). Immunofluorescence staining reveals that the age-related up-
regulation of vimentin is mostly due to expansion of fibroblast population in left ventricle as
shown in Fig. 1E and Fig. 1F). Confocal co-immunofluorescence staining shows that vimentin
could be found in CD31⁺ endothelial cells of blood vessels in the heart of old *wt* but rarely in
age-matched *arg-ii⁻/⁻* mice (**Fig. 3E**). The results suggest at least a partial End-MT occurring in
cardiac aging that is prevented by *arg-ii* ablation. With aging, the heart weight to body weight
ratio (HW/BW) increases to a similar extent between *wt* and *arg-ii⁻/⁻* mice (**Suppl. Fig. 2**),
suggesting no difference in age-related cardiac hypertrophy between *wt* and *arg-ii⁻/⁻* mice.

3. Cellular localization of Arg-II in aging heart

Next, the cellular localization of Arg-II in aging heart are investigated. Since *arg-ii* levels
are upregulated mostly in aged females, heart tissues from old female mice are used mainly for
this purpose. To our surprise, confocal co-immunofluorescence staining does not show Arg-II in
cardiomyocytes as evidenced by lack of Arg-II staining in troponin T⁺-cells, a specific
cardiomyocyte marker (**Fig. 4A**). Arg-II is only found in non-cardiomyocytes (**Fig. 4A**). In line
with this notion, Arg-II is not detectable by immunoblotting in isolated primary cardiomyocytes
from old *wt* mice at baseline or after exposure to hypoxia (1 % O₂, 24 hours, **Fig. 4B**), a well
know strong stimulus for Arg-II expression in many cell types (***Liang et al., 2019; Liang et al.,***
2021; Ren et al., 2022; Zhu et al., 2023). In contrast to cardiomyocytes, cardiac fibroblasts
isolated from old *wt* mice express Arg-II which is up-regulated by hypoxia (**Fig. 4B**). The
absence of *arg-ii* mRNA in cardiomyocytes in contrast to fibroblasts isolated from *wt* old mouse
hearts is also confirmed with qRT-PCR (**Fig. 4C**). Moreover, Arg-II is detected in Mac-2⁺
macrophages (**Fig. 4D**), CD31⁺ endothelial cells (**Fig. 4E**), and PDGF-Rα⁺-fibroblasts (**Fig. 4F**),
but not in α-SMA⁺ vascular smooth muscle cells (**Fig. 4G**). These results demonstrate that Arg-II
in aging mouse heart is mostly expressed in non-cardiomyocytes cells such as macrophages,
endothelial cells and fibroblasts. Similar to mouse heart, in human heart (obtained from a 66-
year-old woman, and a 76-year-old man with no significant cardiac pathology), Arg-II is also
absent in cardiomyocytes as proved by lack of co-localization with troponin-T⁺-cells (**Fig. 4H**),
and is expressed in CD31⁺-endothelial cells (**Fig. 4I**), CD68⁺-macrophages (**Fig. 4J**), and

vimentin$^+$- (**Fig. 4K)** or α-SMA$^+$- (**Fig. 4L)** fibroblasts/myofibroblasts. In contrast to mouse

heart, Arg-II could be found in very few vascular smooth muscle cells in human heart (**Fig. 4M**).

4. Role of Arg-II in crosstalk between macrophages and cardiomyocytes through IL-1β

Given that age-associated cardiomyocyte apoptosis is prevented in *arg-ii$^{-/-}$* mice despite

absence of Arg-II in the cardiomyocytes, the non-cell-autonomous effects of Arg-II on

cardiomyocytes are investigated. As reported above, macrophage numbers and *il-1β* expression

are increased in aging heart and reduced in *arg-ii$^{-/-}$* mice (Fig. 2A and 2C). Both immunoblotting

and immunofluorescence staining confirm that Il-1β protein levels are higher in old *wt* heart

tissues when compared to age-matched *arg-ii$^{-/-}$* (**Fig. 5A to 5D**). Further co-immunofluorescence

staining reveals that Il-1β is localized in Mac2$^+$- macrophages (**Fig. 5E**). Therefore, the role of

IL-1β in crosstalk between macrophages and cardiac cells such as cardiomyocytes, fibroblasts

and endothelial cells is further explored. Considering that ablation of *arg-ii* gene specifically

reduces the number of TUNEL-positive cardiomyocytes (Fig. 2G and 2H), we hypothesized that

Arg-II in macrophages may prime cardiomyocyte apoptosis. To test this hypothesis, splenic

macrophages are isolated from young and old *wt* and age-matched *arg-ii$^{-/-}$* mice, and conditioned

medium from the cells are collected. Elevated Arg-II and IL-1β protein levels are observed in old

wt macrophages as compared to the cells from young animals (**Fig. 6A to 6C**). In line with this

finding, IL-1β levels in conditioned medium from old *wt* mouse macrophages are higher as

compared to the conditioned medium from young macrophages (**Fig. 6D**). Interestingly, *arg-ii*

ablation substantially reduces IL-1β levels in the macrophages (**Fig. 6A, 6C,** and **6D).**

Importantly, cardiomyocytes isolated from adult (five-months) female *wt* mice as bioassay cells,

when incubated with conditioned medium from old (but not from young) *wt* macrophages for 24

hours (**Fig. 6E**), display more apoptosis as monitored by TUNEL assay (**Fig. 6F** and **6G**). This

age-associated cardiac damaging effect of macrophages is not observed with *arg-ii$^{-/-}$* macrophage

conditioned medium (**Fig. 6F** and **6G**), suggesting a role of Arg-II expressing macrophages in

mediating cardiomyocyte apoptosis in aging. To further examine whether IL-1β from

macrophages is the paracrine mediator for cardiomyocyte apoptosis, the above described

experiment is performed in presence of the IL-1 receptor antagonist IL-Ra. Indeed,

cardiomyocyte apoptosis induced by the conditioned medium from old *wt* macrophages is

abolished by IL-Ra (**Fig. 6F** and **6G**).

Similar results are obtained with Raw264.7 cells, a widely used mouse macrophage model.

Raw264.7 cells stimulated with lipopolysaccharide (LPS; 100 nmol/L) for 24 hours, show a

significant increase in both Arg-II and IL-1β, which is inhibited by silencing *arg-ii* (**Suppl. Fig.**

3A to **Suppl. Fig. 3C**), confirming the role of Arg-II in the production of IL-1β in macrophages

(**Suppl. Fig. 3A** and **Suppl. Fig. 3C**). In agreement with the results shown in Fig. 6, conditioned

medium from LPS-treated Raw264.7 cells significantly enhances cardiomyocyte apoptosis as

assessed by TUNEL staining, which is reduced by either *arg-ii* gene silencing or by IL-1 receptor

blocker ILRa (**Suppl. Fig. 3D** and **Suppl. Fig. 3E**). It is of note that that LPS alone does not

induce cardiomyocyte apoptosis at the concentration tested (data not shown), confirming the role

of Arg-II-mediated IL-1β release from macrophages in cardiomyocyte apoptosis.

5. Role of Arg-II in crosstalk between macrophages and cardiac fibroblasts through IL-1β

Since cardiac fibrosis is accompanied with macrophage accumulation in our aging heart

model, a crosstalk between macrophages and fibroblasts is then investigated. Cardiac fibroblasts

isolated from adult (five-months) female *wt* mice as bioassay cells, produce more hydroxyproline

when they are incubated with conditioned medium from old as compared to young *wt*

macrophages (**Suppl. Fig. 4A**). This effect of old macrophages on fibroblasts is reduced with

conditioned medium from *arg-ii$^{-/-}$* macrophages (**Fig. 7A**). In parallel, expression levels of genes

involved in extracellular matrix remodeling such as *fibronectin, collagen IIIα, tgf-β1* (**Fig. 7B** to

Fig. 7D) as well as *mmp2* and *mmp9* (**Suppl. Fig. 4B** and **Suppl. Fig. 4C**) are reduced in

fibroblasts stimulated with conditioned medium from *arg-ii$^{-/-}$* macrophages as compared to the

old *wt* macrophages, while expression of *collagen Iα* is not altered under this condition (**Suppl.**

Fig. 4D). Importantly, the increase in hydroxyproline levels in cardiac fibroblasts incubated with

conditioned medium from old *wt* macrophages is inhibited by ILRa (**Fig. 7E**).

6. Cell-autonomous effects of Arg-II in cardiac fibroblasts: role of mtROS

Since Arg-II is expressed in fibroblasts, a cell-autonomous effect of Arg-II in these cells is

also investigated in cultured human cardiac fibroblasts (HCF). Overexpression of *arg-ii* in the

cells (**Fig. 7F nd 7G**) increases *vimentin* and *collagen IIIα* expression levels (**Fig. 7H and 7I),**

demonstrating a cell-autonomous effect of Arg-II in cardiac fibroblasts. Interestingly,

upregulation of *collagen IIIα* expression by *arg-ii* is accompanied with increased mitochondrial

ROS (mtROS) generation, and inhibition of mtROS by mito-TEMPO (10 μmol/L) prevents *col-*

IIIa upregulation (**Fig. 7J, 7K and 7L**), suggesting a role of Arg-II-induced mtROS in activation

of fibroblasts.

7. Role of Arg-II in crosstalk between macrophages and endothelial cells through IL-1β
243

Results in Fig. 3 suggests EndMT in aging heart, which is regulated by Arg-II. The role of

Arg-II-expressing macrophages in EndMT of endothelial cells is then investigated in cultured

cells. Human endothelial cells exposed to macrophage-derived conditioned medium from old

(not young) *wt* mice for 96 hours has no effect on VE-cadherin but enhances N-cadherin and

vimentin (mesenchymal markers), and also Arg-II protein levels (**Fig. 8A** to **Fig. 8E**), suggesting

a partial End-MT induced by aged macrophages. Interestingly, these effects of macrophages are

prevented in old *arg-ii$^{-/-}$* mice (**Fig. 8A** to **Fig. 8E**) or abolished in the presence of IL-1 receptor

antagonist IL-Ra (**Fig. 8F** to **Fig. 8J)**. The results suggest a role of IL-1β from old *wt*

macrophages in EndMT process.

8. Arg-II gene ablation reduces ischemic damage in aging heart

To study the specific role of Arg-II in heart ischemia/reperfusion (I/R) injury during aging,

ex vivo experiments with Langendorff-perfused heart from old *wt* and *arg-ii$^{-/-}$* mice were

performed followed by cardiac functional analysis. The experimental protocol of global

ischemia/reperfusion is shown in **Fig. 9A**. Under normoxic baseline condition, the cardiac

functions, i.e., dP/dt$_{max.}$ (maximal rate of contraction), dP/dt$_{min.}$ (maximal rate of relaxation), and

left ventricular developed pressure (LVDP) are comparable between *wt* and *arg-ii$^{-/-}$* old mice

(**Suppl. table 3**). *Arg-ii* ablation improves cardiac function following I/R injury, i.e., the

inotropic contractile function as measured by LVDP and dP/dt$_{max}$, lusitropic dP/dt$_{min}$ (**Fig. 9B**).

The post I/R recovery is significantly improved in *arg-ii$^{-/-}$* mice when compared to age-matched

wt animals (**Fig. 9B**). In addition to functional recovery, infarct size after I/R-injury was

analyzed by triphenyl tetrazolium chloride (TTC) staining in old mice. Under the

ischemic/reperfusion condition (20 minutes ischemia and 30 minutes reperfusion), no myocardial

infarct was detected (**Suppl. Fig. 5**), while longer ischemia and reperfusion time (30 minutes and

45 minutes respectively) is required to induce myocardial infarct, i.e., $10.5 \pm 2.9\%$ in *wt* old

mice, which is significantly reduced in the age-matched *arg-ii$^{-/-}$* mice ($3.2 \pm 0.9\%$, **Fig. 9C** and

Fig. 9D). These results reveal a role of Arg-II in vulnerability of aging heart to I/R injury.

Discussion

Previous studies have reported the presence of both Arg-I and Arg-II isoenzymes in the cardiac tissues of various animal models including feline, pig, rats, rabbit, and mice (***Gonon et al., 2012; Heusch et al., 2010; Jung et al., 2006; Jung et al., 2010; Steppan et al., 2006***). These isoenzymes have been implicated in cardiac ischemic injury and cardiac aging (***Jung et al., 2010; Xiong, Yepuri, Montani, et al., 2017***). It has been widely assumed that Arg-I and Arg-II are expressed in cardiomyocytes, where they play a role in cardiac injury via a cell-autonomous effect on the cardiomyocytes. However, the cellular localization of arginase isoforms has not been convincingly demonstrated due to the following two reasons: species-specific expression of arginase isoforms and lack of including negative controls with gene knockout samples.

Our current study systematically investigated the expression and cellular localization of the mitochondrial Arg-II in the heart of a natural aging mouse model and in human biopsies. We found an age-associated increase in *arg-ii* but not *arg-i* in heart of male and female mice. Data extracted from a high throuput sequencing database of male mice (no data on females are available, unfortunately) confirms our findings showing the age-dependent enhancement of *arg-ii* but not *arg-i* expression in the heart (GSE201207) ((***Wolff et al., 2023***). Surprisingly, we found that neither *arg-ii* gene expression nor its protein levels were detectable in cardiomyocytes from either young or old mice. Despite increased *arg-ii* mRNA (but not *arg-i* mRNA) expression in the aging heart, mitochondrial Arg-II was absent from cardiomyocytes. This observation aligns with previous findings in vascular endothelial cells from mice and humans, where Arg-II (not Arg-I) is the predominant isoenzyme (***Ming et al., 2004***). Our study further reveals that Arg-II is not detectable in cardiomyocytes but is instead expressed in non-cardiomyocyte cells such as macrophages, fibroblasts, and endothelial cells in aged mouse heart and human cardiac biopsies. These results suggest that Arg-II exerts non-cell-autonomous effects on aging cardiomyocytes.

It is well known that the aging heart is associated with increased cardiomyocyte apoptosis, chronic inflammation, oxidative stress, and cardiac fibrosis (***Vakka et al., 2023***). In the current study, we observe that *arg-ii*$^{-/-}$ mice are protected from cardiac aging phenotype and reveal a decrease in apoptotic cells, particularly cardiomyocytes but also non-cardiomyocytes. Since cardiomyocytes do not express Arg-II in aging and Arg-II could not even be induced in these cells under hypoxic condition, a well-known strong inducer of Arg-II via HIFs (***Liang et al., 2021***), a non-cell-autonomous effect of Arg-II on cardiomyocyte apoptosis must be involved, meaning that the increased cardiomyocyte apoptosis in aging must be due to paracrine effects of other cell types such as macrophages in which Arg-II levels are elevated in aging. It is evident that crosstalk between cardiomyocytes and non-cardiomyocytes are critical in cardiac functional changes and remodeling (***Hulsmans et al., 2016***). Substantial evidence demonstrates that immune cells,

particularly, monocytes and/or infiltrated macrophages, are important contributors to cardiac remodeling in aging and disease conditions (**Hulsmans et al., 2016; Jimenez et al., 2022; Wang et al., 2020**). Under acute and chronic ischemic heart disease conditions, blood monocytes can be mobilized from bone marrow and spleen, and recruited into the injury site of the heart where they play an important role in cardiac remodeling (**Honold et al., 2018; Swirski et al., 2009**). An increased accumulation of myeloid cells including in aging heart has been reported and contributes to cardiac inflammaging (**Esfahani et al., 2021**). In line with these studies, we demonstrate an increased macrophage accumulation in aging heart associated with elevated expression of numerous inflammatory cytokines in macrophages. This age-associated cardiac inflammation is inhibited in *arg-ii*[-/-] animals. Among these inflammatory cytokines, an increase in IL-1β levels in cardiac tissue and macrophages or monocytes/macrophages derived from spleen of old mice are demonstrated accompanied with elevated Arg-II levels as compared to young animals. Importantly, we showed that Arg-II in the aged macrophages enhances IL-1β expression and release mediating cardiomyocyte apoptosis, since the cardiomyocyte apoptosis caused by the aged macrophages is inhibited in *arg-ii*[-/-] mice and prevented by IL-1β receptor antagonist. This results further confirm our previous findings demonstrating a pro-inflammatory role of Arg-II in infiltrated macrophages in vascular atherosclerosis and high-fat-diet induced type 2 diabetes mouse models (**Ming et al., 2012**). This conclusion is also further supported by the findings with a mouse macrophage cell line RAW264.7. The question what are the underlying mechanisms that upregulate Arg-II levels in macrophages in aging remains to be investigated. Moreover, we also found increased apoptosis of other cell types in aging heart, which is also prevented in *arg-ii*[-/-] mice. What are these cells, how this cell apoptosis is triggered, and what is the impact on age-associated cardiac vulnerability to stressors require further investigation.

One of the important features of cardiac aging is fibrosis which is due to chronic inflammation process (**Lu et al., 2017**). Studies provide evidence that with aging process, cardiac resident macrophages decrease and are progressively replaced by pro-inflammatory recruited macrophages with an increase in inflammatory microenvironment (**Molawi et al., 2014**). There are convincing evidences demonstrating that chronic inflammation facilitates development of myocardial fibrosis in heart failure which is associated with aging (**Lillo et al., 2023**). A crosstalk between macrophages and fibroblasts has been demonstrated to be important for cardiac fibrosis in diseased conditions (**Hartupee et al., 2016**). Studies show that monocyte-derived macrophages from bone marrow and spleen are recruited in advanced age, contributing to cardiac fibrosis and myocardial dysfunction (**Hulsmans et al., 2018**). In our present study using an in vitro cellular model, we show a pivotal role of Arg-II-expressing macrophages in activation of cardiac fibroblasts in aging heart. First, Arg-II protein is present in the cardiac macrophages and enhanced in

monocyte-derived macrophages in aging mice; second, conditioned medium from aged *arg-ii⁻/⁻* macrophages exhibits decreased potential of stimulating cardiac fibroblasts to produce components that are essentially involved in fibrosis such as hydroxyproline, *fibronectin, coll-IIIa,* and *tgfb1.* Interestingly, the enhanced hydroxyproline levels stimulated by old *wt* mouse macrophage-derived conditioned medium is inhibited by the IL-1β receptor blocker, demonstrating a paracrine effect of IL-1β release from aged macrophages on cardiac fibroblast activation. Moreover, the age-associated cardiac fibrosis and increase in fibroblast number in the heart is prevented in *arg-ii⁻/⁻* mice, demonstrating a crosstalk between macrophage and fibroblasts in cardiac aging.

The fact that cardiac fibroblasts in aging heart express elevated Arg-II, indicates a contribution of a cell-autonomous effect of Arg-II in this cell type to cardiac fibrosis in aging. This concept is confirmed by the experiments showing that overexpression of *arg-ii* in cardiac fibroblasts enhances vimentin levels associated with increased expression of *col-IIIa* and mitochondrial ROS generation. Importantly, inhibition of mitochondrial ROS by TEMPO abolished the effect of *arg-ii* overexpression in the fibroblasts, demonstrating that Arg-II exerts its cell-autonomous stimulating effects in fibroblasts through mitochondrial ROS. This finding is supported by the studies showing that ROS favors cardiac fibroblast proliferation and activation to produce collagen (***Alili et al., 2014; Ohtsu et al., 2005***). With these results, we demonstrate that cardiac fibroblasts are activated in aging by a paracrine effect of macrophages, that is mediated by Arg-II / IL-1β and also by a cell-autonomous effect of Arg-II through mtROS.

Endothelial cells play a key role in regulation of blood flow to organs through eNOS pathway, one of the most important regulatory mechanisms in cardiovascular system (***Godo et al., 2017***). Previous studies demonstrated a causal role of Arg-II in promotion of vascular endothelial aging through eNOS uncoupling (***Yepuri et al., 2012***). In line with this, our current study showed that Arg-II is expressed in the coronary vascular endothelium of old mice and also in human heart. Emerging evidence also suggests that endothelial cells have a significant capacity for plasticity and can change their phenotype, including endothelial-to-mesenchymal transition (EndMT), a process that endothelial cells undergoing extensive alterations, including changes in gene and protein expression, loss of the endothelial phenotype, and switch to mesenchymal phenotype such as increased cell migration, proliferation, and increased collagen production, contributing to cardiovascular disease and fibrosis (***Murdoch et al., 2014; Zeisberg et al., 2007; Zhang et al., 2021***). Although EndMT in cardiac fibrotic models, particularly in pro-atherosclerotic mouse models are at least partially evidenced, it is less known in cardiovascular aging (***Evrard et al., 2016; Kidder et al., 2023; Xu et al., 2023***). When compared to young animals, aged *wt* heart tissues show enhanced levels of SNAIL, the master regulator of End-MT process. Moreover, immunofluorescence reveals that CD31[+] endothelial cells also co-express vimentin (mesenchymal

marker). No changes of endothelial marker expression such as CD31 and vascular-endothelial cadherin (data not shown) are appreciated, suggesting that partial End-MT occurs during aging. Ablation of *arg-ii* blunts the age-related elevation of both vimentin and SNAIL, thereby reducing the number of CD31$^+$/vimentin$^+$ endothelial cells. This data hints that lower cardiac tissue fibrosis in *arg-ii*$^{-/-}$ is at least partially contributed by dampened End-MT process. Indeed, conditioned medium from old *wt* macrophages is capable of enhancing mesenchymal markers N-cadherin and vimentin, as well as Arg-II protein levels in endothelial cells without changes in endothelial markers such as VE-cadherin, confirming a partial End-MT induced by aged macrophages. Interestingly, this effect of macrophages is prevented when *arg-ii* was ablated. Furthermore, the paracrine effects of the old macrophages on EndMT are prevented by IL-1 receptor antagonist ILRa, demonstrating a role of IL1β from old *wt* macrophages expressing Arg-II in EndMT process. Moreover, Arg-II is expressed and upregulated in endothelial cells with aging (***Yepuri et al., 2012***). Overexpression of *arg-ii* in the endothelial cells did not affect the End-MT markers (data not shown), suggesting that Arg-II alone is not sufficient to induce a cell- autonomous effect in End-MT. Finally, we demonstrate that *arg-ii*$^{-/-}$ mice in aging exhibit improved cardiac function and are more resistant to ischemic/reperfusion stress and reveal a better recovery and less myocardial infarct area as compared to the old *wt* animals. This protection against age-related cardiac dysfunction and vulnerability to ischemic stress in *arg-ii*$^{-/-}$ animals could be explained by the above described cardiac aging phenotype such as increase in apoptosis of cardiac cells, decrease in macrophage-mediated inflammation, endothelial dysfunction and EndMT, and fibroblast activation.

Previous studies provided sex-related difference in Arg-II levels at both mRNA (***Xiong, Yepuri, Montani, et al., 2017***) and protein levels in various organs (***Huang et al., 2021; Xiong, Yepuri, Necetin, et al., 2017***). We further confirmed the observation that in female mice, there is a greater age-associated upregulation of *arg-ii* levels in the heart, which is specifically found in non-cardiomyocytes but not in cardiomyocytes. A possible explanation for this sex-specific phenomenon may lie in differences in hormonal patterns between male and female animals throughout the aging process. Indeed, sex hormones, including estrogen and testosterone, have been shown to regulate *arg-ii* expression in various tissues. Estrogen seems to downregulate Arg-II (***Hayashi et al., 2006***). The age-related greater increase in *arg-ii* in the female mice might be due to loss or lack of estrogen in old female animals, which requires further investigation. However, testosterone has been reported to upregulate *arg-ii* expression (***Levillain et al., 2005***). This observation, however, does not seem to support the hypothesis that increased *arg-ii* expression in aged male mouse heart is related to sex hormones. Nevertheless, how age enhances Arg-II in sex-specific manner remains a topic of further investigation. Of note, previous studies showed that sex-related difference in pathological phenotypes in the aging kidney, pancreas, and lung are associated with higher levels of Arg-II in the

females (*Huang et al., 2021; Xiong, Yepuri, Necetin, et al., 2017; Zhu et al., 2023*).

In conclusion, our study demonstrates that age-related upregulation of Arg-II in macrophage,

fibroblasts, and endothelial cells contributes to the cardiac aging phenotype. This includes

manifestations, such as cardiac inflammatory response, partial EndMT, fibroblast activation,

resulting in cardiomyocyte apoptosis, cardiac fibrosis, and ultimately, myocardial dysfunction.

Moreover, it enhances the vulnerability of aging heart to ischemic/reperfusion injury, primarily

through a non-cell-autonomous paracrine release of IL-1β from macrophages acting on

cardiomyocytes and endothelial cells. This effect is in addition to a cell-autonomous effect of Arg-II

in cardiac fibroblasts (**Fig. 10**). Future work shall investigate the cell-autonomous and non-cell-

autonomous effects of *arg-ii* using cell-specific conditional knockout animal models. Nevertheless,

our study strongly suggests that targeting Arg-II in macrophages, fibroblasts, and endothelial cells

emerges as a promising approach to prevent or reduce age-related cardiac dysfunction and protect

heart under diseased conditions.

 Materials and methods

Arg-II$^{-/-}$ mouse and sample preparation

Wild type (*wt*) and *arg-ii* knockout (*arg-ii$^{-/-}$*) mice were kindly provided by Dr. William O'Brien

(**Shi et al., 2001**) and back crossed to C57BL/6 J for more than 10 generations. Genotypes of mice

were confirmed by polymerase chain reaction (PCR) as previously described. Offspring of *wt* and

arg-ii$^{-/-}$ mice were generated by interbred from hetero/hetero cross. Mice were housed at 23°C with

a 12-h-light-dark cycle. Animals were fed a normal chow diet and had free access to water and

food. The animals were selected and included in the study according to age, gender, and genotype.

Only healthy animals, based on scoring of activity, posture, general appearance (coat, skin),

locomotion, eyes/nose, body weight loss, were used. In this study, confounders such as the

animal/cage location were controlled. All experimenters were aware of the group allocation at the

different stages of the experiments. Male and female mice at the age of 3–4 months (young) or 20–

22 months (old) were euthanized under deep anesthesia (i.p. injection of a mixture of

ketamine/xylazine 50 mg/kg and 5 mg/kg, respectively) and death was confirmed by absence of all

the reflexes and by exsanguination. The heart was then isolated and either snap frozen in liquid

nitrogen and kept at −80 °C until use or fixed with 4% paraformaldehyde (pH 7.0), and then

embedded in paraffin for immunofluorescence staining experiments. Experimental work with

animals was approved by the Ethical Committee of the Veterinary Office of Fribourg Switzerland

(2020-01-FR) and performed in compliance with guidelines on animal experimentation at our

institution and in accordance with the updated ARRIVE guidelines (**Percie du Sert et al., 2020**).

Experiments with human heart biopsies were approved by the Ethics Committee of Northwestern

and Central Switzerland (Project-ID 2021-01445).

Langendorff-perfused mouse heart preparation

Ex vivo functional assessments were performed by using an isolated heart system (Langendorff

apparatus – EMKA technologies, France). Briefly, after euthanasia as above described, the hearts

were excised, cannulated and retrogradely perfused on the Langendorff system. 150 µL of heparin

(2,000 units/ml) was injected 15 minutes before the heart removal to prevent blood coagulation.

Hearts trimmed of extracardiac tissue were weighed immediately after surgical removal prior

cannulation and heart weight to body weight ratio (HW/BW) was calculated as hypertrophy index.

The aorta was cannulated on a shortened and blunted 21-gauge needle and perfusion initiated at a

constant pressure of 80 mmHg. The hearts were perfused with a modified Krebs–Henseleit solution

composed of (mmol/L): 118 NaCL, 4.7 KCl, 1.2 MgSO$_4$, 1.2 KH$_2$PO$_4$, 25 NaHCO$_3$, 11 Glucose, 2

CaCl$_2$. The fluid was bubbled with a mix of 95% O$_2$ and 5% CO$_2$ at 37°C to give a pH of 7.4 and

was filtered through an in-line 0.45 μm Sterivex-HV filter before delivery to the heart. A fluid-filled balloon was introduced into the left ventricle through an incision in the left atrial appendage. The ventricular balloon was connected via fluid-filled tubing to a pressure transducer for continuous assessment of ventricular function. The balloon was inflated to yield a left ventricular end-diastolic pressure between 5 to 15 mmHg. Hearts were immersed in warmed perfusate in a jacketed bath maintained at 37°C, and perfusate delivered to the coronary circulation was maintained at the same temperature. After 20 minutes stabilization, hearts were subjected to 20 to 30 minutes of no-flow ischemia followed by 30-45 minutes of reperfusion. Recovery of contraction and relaxation rates (dt/dp$_{min}$) as well as the recovery of the left ventricle developed pressure (LV-DP) and left ventricular end diastolic pressure (LV-EDP) were measured to investigate tolerance to ischemia/reperfusion (I/R)-injury. In addition to functional recovery, triphenyl tetrazolium chloride (TTC) staining was also performed to investigate a potential reduction in infarct size upon I/R-injury in *arg-ii$^{-/-}$* animals. After completing the I/R protocol, heart was carefully removed from the experimental set-up. After a freeze-thaw cycle (-20°C for 1 hour), the heart was cross-sectioned from the apex to the atrioventricular groove into four 2.5 mm thick slices. The slices were first incubated in TTC (20 minutes at 37°C), and then fixed in 10% formalin for 30 minutes. A cover glass was then placed over the tissues and images were acquired (***Redfors et al., 2012***). ImageJ software was used to analyze the images and assess the infarcted area.

Real-time quantitative RT-PCR

mRNA expression level of genes from mouse and human origin was measured by two-step quantitative Real Time-PCR as described previously (***Ming et al., 2012***). Ribosomal Protein S12 (*rps12*), and glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) were used as reference genes. Total RNA from mouse tissue, splenic macrophages, and human cardiac fibroblasts was extracted with Trizol Reagent (TR-118, Molecular Research Center) following the manufacturer's protocol. Real-time PCR reaction was performed with the GOTaq® qPCR Master Mix (A6001, Promega) and CFX96 Real-Time PCR Detection System (Bio-Rad). The mRNA expression level of all genes was normalized to the reference gene *rps12* or *gapdh*. All the RT-PCR primer sequences are shown in the **Suppl. Table 1**.

Confocal immunofluorescence microscope

Immunofluorescence staining was performed with heart tissues. Briefly, mouse hearts or human cardiac biopsies were fixed with 4% paraformaldehyde (pH 7.0) and embedded in paraffin. After deparaffinization in xylene (2 times of 10 minutes), the sections were treated in ethanol (twice in 100% ethanol for 3 minutes and twice in 95% ethanol and once in 80%, 75%, 50% ethanol for

1 minute sequentially) followed by antigen retrieval (EDTA buffer, pH 8.0) for Arg-II, troponin-T (TropT), CD31, Mac-2, IL-1β, vimentin (Vim), CD-68, PDGF-Rα and α-smooth muscle actin (αSMA) in a pressure cooker. For co-immunofluorescence staining of Arg-II/TropT, Arg-II/CD31, Arg-II/Vim, Arg-II/CD-68, Arg-II/Mac-2, Arg-II/PDGF-Rα, Arg-II/αSMA, IL-1β/Mac-2 and CD31/Vim, primary antibodies of different species were used. Transverse sections (5 μm) were blocked with mouse Ig blocking reagent (M.O.M, Vector laboratories) for 2 hours and then with PBS containing 1% BSA and 10% goat serum for 1 hour. For CD31 and PDGF-Rα (both polyclonal Goat IgGs), a blocking reagent containing 10% BSA was used. The sections were then incubated overnight at 4°C in a dark/humidified chamber with target primary antibodies, and subsequently incubated for 2 hours with the following secondary antibodies: Alexa Fluor 488–conjugated goat anti-rabbit IgG (H + L) and Alexa Fluor 568-conjugated goat anti-mouse IgG (H + L); Alexa Fluor 488–conjugated goat anti-mouse IgG (H + L) and Alexa Fluor 594-conjugated goat anti-rabbit IgG (H + L); Alexa Fluor 488–conjugated donkey anti-goat IgG (H + L) and Alexa Fluor 594-conjugated goat anti-rabbit IgG (H + L). All sections were finally counterstained with 300 nmol/L DAPI for 5 minutes. Immunofluorescence signals were visualized under Leica TCS SP5 confocal laser microscope. The antibodies used are shown in **Suppl. Table 2.**

Isolation and cultivation of ventricular cardiomyocytes

Ventricular myocytes were isolated from five-month old female mice according to an established protocol (*Louch et al., 2011*). Briefly, the hearts were excised, cannulated and retrogradely perfused on a Langendorff system as described above. 150 μl of heparin (2,000 units/ml) was injected 15 minutes before the heart removal to prevent blood coagulation. To isolate the cardiomyocytes, hearts were perfused at 37°C for 20 minutes with a Ca^{2+}-free solution composed of (in mmol/ liter): 130 NaCl, 5 KCL, 0.5 NaH_2PO_4, 10 HEPES, 10 Glucose, 10 2,3 butanedione monoxime (BDM), 10 Taurine, 1 $MgCl_2$ (pH 7.4, adjusted with NaOH). Cells were enzymatically dissociated using a cocktail of collagenase type II (210 U/ml, Worthington), collagenase type IV (260 U/ml, Worthington) and protease type XIV (0.21 U/ml; Sigma-Aldrich). After isolation, cardiomyocytes were separated by gravity settling and Ca^{2+} was restored to a final concentration of 1 mmol/L. The cardiomyocytes were finally plated onto laminin-coated coverslips (5 μg/ml, Thermo Scientific) and cultured in the M-199 medium supplemented with bovine serum albumin (BSA; 0.1 % - A1470, Sigma-Aldrich), BDM (10 mmol/L), insulin, transferrin, selenium (ITS; I3146 - Sigma-Aldrich), CD lipid (11905-031 - Sigma-Aldrich) and penicillin / streptomycin mix (1 %).

Cell culture and adenoviral transduction

Human cardiac fibroblasts (HCF) from adult human heart tissue were purchased from Innoprot (P10452) and cultured in the proprietary fibroblast culture medium (P60108-2, Innoprot) composed as follow: 500 ml of fibroblast basal medium, 25 ml of FBS, 5 ml of fibroblast growth supplement-2, and 5 ml of penicillin/streptomycin in the Poly-L-Lysine (PLL) coated flasks and dishes. The cells were maintained at 37°C in a humidified incubator (5% CO_2 and 95% atmosphere air). To overexpress *arg-ii*, the cells were seeded on 6-cm dishes for 24 hours and transduced with rAd-CMV-Con/*arg-ii* for 48 hours as described previously (***Ming et al., 2012***).

Human umbilical vein endothelial cells (HUVEC) cells were cultured in RPMI-1640 (with 25 mmol/L HEPES and stable glutamine; Thermo Fisher Scientific) containing 5 % fetal calf serum (FCS; Gibco) and 1% streptomycin and penicillin in gelatin (1%) coated dishes.

Raw 264.7 cells (mouse macrophage cell line) are cultured in Dulbecco modified Eagle medium/F12 (DMEM/F12; Thermo Fisher Scientific) containing 10% fetal bovine serum (FBS; Gibco) and 1% streptomycin and penicillin. To silence *arg-ii*, the cells were seeded on 6-cm dishes for 24 hours and transduced first with the rAd at titers of 100 Multiplicity of Infection (MOI) and cultured in the complete medium for 2 days and then in serum-free medium for another 24 hours before experiments. The rAd expressing shRNA targeting *arg-ii* driven by the U6 promoter (rAd/U6-*arg-ii*-shRNA) and control rAd expressing shRNA targeting *lacz* (rAd/U6-*lacz*-shRNA) were generated as described previously (***Ming et al., 2009***).

TUNEL Staining

Mouse hearts were isolated and fixed with 4% paraformaldehyde (pH 7.0), embedded in paraffin and sliced at 5 μm of thickness. After deparaffinization (xylene, 2 times for 10 minutes) and rehydration (sequential washing in 100%, 95%, 80%, 75% and 50% ethanol), TUNEL staining was performed using a commercial kit (in situ Cell Death Detection Kit, TMR red – Roche, Switzerland). Briefly, 50 μl of TUNEL mixture, containing TdT and dUTP in reaction buffer were added onto the sections and incubated in a humidified chamber for 60 minutes at 37°C in darkness. Positive controls were generated by incubating samples at room temperature (RT) for 10 minutes with DNase in PBS (1 mg/ml) to induce strand breaks. Negative controls were generated by incubation with TUNEL mixture lacking TdT. DAPI counterstaining was finally performed (300 nmol/l for 5 minutes).

TUNEL staining was also performed in isolated ventricular cardiomyocytes. Upon plating onto murine laminin-coated coverslips, the cardiomyocytes were first incubated for 24 hours with either splenic macrophage conditioned media (CM; see below) or with RAW cells CM, and the following protocol was applied: cells are fixed using 4% paraformaldehyde for 1 hour at RT, and then permeabilized via Triton X (0.1 %). As for tissues, 50 μl of TUNEL mixture, containing TdT

and dUTP in reaction buffer were added onto each slide, and then incubated at 37°C for 60 minutes in darkness. DAPI counterstaining was then performed.

For experiments aiming to define cell borders, TUNEL-protocol was followed by a staining with Wheat Germ Agglutinin (WGA)-Alexa Fluor 488 (10 µg/ml for 30 minutes). For heart tissue, five representative images of each section are taken using a Leica TCS SP5 confocal system, and WGA staining was performed to differentiate cardiomyocytes and non-cardiomyocytes according to cell size and presence or absence of striations (characteristics of cardiomyocytes). Data was reported as number of TUNEL-positive nuclei over the total number of nuclei in the images.

Masson's trichrome staining

To analyze cardiac fibrosis, heart sections (5 µm) were subjected to Masson's trichrome (ab150686, Abcam) staining according to the manufacturer's instructions (*de Guzman et al., 2023*).

Hydroxyproline colorimetric assay

Collagen content was analyzed by determination of hydroxyproline levels in left ventricles or in cell homogenates using the Hydroxyproline Assay kit (MAK008, Sigma) according to the manufacturer's instructions.

Isolation of splenic macrophages

Splenic cells were isolated as previously described (*Wang et al., 2013*). Briefly, young and old *wt* and *arg-ii*[-/-] female mice were euthanized as described and spleens were dissected from abdominal cavity and filtered through a 70-µm nylon strainer. Red blood cell lysis buffer was used to remove red blood cells. The cell suspension obtained was plated onto gelatin-coated dishes at a density of 2 x 10^6 cells/ml. After 3 days, cells were washed to ensure macrophage enrichment. Following washing, the macrophages were cultured for further 4 days in RPMI culture media with 20% of L929 cell line conditioned medium (source of the macrophage growth factor M-CSF) to allow cell maturation. Medium was changed every two days.

Crosstalk between macrophages and heart cells

For crosstalk experiments of splenic macrophages from *wt* and *arg-ii*[-/-] mice with cardiac cells, conditioned medium from the isolated splenic macrophages was collected and transferred to cardiomyocytes, fibroblasts and endothelial cells in 1:1 dilution with the specific cell culture medium for the indicated time. To study effects of IL-1β in the conditioned medium of splenic macrophages (CM-SM), cardiac cells, fibroblasts or endothelial cells were pre-treated with IL-1

receptor antagonist IL-1ra (50 ng/ml) for 2 hours followed by the specific experimental protocols

for analysis of cardiac apoptosis, fibroblast activation, and EndMT transition, respectively.

For the experiments with mouse macrophage cell line Raw 264.7, the cells were seeded on

six-well plates with a density of 2×10^5 cells per well. Upon transduction with rAd expressing *arg-*

ii^{shRNA} (see above) and over-night serum starvation, the mouse macrophages were polarized toward

a pro-inflammatory secretion profile by stimulation with lipopolysaccharide (LPS; 100 nmol/L) for

24 hours. The conditioned medium (CM) from non-treated control and LPS-activated RAW cells

(referred to as CM-RAW) was then filtered and transferred onto cardiomyocytes for 24 hours to

investigate cell apoptosis. To study effects of IL-1β in CM-RAW on cardiac cells apoptosis,

myocytes were pre-treated with IL-1 receptor antagonist IL-1ra (280-RA, R&D systems) for 2

hours.

Hypoxia experiments with murine cardiac fibroblasts and cardiomyocytes

Cardiac cells were isolated from *wt* mice according to an established protocol (***Louch et al., 2011***).

Briefly, after euthanasia of mice, hearts were excised, cannulated and retrogradely perfused on a

Langendorff system. Cells were enzymatically dissociated using a cocktail of collagenase type II,

collagenase type IV and protease type XIV. After isolation, cardiomyocytes were separated from the

pull of cardiac cells by gravity settling and plated onto laminin-coated coverslips. After 2 hours, the

cardiomyocytes were washed to remove non-adhering cells and incubated for 24 hours at 1%

oxygen level in a Coy In Vitro Hypoxic Cabinet System (The Coy Laboratory Products, Grass

Lake, MI USA). The rest of cardiac cells were plated on 12-well plates and cultured in DMEM

(10% FBS) for 3 days, allowing cardiac fibroblasts to adhere. The fibroblasts were passaged once

and subjected to hypoxia as for the cardiomyocytes.

ELISA for IL-1β detection in splenic cell conditioned media

IL-1β concentrations in the conditioned medium from murine splenic cells were measured by

ELISA kits (mouse IL-1β, 432604, BioLegend, San Diego, USA) according to the manufacturer's

instructions.

Immunoblotting

Heart tissue or cell lysate preparation, SDS-PAGE and immunoblotting, antibody incubation, and

signal detection were performed as described previously (***Ming et al., 2012***). To prepare heart

homogenates, frozen cardiac tissues were crushed into the fine powder using a mortar and pestle in

liquid nitrogen on ice. A portion of the fine powder was then homogenized (XENOX-

Motorhandstück MHX homogenizer) on ice in 150 µl of ice-cold lysis buffer with the following

635 composition: 10 mmol/L Tris-HCl (pH 7.4), 0.4 % Triton X-100, 10 μg/ml leupeptin, and 0.1

636 mmol/l phenylmethylsulfonyl fluoride (PMSF), protease inhibitor cocktail (B14002, Bio-Tool) and

637 phosphatase inhibitor cocktail (B15002; Bio-Tool). Homogenates were centrifuged (Sorvall Legend

638 Micro 17R) at $13,000 \times g$ for 15 minutes at $4\,°C$. Protein concentrations of supernatants were

639 determined by Lowry method (500-0116, Bio-Rad). Equal amount of protein from each sample was

640 heated at 75°C for 15 minutes in loading buffer and separated by SDS-PAGE electrophoresis.

641 Proteins in the SDS-PAGE gel were then transferred to PVDF membranes which are blocked with

642 PBS-Tween-20 supplemented with 5% skimmed milk. The membranes were then incubated with

643 the corresponding primary antibody overnight at 4°C with gentle agitation. After washing with

644 blocking buffer, the membranes were then incubated with the corresponding anti-mouse or anti-

645 rabbit secondary antibody. Signals were visualized using the Odyssey Infrared Imaging System (LI-

646 COR Biosciences), or the FUSION FX Imaging system (Witec AG) for chemiluminescence, and

647 quantified by Image Studio Lite (5.2, LI-COR Biosciences). The antibodies used in this study are

648 listed in the **Suppl. Table 2.**

650 *Statistical analysis*

651 Data were presented as mean ± SD. Data distribution was determined by Kolmogorov-Smirnov test

652 and statistical analysis for normally distributed values was performed with Student's unpaired t-test

653 or analysis of variance (ANOVA) with Bonferroni post hoc test. For non-normally distributed

654 values, Mann–Whitney test or the Kruskal–Wallis test were used. Differences in mean values were

655 considered statistically significant at a two tailed p-value ≤ 0.05.

Author contributions

DMP, XC, GA, AB, M-NG, AF, and SC performed experiments, data acquisition, analysis, and

interpretation, KDM provided human samples; DMP, ZY and X-FM designed the project and

prepared figures and drafted the manuscript; ZY and X-FM received funding for the study; All the

authors discussed the results and critically revised and approved the manuscript for important

intellectual contents.

Conflict of interest: The authors declare no conflicts of interest.

Funding: This work was supported by grants from the Swiss National Science Foundation

(31003A_179261/1 to ZY) and Swiss Heart Foundation (FF19033 to X-FM).

Availability of data and materials: Data supporting the present study are available from the

corresponding author upon reasonable request.

                                                    Figures legend

Fig. 1. Age-associated increase in Arg-II levels in mouse heart. (**A**) *arg-ii* mRNA levels of male

and female young (3-4 months) and old (20-22 months) wild type (*wt*) heart tissues analyzed by

qRT-PCR. *rps12* served as the reference (n=6 to 7 animals per group); (**B**) Representative

histological images of heart interstitial and perivascular fibrosis in young and old *wt* and *arg-ii*$^{-/-}$

female mice. Fibrosis is shown by the blue-colored Trichrome Masson's staining. Scale bar = 50

μm. (n=5 to 7 mice per group); (**C**) Quantification of total fibrotic area in cardiac tissue (% of total

area); (**D**) Hydroxyproline content of mouse heart from young and old *wt* and *arg-ii*$^{-/-}$ female mice.

(n=4 mice in each group); (**E**) Representative confocal images showing immunofluorescence

staining of PDGF-Rα (green, fibroblasts marker) in young and old *wt* and *arg-ii*$^{-/-}$ heart tissue. DAPI

(blue) is used to stain nuclei. Scale bar = 50 μm; (**F**) Relative PDGF-Rα signal quantification of

confocal images (n= 4 per each group). The values shown are mean ± SD. Data are presented as the

fold change to the young-*wt* group, except for panel D. *P ≤ 0.05, **P ≤ 0.01 between the indicated

groups. *wt*, wild-type mice; *arg-ii*$^{-/-}$, *arg-ii* gene knockout mice.

Fig. 2. Age-associated elevation in inflammatory cytokines and apoptosis in heart is prevented

in female *arg-ii*$^{-/-}$ mice. mRNA expression levels of (**A**) *f4/80*, (**B**) *mcp-1*, (**C**) *il-1β* and (**D**) *tnf-α* in

young and old *wt* and *arg-ii*$^{-/-}$ female mouse hearts were analyzed by qRT-PCR. *rps12* served as the

reference. (n=6-8 mice per group); (**E-F**) Representative confocal images and relative quantification

of apoptotic cardiac cells in young and old *wt* and *arg-ii*$^{-/-}$ heart tissue. DAPI (blue) is used to stain

nuclei. Scale bar = 50 μm; (**G**) wheat germ agglutinin (WGA)-Alexa Fluor 488-conjugate was used

to stain cell membrane, and separate cardiomyocytes and non-cardiomyocytes apoptotic cells. Cell

distinction was based on cell size and shape. Scale bar = 10 μm; (**H**) graphs showing the

quantification of the TUNEL-positive cardiomyocytes (CM) and non-myocytes (NCM) in old *wt*

and *arg-ii*$^{-/-}$ hearts. The values shown are mean ± SD. Data are expressed as fold change to the

young *wt* group, expect for panel H. *p ≤ 0.05, **p≤ 0.01, ***p ≤ 0.005 and ****p ≤ 0.001

between the indicated groups. *wt*, wild-type mice; *arg-ii*$^{-/-}$, *arg-ii* gene knockout mice.

Fig. 3. **Age-related endothelial-to-mesenchymal transition (EndMT) in heart tissue**. (**A**)

Immunoblotting analysis of CD31 (endothelial marker), vimentin (mesenchymal marker) and

SNAIL (master regulator of EndMT) in the heart of *wt* and *arg-ii*$^{-/-}$ female young and old mice;

tubulin and vinculin served as protein loading controls. Molecular weight (kDa) is indicated at the

side of the blots. The plot graphs show the quantification of the SNAIL (**B**), vimentin (**C**) and CD31

(**D**) signals on immunoblots (n=6-10 mice in each group); (**E**) Representative confocal images

showing co-localization of CD31 (green) and vimentin (red) in young and old *wt* and *arg-ii⁻/⁻* heart tissues. DAPI (blue) is used to stain nuclei. Scale bar = 20 μm. *p≤0.05 and **p ≤ 0.01, between the indicated groups. *wt*, wild-type mice; *arg-ii⁻/⁻*, *arg-ii* gene knockout mice.

Fig. 4. Cellular localization of Arg-II in aging heart of female mice. (**A**) Confocal microscopy illustration of immunofluorescence double staining of Arg-II (red) and tropoin-t (TnT, green; cardiomyocytes marker). Scale bar = 25 μm; (**B**) Bright-field microscopy images of isolated *wt* cardiomyocytes and primary cardiac fibroblasts. The immunoblot shows the level of Arg-II in both cardiomyocytes and fibroblasts upon exposure to hypoxia (1 % O_2) for 24 h. C indicates freshly isolated cardiomyocytes used as control, N and H indicates respectively normoxia and hypoxia conditions, and K indicates kidney tissue extract used as positive control. Tubulin served as protein loading control; (**C**) mRNA expression levels of *arg-ii* in cardiomyocytes (card) and non-cardiomyocytes (non-c) cells isolated from old *wt* and *arg-ii⁻/⁻* female mouse hearts. *gapdh* served as the reference. (n=3-5 mice per group); (**D** to **G**) Representative confocal images of old *wt* mouse heart showing co-localization of (**D**) Arg-II (red) and Mac-2 (green, mouse macrophage marker), (**E**) Arg-II (red) and CD31 (green, endothelial marker), (**F**) Arg-II (red) and PDGF-Rα (green, fibroblasts marker), and (**G**) Arg-II and α-smooth muscle actin (α-SMA; green, smooth muscle cell/myofibroblasts marker); (**H** to **M**) Representative confocal images of human heart tissue showing co-localization of (**H**) Arg-II (green) and tropoin-t (red, cardiomyocytes marker), (**I**) Arg-II (red) and CD31 (green, endothelial marker), (**J**) Arg-II (red) and CD-68 (green, macrophage marker), (**K**) Arg-II (red) and vimentin (green, fibroblast marker), and (**L-M**) Arg-II (red) and α-smooth muscle actin (α-SMA; green, (**L**) myofibroblasts and (**M**) smooth muscle cell marker). DAPI (blue) stains cell nuclei. Scale bar = 20 μm. Each experiment was repeated with 3 to 5 animals.

Fig. 5. **Arg-II ablation reduces Il-1β protein levels in aging heart.** (**A**) Immunoblotting analysis of Il-1β (pro-inflammatory cytokine) in the heart of old *wt* and *arg-ii⁻/⁻* female mice; tubulin served as protein loading control. Molecular weight (kDa) is indicated at the side of the blots; (**B**) The plot graph shows quantification of the Il-1β signals on immunoblots (n=6-7 mice in each group); (**C**) Representative confocal images showing Il-1β localization in old *wt* and *arg-ii⁻/⁻* heart tissues. DAPI (blue) is used to stain nuclei. Scale bar = 50 μm; (**D**) Relative Il-1β signal quantification of confocal images (n= 9 per each group); (**E**) Representative confocal images showing co-localization of Mac-2 (green, mouse macrophage marker), and Il-1β (red) in *wt* heart tissues. This experiment was repeated with 3 animals. Scale bar = 10 μm. *p≤0.05 and ***p ≤ 0.005, between the indicated groups. *wt*, wild-type mice; *arg-ii⁻/⁻*, *arg-ii* gene knockout mice.

Fig. 6. *In vitro* **study of crosstalk between aging splenic macrophages and cardiomyocytes.** (**A**) Immunoblotting analysis of Arg-II and Il-1β in mouse splenic macrophages isolated from young and old *wt* and *arg-ii$^{-/-}$* female mice; tubulin served as protein loading control. Molecular weight (kDa) is indicated at the side of the blots. The plot graphs show the quantification of Arg-II (**B**) and Il-1β (**C**) protein signals on immunoblots (n=3 mice in each group); (**D**) IL-1β levels in the conditioned medium from young and old *wt* and *arg-ii$^{-/-}$* splenic macrophages was measured by ELISA; (**E**) Schematic representation of *in vitro* crosstalk study; (**F**) Representative confocal images of adult isolated mouse cardiomyocytes stimulated with conditioned media (CM) from young and old, *wt* and *arg-ii$^{-/-}$* splenic cells (24 h incubation). Wheat Germ Agglutinin (WGA)-Alexa Fluor 488-conjugate was used to stain cell membrane, and TUNEL was performed to identify apoptotic cells. DAPI is used to stain nuclei. Interleukin receptor antagonist (ILRa; 50 ng/mL) is used to prevent IL-1β binding to its receptor; (**G**) Quantification of TUNEL-positive cardiomyocytes (% in respect to total number of cells). Scale bar = 20 μm. *p≤0.05, **p ≤ 0.01, ***p ≤ 0.005 and ****p ≤ 0.001 between the indicated groups. MØ, splenic macrophage; Y, young; O, old; *wt*, wild-type mice; *arg-ii$^{-/-}$*, *arg-ii* gene knockout mice.

Fig. 7. **Crosstalk between splenic macrophages and cardiac fibroblasts and cell-autonomous effect of Arg-II in cardiac fibroblasts.** (**A**) Collagen production measured as hydroxyproline content in mouse *wt* fibroblasts treated with conditioned media (CM) from old, *wt* and *arg-ii$^{-/-}$* splenic cells (96 h incubation) (n=3 mice in each group). The values shown are mean ± SD. mRNA expression levels of (**B**) *fibronectin*, (**C**) *collagen-IIIα (col-IIIα)* and (**D**) *tgf-β1* in *wt* fibroblasts treated with conditioned media (CM) from old *wt* and *arg-ii$^{-/-}$* splenic cells (96 h incubation) were analyzed by qRT-PCR. *gapdh* served as the reference. (n=4 mice per group); (**E**) Hydroxyproline content in fibroblasts treated with CM from old *wt* splenic cells (96 h incubation). ILRa (50 ng/mL) is used to prevent IL-1β binding to its receptor (n=4 independent experiments). (**F**) Immunoblotting analysis of Arg-II and vimentin in human cardiac fibroblasts (HCFs) upon *arg-ii* gene overexpression. GAPDH served as protein loading control; (**G**) qRT-PCR analysis of mRNA expression levels *arg-ii* in HCF cells; (**H**) The plot graph shows the quantification of the vimentin signals on immunoblots shown in panel F. (n=3 independent experiments); (**I**) qRT-PCR analysis of mRNA expression levels of *collagen-IIIα* in HCF cells. *gapdh* served as the reference. (n=6 independent experiments); (**J**) Representative confocal images of human cardiac fibroblasts (HCF) upon transfection with rAd-CMV-Con/Arg-II for 48 h. MitoSOX (Red) is used to stain mitochondrial ROS (mtROS). TEMPO (10 μmol/L) is used to prevent mtROS generation. DAPI (blue) stains cell nuclei. Scale bar = 50 μm. (**K**) Mitosox signal quantification (n=3 independent

experiments); (**L**) qRT-PCR analysis of mRNA expression levels of *col-IIIα* in HCF cells treated as indicated. *gapdh* served as the reference. (n=5 independent experiments). Data are expressed as fold change to respective control group. *p≤0.05, **p ≤ 0.01, ***p ≤ 0.005 and ****p ≤ 0.001 between the indicated groups. MØ, splenic macrophage; Con, control.

Fig. 8. **Aging *wt* macrophages induces EndMT in HUVEC. (A)** Immunoblotting analysis of VE-Cadherin (endothelial marker), N-Cadherin and vimentin (both mesenchymal markers), and Arg-II in HUVEC cells upon incubation with conditioned media (CM) from young and old *wt* and *arg-ii^{-/-}* splenic cells (4 days incubation); GAPDH served as protein loading control. Molecular weight (kDa) is indicated at the side of the blots. The plot graphs show the quantification of Arg-II (**B**), N-Cadherin (**C**), vimentin (**D**), and VE-Cadherin (**E**) protein levels (n=3 independent experiments); (**F**) Immunoblotting analysis of VE-Cadherin, N-Cadherin, vimentin and Arg-II in HUVEC cells upon incubation with CM from old *wt* splenic cells (4 days incubation); Interleukin receptor antagonist (IL-Ra; 50 ng/mL) is used to prevent IL-1β effect. GAPDH served as protein loading control. Molecular weight (kDa) is indicated at the side of the blots. The plot graphs show the quantification of Arg-II (**G**), N-Cadherin (**H**), vimentin (**I**), and VE-Cadherin (**J**) protein signals on immunoblots (n=4 independent experiments); Data are expressed as fold change to respective control group. *p≤0.05, **p ≤ 0.01 and ***p ≤ 0.005 between the indicated groups. MØ, splenic macrophage; *wt*, wild-type mice; *arg-ii^{-/-}*, *arg-ii* gene knockout mice.

Fig. 9. Arg-II ablation improves heart function recovery from global ischemia/reperfusion injury (I/R-I). (**A**) Protocol applied for e*x vivo* Langendorff-heart functional assessment of old *wt* and age-matched *arg-ii^{-/-}* female mice. After baseline recordings, 20 minutes global ischemia is followed by 30 minutes of reperfusion; (**B**) *Ex vivo* Langendorff-heart assessment of old female *wt* (black line) and *arg-ii^{-/-}* (red line) hearts. The graphs show the functional recovery of the left ventricular developed pressure (LVDP), left ventricular end diastolic pressure (LVED) and maximal rate of contraction (dP/dt$_{max}$) and relaxation (dP/dt$_{min}$). The data are expressed as % of recovery in respect to baseline values and represent the mean ± SD of data from 5 mice per group; (**C**) Representative sections of *wt* and *arg-ii^{-/-}* female hearts stained with 2,3,5-TTC and (**D**) relative quantification of infarcted areas. *p≤0.05 and **p ≤ 0.01 between the indicated groups. *wt*, wild-type mice; *arg-ii^{-/-}*, *arg-ii* gene knockout mice.

Fig. 10. Schematic illustration of the role of Arg-II and underlying mechanisms in cardiac aging.

References

Abdellatif M, Rainer PP, Sedej S, Kroemer G. 2023. Hallmarks of cardiovascular ageing. *Nat Rev Cardiol*. doi:10.1038/s41569-023-00881-3

Alili L, Sack M, Puschmann K, Brenneisen P. 2014. Fibroblast-to-myofibroblast switch is mediated by NAD(P)H oxidase generated reactive oxygen species. *Bioscience Reports* **34**. doi:10.1042/BSR20130091

Bujak M, Kweon HJ, Chatila K, Li N, Taffet G, Frangogiannis NG. 2008. Aging-related defects are associated with adverse cardiac remodeling in a mouse model of reperfused myocardial infarction. *J Am Coll Cardiol* **51**:1384-1392. doi:10.1016/j.jacc.2008.01.011

Caldwell RB, Toque HA, Narayanan SP, Caldwell RW. 2015. Arginase: an old enzyme with new tricks. *Trends in Pharmacological Sciences* **36**:395-405. doi:10.1016/j.tips.2015.03.006

Cevenini E, Monti D, Franceschi C. 2013. Inflamm-ageing. *Curr Opin Clin Nutr Metab Care* **16**:14-20. doi:10.1097/MCO.0b013e32835ada13

de Guzman RC, Meer AS, Mathews AA, Israel AR, Moses MT, Sams CM, Deegan DB. 2023. Reduced fibrous capsule elastic fibers from biologic ECM-enveloped CIEDs in minipigs, supported with a novel compression mechanics model. *Biomed Mater Eng* **34**:289-304. doi:10.3233/BME-221488

Esfahani NS, Wu Q, Kumar N, Ganesan LP, Lafuse WP, Rajaram MVS. 2021. Aging influences the cardiac macrophage phenotype and function during steady state and during inflammation. *Aging Cell* **20**:e13438. doi:10.1111/acel.13438

Evrard SM, Lecce L, Michelis KC, Nomura-Kitabayashi A, Pandey G, Purushothaman KR, d'Escamard V, Li JR, Hadri L, Fujitani K, Moreno PR, Benard L, Rimmele P, Cohain A, Mecham B, Randolph GJ, Nabel EG, Hajjar R, Fuster V, Boehm M, ... 2016. Endothelial to mesenchymal transition is common in atherosclerotic lesions and is associated with plaque instability. *Nat Commun* **7**:11853. doi:10.1038/ncomms11853

Godo S, Shimokawa H. 2017. Endothelial Functions. *Arterioscler Thromb Vasc Biol* **37**:e108-e114. doi:10.1161/ATVBAHA.117.309813

Gonon AT, Jung C, Katz A, Westerblad H, Shemyakin A, Sjoquist PO, Lundberg JO, Pernow J. 2012. Local arginase inhibition during early reperfusion mediates cardioprotection via increased nitric oxide production. *PLoS One* **7**:e42038. doi:10.1371/journal.pone.0042038

Hartupee J, Mann DL. 2016. Role of inflammatory cells in fibroblast activation. *J Mol Cell Cardiol* **93**:143-148. doi:10.1016/j.yjmcc.2015.11.016

Hayashi T, Esaki T, Sumi D, Mukherjee T, Iguchi A, Chaudhuri G. 2006. Modulating role of estradiol on arginase II expression in hyperlipidemic rabbits as an atheroprotective mechanism. *Proc.Natl.Acad.Sci.U.S.A* **103**:10485-10490.

Heusch P, Aker S, Boengler K, Deindl E, van de Sand A, Klein K, Rassaf T, Konietzka I, Sewell A, Menazza S, Canton M, Heusch G, Di Lisa F, Schulz R. 2010. Increased inducible nitric oxide synthase and arginase II expression in heart failure: no net nitrite/nitrate production and protein S-nitrosylation. *Am J Physiol Heart Circ Physiol* **299**:H446-453. doi:10.1152/ajpheart.01034.2009

Honold L, Nahrendorf M. 2018. Resident and Monocyte-Derived Macrophages in Cardiovascular Disease. *Circ Res* **122**:113-127. doi:10.1161/CIRCRESAHA.117.311071

Huang J, Liang X, Ladeiras D, Fellay B, Ming XF, Yang Z. 2021. Role of tubular epithelial arginase-II in renal inflammaging. *NPJ Aging Mech Dis* **7**:5. doi:10.1038/s41514-021-00057-8

Huang XW, Pan MD, Du PH, Wang LX. 2018. Arginase-2 protects myocardial ischemia-reperfusion injury via NF-kappaB/TNF-alpha pathway. *Eur Rev Med Pharmacol Sci*

22:6529-6537. doi:10.26355/eurrev_201810_16067

Hulsmans M, Sager HB, Roh JD, Valero-Munoz M, Houstis NE, Iwamoto Y, Sun Y, Wilson RM,
Wojtkiewicz G, Tricot B, Osborne MT, Hung J, Vinegoni C, Naxerova K, Sosnovik DE, Zile
MR, Bradshaw AD, Liao R, Tawakol A, Weissleder R, ... 2018. Cardiac macrophages
promote diastolic dysfunction. *J Exp Med* **215**:423-440. doi:10.1084/jem.20171274

Hulsmans M, Sam F, Nahrendorf M. 2016. Monocyte and macrophage contributions to cardiac
remodeling. *J Mol Cell Cardiol* **93**:149-155. doi:10.1016/j.yjmcc.2015.11.015

Jimenez J, Lavine KJ. 2022. The Dynamic Role of Cardiac Macrophages in Aging and Disease.
Curr Cardiol Rep **24**:925-933. doi:10.1007/s11886-022-01714-4

Jung AS, Kubo H, Wilson R, Houser SR, Margulies KB. 2006. Modulation of contractility by
myocyte-derived arginase in normal and hypertrophied feline myocardium. *Am J Physiol*
Heart Circ Physiol **290**:H1756-1762. doi:10.1152/ajpheart.01104.2005

Jung C, Gonon AT, Sjoquist PO, Lundberg JO, Pernow J. 2010. Arginase inhibition mediates
cardioprotection during ischaemia-reperfusion. *Cardiovasc Res* **85**:147-154.
doi:10.1093/cvr/cvp303

Kidder E, Pea M, Cheng S, Koppada SP, Visvanathan S, Henderson Q, Thuzar M, Yu X, Alfaidi M.
2023. The interleukin-1 receptor type-1 in disturbed flow-induced endothelial mesenchymal
activation. *Front Cardiovasc Med* **10**:1190460. doi:10.3389/fcvm.2023.1190460

Levillain O, Diaz JJ, Blanchard O, Dechaud H. 2005. Testosterone down-regulates ornithine
aminotransferase gene and up-regulates arginase II and ornithine decarboxylase genes for
polyamines synthesis in the murine kidney. *Endocrinology* **146**:950-959.
doi:10.1210/en.2004-1199

Li Z, Wang L, Ren Y, Huang Y, Liu W, Lv Z, Qian L, Yu Y, Xiong Y. 2022. Arginase: shedding light
on the mechanisms and opportunities in cardiovascular diseases. *Cell Death Discov* **8**:413.
doi:10.1038/s41420-022-01200-4

Liang X, Arullampalam P, Yang Z, Ming XF. 2019. Hypoxia Enhances Endothelial Intercellular
Adhesion Molecule 1 Protein Level Through Upregulation of Arginase Type II and
Mitochondrial Oxidative Stress. *Front Physiol* **10**:1003. doi:10.3389/fphys.2019.01003

Liang X, Potenza DM, Brenna A, Ma Y, Ren Z, Cheng X, Ming XF, Yang Z. 2021. Hypoxia Induces
Renal Epithelial Injury and Activates Fibrotic Signaling Through Up-Regulation of
Arginase-II. *Front Physiol* **12**:773719. doi:10.3389/fphys.2021.773719

Lillo R, Graziani F, Franceschi F, Iannaccone G, Massetti M, Olivotto I, Crea F, Liuzzo G. 2023.
Inflammation across the spectrum of hypertrophic cardiac phenotypes. *Heart Fail Rev*
28:1065-1075. doi:10.1007/s10741-023-10307-4

Louch WE, Sheehan KA, Wolska BM. 2011. Methods in cardiomyocyte isolation, culture, and gene
transfer. *J Mol Cell Cardiol* **51**:288-298. doi:10.1016/j.yjmcc.2011.06.012

Lu L, Guo J, Hua Y, Huang K, Magaye R, Cornell J, Kelly DJ, Reid C, Liew D, Zhou Y, Chen A,
Xiao W, Fu Q, Wang BH. 2017. Cardiac fibrosis in the ageing heart: Contributors and
mechanisms. *Clinical and Experimental Pharmacology and Physiology* **44 Suppl 1**:55-63.
doi:10.1111/1440-1681.12753

Lu MJ, Chen YS, Huang HS, Ma MC. 2012. Erythropoietin alleviates post-ischemic injury of rat
hearts by attenuating nitrosative stress. *Life Sciences* **90**:776-784.
doi:10.1016/j.lfs.2012.04.012

Mariani J, Ou R, Bailey M, Rowland M, Nagley P, Rosenfeldt F, Pepe S. 2000. Tolerance to
ischemia and hypoxia is reduced in aged human myocardium. *J Thorac Cardiovasc Surg*
120:660-667. doi:10.1067/mtc.2000.106528

Meschiari CA, Ero OK, Pan H, Finkel T, Lindsey ML. 2017. The impact of aging on cardiac
extracellular matrix. *Geroscience* **39**:7-18. doi:10.1007/s11357-017-9959-9

Ming X-F, Yang Z. 2013. Functions and Mechanisms of Arginase in Age-Associated Cardiovascular
Diseases. *Curr Transl Geriatr and Exp Gerontol Rep* doi:. DOI 10.1007/s13670-013-0060-
7.

Ming XF, Barandier C, Viswambharan H, Kwak BR, Mach F, Mazzolai L, Hayoz D, Ruffieux J,

Rusconi S, Montani JP, Yang Z. 2004. Thrombin stimulates human endothelial arginase
enzymatic activity via RhoA/ROCK pathway: implications for atherosclerotic endothelial
dysfunction. *Circulation* **110**:3708-3714. doi:10.1161/01.CIR.0000142867.26182.32
Ming XF, Rajapakse AG, Carvas JM, Ruffieux J, Yang Z. 2009. Inhibition of S6K1 accounts
partially for the anti-inflammatory effects of the arginase inhibitor L-norvaline. *BMC*
Cardiovasc Disord **9**:12. doi:10.1186/1471-2261-9-12
Ming XF, Rajapakse AG, Yepuri G, Xiong Y, Carvas JM, Ruffieux J, Scerri I, Wu Z, Popp K, Li J,
Sartori C, Scherrer U, Kwak BR, Montani JP, Yang Z. 2012. Arginase II Promotes
Macrophage Inflammatory Responses Through Mitochondrial Reactive Oxygen Species,
Contributing to Insulin Resistance and Atherogenesis. *J Am Heart Assoc* **1**:e000992.
doi:10.1161/JAHA.112.000992
Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, Pinto AR, Klapproth K, Henri
S, Malissen B, Rodewald HR, Rosenthal NA, Bajenoff M, Prinz M, Jung S, Sieweke MH.
2014. Progressive replacement of embryo-derived cardiac macrophages with age. *J Exp Med*
211:2151-2158. doi:10.1084/jem.20140639
Moretto J, Girard C, Demougeot C. 2019. The role of arginase in aging: A systematic review. *Exp*
Gerontol **116**:54-73. doi:10.1016/j.exger.2018.12.011
Moturi S, Ghosh-Choudhary SK, Finkel T. 2022. Cardiovascular disease and the biology of aging. *J*
Mol Cell Cardiol **167**:109-117. doi:10.1016/j.yjmcc.2022.04.005
Murdoch CE, Chaubey S, Zeng L, Yu B, Ivetic A, Walker SJ, Vanhoutte D, Heymans S, Grieve DJ,
Cave AC, Brewer AC, Zhang M, Shah AM. 2014. Endothelial NADPH oxidase-2 promotes
interstitial cardiac fibrosis and diastolic dysfunction through proinflammatory effects and
endothelial-mesenchymal transition. *J Am Coll Cardiol* **63**:2734-2741.
doi:10.1016/j.jacc.2014.02.572
Ohtsu H, Frank GD, Utsunomiya H, Eguchi S. 2005. Redox-dependent protein kinase regulation by
angiotensin II: mechanistic insights and its pathophysiology. *Antioxid Redox Signal* **7**:1315-
1326. doi:10.1089/ars.2005.7.1315
Percie du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC,
Dirnagl U, Emerson M, Garner P, Holgate ST, Howells DW, Hurst V, Karp NA, Lazic SE,
Lidster K, MacCallum CJ, Macleod M, Pearl EJ, ... 2020. Reporting animal research:
Explanation and elaboration for the ARRIVE guidelines 2.0. *PLoS Biology* **18**:e3000411.
doi:10.1371/journal.pbio.3000411
Redfors B, Shao Y, Omerovic E. 2012. Myocardial infarct size and area at risk assessment in mice.
Exp Clin Cardiol **17**:268-272.
Ren Z, Potenza DM, Ma Y, Ajalbert G, Hoogewijs D, Ming XF, Yang Z. 2022. Role of Arginase-II
in Podocyte Injury under Hypoxic Conditions. *Biomolecules* **12**. doi:10.3390/biom12091213
Ruiz-Meana M, Bou-Teen D, Ferdinandy P, Gyongyosi M, Pesce M, Perrino C, Schulz R, Sluijter
JPG, Tocchetti CG, Thum T, Madonna R. 2020. Cardiomyocyte ageing and
cardioprotection: consensus document from the ESC working groups cell biology of the
heart and myocardial function. *Cardiovasc Res* **116**:1835-1849. doi:10.1093/cvr/cvaa132
Shi O, Morris SM, Jr., Zoghbi H, Porter CW, O'Brien WE. 2001. Generation of a mouse model for
arginase II deficiency by targeted disruption of the arginase II gene. *Mol. Cell Biol.* **21**:811-
813.
Sin YY, Baron G, Schulze A, Funk CD. 2015. Arginase-1 deficiency. *J Mol Med (Berl)* **93**:1287-
1296. doi:10.1007/s00109-015-1354-3
Steppan J, Ryoo S, Schuleri KH, Gregg C, Hasan RK, White AR, Bugaj LJ, Khan M, Santhanam L,
Nyhan D, Shoukas AA, Hare JM, Berkowitz DE. 2006. Arginase modulates myocardial
contractility by a nitric oxide synthase 1-dependent mechanism. *Proceedings of the National*
Academy of Sciences, USA **103**:4759-4764. doi:10.1073/pnas.0506589103
Swirski FK, Nahrendorf M, Etzrodt M, Wildgruber M, Cortez-Retamozo V, Panizzi P, Figueiredo
JL, Kohler RH, Chudnovskiy A, Waterman P, Aikawa E, Mempel TR, Libby P, Weissleder
R, Pittet MJ. 2009. Identification of splenic reservoir monocytes and their deployment to

inflammatory sites. *Science* **325**:612-616. doi:10.1126/science.1175202

Vakka A, Warren JS, Drosatos K. 2023. Cardiovascular aging: from cellular and molecular changes
to therapeutic interventions. *J Cardiovasc Aging* **3**. doi:10.20517/jca.2023.09

Wang C, Yu X, Cao Q, Wang Y, Zheng G, Tan TK, Zhao H, Zhao Y, Wang Y, Harris D. 2013.
Characterization of murine macrophages from bone marrow, spleen and peritoneum. *BMC*
Immunology **14**:6. doi:10.1186/1471-2172-14-6

Wang Y, Sano S, Yura Y, Ke Z, Sano M, Oshima K, Ogawa H, Horitani K, Min KD, Miura-Yura E,
Kour A, Evans MA, Zuriaga MA, Hirschi KK, Fuster JJ, Pietras EM, Walsh K. 2020. Tet2-
mediated clonal hematopoiesis in nonconditioned mice accelerates age-associated cardiac
dysfunction. *JCI Insight* **5**. doi:10.1172/jci.insight.135204

Wolff CA, Gutierrez-Monreal MA, Meng L, Zhang X, Douma LG, Costello HM, Douglas CM,
Ebrahimi E, Pham A, Oliveira AC, Fu C, Nguyen A, Alava BR, Hesketh SJ, Morris AR,
Endale MM, Crislip GR, Cheng KY, Schroder EA, Delisle BP, ... 2023. Defining the age-
dependent and tissue-specific circadian transcriptome in male mice. *Cell Rep* **42**:111982.
doi:10.1016/j.celrep.2022.111982

Xiong Y, Yepuri G, Montani JP, Ming XF, Yang Z. 2017. Arginase-II Deficiency Extends Lifespan
in Mice. *Front Physiol* **8**:682. doi:10.3389/fphys.2017.00682

Xiong Y, Yepuri G, Necetin S, Montani JP, Ming XF, Yang Z. 2017. Arginase-II Promotes Tumor
Necrosis Factor-alpha Release From Pancreatic Acinar Cells Causing beta-Cell Apoptosis in
Aging. *Diabetes* **66**:1636-1649. doi:10.2337/db16-1190

Xu Y, Kovacic JC. 2023. Endothelial to Mesenchymal Transition in Health and Disease. *Annual*
Review of Physiology **85**:245-267. doi:10.1146/annurev-physiol-032222-080806

Yepuri G, Velagapudi S, Xiong YY, Rajapakse AG, Montani JP, Ming XF, Yang ZH. 2012. Positive
crosstalk between arginase-II and S6K1 in vascular endothelial inflammation and aging.
Aging Cell **11**:1005-1016. doi:10.1111/Acel.12001

Zeisberg EM, Tarnavski O, Zeisberg M, Dorfman AL, McMullen JR, Gustafsson E, Chandraker A,
Yuan X, Pu WT, Roberts AB, Neilson EG, Sayegh MH, Izumo S, Kalluri R. 2007.
Endothelial-to-mesenchymal transition contributes to cardiac fibrosis. *Nat Med* **13**:952-961.
doi:10.1038/nm1613

Zhang L, He J, Wang J, Liu J, Chen Z, Deng B, Wei L, Wu H, Liang B, Li H, Huang Y, Lu L, Yang
Z, Xian S, Wang L. 2021. Knockout RAGE alleviates cardiac fibrosis through repressing
endothelial-to-mesenchymal transition (EndMT) mediated by autophagy. *Cell Death Dis*
12:470. doi:10.1038/s41419-021-03750-4

Zhu C, Potenza DM, Yang Y, Ajalbert G, Mertz KD, von Gunten S, Ming XF, Yang Z. 2023. Role of
pulmonary epithelial arginase-II in activation of fibroblasts and lung inflammaging. *Aging*
Cell:e13790. doi:10.1111/acel.13790

1003

Fig. 1

Fig. 2

A

SNAIL — 48

Vimentin — 48

Tubulin — 48

CD31 — 110

Vinculin — 110

wt　*arg-ii^{-/-}*　*wt*　*arg-ii^{-/-}*

Young　　　　Old

B

SNAIL / tubulin (fold change)

wt　*arg-ii^{-/-}*　*wt*　*arg-ii^{-/-}*

Young　　　　Old

C

Vimentin / tubulin (fold change)

wt　*arg-ii^{-/-}*　*wt*　*arg-ii^{-/-}*

Young　　　　Old

D

CD31 / vinculin (fold change)

wt　*arg-ii^{-/-}*　*wt*　*arg-ii^{-/-}*

Young　　　　Old

E

CD31/Vimentin/DAPI

wt

arg-ii^{-/-}

20 μm

Young　　　　Old

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Macrophage

↑ Arg-II

↑ IL-1β

Cardiomyocyte

Fibroblasts

↑ Arg-II

↑ mtROS

Endothelial cells

↑ Arg-II

Apoptosis

Matrix production

EndMT

Cardiac Aging

Fig. 10