**Spatially targeted inhibitory rhythms differentially affect neuronal integration**

Drew B. Headley[1,*], Benjamin Latimer[2,*], Adin Aberbach[2], and Satish S. Nair[2]

**AFFILIATIONS:**
[1] Center for Molecular and Behavioral Neuroscience, Rutgers University – Newark, Newark, NJ, 07102, USA
[2] Electrical Engineering & Computer Science, University of Missouri, Columbia, MO, 65211, USA
[*] Contributed equally to this work

**CORRESPONDENCE:**
Satish S. Nair
Department of Electrical Engineering & Computer Science
University of Missouri, Columbia MO 65211
Phone: (573) 882-2964
E-Mail: nairs@missouri.edu
Website: https://nairs.mufaculty.umsystem.edu/

**ABSTRACT**
Pyramidal neurons form dense recurrently connected networks with multiple types of inhibitory interneurons. A major differentiator between interneuron subtypes is whether they synapse onto perisomatic or dendritic regions. They can also engender local inhibitory rhythms, beta (12-35 Hz) and gamma (40-80 Hz). The interaction between the rhythmicity of inhibition and its spatial targeting on the neuron may determine how it regulates neuronal integration. Thus, we sought to understand how rhythmic perisomatic and distal dendritic inhibition impacted integration in a layer 5 pyramidal neuron model with realistic dendrites supporting $Na^+$, NMDA, and $Ca^{2+}$ spikes. We found that inhibition regulated the coupling between dendritic spikes and action potentials in a location and rhythm-dependent manner. Perisomatic inhibition principally regulated action potential generation, while distal dendritic inhibition regulated the incidence of dendritic spikes and their temporal coupling with action potentials. Perisomatic inhibition was most effective when provided at gamma frequencies, while distal dendritic inhibition functioned best at beta. Moreover, beta modulated responsiveness to distal inputs in a phase-dependent manner, while gamma did so for proximal inputs. These results may provide a functional interpretation for the reported association of soma-targeting parvalbumin positive interneurons with gamma, and dendrite-targeting somatostatin interneurons with beta.

**INTRODUCTION**

Cortical circuits are composed of connected networks of excitatory pyramidal neurons. To maintain a balanced level of excitability, they form dense reciprocal connections with local inhibitory interneurons. The two largest subclasses of interneurons in the cortex are parvalbumin positive (PV) and somatostatin positive (SOM) cells [1]. A major distinction between PV and SOM interneurons is that they synapse onto different parts of the pyramidal neuron. PV interneurons tend to contact the soma and proximal dendrites (perisomatic), while SOMs target distal dendrites [2-5]. Numerous studies have identified differences in how they regulate the responsiveness of local excitatory principal neurons [6-8]. These differences could arise from their connectivity, intrinsic properties, or where they synapse onto pyramidal neurons. Indeed, the location of an inhibitory synapse qualitatively changes its effect on synaptic integration [5, 9-11].

Synaptic integration in pyramidal neurons arises from the interplay between passive and active ion channels and dendritic morphology [9, 12-14]. Dendrites produce regenerative spiking events, namely $Na^+$, NMDA, and $Ca^{2+}$ spikes. Their initiation depends upon the spatiotemporal coordination of excitatory synapses and interaction with the depolarization of dendritic branches. Incorporating just NMDA spikes into a model neuron radically increases the complexity of its dendritic integration [15]. $Na^+$ and NMDA spikes can drive somatic spiking in an *in vivo*-like model neuron [13]. A $Ca^{2+}$ spike in the apical trunk converts a single somatic spike to a burst [9, 16], and may allow pyramidal neurons to act as multi-stage integrators [17, 18]. This complexity is only increased by including inhibition. The emission of bursts of action potentials is controlled by the timing and dendritic location of inhibition [9].

The rhythmicity of inhibition at distal dendrites also affects somatic integration [19]. Reciprocal interactions between excitatory principal neurons and inhibitory interneurons gives rise to rhythmic activities (pyramidal-interneuron network gamma or PING mechanism). To initiate the rhythm, excitatory principal neurons activate inhibitory interneurons that deliver feedback inhibition to principal neurons. This transiently suppresses principal neuron firing. As the inhibition wanes, principal neurons resume their activity and reengage the inhibitory population, starting a new oscillatory cycle. Some evidence suggests that different interneuron subtypes pace rhythms at different frequencies, with gamma oscillations depending on PV [20-22], and beta oscillations on SOM [21]. On the other hand, recent work has also found that beta/low-gamma rhythms in V1 engage both PV and SOM in their generation [23, 24].

Numerous cognitive processes are associated with these inhibitory rhythms in the cortex. Gamma oscillations are especially pronounced during stimulus presentation or behavioral initiation [25-27]. Beta rhythms occur during preparatory states or working memory [28, 29]. But fundamentally, these rhythms probably reflect different modes of local cortical activity and interregional communication [30-32]. For instance, gamma oscillations are associated with feedforward transmission of information in cortical circuits and beta oscillations may mediate feedback [30, 33]; but see [34].

Given all this, it is important to better understand how the various integrative events in the dendritic tree are affected by the location and frequency of inhibition. Perhaps some rhythmic timescales are more effective in the dendrites, and others at the soma. Investigating this is beyond the reach of conventional experimental techniques. Moreover, computational models exploring how beta and gamma influence information processing have mostly relied upon simplified model neurons (e.g., [35, 36]). Thus, using a morphologically and biophysically detailed layer 5 pyramidal neuron model with $Na^+$, NMDA, and $Ca^{2+}$ spikes, we investigated how perisomatic and distal dendritic inhibition impacted dendritic integration, and the modulation of these events by beta and gamma rhythmicity.

**RESULTS**

**Construction of a model cortical layer 5 pyramidal neuron**

To study the effect of inhibitory rhythms on synaptic integration and somatic spiking, we adapted a previously published morphologically and biophysically detailed model of a cortical layer 5 pyramidal neuron (**Fig. 1A**; See methods for details). Modifications to the model were done in accordance with the published literature. In brief, this model featured a multicompartmental dendritic tree that produced dendritic $Na^+$, NMDA, and $Ca^{2+}$ spikes, along with somatic action potentials that could backpropagate (**Fig. 1B**). We distributed conductance-based synapses across the dendritic and somatic compartments with an average density of 2.16 excitatory and 0.22 inhibitory contacts per µm. Presynaptic drivers of excitatory synapses were drawn from a pool of 5,200 point process sources that emulated correlated afferent drive with 2-8 synaptic contacts from the same presynaptic neuron. Inhibitory synapses were divided into two populations, those targeting the soma and proximal 100 µm of the dendrites (referred to as perisomatic), and those synapsing outside that area (referred to as distal). To capture excitatory/inhibitory (E/I) balance, a hallmark of cortical activity, the rate of inhibitory synaptic drive was a rescaled version of the rate of excitatory drive, lagged by 4 ms to emulate feedforward inhibition. Naturalistic presynaptic drive elicited a median firing rate of 5.3 Hz, in agreement with *in vivo* rates in cortex (**Fig. 1A** inset, [37]).

Dendrites were endowed with the following voltage-dependent conductances: a fast-inactivating $Na^+$ current ($I_{NaT}$), muscarinic $K^+$ current ($I_m$), fast non-inactivating $K^+$ current ($I_{Kv3.1}$), high voltage-activated $Ca^{2+}$ current ($I_{Ca\_HVA}$), low voltage-activated $Ca^{2+}$ current ($I_{Ca\_LVA}$), and $Ca^{2+}$ activated $K^+$ current ($I_{SK}$). As a result, the basal and apical dendrites could generate $Na^+$ and NMDA spikes (**Fig. 1B**; [13]). Dendritic $Na^+$ spikes were regenerative events lasting less than 1 ms that were not preceded by somatic action potentials (**Fig. 1C1**; [38]). A $Na^+$ spike was detected when the dendritic $Na^+$ channel conductance (gNa) was higher than 0.3 $mS/cm^2$, except when this threshold was reached within 5 ms after a somatic action potential, to distinguish from backpropagating action potentials.

NMDA spikes occur when adjacent NMDA-bearing synapses were synergistically recruited by a combination of glutamatergic activation and local depolarization (**Fig. 1C2**; [18, 39]). They typically lasted between 20 and 80 ms. They were detected when a compartment's membrane voltage exceeded -40 mV for at least 26 ms and NMDA current exceeded -100 pA ms of charge.

$Ca^{2+}$ spikes are depolarizations generated at the nexus of the apical trunk upon activation of voltage-gated $Ca^{2+}$ channels (**Fig. 1C3**; [40, 41]). They lasted between 20 and 50 ms. To detect them, the membrane potential must exceed -40 mV for at least 26 ms, and the combined $Ca^{2+}$ currents (LVA and HVA) had to be 1.3 times higher than when the voltage criterion was reached ($t_v$). The $Ca^{2+}$ spike ended when this value fell to 1.15 times its value at $t_v$.

Altogether, under conditions that mirror *in vivo* afferent drive, our model reproduces the dendritic spikes of a layer 5 pyramidal neuron.

**Relationship between dendritic and somatic spikes**

Dendritic spikes induced by synaptic activity are the principal drivers of somatic action potentials. Prior experimental and modeling work has found this to be the case for layer 2/3 and layer 5 pyramidal neurons [13, 18, 42-44]. Layer 5 pyramidal neurons have a substantially longer apical trunk, which increases the electrotonic distance of their apical tuft from the soma (**Fig. 2A**), and diminishes the ability of tuft synapses to elicit action potentials. Voltage-gated $Ca^{2+}$ channels at the apical nexus compensate for this by producing a robust $Ca^{2+}$ spike that drives a burst of action potentials at the soma [17, 18]. Underscoring that $Ca^{2+}$ spikes compensate for morphology, pyramidal neurons with shorter apical dendrites have weaker $Ca^{2+}$ spikes that only elicit a single spike [45-47]. We thus assessed how dendritic spikes at different electronic distances from the

soma were related to action potential generation.
Dendritic compartments differed in their degree of passive electrical coupling to the soma
(i.e., electrotonic distance; **Fig. 2A**). We measured this by injecting a 20 Hz sinusoidal current in
each dendritic compartment, and then calculating the ratio of the membrane voltage response at
the soma over that at the dendrite. The apical trunk exhibited a relatively small attenuation ratio
of ~10%. Progressing distally into the apical tuft, attenuation reached 0.1%. Basal dendrites
showed greater attenuation than the apical trunk due to their smaller diameter. However, with an
attenuation ratio reaching ~1%, their distal tips were still electrotonically closer than the apical
tuft. Thus, large, and long-lasting changes in membrane potential, like those produced by dendritic
spikes, would be required to have any effect on the soma.
We examined this by measuring the spike-triggered average between somatic action
potentials and the presence of dendritic spikes across the dendritic tree. To simplify the complex
geometry of our model neuron, dendritic compartments were grouped into deciles by their
electrotonic distance, and whether they were on apical or basal branches. For each time lag from
the somatic action potential, we measured the percent change in dendritic spike incidence from
the mean rate across the entire simulation.
Dendritic $Na^+$ spikes increased 2-3 ms prior to somatic action potentials in both basal and
apical dendrites (**Fig. 2B**). This relationship was strongest for the compartments nearest the
soma, with the rate of $Na^+$ spikes in the apical trunk increasing 300% over baseline prior to
somatic action potentials, and 100% in basal compartments. This relationship fell off as the
dendritic spikes moved farther away from the soma, indicating that $Na^+$ spikes in distal branches
had little direct influence on somatic spiking.
The incidence of NMDA spikes increased ~25 ms prior to somatic action potentials, much
earlier than seen with dendritic $Na^+$ spikes (**Fig. 2C**). But, like dendritic $Na^+$ spikes, NMDA spikes
in the apical branch had a stronger coupling with somatic spiking than those in basal branches,
and this effect dropped with distance from the soma. NMDA spikes in the apical branch persisted
after an action potential, while those in basal dendrites did not, potentially because of the action
potential after-hyperpolarization.
$Ca^{2+}$ spikes originate at the nexus, when the apical trunk first branches into the apical tuft.
This region is electrotonically close to the entire apical trunk, facilitating the propagation of $Ca^{2+}$
spikes (**Fig. 3A**). In our model, $Ca^{2+}$ spike occurrence increased within 20 ms of somatic action
potentials (**Fig. 3B**). Furthermore, we found that NMDA spikes in the apical dendrites tended to
precede $Ca^{2+}$ spikes (**Fig. 3C**). Our ionic current-based detection criteria distinguished between
these phenomena despite their similar membrane voltage profiles. Since NMDA spikes in the
apical tuft normally have a weak relationship to somatic spiking (**Fig. 2C**), they may elicit somatic
spiking indirectly by driving $Ca^{2+}$ spikes. Put another way, the apical nexus may serve as a
thresholded nonlinearity for NMDA spikes in the apical tuft to drive action potentials [18]. To test
this, we measured how a $Ca^{2+}$ spike changed the spike-triggered average between apical tuft
NMDA spikes and action potentials (**Fig. 3D, top**). This revealed that action potentials preceded
by a $Ca^{2+}$ spike (by up to 20 ms) had increased coupling with apical NMDA spikes. No such
change was seen in basal dendrites (**Fig. 3D, bottom**).
181
**Effect of perisomatic and distal dendritic inhibition on controlling response gain**
Subtypes of inhibitory interneurons synapse on distinct dendritic zones. PV interneurons mainly
synapse perisomatically, while SOM interneurons target distal dendrites [2-5]. These differences
may affect their modulation of synaptic integration [48], either by shifting the threshold for evoking
action potentials (a subtractive effect) or altering the slope of the relationship between excitation
and firing rate (a divisive effect). There is some disagreement about the degree to which PV and
SOM interneurons produce either of these effects [6-8], depending on the source of excitation
and local circuitry [49]. So, before we applied beta and gamma rhythmic inhibition in our model,
we studied the effect of tonically activating PV- and SOM-like inhibitory synapses.

We doubled the rate of the inhibitory presynaptic drive onto either the PV-targeted
perisomatic compartments (somatic and dendritic compartments within 100 µm), or the SOM-
targeted distal (>100 µm) dendritic branches. Both decreased the firing rate of the pyramidal cell
from 5.5 Hz to less than 1 Hz (**Fig. 4A**; control: 5.5±0.85 Hz; distal: 0.20±0.15 Hz; perisomatic:
0.70±0.31 Hz; mean±SD). These comparable changes may reflect either subtractive or divisive
effects and could derive from different mechanisms.
To isolate these factors, we first assessed how doubling inhibition affected action potential
initiation at the soma. A series of current pulses were injected into the soma to measure the
relationship between firing rate and injected current (f-I curve), which captures the gain function
of the neuron (**Fig. 4B**). This revealed that both perisomatic and dendritic inhibition shifted the
current threshold for action potential initiation. Such an effect is subtractive. In addition,
perisomatic inhibition decreased the slope of the f-I relationship compared with the control, which
is consistent with a divisive effect.
Although perisomatic inhibition produced the strongest subtractive effect, distal dendritic
inhibition reduced firing rate the most (**Fig. 4A**). How can we reconcile these discordant findings?
One possibility is that distal inhibition substantially reduces $Na^+$, NMDA, or $Ca^{2+}$ spikes. To
determine this, we examined the overall rate of dendritic spike events. Perisomatic inhibition did
not affect dendritic events compared to the control condition (**Fig. 4C-E**). By contrast, dendritic
inhibition decreased NMDA and $Ca^{2+}$ spikes (**Fig. 4D,E**). $Na^+$ spikes were relatively unaffected
(**Fig. 4C**). This lack of effect may arise from a shortening of the inactivation for voltage-gated $Na^+$
channels balancing out the loss of excitatory drive.
212
**Distinct excitation/inhibition balance effects of perisomatic and distal dendritic inhibition**
The previous analysis found distinct effects on neuronal gain and dendritic spiking from tonic
changes in perisomatic and distal dendritic inhibition. But *in vivo* inhibition is dynamic and time
varying. PV and SOM interneurons form dense interconnections with local pyramidal neurons,
supplying time-lagged feedback inhibition. This helps maintain the balance of excitation and
inhibition (E/I) in the network by ensuring that an overall increase in excitatory activity is rapidly
counterbalanced by proportionate inhibition.
To emulate situations where E/I balance is important, we increased the dynamic variation
in excitatory drive (see Methods). As with the previous simulations, the rate of inhibitory synaptic
drive was a lagged and rescaled version of the overall excitation rate. Normally that lag is 4 ms,
in line with experimental estimates [50]. To probe whether perisomatic or distal dendritic inhibition
have distinct effects on E/I balance, we independently varied their lags (**Fig. 4F-I**). One was kept
at the nominal 4 ms lag, while the other was extended. Increasing either of their lags by 500 ms
produced obvious differences in the emission of dendritic spikes and their coordination with action
potentials (**Fig. 4F**). Normally dendritic spikes are evenly distributed in time and drive somatic
action potentials. Lagging distal dendritic inhibition clustered dendritic spikes together in time,
during which action potentials were emitted. Lagging perisomatic inhibition did not affect the
spacing of dendritic spikes, but decreased their coupling with somatic spiking.
We systematically characterized these lag effects for the following spiking events
modulated by tonic changes in inhibition: action potentials, $Ca^{2+}$, and NMDA spikes (**Fig. 4G**).
Increasing the lag of perisomatic inhibition lowered action potential firing, while for distal dendritic
inhibition, the firing rate decreased out to a lag of 125 ms, and then returned to normal at 500 ms.
To better understand these effects, we calculated the cross-correlation between dendritic spikes
and action potentials. Increasing the lag decreased the coordination between $Ca^{2+}$ and somatic
spikes (**Fig. 4H**). For distal dendritic inhibition, this decrease was accompanied by a broadening
of their temporal relationship, while a sharp temporal relationship was maintained in the
perisomatic case. A similar pattern was observed with NMDA spikes on apical branches (**Fig. 4I**).
Basal branches, on the other hand, were relatively unaffected.

In summary, distal and perisomatic inhibition modulate the coupling between synaptic
drive and spiking through distinct processes. And while for both cases, the normal 4 ms E/I lag
produced maximal conversion of dendritic spiking events into action potentials, extending this lag
exerted distinct effects.
245
**Effect of beta and gamma rhythmic inhibition on neuronal integration**
The mechanisms modulating neuronal responsiveness during tonic inhibition of somatic and
dendritic compartments may extend to rhythmic inhibition. Thus, we emulated beta and gamma
rhythmic input (**Fig. 5A,F**). Depths of modulation were set to similarly entrain action potentials
(**Fig. 5B,G**), and were comparable to spontaneous and optogenetically-induced gamma and beta
bursts seen *in vivo* [51-54]. Beta rhythmic inhibition was modeled as a 16 Hz sinusoidally
modulated rate (20% depth) of the Poisson processes driving inhibitory synapses. Gamma
rhythmic inhibition was a 64 Hz sinusoidal modulation (40%) of inhibitory synapses. For both
cases, excitatory synaptic drive was a stable Poisson process.
Initially, we delivered beta rhythmic inhibition to the distal dendritic compartments, and
gamma to the perisomatic. Even though both rhythms produced similar depths of modulation of
somatic action potentials, the underlying causes were distinct. The phase of beta modulated the
occurrence of $Ca^{2+}$, NMDA, and $Na^+$ spikes, with each showing an ~75% depth of modulation with
respect to their mean level (**Fig. 5C-E**). These changes were seen across the entire dendritic tree,
spanning sites electrotonically close to and far from the soma. In addition, the impact on $Na^+$
spikes was unexpected (**Fig. 5E**), since delivery of the same inhibition tonically had little effect.
By contrast, gamma had virtually no effect on dendritic spikes (**Fig. 5F-H**). Its strongest impact
was on $Na^+$ spikes in basal dendritic compartments that were electronically close to the soma.
To uncover the modulation of action potentials by gamma, we turned our attention to
somatic action potential initiation (**Fig. 6**). For these analyses we divided the inhibitory rhythm into
two phases. During the *peak* phase, inhibition was greater than its mean rate, while during the
*trough* phase inhibition was lower (see **Fig. 5A,F**). We found that the somatic action potential
voltage threshold shifted lower during the 'trough' phase of gamma, when inhibition was at its
weakest (**Fig. 6A1**), and without any change in the mean membrane voltage (**Fig. 6A2**). This is
consistent with gamma phase modulating the shunting of voltage-gated $Na^+$ channel currents,
which occurs when GABAergic synapses co-locate with the channels mediating action potential
initiation [55]. We also observed changes in action potential initiation to beta rhythmic inhibition,
but through a different mechanism. During the 'peak' phase of beta, when inhibition was maximal,
the threshold for evoking an action potential increased, which may reflect an 'off-path' shunting of
excitatory current away from the soma and towards the dendrites (**Fig. 6B1;** [56]). Additionally,
there was a decrease in membrane voltage during the peak phase, which may correspond to
decreased excitation arising from the suppression of dendritic spikes (**Fig. 6B2**).
Accompanying these effects were changes in the rate of dendritic spikes compared with
the Poisson inhibition case. Beta increased the rate of $Na^+$ (+16.6%) and $Ca^{2+}$ spikes (+15.1%)
but decreased the rate of NMDA spikes (-10.7%). Gamma caused no change in NMDA (+0.2%)
and a weak increase in $Na^+$ spikes (+6.4%), but a robust gain in $Ca^{2+}$ spikes (+37.9%).
We next switched the locations on the pyramidal neuron targeted by the beta and gamma
rhythms to disassociate the frequency of inhibition from its location. Beta rhythmic inhibition was
delivered perisomatically and gamma rhythmic inhibition to distal dendrites. While phase
modulation of firing rate was maintained with both rhythms, the overall level of spiking was
dramatically reduced (**Fig. 5S1A,E**). Neither rhythm modulated $Ca^{2+}$ or NMDA spikes (**Fig.
5S1B,C,F,G**). It is likely that the slow time scale of $Ca^{2+}$ and NMDA spikes, ~50 ms, is not optimal
for the fast periodicity of the gamma rhythm, which cycles every ~15 ms. In agreement with this,
$Na^+$ spikes, which last less than 1 ms, did show modulation by gamma rhythms delivered to the
distal dendrites (**Fig. 5S1D,H**).

Swapping the location of beta and gamma synapses altered their effects on somatic excitability. Gamma rhythmic inhibition on the dendrites had minimal or no impact on action potential threshold, but did shift the somatic membrane potential more negative (**Fig. 6S1A**). This hyperpolarization was not dependent on gamma phase, and likely reflected an overall decrease in the rate of dendritic spikes that could supply excitatory drive to the soma (NMDA: -26.1%, $Na^+$: -12.2%, and $Ca^{2+}$: -39.9% compared with the Poisson inhibition case). By contrast, delivering beta rhythmic inhibition to the soma raised the action potential threshold and hyperpolarized the membrane potential during the peak phase (**Fig. 6S1B**). It also reduced the incidence of dendritic spikes (NMDA: -17.1%, $Na^+$: -11.5%, and $Ca^{2+}$: -7.6%).

Putting all this together, the effectiveness of rhythmic inhibition depends on where it impinges upon the neuron. Beta rhythms targeting the distal dendrites modulate the incidence of dendritic spikes in a phase-dependent manner, while gamma rhythms delivered perisomatically phase modulate somatic excitability. Swapping the locations of these rhythms diminishes these effects and lowers overall excitability.

**Frequency specific effects of rhythmic inhibition on neuronal integration**

Having demonstrated that the location of beta and gamma rhythmic inhibition impacts their effectiveness, we next determined its frequency specificity. To do this, we varied the frequency of rhythmic inhibition between 0.5 and 80 Hz on either the perisomatic or distal dendritic neuronal compartments. Starting with distal dendrites, increasing inhibition frequency above 20 Hz diminished its entrainment of NMDA, $Na^+$, and $Ca^{2+}$ spike onsets (**Fig. 7A**). The falloff in entrainment was most pronounced above 20 Hz. Curiously, $Na^+$ spikes exhibited a preferential entrainment at 20 Hz. In general, entrainment was strongest in the apical dendrites.

The previous analysis considered the entrainment of dendritic spike *onsets*, but NMDA and $Ca^{2+}$ spikes also exhibit *offsets* that could also be modulated by rhythmic inhibition. Indeed, examination of voltage traces in the dendrites during beta rhythmic inhibition revealed that NMDA and $Ca^{2+}$ spike onsets tended to occur during the trough, while offsets happened during the peaks (**Fig. 7B**). To quantify this, we plotted the percent change in the probability of dendritic spike onsets and offsets with respect to both the phase and frequency of the inhibitory rhythm (**Fig. 7C**). For frequencies less than 5 Hz, there was minimal phase separation between the onsets and offsets of $Ca^{2+}$ or NMDA spikes, both events occurred near the rhythm trough, when inhibition was at its weakest. As frequency increased up to 20 Hz, the preferred phase of dendritic spike offsets migrated toward the peak phase, where inhibition is strongest. This phase separation effect was strongest for $Ca^{2+}$ spikes. Thus, beta band frequencies exhibit unique coordination with dendritic spikes: they are the fastest rhythm capable of entraining them and align with their initiation and cessation in a phase-dependent manner.

Turning to perisomatic inhibition, we again varied the frequency of the inhibitory rhythm. Lower frequency inhibition produced phase-dependent shifts in the mean membrane potential (**Fig. 8A**). The trough of inhibition depolarized the soma, while the peak of inhibition had the opposite effect. As the frequency increased, this phase dependent difference went away, vanishing above 50 Hz. By contrast, as frequency increased the bias in momentary changes in the membrane potential diverged between peaks and troughs (**Fig. 8B**). During the peak phase membrane voltage fluctuations were biased negative, while during the trough they were biased positive. This effect increased with frequency, peaking at 50 Hz, and then declining modestly. Together, these effects make gamma frequencies unique in keeping the mean membrane potential equivalent between phases but biasing its fluctuations towards depolarizing or hyperpolarizing with phase. During the trough there was an excess of depolarizing membrane potential fluctuations. Since the rate of excitatory synaptic drive was independent of phase, this suggests that its responsiveness to excitatory inputs increased in a phase-dependent manner with gamma.

**Modulation of dendritic spikes during oscillatory bursts**

*In vivo*, beta and gamma rhythms occur as bursts lasting less than a few hundred milliseconds. Since the previous analyses relied on tonically delivered rhythms, we verified that similar effects were observed with oscillatory bursts. Gamma and beta bursts were delivered to the same model with mean depth of modulation like the tonic case (**Fig. 9A,F**; see Methods for details).

Under the burst regime, both rhythms mirrored their behavioral effects. Gamma bursts entrained spiking, with entrainment strongest during the middle of the burst (**Fig. 9B**). As with the tonically imposed rhythm, there was none or minimal modulation of $Ca^{2+}$ (**Fig. 9C**), NMDA (**Fig. 9D**), and $Na^+$ spikes (**Fig. 9E**). Beta rhythms entrained somatic action potentials (**Fig. 9G**), $Ca^{2+}$ spikes (**Fig. 9H**), NMDA (**Fig. 9I**), and $Na^+$ spikes (**Fig. 9J**). These modulations were evident within the first few cycles of a burst, suggesting that they did not require a buildup or evolving entrainment of an underlying process.

**Effect of beta and gamma rhythms on responding to clustered synaptic drive**

The results so far suggest that beta and gamma rhythms modulate synaptic integration through different mechanisms depending on their phase and the location of the synapses on the dendritic tree. To examine this further, we added patches of concentrated excitatory synaptic inputs onto either the distal or proximal dendrites (**Fig. 10A**), with densities similar to functional clusters *in vivo* [57, 58]. Coactivated inputs were simulated by driving each synapse with a jittered (2 ms) Poisson process. We ran six separate simulations, either with the distal or proximal clusters engaged, and under conditions of Poisson, beta, or gamma rhythmic inhibition.

To capture how beta and gamma influence synaptic integration we measured the cross-correlation between presynaptic activations at the clustered input and somatic spiking. Separate cross-correlograms were calculated depending on whether the presynaptic spikes occurred during the peak or trough phase of the inhibitory rhythm. This necessarily introduced spurious periodicities into the cross-correlogram that were compensated (see Methods for details). Relative to the arhythmic Poisson inhibition case, beta rhythms enhanced the transmission of distal inputs when inhibition was low (trough phase) and suppressed them when inhibition was high (peak phase, **Fig. 10B, top left**). Proximal inputs were either unaffected or moderately suppressed during the trough and suppressed during the peak (**Fig. 10B, bottom left**). The opposite was the case for gamma. It barely affected or moderately suppressed distal inputs (**Fig. 10B top right**), while proximal inputs were enhanced during the trough and suppressed during the peak (**Fig. 10B bottom right**).

Summarizing these results (**Fig. 10C**), we found that somatic spiking driven by clustered proximal synapses was bidirectionally modulated by gamma rhythms and suppressed by beta. On the other hand, spiking driven by distal clusters was bidirectionally modulated by the beta rhythm and suppressed by gamma. Thus, both rhythms regulate the sensitivity of pyramidal neurons to afferents throughout the dendritic tree, but in a counterposed location-dependent manner.

**DISCUSSION**

Arising from multiple interneuron subtypes, inhibition sculpts pyramidal neuron activity by acting at different membrane regions and distinct rhythmic frequencies (**Fig. 11A**). Little was known about how these factors interact with the complexity of dendritic integration. To address this, we characterized the interaction between the location and rhythmicity of inhibition on integration in a morphologically and biophysically detailed layer 5 pyramidal neuron model with active dendrites. We found that distal dendritic inhibition modulated the occurrence of dendritic spikes, while perisomatic inhibition altered action potential generation. This translated into location-specific differences in the effectiveness of inhibitory rhythms. Beta rhythmic inhibition entrained dendritic spikes, focusing them into the phase when inhibition was at a minimum, but only when delivered to the distal dendrites (**Fig. 11B**). In contrast, gamma modulated the threshold for action potential initiation, but only when delivered perisomatically (**Fig. 11C**). The effects of these rhythms were frequency specific, with the timing of beta and gamma aligning preferentially with phase dependent effects on neuronal integration. As a likely result, beta oscillations bidirectionally controlled transmission in distal dendrites and suppressed those onto proximal dendrites, while gamma oscillations did the opposite.

This indicates that the spatial targeting and timing of inhibition go hand in hand. Their alignment is thus fundamental to their distinct effects on neuronal integration. And in turn, this affects function at multiple levels, from synaptic responsiveness to neural coding, and microcircuit operation.

**Neuronal responsiveness**

Oscillatory rhythms potentially affect synaptic integration through several mechanisms. Early theorizing held that oscillations synchronize populations of excitatory neurons such that their efferent synapses benefit from spatiotemporal summation in eliciting action potentials [59]. Later it was proposed that if neurons in two regions exhibited coherent variations in their somatic membrane potentials, bringing them closer to or farther from action potential threshold, they could preferentially exchange activities [60]. Modeling and *in vitro* studies have shown that the phase of a sinusoidal current injected at the soma modulates the timing of evoked action potentials [61] or their probability of occurrence [62]. Similar gamma phase dependence has been found *in vivo* [63].

Rhythmic inhibition may also periodically modulate the sensitivity of the neuron to synaptic input. Inhibitory synapses achieve this by lowering the membrane resistance i.e., shunting [64]. This makes it harder for an excitatory synaptic current to drive the membrane voltage toward the action potential threshold. Indeed, optogenetically stimulating PV interneurons at gamma frequencies modulates the responsiveness of cortical neurons to sensory inputs in a phase-dependent manner [65]. However, normally PV interneurons are driven by local principal cells, so the inhibition they impose reflects the aggregate activity in the local network (leading to E/I balance, [66]). Model networks wired in this manner emit spontaneous gamma bursts that produce phase-dependent modulation of the relationship between EPSP amplitude and spiking probability, with a strong positive relationship during the trough of gamma, and a weak one during the peak [36]. Consistent with these results, putative monosynaptically connected pairs of single units recorded *in vivo* show an increase in spike transmission probability during the trough of gamma [67].

To summarize the present possibilities, oscillations could modulate a neuron's responsiveness by (1) synchronizing the synaptic inputs impinging upon it, (2) modulate how close the membrane potential is to the action potential threshold, or (3) vary its sensitivity to synaptic inputs via shunting inhibition.

Our results fill out this picture in two ways. First, we add a fourth mechanism, which is (4) the modulation of dendritic spiking events. Second, we found that beta and gamma oscillations differentially engaged these mechanisms. Beta oscillations primarily caused 2 via 4, and affected

433   3. Gamma, on the other hand, operated primarily through 3. It should be noted that to isolate
434   these effects, our model had Poisson excitatory synaptic inputs, which precluded mechanism 1,
435   afferent synchrony. However, it is likely the degree of synchrony would impact both rhythms.
436   Supporting this, *in vivo* whole-cell recordings from neurons in mouse visual cortex exhibit brief
437   membrane depolarizations in synchrony with gamma [68].
438
**Relevance to coding**
Given that beta and gamma rhythms influence spiking via distinct mechanisms, they may also
differentially impact coding. During gamma bursts in V1, neurons show a stronger modulation of
their firing rate by the contrast of grating stimuli [69]. Gamma phase also modulates orientation
selectivity and correlated variability between neurons [70]. This enhancement is modulated by
attention [71]. Our results indicate that these effects reflect shifts in somatic sensitivity to
excitatory synaptic inputs.
Much less is known about the influence of beta oscillations on coding. In the visual cortex
of mice, activation of SOM interneurons increased with the visual stimulus size and homogeneity
([22], note that this study referred to the rhythm as low gamma). In a different study, using a
sequence working memory task in primates, the order of presented items corresponded to their
phase in the beta rhythm [72]. However, this does not indicate how coding varies with beta phase.
Considering our finding that beta synchronized dendritic spikes, it may operate by enabling the
summation of normally asynchronous afferents on distal dendrites, facilitating their ability to drive
somatic spiking.
One interpretation of rhythms arising from local inhibitory feedback is that they maintain
the balance between excitation and inhibition. This can be thought of as a normalization operation
that maintains activity within a set range. Normalization can be achieved either through a
subtractive effect that raises the threshold for initiating an action potential, or a multiplicative effect
that lowers the slope of the relationship between excitation and action potential firing rate. When
considered at the population level, these normalization effects impact coding in different ways.
Subtractive normalization increases sparsity by dropping out neurons whose excitation is below
the raised threshold. Multiplicative normalization, however, encourages dense codes by scaling
down firing rates and compressing the range of firing rates. This study found that while both
perisomatic and distal dendritic inhibition produced subtractive effects, only perisomatic had a
multiplicative effect. Tying this to beta and gamma, beta rhythms may encourage sparse
population codes while gamma allows for dense.
466
**Interaction with microcircuitry**
Pyramidal neurons are embedded in a cortical column where they are sparsely connected
amongst themselves and densely connected with local interneurons. The dense connectivity with
interneurons is crucial for the generation of local beta and gamma rhythms [21, 22]. How might
this circuitry interact with the differential regulation of dendritic integration by beta and gamma?
We found that beta rhythms preferentially modulated a pyramidal neuron's response to
inputs on its distal dendrites. The apical tuft in the superficial layers of cortex primarily receives
long-range inputs from regions higher up in the cortical hierarchy [73, 74]. This makes beta
rhythms ideally positioned to modulate these top-down/feedback signals. Since the rhythm likely
derives from local activation of SOM interneurons, it is worth considering what situations would
activate those and how they might relate to the functioning of apical dendrites. SOM interneurons
receive facilitating synapses from local pyramidal neurons [75, 76], which would make them
especially sensitive to bursts of action potentials arising from $Ca^{2+}$ spikes at the apical nexus [18].
Since $Ca^{2+}$ spikes are driven by synaptic activation in the apical tuft, it is likely that beta rhythms
regulate the generation of action potential bursts arising from long-range inputs.
Particularly in the visual cortex, SOM interneurons can generate a rhythm in the 25-30 Hz
range [22]. We found this to be at the upper end of the frequency range for dendritic inhibitory

rhythms to be effective in modulating NMDA and $Ca^{2+}$ spikes. If this rhythm solely recruited SOM
interneurons, its effectiveness would be marginal. Potentially compensating for this, recent work
has found that PV interneurons also participate in beta/low-gamma [23, 24] (but see [21, 22]). In
our model, on its own when beta rhythmic inhibition was delivered perisomatically we found that
it was less able to entrain spiking and had an overall hyperpolarizing effect. However, if delivered
in conjunction with the distal dendritic inhibition arising from SOM interneurons, this may
strengthen entrainment.
Turning to gamma, our results indicated that it mainly affects processing in the soma and
proximal dendrites. These are preferentially targeted by layer 5 pyramidal neurons, which
sparsely interconnect with projection-type specificity [77-79]. A confluence of evidence implicates
PV interneurons in the production of high-frequency inhibitory rhythms such as gamma [80, 81].
Given that PV interneurons diffusely interconnect with local pyramidal neurons [82, 83], and the
synapses they receive from them are depressing [75, 76], they likely regulate abrupt increases in
local ensemble activity. Interactions within ensembles that precede activating the PV population
will be boosted, while those following will be attenuated.
499
**Interneuron specializations and rhythm timescales**
Lastly, it is worth considering why beta and gamma rhythms are primarily mediated by different
types of interneurons (but see [24]). One possibility is that the differing pace merely arises from
the electrotonic timescales of their feedback inhibition; SOM cells target distal dendrites and thus
produce a slower inhibitory feedback signal, versus PV cells delivering rapid perisomatic
inhibition. Indeed, we found that swapping the cellular compartments receiving beta and gamma
rhythmic inhibition impaired their effectiveness and lowered overall excitability.
So, while our results suggest that spatial targeting of SOM and PV interneurons aligns
with the timescales of their *network-level* rhythms, it could also be that their timing and subcellular
localization interact to produce specialized *neuron-level* functions [84]. For instance, NMDA and
$Ca^{2+}$ spikes in the distal dendrites last for ~50 ms, making the slower beta rhythm more
appropriate for bidirectionally controlling them. Both can be described as dynamical systems with
distinct phases with differing sensitivity to inhibition. $Ca^{2+}$ spikes are dynamical events comprised
of an initiation, plateau, and termination phase. Inhibition delivered during the plateau phase
shortens their duration [85]. If the beta rhythm is comprised of cycling between periods of elevated
excitation (increased NMDA spike generation) followed by elevated inhibition, then $Ca^{2+}$ spike
initiation will tend to occur during the excitatory phase, and its plateau during the subsequent
inhibitory phase. A plateau during the inhibitory phase will more quickly enter termination. This is
bidirectional control. On the other hand, slower rhythms (e.g., 1 Hz) initiate $Ca^{2+}$ spikes during the
excitatory phase that plateau and enter termination autonomously, before the inhibitory phase is
reached. The same principle holds for NMDA spikes [86]. As a result, rhythms in the range from
15-30 Hz are optimal for synchronizing the onsets and offsets of dendritic spikes across a
population of neurons.
The integrative effects of gamma (>40 Hz) are also specialized. Low frequency inhibitory
rhythms delivered to the soma tended to shift the membrane potential higher or lower with the
rhythm's phase, effectively bringing it closer or farther from action potential threshold but not
changing the neuron's sensitivity to fast synaptic inputs. In the gamma frequency range, this is
reversed, with the mean membrane potential not varying with rhythm phase but with a shifting
bias to positive or negative membrane potential fluctuations. In addition, the trough phase of
gamma lowers the threshold for action potential initiation, while slower rhythms like beta only raise
the threshold. Consequently, the timing of gamma is ideal for increasing the sensitivity of the
neuron to rapid excitation. This agrees with the observation that gamma oscillations accompany
rapid excitation-inhibition balancing [87].
533
**Conclusion**

For the most part, the study of beta and gamma rhythms has focused on their correlation
with task-related events and entrainment of neuronal firing. Fundamentally, these effects derive
from their influence on neuronal integration, which has multiple stages, from modulating the post-
synaptic response, inducing a dendritic spike, and triggering an action potential. Since beta and
gamma rhythms differentially impacted these processes, this invites a reappraisal of their role in
coding and communication, along with unveiling a new hypothesis space for understanding their
function.
542
543

544 **METHODS**
545
We adapted a previously published model of a L5 pyramidal neuron [12, 88] to the Brain
Modeling Toolkit [89] format to facilitate reproducibility. This open-source software utilized
NEURON 7.7 [90] with an integration time step of 0.1 ms for simulating the membrane potential.
We added synapses and designed inputs in accordance with the literature to study synaptic
integration in vivo. Details of the passive membrane properties and active conductances are
only described briefly (full details can be found in [88]). Our focus here is on describing the
enhancements we incorporated to mimic in vivo features in the model neuron.
553
*L5 pyramidal cell compartmental model*
Briefly, the cell was built on a morphological reconstruction of a L5 pyramidal tract neuron and
included 10 active Hodgkin-Huxley conductances with kinetics taken from the literature [2].
These channels were a fast-inactivating Na$^+$ current ($I_{NaT}$), persistent Na$^+$ current ($I_{NaP}$), non-
specific cation current ($I_h$), muscarinic K$^+$ current ($I_m$), slow inactivating K$^+$ current ($I_{Kp}$), fast
inactivating K$^+$ current ($I_{Kt}$), fast non-inactivating K$^+$ current ($I_{Kv3.1}$), high voltage-activated Ca$^{2+}$
current ($I_{Ca\_HVA}$), low voltage-activated Ca$^{2+}$ current ($I_{Ca\_LVA}$), and Ca$^{2+}$ activated K$^+$ current ($I_{SK}$).
The somatic compartments contained all of these channels except $I_m$. Both the apical and basal
dendrite compartments contained $I_{Ca\_LVA}$, $I_{Ca\_HVA}$, $I_{SK}$, $I_{Kv3.1}$, $I_{NaT}$, $I_m$, and $I_h$. The membrane
capacitance was set to 1 $\mu$F/cm$^2$ for the soma and axon and 2 $\mu$F/cm$^2$ for the basal and apical
dendrites to correct for dendritic spine area. Axial resistance was set to 100 W-cm and leak
reversal potential was -90 mV.
566
*Divergence, probability of transmission, and functional clustering*
Experimental studies indicate that each presynaptic excitatory neuron forms 2-8 synapses with
its pyramidal partners, with a release probability between 0.16 and 0.9 [91]. Further, the 2-8
synaptic contacts occur over a span of 20-100 $\mu$m [92]). We constrained our model to have
these properties. In addition, since synapses may cluster on dendritic branches such that
functional presynaptic cell assemblies activate the same branch within 5-10 $\mu$m [93, 94], we
forced synapses coming from the same presynaptic cell assembly to cluster in this way. In
summary, a single presynaptic input fiber could have 2-8 synapses, drawn from a uniform
distribution with a release probability between 0.16 and 0.9, also drawn from a uniform
distribution 91]. And the synapses from a functional group formed clusters within 5-10$\mu$m on the
same dendritic branch over a span of 100 $\mu$m.
578
*Synaptic conductances*
We randomly distributed excitatory synaptic conductances following previous modeling work
[13]. Maximum conductance values followed a log-normal distribution with a mean of 0.2 nS and
standard deviation of 0.345 nS for excitatory synaptic conductances. Inhibitory synaptic
conductances were fixed at 1 nS [12]. We validated these conductances against excitatory and
inhibitory post-synaptic currents measured *in vitro.* The model's values are compared with
experimental values in Table 1.
586
*Synapse density*
Synapse density was based on a comprehensive report [95] that not only quantified the number
of spines but also counted the synapses by identifying vesicle clouds and postsynaptic density
using serial block-face electron microscopy. The authors found the apical tuft synapse density
per micron as 2.16 ± 0.16 for excitatory and 0.22 ± 0.02 for inhibitory synapses (mean ±
standard error of the mean). Other studies that only quantified spine density may undercount
the true number of synapses by about 3 times because one can only count lateral spines, not

those in the z-axis [96]. With that adjustment, the other values reported – 0.47 ± 0.01 spines/µm
[97]  and 0.53 ± 0.05 spines/µm [98]– are roughly in alignment with that in [95]. We used the
values from [95] for excitatory synapse density because we found the higher value for excitatory
synapse density was needed to achieve a reasonable baseline firing rate.
Morphological measurements (length of dendritic cable in µm) combined with these
synapse density estimates yield the following estimates for total number of synapses:
Excitatory: 16070 (2.16 syns/µm * 7440 µm) on apical dendrites and 10042 (2.16 syns/µm *
4649 µm) on basal dendrites. Inhibitory: 150 on soma [12, 99, 100] + (0.22 syns/µm *483 mm) =
256 perisomatic, 1637 (0.22 syns/µm * 7440 µm) on apical dendrites, 1023 (0.22 syns/µm *
4649 µm) on basal dendrites.
604
*Synaptic kinetics and short-term plasticity*
All excitatory transmission was mediated by AMPA/NMDA receptors and inhibitory transmission
by GABA$_A$ receptors as in our prior models. The rise/decay constants for the excitatory
connections were as follows: Rise/decay time constants of 0.6/6.9 ms for AMPA, and 3.7/125
ms for NMDA synapses, and a reversal potential of 0 for both. And for the inhibitory GABA-ergic
connections, the rise/decay time constants were 0.5 and 6.8 ms, respectively, with a reversal
potential of -75 mV. We implemented short-term depression in both the excitatory and inhibitory
synapses [76] via an exponential decay of synaptic weight over short time scales [101].
Following [76] we calculated a short-term plasticity (STP) train-induction value by applying 8
pulses at 50 Hz to the synapse while in voltage clamp and measured the resulting postsynaptic
current (PSC). The 6$^{th}$, 7$^{th}$, and 8$^{th}$ PSCs were averaged, and the 1$^{st}$ PSC was subtracted from
the result. This value was divided by the max PSC current to yield the STP train-induction value.
Excitatory synapses to L5 pyramidal neurons had an STP train-induction value of -0.34.
Inhibitory synapses arising from PV+ basket interneurons making perisomatic contact had a
value of -0.60 and inhibitory synapses arising from SOM+ interneurons making dendritic contact
had a value of -0.10. The equations to model the synaptic currents and short-term depression
are provided in the next section.
622
623   *Modeling of synaptic currents*
624   The corresponding synaptic currents were modeled by dual exponential functions [102, 103]. All
625   excitatory transmission was mediated by AMPA/NMDA receptors and inhibitory transmission by
626   GABA$_A$ receptors. The corresponding synaptic currents were modeled by dual exponential
627   functions [102, 103], as shown in eqns. 1-3,
628

$$I_{AMPA} = w \times G_{AMPA} \times (V - E_{AMPA})$$
$$G_{AMPA} = g_{AMPA,max} \times D_{AMPA} \times r_{AMPA}$$
$$r_{AMPA}' = \alpha Tmax_{AMPA} \times ON_{AMPA} \times (1 - r_{AMPA}) - \beta_{AMPA} \times r_{AMPA} \qquad (1)$$

$$I_{NMDA} = w \times G_{NMDA} \times (V - E_{NMDA})$$
$$G_{NMDA} = g_{NMDA,max} \times D_{NMDA} \times s(V) \times r_{NMDA}$$
$$r_{NMDA}' = \alpha Tmax_{NMDA} \times ON_{NMDA} \times (1 - r_{NMDA}) - \beta_{NMDA} \times r_{NMDA} \qquad (2)$$

$$I_{GABAa} = w \times G_{GABAa} \times (V - E_{GABAa})$$
$$G_{GABAa} = g_{GABAa,max} \times D_{GABAa} \times r_{GABAa}$$
$$r_{GABAa}' = \alpha Tmax_{GABAa} \times ON_{GABAa} \times (1 - r_{GABAa}) - \beta_{GABAa} \times r_{GABAa} \qquad (3)$$

where *V* is the membrane potential (mV) of the compartment (dendrite or soma) where the
synapse is located, *I* is the current injected into the compartment (nA), *G* is the synaptic

conductance (µS), $w$ is the synaptic weight (unitless), and $E$ is the reversal potential of the
synapse (mV). $g_{x,max}$ is the maximal conductance (µS), $D$ implements short-term plasticity as
defined below, and $r_x$ determines the synaptic current rise and decay time constants based on
the terms $\alpha Tmax$ and $\beta$ [102]. The voltage-dependent variable $s(V)$ which implements the Mg$^{2+}$
block was defined as: $s(V) = [1 + 0.33 \exp(-0.06\ V)]^{-1}$ [104]. The terms $ON_{NMDA}$ and $ON_{AMPA}$ are
set to 1 if the corresponding receptor is open, else to 0. To calculate the factor D for depression:
$\tau\_D^*dD/dt=1-D$ and D constrained to be ≤ 1. After each stimulus, D was multiplied by a constant
d (≤ 1) representing the amount of depression per presynaptic action potential and updated as
D→D*d. Between stimuli, D recovered exponentially back toward 1. We modelled depression
using two factors d1 and d2 with d1 being fast and d2 being slow subtypes, and d=d1*d2 was
constrained to be ≥ 1.
654
*Synapse distribution and extrinsic inputs*
We distributed 26,112 excitatory synapses over the cell. With an average of 5 synapses per pre-
synaptic neuron, this yielded 5,222 presynaptic nodes. To represent pre-synaptic cell
assemblies, we defined "functional groups" of presynaptic nodes. We put 100 nodes in each
functional group, yielding 52 functional groups. We found that varying the number of neurons in
the cell assembly did not qualitatively affect our results. To have exactly 100 nodes per
functional group we rounded the total number of nodes down to 5,200. Inhibitory inputs were not
defined by functional groups, but followed the literature cited in Table 1, with more than 1
synapse per pre-synaptic node. For example, with 1637 inhibitory synapses on the apical
dendrites and an average of 12 synapses per pre-synaptic node, there were about 136
inhibitory nodes that contacted the apical dendrites. The numbers and distributions of model
synapses along the dendrites are shown in Table 1.
The extrinsic inputs to the model were represented by spike trains arriving at the
corresponding synapses along the dendritic tree. The L5 pyramidal model is spatially divided
into the following sub-divisions: somatic, perisomatic, and proximal and distal components of the
apical and basal trees. We used standard definitions for these sub-divisions except for
perisomatic which is typically defined as the dendritic region within 100 µm of the soma [105].
Tonic inputs (Figs. 1-3): The firing rate of each excitatory spike train was randomly sampled
from an experimentally recorded distribution (Headley, personal communication). To reproduce
the 1/f spectrum that is widely found in neuronal spike trains, the 100 spike trains in each
functional group were modulated by a unique normalized (from 0.5 to 1.5) firing rate time series
that had a 1/f spectrum (more slow modulation of firing than fast) that was created by filtering
white noise using standard techniques. The normalized firing rate time series was then
multiplied with the mean firing rate of each of the 100 pre-synaptic nodes. Spike trains are then
generated by sampling from a Poisson distribution using the firing rate time series of each
presynaptic node of the functional group. In this way, spike trains that received the same
modulation trace would be statistically correlated, representing functional presynaptic cell
assemblies. Since feed-forward inhibition follows excitation in time, we averaged the firing rate
time series used to design the excitatory spike trains, shifted it forward in time by 4 milliseconds,
and used this to modulate the firing rate of the inhibitory spike trains.
Enhanced tonic inhibition (Fig. 4): Inhibitory synapses were divided into two groups, those
targeting perisomatic region that originate mainly from PV+ interneurons, and those on more
distal dendritic regions arising mainly from SOM+ interneurons [1, 4]. Perisomatic synapses
were defined as those within 100 µm of the soma with the properties specified in Table 1 under
"Inhibitory (Perisomatic and somatic)." Distal inhibitory synapses were greater than 100 µm from
the soma and had their properties specified in sections "Inhibitory (Basal)" and "Inhibitory
(Apical)" of the table To investigate if these inhibitory sources differentially regulated action
potential initiation and dendritic spiking, we doubled the rate of presynaptic events to one of
these inhibitory populations.

Investigation of E/I lag (Fig. 4F-I): We used the same scheme to design the extrinsic inputs as in
Figs. 1-3. However, to investigate the effect of E/I lag, we increased the variance of the overall
excitatory firing rate. So, while the mean rate stayed the same, we raised the modulatory trace
to the fourth power so that large deviations from the mean were exaggerated. This excitatory
firing rate trace was used to modulate the inhibitory firing rates (same as for Figs. 1-3). The
perisomatic and distal traces were shifted independently. One was fixed at 4 ms while the other
was shifted by 4, 125, 250, or 500 ms.
Rhythmic inhibition (Figs. 5 & 6): Gamma rhythms depend on PV+ interneurons, while beta
rhythms arise from SOM+ interneurons  [21]. Thus, to emulate an inhibitory rhythm we
rhythmically modulated inhibitory synapses in the respective dendritic region (see description
above in "Enhanced tonic inhibition") at the appropriate frequency. Beta rhythms were emulated
by modulating distal inhibitory synapses at 16 Hz with a depth of 20%. Gamma rhythms were
emulated by modulating perisomatic synapses at 64 Hz with a depth of 40%. The different
depths of modulation ensured that these rhythms produced similar entrainment of action
potentials.
To generate inhibitory spike trains that produce rhythmic inhibition during oscillations, we
modulated their collective firing rate with a sine wave:

$$r[t] = A * (\sin(2 * \pi * f * t) + \phi) + off \qquad (4)$$

where $A = \frac{off}{\frac{1}{d}-1}$, $f$ is the frequency, $t$ is a vector representing time, $\phi$ is the phase, $off$ is the
offset firing rate of the spike train being modulated, and $0 < d < 1$ is the depth of modulation
which represents the amplitude of the sine wave relative to $off$. To generate the spike train, a
random vector x[t] was created with values uniformly distributed between 0 and 1. A spike was
generated if $x[t] \leq r[t]dt$ where dt in our case was 1/1000.
Varying frequency of rhythmic inhibition (Figs. 7 & 8): We reran the simulations used in Figures
5 and 6, but across a range of frequencies that extended above and below the beta and gamma
frequencies we used in all other simulations (0.5, 2, 5, 10, 16, 20, 30, 40, 50, 60, 80 Hz).
Bursty inhibition (Fig. 9): We applied an envelope to the rhythmic inhibitory modulation trace to
simulate the bursty temporal nature of rhythms *in vivo*. The underlying sine wave was produced
in the same way as in the previous section. The envelope was a Gaussian kernel with a
standard deviation of two oscillatory cycles.
Clustered synaptic input (Fig. 10): For this experiment we kept everything the same as in the
rhythmic inhibition simulations, but introduced additional synapses driven by 40 pre-synaptic
spike trains. These spike trains were created by choosing one Poisson process and jittering the
spike times by ±2 ms. Each spike train drove an average of 4.6 synapses for a total of 187
synapses. These were placed either on the apical dendrites near the nexus or the basal
dendrites near the soma. For both nexus and basal clustered synapses, we targeted 140 $\mu$m of
dendritic cable length with 187 synapses yielding a synaptic density of 1.3 synapses per micron.
For the nexus case we targeted the 140 µm region starting at the bifurcation going toward the
apical dendrites. For the basal case, we targeted 3 basal dendrites that were directly connected
to the soma each one had a length of 40-50 µm adding up to 140 $\mu$m total (Figure 7A). The
synaptic weights were drawn from the distribution in Table 1.
735
736    *Detection of action potentials*
737    Action potentials were detected by the NEURON software as being any time point when the
738    somatic membrane potential crossed -10 mV with a positive first derivative.
739
*Detection of dendritic spikes*
Na$^+$ and NMDA dendritic spike events were detected following the definitions developed in [13].
We developed our own detection algorithm for Ca$^{2+}$ spikes since they were not previously
studied. Na$^+$ spikes were detected if the voltage-gated sodium conductance density for a

compartment exceeded 0.3 mS/cm$^2$. If sodium spikes were within 2 ms of a somatic action
potential, they were assumed to be a consequence of a backpropagating action potential and
were not counted. An example of a Na$^+$ spike can be seen in Fig. 1C1.
NMDA spikes were detected by two criteria – voltage and current. The voltage criteria
were met if the membrane potential rose above -40 mV and remained there for 26 ms or more.
The current criteria was met if the NMDA current exceeded 130% of the current at the -40 mV
crossing and continued to be met until the current fell to 115% of that value.  Whenever both the
voltage and current criteria were met, an NMDA spike was recorded. An example of an NMDA
spike is shown in Fig. 1C2.
Ca$^{2+}$ spikes were detected in the same way as NMDA spikes – using a voltage and
current threshold. The only difference was instead of NMDA current for the current criteria we
used the sum of HVA and LVA calcium currents. An example is shown in Fig. 1C3.
756
*Determination of electrotonic distance*
We calculated electrotonic distance using NEURON's built-in Impedance method. The method
calculates signal attenuation between two compartments at a given frequency. We used 20 Hz
because the wavelength is on a time scale comparable to the membrane time constant (10s of
milliseconds). However, changing this frequency 10 Hz in either direction did not qualitatively
affect our results.
763
*Spike-triggered average (STA) of dendritic spikes (Figs. 2, 3)*
For each dendritic compartment, we created a binary time series with 2 ms bins that was true
during periods where a dendritic spike was present (based on their detected start and stop
times). This was done separately for each type of dendritic spike (Na$^+$, NMDA, and Ca$^{2+}$ spikes).
Each then had its spike-triggered average taken around the times of somatic action potentials
and their percent change from the mean across the entire simulation calculated. The STAs were
then grouped together based on their electrotonic distance percentile (10% bins) from the soma
and type of dendrite (apical or basal), and the median taken for each time lag.
772
*Generation of F-I curves (Fig. 4)*
The model with Poisson excitation and inhibition was used. In the *perisomatic* condition, we
doubled the rate of inhibition to the somatic and proximal dendritic compartments (<100µm from
soma). In the *distal* case, we doubled the rate of inhibition to the distal dendritic compartments
(>100µm from soma). In each case, we injected 20 s current steps ranging from -1 nA to 2.4 nA
in 200 pA intervals.
779
*Phase histogram of dendritic spikes (Fig. 5)*
Entrainment of dendritic spikes to inhibitory rhythms was calculated by first converting dendritic
spikes to a binary time series as was done for the STA calculation. The inhibitory rhythm was
then binned by phase, with bin widths of π/4, yielding 8 phase bins. The mean of the dendritic
spike binary series was taken for each phase bin. These were converted to percent change
from mean and their median taken across compartments by their electronic distance and
dendritic type (the same as in the STA calculation).
787
788  *Determination of action potential threshold and somatic membrane voltage distribution (Fig. 6)*
789  Action potential threshold was taken to be the voltage 1 ms prior to the peak of the spike.
790  Voltage distributions were calculated using 1 ms bins.
791
*Entrainment of dendritic spike onsets (Fig. 7)*
We measured the entrainment of dendritic spike onsets to the inhibitory rhythm using the pair-
wise phase consistency index, an unbiased metric [106].

*Phase histogram of dendritic spikes (Fig. 9)*
For calculating entrainment of dendritic spikes to the rhythm, we followed the same procedure
as detailed above for Fig. 5.
798
*Cross-correlation (CC) between presynaptic spikes and action potentials (Fig. 10)*
We measured the effect of inhibitory rhythms on the coupling between concentrated synaptic
inputs and action potentials using a phase-stratified CC. Presynaptic spike times that drove the
concentrated synaptic inputs were grouped together and separated based on whether they
occurred during the positive or negative phase of the imposed inhibitory rhythm. Their CC with
somatic action potentials was then calculated using 1 ms bins. Since both the presynaptic and
action potential times had inherent periodicities, these had to be corrected for [25]. This was
performed by converting the CC and both autocorrelations to the frequency domain using the
fast Fourier transform, setting the coefficients at the inhibitory modulation rhythm to 0,
performing element-wise division of the CC by the square-root of the autocorrelations multiplied
together, and then converting back to the time domain using the inverse fast Fourier transform
on that result. To measure the strength of the cross-correlation peak, we identified the positive
peak of the CC nearest to zero time lag, and then integrated over its area with the edges
defined by when the CC crossed from positive to negative values.
813
814 **CODE AND DATA AVAILABILITY**
815 Simulation code is deposited at ModelDB at https://modeldb.science/2019883. The raw
816 simulation data are available from DBH upon request. Analysis code is posted as a GitHub repo
817 at https://github.com/dbheadley/InhibOnDendComp.
818
819

820  **Table 1.** Inputs to L5 PN. L5 PN dendrites can course up to layer 1 and receive both excitatory
821  and inhibitory inputs to their dendrites. Here we quantified, where possible, the experimental
822  values for synaptic magnitude, firing rate, divergence, and release probability. We matched the
823  model parameters to the experimental values as closely as possible while preserving a
824  reasonable basal firing rate.
825

| Synapse type | Characteristics | Model | Experimental |
|---|---|---|---|
| Excitatory (Basal) | Magnitude of EPSCs | 37.0 ± 32.3 pA | **30.6 ± 29.9 pA** [78] |
| | Firing rate | 4.43 ± 2.9 Hz | **4.43 ± 2.9 Hz** (Headley personal comm.) |
| | Divergence | 2-8 | **2-8**  [92, 100, 107] |
| | Number of synapses | 10042 | **10042** [95] |
| | Release probability | 0.53 ± 0.22 | **0.53 ± 0.22** [108] |
| Excitatory (Apical) | Magnitude | 25.9 ± 24.9 pA | **30.6 ± 29.9 pA** [78] |
| | Firing rate | 4.43 ± 2.9 Hz | **4.43 ± 2.9 Hz** (Headley personal comm.) |
| | Divergence | 2-8 | **2-8**  [92, 100, 107] |
| | Number of synapses | 16070 | **16070** [95] |
| | Release probability | 0.53 ± 0.22 | **0.53 ± 0.22** [108] |
| Inhibitory (Perisomatic and somatic) | Magnitude | 162.5 ± 103.1 pA | **208.3 ± 58.7 pA** [109] |
| | Firing rate | 16.9 ± 14.3 Hz | **16.9 ± 14.3 Hz** [110] |
| | Divergence | 2.8 ± 1.9 | **2.8 ± 1.9** |
| | Number of synapses | 406 | **406** [95] |
| | Release probability | 0.88 ± 0.05 | **0.88 ± 0.05** [109] |
| Inhibitory (Basal) | Magnitude | 24.3 ± 18.4 pA | **26.5 ± 1.6 pA** [109] |
| | Firing rate | 3.9 ± 4.9 Hz | **3.9 ± 4.9 Hz**  [110] |
| | Divergence | 2.7 ± 1.6 | **2.7 ± 1.6** [111, 112] |
| | Number of synapses | 1023 | **1023** [95] [10] |
| | Release probability | 0.72 ± 0.10 | **0.72 ± 0.10**  [109] |
| Inhibitory (Apical) | Magnitude | 24.3 ± 33.1 pA | **26.5 ± 1.6 pA** [109] |
| | Firing rate | 3.9 ± 4.9 Hz | **3.9 ± 4.9 Hz** [110] |
| | Divergence | 12 ± 3 | **12 ± 3** [113][33] |
| | Number of synapses | 1637 | **1637** [95] |
| | Release probability | 0.30 ± 0.08 | **0.30 ± 0.08** [113] |

826
827

828　**Table 2.** Short term pre-synaptic plasticity. Parameters were tuned to match experimental
829　recordings reported in [76].
830

| Synapse type | tauD1 (ms) | d1 | tauD2 (ms) | d2 | Synaptic current @ 20 Hz | Train induction vs. frequency |
|---|---|---|---|---|---|---|
| Excitatory | 35 | 0.95 | 250 | 0.8 |  |  |
| Perisomatic inh. | 40 | 0.7 | 500 | 0.7 |  |  |
| Dendritic inh. | 200 | 0.8 | 1 | 1 |  |  |

831
832
833

**REFERENCES**

1.  Tremblay, R., S. Lee, and B. Rudy, *GABAergic interneurons in the neocortex: From cellular properties to circuits.* Neuron, 2016. **91**(2): p. 260–92.
2.  Kawaguchi, Y. and Y. Kubota, *GABAergic cell subtypes and their synaptic connections in rat frontal cortex.* Cereb Cortex, 1997. **7**(6): p. 476–86.
3.  Wang, Y., et al., *Anatomical, physiological and molecular properties of Martinotti cells in the somatosensory cortex of the juvenile rat.* J Physiol, 2004. **561**(Pt 1): p. 65–90.
4.  Kubota, Y., *Untangling GABAergic wiring in the cortical microcircuit.* Curr Opin Neurobiol, 2014. **26**: p. 7–14.
5.  Kubota, Y., et al., *Functional effects of distinct innervation styles of pyramidal cells by fast spiking cortical interneurons.* Elife, 2015. **4**.
6.  Wilson, N.R., et al., *Division and subtraction by distinct cortical inhibitory networks in vivo.* Nature, 2012. **488**(7411): p. 343–8.
7.  Lee, S.H., et al., *Activation of specific interneurons improves V1 feature selectivity and visual perception.* Nature, 2012. **488**(7411): p. 379–83.
8.  Atallah, B.V., et al., *Parvalbumin-expressing interneurons linearly transform cortical responses to visual stimuli.* Neuron, 2012. **73**(1): p. 159–70.
9.  Leleo, E.G. and I. Segev, *Burst control: Synaptic conditions for burst generation in cortical layer 5 pyramidal neurons.* PLOS Computational Biology, 2021. **17**(11): p. e1009558.
10. Jadi, M., et al., *Location-dependent effects of inhibition on local spiking in pyramidal neuron dendrites.* PLoS Comput Biol, 2012. **8**(6): p. e1002550.
11. Ferguson, K.A. and J.A. Cardin, *Mechanisms underlying gain modulation in the cortex.* Nat Rev Neurosci, 2020. **21**(2): p. 80–92.
12. Egger, R., et al., *Cortical output is gated by horizontally projecting neurons in the deep layers.* Neuron, 2020. **105**(1): p. 122–137 e8.
13. Goetz, L., A. Roth, and M. Häusser, *Active dendrites enable strong but sparse inputs to determine orientation selectivity.* Proceedings of the National Academy of Sciences, 2021. **118**(30): p. e2017339118.
14. Ujfalussy, B.B., et al., *Global and multiplexed dendritic computations under in vivo-like conditions.* Neuron, 2018. **100**: p. 579–592.e5.
15. Beniaguev, D., I. Segev, and M. London, *Single cortical neurons as deep artificial neural networks.* Neuron, 2021. **109**(17): p. 2727–2739.e3.
16. Larkum, M.E., J.J. Zhu, and B. Sakmann, *A new cellular mechanism for coupling inputs arriving at different cortical layers.* Nature, 1999. **398**(6725): p. 338–41.
17. Poirazi, P., T. Brannon, and B.W. Mel, *Pyramidal neuron as two-layer neural network.* Neuron, 2003. **37**(6): p. 989–99.
18. Larkum, M.E., et al., *Synaptic integration in tuft dendrites of layer 5 pyramidal neurons: a new unifying principle.* Science, 2009. **325**(5941): p. 756–60.
19. Li, X., et al., *Control of layer 5 pyramidal cell spiking by oscillatory inhibition in the distal apical dendrites: a computational modeling study.* J Neurophysiol, 2013. **109**(11): p. 2739–56.
20. Sohal, V.S., et al., *Parvalbumin neurons and gamma rhythms enhance cortical circuit performance.* Nature, 2009. **459**: p. 698–702.
21. Chen, G., et al., *Distinct inhibitory circuits orchestrate cortical beta and gamma band oscillations.* Neuron, 2017. **96**(6): p. 1403–1418.e6.
22. Veit, J., et al., *Cortical gamma band synchronization through somatostatin interneurons.* Nat Neurosci, 2017.
23. Onorato, I., et al., *Distinct roles of PV and Sst interneurons in visually induced gamma oscillations.* Cell Rep, 2025. **44**(3): p. 115385.

884    24.    Tahvili, F., M. Vinck, and M. di Volo, *PV and SOM cells play distinct causal roles in*
885           *controlling network oscillations and stability.* Cell Rep, 2025. **44**(8): p. 116131.

886    25.    Headley, D.B. and N.M. Weinberger, *Fear conditioning enhances γ oscillations and their*
887           *entrainment of neurons representing the conditioned stimulus.* J Neurosci, 2013. **33**(13):
888           p. 5705–17.

889    26.    Murty, D., et al., *Large visual stimuli induce two distinct gamma oscillations in primate*
890           *visual cortex.* J Neurosci, 2018. **38**(11): p. 2730–2744.

891    27.    Muthukumaraswamy, S.D., *Functional properties of human primary motor cortex gamma*
892           *oscillations.* J Neurophysiol, 2010. **104**(5): p. 2873–85.

893    28.    Sanes, J.N. and J.P. Donoghue, *Oscillations in local field potentials of the primate motor*
894           *cortex during voluntary movement.* Proc Natl Acad Sci U S A, 1993. **90**(10): p. 4470–4.

895    29.    Lundqvist, M., et al., *Gamma and beta bursts underlie working memory.* Neuron, 2016.
896           **90**(1): p. 152–164.

897    30.    Bastos, A.M., et al., *Layer and rhythm specificity for predictive routing.* Proc Natl Acad
898           Sci U S A, 2020. **117**(49): p. 31459–31469.

899    31.    Bastos, A.M., et al., *Visual areas exert feedforward and feedback influences through*
900           *distinct frequency channels.* Neuron, 2015. **85**(2): p. 390–401.

901    32.    Michalareas, G., et al., *Alpha-beta and gamma rhythms subserve feedback and*
902           *feedforward influences among human visual cortical areas.* Neuron, 2016. **89**(2): p. 384–
903           97.

904    33.    Vezoli, J., et al., *Brain rhythms define distinct interaction networks with differential*
905           *dependence on anatomy.* Neuron, 2021. **109**(23): p. 3862–3878.e5.

906    34.    Vinck, M., et al., *Principles of large-scale neural interactions.* Neuron, 2023. **111**(7): p.
907           987–1002.

908    35.    Tiesinga, P.H., et al., *Inhibitory synchrony as a mechanism for attentional gain*
909           *modulation.* J Physiol Paris, 2004. **98**(4-6): p. 296–314.

910    36.    Feng, F., et al., *Gamma oscillations in the basolateral amygdala: Biophysical*
911           *mechanisms and computational consequences.* eNeuro, 2019. **6**(1): p. ENEURO.0388–
912           18.2018.

913    37.    Saiki, A., et al., *In vivo spiking dynamics of intra- and extratelencephalic projection*
914           *neurons in rat motor cortex.* Cereb Cortex, 2018. **28**(3): p. 1024–1038.

915    38.    Golding, N.L. and N. Spruston, *Dendritic sodium spikes are variable triggers of axonal*
916           *action potentials in hippocampal CA1 pyramidal neurons.* Neuron, 1998. **21**(5): p. 1189–
917           200.

918    39.    Schiller, J., et al., *NMDA spikes in basal dendrites of cortical pyramidal neurons.* Nature,
919           2000. **404**(6775): p. 285–9.

920    40.    Schiller, J., et al., *Calcium action potentials restricted to distal apical dendrites of rat*
921           *neocortical pyramidal neurons.* J Physiol, 1997. **505 ( Pt 3)**(Pt 3): p. 605–16.

922    41.    Larkum, M.E. and J.J. Zhu, *Signaling of layer 1 and whisker-evoked Ca2+ and Na+*
923           *action potentials in distal and terminal dendrites of rat neocortical pyramidal neurons in*
924           *vitro and in vivo.* J Neurosci, 2002. **22**(16): p. 6991–7005.

925    42.    Smith, S.L., et al., *Dendritic spikes enhance stimulus selectivity in cortical neurons in*
926           *vivo.* Nature, 2013. **503**(7474): p. 115–20.

927    43.    Helmchen, F., et al., *In vivo dendritic calcium dynamics in deep-layer cortical pyramidal*
928           *neurons.* Nat Neurosci, 1999. **2**(11): p. 989–96.

929    44.    Xu, N.L., et al., *Nonlinear dendritic integration of sensory and motor input during an*
930           *active sensing task.* Nature, 2012. **492**(7428): p. 247–51.

931    45.    Fletcher, L.N. and S.R. Williams, *Neocortical topology governs the dendritic integrative*
932           *capacity of layer 5 pyramidal neurons.* Neuron, 2019. **101**(1): p. 76–90.e4.

933    46.    Gidon, A., et al., *Dendritic action potentials and computation in human layer 2/3 cortical*
934           *neurons.* Science, 2020. **367**(6473): p. 83–87.

47.    Larkum, M.E., et al., *Dendritic spikes in apical dendrites of neocortical layer 2/3*
*pyramidal neurons.* J Neurosci, 2007. **27**(34): p. 8999–9008.

48.    Doiron, B., et al., *Subtractive and divisive inhibition: effect of voltage-dependent*
*inhibitory conductances and noise.* Neural Comput, 2001. **13**(1): p. 227–48.

49.    Seybold, B.A., et al., *Inhibitory actions unified by network integration.* Neuron, 2015.
**87**(6): p. 1181–1192.

50.    Wehr, M. and A.M. Zador, *Balanced inhibition underlies tuning and sharpens spike*
*timing in auditory cortex.* Nature, 2003. **426**(6965): p. 442–6.

51.    Amir, A., et al., *Vigilance-associated gamma oscillations coordinate the ensemble*
*activity of basolateral amygdala neurons.* Neuron, 2018. **97**(3): p. 656–669.e7.

52.    Onorato, I., et al., *A distinct class of bursting neurons with strong gamma*
*synchronization and stimulus selectivity in monkey V1.* Neuron, 2020. **105**(1): p. 180–
197.e5.

53.    Adesnik, H., *Layer-specific excitation/inhibition balances during neuronal*
*synchronization in the visual cortex.* J Physiol, 2018. **596**(9): p. 1639–1657.

54.    Murthy, V.N. and E.E. Fetz, *Coherent 25- to 35-Hz oscillations in the sensorimotor*
*cortex of awake behaving monkeys.* Proc Natl Acad Sci U S A, 1992. **89**(12): p. 5670–4.

55.    Rojas, P., et al., *Differential effects of axon initial segment and somatodendritic GABAA*
*receptors on excitability measures in rat dentate granule neurons.* J Neurophysiol, 2011.
**105**(1): p. 366–79.

56.    Gidon, A. and I. Segev, *Principles governing the operation of synaptic inhibition in*
*dendrites.* Neuron, 2012. **75**(2): p. 330–41.

57.    Iacaruso, M.F., I.T. Gasler, and S.B. Hofer, *Synaptic organization of visual space in*
*primary visual cortex.* Nature, 2017. **547**(7664): p. 449–452.

58.    Fu, M., et al., *Repetitive motor learning induces coordinated formation of clustered*
*dendritic spines in vivo.* Nature, 2012. **483**(7387): p. 92–5.

59.    Salinas, E. and T.J. Sejnowski, *Impact of correlated synaptic input on output firing rate*
*and variability in simple neuronal models.* J Neurosci, 2000. **20**(16): p. 6193–209.

60.    Fries, P., *A mechanism for cognitive dynamics: neuronal communication through*
*neuronal coherence.* Trends in Cognitive Sciences, 2005. **9**(10): p. 474–480.

61.    Hopfield, J.J., *Pattern recognition computation using action potential timing for stimulus*
*representation.* Nature, 1995. **376**(6535): p. 33–6.

62.    Volgushev, M., M. Chistiakova, and W. Singer, *Modification of discharge patterns of*
*neocortical neurons by induced oscillations of the membrane potential.* Neuroscience,
1998. **83**(1): p. 15–25.

63.    Vinck, M., et al., *Gamma-phase shifting in awake monkey visual cortex.* J Neurosci,
2010. **30**(4): p. 1250–7.

64.    Prescott, S.A. and Y. De Koninck, *Gain control of firing rate by shunting inhibition: roles*
*of synaptic noise and dendritic saturation.* Proc Natl Acad Sci U S A, 2003. **100**(4): p.
2076–81.

65.    Cardin, J.A., et al., *Driving fast-spiking cells induces gamma rhythm and controls*
*sensory responses.* Nature, 2009. **459**: p. 663–7.

66.    Ferguson, B.R. and W.J. Gao, *PV interneurons: Critical regulators of E/I balance for*
*prefrontal cortex-dependent behavior and psychiatric disorders.* Front Neural Circuits,
2018. **12**: p. 37.

67.    Headley, D.B., et al., *Gamma oscillations in the basolateral amygdala: localization,*
*microcircuitry, and behavioral correlates.* J Neurosci, 2021. **41**(28): p. 6087–101.

68.    Perrenoud, Q., C.M. Pennartz, and L.J. Gentet, *Membrane potential dynamics of*
*spontaneous and visually evoked gamma activity in V1 of awake mice.* PLoS Biol, 2016.
**14**(2): p. e1002383.

69. Perrenoud, Q., et al., *Flexible perceptual encoding by discrete gamma events.* bioRxiv,
2022: p. 2022.05.13.491832.
70. Womelsdorf, T., et al., *Orientation selectivity and noise correlation in awake monkey
area V1 are modulated by the gamma cycle.* Proc Natl Acad Sci U S A, 2012. **109**(11):
p. 4302–7.
71. Lisitsyn, D., et al., *Visual stimulus content in V4 is conveyed by gamma-rhythmic
information packages.* J Neurosci, 2020. **40**(50): p. 9650–9662.
72. Siegel, M., M.R. Warden, and E.K. Miller, *Phase-dependent neuronal coding of objects
in short-term memory.* Proc Natl Acad Sci U S A, 2009. **106**(50): p. 21341–6.
73. D'Souza, R.D. and A. Burkhalter, *A laminar organization for selective cortico-cortical
communication.* Front Neuroanat, 2017. **11**: p. 71.
74. Harris, J.A., et al., *Hierarchical organization of cortical and thalamic connectivity.* Nature,
2019. **575**(7781): p. 195–202.
75. Beierlein, M., J.R. Gibson, and B.W. Connors, *Two dynamically distinct inhibitory
networks in layer 4 of the neocortex.* J Neurophysiol, 2003. **90**(5): p. 2987–3000.
76. Campagnola, L., et al., *Local connectivity and synaptic dynamics in mouse and human
neocortex.* Science, 2022. **375**(6585): p. eabj5861.
77. Kawaguchi, Y., *Pyramidal cell subtypes and their synaptic connections in layer 5 of rat
frontal cortex.* Cereb Cortex, 2017. **27**(12): p. 5755–5771.
78. Morishima, M., et al., *Highly differentiated projection-specific cortical subnetworks.* J
Neurosci, 2011. **31**(28): p. 10380–91.
79. Morishima, M. and Y. Kawaguchi, *Recurrent connection patterns of corticostriatal
pyramidal cells in frontal cortex.* J Neurosci, 2006. **26**(16): p. 4394–405.
80. Buzsáki, G. and X.-J. Wang, *Mechanisms of gamma oscillations.* Annual Review of
Neuroscience, 2012. **35**(1): p. 203–225.
81. Fernandez-Ruiz, A., et al., *Over and above frequency: Gamma oscillations as units of
neural circuit operations.* Neuron, 2023. **111**(7): p. 936–953.
82. Packer, A.M. and R. Yuste, *Dense, unspecific connectivity of neocortical parvalbumin-
positive interneurons: a canonical microcircuit for inhibition?* J Neurosci, 2011. **31**(37): p.
13260–71.
83. Morishima, M., et al., *Segregated excitatory-inhibitory recurrent subnetworks in layer 5 of
the rat frontal cortex.* Cereb Cortex, 2017. **27**(12): p. 5846–5857.
84. Lovett-Barron, M., et al., *Regulation of neuronal input transformations by tunable
dendritic inhibition.* Nat Neurosci, 2012. **15**(3): p. 423–30, S1–3.
85. Dudai, A., et al., *Synaptic input and ACh modulation regulate dendritic Ca(2+) spike
duration in pyramidal neurons, directly affecting their somatic output.* J Neurosci, 2022.
**42**(7): p. 1184–1195.
86. Doron, M., et al., *Timed synaptic inhibition shapes NMDA spikes, influencing local
dendritic processing and global I/O properties of cortical neurons.* Cell Rep, 2017. **21**(6):
p. 1550–1561.
87. Atallah, B.V. and M. Scanziani, *Instantaneous modulation of gamma oscillation
frequency by balancing excitation with inhibition.* Neuron, 2009. **62**(4): p. 566–77.
88. Hay, E., et al., *Models of Neocortical Layer 5b Pyramidal Cells Capturing a Wide Range
of Dendritic and Perisomatic Active Properties.* PLOS Computational Biology, 2011. **7**(7):
p. e1002107.
89. Dai, K., et al., *Brain Modeling ToolKit: An open source software suite for multiscale
modeling of brain circuits.* PLoS Comput Biol, 2020. **16**(11): p. e1008386.
90. Carnevale Nt, H.M.L., *The NEURON Book.* Cambridge University Press, 2009.
91. Branco, T. and K. Staras, *The probability of neurotransmitter release: variability and
feedback control at single synapses.* Nat Rev Neurosci, 2009. **10**(5): p. 373–83.

| 1035 | 92. | Markram, H., et al., *Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex.* J Physiol, 1997. **500 ( Pt 2)**(Pt 2): p. 409–40. |
| 1038 | 93. | Chen, X., et al., *Functional mapping of single spines in cortical neurons in vivo.* Nature, 2011. **475**: p. 501–505. |
| 1040 | 94. | Takahashi, N., et al., *Locally Synchronized Synaptic Inputs.* Science, 2012. **335**: p. 353–357. |
| 1042 | 95. | Karimi, A., et al., *Cell-type specific innervation of cortical pyramidal cells at their apical dendrites.* eLife, 2020. **9**: p. e46876. |
| 1044 | 96. | Horner, C.H. and E. Arbuthnott, *Methods of estimation of spine density--are spines evenly distributed throughout the dendritic field?* J Anat, 1991. **177**: p. 179–84. |
| 1046 | 97. | Tjia, M., et al., *Pyramidal Neurons in Different Cortical Layers Exhibit Distinct Dynamics and Plasticity of Apical Dendritic Spines.* Front Neural Circuits, 2017. **11**: p. 43. |
| 1048 | 98. | Davidson, A.M., et al., *Dendritic Spine Density and Dynamics of Layer 5 Pyramidal Neurons of the Primary Motor Cortex Are Elevated With Aging.* Cereb Cortex, 2020. **30**(2): p. 767–777. |
| 1051 | 99. | Markram, H. and A.e. al., *Reconstruction and Simulation of Neocortical Microcircuitry.* Cell, 2015. **163**(2): p. 456–492. |
| 1053 | 100. | Reimann, M.W., et al., *An algorithm to predict the connectome of neural microcircuits.* Frontiers in Computational Neuroscience, 2015. **Volume 9 - 2015**. |
| 1055 | 101. | Varela, J.A., et al., *A quantitative description of short-term plasticity at excitatory synapses in layer 2/3 of rat primary visual cortex.* J Neurosci, 1997. **17**(20): p. 7926–40. |
| 1057 | 102. | Destexhe, A., Z.F. Mainen, and T.J. Sejnowski, *An Efficient Method for Computing Synaptic Conductances Based on a Kinetic Model of Receptor Binding.* Neural Computation, 1994. **6**(1): p. 14–18. |
| 1060 | 103. | Durstewitz, D., J.K. Seamans, and T.J. Sejnowski, *Dopamine-mediated stabilization of delay-period activity in a network model of prefrontal cortex.* J Neurophysiol, 2000. **83**(3): p. 1733–50. |
| 1063 | 104. | Zador, A., C. Koch, and T.H. Brown, *Biophysical model of a Hebbian synapse.* Proc Natl Acad Sci U S A, 1990. **87**(17): p. 6718–22. |
| 1065 | 105. | Freund, T.F. and I. Katona, *Perisomatic inhibition.* Neuron, 2007. **56**(1): p. 33–42. |
| 1066 | 106. | Vinck, M., et al., *The pairwise phase consistency: a bias-free measure of rhythmic neuronal synchronization.* Neuroimage, 2010. **51**(1): p. 112–22. |
| 1068 | 107. | Deuchars, J., D.C. West, and A.M. Thomson, *Relationships between morphology and physiology of pyramid-pyramid single axon connections in rat neocortex in vitro.* J Physiol, 1994. **478 Pt 3**(Pt 3): p. 423–35. |
| 1071 | 108. | Brémaud, A., D.C. West, and A.M. Thomson, *Binomial parameters differ across neocortical layers and with different classes of connections in adult rat and cat neocortex.* Proc Natl Acad Sci U S A, 2007. **104**(35): p. 14134–9. |
| 1074 | 109. | Xiang, Z., J.R. Huguenard, and D.A. Prince, *Synaptic inhibition of pyramidal cells evoked by different interneuronal subtypes in layer v of rat visual cortex.* J Neurophysiol, 2002. **88**(2): p. 740–50. |
| 1077 | 110. | Yu, J., et al., *Recruitment of GABAergic Interneurons in the Barrel Cortex during Active Tactile Behavior.* Neuron, 2019. **104**: p. 412–427.e4. |
| 1079 | 111. | Tanaka, Y.H., et al., *Local connections of layer 5 GABAergic interneurons to corticospinal neurons.* Front Neural Circuits, 2011. **5**: p. 12. |
| 1081 | 112. | Thomson, A.M., J. Deuchars, and D.C. West, *Neocortical local synaptic circuitry revealed with dual intracellular recordings and biocytin-filling.* J Physiol Paris, 1996. **90**(3-4): p. 211–5. |
| 1084 | 113. | Silberberg, G. and H. Markram, *Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells.* Neuron, 2007. **53**(5): p. 735–46. |

## FIGURES

**Figure 1:** A model layer 5 pyramidal neuron with active dendrites. **(A)** The morphology of the neuron. Virtual recordings can be obtained from any desired compartment (colored pipettes). Inset, naturalistic presynaptic activity drives firing rates in our model like those in vivo. Each black cross is the mean rate for a different simulation. **(B)** Examples of membrane potentials recorded simultaneously across the dendritic tree (in color) and soma (black) during naturalistic drive. Regenerative events are indicated with arrows or text (AP: action potential, bAP: backpropagating action potential). **(C1-3)** Demonstration of our detection of dendritic spike events (top) and characterization of their properties (bottom). Events are binned according to the properties that were used in their detection. Bin edges for the event durations were not evenly set for panels **C2** and **C3**.

**Figure 2:** Influence of $Na^+$ and NMDA spikes on action potential generation. **(A)** Electrotonic distance between each dendritic compartment and the soma. **(B)** Dendritic compartments were grouped by their type (apical or basal) and electrotonic distance (percentile) from the soma. The percent change in $Na^+$ spike presence in those compartments relative to somatic spiking. $Na^+$ spikes increased immediately prior to action potentials in dendritic compartments that were electrotonically close to the soma. **(C)** Same format as **B**, but for NMDA spikes. These showed a similar degree of change, but a broader temporal coupling.

**Figure 3:** Influence of $Ca^{2+}$ spikes on action potential generation, and their behavior as a second integrative mechanism. **(A)** Electrotonic distance between dendritic compartments and the apical nexus, where $Ca^{2+}$ spikes are generated. **(B)** Change in the incidence of $Ca^{2+}$ spikes at the nexus surrounding action potentials. **(C)** Percent change in NMDA spike presence in the apical dendrites centered on $Ca^{2+}$ spike initiation. **(D)** Percent change in NMDA spike coupling with action potentials during $Ca^{2+}$ spikes. Top, NMDA spikes in apical dendrites were more strongly coupled with action potentials during $Ca^{2+}$ spikes. Bottom, this was not the case for basal dendrites.

**Figure 4:** Distal dendritic and perisomatic inhibition reduce action potential generation through different mechanisms. **(A)** Action potential rate during periods with normal inhibitory tone (Control), double rate on Distal branches, or double rate on Perisomatic. Both increases in inhibition dramatically reduced the firing of somatic action potentials. **(B)** Somatic excitability was measured by delivering current steps during the control, distal, and perisomatic inhibition states. Left, example somatic voltage responses to current steps. Right, spike frequency versus current (f-I) curve for each condition. The threshold for evoking an action potential shifted with 2X distal and perisomatic inhibition. Perisomatic, but not dendritic, inhibition changed the f-I slope (compare dashed lines with solid black). **(C)** Impact of altered dendritic inhibition on rate of $Na^+$ spikes in apical and basal dendrites. **(D)** Same format as **c**, but for NMDA spikes. **(E)** Rate of $Ca^{2+}$ spikes in the apical dendrites. Basal dendrites lacked $Ca^{2+}$ spikes and were excluded. All error bars are mean ± standard deviation. **(F)** Examples of membrane potential recorded in control (top), and both distal (middle) and proximal (bottom) inhibition lagged by 500 ms. **(G)** Change in firing rate for control (black dot) and for perisomatic (blue) and distal (red) lags in inhibition from 0 to 500 ms. **(H)** Change in incidence of $Ca^{2+}$ spikes for distal (red, top) and proximal (blue, bottom) inhibition. The control case is shown in black in both panels. **(I)** Same as (H) but for NMDA spikes.

**Figure 5:** Phase-dependent effects of beta and gamma rhythmic inhibition on dendritic spikes. **(A)** Example data from the beta rhythmic inhibition simulation. Top, presynaptic spike counts. Bottom, voltage traces from somatic and dendritic compartments. Grayed periods are when inhibitory presynaptic spikes are peaking. **(B)** Action potential rate as a function of the phase of

the beta rhythm. Dashed gray line shows the modulation of inhibitory drive with respect to phase.
**(C)** Percent change in $Ca^{2+}$ spike presence at apical nexus by beta phase. (**D1-2)** Percent change
of NMDA spike presence in apical (1) and basal (2) dendrites stratified by electronic distance from
the soma. **(E1-2)** Same as **C**, but for $Na^+$ spikes. **(F, G, H, I1-2, J1-2)** Same format as above, but
with events binned by the phase of gamma rhythmic inhibition. For all graphs phase is given in
radians with inhibition at a minimum for -π and maximum at 0.
1143
**Figure 5, Supplementary 1:** Phase-dependent effects on dendritic spikes of beta and gamma
rhythmic inhibition delivered to opposite areas of the neuron. Beta was delivered perisomatically,
while gamma was supplied to the distal dendrites. **(A)** Action potential rate as a function of the
phase of the beta rhythm. Dashed gray line shows the modulation of inhibitory drive with respect
to phase. **(B)** Percent change in $Ca^{2+}$ spike presence at apical nexus by beta phase. (**C1-2)**
Percent change of NMDA spike presence in apical (1) and basal (2) dendrites stratified by
electronic distance from the soma. **(D1-2)** Same as **C**, but for $Na^+$ spikes. **(E, F, G1-2, H1-2)** Same
format as above, but with events binned by the phase of gamma rhythmic inhibition. For all graphs
phase is given in radians with inhibition at a minimum for -π and maximum at 0.
1153
**Figure 6:** Phase-dependent effects of gamma and beta rhythmic inhibition on somatic excitability.
**(A1)** A cumulative probability plot of the distribution of somatic membrane potentials 1ms prior to
an action potential, sorted by whether they occurred during the gamma phase with maximal
(Peak, red line) or minimal (Trough, blue line) inhibitory drive. Poisson (black) had no rhythmic
modulation, but the same mean inhibitory rate. **(A2)** Probability distribution of somatic membrane
voltage as a function gamma phase, normalized to the peak probability value. Lines have the
same color scheme as in **A1**. **(B1)** Same format as **A1**, but for the beta rhythm. **(B2)** Same format
as **A2**, but for the beta rhythm.
1162
**Figure 6, Supplementary 1:** Phase-dependent effects on somatic excitability of beta and gamma
rhythmic inhibition delivered to opposite areas of the neuron. Beta was delivered perisomatically,
while gamma was supplied to the distal dendrites. **(A1)** A cumulative probability plot of the
distribution of somatic membrane potentials 1ms prior to an action potential, sorted by whether
they occurred during the gamma phase with maximal (Peak, red line) or minimal (Trough, blue
line) inhibitory drive. Poisson (black) had no rhythmic modulation, but the same mean inhibitory
rate. **(A2)** Probability distribution of somatic membrane voltage as a function gamma phase,
normalized to the peak probability value. Lines have the same color scheme as in **A1**. **(B1)** Same
format as **A1**, but for the beta rhythm. **(B2)** Same format as **A2**, but for the beta rhythm.
1172
**Figure 7:** Frequency- and phase-dependent effects of inhibitory rhythms on the distal dendrites.
**(A)** Entrainment to an inhibitory rhythm delivered to the distal dendrites varied with its
frequency. Higher frequencies were less able to entrain dendritic spikes. Entrainment tended to
be strongest for apical (blue) over basal (orange) compartments. **(B)** Example voltage traces
from dendritic compartments in either the distal basal or apical branches. Gray shading denotes
period where the inhibitory rhythm troughs occurred. For the basal segment, NMDA spikes were
shaded in purple, while in the apical segment $Ca^{2+}$ spikes were shaded in green. Dendritic spike
onsets denote with red lines, and offsets with blue lines. **(C)** Percent change from the mean in
the rate of dendritic spike onsets (red gradient) and offsets (blue gradient) as a function of
rhythm frequency and phase. Purple regions denote phase/frequency combinations where both
onsets and offsets were elevated, while regions with either just blue or red indicate that offsets
or onsets preferentially occurred, respectively. We did not determine an offset for Na+ spikes
due to their transience (~ 1 ms).
1186

**Figure 8:** Frequency- and phase-dependent effects of inhibitory rhythms on the perisomatic region. **(A)** The mean somatic membrane potential during either the trough or peak phase of the inhibitory rhythm. **(B)** Mean of the distribution of somatic membrane potential fluctuations as a function rhythm phase and frequency. Fluctuations were measured across the entire simulation time as the difference in membrane potential at 1 ms delays. For both graphs red lines are peaks and blue lines are troughs.

**Figure 9:** Phase-dependent effects of gamma and beta bursts on dendritic spikes. **(A)** Example data from the gamma rhythmic inhibition simulation. Top, Somatic potential (black line), with firing rate of perisomatic inhibitory synapses (blue line). Middle, voltage trace of apical compartment. Bottom, voltage trace of basal compartment. **(B)** Action potential rate as a function of the phase of the gamma rhythm. Blue line shows the modulation of inhibitory drive with respect to phase. **(C)** Percent change in $Ca^{2+}$ spike presence at apical nexus by gamma phase. (**D1-2**) Percent change of NMDA spike presence in apical (1) and basal (2) dendrites stratified by electronic distance from the soma. **(E1-2)** Same as **C**, but for $Na^+$ spikes. **(F, G, H, I1-2, J1-2)** Same format as above, but with events binned by the phase of beta rhythmic inhibition. For all graphs cycle number is given relative to the amplitude peak of the burst.

**Figure 10:** Effect of beta and gamma rhythms on responsiveness to synaptic inputs targeting distinct regions of the dendritic tree. **(A)** Schematic of the location for clustered excitatory synaptic inputs. **(B)** Normalized cross-correlation between synaptic drive onto a clustered input and spiking at the soma, stratified by whether the presynaptic spike arrived during the Peak (red line) or Trough (blue line) of the rhythm. Solid bars correspond to the Poisson stimulation case where inhibition was not rhythmically modulated. Top left, effect of beta on distal inputs. Top right, effect of gamma on distal inputs. Bottom left, effect of beta on proximal inputs. Bottom right, effect of gamma on proximal inputs. **(C)** Summary of effects in panel **B** where the strength of each normalized cross-correlation was measured as its area under the curve. Dots are connected by dashed gray lines if the data points came from the same simulation but at different phases of the rhythm. Solid horizontal lines reflect the cross-correlation strength in the Poisson inhibitory case (no rhythmicity).

**Figure 11:** A summary schematic of the principal findings. **(A)** The microcircuitry that was simulated in this study. **(B)** Beta rhythmic inhibition to the distal dendrites modulated dendritic spikes. **(C)** Gamma rhythmic inhibition to the perisomatic region modulated action potential initiation. AP stands for action potentials.

**A** Morphology of Layer 5 neuron

**B** Membrane potentials recorded throughout the neuron during naturalistic synaptic drive exhibit regenerative events

Apical tuft

Apical nexus

Apical trunk

Basal dendrites

Spontaneous firing rates

bAP

AP

▼ NMDA spike
▼ Na+ spike
▼ Ca2+ spike

**C1** Na+ spike example

Soma voltage
Dendrite voltage
NaP current

Na+ spike properties

Conductance (mS/cm²)

Duration (ms)

**C2** NMDA spike example

Soma voltage
Dendrite voltage
NMDA current

NMDA spike properties

Charge (nA ms)

Duration (ms)

**C3** Ca2+ spike example

Soma voltage
Dendrite voltage
HVA current
LVA current

Ca2+ spike properties

Charge (nA ms)

Duration (ms)

Percentage of events

**A** Electrotonic distance across the dendritic tree

Percent of signal at soma

**B** Na⁺ spikes on apical branches

Na⁺ spikes on basal branches

Electrotonic distance from soma (percentile rank)

Time from action potential (ms)

Percent change

**C** NMDA spikes on apical branches

NMDA spikes on basal branches

Time from action potential (ms)

Percent change

**A** Electrotonic distance from apical nexus

100μm
100μm

Percent of signal at nexus
100
10
1

**B** Ca²⁺ spikes on apical dendrites

Percent change

Time from action potential (ms)

**C** NMDA spikes on apical branches

Electrotonic distance from nexus (percentile rank)
Far
Mid
Close

Time from Ca²⁺ spike at nexus (ms)

Percent change

**D** Modulation of NMDA spike/action potential coordination by Ca²⁺ spikes

NMDA spikes on apical branches

Electrotonic distance from soma (percentile rank)
Far
Mid
Nexus
Close

Time from action potential (ms)

Percent change
>500
0
<−500

NMDA spikes on basal branches

Electrotonic distance from soma (percentile rank)
Far
Mid
Close

Time from action potential (ms)

Percent change
>500
0
<−500

**A** Inhibitory modulation of action potentials

**B** f-I curve to characterize response gain

**C** Inhibitory modulation of Na⁺ spikes

**D** Inhibitory modulation of NMDA spikes

**E** Inhibitory modulation of Ca²⁺ spikes

**F** Voltage traces with different E/I lags

**G** Firing rate vs. E/I lag

**H** Ca²⁺ spikes vs. E/I lag

**I** NMDA spikes vs. E/I lag

**A** Modulation by beta rhythmic inhibition

Perisomatic presynaptic spikes

Distal dendritic presynaptic spikes

Membrane voltage traces

**B** Action potentials

**C** Ca²⁺ spikes

**D1** NMDA Apical

**D2** NMDA Basal

**E1** Na⁺ Apical

**E2** Na⁺ Basal

**F** Modulation by gamma rhythmic inhibition

Perisomatic presynaptic spikes

Distal dendritic presynaptic spikes

Membrane voltage traces

**G** Action potentials

**H** Ca²⁺ spikes

**I1** NMDA Apical

**I2** NMDA Basal

**J1** Na⁺ Apical

**J2** Na⁺ Basal

# Modulation by perisomatic beta rhythmic inhibition

**A** Action potentials

**B** Ca²⁺ spikes

**C1** NMDA Apical

**C2** NMDA Basal

**D1** Na⁺ Apical

**D2** Na⁺ Basal

# Modulation by distal dendritic gamma rhythmic inhibition

**E** Action potentials

**F** Ca²⁺ spikes

**G1** NMDA Apical

**G2** NMDA Basal

**H1** Na⁺ Apical

**H2** Na⁺ Basal

**A** Variation of dendritic spike entrainment with the frequency of rhythmic inhibition

Ca²⁺ spikes | NMDA spikes | Na⁺ spikes

Entrainment (PPC)

Rhythm frequency

**B** Membrane voltage traces

Basal -39 mv

Trough · Peak · T · P · T · P · T · P · T · P · T · P · T · P · T · P · T · P · T

NMDA spikes

50 mV

Ca²⁺ spikes
on · off

Apical -73 mv

100 ms

**C** Phase/frequency-dependent modulation of dendritic spike onsets and offsets

Ca²⁺ spikes | NMDA spikes | Na⁺ spikes

Phase of inhibition (radians)

Inhibition modulation

Rhythm frequency

Offset (% change)
Onset (% change)

**A** Phase-dependent modulation of mean membrane voltage

**B** Phase-dependent modulation of changes in membrane voltage

**A** Modulation by gamma rhythmic inhibition

Inhibitory mod.
Soma
Apical
Basal

**F** Modulation by beta rhythmic inhibition

Inhibitory mod.
Soma
Apical
Basal

50 mV

200 ms

**B** Action potentials

Firing rate (Hz)

Burst cycle

**C** Ca²⁺ spikes

Percent change

Burst cycle

**G** Action potentials

Firing rate (Hz)

Burst cycle

**H** Ca²⁺ spikes

Percent change

Burst cycle

**D1** NMDA Apical

Electrotonic distance from soma

Burst cycle

**E1** Na⁺ Apical

Burst cycle

**D2** NMDA Basal

Burst cycle

**E2** Na⁺ Basal

Burst cycle

**I1** NMDA Apical

Electrotonic distance from soma

Burst cycle

**J1** Na⁺ Apical

Burst cycle

**I2** NMDA Basal

Burst cycle

**J2** Na⁺ Basal

Burst cycle

Percent change
−100  0  100

Percent change
−100  0  100

Percent change
−100  0  100

Percent change
−100  0  100

**A** Positioning of clustered excitatory synapses

Distal excitatory inputs

100um

100um

Proximal excitatory inputs

**B** Effect of rhythmic inhibition phase on the cross-correlation between afferent excitation and somatic action potentials

Beta rhythmic inhibition

Gamma rhythmic inhibition

Distal excitation

Peak
Trough
Poisson

Cross-correlation (normalized × 10²)

Proximal excitation

Peak
Trough
Poisson

Time from presynaptic spike (ms)

**C** Response strength by input location, inhibition type, and phase

Peak
Trough
Proximal poisson
Distal poisson

Area under cross-correlation peak

16 Hz    64 Hz    16 Hz    64 Hz
Proximal         Distal

**A** Microcircuitry of beta/gamma

Beta rhythm (15-35 Hz)

SOM

PN

PV

Gamma rhythm (>35 Hz)

**B** Beta modulation of dendritic spikes

SOM

NMDA

Ca²⁺

AP

Na⁺

| P | T | P | T | P | Peak/trough |

**C** Gamma modulation of soma responsiveness

PV

NMDA

Ca²⁺

AP

Na⁺

PTPTPTPTPTPTPTPT Peak/trough