Intestinal barrier function in the naked mole-rat: an emergent model for gastrointestinal insights

Javier Aguilera-Lizarraga[1]*, Anne Ritoux[1], David C. Bulmer[1], Ewan St. John Smith[1]*

[1] Department of Pharmacology, University of Cambridge, Cambridge, United Kingdom

* To whom correspondence should be addressed: Dr. Javier Aguilera-Lizarraga: ja783@cam.ac.uk; Prof. Ewan St. John Smith: es336@cam.ac.uk.

Abstract:

The intestinal barrier plays a crucial role in homeostasis, both by facilitating absorption of nutrients and fluids, and providing a tight shield to prevent the invasion by either pathogen or commensal microorganisms. Intestinal barrier malfunction is associated with systemic inflammation, oxidative stress, and decreased insulin sensitivity, which may lead to the dysregulation of other tissues. Therefore, a deeper understanding of physiological aspects related to an enhanced barrier function is of significant scientific and clinical relevance. The naked mole-rat has many unusual biological features, including attenuated colonic neuron sensitivity to acid and bradykinin, and resistance to chemical-induced intestinal damage. However, insight into their intestinal barrier physiology is scarce. Here, we observed notable macroscopic and microscopic differences in intestinal tissue structure between naked mole-rats and mice. Moreover, naked mole-rats showed increased number of larger goblet cells and elevated mucus content. In measuring gut permeability, naked mole-rats showed reduced permeability compared to mice, measured as transepithelial electrical resistance, especially in ileum. Furthermore, intestinal ion secretion induced by serotonin, bradykinin, histamine, and capsaicin was significantly reduced in naked mole-rats compared to mice, despite the expression of receptors for all these agonists. In addition, naked mole-rats exhibited reduced pro-secretory responses to the non-selective adenylate cyclase activator forskolin. Collectively, these findings indicate that naked mole-rats possess a robust and hard-to-penetrate gastrointestinal barrier, that is resistant to environmental and endogenous irritants. Naked mole-rats may therefore provide valuable insights into the physiology of the intestinal barrier and set the stage for the development of innovative and effective therapies.

Keywords: gut; intestinal function; naked mole-rat; intestinal permeability; mucosal ion secretion.

35 **Graphical abstract**:

Naked mole-rats exhibit:

- A robust gut tissue (thicker intestinal layers, longer villi, bigger crypts), despite shorter small intestine length.
- More and larger goblet cells, with increased mucus granule content.
- Remarkable intestinal tissue integrity ("pore" pathway), especially in the ileum.
- Reduced intestinal ion secretion in response to serotonin, bradykinin, histamine, capsaicin (TRPV1 agonist), and forskolin (adenylate cyclase activator).

1. Introduction:

Digestive pathologies stand out as the most common type of disease worldwide (*1*). Importantly, they result in a significant reduction in the quality of life for patients and represent a multi-billion healthcare burden globally (*2–5*). Intestinal barrier dysfunction is a common feature of most gastrointestinal diseases, including coeliac disease, inflammatory bowel disease, metabolic dysfunction-associated liver disease and disorders of gut-brain interaction (*6*). Moreover, numerous pathologies unrelated to the gastrointestinal tract, such as respiratory diseases (*7*), skin conditions (*8*) and neurodegenerative diseases (*9*), have also been associated with impaired intestinal barrier function. The common denominator of intestinal barrier perturbation led to the "leaky gut" hypothesis (*10*), which states that enhanced intestinal permeability, caused by endogenous or exogenous factors, may increase the passage of food antigens, commensal microorganisms and/or pathogens into the underlying tissue and the systemic circulation, leading to aberrant immune activation described in different disease conditions. Although increased permeability may not necessarily be deleterious (*11*), and whether intestinal barrier dysregulation is the cause or consequence in most disorders is unknown, it is broadly accepted that adequate functioning of the intestinal barrier is required to maintain gut and inter-organ homeostasis. Consequently, identifying mechanisms leading to enhanced intestinal barrier function is highly relevant for both basic and clinical sciences.

In addition to the use of human intestinal mucosal biopsies (*12*), many models have been employed to investigate the (patho)physiology of the intestinal barrier. Among these, human gut-derived cell lines (*13*), such as Caco-2 cell monolayers (*14*), and animal models, primarily rodents, but also porcine, equine and avian models (*15*), have been most commonly utilised. Furthermore, human organoids have recently been implemented to model intestinal epithelial barrier function (*16, 17*). Despite important advances in our understanding of intestinal barrier function, there has been little to no reduction in the global incidence and prevalence of gastrointestinal conditions over the last 30+ years (*1*). There is thus an imperative need for further research, including the development of novel models, to enhance understanding of intestinal barrier function.

In recent years, naked mole-rats (NMR, *Heterocephalus glaber*) have attracted considerable attention from biomedical scientists as a non-traditional animal model due to their highly unusual biological features (*18*). Notably, these animals show an outstanding tolerance to environmental stressors associated with human diseases. Indeed, these eusocial rodents, which have an unusually long lifespan given their 40-80 g size (>35 years) (*19*), have been used for investigating 'healthy' aging (*20*), cancer resistance (*21*), and protection against neurodegeneration (*22*). However, the gastrointestinal tract of these animals remains poorly characterised. Of interest, a recent study revealed that the intestinal barrier of NMRs is resistant to tissue damage induced by dextran sodium sulphate (DSS) (*23*), a widely used colitogenic chemical for modelling inflammatory bowel disease. Although the authors employed a modified version of the traditional DSS protocol (*24*), NMRs displayed no signs of histological damage, even when exposed to 8.75% DSS, albeit for only 3-days. Although several factors may be involved in the increased resistance to DSS observed in these animals, such as a thicker mucus layer (*23*), differences in the gut microbiota (*25*), a distinct

immune system (*26*), etc., this observation suggests that the gut physiology of NMRs may
possess characteristics of potential biological interest. We therefore hypothesised that NMRs
would display enhanced intestinal function, that is, improved permeability regulation and
resistance to intestinal irritants. To answer this question, we explored the macroscopic and
microscopic features of the NMR gut, intestinal permeability, changes in mucosal ion
secretion, and expression of intestinal markers of biological interest. Investigating the
intestinal barrier function in NMRs may enable a better understanding of the underlying
mechanisms that contribute to better mucosal protection.

2. **Material and Methods**:

2.1. Animals

All animal experiments were performed in accordance with the Animals (Scientific
Procedures) Act 1986 Amendment Regulations 2012 under Project Licenses granted to
E.St.J.S. (P7EBFC1B1 and PP5814995) by the Home Office and approved by the University
of Cambridge Animal Welfare Ethical Review Body. Experiments were performed using a
mixture of male and female C57BL6/J mice (10-15 weeks old; IQR 10-12) and a mixture of
male and female, non-breeder, NMRs (23-145 weeks; IQR 48-54). Mice were purchased
from Envigo and housed conventionally with nesting material and a red plastic shelter in a
temperature-controlled room at 21 °C, with a 12-hour light/dark cycle and access to food and
water *ad libitum*. NMRs were bred in-house and maintained in an interconnected network of
cages in a humidified (~55%) temperature-controlled room at 28-32 °C with red lighting
(08:00−16:00). NMRs had access to food *ad libitum*. In addition, a heat cable provided extra
warmth under two to three cages/colony. NMRs used in this study came from five different
colonies. Animals were humanely killed by CO_2 exposure followed by cervical dislocation
(mice) or decapitation (NMRs).

2.2. Intestinal tissue processing for histology and histomorphometric assessment

Intestinal biopsies (ileum and colon) were fixed and embedded using conventional
techniques. In brief, samples were incubated, after dissection, in 4% paraformaldehyde
(PFA) at 4 °C overnight. Next, samples were washed in PBS, transferred to sucrose (0.3 g ·
ml^{-1}) and stored at 4 °C overnight. Finally, samples were embedded in Shandon M-1
Embedding Matrix (Thermo Fisher Scientific), snap-frozen and stored at −80°C until
processing. Cryostat (Leica) sections (12 µm) were collated across Superfrost Plus slides
(Thermo Fisher Scientific) and stained with haematoxylin and eosin (H&E) using
conventional techniques (*27, 28*). Villi length and width, crypts length and width, and
thickness of the different layers (submucosa and muscularis) were measured in ileum and
colon samples (Supplementary Figure 1a, b) using ImageJ software. For analyses of villi and
crypts, only intestinal units that were overtly complete and full-sized, and showed no signs of
bending or mechanical damage, were chosen in each sample. A total of 5-10 units of each
kind were carefully selected, and their average calculated.

2.3. Alcian blue/periodic acid-Schiff (AB/PAS) staining

For goblet cell identification and quantification of their mucus content, 12-µm tissue sections
were subjected to Alcian Blue and Periodic Acid Schiff's (AB/PAS) staining. In brief, sections
were hydrated in double distilled water and then stained with Alcian Blue solution (Sigma,
1% w/v Alcian Blue in 3% v/v aqueous acetic acid) for 15 minutes. Next, sections were
rinsed in double distilled water before being treated for 5 minutes with 0.5% periodic acid
(Sigma). Thereafter, tissue sections were washed again in double distilled water and stained
for 10 minutes in Schiff's reagent (Sigma). Finally, sections were washed for 5 minutes in tap
water, mounted, and imaged using a NanoZoomer S360 (Hamamatsu).

For analyses, images were loaded into the QuPath software (29). In each image, regions of
interest (ROI) for crypts and villi were manually annotated. Within these, the cell detection
tool was used to segment individual cells using a manually selected threshold. To ensure
that the segmentation correctly circled cells, ROIs were visually inspected. After colour
deconvolution, measurements for average pixel intensity across each cell ROI and cell area
were analysed.

2.4. Ussing chambers

Post-mortem, for both species, the gastrointestinal tract (from pyloric sphincter to anus) was
carefully removed, cleaned of faecal pellets, and maintained in ice-cold Krebs solution. For
experiments where transepithelial electrical resistance (*TEER*) was measured, full-thickness
segments of ileum and colon were mounted in Ussing chambers without removal of the
muscularis propria or the serosa. For experiments where short-circuit current (I_{sc}) was
measured, ileum samples were pinned in Sylgard plates with the lamina propria facing down
and muscularis externa and serosal layers gently peeled off with forceps to prepare lamina
propria-submucosal plexus intestinal strips. Multichannel DVC 1000 Voltage/Current clamp
apparatus (World Precision Instruments) was used to measure changes in intestinal tissue
function. Each channel was connected to a DVC-3 preamplifier (World Precision
Instruments), connected to two current and two voltage electrodes filled with 1.5% (w/v) agar
(Millipore) dissolved in 3M KCl. The voltage and fluid resistance were zeroed to eliminate
electrical bias and the voltage clamp circuitry was set to standby mode and the intestinal
preparation was mounted in the chamber with fresh superfusate. After calibration, intestinal
tissues were then mounted in Ussing chambers and the mucosal and serosal compartments
were bathed independently with 10 mL of Krebs-Ringer bicarbonate buffer (pH 7.4)
containing (in mM): 118 NaCl; 4.7 KCl; 1.2 $CaCl_2$; 1.2 $MgSO_4$; 1.2 NaH_2PO_4; 25 $NaHCO_3$).
Glucose (10 mM) was present in the medium that bathed the serosal surface of the tissue
and mannitol (10 mM) was substituted for glucose in the medium that bathed the mucosal
surface. Media were oxygenated (95% O_2, 5% CO_2) throughout the experiment via a gas-
uplift recirculation system and maintained at 37°C by water-jacketed reservoirs. Although
NMRs are cold blooded and maintained at 28-32 °C, to minimise the number of variables
impacting our results, all experiments were conducted with media maintained at 37 °C. For
mouse tissue, Ussing chambers with an exposed area of 0.071 cm^2 were used, while an
exposed area of 0.196 cm^2 was used for NMR tissue.

Transepithelial electrical resistance (TEER) experiments: The transepithelial electrical
resistance was calculated according to Ohm's law (resistance = voltage/current,
normalised by the exposed window area; TEER = $R_{tissue}(\Omega) \cdot$ Area (mm^2)) from the voltage

deflections induced by bipolar current pulses of 50 µA every 50 s with a duration of
500 ms. After 30 min of tissue stabilisation, TEER was measured over a period of 2 h.

Short-circuit current (I_{sc}) experiments: After mounting, tissue was allowed to equilibrate for
15-20 min. Thereafter, preparations were short-circuited (voltage-clamped at zero potential)
and the short-circuit current (I_{sc}) required to maintain the 0 mV potential was monitored over
time. After short-circuiting, intestinal tissue was allowed to equilibrate for 10-15 min to allow
bioelectric stabilisation (baseline). All I_{sc} responses were normalised to µA · cm^{-2} and
changes in I_{sc} (ΔI_{sc}) were recorded after application of compounds to the serosal side, except
forskolin which was added to both the apical and the serosal sides.

2.5. Chemicals

178 The following drugs were used for short-circuit current (I_{sc}) experiments: capsaicin (Sigma-
179 Aldrich), bradykinin (Tocris), 5-hydroxytryptamine (5-HT, serotonin) (Sigma-Aldrich),
180 histamine (Sigma-Aldrich) and forskolin (Tocris). Capsaicin (1 mM) was dissolved in 100%
181 ethanol, forskolin was dissolved in dimethyl sulfoxide (DMSO, 10 mM), and bradykinin (1
182 mM), 5-HT (10 mM) and histamine (10 mM) were dissolved in double distilled water.

2.6. RNA extraction from intestinal tissue and cDNA preparation

185 Following dissection and cleaning, intestinal samples (ileum and colon) were stored in
186 RNAprotect Tissue Reagent (Qiagen) at 4°C until processing. Next, total RNA was isolated
187 using the RNeasy Mini Kit (Qiagen) following the manufacturer's instructions and including a
188 DNase digestion step. Subsequently, cDNA was transcribed using qScript cDNA Synthesis
189 Kit (Quantabio) following the manufacturer's instructions.

2.7. Real-time quantitative PCR (RT-qPCR):

Reactions were prepared with FastStart Sybr Green Master (Merk) following the
manufacturer's instructions. In brief, 10 µL reaction mix per well, containing cDNA (20 ng)
oligonucleotides (200 nM) and UltraPure distilled water (DNase/RNase free; Invitrogen),
were loaded onto 96-well plates. Thermocycling was performed using a StepOnePlus^{TM}
Real-Time PCR System (Thermo Fisher). A melt curve was performed at the end of
thermocycling to assess the specificity of primers. Single peaks were obtained for all primer
pairings.

To determine primer efficiency, a standard curve was generated for each pair of primers at
cDNA concentrations of 50, 5, 0.5, 0.05 and 0.005 ng/µL and plotting C_t against log
concentration. Efficiency was calculated by the following equation: Efficiency = $10^{-(1/slope)}$-1
(*30, 31*). Efficiency of reaction values between 90 and 110% were considered acceptable.

Gene expression was normalised to the endogenous reference gene *Tbp* for both species.
This gene encodes the TATA-binding protein. *Tbp* was chosen as it has been proposed as
an ideal reference gene for the normalisation of gene expression in intestinal samples of
mice and other animals (*32–36*). Relative gene expression was calculated as $2^{-\Delta\Delta Ct}$ (*37*). The
primer pairs used are described in Supplementary Table 1.

2.8. Measurement of Substance P from intestinal explants

Post-mortem, the small intestine and colon were carefully removed, cleaned of faecal pellets and maintained in ice-cold Krebs solution. Next, 30-50 mg tissue sections (terminal ileum and distal colon) were dissected and transferred into 24-well plates containing 500 µL per well of L-15 Medium (GlutaMAX) medium (Gibco). Explants from both species were incubated for 5 min at 32 °C (5% CO_2, 3%O_2). Thereafter, culture supernatant was collected and stored at -80˚C until use. Substance P was measured using an enzyme-linked immunosorbent assay (Substance P ELISA Kit, Cayman Chemicals) according to the manufacturer's instructions. Substance P levels were normalised to protein content upon quantification using the Bradford Assay (Thermo Fisher) following the manufacturer's instructions.

2.9. Statistical analyses:

Tissue was taken from non-manipulated healthy animals being used for other experiments, thus meeting the Reduction aim of the 3Rs. While there were no upfront power calculations to determine the sample size, we included sample sizes commonly used for the types of assays used in this study (*38, 39*).

Normal (Gaussian) distribution was determined for all datasets using the Shapiro–Wilk normality test. Non-parametric tests were used for datasets where ≥1 group did not pass the normality test (assuming α = 0.05). Otherwise, parametric tests were used. The type of statistical test and the sample sizes for each experiment are provided in figure legends. Data were plotted and statistical analyses performed with Prism (GraphPad Software version 10.2.1). A *p*-value <0.05 was considered statistically significant. Finally, effect sizes were calculated post *hoc* using the statistical software G*Power (Version 3.1.9.6).

For histological assessment of mouse and NMR samples, 2-4 sections per animal, separated by at least 150 µm, were randomly captured, and analysed by an experimenter in a blinded manner. The experimenter was unblinded for statistical analysis.

3. Results:

3.1. Macroscopic features of the naked mole-rat gastrointestinal tract.

NMRs are small rodents whose body mass is significantly larger compared to mice (data from this study: mouse 21.05 ± 1.71 g vs. NMR 47.07 ± 12.03 g, $p < 0.0001$, Figure 1a), yet substantially smaller than typically used laboratory adult rats (Sprague-Dawley, ∼300-400 g; Wistar rats, ∼250-500 g) (*40, 41*) or guinea pigs (Hartley, ∼625-1,200 g) (*42*). Despite the larger size of NMRs compared to mice, NMRs had a noticeably shorter gut length, measured from the pyloric sphincter to the anus (mouse 40.09 ± 2.35 cm vs. NMR 22.17 ± 1.94 cm, $p < 0.0001$; Figure 1b, c). Further analyses revealed that these differences were driven by a shorter small intestine in NMR than that of mice (small intestine: mouse 32.69 ± 1.92 cm vs. NMR 14.95 ± 1.88 cm, $p < 0.0001$; colon: mouse 7.40 ± 0.65 cm vs. NMR 7.28 ± 0.59 cm, $p = 0.6425$; Figure 1d, e), in keeping with previous studies (*23, 43*). Of interest, while the ratio of the small intestine/colon length in mice was comparable to that in humans (*44−46*), this was reduced by ∼2-fold in NMRs (human 3.956 ± 1.36 vs. mouse 4.52 ± 0.99 vs. NMR 2.07

± 1.20, *p* < 0.0001; Figure 1f). The lengths of the NMR small intestine and colon were
comparable between male and female (small intestine: male 15.50 ± 6.60 cm vs. female
15.05 ± 2.50 cm, *p* = 0.5626; colon: male 7.10 ± 2.00 cm vs. female 7.55 ± 0.20 cm, *p* =
0.2265; Supplementary Figure 1c, d). Given the considerable age range of the NMRs used
in our study (minimum age: 23 weeks old; maximum age: 145 weeks old) and their
attainment of adulthood at ~1 year old, we analysed the relationship between gut length and
both age and weight. The colon length was not associated with the age or weight of the
NMR (age vs. length: r = 0.0581, p = 0.8248, weight vs. length: r = 0.0245, p = 0.9339;
Supplementary Figure 1e, f). However, we did find statistically significant correlations
between the small intestine length and the age and weight of NMRs (age vs. length: r =
0.7293, p = 0.0012, weight vs. length: r = 0.6258, p = 0.0282; Supplementary Figure 1g, h).
These findings parallel human anatomical data, that is, the length of the colon in
young/adolescent individuals is similar to that in adults (*47*), while the small intestine only
reaches its maximum length in adulthood (*48*). Overall, these results show gross anatomical
differences in the gut of NMRs compared to humans and mice, especially regarding the
short length of the small intestine.

Figure 1. Anatomical features of gastrointestinal tracts of NMRs compared to mice. Weight of mice (n = 20)
compared to NMR (n = 25) (**a**). **b**, Representative pictures of the gastrointestinal tract (from stomach to anus) of
NMR (upper panel) and mouse (lower panel). Gastrointestinal length measured from the pyloric sphincter to the
anus (**c**), small intestine (**d**) and colon (**e**) in mice (n= 8) and NMRs (n = 17). **f)** Length ratio of the small intestine
compared to the colon in humans (data extracted from *(44–46)*), mice and NMR. Human* data reflect median of
the means provided by 3 studies that analysed 100, 100 and 200 individuals, respectively. *P* values are shown in
plots. **c**, **d** and **e**, unpaired t-test (data shown as mean ± SD); **f**, Kruskal-Wallis test followed by Dunn's multiple
comparisons test (data shown as median ± IQR). The NMR and mouse illustrations in **a** were created with
BioRender.com).

3.2. Intestinal histology in naked mole-rats

Next, we explored the microscopic architecture of the small intestine (ileum) and colon in NMRs, comparing them to those of mouse tissue (Figure 2a-d). To this end, we measured the histological properties of the intestinal villi and crypts, as well as the thickness of different layers. The ileum of NMRs showed thicker muscularis (mouse 37.56 ± 8.24 µm vs. NMR 60.45 ± 23.43 µm, p = 0.0055; Figure 2e) and submucosal layers compared to mice (mouse 18.90 ± 2.02 µm vs. NMR 40.46 ± 11.22 µm, p < 0.0001; Figure 2f). Furthermore, NMR ileal villi were substantially longer (mouse 344.4 ± 180.8 µm vs. NMR 801.0 ± 394.3 µm, p < 0.0001; Figure 2g) and wider (mouse 90.95 ± 10.00 µm vs. NMR 135.3 ± 36.63 µm, p = 0.0008; Figure 2h) compared to those of mice, and NMR ileal crypts were significantly deeper (mouse -135.60 ± 66.81 µm vs. NMR -157.10 ± 47.49 µm, p = 0.0039; Figure 2i) and wider (mouse 35.72 ± 6.45 µm vs. NMR 57.86 ± 10.39 µm, p < 0.0001; Figure 2j). In the colon, the muscularis was comparable in both species (mouse 128.2 ± 78.31 µm vs. NMR 139.3 ± 83.82 µm, p = 0.2188; Figure 2k), but the colonic submucosa was thicker in NMRs (mouse 63.13 ± 10.18 µm vs. NMR 80.27 ± 10.26 µm, p = 0.0012; Figure 2l). Finally, colonic crypts were similarly deep in both species (mouse -219.0 ± 32.57 µm vs. NMR -198.0 ± 12.64 µm, p = 0.0837; Figure 2m), but those of NMRs were significantly wider compared to mice (mouse 34.07 ± 3.35 µm vs. NMR 56.79 ± 7.24 µm, p < 0.0001; Figure 2n).

Figure 2. Intestinal histology in NMRs. Representative images of H&E staining sections from mouse (**a**) and NMR (**b**) ileum, and mouse (**c**) and NMR (**d**) colon. Ileum muscularis (**e**) and submucosal (**f**) layers thickness in mouse (n = 11, 2-3 sections/animal, 4 mice) and NMR (n = 13, 3-4 sections/animal, 4 NMR). **g**, average villi height (5-10 villi/sample) in mice (n = 11, 2-3 sections/animal, 4 mice) and NMRs (n = 12). **h**, average villi width (5-10 villi/sample) in mice (n = 11, 2-3 sections/animal, 4 mice) and NMRs (n = 13, 3-4 sections/animal, 4 NMR). **i**, average crypt depth (5-10 crypts/sample) in mice (n = 11, 2-3 sections/animal, 4 mice) and NMRs (n = 13, 3-4 sections/animal, 4 NMR). **j**, average crypt width (5-10 crypts/sample) in mice (n = 11) and NMRs (n = 13). Colon muscularis (**k**) and submucosal (**l**) layers thickness in mouse (n = 12, 3 sections/animal, 4 mice) and NMR (n = 9, 3 sections/animal, 3 NMRs). **m**, average crypt depth (5-10 crypts/sample) in mice (n = 12, 3 sections/animal, 4 mice) and NMRs (n = 9, 3 sections/animal, 3 NMRs). **n**, average crypt width (5-10 crypts/sample) in mice (n = 12, 3 sections/animal, 4 mice) and NMRs (n = 9, 3 sections/animal, 3 NMRs). *P* values are shown in plots. **e**, **f**, **h**, **j**, **l**, **m** and **n**, unpaired t-test (data shown as mean ± SD); **g**, **i** and **k**, Mann-Whitney test (data shown as median ± IQR).

We next used Alcian Blue/Periodic Acid-Schiff (AB/PAS) staining to identify goblet cells and quantify mucus granule content in small intestine and colon samples. Ileum samples (Figure 3a, b) of NMRs showed similar numbers of AB$^+$ mucin-producing goblet cells in the villi compared to mice (mouse 80.07 ± 23.25 cells/villus vs. NMR 93.07 ± 20.41 cells/villus, *p* = 0.2254; Figure 3c), and slightly higher numbers in the crypts (mouse 28.66 ± 7.97 cells/crypt vs. NMR 38.98 ± 8.86 cells/crypt, *p* = 0.0194; Figure 3c). However, the size of the goblet

cells present in the NMR ileum was significantly larger than that of mouse ileum goblet cells,
both in the villi and crypts (mouse villi 33.91 ± 3.42 µm^2 vs. NMR villi 41.79 ± 7.62 µm^2, $p =$
0.012; mouse crypts 29.25 ± 3.71 µm^2 vs. NMR crypts 46.53 ± 5.79 µm^2, $p < 0.0001$; Figure
3d). In addition, the goblet cell-mucus granule content (measured as AB/PAS staining
intensity per cell) in NMR ileum, in both villi- and crypts-residing cells, was significantly
higher compared to that of mice (mouse villi 0.51 ± 0.04 a.u. vs. NMR villi 0.74 ± 0.10 a.u., p
< 0.0001; mouse crypts 0.48 ± 0.06 a.u. vs. NMR crypts 0.89 ± 0.12 a.u., $p < 0.0001$; Figure
3e). In the colon (Figure 3f, g), the number of AB$^+$ goblet cells was significantly elevated in
the crypts of NMRs compared to in mice (mouse 14.07 ± 1.51 cells vs. NMR 23.97 ± 4.54
cells, $p < 0.0001$; Figure 3h). Furthermore, as in the ileum, the size of these goblet cells was
significantly bigger in NMRs compared to mice (mouse crypts 65.98 ± 14.77 µm^2 vs. NMR
crypts 170.2 ± 21.43 µm^2, $p < 0.0001$; Figure 3d). Finally, again, in keeping with the results
obtained in the ileum, the mucus granule content in NMR colonic crypts was significantly
larger than that of mouse colonic crypts (mouse 1.00 ± 0.05 a.u. vs. NMR 1.42 ± 0.19 a.u., p
< 0.0001; Figure 3j). Interestingly, differences in the AB/PAS staining colouration between
these two species may suggest that their mucin compositions differ, but further investigation
is required to substantiate this.

Figure 3. Goblet cells and mucus content in the gut of NMRs. Representative images of mouse (**a**) and NMR
(**b**) ileum samples stained with AB/PAS. **c**, number of AB$^+$ cells (goblet cells) in the ileum villi (left) and crypts
(right) in mice (n = 9, 3 sections/animal, 3 mice) and NMRs (n = 9, 3 sections/animal, 3 NMRs). **d**, area of AB$^+$
cells in the ileum villi (left) and crypts (right) in mice (n = 9, 3 sections/animal, 3 mice) and NMRs (n = 9, 3
sections/animal, 3 NMRs). **e**, pixel intensity measurement AB/PAS staining in the ileum villi (left) and crypts (right)
in mice (n = 9, 3 sections/animal, 3 mice) and NMRs (n = 9, 3 sections/animal, 3 NMRs). Representative images
of mouse (**f**) and NMR (**g**) colon samples stained with AB/PAS **h**, number of AB$^+$ cells in the colon crypts in mice
(n = 9, 3 sections/animal, 3 mice) and NMRs (n = 9, 3 sections/animal, 3 NMRs). **i**, area of AB$^+$ cells in crypts in

Altogether, these findings indicate that, despite having a shorter small intestine, the NMR ileum exhibits features that maximise the surface area for nutrient and water absorption, and ionic transport, such as ~2.5-fold larger villi compared to mice (Cohen's d size effect = 5.3). Additionally, the larger crypts found in NMRs may be reflective of differences in tissue turnover in these animals compared to mice. Furthermore, NMR show increased numbers of larger goblet cells both in the ileum and in the colon, with a significantly higher mucus content compared to murine goblet cells. This likely confers enhanced protection upon these animals against the infiltration of microorganisms, digestive enzymes and acids, digested food particles, as well as luminal by-products and toxins.

3.3. Intestinal integrity: ileum and colon.

Because of the clear differences in NMR intestinal structures compared to those of mice, we next aimed to examine the physical properties of this tissue. To this end, we used the gold standard technique for this purpose, i.e., the Ussing chamber for measuring mucosal membrane properties. We measured the transepithelial electrical resistance (TEER), as a measure of intestinal barrier integrity. High TEER values are indicative of decreased permeability to ions. Interestingly, the NMR terminal ileum showed a remarkably higher TEER, i.e. reduced permeability, compared to the mouse terminal ileum (mouse 45.41 ± 19.71 $\Omega \cdot cm^2$ vs. NMR 145.8 ± 100.8 $\Omega \cdot cm^2$, *p* = 0.0001; Cohen's *d* size effect = 6.27; Figure 4a, b). The NMR distal colon also showed reduced permeability compared to the mouse distal colon, but the difference was less pronounced (mouse 52.35 ± 20.27 $\Omega \cdot cm^2$ vs. NMR 65.65 ± 24.63 $\Omega \cdot cm^2$, *p* = 0.0062; Cohen's *d* size effect = 1.76; Figure 4c, d). While the terminal ileum in mice exhibited greater permeability compared to their colon (ileum 45.41 ± 19.71 $\Omega \cdot cm^2$ vs. colon 52.35 ± 20.27 $\Omega \cdot cm^2$, *p* = 0.0197; Cohen's *d* size effect = 1.24; Supplementary Figure 2a), which aligns with prior studies in mice (*49*) and is also noted in rats (*50, 51*) and humans (*51, 52*), the terminal ileum of NMRs was extraordinarily less permeable than their distal colon (ileum 145.8 ± 100.8 $\Omega \cdot cm^2$ vs. colon 65.65 ± 24.63 $\Omega \cdot cm^2$, *p* < 0.0001; Cohen's *d* size effect = 4.63; Supplementary Figure 2b). Altogether, these results indicate that the integrity of the gastrointestinal tract in NMRs is increased compared to mice, with the NMR small intestine (ileum) showing a strikingly low tissue permeability.

378
379 **Figure 4. Increased intestinal integrity in NMR. a**, **b)** transepithelial electrical resistance (TEER) in the terminal
380 ileum of mice (n = 7) and NMRs (n = 10). **c**, **d)** TEER in the distal colon of mice (n = 8) and NMRs (n = 8). Data in
381 **b** and **d** were calculated as the average TEER values between 60 min and 120 min and plotted as individual
382 values. *P* values are shown in plots. **a**, **c**, 2-way ANOVA and Šidak's multiple comparison test (data shown as
383 mean, shadow area represents the SD); **b**, **d**, Mann-Whitney test (data shown as median ± IQR).

384
3.4. Ion transport in the terminal ileum.

Given the different mucosal integrity properties observed in NMR vs. mouse ileal tissue, we
next utilised this region to assess intestinal ion secretion. To this end, we measured the
short-circuit current (I_{sc}), as this reflects the net ionic transport, mainly Cl^- and Na^+ in the
small intestine, through the mucosa (*6*). An increase in I_{sc} is primarily attributable to ion
secretion to the intestinal lumen. We first selected four compounds known to evoke ion
secretion in rodents and/or humans, such as 5-hydroxytryptamine (5-HT, serotonin) (*53, 54*),
bradykinin (*55, 56*), histamine (*57, 58*) and capsaicin (*59, 60*). In keeping with previous
studies, we confirmed that 5-HT, bradykinin, histamine, and capsaicin evoke a prominent ion
secretion to the ileal mucosa in mice (Figure 5a-d). By contrast, ileum preparations from
NMRs showed markedly reduced responses to 5-HT (mouse 179.8 ± 58.15 $\Delta\mu A \cdot cm^{-2}$ vs.
NMR 13.23 ± 17.41 $\Delta\mu A \cdot cm^{-2}$, *p* < 0.0001; Figure 5a) and bradykinin (mouse 78.64 ± 22.04
$\Delta\mu A \cdot cm^{-2}$ vs. NMR 17.98 ± 17.05 $\Delta\mu A \cdot cm^{-2}$, *p* < 0.0001; Figure 5b), and were virtually
unresponsive to histamine (mouse 22.28 ± 24.93 $\Delta\mu A \cdot cm^{-2}$ vs. NMR 3.08 ± 3.16 $\Delta\mu A \cdot cm^{-2}$,
p = 0.0006; Figure 5c) and capsaicin (mouse 42.56 ± 20.20 $\Delta\mu A \cdot cm^{-2}$ vs. NMR -0.74 ± 1.98
$\Delta\mu A \cdot cm^{-2}$, *p* = 0.0002; Figure 5d).

Next, we aimed to further evaluate the pro-secretory capacity of the NMR ileum via receptor-
independent pathways. To this end, we used the non-selective adenylate cyclase activator
forskolin, which elevates the intracellular cyclic adenosine monophosphate (cAMP)
concentration (*61*) and leads to increased intestinal ion secretion (Figure 5e) (*62*).
Remarkably, NMR ileum tissue showed significantly reduced forskolin-evoked responses
compared to mice (mouse 330.1 ± 166.1 $\Delta\mu A \cdot cm^{-2}$ vs. NMR 54.41 ± 12.96 $\Delta\mu A \cdot cm^{-2}$, *p* =
0.0009; Figure 5e).

Altogether, these results suggest that the cAMP signalling pathway is altered in the intestinal
epithelium of NMRs, likely influencing their intestinal ion pro-secretory pathways. Moreover,
NMRs display a pronounced resistance to intestinal ion secretion in response to some
neurotransmitters, inflammatory mediators, and chemical irritants.

Figure 5. Reduced ion secretion responses in NMRs. a) Short-circuit current increase (ΔI_{sc}) in response to 5-
HT in mice (n = 10 ileum sections, N = 3 animals) and NMRs (n = 11 ileum sections, N = 3 animals). **b)** ΔI_{sc} in
response to bradykinin in mice (n = 10 ileum sections, N = 3 animals) and NMRs (n = 11 ileum sections, N = 3
animals). **c)** ΔI_{sc} in response to histamine in mice (n = 7 ileum sections, N = 3 animals) and NMRs (n = 7 ileum
sections, N = 2 animals). **d)** ΔI_{sc} in response to capsaicin in mice (n = 6 ileum sections, N = 3 animals) and NMRs
(n = 10 ileum sections, N = 3 animals). **e)** ΔI_{sc} in response to forskolin in mice (n = 7 ileum sections, N = 2
animals) and NMRs (n = 7 ileum sections, N = 2 animals). For clarity, circles refer to mouse samples and squares
to NMRs samples. P values are shown in plots. All data were analysed with Mann-Whitney test (data shown as
median ± IQR).

3.5. Expression of receptors involved on intestinal ion transport.

Because NMRs showed altered ion secretion responses to 5-HT, bradykinin, histamine, and
capsaicin, we next asked whether NMRs express receptors for these compounds in their

intestinal tract. To this end, we used RT-qPCR to evaluate the gene expression of ion channels and receptors associated to these agonists. Importantly, some of these substances can activate several receptors (such as 5-HT, bradykinin, and histamine). Hence, we selected genes that encode receptors known to be involved in intestinal ion secretion in humans and/or rodents. Thus, we assessed the gene expression of the 5-HT receptor 4 (*Htr4*) (*53*), the bradykinin receptor B$_2$ (*Bdkrb2*) (*55*), histamine H$_1$ receptor (*Hrh1*) (*58*) and the capsaicin receptor, transient receptor potential vanilloid 1 (*Trpv1*). Notably, both the terminal ileum and the distal colon of NMR expressed all the genes evaluated. Ileum samples from NMR showed a reduced gene expression compared to mouse of *Bdkrb2* (mouse 0.014 ± 0.039 relative expression vs. NMR 0.001 ± 0.007 relative expression, *p* = 0.0002; Figure 6) and *Hrh1* (mouse 0.11 ± 0.73 relative expression vs. NMR 0.007 ± 0.022 relative expression, *p* = 0.0003; Figure 6), whereas similar levels of these genes were expressed in the colon of both species (*Bdkrb2*: mouse 0.01 ± 0.02 relative expression vs. NMR 0.01 ± 0.07 relative expression, *p* = 0.7984; *Hrh1*: mouse 0.01 ± 0.008 relative expression vs. NMR 0.01 ± 0.005 relative expression, *p* = 0.2220; Figure 6). By contrast, colon tissue of NMRs displayed a lower expression of *Htr4* compared to mice (mouse 0.14 ± 0.33 relative expression vs. NMR 0.01 ± 0.03 relative expression, *p* = 0.0002; Figure 6), although no such statistical difference was observed in the ileum across species for *Htr4* (mouse 0.04 ± 0.26 relative expression vs. NMR 0.18 ± 0.42 relative expression, *p* = 0.2785; Figure 6). These results indicate that, while changes in the gene expression of these receptors could partially account for the altered intestinal ion secretion observed in response to some of the compounds evaluated, NMRs express receptors for all these agents along their gastrointestinal tract. Further studies are needed to characterise intestinal secretory responses in these animals.

Figure 6. Intestinal gene expression of receptors associated with increased ion secretion and intestinal secretion of substance P. a) Heatmap of the relative gene expression in terminal ileum and distal colon samples in mice (n = 8) and NMRs (n = 8). **b**) Intestinal levels of substance P (normalised by the content of total protein) in mouse (n = 8; 4 ileum samples and 4 colon samples) and NMR (n = 6; 3 ileum samples and 3 colon samples). White circles refer to terminal ileum samples and grey squares to colon samples. All data were analysed with Mann-Whitney test (in **a**, data shown as colour gradient scale of the relative expression; and, in **b**, data shown as median ± IQR). Each row in **a** represents the median of the side-by-side replicates. ****P* < 0.001.

3.6. Substance P levels in the gut of naked mole-rats

Finally, because of its role on intestinal ion secretion (*63, 64*), we aimed to evaluate the
expression of the neuropeptide substance P in the gut of NMRs. Of interest, NMRs do not
express substance P and calcitonin gene-related peptide (CGRP) in cutaneous C-fibres (*65*),
in contrast to other mammalian species, including mice. Nevertheless, NMRs do produce
both substance P and CGRP in longitudinal lanceolate endings associated with follicle-sinus
complexes and body hair follicles (*65*), likely through Aβ-fibres. In keeping with this, we
previously observed CGRP$^+$ fibres in myenteric neurons in the colorectum of NMR (*43*).
Therefore, we asked whether substance P is also synthetised in the NMR gut. To this end,
we collected supernatants from intestinal explants (terminal ileum and distal colon) and
analysed the presence of substance P by enzyme immunoassay. The levels of substance P
under basal conditions were similar between NMRs and mice (mouse 1.38 ± 2.033 pg
substance P · mg protein^{-1} vs. NMR 1.09 ± 1.38 pg substance P · mg protein^{-1}, *p* > 0.9999;
Figure 6b). Further studies will investigate the capacity of different enteric cells to produce
this neuropeptide (such as neurons or enterochromaffin cells) and how this molecule and
other neuropeptides modulate the gut physiology of NMRs.

Discussion*:*

In the present study, we investigated some aspects of the intestinal barrier function in NMRs
as a potential emergent animal model in gastroenterology. Intestinal barrier dysfunction is
considered one of the potential underlying pathophysiological mechanisms of many
gastrointestinal diseases (*6*), and thus deemed to be of great biological and clinical
relevance (*11*). The persisting incidence and prevalence of gastrointestinal conditions in the
last decades (*1*) reflects the unmet need for novel strategies in the field. Here, we used
NMRs as a non-traditional animal model due to their exceptional biological features (*66*) and
studied their intestinal physiology. We found that the NMR small intestine was significantly
shorter compared to that of mice, although it displayed a more robust tissue architecture
(thicker intestinal layers, longer ileum villi, bigger crypts, and elevated mucus content). In
keeping with this, the integrity (measured as TEER) of the NMR ileum was remarkably
increased compared to the mouse ileum. The colon of NMR, on the other hand, showed
more structural similarities to mice, but again displayed enhanced mucus content and
superior tissue integrity. Of particular interest, we found that the ileum of NMRs showed
reduced ion secretion responses to 5-HT, bradykinin, histamine, and capsaicin, even though
receptors for these agonists were expressed in the gastrointestinal tract. Taken together,
NMRs represent an animal model of considerable interest for gastroenterology and,
particularly, for studying intestinal barrier function.

We first characterised the morphoanatomical features of the intestinal tract in NMRs. The
NMR's notably shorter small intestinal length has likely led to the simultaneous development
of longer villi compared to those mice, or rats (*67*). NMRs do not drink water and fulfil their
hydration requirements entirely through their solid diet. Therefore, we presume that longer
villi enable NMRs to increase the surface area to volume ratio for nutrient and water
absorption, which has led to water consumption being redundant. In keeping with this, a
recent study showed that long villi are found along the entire small bowel of NMR (*23*), as
well as enterocytes involved in the digestion and absorption of nutrients and water (*23, 68*).
Furthermore, NMRs exhibited a greater number of goblet cells in the ileum and colon.

Additionally, the size of these cells was substantially larger in NMRs and had increased
mucus granule content compared to mice. These results strongly suggest that goblet cells in
NMRs have an enhanced capacity to produce mucus/mucins. In keeping with this, it was
recently reported that NMRs have a thicker mucus layer and elevated expression of *Muc2*
(gene encoding for the oligomeric mucus gel-forming protein Mucin 2) compared to mice
(*23*), likely conferring an increased protection against environmental chemicals (*23*).

Notably, the histological features of the NMRs small intestine and colon more closely
resemble those of bigger rodents, such as rats (*69*) and guinea pigs (*70, 71*). This is of
particular interest because the altered intestinal architecture found in NMRs would not
completely explain the increased ileum tissue integrity (reflected as increased in TEER)
observed in these animals compared to mice. In fact, previous studies utilising Ussing
chambers have reported >2-fold lower TEER values in the ileum of guinea pigs (*72*) and rats
(*73, 74*) compared to our results in NMRs, and similar values in the colon of rats (*73−75*)
compared to our findings. This strongly suggests that the small intestine of NMRs possesses
strong tight junction structures that increase the integrity of this barrier. It is important to note,
however, that flux across tight junctions occurs by two distinct routes, referred to as the
"pore" and "leak" pathways. In the current study, we measured TEER as a proxy for intestinal
integrity, but this does not discriminate between these two different pathways (*6*). The pore
pathway is a high-capacity route that selectively allows passage of molecules based on
charge and size, with a maximum diameter of ~4-6 Å. On the other hand, the low-capacity

leak pathway is less selective and allows the passage of larger molecules (up to ~100 Å),
irrespective of the charge. Future studies are needed to evaluate in detail differences in the
pore and leak pathways between the small intestine and colon of NMRs. Moreover, due to
the crucial role of different tight junctions to selectively regulate each of these pathways, the
regulation of these barrier components is of great relevance. Thus, future studies will assess
the expression, organisation and interprotein interactions of tight junctions in the NMR small
intestine and how these control the pore and leak pathways. Overall, our data support that
the specific characteristics of the NMR small intestine may represent an interesting model for
studying intestinal barrier dysfunction in coeliac disease (*76*) and irritable bowel syndrome
(*77*), amongst other conditions(*23*).

Another crucial function of the intestinal barrier is the transport of ions and fluids across the
epithelium. Disruption of this well-coordinated process may lead to diarrhoea when the gut's
capacity for fluid absorption is exceeded. Of note, we found that the NMR ileum displays
reduced ion secretion responses to four different compounds: 5-HT, bradykinin, histamine,
and capsaicin. The bradykinin result aligns with our prior observation that this inflammatory
mediator fails to activate colonic afferent nerves in NMRs (*43*). However, we did observe that
bradykinin led to mechanical sensitisation of these fibres (*43*), suggesting that they express
functional bradykinin B_2 receptors. In the present study, we confirmed that NMRs indeed
express *Bdkrb2* (gene encoding the bradykinin B_2 receptor) in both the colon and ileum.
However, differences in the functionality of these receptors and the downstream effector
signalling in NMRs remain to be elucidated. Future studies will also explore the different cells
expressing this marker (epithelial cells (*78*) and nerve fibres (*79*)) in NMRs, and how these
modulate visceral pain and mucosal ion secretion. The role of 5-HT in the NMR
gastrointestinal tract has not been previously characterised. While fibroblasts isolated from

the back skin or the lungs of NMRs express high levels of 5-HT (*80*), plasma circulating
levels of this neurotransmitter are significantly lower compared to mice (*81*). In the gut, we
showed that the ion secretion induced by 5-HT in the ileum was decreased in NMRs
compared to mice, although the expression of *Htr4* in this tissue was similar in both species.
Further studies are therefore required to determine the functionality of this and other
serotonergic receptors, such as 5-HT receptors 1, 2 and 3 (*82*), in different cell types in the
NMR intestine. Finally, the lack of cutaneous neuropeptides and unusual connectivity of
cutaneous afferent neurons in the spinal cord in NMRs have been proposed to explain the
absence capsaicin-induced nocifensive behaviour and associated thermal hypersensitivity
(*83*), as well as the lack of histamine-induced scratching behaviour (*84*) in these animals.
However, it should be noted that the lack of substance P and CGRP alone cannot
completely explain the lack of pain- and itch-associated behaviours induced by these
compounds, as *Tac1-Calca* double knockout mice (therefore lacking both substance P and
CGRP◌ peptides) display intact responses to mechanical, thermal, chemical, visceral pain,
itch, and neurogenic inflammation (*85*). In the present study, we showed that despite the
virtually complete absence of ion secretion induced by histamine and capsaicin in NMRs,
these animals display intestinal basal levels of substance P comparable to mice. Future
studies will investigate the role of substance P and other neurotransmitters/neuropeptides in
the gut physiology of NMRs, as well as their cellular sources. Overall, we propose that
NMRs may be a promising model for the pathogenesis of inflammation-associated diarrhoea
in inflammatory bowel disease (*86*) and serotonin- and histamine-associated diarrhoea in
irritable bowel syndrome (*87, 88*).

The most notable observation regarding intestinal ion secretion is the reduced response to
the cAMP pathway-activator forskolin observed in NMR tissue. cAMP is a ubiquitous second
messenger found across all living domains that is generated by adenylate cyclases upon
conversion of adenosine triphosphate (ATP) to cAMP (*89*). Forskolin activates all adenylate
cyclases except AC9 and sAC (*90*), dramatically increasing the levels of cAMP in virtually all
tissues (*61*). Our findings raise numerous questions, not only concerning the intestinal ion
pro-secretory pathways of NMRs, but also more broadly about their cell biology. In the
context of intestinal ion secretion, the conventional paradigm indicates that activation of the
cAMP/protein kinase A (PKA) pathway activates the cystic fibrosis transmembrane
conductance regulator (CFTR) protein, representing the primary route for epithelial ion
secretion (*91, 92*). Of interest, a study (*93*) showed that adenylate cyclase 6 (AC6) is the
most abundant isoform in the gut of mouse and humans and is physically and functionally
associated with CFTR at the apical surface of intestinal epithelial cells. Interestingly, these
authors found that gut epithelial cells lacking AC6 fail to generate cAMP in response to
forskolin and AC6-deficient tracheal cells showed reduced forskolin-evoked Cl⁻ secretion
compared to preparations from littermate controls (*93*). Moreover, AC6-deficient mice were
protected from cholera toxin-induced diarrhoea (*93*), suggesting an important role of this
isoform in ion and water secretion. Whether AC6 is present in the NMR gut or if the
functionality of this and other adenylate cyclases is altered in this species remains to be
investigated. Furthermore, the functionality of the CFTR protein and associated signalling
pathways in this species should be investigated in detail. Finally, given that the cAMP
system plays fundamental roles in the regulation of the signalling pathways of serotonin (*94*),
bradykinin (*95*), and TRPV1 (*96*), this pathway could also interfere with the mechanisms

leading to receptor-dependent ion secretion studied here. Therefore, further studies are
needed to characterise the cAMP signalling system in NMRs.

We recognise that our study has some limitations. Although approximately physiologically
age-matched animals for both species were used, intestinal permeability typically changes
throughout the lifetime. However, it should be noted that differences in intestinal permeability
between mice and NMRs are in the opposite direction of what one would expect, that is, an
increased permeability in older animals (*97*). Interestingly, we found that the ileum
permeability was similar in young NMRs compared to older animals (young 153.0 ± 57.09 Ω
· cm^2 vs. older 144.2 ± 42.75 Ω · cm^2, *p* = 0.7000; Supplementary Figure 2c), thus
demonstrating that the differences observed between species are not related to age.
Furthermore, all NMRs used in the present study were non-breeders and thus we cannot
comment on whether or findings would hold true in sexual mature, breeding animals. Sexual
maturity is of relevance to our work because it has been shown, for example, that
progesterone decreases gut permeability in humans and Caco-2 cells (*98*) and that
oestrogen modulates ion secretion in intestinal epithelial cells (*99*). Most of the NMRs used
in our study were males (only 1 female was used in TEER experiments and another 1 in
short-circuit current experiments), and thus we are unable to draw firm conclusions with
regard to any potential sex differences, but being non-breeder animals, all were sexually
immature. However, we do not believe that sex and/or state of sexual maturity are likely
responsible for the significant differences observed in the NMR ileum, as the relevant
hormones would similarly affect the colon, where we found similar TEER values as
previously reported in rats (*73–75*). It should be noted as well that the Ussing chambers
experiments were performed at 37°C, that is, 5-7°C above the NMRs physiological
temperature (*100*). However, an increase in the physiological temperature is typically
associated with an increase in mucosal permeability (*101, 102*). It remains to be elucidated
whether changes in physiological temperature affect intestinal function in NMRs, although
one would normally expect changes in the opposite direction to our findings. Finally, other
caveats that must be considered are the differences in diet and microbiota between mice
and NMRs, which can affect both the intestinal integrity (*103*) and the ion secretion (*104*).
Whether specific gut bacteria found in NMRs (*25*) or compounds of their diet are important to
improve intestinal barrier function remain to be elucidated.

In conclusion, our work illustrates a remarkably low permeability in the NMR small intestine.
Moreover, it reveals that these animals have reduced intestinal ion secretion responses to 5-
HT, bradykinin, histamine, and capsaicin. Overall, this study adds further evidence to the
uncommon biological features exhibited by this species. The NMR gut may therefore offer
particular insights into the physiology relevant to gastrointestinal conditions and aid in
devising strategies for their treatments. Thus, we strongly support the use of this
unconventional animal model in further investigations of intestinal barrier function.

4. Acknowledgements:

JA-L and EStJS acknowledge funding from the Medical Research Council (MRC)
(MR/W002426/1) and AR was supported by AstraZeneca PhD Studentship (G115018).

5. Disclosures:

The authors declare no competing interests.

6. Author contribution:

JA-L conceptualised and designed the research study, conducted the experiments (Ussing chamber experiments, gene expression, enzyme immunoassays), acquired and analysed the data and wrote the manuscript. AR conducted the experiments (histology, AB/PAS staining and IHC) and acquired the data. DCB provided experimental support and discussed the data. EStJS provided experimental support, reviewed, and discussed the data and revised the manuscript. All authors corrected and approved the final version of the manuscript.

7. Data availability:

Data sets supporting the conclusions of this article are available in University of Cambridge Apollo Repository (http://doi.org/10.17863/CAM.106848).

8. ORCID iD:

Javier Aguilera-Lizarraga: https://orcid.org/0000-0002-8135-9933.

Anne Ritoux: https://orcid.org/0000-0002-5760-6172

David C. Bulmer: https://orcid.org/0000-0002-4703-7877

Ewan St. J. Smith: https://orcid.org/0000-0002-2699-1979.

9. References:

1. Y. Wang, Y. Huang, R. C. Chase, T. Li, D. Ramai, S. Li, X. Huang, S. O. Antwi, A. P. Keaveny, M. Pang, Global Burden of Digestive Diseases: A Systematic Analysis of the Global Burden of Diseases Study, 1990 to 2019. *Gastroenterology* **165**, 773-783.e15 (2023).

2. T. C. Rose, A. Pennington, C. Kypridemos, T. Chen, M. Subhani, J. Hanefeld, L. Ricciardiello, B. Barr, Analysis of the burden and economic impact of digestive diseases and investigation of research gaps and priorities in the field of digestive health in the European Region—White Book 2: Executive summary. *United European Gastroenterol J* **10**, 657–662 (2022).

3. V. C. Goodoory, C. E. Ng, C. J. Black, A. C. Ford, Direct healthcare costs of Rome IV or Rome III-defined irritable bowel syndrome in the United Kingdom. *Aliment Pharmacol Ther* **56**, 110–120 (2022).

4. R. S. Sandler, J. E. Everhart, M. Donowitz, E. Adams, K. Cronin, C. Goodman, E. Gemmen, S. Shah, A. Avdic, R. Rubin, The burden of selected digestive diseases in the United States. *Gastroenterology* **122**, 1500–1511 (2002).

5. L. A. Pattison, A. Cloake, S. Chakrabarti, H. Hilton, R. H. Rickman, J. P. Higham, M. Y. Meng, L. W. Paine, L. Qiu, A. Ritoux, D. C. Bulmer, G. Callejo, E. St John Smith, Digging

deeper into pain – an ethological behavior assay correlating well-being in mice with
human pain experience. *Pain*, doi: 10.1101/2023.08.18.553862 (2024).

6. T. Vanuytsel, J. Tack, R. Farre, The Role of Intestinal Permeability in Gastrointestinal
Disorders and Current Methods of Evaluation. *Front Nutr* **8**, 717925 (2021).

7. H. Majamaa, E. Isolauri, Evaluation of the gut mucosal barrier: Evidence for increased
antigen transfer in children with atopic eczema. *J Allergy Clin Immunol* **97**, :985-90
(1996).

8. A. Benard, P. Desreumeaux, D. Huglo, A. Hoorelbeke, A.-B. Tonnel, B. Wallaert, iD Lille,
Increased intestinal permeability in bronchial asthma. *J Allergy Clin Immunol* **97**,
1173–1181 (1996).

9. A. Fasano, N. P. Visanji, L. W. C. Liu, A. E. Lang, R. F. Pfeiffer, Gastrointestinal
dysfunction in Parkinson's disease. Lancet Publishing Group [Preprint] (2015).
https://doi.org/10.1016/S1474-4422(15)00007-1.

10. M. P. Fink, Leaky gut hypothesis: a historical perspective. [Preprint] (1990).
https://doi.org/10.1097/00003246-199005000-00024.

11. M. Camilleri, Leaky gut: mechanisms, measurement and clinical implications in
humans. BMJ Publishing Group [Preprint] (2019). https://doi.org/10.1136/gutjnl-
2019-318427.

12. H. Vanheel, M. Vicario, T. Vanuytsel, L. Van Oudenhove, C. Martinez, À. V. Keita, N.
Pardon, J. Santos, J. D. Soderholm, J. Tack, R. Farré, Impaired duodenal mucosal
integrity and low-grade inflammation in functional dyspepsia. *Gut* **63**, 262–71 (2014).

13. A. Fedi, C. Vitale, G. Ponschin, S. Ayehunie, M. Fato, S. Scaglione, In vitro models
replicating the human intestinal epithelium for absorption and metabolism studies: A
systematic review. *Journal of Controlled Release* **335**, 247–268 (2021).

14. J. Fogh, T. Orfeo, J. Tiso, F. E. Sharkey, Establishment of human colon carcinoma lines
in nude mice. *Exp Cell Biol* **47**, 136–144 (1979).

15. M. González-González, C. Díaz-Zepeda, J. Eyzaguirre-Velásquez, C. González-Arancibia,
J. A. Bravo, M. Julio-Pieper, Investigating gut permeability in animal models of disease.
Frontiers Media S.A. [Preprint] (2019). https://doi.org/10.3389/fphys.2018.01962.

16. J. Foulke-Abel, J. In, J. Yin, N. C. Zachos, O. Kovbasnjuk, M. K. Estes, H. De Jonge, M.
Donowitz, Human Enteroids as a Model of Upper Small Intestinal Ion Transport
Physiology and Pathophysiology. *Gastroenterology* **150**, 638-649.e8 (2016).

17. F. Weiß, D. Holthaus, M. Kraft, C. Klotz, M. Schneemann, J. D. Schulzke, S. M. Krug,
Human duodenal organoid-derived monolayers serve as a suitable barrier model for
duodenal tissue. *Ann N Y Acad Sci* **1515**, 155–167 (2022).

18. A. J. Barker, U. Koch, G. R. Lewin, S. J. Pyott, *The Extraordinary Biology of the Naked
Mole-Rat* (2021)vol. 1319.

19. R. Buffenstein, The Naked Mole-Rat: A New Long-Living Model for Human Aging
Research. *J Gerontol A Biol Sci Med Sci* **60**, 1369–77 (2005).

20. K. Oka, M. Yamakawa, Y. Kawamura, N. Kutsukake, K. Miura, The Naked Mole-Rat as a
Model for Healthy Aging. *Annu Rev Anim Biosci* **11**, 207–226 (2023).

21. G. Maldonado, G. Hernández, Translational control in the naked mole-rat as a model
highly resistant to cancer. [Preprint] (2021).
https://doi.org/10.1016/j.bbcan.2020.188455.

22. Y. H. Edrey, D. X. Medina, M. Gaczynska, P. A. Osmulski, S. Oddo, A. Caccamo, R.
Buffenstein, Amyloid beta and the longest-lived rodent: the naked mole-rat as a

model for natural protection from Alzheimer's disease. *Neurobiol Aging* **34**, 2352–60
(2013).

23. S. Montazid, S. Bandyopadhyay, D. W. Hart, N. Gao, B. Johnson, S. G. Thrumurthy, D. J.
Penn, B. Wernisch, M. Bansal, P. M. Altrock, F. Rost, P. Gazinska, P. Ziolkowski, B. H.
Hayee, Y. Liu, J. Han, A. Tessitore, J. Koth, W. F. Bodmer, J. E. East, N. C. Bennett, I.
Tomlinson, S. Irshad, Adult stem cell activity in naked mole rats for long-term tissue
maintenance. *Nat Commun* **14**, 8484 (2023).

24. S. Wirtz, V. Popp, M. Kindermann, K. Gerlach, B. Weigmann, S. Fichtner-Feigl, M. F.
Neurath, Chemically induced mouse models of acute and chronic intestinal
inflammation. *Nat Protoc* **12**, 1295–1309 (2017).

25. T. Debebe, E. Biagi, M. Soverini, S. Holtze, T. B. Hildebrandt, C. Birkemeyer, D.
Wyohannis, A. Lemma, P. Brigidi, V. Savkovic, B. König, M. Candela, G. Birkenmeier,
Unraveling the gut microbiome of the long-lived naked mole-rat. *Sci Rep* **7**, 9590
(2017).

26. H. G. Hilton, N. D. Rubinstein, P. Janki, A. T. Ireland, N. Bernstein, N. L. Fong, K. M.
Wright, M. Smith, D. Finkle, B. Martin-McNulty, M. Roy, D. M. Imai, V. Jojic, R.
Buffenstein, Single-cell transcriptomics of the naked molerat reveals unexpected
features of mammalian immunity. *PLoS Biol* **17**, e3000528 (2019).

27. T. S. Taylor, P. Konda, S. S. John, D. C. Bulmer, J. R. F. Hockley, E. S. J. Smith, Galanin
suppresses visceral afferent responses to noxious mechanical and inflammatory
stimuli. *Physiol Rep* **8**, e14326 (2020).

28. S. U. Mondelaers, S. A. Theofanous, M. V. Florens, E. Perna, J. Aguilera-Lizarraga, G. E.
Boeckxstaens, M. M. Wouters, W. M. Mondelaers SU, Theofanous SA, Florens MV,
Perna E, Aguilera-Lizarraga J, Boeckxstaens GE, S. U. Mondelaers, S. A. Theofanous, M.
V. Florens, E. Perna, J. Aguilera-Lizarraga, G. E. Boeckxstaens, M. M. Wouters, Effect of
genetic background and postinfectious stress on visceral sensitivity in Citrobacter
rodentium-infected mice. *Neurogastroenterology and Motility* **28**, 647–658 (2016).

29. P. Bankhead, M. B. Loughrey, J. A. Fernández, Y. Dombrowski, D. G. McArt, P. D. Dunne,
S. McQuaid, R. T. Gray, L. J. Murray, H. G. Coleman, J. A. James, M. Salto-Tellez, P. W.
Hamilton, QuPath: Open source software for digital pathology image analysis. *Sci Rep*
7, 16878 (2017).

30. K. R. Rogers-Broadway, E. Karteris, Amplification efficiency and thermal stability of
qPCR instrumentation: Current landscape and future perspectives. *Exp Ther Med* **10**,
1261–1264 (2015).

31. D. Svec, A. Tichopad, V. Novosadova, M. W. Pfaffl, M. Kubista, How good is a PCR
efficiency estimate: Recommendations for precise and robust qPCR efficiency
assessments. *Biomol Detect Quantif* **3**, 9–16 (2015).

32. N. Eissa, H. Hussein, H. Wang, M. F. Rabbi, C. N. Bernstein, J. E. Ghia, Stability of
reference genes for messenger RNA quantification by real-time PCR in mouse dextran
sodium sulfate experimental colitis. *PLoS One* **11**, e0156289 (2016).

33. N. Eissa, L. Kermarrec, H. Hussein, C. N. Bernstein, J. E. Ghia, Appropriateness of
reference genes for normalizing messenger RNA in mouse 2,4-dinitrobenzene sulfonic
acid (DNBS)-induced colitis using quantitative real time PCR. *Sci Rep* **7**, 42427 (2017).

34. F. Wang, J. Wang, D. Liu, Y. Su, Normalizing genes for real-time polymerase chain
reaction in epithelial and nonepithelial cells of mouse small intestine. *Anal Biochem*
399, 211–217 (2010).

35. J. Zhang, Y. Y. Gao, Y. Q. Huang, Q. Fan, X. T. Lu, C. K. Wang, Selection of housekeeping
genes for quantitative gene expression analysis in yellow-feathered broilers. *Ital J*
Anim Sci **17**, 540–546 (2018).
36. A. B. Nygard, C. B. Jørgensen, S. Cirera, M. Fredholm, Selection of reference genes for
gene expression studies in pig tissues using SYBR green qPCR. *BMC Mol Biol* **8** (2007).
37. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, doi:
10.1006/meth.2001.1262 (2001).
38. J. Aguilera-Lizarraga, M. V. Florens, M. F. Viola, P. Jain, L. Decraecker, I. Appeltans, M.
Cuende-Estevez, N. Fabre, K. Van Beek, E. Perna, D. Balemans, N. Stakenborg, S.
Theofanous, G. Bosmans, S. U. Mondelaers, G. Matteoli, S. Ibiza Martínez, C. Lopez-
Lopez, J. Jaramillo-Polanco, K. Talavera, Y. A. Alpizar, T. B. Feyerabend, H. R. Rodewald,
R. Farre, F. A. Redegeld, J. Si, J. Raes, C. Breynaert, R. Schrijvers, C. Bosteels, B. N.
Lambrecht, S. D. Boyd, R. A. Hoh, D. Cabooter, M. Nelis, P. Augustijns, S. Hendrix, J.
Strid, R. Bisschops, D. E. Reed, S. J. Vanner, A. Denadai-Souza, M. M. Wouters, G. E.
Boeckxstaens, Local immune response to food antigens drives meal-induced
abdominal pain. *Nature* **590**, 151–156 (2021).
39. S. De Schepper, S. Verheijden, J. Aguilera-Lizarraga, M. F. Viola, W. Boesmans, N.
Stakenborg, I. Voytyuk, I. Smidt, B. Boeckx, I. Dierckx de Casterlé, V. Baekelandt, E.
Dominguez Gonzalez, M. Mack, I. Depoortere, B. De Strooper, B. Sprangers, U.
Himmelreich, S. Soenen, M. Guilliams, P. Vanden Berghe, E. Jones, D. Lambrechts, G.
Boeckxstaens, Self-Maintaining Gut Macrophages Are Essential for Intestinal
Homeostasis. *Cell* **175**, 400-415.e13 (2018).
40. C. Wang, R. Weindruch, J. R. Fernández, C. S. Coffey, P. Patel, D. B. Allison, Caloric
restriction and body weight independently affect longevity in Wistar rats.
International journal of obesity and related metabolic disorders: journal of the
International Association for the Study of Obesity. **28**, 357–362 (2004).
41. M. Altun, E. Bergman, E. Edström, H. Johnson, B. Ulfhake, Behavioral impairments of
the aging rat. *Physiol Behav* **92**, 911–923 (2007).
42. B. Wiggs, A. Churg, J. L. Wright, Influence of weight on pulmonary function in the
adult guinea pig. *Respiration* **58**, 37–41 (1991).
43. J. R. F. Hockley, K. H. Barker, T. S. Taylor, G. Callejo, Z. M. Husson, D. C. Bulmer, E. S. J.
Smith, Acid and inflammatory sensitisation of naked mole-rat colonic afferent nerves.
Mol Pain **16**, 1744806920903150 (2020).
44. F. Treves, Anatomy of the Intestinal Canal and Peritoneum in Man. *British Medical*
Journal **1**, 580–583 (1885).
45. B. M. L. Underhill, Intestinal length in man. *British Medical Journal* **2**, 1243–1246
(1955).
46. G. Hounnou, C. Destrieux, J. Desmé, P. Bertrand, S. Velut, Anatomical study of the
length of the human intestine. *Surgical and Radiologic Anatomy* **24**, 290–294 (2002).
47. S. Sadahiro, T. Ohmura, Y. Yamada, T. Saito, Y. Taki, Analysis of length and surface area
of each segment of the large intestine according to age, sex and physique. *Surg Radiol*
Anat **14**, 251–257 (1992).
48. L. T. Weaver, S. Austin, T. J. Cole, M. Dunn, Small intestinal length: a factor essential for
gut adaptation. *Gut* **32**, 1321–1323 (1991).
49. N. Hanning, R. Verboven, J. G. De Man, H. Ceuleers, H. U. De Schepper, A. Smet, B. Y.
De Winter, Single-day and multi-day exposure to orogastric gavages does not affect

intestinal barrier function in mice. *American Journal of Physiology-Gastrointestinal*
and Liver Physiology **324**, G281–G294 (2023).

50. L. Feighery, A. Smyth, S. Keely, A. W. Baird, W. T. O'Connor, J. J. Callanan, D. J. Brayden,
Increased intestinal permeability in rats subjected to traumatic frontal lobe
percussion brain injury. *Journal of Trauma - Injury, Infection and Critical Care* **64**, 131–
137 (2008).

51. P. Nejdfors, M. Ekelund, B. Jeppsson, B. R. Weström, Mucosal in vitro permeability in
the intestinal tract of the pig, the rat, and man: Species- and region-related
differences. *Scand J Gastroenterol* **35**, 501–507 (2000).

52. Å. Sjöberg, M. Lutz, C. Tannergren, C. Wingolf, A. Borde, A. L. Ungell, Comprehensive
study on regional human intestinal permeability and prediction of fraction absorbed
of drugs using the Ussing chamber technique. *European Journal of Pharmaceutical*
Sciences **48**, 166–180 (2013).

53. M. R. Budhoo, R. P. Harris, J. M. Kellum, 5-Hydroxytryptamine-induced Cl- transport is
mediated by 5-HT3 and 5-HT4 receptors in the rat distal colon. *Eur J Pharmacol* **298**,
137–144 (1996).

54. M. R. Budhoo, J. M. Kellum, E. Woltering, S. Ashley, B. Bass, The 5-HT4 receptor
mediates 5-hydroxytryptamine-induced rise in short circuit current in the human
jejunum in vitro. *Surgery* **116**, 396–400 (1994).

55. A. W. Baird, M. M. Skelly, D. P. O'Donoghue, K. E. Barrett, S. J. Keely, Bradykinin
regulates human colonic ion transport in vitro. *Br J Pharmacol* **155**, 558–566 (2008).

56. D. C. Manning, S. H. Snyder, J. F. Kachur, R. J. Miller, M. Field, Bradykinin receptor-
mediated chloride secretion in intestinal function. *Nature* **299**, 256–259 (1982).

57. Y. Wang, H. J. Cooke, R. Fertel, H. Su, Histamine augments colonic secretion in guinea
pig distal colon. *American journal of physiology* **258**, G432–G439 (1990).

58. G. Schultheiss, B. Hennig, W. Schunack, G. Prinz, M. Diener, Histamine-induced ion
secretion across rat distal colon: Involvement of histamine H1 and H2 receptors. *Eur J*
Pharmacol **546**, 161–170 (2006).

59. S. Yarrow, J. A. Ferrar, H. M. Cox, The effects of capsaicin upon electrogenic ion
transport in rat descending colon. *Naunyn-Schmiedeberg's Arch Pharmacol* **344**, 557–
563 (1991).

60. E. Weber, M. Neunlist, M. Schemann, T. Frieling, Neural components of distension-
evoked secretory responses in the guinea-pig distal colon. *Journal of Physiology* **536**,
741–751 (2001).

61. K. B. Seamon, W. Padgett, J. W. Daly, Forskolin: Unique diterpene activator of
adenylate cyclase in membranes and in intact cells. *Proc Natl Acad Sci U S A* **78**, 3363–
7 (1981).

62. R. J. Bridges, W. Rummel, B. Simon, Forskolin induced chloride secretion across the
isolated mucosa of rat colon descendens. *Naunyn Schmiedebergs Arch Pharmacol*
323, 355–60 (1983).

63. L. Wang, A. M. Stanisz, B. K. Wershil, S. J. Galli, M. H. Perdue, Substance P induces ion
secretion in mouse small intestine through effects on enteric nerves and mast cells.
Am J Physiol Gastrointest Liver Physiol **269**, G85-92 (1995).

64. M. Riegler, I. Castagliuolo, P. T. C. So, M. Lotz, C. Wang, M. Wlk, T. Sogukoglu, E.
Cosentini, G. Bischof, G. Hamilton, B. Teleky, E. Wenzl, J. B. Matthews, C. Pothoulakis,
Effects of substance P on human colonic mucosa in vitro. *Am J Physiol Gastrointest*
Liver Physiol **276**, G1473-83 (1999).

65. T. J. Park, C. Comer, A. Carol, Y. Lu, H. S. Hong, F. L. Rice, Somatosensory organization
and behavior in naked mole-rats: II. Peripheral structures, innervation, and selective
lack of neuropeptides associated with thermoregulation and pain. *Journal of*
Comparative Neurology **465**, 104–120 (2003).

66. R. Buffenstein, V. Amoroso, B. Andziak, S. Avdieiev, J. Azpurua, A. J. Barker, N. C.
Bennett, M. A. Brieño-Enríquez, G. N. Bronner, C. Coen, M. A. Delaney, C. M. Dengler-
Crish, Y. H. Edrey, C. G. Faulkes, D. Frankel, G. Friedlander, P. A. Gibney, V. Gorbunova,
C. Hine, M. M. Holmes, J. U. M. Jarvis, Y. Kawamura, N. Kutsukake, C. Kenyon, W. T.
Khaled, T. Kikusui, J. Kissil, S. Lagestee, J. Larson, A. Lauer, L. A. Lavrenchenko, A. Lee, J.
B. Levitt, G. R. Lewin, K. N. Lewis Hardell, T. H. D. Lin, M. J. Mason, D. McCloskey, M.
McMahon, K. Miura, K. Mogi, V. Narayan, T. P. O'Connor, K. Okanoya, M. J. O'Riain, T. J.
Park, N. J. Place, K. Podshivalova, M. E. Pamenter, S. J. Pyott, J. Reznick, J. G. Ruby, A.
B. Salmon, J. Santos-Sacchi, D. K. Sarko, A. Seluanov, A. Shepard, M. Smith, K. B.
Storey, X. Tian, E. N. Vice, M. Viltard, A. Watarai, E. Wywial, M. Yamakawa, E. D.
Zemlemerova, M. Zions, E. S. J. Smith, The naked truth: a comprehensive clarification
and classification of current 'myths' in naked mole-rat biology. *Biological Reviews* **97**,
115–140 (2022).

67. G. E. Holle, Changes in the structure and regeneration mode of the rat small intestinal
mucosa following benzalkonium chloride treatment. *Gastroenterology* **101**, 1264–73
(1991).

68. N. Miron, V. Cristea, Enterocytes: Active cells in tolerance to food and microbial
antigens in the gut. [Preprint] (2012). https://doi.org/10.1111/j.1365-
2249.2011.04523.x.

69. A. Bosi, D. Banfi, M. Bistoletti, L. M. Catizzone, A. M. Chiaravalli, P. Moretto, E. Moro,
E. Karousou, M. Viola, M. C. Giron, F. Crema, C. Rossetti, G. Binelli, A. Passi, D. Vigetti,
C. Giaroni, A. Baj, Hyaluronan Regulates Neuronal and Immune Function in the Rat
Small Intestine and Colonic Microbiota after Ischemic/Reperfusion Injury. *Cells* **11**,
3370 (2022).

70. M. Mitjans, R. Ferrer, Morphometric Study of the Guinea Pig Small Intestine during
Development. *Microsc Res Tech* **63**, 206–214 (2004).

71. J. Zhao, D. Liao, J. Yang, H. Gregersen, Biomechanical remodelling of obstructed
guinea pig jejunum. *J Biomech* **43**, 1322–1329 (2010).

72. R. Moore, C. Pothoulakis, J. Thomas Lamont, S. Carlson, J. L. Madara, J. L. MADARA C
difficile, C. difficile toxin A increases intestinal permeability and induces Cl- secretion.
Am J Physiol **259**, G165-72 (1990).

73. L. Feighery, A. Smyth, S. Keely, A. W. Baird, W. T. O'Connor, J. J. Callanan, D. J. Brayden,
Increased intestinal permeability in rats subjected to traumatic frontal lobe
percussion brain injury. *Journal of Trauma - Injury, Infection and Critical Care* **64**, 131–
137 (2008).

74. B. I. Polentarutti, A. L. Peterson, À. K. Sjöberg, E. K. I. Anderberg, L. M. Utter, A. L. B.
Ungell, Evaluation of viability of excised rat intestinal segments in the Ussing
chamber: Investigation of morphology, electrical parameters, and permeability
characteristics. *Pharm Res* **16**, 446–54 (1999).

75. A. A. Livanova, A. A. Fedorova, A. V. Zavirsky, I. I. Krivoi, A. G. Markov, Dose- and
Segment-Dependent Disturbance of Rat Gut by Ionizing Radiation: Impact of Tight
Junction Proteins. *Int J Mol Sci* **24**, 1753 (2023).

76. R. M. Van Elburg, J. J. Uil, C. J. J. Mulder, H. S. A. Heymans, Intestinal permeability in
patients with coeliac disease and relatives of patients with coeliac disease. *Gut* **34**,
354–7 (1993).

77. A. M. González-Castro, C. Martínez, E. Salvo-Romero, M. Fortea, C. Pardo-Camacho, T.
Pérez-Berezo, C. Alonso-Cotoner, J. Santos, M. Vicario, Mucosal pathobiology and
molecular signature of epithelial barrier dysfunction in the small intestine in irritable
bowel syndrome. [Preprint] (2017). https://doi.org/10.1111/jgh.13417.

78. A. W. Baird, M. M. Skelly, D. P. O'Donoghue, K. E. Barrett, S. J. Keely, Bradykinin
regulates human colonic ion transport in vitro. *Br J Pharmacol* **155**, 558–66 (2008).

79. J. Yi, Z. Bertels, J. S. Del Rosario, A. J. Widman, R. A. Slivicki, M. Payne, H. M. Susser, B.
A. Copits, R. W. Gereau, Bradykinin receptor expression and bradykinin-mediated
sensitization of human sensory neurons. *Pain* **165**, 202–215 (2024).

80. Y. Kawamura, K. Oka, T. Semba, M. Takamori, Y. Sugiura, R. Yamasaki, Y. Suzuki, T.
Chujo, M. Nagase, Y. Oiwa, S. Fujioka, S. Homma, Y. Yamamura, S. Miyawaki, M. Narita,
T. Fukuda, Y. Sakai, T. Ishimoto, K. Tomizawa, M. Suematsu, T. Yamamoto, H. Bono, H.
Okano, K. Miura, Cellular senescence induction leads to progressive cell death via the
INK4a-RB pathway in naked mole-rats. *EMBO J* **42**, e111133 (2023).

81. K. N. Lewis, N. D. Rubinstein, R. Buffenstein, A window into extreme longevity; the
circulating metabolomic signature of the naked mole-rat, a mammal that shows
negligible senescence. *Geroscience* **40**, 105–121 (2018).

82. M. B. Hansen, A. B. Witte, The role of serotonin in intestinal luminal sensing and
secretion. [Preprint] (2008). https://doi.org/10.1111/j.1748-1716.2008.01870.x.

83. T. J. Park, Y. Lu, R. Jüttner, E. S. J. Smith, J. Hu, A. Brand, C. Wetzel, N. Milenkovic, B.
Erdmann, P. A. Heppenstall, C. E. Laurito, S. P. Wilson, G. R. Lewin, Selective
inflammatory pain insensitivity in the African naked mole-rat (Heterocephalus glaber).
PLoS Biol **6**, 0156–0170 (2008).

84. E. S. J. Smith, G. R. C. Blass, G. R. Lewin, T. J. Park, Absence of histamine-induced itch
in the African naked mole-rat and "rescue" by Substance P. *Mol Pain* **6**, 29 (2010).

85. D. I. Macdonald, M. Jayabalan, J. Seaman, A. Nickolls, A. Chesler, Pain persists in mice
lacking both Substance P and CGRPα signaling. *Elife* **13**, RP93754 (2024).

86. A. N. Anbazhagan, S. Priyamvada, W. A. Alrefai, P. K. Dudeja, Pathophysiology of IBD
associated diarrhea. *Tissue Barriers* **6**, e1463897 (2018).

87. W. D. Chey, J. Kurlander, S. Eswaran, Irritable bowel syndrome: A clinical review.
American Medical Association [Preprint] (2015).
https://doi.org/10.1001/jama.2015.0954.

88. J. D. Wood, Neuropathophysiology of the Irritable Bowel Syndrome. *J Clin*
Gastroenterol **35**, S11-22 (2002).

89. K. F. Ostrom, J. E. Lavigne, T. F. Brust, R. Seifert, C. W. Dessauer, V. J. Watts, R. S.
Ostrom, Physiological roles of mammalian transmembrane adenylyl cyclase isoforms.
[Preprint] (2022). https://doi.org/10.1152/physrev.00013.2021.

90. S. Pierre, T. Eschenhagen, G. Geisslinger, K. Scholich, Capturing adenylyl cyclases as
potential drug targets. [Preprint] (2009). https://doi.org/10.1038/nrd2827.

91. T. V. Strong, K. Boehm, F. S. Collins, Localization of cystic fibrosis transmembrane
conductance regulator mRNA in the human gastrointestinal tract by in situ
hybridization. *Journal of Clinical Investigation* **93**, 319–27 (1994).

92. R. A. Frizzell, J. W. Hanrahan, Physiology of epithelial chloride and fluid secretion. *Cold*
Spring Harb Perspect Med **2**, a009563 (2012).

93. A. Thomas, Y. Ramananda, K. S. Mun, A. P. Naren, K. Arora, AC6 is the major adenylate
cyclase forming a diarrheagenic protein complex with cystic fibrosis transmembrane
conductance regulator in cholera. [Preprint] (2018).
https://doi.org/10.1074/jbc.RA118.003378.

94. X. Wang, K. Wang, X. Wu, W. Huang, L. Yang, Role of the cAMP–PKA–CREB–BDNF
pathway in abnormal behaviours of serotonin transporter knockout mice. *Behavioural*
Brain Research **419**, 113681 (2022).

95. B. S. Dixon, Cyclic AMP selectively enhances bradykinin receptor synthesis and
expression in cultured arterial smooth muscle: Inhibition of angiotensin II and
vasopressin response. *Journal of Clinical Investigation* **93**, 2535–44 (1994).

96. G. Bhave, W. Zhu, H. Wang, D. J. Brasier, G. S. Oxford, R. W. Gereau IV, cAMP-
dependent protein kinase regulates desensitization of the capsaicin receptor (VR1) by
direct phosphorylation. *Neuron* **35**, 721–31 (2002).

97. N. Thevaranjan, A. Puchta, C. Schulz, A. Naidoo, J. C. Szamosi, C. P. Verschoor, D.
Loukov, L. P. Schenck, J. Jury, K. P. Foley, J. D. Schertzer, M. J. Larché, D. J. Davidson, E.
F. Verdú, M. G. Surette, D. M. E. Bowdish, Age-Associated Microbial Dysbiosis
Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage
Dysfunction. *Cell Host Microbe* **21**, 570 (2018).

98. Z. Zhou, C. Bian, Z. Luo, C. Guille, E. Ogunrinde, J. Wu, M. Zhao, S. Fitting, D. L. Kamen,
J. C. Oates, G. Gilkeson, W. Jiang, Progesterone decreases gut permeability through
upregulating occludin expression in primary human gut tissues and Caco-2 cells. *Sci*
Rep **9**, 8367 (2019).

99. X. Yang, Y. Guo, J. He, F. Zhang, X. Sun, S. Yang, H. Dong, Estrogen and estrogen
receptors in the modulation of gastrointestinal epithelial secretion. *Oncotarget* **8**,
97683–97692 (2017).

100. V. M. Kovalzon, O. A. Averina, V. A. Minkov, A. A. Petrin, M. Yu. Vysokikh, Unusual
Correlation between Rest–Activity and Body Temperature Rhythms in the Naked Mole
Rat (Heterocephalus glaber) as Compared to Five Other Mammalian Species. *J Evol*
Biochem Physiol **56**, 451–458 (2020).

101. G. P. Lambert, C. V. Gisolfi, D. J. Berg, P. L. Moseley, L. W. Oberley, K. C. Kregel, Selected
contribution: Hyperthermia-induced intestinal permeability and the role of oxidative
and nitrosative stress. *J Appl Physiol* **92**, 1750–61 (2002).

102. K. Dokladny, P. L. Moseley, T. Y. Ma, Physiologically relevant increase in temperature
causes an increase in intestinal epithelial tight junction permeability. *Am J Physiol*
Gastrointest Liver Physiol **290**, G204-12 (2006).

103. O. Inczefi, P. Bacsur, T. Resál, C. Keresztes, T. Molnár, The Influence of Nutrition on
Intestinal Permeability and the Microbiome in Health and Disease. [Preprint] (2022).
https://doi.org/10.3389/fnut.2022.718710.

104. A. C. Engevik, M. A. Engevik, Exploring the impact of intestinal ion transport on the
gut microbiota. *Comput Struct Biotechnol J* **19**, 134–144 (2021).

10. Supplementary information:

Supplementary Table 1. Primer sequences for gene detection by RT-qPCR.

Species	Sequence (5' → 3')	Target marker	NCBI Ref Seq
Mouse	CACCCCCTTGTACCCTTCAC	TATA-box binding protein (*Tbp*)	NM_013684.3
	CAGTTGTCCGTGGCTCTCTT		NM_013684.3
Mouse	CCGGCTCGCTTGAGAAAAGG	Bradykinin receptor β2 (*Bdkrb2*)	NM_009747.2
	CCGACAGAAACCCGAGTAGC		NM_009747.2
Mouse	ACGAGTGAAACCCGATGCTT	Histamine receptor H1 (*Hrh1*)	NM_001252642.3
	CTCTGGCTGTCCTAGGGAGT		NM_001252642.3
Mouse	CCTCACAGCAACTTCTCCTT	5-hydroxytryptamine (serotonin) receptor 4 (*Htr4*)	NM_008313.4
	TCCCCTGACTTCCTCAAATA		NM_008313.4
Mouse	GAGTATGCCCAGAGCCCATC	Transient receptor potential vanilloid 1 (*Trpv1*)	NM_001001445.2
	CAGGCTGCTGTGTGGTAAGA		NM_001001445.2
NMR	TTAATTTCGTTTCCGCGGGC	TATA-box binding protein (*Tbp*)	XM_004868859.3
	GCATAAGGTGGGAGGCTGTT		XM_004868859.3
NMR	CTGGGAGGGAGCTGTAGTCA	Bradykinin receptor β2 (*Bdkrb2*)	XM_021244763.1
	CCTCCTAGGCACTTCCTCCT		XM_021244763.1
NMR	CCCACACGTGGCTTAGAGTT	Histamine receptor H1 (*Hrh1*)	XM_004874845.3
	TTGCTTTGAACATGCAGCCC		XM_004874845.3
NMR	ATGTGGCTCACCACCTTAGC	5-hydroxytryptamine (serotonin) receptor 4 (*Htr4*)	XM_021266180.1
	GTGCCTCATGCCTCACTGAT		XM_021266180.1
NMR	TGTGCAATCGCTGTTCATGC	Transient receptor potential vanilloid 1 (*Trpv1*)	NM_001279859.1
	TGGAAGCCTCGGGTGTAGTA		NM_001279859.1

1001

Supplementary Figure 1. Extended anatomical analyses of the NMR gastrointestinal tract. Representative
images of the histological measurements performed in ileum (**a**) and colon (**b**) samples. **c)** Colon length in male
(n = 13) and female (n = 4) NMRs. **d)** Small intestine length in male (n = 13) and female (n = 4) NMRs. **e)**
Correlation between age and colon length in NMRs (n = 17). **f)** Correlation between weight and colon length in
NMRs (n = 17). **g)** Correlation between age and small intestine length in NMRs (n = 17). **h)** Correlation between
weight and small intestine length in NMRs (n = 16). *P* values are shown in plots. **c** and **d**, Mann Whitney test

(data shown as median ± IQR); **e**, **f**, **g** and **h**, two-tailed Pearson's correlation (data shown as individual data, and
95% CI).

1010

1011

1012　**Supplementary Figure 2. Extended analyses of the tissue integrity in NMR and mice**. **a)** TEER in the
1013　terminal ileum (n = 7) and colon (n = 8) in mice. **b)** TEER in the terminal ileum (n = 10) and colon (n = 8) in
1014　NMRs. **c)** TEER in the terminal ileum of young (<25 weeks old; n = 3) and old (>40 weeks old; n = 7) NMRs. Data
1015　were calculated as the average TEER values between 60 min and 120 min and plotted as individual values. *P*
1016　values are shown in plots. **a**, **b** and **c**, Mann-Whitney test (data shown as median ± IQR).