Genetic architecture of heritable leaf microbes

Julia A. Boyle[1], Megan. E. Frederickson[1], John R. Stinchcombe[1,2]

1. Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St.,

Toronto, ON, Canada, M5S3B2

2. Swedish Collegium for Advanced Study, SE-752 38 Uppsala, Sweden

julia.boyle@mail.utoronto.ca, m.frederickson@utoronto.ca, john.stinchcombe@utoronto.ca

Corresponding author information: Julia Boyle, julia.boyle@mail.utoronto.ca, Department of
Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St., Toronto, ON,
Canada, M5S3B2

Abstract

Background

Host-associated microbiomes are shaped by both their environment and host genetics, and often impact host performance. The scale of host genetic variation important to microbes is largely unknown, yet fundamental to the community assembly of host-associated microbiomes, and with implications for the eco-evolutionary dynamics of microbes and hosts. Using *Ipomoea hederacea*, Ivy-leaved morning glory, we generated matrilines differing in quantitative genetic variation and leaf shape, which is controlled by a single Mendelian locus. We then investigated the relative roles of Mendelian and quantitative genetic variation in structuring the leaf microbiome, and how these two sources of genetic variation contributed to microbe heritability.

Results

We found that despite large effects of the environment, both Mendelian and quantitative genetic host variation were important in contributing to microbe heritability, and that the cumulative small effect genomic differences due to matriline explained as much or more microbial variation than a single large effect locus. Furthermore, our results are the first to suggest that leaf shape itself contributes to variation in the abundances of some microbes in the leaf microbiome.

Conclusions

Our results demonstrate the roles of different scales of host genetic variation in the assembly of a natural microbiome. The genetic architecture of plant-associated microbiomes depends on both quantitative genetic variation and Mendelian traits, with similar contributions to microbe heritability. The presence of Mendelian and quantitative genetic variation for host associated microbes suggests that evolution in plant hosts-- at the leaf shape locus, and other quantitative genetic loci-- has the potential to affect microbiome distribution, abundance, and composition.

Keywords: leaf shape, phyllosphere, *Ipomoea hederacea*, microclimate, host age, leaf

microbiome, heritability, matriline, Mendelian, quantitative genetics

Background

The community assembly of microbes on a host depends on the environment, the host's

traits, and the microbes [1]. Microbes that colonize leaves and other aboveground plant parts,

known as the phyllosphere, can disperse to a leaf via air [2], rain [3], or soil [4], after which they

experience selection due to conditions on the leaf surface and microbe-microbe interactions [1,

5]. Host genetics play a role in the leaf microbiome; the similarity of leaf microbiomes across

species depends on host phylogenetic relatedness [6, 7], and the similarity of microbiomes

between individuals of the same species can depend on host intraspecific genetic variation [8–

11]. Here we test how a single leaf shape gene affects leaf bacterial communities in *Ipomoea*

hederacea, and compare the magnitude of effect from that single locus to that of many small

effect loci.

The genetically-based physiological, morphological, or immune traits of hosts can lead to

consistent associations between the abundance or community composition of microbes and host

genotypes [12–15]. When consistent host-microbe associations are present, the host's microbial

phenotype can be considered a heritable host trait, even when the microbes are environmentally-

acquired [16–18]. In other words, horizontally-transmitted microbes can be heritable in host

populations when host traits have predictable effects on microbial recruitment and alleles for

these host traits are transmitted from host parents to host offspring. The heritability of a host's

microbial phenotype is important to understanding host evolution and ecology due to microbial

effects on host fitness [19–21], and potential reciprocal selection between hosts and microbiomes

[17, 22, 23]. The variation in microbe heritability may have interesting evolutionary

repercussions on hosts, given that the phenotypic variance of many host traits can be microbially

mediated [17, 18]. Furthermore, if a host's microbial phenotype is due to heritable host genetic

variation, then the evolutionary forces affecting those host traits-- selection, drift, mutation,

migration, non-random mating-- can in turn affect microbial abundances, distributions, and

microbiome composition [24]. Like all heritable traits, host microbial phenotypes can be

influenced by few loci of large effect or the cumulative effects of many small effect loci, and

understanding microbe heritability necessitates we determine the scale of host variation that

matters to microbes.

Mechanisms producing genotype-specific phyllosphere microbiomes are complex and

varied. In *Arabidopsis thaliana*, a genome-wide association study of the phyllosphere microbial

community found that loci related to defense and plant cell wall integrity affect microbial

community variation, while species richness was affected by loci involved with viral

reproduction, trichomes, and morphogenesis [9]. Host genotypes may differ in immune genes

and disease resistance, which have been linked to differences in switchgrass leaf fungal

community structure [25] and inconsistent effects on bacterial and fungal maize phyllospheres

[11]. Plant genotypes can also vary in leaf morphology and other leaf attributes; for example,

mutations in cuticle formation and ethylene production also affected the microbiome in a

synthetic *A. thaliana* phyllosphere [26]. The microclimate of a leaf is determined by many leaf

properties including surface temperature, surface area, thickness of the boundary layer of air,

chemical composition, trichomes, gas exchange, and nutrients [27]. The boundary layer of air is

closest to the leaf surface and has warmer, humid air; non-segmented leaves with larger surface

area often have a thicker boundary layer, and thicker boundary layers can impede gas exchange

from the leaves [28]. Compounds and water released by leaves during gas exchange can be an

important source of nutrients and resources for phyllosphere microbes [5], and they can impose

selection on which microbial species can establish. These findings suggest that leaf morphology,

and its effects on microclimates and resources, can potentially affect phyllosphere microbes.

Factors driving phyllosphere microbiome composition may depend strongly on the

environment and the interaction between environment and genotype, since the leaf microbiome

is primarily environmentally acquired [29]. Environmental factors like weather and seasonality

influence which microbes are available to disperse to leaves and with what frequency [29].

Phyllosphere microbes may differ over time due to changing moisture and temperature

conditions abiotically filtering for microbes in the air and soil [30], and biotic interactions that

emerge at different times, e.g. changing microbe-microbe interactions and new microbes from

insects [31] or other plants [32]. Determining the relative roles of environment, phenology, and

plant genotype on the microbial community composition and abundance in the phyllosphere is an

unresolved empirical challenge.

One unstudied potential source of host genotype influence on the leaf microbiome is leaf

shape. Leaf shape is a key morphological difference between plant species [33] and could

contribute to interspecific variation in leaf microbiomes. The shape of a leaf determines the

temperature and humidity conditions it experiences and even its photosynthetic and gas exchange

abilities [34, 35], potentially affecting microbial communities. Comparing the effects of leaf

shape on the microbiome between plant species is confounded by other genetic and ecological

differences between taxa, and for this reason, it is preferable to use a single species that exhibits

intraspecific variation in leaf shape. *Ipomoea hederacea*, Ivy-leaved morning glory, is a

flowering annual plant whose leaf shape is determined by a single Mendelian locus [36–38];

homozygous genotypes are either fully lobed or whole, with heterozygotes being partially lobed

(Figure 1). In *I. hederacea,* homozygous lobed genotypes have been shown to more protected

against extreme temperature changes compared to whole leaves; homozygous lobed leaves

remained warmer at night by a mean difference of 0.16-0.22°C, likely due to their altered

boundary layer resulting in leaves being more coupled with ambient air temperature [39, 40].

The differences in *Ipomoea hederacea* leaf morphology, and its effects on temperature and

boundary layers that affect leaf microclimates, suggest that different leaf shape genotypes could

directly determine which colonizing microbes establish on the leaf, indirectly affecting microbe-

microbe interactions, and structuring leaf microbial communities as a whole.

To assess the relative contributions of quantitative and Mendelian genetic variation we

first examined if a single large effect Mendelian locus underlying leaf shape generates leaf

microbiome differences within a host species. Then, we estimated the heritability of microbes

due to many small effect loci that differ between plant lines– in other words, the amount of

microbiome variation attributable to quantitative genetic causes. Finally, we compared the

relative importance of host genotype to other factors such as germination time, and therefore host

age, to assess the role of the environment in structuring leaf microbiomes.

Methods

Study system and crossing design

Ipomoea hederacea, Ivy-leaved morning glory, is a flowering annual plant, commonly

found in eastern North America in roadside ditches and agricultural fields. It is predominantly

selfing [41], and dies with the first hard frost in autumn. We used seeds derived from a cross by

Campitelli and Stinchcombe [39], where individuals from the two alternate homozygous leaf

shape phenotypes (i.e., fully lobed or whole) were collected from North Carolina, USA, selfed

for seven generations to generate homozygous parents (P1), and then crossed with each other

(Figure 1). A single F1 was allowed to self-fertilize, producing F2 progeny. We scored F2 plants

for leaf shape, and allowed them to self-fertilize; we refer to all the selfed progeny of an

individual F2 plant as a "matriline." We used F3 seeds, set by F2 plants we had identified as

heterozygous for leaf shape, as our experimental seeds. In total, we used seeds from 82

matrilines. As expected, genotype frequencies did not significantly differ from a 1:2:1 ratio for

leaf shape ($\chi^2_{(2,\,218)}= 3$, $p = 0.22$), with 55 plants with whole leaves, 112 plants with partially

lobed leaves, and 51 plants with fully lobed leaves.

Our breeding design has three consequences for leaf microbiomes, which we return to in

the Discussion. First, variation between matrilines is due to the combined effects of loci that

differed between the original parents of the cross and the effects of recombination. Second,

because of the effects of recombination, significant differences between leaf shape genotypes are

due to either the effects of the leaf shape locus itself or linked loci not broken up by

recombination. Third, within a matriline, we expect 25% of the loci that differed between the

parents to still be heterozygous, which is considerably more within-matriline genetic variation

than would be found in inbred lines or recombinant inbred lines (RILs), potentially reducing our

power and making our estimates of microbe heritability more conservative.

Figure 1. *Ipomoea hederacea* leaf polymorphism and crossing design. Leaf shape is determined by a single Mendelian locus. First, both homozygous genotypes were collected from North Carolina, USA and selfed for seven generations to create the parents (P1). To generate the plants used in the field experiment, the two homozygous P1 individuals were crossed, then heterozygote offspring were selfed for two more generations. Our experiment used F3 plants from 82 matrilines. Photos taken by Julia Boyle, with brightness and colors edited for style and clarity.

Field site and experimental design

In 2021, we planted a total of 240 *I. hederacea* seeds from 82 matrilines (2-3 seeds/matriline) in a common garden at Koffler Scientific Reserve (www.ksr.utoronto.ca) in Ontario, Canada. We plowed and disked an old field for the common garden site, and we assigned morning glories in 3 blocks of 80 plants each, planted 1 m apart with a 6 ft tall bamboo

pole to climb. We scarified seeds then planted them in a greenhouse in peat pots containing soil

from the field. We planted the first cohort on June 4th and a second cohort on July 2nd, because

of poor germination in the first cohort. After the majority of plants in a given cohort germinated

(approximately one week), we transplanted the pots into the field. The 218 surviving plants

consisted of 83 individuals from the first cohort, and 135 individuals from the second cohort.

There were no significant differences in the frequencies of leaf shape genotypes between the two

cohorts ($\chi^2_{(2, 218)}$= 0.31, p = 0.86; Table S1). The common garden was moderately weeded until

the *I. hederacea* plants had established; neighboring plants in the common garden were

predominantly *Cirsium arvense*, Canada thistle. On September 1st, we collected similarly-sized

leaves within one foot from the ground by cutting them with sterilized scissors into a sterile

plastic bag, after which they were stored in a -80°C freezer until DNA extraction. We avoided

leaves that looked very young or very old, as well as leaves with herbivory damage. At the time

of leaf collection, plants in the first germination cohort were 89 days old with 96% of plants

flowering, and plants in the second germination cohort were 61 days old with 49% of plants

flowering. We photographed collected leaves against a 1cm^2 grid background, and measured leaf

surface area and shape using ImageJ [42]. From these photographs, we calculated leaf circularity

as 4π(Area/Perimeter^2), which has a value between 0 and 1, with 1 indicating a perfect circle.

Sequencing and QIIME2 analysis

We performed extractions using the whole leaves with QIAGEN DNeasy® PowerSoil®

Pro Kits; both epiphytic and endophytic microbes were extracted. We sent samples to Génome

Québec (Montréal, Canada) for Illumina MiSeq PE 250bp 16S rRNA gene amplicon sequencing

on the conserved hypervariable V4 region (primer pair 515F-806R). We used Quantitative

Insights Into Microbial Ecology 2 (QIIME2) v.2022.2 [43] to trim the sequences for quality and

we denoised the sequences with DADA2 [44]. Samples had a median read frequency of 35,436

reads. Using QIIME2, we removed amplicon sequence variants (ASVs) that had fewer than 10

reads across all samples, and assigned taxonomy using the 'sklearn' feature classifier and

Greengenes 16S rRNA gene V4 region reference [45], then filtered out reads assigned as

cyanobacteria and mitochondria to remove plant DNA. After these steps, we had 4,934 ASVs

across the 218 leaf samples, with samples having a median read frequency of 4,523 reads. After

visualizing a rarefaction curve, we created a rarefied dataset with 4,695 ASVs total across 183

samples, with 1809 reads/sample; the first and second cohorts retained 77 and 106 samples

respectively. Finally, we constructed a phylogeny using QIIME2's MAFFT [46] and FastTree 2

[47] to obtain a rooted tree.

Statistical analysis

For statistical analysis, we used R v4.2.0 [48], with the tidyverse [49], lmerTest [50],

phyloseq [51], vegan [52], and microbiome [53] packages. The general model structure we used

included leaf shape genotype and germination cohort as fixed effects, and random effects of

block and matriline, unless otherwise specified. All linear models were adjusted with type III

ANOVAs, calculated in the car package [54].

First, we compared leaf shape surface area and circularity between genotypes using linear

mixed effect models. Next, we identified the core microbiota by filtering the rarefied dataset to

genera found in at least 80% of samples. To assess the effect of leaf shape and germination

cohort on microbial composition, we calculated a weighted UniFrac distance matrix using

rarefied data, then we used an adonis2 PERMANOVA (999 permutations) followed by a

permutational test of dispersion on significant groups using betadisper. The vegan package does

not allow for random effects, so we did not include block and matriline. To confirm that

germination cohort sample sizes did not affect our results, we also downsampled to equal cohort

sizes 100 times, and used the same permutation tests for composition and dispersion (Figure S3).

To visualize the community, we used a weighted UniFrac Principal Coordinate Analysis (PCoA)

using rarefied relative abundance data.

We estimated broad-sense heritability (H^2) of microbial community phenotypes and

taxonomic groups using linear mixed effects models, taking a similar approach as Wagner et al.

[10] and Grieneisen et al. [14]. Here, the random effects of block and matriline explain variance

in the microbe abundance phenotype after accounting for the mean effects of the germination

cohort and leaf shape, meaning our estimates of H^2 are conditioned on the fixed effects of the

germination cohort and leaf shape (as is commonly the case [55]). To estimate H^2 from the

random effects, we calculated the ratio of matriline genetic variance (V_G) to the sum of matriline,

spatial block, and residual variance ($V_P = V_G + V_{Block} + V_E$). We generated the community

phenotypes using rarefied data and included observed species richness (base 10 log-

transformed), evenness, Shannon diversity, and the first three axes of the weighted UniFrac

PCoA as the response variables in individual linear mixed effects models. To estimate the

heritability of taxonomic groups in a more compositionally aware way [56], we aggregated the

non-rarefied dataset to genera that were in at least 30% of samples, added 1 to each abundance,

and center log ratio transformed the abundance matrix. Next, we fit a linear mixed effects model

to the abundance of each genus that was in at least 30% of samples. To assess the statistical

significance of the matriline effect, we compared the log-likelihood of our models with and

without line as a random effect; the difference in log-likelihoods is X^2 distributed, with one

degree of freedom [57]. Finally, solely for the purposes of a qualitative comparison of the

magnitude of variance explained by the leaf shape locus, matriline effects, spatial blocks, and

residual variation, we used the same mixed effects model except with leaf shape as a random

effect.

Results

Genotype morphology

Leaf shape genotypes of *I. hederacea* differed quantitatively. Leaf shape genotypes

differed significantly in measures of circularity (*Wald $X^2_{(2,218)}$*=1520, *p*<0.001; Table S2), with

the whole genotype having the highest circularity followed by heterozygotes (Figure S1). While

there was a trend for leaf surface area to be the largest in whole leaves and lowest in the lobed

leaves, the difference was not significantly predicted by leaf shape genotype (*Wald $X^2_{(2,218)}$*=4.15,

p>0.05; Table S2) or germination cohort (*Wald $X^2_{(1,218)}$*=2.22, *p*>0.05; Table S2) (Figure S1). As

we specifically chose to sample similarly sized leaves, with the intent of minimizing differences

in total leaf area, these results suggest our sampling strategy worked and that we can attribute

differences in microbes due to leaf shape as not simply being the result of differences in leaf

area.

Community composition and diversity

The core *I. hederacea* leaf microbes were the genera *Methylobacterium*, *Sphingomonas*,

Deinococcus, *Pseudomonas*, *Arthrobacter*, and *Hymenobacter*. Communities were structured by

the high relative abundances of *Methylobacterium* and *Sphingomonas* genera (Figure 3 and 4),

and their relative abundances were negatively correlated (Figure S2). The first PCoA axis was

strongly positively correlated with *Methylobacterium* relative abundance while the second axis

was strongly negatively correlated with *Sphingomonas* relative abundance (Figure S2).

Germination cohort was the main driver of overall microbial community composition.

While microbiomes from both cohorts overlapped in composition, they were significantly

different from each other ($F_{(1,182)}$=10.4, p=0.001; Table S3); older plants from the first cohort had

more *Methylobacterium*, and younger plants from the second cohort had more *Sphingomonas*

(Figure 4ACD). The younger cohort's microbiomes were more distinct from one another,

creating significantly higher dispersion ($F_{(1,182)}$=10.4, p=0.001; Table S3). Germination cohorts

always remained significantly different in composition and dispersion when cohort sample sizes

were equalized (Figure S3), suggesting this is a true biological phenomenon. Germination cohort

significantly predicted observed richness (*Wald* $X^2_{(1,182)}$=10.1, p=0.001; Table S4), with older

plants having higher ASV richness than younger plants, but it did not predict Shannon diversity

(*Wald* $X^2_{(1,182)}$=3.33, p>0.05; Table S4) or evenness (*Wald* $X^2_{(1,182)}$=2.96, p>0.05; Table S4).

Despite differences in morphology and microclimate that we *a priori* predicted to affect

the leaf microbiome, we did not detect strong leaf shape effects on community composition as a

whole. Leaf shape did not significantly structure composition ($F_{(2,182)}$=1.83, p>0.05; Table S3),

or affect Shannon diversity, evenness, or observed richness (*Wald* $X^2_{(2,182)}$<3.97, p>0.05 for all;

Table S4). While whole community composition was not affected, we still identified microbial

genera significantly affected by leaf shape, as discussed below.

Heritable microbes and community phenotypes

Microbial community phenotypes did not have a strong genetic basis, yet we still

discovered significant heritable variation in some microbial traits. Community phenotypes had

low broad-sense heritability ($0 \leq H^2 \leq \sim 0.05$) with the third axis of the PCoA having the highest H^2,

followed by Shannon diversity (Figure 2A). All community phenotypes with non-zero H^2 were

significantly heritable, as the models' log-likelihoods were improved by including matriline;

community phenotypes with no H^2 (evenness and PCoA axis 2) showed no difference in fit when

matriline was included (Figure S4). The main effect of the germination cohort significantly

affected observed richness, as previously described, as well as the first two axes of the PCoA

(*Wald X*$^2_{(1,182)}$>6.71, *p*<0.01 for both axes; Table S4), whereas leaf shape did not significantly

affect community phenotypes (*Wald X*$^2_{(2,182)}$<4.85, *p*>0.05 for all; Table S4). In our qualitative

comparison of the magnitudes of variances explained, plant matriline had a larger effect on

community phenotypes' H^2 than leaf shape, with the exception of observed richness (Figure 2C,

Table S5).

Host genetics had a stronger influence on the abundance of microbial genera than

community phenotypes, with both quantitative genetic and Mendelian traits affecting microbe

abundance. Considering the genera in at least 30% of samples, we identified eight genera with

non-zero and significant broad-sense heritability. The most heritable genera ($0.05 \leq H^2 \leq 0.07$)

were *Nocardioides*, *Kineococcus*, *Pseudomonas*, and *Agrobacterium* (Figure 2B). Our likelihood

ratio test results showed that including plant matriline improved model fit for 10 out of 15 genera

(which included all genera with non-zero H^2), there was no difference in fit for 3 genera

(*Deinococcus, Hymenobacter, Sphingomonas*), and slightly reduced model fit for 2 genera

(*Actinotelluria*, *Rubellimicrobium*) (Figure S4). The germination cohort significantly affected the

abundance of more than half of the most common genera (*Wald X*$^2_{(1,218)}$>3.83, *p*<0.05 for all;

Table S6, Figure 2B). Four genera, *Sphingomonas*, *Nocardioides*, *Methylobacterium* and

Agrobacterium, had significant main effects of leaf shape on their phenotype (*Wald*

$X^2_{(2,218)}$>7.03, *p*<0.05 for all; Table S6, Figure 2B). When we examined whether the magnitude

of plant matriline variance was qualitatively comparable to leaf shape locus, we found that these

four genera had equal to larger magnitudes of variance attributable to leaf shape than plant

matriline (Figure 2D, Table S7). In heritable genera with a non-significant main effect of leaf

shape, plant matriline explained more variance than leaf shape (Figure 2D, Table S7), as

expected.

Figure 2. Broad-sense heritability (H^2) of community phenotypes and common microbial

genera. Genera are ordered by percent variance explained by plant matriline in the H^2 model. A

and B: H^2 of community phenotypes and genera in at least 30% of samples. H^2 was calculated as

the percent variation explained by genetic line compared to the total of matriline, spatial block,

and residual variances. Significant fixed effects of germination cohort and leaf shape are

indicated by diamonds and triangles respectively. C and D: H^2 of community phenotypes and

genera in at least 30% of samples, but with leaf shape as a random effect to directly compare the

magnitude of variance explained to that of matriline.

Figure 3. Relative abundance of bacterial families in the *Ipomoea hederacea* leaf microbiome,

indicated by color. Samples are ordered by *Sphingomonadaceae* abundance and grouped by

germination cohort. The first germination cohort (A) was 89 days old (n=83) and the second

germination cohort (B) was 61 days old (n=135) at the time of sampling. For this visualization

we included only families in at least 30% of samples and at least 1% relative abundance.

Figure 4. Weighted UniFrac PCoA of *I. hederacea* leaf microbiomes. Germination cohort

significantly predicted community composition (A), while leaf shape was non-significant (B).

Core microbiota structured communities (C-F). For C-F, note that the relative abundance legend

scales differ.

Discussion

Microbes can be considered an extended host phenotype with potentially adaptive

functions for the host [58], and heritable microbes are more likely to consistently affect host

phenotype and fitness through time. Here we examined the relative contributions of Mendelian

and quantitative genetic variation to the heritability of host-associated microbes to determine

what types of host genetic variation mattered to microbial ecology. We found that both

Mendelian and quantitative genetic host variation contribute to microbe heritability, and that the

cumulative small effect genomic differences due to matriline explained as much or more

variation than a single large effect locus. We were able to identify heritable microbial variation

despite a large effect of germination timing, which incorporated host age and initial microbial

environment, and which was a major factor structuring community composition. We discuss the

implications of our results below.

Genetic influence on the microbiome

There was genetic variation at several scales mediating how well microbes can establish

and perform in the phyllosphere. It is important to note that, as always, our estimates of

heritability are specific to the population [59] and conditioned on fixed effects in the model [55],

in our case, the germination cohort and leaf shape. A single leaf shape gene significantly

explained the abundance of several common microbes, suggesting the genetic differences in leaf

microclimate (or other, unknown features of the leaf shape genotypes) impact microbial

establishment and persistence, and can contribute to intraspecific differences in the phyllosphere.

When comparing the magnitude of effect, both leaf shape and matriline were important factors

influencing microbe abundance, although plant matriline affected more of the microbes tested.

Community phenotypes like species diversity and weighted UniFrac distances showed lower

heritability than individual genera, but since the majority of microbes are not likely to be

heritable in an environmentally acquired microbiome, very low heritability at a community scale

was perhaps unsurprising. Our estimates of heritability are low to commensurate compared to

other heritable microbiome studies (summarized in Table S8; [10, 12–14, 16, 60–64]; while the

range of heritabilities for plant-associated microbes across other studies was as broad as H^2=0-1,

the average significant heritability of microbes and community phenotypes was usually low

(H^2≤0.10) (Table S8). Our results therefore follow the trend of low heritability for

environmentally acquired microbes, with a strong influence of the environment.

In our experiment, each plant line represented a different mosaic of the original parental

genotypes, similar in concept to recombinant inbred lines (RILs). However, because the seeds we

used in our experiment were F3 individuals, approximately 25% of the loci initially heterozygous

in the F1 (i.e., those that differed between the parents) remain heterozygous within matrilines.

One consequence of this is that there is genetic variation between matrilines due to the effects of

recombination, and genetic variation within matrilines (due to heterozygosity and segregation

within a matriline). These effects make it more difficult to detect quantitative genetic variation in

microbial phenotypes, especially compared to RIL populations that have been made homozygous

within lines. Our results may thus underestimate the prevalence and magnitude of H^2 due to

reduced power. Furthermore, our crossing design included only two parental genotypes,

potentially limiting the amount of genetic variation segregating in the cross. For instance, had we

sampled more genotypes to create MAGIC or Nested Association Mapping lines [65, 66], more

genetic variation would be captured in our population. It is difficult to determine, *a priori*,

whether this would lead to increased or decreased estimates of genetic variation. On the one

hand, two parental genotypes is a limited sample of the genetic variation found in many

populations. Although putatively neutral genetic variation in *I. hederacea* populations is often

quite low [67], there is frequently quantitative genetic variation [68, 69]. On the other hand, the

selfing rate in *I. hederacea* is quite high (~93%) [67, 70]. Rare outcrossing events, followed by

selfing, may produce recombinant populations not that dissimilar from our experimental

population. Similar arguments have been made for *A. thaliana* [71, 72], which is also highly

selfing. As a qualitative investigation of the influence of including limited genetic variation in

our crossing design, we gathered estimates of H^2 for life history and quantitative traits in

Arabidopsis thaliana, specifically comparing studies that used either multiple accessions or

RILs. These data (Table S9), while an imperfect comparison, suggest that if anything, our

estimates of heritability could be under-estimated from using two parental lines in the initial

cross.

Evolutionary implications

We found similar magnitudes of variance in microbial phenotypes attributable to plant

line and leaf shape genotype for several common bacterial genera, which is important because

there is the potential for different evolutionary forces to act on quantitative genetic loci versus

Mendelian loci. For example, Mendelian traits like leaf shape have high potential to show

sampling effects due to drift, especially in small populations [73]. In contrast, drift is much less

likely to produce a change in quantitative traits which are influenced by many genes [73]. Thus,

drift can potentially lead to bigger effects on leaf shape-associated microbes. Furthermore, high

selfing in *I. hederacea* will reduce the frequency of heterozygous leaf shape genotypes by as

much as 50% per generation. In this way, the mating system may have outsized effects on

microbes linked to Mendelian traits. If natural selection acts on host traits, as it appears to with *I.*

hederacea's leaf shape [38], there may be indirect effects of selection on the microbial

phenotype in the host population.

The microbes we identified as having heritable variation in abundance are also linked to

host performance and fitness in other systems. These findings from other systems suggest

intriguing hypotheses for further study. For example, one of the most heritable genera that was

also significantly affected by leaf shape, *Agrobacterium*, has pathogens linked to causing tumor-

like galls in plants [74], suggesting a potential link of leaf shape to disease risk.

Methylobacterium spp. are linked to improving plant growth [75], and increased growth rate

improves fitness in *I. hederacea* [39]. *Pseudomonas* spp. have a wide range of positive or

negative effects on plant performance [76]. The most heritable genus *Nocardioides* is commonly

endophytic on plant leaves and stems [77, 78] and has species capable of fixing nitrogen [79] and

reducing nitrate [80]. Evaluating whether any of these microbial genera which have effects on

host performance in other systems do so in *I. hederacea* will require further empirical work. If

heritable microbes affect plant fitness, then there is higher potential for reciprocal selection

between the microbes and plant hosts.

Environmental and phenological effects on the leaf microbiome

Community-wide differences between germination cohorts could be due to seasonal

differences in what microbes were present in the plant's early life and the amount of time

selection had to act on the microbial communities. Input of new microbes to the phyllosphere can

be low relative to environments like soil rhizospheres, and these sources may change across the

growing season [29]. The cumulative input of microbes may matter as well, since we found that

the older plant phyllospheres had a higher species richness than younger plants; this is possibly

because the leaves themselves were older and had more time to collect rare microbes dispersing

from the environment. Additionally, as plants age and change phenologically, this may change

the leaf environment for microbes [81–83]. Whilst in a time of growth and development, young

plants have weaker immune responses [84], meaning they may impose weaker selection on their

phyllospheres. Thus, the disparate microbiomes on young plants could be due to weaker

selection from the environment and host over a shorter time. In contrast, older plants showed

more similar and clustered microbiomes suggesting that over time there is selection on microbes

by the *I. hederacea* leaf microclimate, and the microbial community converges. The merging

community compositions between young and old plants could suggest microbial succession

occurred in the phyllosphere [85, 86].

The large effect of germination cohort is not unexpected, given that most studies on

microbiome heritability find a very large effect of environment and host age on the microbiome

[10, 12, 16, 87]. For example, Walters et al. [16] found that plant age was the largest driver of

the maize rhizosphere, but in one year they nonetheless found 143 heritable root microbes with

H^2=0.15-0.25. While our results suggest genera like *Methylobacterium* and *Sphingomonas* were

significantly affected by host age and initial microbial environment, host genotype still

significantly mediated these environmental and phenological changes in microbe abundance. Our

results add to recent evidence that the assembly process of host-associated microbiomes is

governed by both stochastic forces and host-based selective forces [87, 88].

Conclusions

Our results show that while *I. hederacea* leaf microbe composition differences were

primarily shaped by host age and the environment, there exists a heritable subset of core

microbes in the *I. hederacea* microbiome. A Mendelian trait and quantitative genetic variation

across matrilines explained similar amounts of variation in microbial abundance, with

implications for plant-microbe eco-evolutionary dynamics across time. Furthermore, we show

for the first time that leaf shape itself may contribute to differences in phyllosphere microbial

abundances within a species.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

Funding sources include NSERC graduate funding to JAB and NSERC discovery grants to MEF

and JRS.

Authors' contributions

JAB, MEF, and JRS collaborated on the ideas; JAB, MEF, and JRS designed the methodology,
JAB performed the experiments, and JAB, MEF, and JRS wrote and edited the manuscript. JAB,
MEF, and JRS analyzed the data. All authors contributed to this manuscript and approved its
publication.

Acknowledgements

We thank Brandon Campitelli for generating the morning glory lines and the staff of Koffler

Scientific Reserve in 2021 for helping to maintain the field site. We acknowledge Génome

Québec (Montréal, Canada) for sequencing samples. Finally, we gratefully acknowledge the

Swedish Collegium for Advanced Study, and Jennifer Mack, Natuschka Lee, Iva Lučić, and

Arthur Asseraf, for support.

References

1. Leveau JH. A brief from the leaf: latest research to inform our understanding of the

phyllosphere microbiome. Curr Opin Microbiol. 2019;49:41–9.

2. Vokou D, Vareli K, Zarali E, Karamanoli K, Constantinidou HIA, Monokrousos N, et al.

Exploring Biodiversity in the Bacterial Community of the Mediterranean Phyllosphere and its

Relationship with Airborne Bacteria. Microb Ecol. 2012;64:714–24.

3. Lindow SE. Role of Immigration and Other Processes in Determining Epiphytic Bacterial

Populations. In: Morris CE, Nicot PC, Nguyen-The C, editors. Aerial Plant Surface

Microbiology. Boston, MA: Springer, Boston, MA; 1996. p. 155–68.

4. Massoni J, Bortfeld-Miller M, Widmer A, Vorholt JA. Capacity of soil bacteria to reach the

phyllosphere and convergence of floral communities despite soil microbiota variation. Proc Natl

Acad Sci U S A. 2021;118:e2100150118.

5. Dastogeer KMG, Tumpa FH, Sultana A, Akter MA, Chakraborty A. Plant microbiome–an

account of the factors that shape community composition and diversity. Curr Plant Biol.

2020;23:100161.

6. Redford AJ, Bowers RM, Knight R, Linhart Y, Fierer N. The ecology of the phyllosphere:

geographic and phylogenetic variability in the distribution of bacteria on tree leaves. Environ

Microbiol. 2010;12:2885–93.

7. Kembel SW, O'Connor TK, Arnold HK, Hubbell SP, Wright SJ, Green JL. Relationships

between phyllosphere bacterial communities and plant functional traits in a neotropical forest.

Proc Natl Acad Sci. 2014;111:13715–20.

8. Bodenhausen N, Horton MW, Bergelson J. Bacterial Communities Associated with the Leaves

and the Roots of Arabidopsis thaliana. PLoS One. 2013;8:e56329.

9. Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD, Subramanian S, et al.

Genome-wide association study of Arabidopsis thaliana leaf microbial community. Nat

Commun. 2014;5:5320.

10. Wagner MR, Lundberg DS, Del Rio TG, Tringe SG, Dangl JL, Mitchell-Olds T. Host

genotype and age shape the leaf and root microbiomes of a wild perennial plant. Nat Commun.

2016;7:12151.

11. Wagner MR, Busby PE, Balint☐Kurti P. Analysis of leaf microbiome composition of

near☐isogenic maize lines differing in broad☐spectrum disease resistance. New Phytol.

2019;225:2152–65.

12. Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, et al. Diversity and heritability of

the maize rhizosphere microbiome under field conditions. Proc Natl Acad Sci U S A.

2013;110:6548–53.

13. Deng S, Caddell DF, Xu G, Dahlen L, Washington L, Yang J, et al. Genome wide

association study reveals plant loci controlling heritability of the rhizosphere microbiome. ISME

J 2021 1511. 2021;15:3181–94.

14. Grieneisen L, Dasari M, Gould TJ, Björk JR, Grenier JC, Yotova V, et al. Gut microbiome

heritability is nearly universal but environmentally contingent. Science (80-). 2021;373:181–6.

15. Wagner MR, Tang C, Salvato F, Clouse KM, Bartlett A, Vintila S, et al. Microbe-dependent

heterosis in maize. Proc Natl Acad Sci. 2021;118:e2021965118.

16. Walters WA, Jin Z, Youngblut N, Wallace JG, Sutter J, Zhang W, et al. Large-scale

replicated field study of maize rhizosphere identifies heritable microbes. Proc Natl Acad Sci U S

A. 2018;115:7368–73.

17. Henry LP, Bruijning M, Forsberg SKG, Ayroles JF. The microbiome extends host

evolutionary potential. Nat Commun. 2021;12:5141.

18. Mueller UG, Linksvayer TA. Microbiome breeding: conceptual and practical issues. Trends

Microbiol. 2022;30:997–1011.

19. Wagner MR, Lundberg DS, Coleman-Derr D, Tringe SG, Dangl JL, Mitchell-Olds T. Natural

soil microbes alter flowering phenology and the intensity of selection on flowering time in a wild

Arabidopsis relative. Ecol Lett. 2014;17:717–26.

20. Petersen C, Hamerich IK, Adair KL, Griem-Krey H, Torres Oliva M, Hoeppner MP, et al.

Host and microbiome jointly contribute to environmental adaptation. ISME J. 2023;17:1953–65.

21. Pogoda CS, Keepers KG, Reinert S, Talukder ZI, Smart BC, Attia Z, et al. Heritable

differences in abundance of bacterial rhizosphere taxa are correlated with fungal necrotrophic

pathogen resistance. Mol Ecol. 2023;00:1–14.

22. Baltrus DA. Adaptation, specialization, and coevolution within phytobiomes. Curr Opin

Plant Biol. 2017;38:109–16.

23. Koskella B, Bergelson J. The study of host–microbiome (co)evolution across levels of

selection. Philos Trans R Soc B. 2020;375:20190604.

24. Johnson MTJ, Stinchcombe JR. An emerging synthesis between community ecology and

evolutionary biology. Trends Ecol Evol. 2007;22:250–7.

25. VanWallendael A, Benucci GMN, da Costa PB, Fraser L, Sreedasyam A, Fritschi F, et al.

Host genotype controls ecological change in the leaf fungal microbiome. PLOS Biol.

2022;20:e3001681.

26. Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. A Synthetic Community

Approach Reveals Plant Genotypes Affecting the Phyllosphere Microbiota. PLOS Genet.

2014;10:e1004283.

27. Pincebourde S, Woods HA. Climate uncertainty on leaf surfaces: the biophysics of leaf

microclimates and their consequences for leaf-dwelling organisms. Funct Ecol. 2012;26:844–53.

28. Runkle E. The Boundary Layer and Its Importance. Greenhouse Product News. 2016;:34.

29. Vacher C, Hampe A, Porté AJ, Sauer U, Compant S, Morris CE. The Phyllosphere:

Microbial Jungle at the Plant–Climate Interface. Annu Rev Ecol Evol Syst. 2016;47:1–24.

30. Núñez A, García AM, Moreno DA, Guantes R. Seasonal changes dominate long-term

variability of the urban air microbiome across space and time. Environ Int. 2021;150:106423.

31. Humphrey PT, Nguyen TT, Villalobos MM, Whiteman NK. Diversity and abundance of

phyllosphere bacteria are linked to insect herbivory. Mol Ecol. 2014;23:1497–515.

32. Meyer KM, Porch R, Muscettola IE, Luisa A, Vasconcelos S, Sherman JK, et al. Plant

neighborhood shapes diversity and reduces interspecific variation of the phyllosphere

microbiome. ISME J 2022. 2022;:1–12.

33. Nicotra AB, Leigh A, Boyce CK, Jones CS, Niklas KJ, Royer DL, et al. The evolution and

functional significance of leaf shape in the angiosperms. Functional Plant Biology. 2011;38:535–

52.

34. Givnish TJ. Comparative Studies Of Leaf Form: Assessing The Relative Roles Of Selective

Pressures And Phylogenetic Constraints. New Phytol. 1987;106:131–60.

35. Yates MJ, Verboom GA, Rebelo AG, Cramer MD. Ecophysiological significance of leaf size

variation in Proteaceae from the Cape Floristic Region. Funct Ecol. 2010;24:485–92.

36. Elmore CD. Mode of Reproduction and Inheritance of Leaf Shape in Ipomoea hederacea .

Weed Sci. 1986;34:391–5.

37. Bright KL, Rausher MD. Natural selection on a leaf-shape polymorphism in the ivyleaf

morning glory (Ipomoea hederacea). Evolution (N Y). 2008;62:1978–90.

38. Campitelli BE, Stinchcombe JR. Natural selection maintains a single-locus leaf shape cline in

Ivyleaf morning glory, Ipomoea hederacea. Mol Ecol. 2013;22:552–64.

39. Campitelli BE, Stinchcombe JR. Testing potential selective agents acting on leaf shape in

Ipomoea hederacea: Predictions based on an adaptive leaf shape cline. Ecol Evol. 2013;3:2409–

23.

40. Campitelli BE, Gorton AJ, Ostevik KL, Stinchcombe JR. The effect of leaf shape on the

thermoregulation and frost tolerance of an annual vine, Ipomoea hederacea (Convolvulaceae).

Am J Bot. 2013;100:2175–82.

41. Hull-Sanders HM, Eubanks MD, Carr DE. Inbreeding depression and selfing rate of Ipomoea

hederacea var. integriuscula (Convolvulaceae). Am J Bot. 2005;92:1871–7.

42. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.

Nat Methods. 2012;9:671–5.

43. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.

Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat

Biotechnol 2019 378. 2019;37:852–7.

44. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-

resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13:581–3.

45. McDonald D, Price MN, Goodrich J, Nawrocki EP, Desantis TZ, Probst A, et al. An

improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of

bacteria and archaea. ISME J . 2011;6:610–8.

46. Katoh K, Misawa K, Kuma KI, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30:3059–66.

47. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for

Large Alignments. PLoS One. 2010;5:e9490.

48. R Core Team. R: A language and environment for statistical computing. 2022.

49. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the

Tidyverse. J Open Source Softw. 2019;4:1686.

50. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed

Effects Models . J Stat Softw. 2017;82:1–26.

51. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. PLoS One. 2013;8:e61217.

52. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, et al. Package

"vegan" Version 2.5-7. 2020.

53. Lahti L, Shetty S. microbiome R package. 2019.

https://bioconductor.org/packages/release/bioc/html/microbiome.html. Accessed 3 Apr 2023.

54. Fox J, Weisberg Sanford. An R Companion to Applied Regression. 3rd edition. Sage

Publications; 2019.

55. Wilson AJ. Why h2 does not always equal Va / Vp□? J Evol Biol. 2008;21:647–50.

56. Bruijning M, Ayroles JF, Henry LP, Koskella B, Meyer KM, Metcalf CJE. Relative

abundance data can misrepresent heritability of the microbiome. Microbiome. 2023;11:222.

57. Buse A. The Likelihood Ratio, Wald, and Lagrange Multiplier Tests: An Expository Note.

Am Stat. 1982;36:153–7.

58. Friesen ML, Porter SS, Stark SC, Von Wettberg EJ, Sachs JL, Martinez-Romero E.

Microbially Mediated Plant Functional Traits. Annu Rev Ecol Evol Syst. 2011;42:23–46.

59. Falconer D, Mackay T. Introduction to Quantitative Genetics. 4th edition. England: Prentice

Hall, Harlow; 1996.

60. Schweitzer JA, Bailey JK, Fischer DG, LeRoy CJ, Lonsdorf E V., Whitham TG, et al. Plant–

Soil–Microorganism Interactions: Heritable Relationship Between Plant Genotype And

Associated Soil Microorganisms. Ecology. 2008;89:773–81.

61. Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, et al. Human Genetics

Shape the Gut Microbiome. Cell. 2014;159:789–99.

62. Wallace JG, Kremling KA, Kovar LL, Buckler ES. Quantitative genetics of the maize leaf

microbiome. Phytobiomes J. 2018;2:208–24.

63. John Wallace R, Sasson G, Garnsworthy PC, Tapio I, Gregson E, Bani P, et al. A heritable

subset of the core rumen microbiome dictates dairy cow productivity and emissions. Sci Adv.

2019;5:8391–4.

64. Edwards JA, Saran UB, Bonnette J, MacQueen A, Yin J, Nguyen T uyen, et al. Genetic

determinants of switchgrass-root-associated microbiota in field sites spanning its natural range.

Curr Biol. 2023;33:1926-1938.e6.

65. Cavanagh C, Morell M, Mackay I, Powell W. From mutations to MAGIC: resources for gene

discovery, validation and delivery in crop plants. Curr Opin Plant Biol. 2008;11:215–21.

66. Yu J, Holland JB, McMullen MD, Buckler ES. Genetic Design and Statistical Power of

Nested Association Mapping in Maize. Genetics. 2008;178:539–51.

67. Campitelli BE, Stinchcombe JR. Population dynamics and evolutionary history of the weedy

vine Ipomoea hederacea in North America. G3 Genes, Genomes, Genet. 2014;4:1407–16.

68. Stock AJ, Campitelli BE, Stinchcombe JR. Quantitative genetic variance and multivariate

clines in the Ivyleaf morning glory, Ipomoea hederacea. Philos Trans R Soc B Biol Sci.

2014;369:20130259.

69. Henry GA, Stinchcombe JR. G-matrix stability in clinally diverging populations of an annual

weed. Evolution (N Y). 2023;77:49–62.

70. Ennos RA. Quantitative studies of the mating system in two sympatric species of Ipomoea

(Convolvulaceae). Genetica. 1981;57:93–8.

71. Weinig C, Dorn LA, Kane NC, German ZM, Halldorsdottir SS, Ungerer MC, et al.

Heterogeneous Selection at Specific Loci in Natural Environments in *Arabidopsis thaliana*.

Genetics. 2003;165:321–9.

72. Weinig C, Stinchcombe JR, Schmitt J. Evolutionary Genetics Of Resistance And Tolerance

To Natural Herbivory In Arabidopsis thaliana. Evolution (N Y). 2003;57:1270–80.

73. Vasemägi A. The Adaptive Hypothesis of Clinal Variation Revisited: Single-Locus Clines

as a Result of Spatially Restricted Gene Flow. Genetics. 2006;173:2411–4.

74. Lippincott JA, Lippincott BB. Tumour-initiating Ability and Nutrition in the Genus

Agrobacterium. J Gen Microbiol. 1969;59:57–75.

75. Lee HS, Madhaiyan M, Kim CW, Choi SJ, Chung KY, Sa TM. Physiological enhancement

of early growth of rice seedlings (Oryza sativa L.) by production of phytohormone of N2-fixing

methylotrophic isolates. Biol Fertil Soils. 2006;42:402–8.

76. Wang NR, Wiesmann CL, Melnyk RA, Hossain SS, Chi M-H, Martens K, et al. Commensal

Pseudomonas fluorescens Strains Protect Arabidopsis from Closely Related Pseudomonas

Pathogens in a Colonization-Dependent Manner. MBio. 2022;13:e02892-21.

77. Kämpfer P, Glaeser SP, McInroy JA, Busse H-J. Nocardioides zeicaulis sp. nov., an

endophyte actinobacterium of maize. Int J Syst Evol Microbiol. 2016;66:1869–74.

78. Wang P, Kong X, Chen H, Xiao Y, Liu H, Li X, et al. Exploration of Intrinsic Microbial

Community Modulators in the Rice Endosphere Indicates a Key Role of Distinct Bacterial Taxa

Across Different Cultivars. Front Microbiol. 2021;12:629852.

79. Nafis A, Raklami A, Bechtaoui N, El Khalloufi F, El Alaoui A, Glick BR, et al.

Actinobacteria from Extreme Niches in Morocco and Their Plant Growth-Promoting Potentials.

Diversity. 2019;11:139.

80. Dastager SG, Lee J-C, Pandey A, Kim C-J. Nocardioides mesophilus sp. nov., isolated from

soil. Int J Syst Evol Microbiol. 2010;60:2288–92.

81. Escudero A, Mediavilla S. Decline in photosynthetic nitrogen use efficiency with leaf age

and nitrogen resorption as determinants of leaf life span. J Ecol. 2003;91:880–9.

82. England JR, Attiwill PM. Changes in leaf morphology and anatomy with tree age and height

in the broadleaved evergreen species, Eucalyptus regnans F. Muell. Trees. 2006;20:79.

83. Pantin F, Simonneau T, Muller B. Coming of leaf age: control of growth by hydraulics and

metabolics during leaf ontogeny. New Phytol. 2012;196:349–66.

84. van Butselaar T, Van den Ackerveken G. Salicylic Acid Steers the Growth–Immunity

Tradeoff. Trends Plant Sci. 2020;25:566–76.

85. Redford AJ, Fierer N. Bacterial succession on the leaf surface: a novel system for studying

successional dynamics. Microb Ecol. 2009;58:189–98.

86. Copeland JK, Yuan L, Layeghifard M, Wang PW, Guttman DS. Seasonal Community

Succession of the Phyllosphere Microbiome. Mol Plant-Microbe Interact. 2015;28:274–85.

87. Dove NC, Veach AM, Muchero W, Wahl T, Stegen JC, Schadt CW, et al. Assembly of the

Populus Microbiome Is Temporally Dynamic and Determined by Selective and Stochastic

Factors . mSphere. 2021;6.

88. Xu CCY, Lemoine J, Albert A, Whirter É Mac, Barrett RDH. Community assembly of the

human piercing microbiome. Proc R Soc B Biol Sci. 2023;290.

688