1   **Czech Republic butterfly barcoding reveals that distribution of genetic lineages depends**
2   **on species traits**
3
4
5   **Running title:**
6   Czech butterfly DNA barcoding
7
8

9   Alena SUCHÁČKOVÁ BARTOŇOVÁ[1*], Patrik ŠKOPEK[3], Martin KONVIČKA[1,2], Jiří
10 BENEŠ[1], Lukáš SPITZER[1,4], Claudio SBARAGLIA[1,2], Vladimír VRABEC[3], Jana PAPP
11 MAREŠOVÁ[1,2], Hana KONVIČKOVÁ[1,2] & Zdeněk FALTÝNEK FRIC[1,3]
12
13
[1] Biology Centre of the Czech Academy of Sciences, Institute of Entomology, Branišovská
31, 370 05 České Budějovice, Czech Republic
[2] University of South Bohemia in České Budějovice, Faculty of Science, Branišovská 1760,
370 05 České Budějovice, Czech Republic
[3] Czech University of Life Sciences Prague, Faculty of Agrobiology, Food and Natural
Resources, Kamýcká 129, 165 00 Praha – Suchdol, Czech Republic
[4] Museum of the Moravian Wallachia Region, Horní náměstí 2, 755 01 Vsetín, Czech
Republic
22
23 *Correspondence: Alena Sucháčková Bartoňová, al.bartonova@gmail.com, +420731856818

24
25
26

27 **ACKNOWLEDGEMENTS**
28
The work was supported by the Technology Agency of the Czech Republic (SS01010526).
We would like to thank the Nature Conservation Agency of the Czech Republic and Ministry
of the Environment of the Czech Republic, as well as all the local conservation authorities
(National Parks and Protected Landscape Areas). The following colleagues helped us with
material: Oldřich Čížek, Marek Fišer, Vladimír Hula, Tomáš Kadlec, František Kopeček,
Tomáš Kuras, Dan Leština, Zdeněk Mráček, Alois Pavlíčko, David Ričl, Ondřej Sedláček,
Pavel Skala, Přemysl Tájek, Pavel Vrba, and Jan Walter. We would also like to thank Petr
Šmilauer for consultation on statistics, Pedro de G. Ribeiro and Marie Drábková for
comments on the manuscripts, and Matthew Sweney for English correction.

ABSTRACT

Aim: The distribution of within-species lineages has been affected by Quaternary climate changes, and population differentiation has been influenced by species life histories. We investigated whether the distribution of individual mitochondrial genetic lineages reflects the constituent species' traits. Using the functionally diverse group of butterflies, we examined which lineages are present in Central Europe, an important suture zone.

Location: Czech Republic and Western Palearctic.

Taxon: A total of 140 butterfly species.

Methods: We sequenced DNA barcodes (cytochrome c oxidase 1) (959 sequences) of the entire Czech Republic butterfly fauna and used BOLD data to visualize the species' biogeographic patterns across Europe. We categorised the distribution patterns of lineages inhabiting the Czech Republic, and used multivariate statistics to interpret these categories by the butterflies' habitats, life histories, and threat levels.

Results: Open habitat dwellers with specialist traits belonged to Eastern, Southern, and temperate lineages. Habitat generalists and woodland dwellers belonged to the Western lineage, formed several lineages, or displayed low genetic diversity; they often developed on woody plants, were large-winged, and had long flight periods. The most threatened species were the specialists of Southern, Eastern, and temperate lineages.

Main conclusions: The distribution of lineages in Central Europe reflects the history of Quaternary ecosystems: during cold periods of the Pleistocene, the diverse open habitats prevailed, and species could expand westwards. Such species also suffer the most under the current anthropogenic habitat alteration. On the other hand, the mobile generalists and woodland dwellers expanded to Central Europe during the Holocene. Our approach of linking the distribution of lineages with species traits can be transferred to other study systems, and we show that DNA barcoding of under-sampled areas represents a powerful tool for discovering the driving forces of biogeography.

KEYWORDS

Barcoding library, COI, insects, Lepidoptera, life history, mitochondrial DNA, phylogeography, refugia theory

INTRODUCTION

Population genetic differentiation is affected by species' range sizes, life histories, and varies across geographic regions (Dapporto et al., 2019; Gamba & Muchhala, 2020; Miller et al., 2021). In temperate regions, the distribution of different genetic lineages, i.e., the groups of sequences with shared history, has been massively influenced by Quaternary climate change (Hewitt, 2000; 2004; Schmitt, 2007; Hofreiter & Stewart, 2009). Thus, a population inhabiting an area might be the result of combining the landscape history with trait-mediated divergence.

Under the Quaternary's dynamic conditions, Europe's unique geographic idiosyncrasy affected the distribution of genetic lineages. In Europe, there are three peninsulas of a limited area with a Mediterranean climate functioning as speciation centres, east-west oriented mountain ridges acting as barriers, and a gradient between oceanic climate in the west and continental climate in the east. These factors influenced species' ranges: during the shorter warm stages of the Quaternary, warm-adapted species expanded from Mediterranean peninsulas or other spatially restricted areas (refugia), while cold-adapted species retreated uphill and northwards (de Lattin, 1967; Hewitt, 1996, 1999; Schmitt, 2007; Schmitt & Varga, 2012). Simultaneously, continental species tended to retreat eastwards (Stewart, Lister, Barnes, & Dalén, 2010). Notably, Central Europe represents a crossroad of different lineages (Janoušek et al., 2012; Pfäffle, Bolfíková, Hulva, & Petney, 2014; Nürnberger, Lohse, Fijarczyk, Szymura, & Blaxter, 2016). This area went through immense changes during the Quaternary, with an exchange of biota between the cold and warm stages (Horáček & Ložek, 1988), while the landscape varied from a cold steppe-tundra to open park-like woodlands (Kahlke, 2014; Sandom, Ejrnæs, Hansen, & Svenning, 2014; Vera, 2000; Pearce et al., 2023).

A feasible method to evaluate genetic differentiation and lineage distribution on large-scale data is DNA barcoding, i.e., sequencing a short standardized gene fragment, which originally aimed for a simple tool for species identification and discovery (Hebert, Cywinska, Ball, & deWaard, 2003). An important step towards this goal was building well-curated world-wide databases covering as many species as possible, such as the Barcoding of Life Data System (BOLD; Ratnasingham & Hebert 2007). The DNA barcode widely used in various animal groups, mitochondrial gene cytochrome c oxidase subunit I (COI), proved effective not only for species identification (Hebert, Ratnasingham, & de Waard, 2003), but also for uncovering cryptic diversity including discovery of new species (Hernández-Roldán et al., 2016), and phylogeographic and population genetic studies (Kühne, Kosuch, Hochkirch, & Schmitt, 2017; Maresova et al., 2021). The existence of large database data allowed testing the DNA barcoding performance across large scales on entire faunal groups, summarizing genetic diversity and uncovering biogeographic patterns (e.g., Meier et al., 2006; Geiger et al., 2014; Hendrich et al., 2015; Weigand et al., 2019; Dincă et al., 2021; D'Ercole et al., 2021; Galimberti et al., 2021; Dapporto et al., 2022).

European butterflies are one of the best DNA barcoded groups of organisms, represented by several national and international DNA barcoding libraries (Dincă, Zakharov, Hebert, & Vila, 2011; Dincă et al., 2015, 2021; Hausmann et al., 2011; Huemer, Mutanen, Sefc, & Hebert, 2014; Huemer et al., 2018; Litman et al., 2018; Dapporto et al., 2022). Geographic patterns of diversity and differentiation are extraordinarily well-explored, especially for south-western Europe (Dapporto et al., 2019, 2022; Dincă et al., 2021).

In this study, we focused on the functionally diverse butterfly fauna of the Czech Republic and provide the DNA barcoding library for the country. We view the country, situated in Central Europe, as a sieve of species that either reached the area during the Holocene, or survived there since the last glaciation. The Czech butterfly fauna consists of both generalists and specialists of diverse habitats, dry and wet, lowland and highland, and of

different stages of habitat openness, whose life histories are known in considerable detail
(Bartonova, Benes, & Konvicka, 2014; Macek, Laštůvka, Beneš, & Traxler, 2015), and at the
same time have one of the highest threat values in Europe (Warren et al., 2021). We DNA
barcoded samples of Czech populations of 140 species (~98% of extant Czech butterfly
fauna), accessed BOLD data for these species across their ranges, analysed their
mitochondrial phylogeographic structure, and related them to species-specific traits. We
predict that lineages do not inhabit the region randomly, but do so with respect to individual
species' habitat affinities and life histories. In particular, we expect differences between
generalist and specialist species, and between species associated with closed (woodland) and
open (grasslands, steppe, wetlands) habitats. Moreover, the lineage presence can be related to
the species' current threat levels, thus, the genetic backgrounds can predetermine individual
species' future.
138
139 **MATERIAL AND METHODS**
140
141 **Sampling and DNA sequencing**
142
Czech butterfly fauna consists of 141 species including migrants, plus six irregular vagrants,
and a further 16 species that are considered extinct (Benes et al., 2002; Wiemers et al., 2018).
We collated samples from 140 butterfly species originating from the Czech Republic (CZ)
(Figure 1a), from the years 2004–2022 (Supplementary Materials: figshare repository DOI
10.6084/m9.figshare.24559852). The dataset comprises all the extant species except for
Melitaea phoebe (recent colonizer), *Parnassius apollo* (existing as a conservation-dependent
reintroduced population), and *Hyponephele lycaon* (recently extinct in the country), but
including *Pieris mannii*, which newly recolonized the country, and a rare vagrant, *Leptotes*
pirithous. We also sequenced one specimen of *Lampides boeticus*, imported with fruit from
Spain (not included in analyses).
We extracted DNA from the butterfly legs using Genomic DNA Mini Kit (Tissue)
(Geneaid Biotech Ltd.). We amplified the barcode, first part of COI, using primers hybLCO
(5ʹ-TAATACGACTCACTATAGGGGGTCAACAAATCATAAAGATATTGG-3ʹ) and
hybHCO (5ʹ-ATTAACCCTCACTAAAGGGTAAACTTCAGGGTGACCAAAAAATCA-3ʹ)
(Folmer, Black, Hoeh, Lutz, & Vrijenhoek, 1994). In the case of fragmented DNA, we used
two pairs of primers: [LCO-K699] + [Ron-HCO] (K699: 5ʹ-
ATTAACCCTCACTAAAGGGWGGGGGGGTAAACTGTTCATCC-3ʹ; Ron: 5ʹ-
TAATACGACTCACTATAGGGGGATCACCTGATATAGCATTCCC-3ʹ)
(www.nymphalidae.net; Monteiro & Pierce 2001). The mixture for each sample consisted of
4 µl of PCR H$_2$O, 0.625 µl of each primer, 6.25 µl of Bioline 2x MyTaq HS Red Mix
(Meridian Bioscience Inc.), and 2 µl of extracted DNA. The polymerase chain reaction
protocol consisted of 95°C for 5 min; followed by 40 cycles of 94°C for 30 s, 50°C for 30 s,
and 72°C for 90 s; with a final extension of 72°C for 10 min. PCR products were cleaned
using FastAP and Exo I enzymes (Thermo Fisher Scientific). Sanger sequencing was
performed in Macrogen Inc. (Amsterdam, Netherlands) on 3730xl DNA Analyzer, in one
direction from the 5ʹ end. The sequences were checked, trimmed, and aligned in Geneious v.
8.0.5 (Kearse et al., 2012). We downloaded all COI sequences for each species occurring in
the Czech Republic from the BOLD database (Ratnasingham & Hebert, 2007) (for 12 October
2022) which were georeferenced and sequenced from the 5ʹ end. We aligned the samples
together with the newly sequenced data.
We analysed each species dataset separately. Members of species pairs known to share
haplotypes or not monophyletic in the DNA barcode (Dincă et al., 2011) (specifically *Colias*
crocea/erate, *Cupido decoloratus/alcetas*, *Erebia euryale/ligea*, *Pieris napi/bryoniae*,

Polyommatus bellargus/coridon, *Pseudophilotes baton/vicrama*) entered the analyses as
separate species because of their unique species traits.
178
179 **Phylogeographic patterns**
180
For each species, we performed two phylogeographic analyses: (1) Bayesian Analysis of
Population Structure (BAPS) (Cheng, Connor, Sirén, Aanensen, & Corander, 2013), which
sorts data into clusters of related sequences (*genetic lineages*), in R package 'rhierbaps'
(Tonkin-Hill, Lees, Bentley, Frost, & Corander, 2018), on level 1, automatically estimating
the number of clusters; and (2) TCS haplotype networks in the program POPART (Leigh &
Bryant, 2015), a parsimony method which, additionally, shows putative evolutionary
relationships among samples. In both cases, we checked the results visually and cleaned the
dataset as to omit database samples which were either shorter than 500 bp, or potentially
misidentified (dissimilarity of sequences represented by long branches in haplotype networks
was evaluated using the BLAST algorithm, Altschul et al., 1990, in the NCBI database).
We plotted the distribution of BAPS clusters on a map of Europe using the geographic
coordinates of the samples. Then, we manually assigned the lineages present in the Czech
Republic to one of the following categories, or to two, if two specific lineages were present:
(1) *BAPS-low*: low genetic diversity – only a single BAPS cluster in the entire species; (2)
BAPS-wide: a lineage widely distributed across Europe also present in CZ (but more lineages
are present within the species); (3) *BAPS-mix*: two or more widespread European lineages are
present in CZ; (4) *BAPS-West*: the CZ samples related to samples from Western Europe; (5)
BAPS-Italy-Balkans: the CZ samples related to those from Italy or Italy plus the Balkans; (6)
BAPS-East-Balkans: the CZ samples are related to those from the Balkans or Balkans plus
Eastern Europe/Asia; (7) *BAPS-mix-East*: CZ samples consist of more lineages distributed in
Italy/Balkans or East/Balkans; (8) *BAPS-Central*: the Central European mountains or
Pannonian lowlands produced a separate lineage to which CZ samples belong; and (9) *BAPS-*
temperate: the CZ samples are related to lineages inhabiting a similar latitudinal belt, not
present in the Mediterranean peninsulas.
The results of haplotype networks were scored based on the geographic area inhabited
by haplotypes found in CZ and haplotypes distant from them with maximum of two
mutations. When the Czech samples were part of a widespread lineage, the supposed
demographic scenario from the net was visually estimated. The categories were: (1) *HAP-low*:
lack of differentiation (either low diversity or expansion - star-like haplotype structure) in the
entire species. Then, in categories (2)–(4), the Czech samples are part of a widespread
European lineage (another, diverged lineage exists in the network): (2) *HAP-wide-low*: CZ
samples belong to a widespread European lineage with low haplotype diversity (supposed
bottleneck; maximum of 10 closely related haplotypes); (3) *HAP-wide-expand*: part of
expanding widespread European lineage (star-like structure, with eight and more haplotypes
diverging from the main one); (4) *HAP-wide-complex*: part of a complex widespread lineage
(showing either a stable network with several connections and even sequences in haplotypes,
or combination of a stable network and expansion. (5) *HAP-mix*: divergent haplotypes were
present in CZ with distance >2 mutations. The following categories were defined similarly to
BAPS classification: (6) *HAP-West*; (7) *HAP-Italy-Balkans*; (8) *HAP-East-Balkans*; (9)
HAP-Central; and (10) *HAP-temperate.*
221
222 **Species traits**
223
We prepared data on the following three groups of species traits: (1) Habitat affinity, defining
nine categories: *ubiquitous* (generalist), *mesophilic 1* (grassland), *mesophilic 2* (shrubland),

mesophilic 3 (woodland), *xerothermophilic 1* (steppe), *xerothermophilic 2* (dry shrubland),
hydrophilic (wetland), *tyrphophilic* (bog/peatland), and *alpine* (Benes et al., 2002). If two
categories applied for a species, these were assigned as "0.5" per category.
(2) Life history traits linked to dispersal and landscape scale survival (Bartonova,
Benes, Fric, Chobot, & Konvicka, 2016): *density*, the number of individuals which can occur
per area of habitat (adapted from area demand) [ranked 1–9, from sparse to dense] (Reinhardt,
Sbieschne, Settele, Fischer, & Fiedler, 2007); *feeding index*, the trophic range, defined as an
index that weights the number of consumed host plant families [F] and genera [G] in the
Czech Republic: $(G*F^a)^{1/2}$ where a=F/2G (Benes et al., 2002; index modified after Garcia-
Barros 2000); *fertility*, the number of eggs per female at eclosion [ranked 1–9] (Reinhardt et
al., 2007); *flight period length*, the number of months of adult flight, summed across
generations and excluding hibernation (Benes et al., 2002); *forewing length,* approximation
for body size [mm] (Higgins & Riley, 1970); *host plant form*, express prevailing host plant
apparency [ranked 1–4, small forbs, large forbs/grasses, bushes, trees] (Cizek, Fric, &
Konvicka, 2006); *mobility*, the propensity to disperse [ranked 1–9, from sedentary to mobile]
(Reinhardt et al., 2007); *overwintering stage* [ranked 1–5, from egg to migrating adult]
(Tolman & Lewington, 2008); and *voltinism*, average number of generations per year in the
Czech Republic (Benes et al., 2002). In the case of species not covered in Reinhardt et al.
(2007) (N=39), missing values of density and fertility were substituted by averages, and
missing values of mobility were scored based on authors' expertise (similarly to Bartonova et
al.,2016).
(3) Threat status, based on national Red List categories (ordinal scale: 1 – least
concern, 2 – near threatened, 3 – vulnerable, 4 – endangered, and 5 – critically endangered)
(Hejda, Farkač, & Chobot, 2017).
250
251   **Relating phylogeographic patterns to species traits**
252
To investigate the relationships between phylogeographic patterns and traits, we used a
multivariate approach, the unimodal canonical correspondence analyses (CCA) in CANOCO
v.5.0 (Ter Braak & Šmilauer, 2012), which extracts variation in multivariate dependent
variables even if obscured by other larger sources of variation (Ter Braak & Verdonschot,
1995), and tests the significance of the ordinations via the Monte-Carlo permutation tests (999
runs). In these analyses, individual species were the samples, either BAPS or haplotype
network categories were responses, and species traits were predictors. The response variables
were categorical as they identified the group to which a sample (butterfly species) belong.
Since response variables could have been represented by more categories, they entered the
analysis in the form of dummy variables with fuzzy coding (0, 0.5 or 1), and CCA in this case
represented a generalization of linear discriminant analysis (LDA), optimal for categorical
variables, with chi-square distances as generalization of Mahalanobis distances among
observations (Ter Braak & Verdonschot, 1995; Ter Braak & Šmilauer, 2012). CANOCO v.5.0
automatically applies standardization to unit variance (bringing their means to zero and
variances to one) to control for the different dimensions of the predictors. In the case of
habitat affinities and life history traits, two tests were performed: (a) global test, using all
variables and (b) a forward selection procedure, picking a combination of traits that
significantly fit the phylogeographic patterns. Threat status, representing a single explanatory
variable, was also tested using CCA.
272
273   **RESULTS**
274
275   **Dataset**

276
The final dataset consisted of 1,110 sequences of the 140 species found in the Czech Republic
(Fig. 1a) (excluding *L. boeticus*); 151 sequences from BOLD and GenBank, and 959 newly
generated (mean=8±5.1 SE sequence per species, range 1–83), which can be found in BOLD
as project BBCZ. We added 20,695 sequences for these species from BOLD originating in
other locations. Together, we used 21,805 sequences (mean=156±86.5 SE sequence per
species, range 18–854) (Supplementary Materials). The number of species assigned to each
BAPS and haplotype network category are summarized in Fig. 1b and c, and the categories
assigned for each species are found in Supporting Information 1 and Supplementary Materials
. The BAPS category (Fig. 2) with the highest number of species was *BAPS-low* (N=36),
followed by *BAPS-wide* (N=285). The haplotype network category (Fig. 3) represented by
highest number of species was *HAP-temperate* (N=19.5), followed by *HAP-wide-complex*
(N=18), *HAP-wide-low* (N=17.5), and *HAP-mix* (N=17).
289
290   **Relating BAPS categories to species traits**
291
In the CCA relating BAPS categories to (1) habitat affinity, the global test revealed a
significant relationship (eigenvalues: 0.23, 0.14, 0.11, 0.09; first axis var. [explained
variation] =3.1%, F=4.2, p=0.046; all axes var.=9.0%, F=1.6, p=0.005). The forward selection
identified *ubiquitous* (i.e., generalist, var.=2.2%, F=3.1, p=0.001) and *xerothermophilic 1*
(i.e., steppe, var.=1.5%, pseudo-F=2.1, p=0.034) affinities (model's canonical eigenvalues:
0.18, 0.09; total var.=3.7%). The first canonical axis in the forward selection CCA
distinguished *BAPS-East-Balkans*, *BAPS-Italy-Balkans*, *BAPS-wide, BAPS-temperate*, and
BAPS-Central patterns from *BAPS-mix*, *BAPS-mix-East*, *BAPS-West*, and *BAPS-low* patterns,
and showed that the former tend to be steppe species, whereas the latter are habitat generalists
(Fig. 4a).
The CCA relating BAPS categories to (2) life history traits also produced significant
ordination (global test eigenvalues: 0.22, 0.20, 0.11, 0.07; first axis var. =3.1%, F=4.1,
p=0.066; all axes var.=9.0%, F=1.4, p=0.015). The forward-selected traits were *host plant*
form (var.=1.7%, F=2.3, p=0.014), *forewing length* (var.=1.6%, F=2.3, p=0.015), and
voltinism (var.=1.6%, F=2.3, p=0.017) (model's canonical eigenvalues: 0.19, 0.12, 0.05; total
var.=4.9%). The first canonical axis in the forward selection CCA separated species
belonging to categories *BAPS-low, BAPS-West, BAPS-mix*, and *BAPS-wide,* from those
belonging to *BAPS-Central, BAPS-East-Balkans, BAPS-temperate,* BAPS-*mix-East,* and
BAPS-Italy-Balkans, indicating that the former tend to develop on woody plants, have more
generations per year and a large wingspan; whereas the latter develop on herbs, have less
generations and a small wingspan (Fig. 4b).
The BAPS categories were also related to (3) threat status (model's canonical
eigenvalue: 0.13; var.=1.8%, F=2.5, p=0.014). The analysis indicated that species with pattern
BAPS-East-Balkans tend to be the most threatened, followed by *BAPS-Italy-Balkans,* and
BAPS-West; and those of *BAPS-mix* are the least endangered (Fig. 4c).
317
Relating haplotype network categories to species traits
The global test relating haplotype network categories to (1) habitat affinity was significant
(global test eigenvalues: 0.23, 0.19, 0.12, 0.06; first axis var.=2.7%, F=3.6, p=0.068; all axes
var.= 8.8%, F=1.4, p=0.012). Two significant habitat affinities were forward-selected:
ubiquitous (var.=2.2%, F=3.1, p=0.002) and *mesophilic 3* (*i.e.*, woodland; var.=2.0%, F=2.9,
p=0.001) (model's canonical eigenvalues: 0.21, 0.16; total var.=4.2%). The first canonical
axis in the forward selection CCA separated species of patterns *HAP-wide-expand, HAP-mix,*
HAP-West, HAP-wide-low, HAP-low, from *HAP-Central, HAP-East-Balkans, HAP-wide-*

complex, *HAP-Italy-Balkans*, and *HAP-temperate*, revealing that the former are habitat generalists and woodland species, whereas the latter are non-woodland specialists (Fig. 4d).

Explaining haplotype network categories by (2) life history traits was significant (global test eigenvalues: 0.35, 0.21, 0.12, 0.10; first axis var.=4.1%, F=5.5, p=0.001; all axes var.=11.2%, F=1.8, p=0.001). The significant forward-selected variables were *host plant form* (var.=2.3%, F=3.2, p=0.001), *flight period length* (var.=1.6%, F=2.3, p=0.02), and *feeding index* (var.=1.3%, F=1.9, p=0.042) (model's canonical eigenvalues: 0.25, 0.15, 0.05; total var.=5.2%). The forward-selected ordination separated patterns *HAP-wide-low*, *HAP-low*, *HAP-wide-expand*, and *HAP-West* from *HAP-Central*, *HAP-East-Balkans*, *HAP-Italy-Balkans*, *HAP-temperate*, *HAP-mix*, and *HAP-wide complex*. The former species have long flight periods, develop on tree or shrub host plans, and have a broader feeding niche, in contrast to the latter species (Fig. 4e).

The haplotype network categories were also explicable by (3) threat status (model's canonical eigenvalue: 0.14; var.=1.7%, F=2.3, p=0.008). Species with patterns *HAP-temperate* and *HAP-East-Balkans* tend to be the most threatened, whereas *HAP-wide-low*, *HAP-mix*, and *HAP-low* are the least threatened (Fig. 4f).

DISCUSSION

We DNA barcoded 140 species of butterflies occurring in the Czech Republic, used database data to visualize the species' biogeographic patterns across Europe, and related the mitochondrial genetic structure of Czech samples to species traits. We showed that species' habitat use and life histories influence which of the mitochondrial lineages inhabit a specific area, and these factors in turn influence their current threat status.

Several authors have related quantitative measures of overall species' genetic diversity and differentiation to species' range sizes, life histories and samples' geographic origins. Genetic diversity varies greatly in European butterflies, as measured by both mtDNA (Dincă et al., 2021) and genome-wide markers (Mackintosh et al., 2019). Mitochondrial COI haplotype diversity itself could not be explained by life history traits determining dispersal and colonization abilities (Dapporto et al., 2019). Generalists do not have greater levels of genomic diversity than specialists; only smaller butterflies tend to have higher genomic diversity (Mackintosh et al., 2019; but see Habel et al., 2013 on landscape level). Population differentiation, on the other hand, is explicable by life histories. Lower differentiation applies to species with longer flight periods and higher number of generations. Accordingly, a higher differentiation was discovered in species with smaller wings, utilising a lower number of host plants, and with short flight periods, i.e., in specialists (Habel et al., 2013; Dapporto et al., 2017, 2019; Scalercio et al., 2020). Widespread species also showed high population differentiation (Dapporto et al., 2019).

We use a more detailed approach here, explaining geographic distribution of individual genetic lineages by species' traits. The Czech Republic well represents temperate Europe as a whole, owing to its diverse butterfly fauna. We are aware that classification of the lineage distribution is tied to the target region. It is also influenced by the amount of available sequence data and their geographical coverage for a given species. The habitat affinities, and such life history traits as flight period length or host plant use, and of course the level of threat, may also vary across species' ranges (e.g., Korb et al., 2016; Lindestad et al., 2022). Changing the target region could reveal different lineages for species analysed here, in parallel with changing regional fauna composition and its life histories. Still, our approach offers a template for analysing the constitution of regional faunas.

Both BAPS and haplotype networks categories of lineages present in the Czech butterfly species revealed matching patterns (Fig. 4) when associated with habitats, and

analyses of life histories contributed to the understanding of the system (cf. Potocký et al.,
2018). In principle, generalists and woodlands dwellers displayed different genetic patterns
than specialists and open habitats species. Specifically, generalists and species of wooded
habitats formed mixed lineages, or displayed shallow genetic patterns, or had Western
affinities (Fig. 4 a, d). Such species frequently develop on shrubs or trees, form multiple
generations per year, and have long flight periods, long wings, and wide host plant ranges
(Fig. 4 b, e). Species of low genetic diversity (and hence low population differentiation in this
case) belong to this group, in agreement with Dapporto et al. (2019). In contrast, open habitats
specialists (i.e., steppe species) displayed Southern and/or Eastern affinities, or had temperate
patterns, or formed Central European genetic lineages. Such species tended to develop on
herbs and grasses and displayed such specialist traits as small wings, low numbers of
generations, short flight periods, and narrow ranges of host plants (Bartonova et al., 2014).
Such species were linked with high genetic differentiation in Dapporto et al. (2019).
The findings for open habitat species can be attributed to the Quaternary history of
biomes in Central Europe. During the mid to late Quaternary, the habitat prevailing in both
time and space was the cold mammoth steppe (Kahlke, 2014), associated with diverse types
of open habitats. Evidence points to *in situ* survival of steppe lineages in Central Europe
through the climatic cycles (Kajtoch et al., 2016; Kirschner et al., 2020; Sucháčková
Bartoňová et al., 2021) including the Holocene (Feurdean, Ruprecht, Molnár, Hutchinson, &
Hickler, 2018). That long-term persistence could have produced higher population
differentiation (Kajtoch, Kubisz, Gutowski, & Babik, 2014; Dapporto et al., 2019;
Hawlitschek et al. 2023). In some butterflies targeted here, we indeed observed the Central
European genetic patterns (e.g., one lineage in *Melanargia galathea* and *Pyrgus malvae*,
Polyommatus thersites).
The association of steppe butterflies with Southern and Eastern lineages is attributable
to the geography of the Czech Republic, with lowland corridors situated in the southeast and
east (e.g., in *Melitaea didyma*, *Polyommatus daphnis*, *Pyrgus carthami*). The Pannonian
lowlands, with natural vegetation formed by forest steppes, are connecting the Czech
Republic with the Balkan Peninsula. An alternative route is through the Moravian Gate,
connecting Czech territory with the lowlands of southern Poland, continuing to the Pontic
steppes (the "Sarmatian route"; Mařan 1958; Sternberg 1998). On the other hand, the warm
lowlands in Bohemia (Elbe river valley) were likely colonized by steppe elements from
Moravia, blocked by mountain chains from similar habitats in Western Europe. This scenario
was documented, e.g., for the ground squirrel *Spermophilus citellus* (Říčanová et al., 2013).
Multiple species had a genetic lineage distributed along a single latitudinal belt
(temperate patterns), lacking the Mediterranean distribution. These were species of open
habitats and habitat specialists. Many genetic lineages could have inhabited the Palearctic
continent alongside a temperature belt (Maresova et al., 2021), with no barriers to dispersal in
the east-west direction. These species might follow the east-west oceanic-continental gradient
of glacial-interglacial faunal interchange related to moisture (Stewart et al., 2010).
Continental species might have expanded west during the arid glacials, establishing stable
widespread populations, with Mediterranean peninsulas serving as areas of endemism rather
than refugia for them (e.g., in *Agriades optilete*, *Boloria euphrosyne*, *Glaucopsyche alexis*).
Woodlands re-expanded across Europe during the Holocene. Thus, most of the
populations of forest species are relatively newly established and shallow genetic structures,
which we observed for woodland inhabitants and tree feeders, are expected (e.g., in the
European woodland species *Pararge aegeria*, Livraghi et al., 2018; or in *Nymphalis
polychloros*, *Celastrina argiolus*). Habitat generalists display similar genetic patterns to the
woodland species, linked especially with a mix of lineages inhabiting the focal area, which
likely diversified in multiple refugia (e.g., *Pieris napi*, *Polyommatus icarus*). Longer flight

periods, multiple generations and larger bodies, expressed as longer wings, might facilitate
such swift expansions, making these species good long-distance dispersers (Stevens, Trochet,
Van Dyck, Clobert, & Baguette, 2012). Species with Western patterns (lineages connected to
Iberia, or Iberia and Italy) possess similar traits (e.g., *Lycaena phlaeas*, *Lasiommata megera*).
Better dispersers might be equipped to cross the mountains (Fric, Hula, Konvička, &
Pavlíčko, 2000), and expansions of woodland butterflies could have been facilitated by the
earlier onset of the Holocene near the Atlantic coast (Heiri et al., 2014), and slowed down by
continental climates surrounding the Balkans, and/or East Asian refugia.
The most threatened species in the Czech Republic are those having multiple
differentiated lineages across Europe with a single lineage inhabiting the focal region (*BAPS-*
East-Balkans, *BAPS-West*, *BAPS-Italy-Balkans* and *BAPS-temperate*) (e.g., *Pyrgus alveus*,
Hipparchia hermione, Polyommatus dorylas, and *Polyommatus damon*), whereas the
generalists (*BAPS-mix, HAP-mix* or *HAP-low*) are the least endangered (Fig. 4 c, f). This can
be related to the dispersal abilities of the species (Dapporto et al., 2019): the higher the
differentiation, the more specialized the butterfly is, and the less likely to survive in simplified
modern landscapes (Bartonova et al., 2016; Ockinger et al., 2010) because of reduced
possibility of rescue effect (Brückmann, Krauss, & Steffan-Dewenter, 2010). This is
supported by the evidence that butterfly species with a higher population diversification
disappeared first from islands (Dapporto et al., 2017). The species that display multiple
lineages across Europe are also threatened in other European countries (Maes et al., 2019).
The existence of distinct lineages across the species ranges was noted already by de
Lattin (1967), and, with the advent of genetic data, investigated in many case studies. Only
now, however, does the vast DNA database data allow us to test predictions on the resolution
of entire faunas. For a better understanding of lineage distribution and suture zones, it would
be necessary to properly cover the whole of Europe, and extend the sampling beyond, to
obtain information for entire species ranges. DNA barcoding of specimens in under-sampled
areas, even if not particularly biodiversity-rich, still represents a powerful tool for discovering
the driving forces of biogeography on large scales, such as the links between phylogeography
and species-specific traits.

 FIGURES

Figure 1. (a) Map of the Czech Republic with samples obtained for Czech butterfly DNA barcoding (sequencing of the mitochondrial gene COI). The points' sizes reflect the number of samples. (b–c) Numbers of species with different genetic patterns, as revealed by **(b)** Bayesian Analysis of Population Structure (BAPS), and **(c)** haplotype networks. The map projection is S-JTSK/Krovak East North.

Figure 2. Examples of Czech Republic butterfly species displaying different mitochondrial
genetic patterns according to Bayesian Analysis of Population Structure (BAPS categories),
depicted on the map of Europe. The method is sorting sequences into clusters of genetically
similar individuals (BAPS1–BAPS5, termed here as *genetic lineages*); the number of clusters
is estimated automatically by the 'rhierbaps' R package (Tonkin-Hill *et al.* 2018) and differs
among species. Examining the map, based on the presence of genetic lineages in the Czech
Republic, we assigned each of the 140 species into one of nine categories (*BAPS-low*, *BAPS-wide*, *BAPS-West*, *BAPS-mix*, *BAPS-Italy-Balkans*, *BAPS-East-Balkans*, *BAPS-mix-East*,
BAPS-Central, and *BAPS-temperate*). Maps were produced in QGIS 3.28.1 (http://qgis.org)
with WGS84 projection using country boundaries available at Natural Earth Free vector and
raster map data (https://www.naturalearthdata.com/).

Figure 3. Examples of the Czech butterfly species displaying different mitochondrial genetic
patterns, obtained by scoring haplotype networks (HAP). Mutations are depicted as black dots
and hatch marks. Examining the network, based on the relations of haplotypes present in the
Czech Republic to the other regions, we assigned each of the 140 species into one of ten
categories (*HAP-low*, *HAP-wide-low*, *HAP-wide-expand*, *HAP-wide-complex*, *HAP-mix*,
HAP-West, *HAP-Italy-Balkans*, *HAP-East-Balkans*, *HAP-Central*, and *HAP-temperate*).

Figure 4. Canonical correspondence analyses (CCA) biplots relating the mitochondrial genetic patterns found in Czech Republic butterflies to individual species traits. The explanatory variables (species traits) were forward-selected. Upper line: (a–c) patterns from Bayesian Analysis of Population Structure (BAPS) interpreted by **(a)** habitats, **(b)** life histories, **(c)** Red List categories (threat status) of individual species. Bottom line: (d–f) patterns from haplotype networks (HAP) interpreted by **(d)** habitats, **(e)** life histories, **(f)** Red List categories (threat status) of the species.

490   **DATA AVAILABILITY**
491
The newly generated sequences for Czech butterfly species are deposited in the BOLD
database as the BBCZ Project [CZBB001–CZBB959], and in the NCBI database [accession
codes OR890444–OR891402]. The Supplementary Materials (specimen metadata, Nexus
alignments for each species, R script for data handling, species traits and assigned
phylogeographic patterns) are stored at the figshare repository DOI
10.6084/m9.figshare.24559852.

REFERENCES

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Bartonova, A., Benes, J., Fric, Z. F., Chobot, K., & Konvicka, M. (2016). How universal are reserve design rules? A test using butterflies and their life history traits. *Ecography*, *39*(5), 456–464.

Bartonova, A., Benes, J., & Konvicka, M. (2014). Generalist-specialist continuum and life history traits of Central European butterflies (Lepidoptera) – Are we missing a part of the picture? *European Journal of Entomology*, *111*(4), 543–553.

Benes, J., Konvicka, M., Dvorak, J., Fric, Z., Havelda, Z., Pavlicko, A., … Weidenhoffer, Z. (2002). *Motýli České republiky: Rozšíření a ochrana I, II* (Vol. 2002). Praha: SOM.

Brückmann, S. V., Krauss, J., & Steffan‐Dewenter, I. (2010). Butterfly and plant specialists suffer from reduced connectivity in fragmented landscapes. *Journal of Applied Ecology*, *47*(4), 799–809.

Cheng, L., Connor, T. R., Sirén, J., Aanensen, D. M., & Corander, J. (2013). Hierarchical and spatially explicit clustering of DNA sequences with BAPS software. *Molecular Biology and Evolution*, *30*(5), 1224–1228.

Cizek, L., Fric, Z., & Konvicka, M. (2006). Host plant defences and voltinism in European butterflies. *Ecological Entomology*, *31*(4), 337–344.

Dapporto, L., Cini, A., Menchetti, M., Vodă, R., Bonelli, S., Casacci, L. P., … Vila, R. (2017). Rise and fall of island butterfly diversity: Understanding genetic differentiation and extinction in a highly diverse archipelago. *Diversity and Distributions*, *23*(10), 1169–1181.

Dapporto, L., Cini, A., Vodă, R., Dincă, V., Wiemers, M., Menchetti, M., … Vila, R. (2019). Integrating three comprehensive data sets shows that mitochondrial DNA variation is linked to species traits and paleogeographic events in European butterflies. *Molecular Ecology Resources*, *19*(6), 1623–1636.

Dapporto, L., Menchetti, M., Vodă, R., Corbella, C., Cuvelier, S., Djemadi, I., … Vila, R. (2022). The atlas of mitochondrial genetic diversity for Western Palaearctic butterflies. *Global Ecology and Biogeography*, *31*(11), 2184–2190.

de Lattin, G. (1967). *Grundriss der Zoogeographie*. Jena: VEB Gustav Fischer Verlag.

D'Ercole, J., Dincă, V., Opler, P. A., Kondla, N., Schmidt, C., Phillips, J. D., … Hebert, P. D. N. (2021). A DNA barcode library for the butterflies of North America. *PeerJ*, *9*, e11157.

Dincă, V., Zakharov, E. V., Hebert, P. D. N., & Vila, R. (2011). Complete DNA barcode reference library for a country's butterfly fauna reveals high performance for temperate Europe. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1704), 347–355.

Dincă, Vlad, Dapporto, L., Somervuo, P., Vodă, R., Cuvelier, S., Gascoigne-Pees, M., … Vila, R. (2021). High resolution DNA barcode library for European butterflies reveals continental patterns of mitochondrial genetic diversity. *Communications Biology*, *4*(1), 1–11.

Dincă, Vlad, Montagud, S., Talavera, G., Hernández-Roldán, J., Munguira, M. L., García-Barros, E., … Vila, R. (2015). DNA barcode reference library for Iberian butterflies enables a continental-scale preview of potential cryptic diversity. *Scientific Reports*, *5*, 12395.

Feurdean, A., Ruprecht, E., Molnár, Z., Hutchinson, S. M., & Hickler, T. (2018). Biodiversity-rich European grasslands: Ancient, forgotten ecosystems. *Biological Conservation*, *228*, 224–232.

Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for
amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan
invertebrates. *Molecular Marine Biology and Biotechnology*, *3*(5), 294–299.
Fric, Z., Hula, V., Konvička, M., & Pavlíčko, A. (2000). A note on the recent distribution of
Aporia crataegi (Linnaeus, 1758) in the Czech republic (Lepidoptera, Pieridae).
Atalanta, *31*(3/4), 453–454.
Galimberti, A., Assandri, G., Maggioni, D., Ramazzotti, F., Baroni, D., Bazzi, G., …
Casiraghi, M. (2021). Italian Odonates in the Pandora's Box: A Comprehensive DNA
Barcoding Inventory Shows Taxonomic Warnings at the Holarctic Scale. *Molecular*
Ecology Resources, *21*(1), 183–200.
Gamba, D., & Muchhala, N. (2020). Global patterns of population genetic differentiation in
seed plants. *Molecular Ecology*, *29*(18), 3413–3428.
Garcia-Barros, E. (2000). Body size, egg size, and their interspecific relationships with
ecological and life history traits in butterflies (Lepidoptera: Papilionoidea,
Hesperioidea). *Biological Journal of the Linnean Society*, *70*(2), 251–284.
Geiger, M. F., Herder, F., Monaghan, M. T., Almada, V., Barbieri, R., Bariche, M., …
Freyhof, J. (2014). Spatial heterogeneity in the Mediterranean Biodiversity Hotspot
affects barcoding accuracy of its freshwater fishes. *Molecular Ecology Resources*,
14(6), 1210–1221.
Habel, J. C., Roedder, D., Lens, L., & Schmitt, T. (2013). The genetic signature of
ecologically different grassland Lepidopterans. *Biodiversity and Conservation*, *22*(10),
2401–2411.
Hausmann, A., Haszprunar, G., Segerer, A. H., Speidel, W., Behounek, G., & Hebert, P. D. N.
(2011). Now DNA-barcoded: The butterflies and larger moths of Germany. *Spixiana*,
34(1), 47–58.
Hawlitschek, O., Bruns, C., Dey, L.S., Nuhlíčková, S., Felix, R., van Kleef, H., … Husemann,
M. (2023). The Genomics of Isolated Populations of Gampsocleis glabra (Orthoptera:
Tettigoniidae) in Central and Western Europe. *Insects*, *14*(12), 946.Hebert, P. D. N.,
Cywinska, A., Ball, S. L., & deWaard, J. R. (2003). Biological identifications through
DNA barcodes. *Proceedings. Biological Sciences*, *270*(1512), 313–321.
Hebert, P. D. N., Ratnasingham, S., & de Waard, J. R. (2003). Barcoding animal life:
Cytochrome c oxidase subunit 1 divergences among closely related species.
Proceedings of the Royal Society of London. Series B: Biological Sciences,
270(suppl_1), S96–S99.
Heiri, O., Brooks, S. J., Renssen, H., Bedford, A., Hazekamp, M., Ilyashuk, B., … Lotter, A.
F. (2014). Validation of climate model-inferred regional temperature change for late-
glacial Europe. *Nature Communications*, *5*, 4914.
Hejda, R., Farkač, J., & Chobot, K. (2017). *Červený seznam ohrožených druhů České*
republiky: BEZOBRATLÍ□: Red List of Threatened Species of Czech Republic:
INVERTEBRATES. Praha: Nature Conservation Agency of the Czech Republic.
Hendrich, L., Morinière, J., Haszprunar, G., Hebert, P. D. N., Hausmann, A., Köhler, F., &
Balke, M. (2015). A comprehensive DNA barcode database for Central European
beetles with a focus on Germany: Adding more than 3500 identified species to BOLD.
Molecular Ecology Resources, *15*(4), 795–818.
Hernández-Roldán, J. L., Dapporto, L., Dincă, V., Vicente, J. C., Hornett, E. A., Šíchová, J.,
… Vila, R. (2016). Integrative analyses unveil speciation linked to host plant shift in
Spialia butterflies. *Molecular Ecology*, *25*(17), 4267–4284.
Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), 907–
913.

Hewitt, G M. (2004). Genetic consequences of climatic oscillations in the Quaternary.
Philosophical Transactions of the Royal Society B: Biological Sciences, *359*(1442),
183–195.
Hewitt, Godfrey M. (1996). Some genetic consequences of ice ages, and their role in
divergence and speciation. *Biological Journal of the Linnean Society*, *58*(3), 247–276.
Hewitt, Godfrey M. (1999). Post-glacial re-colonization of European biota. *Biological
Journal of the Linnean Society*, *68*(1–2), 87–112.
Higgins, L. G., & Riley, N. D. (1970). *A Field Guide to the Butterflies of Britain and Europe*.
London: Collins.
Hofreiter, M., & Stewart, J. (2009). Ecological Change, Range Fluctuations and Population
Dynamics during the Pleistocene. *Current Biology*, *19*(14), R584–R594.
Horáček, I., & Ložek, V. (1988). *Palaeozoology and the Mid-European Quaternary Past:
Scope of the Approach and Selected Results*. Praha: Academia.
Huemer, P., Hebert, P. D. N., Mutanen, M., Wieser, C., Wiesmair, B., Hausmann, A., …
Fiedler, K. (2018). Large geographic distance versus small DNA barcode divergence:
Insights from a comparison of European to South Siberian Lepidoptera. *PLOS ONE*,
13(11), e0206668.
Huemer, P., Mutanen, M., Sefc, K. M., & Hebert, P. D. N. (2014). Testing DNA Barcode
Performance in 1000 Species of European Lepidoptera: Large Geographic Distances
Have Small Genetic Impacts. *PLOS ONE*, *9*(12), e115774.
Janoušek, V., Wang, L., Luzynski, K., Dufková, P., Vyskočilová, M. M., Nachman, M. W.,
… Tucker, P. K. (2012). Genome-wide architecture of reproductive isolation in a
naturally occurring hybrid zone between Mus musculus musculus and M. m.
Domesticus. *Molecular Ecology*, *21*(12), 3032–3047.
Kahlke, R.-D. (2014). The origin of Eurasian Mammoth Faunas (Mammuthus–Coelodonta
Faunal Complex). *Quaternary Science Reviews*, *96*, 32–49.
Kajtoch, Ł., Kubisz, D., Gutowski, J. M., & Babik, W. (2014). Evolutionary units of
Coraebus elatus (Coleoptera: Buprestidae) in central and eastern Europe–implications
for origin and conservation. *Insect Conservation and Diversity*, *7*(1), 41–54.Kajtoch,
Ł., Cieślak, E., Varga, Z., Paul, W., Mazur, M. A., Sramkó, G., & Kubisz, D. (2016).
Phylogeographic patterns of steppe species in Eastern Central Europe: A review and
the implications for conservation. *Biodiversity and Conservation*, *25*(12), 2309–2339.
Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., … Drummond,
A. (2012). Geneious Basic: An integrated and extendable desktop software platform
for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)*,
28(12), 1647–1649.
Kirschner, P., Záveská, E., Gamisch, A., Hilpold, A., Trucchi, E., Paun, O., … Schönswetter,
P. (2020). Long-term isolation of European steppe outposts boosts the biome's
conservation value. *Nature communications, 11*(1), 1968.Korb, S. K., Bolshakov, L.
V., Fric, Z. F., & Bartonova, A. (2016). Cluster biodiversity as a multidimensional
structure evolution strategy: Checkerspot butterflies of the group Euphydryas aurinia
(Rottemburg, 1775) (Lepidoptera: Nymphalidae). *Systematic Entomology*, *41*(2), 441–
457.
Kühne, G., Kosuch, J., Hochkirch, A., & Schmitt, T. (2017). Extra-Mediterranean glacial
refugia in a Mediterranean faunal element: The phylogeography of the chalk-hill blue
Polyommatus coridon (Lepidoptera, Lycaenidae). *Scientific Reports*, *7*, srep43533.
Leigh, J. W., & Bryant, D. (2015). popart: Full-feature software for haplotype network
construction. *Methods in Ecology and Evolution*, *6*(9), 1110–1116.

Lindestad, O., Nylin, S., Wheat, C. W., & Gotthard, K. (2022). Local adaptation of life cycles
645   in a butterfly is associated with variation in several circadian clock genes. *Molecular*
646   *Ecology*, *31*(5), 1461–1475.
647   Litman, J., Chittaro, Y., Birrer, S., Praz, C., Wermeille, E., Fluri, M., … Gonseth, Y. (2018).
648   A DNA barcode reference library for Swiss butterflies and forester moths as a tool for
649   species identification, systematics and conservation. *PLOS ONE*, *13*(12), e0208639.
650   Livraghi, L., Vodă, R., Evans, L. C., Gibbs, M., Dincă, V., Holland, P. W. H., … Breuker, C.
651   J. (2018). Historical and current patterns of gene flow in the butterfly Pararge aegeria.
652   *Journal of Biogeography*, *45*(7), 1628–1639.
653   Macek, J., Laštůvka, Z., Beneš, J., & Traxler, L. (2015). *Motýli a housenky střední Evropy IV.*
654   *Denní motýli*. Praha: Academia.
655   Mackintosh, A., Laetsch, D. R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., &
656   Lohse, K. (2019). The determinants of genetic diversity in butterflies. *Nature*
657   *Communications*, *10*(1), 3466.
658   Maes, D., Verovnik, R., Wiemers, M., Brosens, D., Beshkov, S., Bonelli, S., … Warren, M. S.
659   (2019). Integrating national Red Lists for prioritising conservation actions for
660   European butterflies. *Journal of Insect Conservation*, *23*(2), 301–330.
661   Mařan, J. (1958). Zoogeografické členění Československa. *Sborník Československé*
662   *Společnosti Zeměpisné*, *63*, 89–110.
663   Maresova, J., Suchackova Bartonova, A., Konvicka, M., Høye, T. T., Gilg, O., Kresse, J.-C.,
664   … Faltynek Fric, Z. (2021). The story of endurance: Biogeography and the
665   evolutionary history of four Holarctic butterflies with different habitat requirements.
666   *Journal of Biogeography*, *48*(3), 590–602.
667   Meier, R., Shiyang, K., Vaidya, G., & Ng, P. K. L. (2006). DNA Barcoding and Taxonomy in
668   Diptera: A Tale of High Intraspecific Variability and Low Identification Success.
669   *Systematic Biology*, *55*(5), 715–728.
670   Miller, M. J., Bermingham, E., Turner, B. L., Touchon, J. C., Johnson, A. B., & Winker, K.
671   (2021). Demographic consequences of foraging ecology explain genetic
672   diversification in Neotropical bird species. *Ecology Letters*, *24*(3), 563–571.
673   Monteiro, A., & Pierce, N. E. (2001). Phylogeny of Bicyclus (Lepidoptera: Nymphalidae)
674   inferred from COI, COII, and EF-1alpha gene sequences. *Molecular Phylogenetics*
675   *and Evolution*, *18*(2), 264–281.
676   Nürnberger, B., Lohse, K., Fijarczyk, A., Szymura, J. M., & Blaxter, M. L. (2016). Para-
677   allopatry in hybridizing fire-bellied toads (Bombina bombina and B. variegata):
678   Inference from transcriptome-wide coalescence analyses. *Evolution*, *70*(8), 1803–
679   1818.
680   Ockinger, E., Schweiger, O., Crist, T. O., Debinski, D. M., Krauss, J., Kuussaari, M., …
681   Bommarco, R. (2010). Life-history traits predict species responses to habitat area and
682   isolation: A cross-continental synthesis. *Ecology Letters*, *13*(8), 969–979.
683   Pearce, E. A., Mazier, F., Normand, S., Fyfe, R., Andrieu, V., Bakels, C., ... & Svenning, J. C.
684   (2023). Substantial light woodland and open vegetation characterized the temperate
685   forest biome before Homo sapiens. *Science advances*, *9*(45), 9135.
686   Pfäffle, M., Bolfíková, B. Č., Hulva, P., & Petney, T. (2014). Different Parasite Faunas in
687   Sympatric Populations of Sister Hedgehog Species in a Secondary Contact Zone.
688   *PLOS ONE*, *9*(12), e114030.
689   Potocký, P., Bartoňová, A., Beneš, J., Zapletal, M., & Konvička, M. (2018). Life-history traits
690   of Central European moths: Gradients of variation and their association with rarity and
691   threats. *Insect Conservation and Diversity*, *11*(5), 493–505.
692   Ratnasingham, S., & Hebert, P. D. N. (2007). bold: The Barcode of Life Data System
693   (http://www.barcodinglife.org). *Molecular Ecology Notes*, *7*(3), 355–364.

Reinhardt, R., Sbieschne, H., Settele, J., Fischer, U., & Fiedler, G. (2007). *Tagfalter von*
Sachsen. In B. Klausnitzer & R. Reinhardt (Eds.), Beiträge zur Insektenfauna
Sachsens Band 6. Entomologische Nachrichten und Berichte (vol. 11). Dresden.
Říčanová, Š., Koshev, Y., Říčan, O., Ćosić, N., Ćirović, D., Sedláček, F., & Bryja, J. (2013).
Multilocus phylogeography of the European ground squirrel: Cryptic interglacial
refugia of continental climate in Europe. *Molecular Ecology*, *22*(16), 4256–4269.
Sandom, C. J., Ejrnæs, R., Hansen, M. D. D., & Svenning, J.-C. (2014). High herbivore
density associated with vegetation diversity in interglacial ecosystems. *Proceedings of*
the National Academy of Sciences, *111*(11), 4162–4167.
Scalercio, S., Cini, A., Menchetti, M., Vodă, R., Bonelli, S., Bordoni, A., … Dapporto, L.
(2020). How long is 3 km for a butterfly? Ecological constraints and functional traits
explain high mitochondrial genetic diversity between Sicily and the Italian Peninsula.
Journal of Animal Ecology, *89*(9), 2013–2026.
Schmitt, T. (2007). Molecular biogeography of Europe: Pleistocene cycles and postglacial
trends. *Frontiers in Zoology*, *4*, 11.
Schmitt, T., & Varga, Z. (2012). Extra-Mediterranean refugia: The rule and not the exception?
Frontiers in Zoology, *9*(1), 1–12.
Sternberg, K. (1998). Die postglaziale Besiedlung Mitteleuropas durch Libellen, mit
besonderer Berücksichtigung Südwestdeutschlands (Insecta, Odonata). The postglacial
colonization of Central Europe by dragonflies, with special reference to southwestern
Germany (Insecta, Odonata). *Journal of Biogeography*, *25*(2), 319–337.
Stevens, V. M., Trochet, A., Van Dyck, H., Clobert, J., & Baguette, M. (2012). How is
dispersal integrated in life histories: A quantitative analysis using butterflies. *Ecology*
Letters, *15*(1), 74–86.
Stewart, J. R., Lister, A. M., Barnes, I., & Dalén, L. (2010). Refugia revisited: Individualistic
responses of species in space and time. *Proceedings of the Royal Society of London B:*
Biological Sciences, *277*(1682), 661–671.
Sucháčková Bartoňová, A., Konvička, M., Marešová, J., Bláhová, D., Číp, D., Skala, P., …
Faltýnek Fric, Z. (2021). Extremely Endangered Butterflies of Scattered Central
European Dry Grasslands Under Current Habitat Alteration. *Insect Systematics and*
Diversity, *5*(5). https://doi.org/10.1093/isd/ixab017
Ter Braak, C. J. F., & Šmilauer, P. (2012). *CANOCO Reference Manual and User's Guide:*
Software for Ordination (version 5.0). Ithaca (NY): Biometris.
Ter Braak, C. J. F., & Verdonschot, P. F. (1995). Canonical correspondence analysis and
related multivariate methods in aquatic ecology. *Aquatic sciences*, *57*(3), 255–289.
Tolman, T., & Lewington, R. (2008). *Collins butterfly guide: The Most Complete Field Guide*
to the Butterflies of Britain and Europe. London: Collins.
Tonkin-Hill, G., Lees, J. A., Bentley, S. D., Frost, S. D. W., & Corander, J. (2018).
RhierBAPS: An R implementation of the population clustering algorithm hierBAPS.
Wellcome Open Research, *3*, 93.
Vera, F. W. M. (2000). *Grazing Ecology and Forest History*. Egham: CABI.
Warren, M. S., Maes, D., Swaay, C. A. M. van, Goffart, P., Dyck, H. V., Bourn, N. A. D., …
Ellis, S. (2021). The decline of butterflies in Europe: Problems, significance, and
possible solutions. *Proceedings of the National Academy of Sciences*, *118*(2),
e2002551117.
Weigand, H., Beermann, A. J., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., … Ekrem, T.
(2019). DNA barcode reference libraries for the monitoring of aquatic biota in Europe:
Gap-analysis and recommendations for future work. *Science of The Total*
Environment, *678*, 499–524.

Wiemers, M., Balletto, E., Dincă, V., Fric, Z. F., Lamas, G., Lukhtanov, V., … Verovnik, R.
(2018). An updated checklist of the European Butterflies (Lepidoptera, Papilionoidea).
ZooKeys, *811*, 9–45.

 BIOSKETCH
747

The authors are experts in insect ecology, phylogeny, phylogeography, and distribution
patterns. The authors focus on the history, the current times and a future of Quaternary
ecosystems. They utilize butterflies, and other insects, as examples, to study the impact of
climate change, extinct megafauna, and habitat alterations on communities. They combine the
diverse scientific approaches and knowledge of the past to contribute to current habitat and
insect conservation. Author contributions: ASB designed the study, collected data, performed
analyses, and wrote the first draft of the manuscript. PŠ performed laboratory work and
analyses. MK collected data and contributed to the writing of the first draft. JB, LS, and VV
collected data. CS and HK performed laboratory work. JPM prepared all graphics in the
manuscript and supplements. ZFF designed the study, collected data, and performed analyses.
All authors contributed substantially to revisions and approved the final version of the
manuscript.

760
761

762 **SUPPORTING INFORMATION**

764 **Supporting Information 1.** Detailed summary and results of the Czech Republic butterfly
765 barcoding (sequencing cytochrome c oxidase subunit 1, COI) for each species occurring in the
766 country.