Running head: PbrSYP71 regulates ER accumulation in pollen tube

Corresponding author:

Peng Wang, E-mail: wangpeng@njau.edu.cn;

Juyou Wu, E-mail: juyouwu@njau.edu.cn

PbrSYP71 regulates ER accumulation by interacting with actin during pollen

tube growth in *Pyrus*

Mingliang Zhang[1], Ningyi Zhang[1], Chao Tang[1], Ming Qian[1], Mengjun Sun[1], Zongqi

Liu[1], Zhu Xie[1], Hao Zhang[1], Zhuqin Liu[3], Shaoling Zhang[1], Peng Wang[1]* and Juyou

Wu[1,2]*

1. Sanya Institute of Nanjing Agricultural University, State Key Laboratory of Crop

Genetics & Germplasm Enhancement and Utilization, Nanjing Agricultural University,

Nanjing 210095, China

2. Zhongshan Biological Breeding Laboratory, No.50 Zhongling Street, Nanjing,

Jiangsu 210014, China

3. Ningbo Key Laboratory of Charateristic Horticultural Crops in Quality Adjustment

and Resistance Breeding, Ningbo Academy of Agricultural Science, Ningbo 315040,

Zhejiang, China

*To whom correspondence should be addressed:

Peng Wang, E-mail: wangpeng@njau.edu.cn;

Juyou Wu, E-mail: juyouwu@njau.edu.cn

One-sentence summary: PbrSYP71, a syntaxin of plant protein, regulates ER

accumulation by interacting with actin for pollen tube growth.

Author contributions: Juyou Wu, Mingliang Zhang, and Ming Qian designed the

experiments; Mingliang Zhang, Chao Tang, and Mengjun Sun conducted the

experiments, with the help from Zongqi Liu, Zhu Xie, Hao Zhang, and Peng Wang; Mingliang Zhang, Juyou Wu, and Chao Tang wrote the manuscript; Ningyi Zhang, Peng Wang, Zhuqin Liu and Shaoling Zhang critically read and commented on the manuscript.

Funding information: This work was supported by Jiangsu Agricultural Science and Technology Innovation Fund (CX(22)3044), the National Natural Science Foundation of China (32102358), the Natural Science Foundation of Jiangsu Province (BK20210394), the Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions, and the Earmarked Fund for China Agriculture Research System (CARS-28).

Abstract

The pollen tube secretion system provides essential substances to support polarized pollen tube growth. The endoplasmic reticulum (ER) is a crucial component of the secretory system, and is predominantly concentrated in the pollen tube subapical region. However, the mechanism underlying the regulation of ER accumulation in pollen tube remains unclear. In this study, we identified a syntaxin protein PbrSYP71 that promoted pollen tube growth by regulating ER aggregation at pollen tube tip in pear. We found that PbrSYP71 anchored ER via the transmembrane domain. Reducing PbrSYP71 expression decreased the accumulation of ER in pollen tube. Moreover, overexpression of *PbrSYP71* resulted in bundle-like aggregation of the ER in tobacco leaf cells. Such aggregation state was disappeared when cytoplasmic domain of PbrSYP71 protein was absent or actin cytoskeleton was depolymerized. Additionally, the ER accumulation was decreased when the actin filament was depolymerized in pear pollen tube. We conclude that PbrSYP71 positively regulates ER accumulation in the subapical area of pollen tube by anchoring ER to actin, providing new insights into the underlying mechanisms of ER accumulation during polarized cell growth.

Abbreviations

ER, Endoplasmic reticulum; SNARE, soluble N-ethylmaleimide-sensitive factor attachment protein receptor; F-actin, actin filaments; v-SNAREs, vesicle-associated SNAREs; t-SNAREs, target cell-associated; SYP, syntaxin of plants; s-ODN, sense oligodeoxynucleotide; as-ODN, antisense oligodeoxynucleotide; S4B, pontamine Fast Scarlet 4B; LCI, luciferase complementation imaging; Y2H, Yeast two-hybrid; CB, Cytochalasins B; TM, transmembrane; aa, amino acids

Introduction

Pollen tube delivers male gametes to the ovaries for double fertilization, which is a necessary step for sexual reproduction in flowering plants. The polarized growth of pollen tube is a very fast process, which requires essential materials supported by secretory system consisting of the endoplasmic reticulum (ER), the Golgi apparatus, and vesicles for cell wall synthesis and plasma membrane expansion within the apical region (Steer and Steer, 1989; Campanoni and Blatt, 2007; Ruan *et al*., 2020). Meanwhile, pollen tube requires intense movements of secretory system to ensure the efficient delivery of these materials to the growing tip. However these distribution and movements are not completely randomized, instead, specific classes of organelles and vesicles are aggregated at the pollen tube tip (Cai *et al*., 2015). For example, secretory vesicles and ER respectively accumulate at apical and subapical area of pollen tube (Parton *et al*., 2001; Lovy-Wheeler *et al*., 2007). Exploring the molecular mechanisms of secretion system aggregation in pollen tube is crucial for understanding its rapid polarized growth.

The ER dynamically spreads throughout the cytoplasm and is crucial for cellular health (Westrate *et al*., 2015; Zheng *et al*., 2022). ER accumulation is regulated by the actin cytoskeleton, which is further mediated by actin-linker proteins. For now, two actin-linker proteins are identified, including AtSYP73 (a syntaxin protein of plants) and NET 3B in *Arabidopsis* (Cao *et al*., 2016; Wang and Hussey, 2017). The aggregation state of the ER in non-polarized growing tissues (e.g. leaves) is formed by an increase in the actin-linker protein and is decomposed by the depolymerization of the actin cytoskeleton (Cao *et al*., 2016; Wang and Hussey, 2017). In polarized growing tissues like pollen tube, action cytoskeleton is found to regulate ER aggregation at the subapice of pollen tube (Lovy-Wheeler *et al*., 2007). The role of actin-linker proteins (i.e. AtSYP73 protein that is found to localized at ER) in regulating ER aggregation in pollen tube, however, is still unknown.

AtSYP73 is a member of the plant SYP7 subgroup of the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) proteins, which is important for membrane fusion (Jahn and Scheller, 2006). SNAREs are

divided into vesicle-associated SNAREs (v-SNAREs) and target cell-associated SNAREs (t-SNAREs) based on their subcellular localization, and are divided into R-SNARE and Q-SNARE based on their conserved residues. Q-SNARE is further classified into three types, i.e. Qa-, Qb-, and Qc-SNARE (Lipka *et al.*, 2007). In general, R-SNAREs function as v-SNAREs, while Q-SNAREs act as t-SNAREs. v-SNAREs and t-SNAREs are two important components in exocytosis process. v-SNAREs are localized at vesicle, which need to be attached to "t-SNARE complex", composed of three t-SNARE proteins localized at target membrane, to form a core four-helix SNARE bundle called trans-SNARE complex to finish the membrane fusion (Jahn and Scheller, 2006; Han *et al.*, 2017). For tissues (e.g. leaves) in which ER is randomly distributed, exocytosis occurs at different directions in the tissue, causing non-polarized growth of these tissues (Cao *et al.*, 2016; Schwarz and Blower, 2016; Akita *et al.*, 2017). In some tissues (e.g. pollen tube), ER is aggregated at their specific location (Campanoni and Blatt, 2007; Cheung and Wu, 2007; Lovy-Wheeler *et al.*, 2007; Hepler and Winship, 2015), exocytosis thus potentially causes polarized growth of such tissues.

In "t-SNARE complex", one helix always comes from syntaxin group (Weimbs *et al.*, 1997; Sanderfoot *et al.*, 2000). Syntaxin of plants (SYP) is categorized into eight subgroups (SYP1-8) as part of the Q-SNARE complex. Within *Arabidopsis* the SYP1, 2, 3, 4, and 8 subgroups are associated with Qa-SNARE, while the SYP5, 6, and 7 subgroups are linked to Qc-SNARE (Sanderfoot *et al.*, 2000; Lipka *et al.*, 2007). In SYP7 subgroup in *Arabidopsis*, AtSYP73 regulates the ER accumulation and movement by interacting with actin in hypocotyl epidermal cells (Cao *et al.*, 2016). It remains unclear whether the members of the SYP7 subgroup play a role in the ER distribution in the pollen tube.

Pear (*Pyrus bretschneideri*) is an important orchard crop worldwide, of which the pollination is a very laborious but necessary process due to its self-incompatibility. Breeding self-compatibility cultivars has always been an important goal in pear production, for which understanding the underlying mechanisms of pollen tube growth is very relevant. Here, we chose pear (*Pyrus bretschneideri* cv. 'Dangshansuli')

as the case study due to its well-documented genome information that provides

fundamental database for studying the molecular mechanisms of pollen tube growth

(Wu *et al*., 2013). We identified a gene from the SYP7 subgroup, *PbrSYP71*, which

was highly expressed in pear pollen and promotes pollen tube growth. We found that

the PbrSYP71 protein was anchored to the ER, regulating the accumulation of ER in

the pear pollen tube. Moreover, the accumulation of PbrSYP71 was dependent on

actin cytoskeleton. Meanwhile, the aggregation of ER in pear pollen tube required the

support of F-actin. These results indicate that PbrSYP71 serves as a bridge connecting

ER and F-actin, anchors ER to F-actin to maintain the accumulation of ER in the

subapical area, thereby promoting efficient and rapid transport of materials to the

rapidly growing tip of the pollen tube.

Results

***PbrSYP* genes were identified in pear and *PbrSYP71* was specifically expressed in**

the pollen tube

A total of 43 *SYP* members from the pear genome were identified (Table S1). These

proteins were distributed in 8 subgroups (SYP1-SYP8) based on evolutionary

relationships (Figure S1). The SYP7 subgroup included five members, i.e. *PbrSYP71*,

72, *73*, *74*, and *75*, which had high level of homology and conservative domain and

motif with *AtSYP71*, *72*, and *73* (Figure 1A, B; Figure S2, and Figure S3). Our

RT-qPCR data showed that among all five PbrSYPs, only *PbrSYP71*, *73* and *75* were

expressed in the pollen (in grains and tubes), with PbrSYP71 showing the highest

expression level (Figure 1C). Expressions of PbrSYP72 and 74 were hardly detected

in the pollen. Moreover, *PbrSYP71* was only highly expressed in the pollen but not in

any other tissues e.g. leaves or stems (Figure S4, and Figure S5). These suggested an

important role of *PbrSYP71* in pear pollen tube growth.

***PbrSYP71* positively regulated pear pollen tube growth**

To verify the function of *PbrSYP71* in pear pollen tube growth, the *PbrSYP71*

expression was knocked down using as-ODN silencing technology. The phenotype of

pear pollen tubes was mal-functioning when treated with as-ODN-PbrSYP71 (Figure

2A). The relative expression of *PbrSYP71* in pollen tube treated with

as-ODN-PbrSYP71 reduced remarkably compared with that in s-ODN treated pollen

tube (Figure 2B). The germination rate of pollen grains was decreased when

inhibiting the expression of *PbrSYP71* (Figure 2C). The pollen tube length was

significantly decreased whereas pollen tube diameter was increased in the presence of

as-ODN-PbrSYP71 (Figure 2A, D, E). The cellulose content of the pollen tube was

significantly decreased when the *PbrSYP71* expression was knocked down (Figure

2F-H). These results suggested that knockdown of *PbrSYP71* expression caused

pollen tube defects in terms of pollen size growth and cellulose biosynthesis.

PbrSYP71 protein was localized at ER

The subcellular localization of PbrSYP71 assays were performed in leaf of *N.*

benthamiana. We found that PbrSYP71-GFP localized to cytoplasm and perinuclear

area (Figure 3 A, C, and D), in which perinuclear area was composed of ER (Schwarz

and Blower, 2016). Then, the PbrSYP71-GFP fusion protein and a ER marker,

AtRTP1-mCherry (Pan *et al*., 2016), were co-expressed in leaf epidermal cells.

Results showed that their fluorescence patterns overlapped (Figure 3E), suggesting

that PbrSYP71 was localized to ER. To further verify whether the transmembrane

(TM) domain of the PbrSYP71 protein anchors the ER, we constructed a

PbrSYP71-TM-GFP vector containing the TM domain (236-265 aa) (Figure S6A;

Table S2). When the PbrSYP71-TM-GFP fusion protein and AtRTP1-mCherry were

co-expressed in leaf epidermal cells, the co-localization of PbrSYP71-TM-GFP and

AtRTP1-mCherry was observed (Figure S6B). These results indicated that PbrSYP71

localized to the ER via TM domain.

PbrSYP71 **regulated ER distribution in pear pollen tube**

ER and secretory vesicles labeling results in pear showed that the ER accumulated at

the pollen tube sub-apical area and vesicles accumulated at the pollen tube tip,

forming a V-shaped tip followed by a tail (Figure S7). Similar results were also found

in lily pollen tube (Parton *et al*., 2001; Lovy-Wheeler *et al*., 2007; Hepler and

Winship, 2015). The fluorescence intensity of the ER at sub-apical area (area of black

box) significantly increased (Figure 4A and C) and the apical accumulation of vesicles

was disrupted when treating pollen tube with as-ODN-PbrSYP71 (Figure 4B, D and

E). These results suggested that *PbrSYP71* positively regulated accumulations of ER

and vesicles in pear pollen tube.

The accumulation and movement of PbrSYP71 protein was dependent on actin

cytoskeleton

PbrSYP71 protein was predicted to contain an actin-binding domain (ABD), a

SNARE-Qc domain and a TM domain (Figure 5A). Based on protein sequence

alignment, the ABD domain of PbrSYP71 was highly conserved with ABD domain

(77-95 aa) of the AtSYP73 protein (Figure 5B). Such a structure indicated that

PbrSYP71 was able to bind to actin. To further confirm this interaction, we chose

actin protein PbrActin1 (Pbr032870.1) as a candidate of PbrSYP71 interacting protein.

Y2H assays revealed that the yeast cells co-expressing PbrSYP71-BD and

PbrActin1-AD were able to grow on medium lacking Leu, Trp, Ade, and His (Figure

5C). In the LCI assays, fluorescence was observed when PbrSYP71-cLUC and

PbrAtin1-nLUC were co-expressed in tobacco leaves, as well as when

PbrSYP71-ABD-cLUC and PbrAtin1-nLUC were co-expressed; however, no

fluorescence was observed in the controls in both cases (Figure 5D, E). These results

suggested that PbrSYP71 interacted with PbrActin1 by ABD domain.

To test the function of actin cytoskeleton on the accumulation of PbrSYP71

protein, we examined the subcellular localization of PbrSYP71, TM domain of

PbrSYP71 (PbrSYP71-TM), and Cytoplasmic domain of PbrSYP71 (PbrSYP71-N) in

tobacco leaf cells (Figure 6A). PbrSYP71-GFP distributed around the whole

cytoplasm and presented bundle structure (Figure 6B). When actin cytoskeleton was

depolymerized with the actin cytoskeleton depolymerizing agent Cytochalasins B

(CB), PbrSYP71-GFP also distributed in the whole cytoplasm, but presented cisternae

network structure that was consistent with the structure of the ER (Figure 6C)

(Brandizzi *et al.*, 2002). These results suggested that the shape of PbrSYP71

depended on actin cytoskeleton. Further, we observed that PbrSYP71-TM-GFP

distributed in the whole cytoplasm presenting cisternae network structure (Figure 6D).

Cisternae network structure of PbrSYP71-TM-GFP enlarged when treated with CB

(Figure 6E), which was similar to PbrSYP71-GFP treated with CB (Figure 6C).

PbrSYP71-N-GFP also distributed in the whole cytoplasm, but presented a more

obvious bundle structure than that of PbrSYP71-GFP (Figure 6F). When

PbrSYP71-N-GFP was treated with CB, the bundle structure transitioned into circular

structure that evenly distributed in the cytoplasm (Figure 6G). These results suggested

that the state of aggregation of PbrSYP71 depended on cytoplasmic domain and actin

cytoskeleton. PbrSYP71 and PbrSYP71-TM were located at ER, and their cisternae

network structures were consistent with the ER structure. Thus, the aggregation state

of the ER was confirmed to be regulated by PbrSYP71 protein interaction with actin.

The bundles of PbrSYP71-GFP were continuously moving and changing in

tobacco leaf cells (Movie S1). When the actin cytoskeleton was depolymerized, the

movement of PbrSYP71-GFP was reduced with bundles disappearing (Movie S2).

PbrSYP71-TM-GFP, presenting cisternae network, also kept moving (Movie S3), and

the movement was reduced when treated with CB (Movie S4). We found that the

PbrSYP71-N-GFP presented strong bundles and exhibited rapid movements (Movie

S5), and the movement ceased when actin cytoskeleton was depolymerized (Movie

S6).

F-actin maintained ER distribution for pollen tube growth

To verify the function of F-actin on ER accumulation and vesicle accumulation, the

F-actin was depolymerized by treatment with CB in pear pollen tube. The ER

accumulation disappeared in the sub-apical region and uniformly distributed

throughout the cytoplasm when pollen tube was treated with CB (Figure 7A and C).

The apical accumulation of vesicles was reduced as well (Figure 7B, D and E). pollen

tube evolved abnormal phenotypes including decreased germination rate (Figure 8B),

reduced length (Figure 8A and C), increased diameter (Figure 8A and D) and

decreased cellulose content of cell wall (Figure 8E, F and G). These results suggested

that F-actin maintained the ER and vesicles accumulations, and supported pollen tube

growth.

Discussion

Syntaxin protein PbrSYP71 regulates cell tip growth

Syntaxin proteins are required for eukaryotes development as they play key roles in

cell tip growth (Hong *et al*., 2010; Qi *et al*., 2016). Here, we identified a syntaxin

protein – PbrSYP71 – that was involved in pear pollen tube tip growth by regulating

accumulations of the ER and secretory vesicles, and cell wall cellulose biosynthesis

(Figure 2; Figure 4). Previously, syntaxin proteins are found to regulate cell tip

growth in bacteria, fungi and model plant species *Arabidopsis*. For examples, in yeast,

syntaxin Tlg1p protein mediates polarized growth by mediating delivery of chitin

synthase III (Holthuis *et al*., 1998); in *N. crassa*, syntaxin NSYN1/2 proteins are

involved in hyphal tip growth by regulating vesicle fusion (Gupta *et al*., 2003); in

Arabidopsis, syntaxin of plant SYP121, SYP123, SYP132, and AtSYP81 proteins

promote root tip growth by mediating vesicle fusion (Hirano *et al*., 2023; Wang *et al*.,

2023), SYP124 is involved in pollen tube tip growth (Silva *et al*., 2010), SYP1s and

SYP72 plays a key role in pollen development and germination (Li *et al*., 2019; Zhou

et al., 2022). To our knowledge, we are the first to identify the role of syntaxin protein

in polarized cell growth in important crop plants (i.e. pear). All together, these results

suggest that the function of syntaxin proteins in regulating cell tip growth is

conserved across different species.

PbrSYP71 regulates cell tip growth by affecting ER accumulation and

membrane fusion

We showed that PbrSYP71 protein promoted the ER accumulation in pear pollen tube

(Figure 4), which was necessary for pollen tube growth (Figure 2). Recent studies

have found that the distribution of ER is directly related to cell health and cell polarity.

For examples, in human U2OS cells, when ER distribution is disrupted, the

distribution of other organelles is also affected, resulting in neurologic disorders and

cancer (Zheng *et al*., 2022). ER accumulation at the root hair tip is necessary for the

root hair development (Campanoni and Blatt, 2007). In addition, PbrSYP71 protein,

as t-SNARE protein, was localized at ER (Figure 3). Previously, syntaxin proteins are

found to mediate vesicle fusion in together with ER. For example, ER-located

t-SNARE proteins (i.e. syntaxin 18) mediate the fusion of COPI (coatomer complex I)

vesicles, deriving by Golgi with ER and the fusion of legionella-containing vacuole

with ER (Spang and Schekman, 1998; Hatsuzawa *et al*., 2006; Spang, 2013;

Kawabata *et al*., 2021). Therefore, we propose that PbrSYP71 protein located at ER

could mediate the fusion of vesicles with ER for maintaining the functioning of

secretory system in pear pollen tube. Previous studies on the function of syntaxin in

cell polar growth mainly focus on regulating the fusion of vesicles and plasma

membranes. For examples, syntaxin proteins located in plasma membrane, including

syntaxin Tlg1p (in yeast), syntaxin NSYN1/2 (in *N. crassa*), SYP121 SYP123,

SYP124, and SYP132 (in *Arabidopsis*), are involved in regulating the membrane

fusion of secretory vesicle with plasma membrane for cell tip growth (Holthuis *et al*.,

1998; Gupta *et al*., 2003; Silva *et al*., 2010; Hirano *et al*., 2023). Here, we showed that

PbrSYP71 was located at ER and positively regulate vesicle secretion, which

provided a specific target for studying the function of syntaxin in mediating vesicle

fusion with ER during pollen tube growth. We conclude that syntaxin proteins

regulates ER accumulation in addition to mediating membrane fusion in cell polar

growth.

The cytoplasmic domain of PbrSYP71 protein regulates the ER accumulation

and movement

We found that the ER accumulation in pear pollen tube depended on the interaction

between cytoplasmic domain of PbrSYP71 protein and actin (Figure 5; Figure 6;

Figure 7). Similar functions of syntaxin proteins exist in leaf tissues and animal cells.

For examples, in *Arabidopsis*, cytoplasmic domain of AtSYP73 protein interacts with

actin, leading to the ER accumulation (Cao *et al*., 2016). In INS-1 and CHO cells,

H$_{abc}$ domains of syntaxin 1A determine its cluster distribution without affecting its

localization on plasma membrane (Yang *et al*., 2006; Fan *et al*., 2007). These results

showed that cytoplasmic domain of syntaxin plays a key role in ER accumulation.

Interestingly, we found that cytoplasmic domain of PbrSYP71 moved on the actin

cytoskeleton independent of ER movement (Movie S1-6), suggesting that the ER

movement may be driven by the movement of PbrSYP71 cytoplasmic domain. In

Arabidopsis, AtSYP73 protein has been found to regulate the ER flow rate (Cao *et al*.,

2016). Our results clarified the role of PbrSYP71 cytoplasmic domain in regulating

ER accumulation and movement, further broadening our understanding of the

functions of syntaxin proteins in exocytosis process during cellular polar growth.

314 The intracellular localization of PbrSYP71 protein is dependent on TM domain

Different syntaxin proteins have different subcellular localizations. For examples,

syntaxin 1 and SYP131 (in mammals and *Arabidopsis*) are located on plasma

membrane (Teng *et al*., 2001; Ichikawa *et al*., 2015); syntaxin 5 and SYP31 (in

mammals and *Arabidopsis*) are presenting on the Golgi apparatus (Teng *et al*., 2001;

Rui *et al*., 2021); syntaxin 18 and SYP73 (in mammals and *Arabidopsis*), as well as

PbrSYP71 in pear found in our study, are located at ER (Figure 3) (Teng *et al*., 2001;

Cao *et al*., 2016). The domains of syntaxin proteins that determine their specific

subcellular localizations are also different for different syntaxin members (Kasai and

Akagawa, 2001). We found that PbrSYP71 anchored the ER via TM domain (Figure

S6). Additionally, the length of the TM domain determines the subcellular localization

of syntaxins 3 (Watson and Pessin, 2001). The trans-Golgi network localization of

syntaxin 6 depends on H2 domain and YGRL motif (Watson and Pessin, 2000), and

the Golgi localization of syntaxin Sed5p relies on its the TMD amino acid, the COPI

coatomer binding motif, and the folded-back conformation (Gao and Banfield, 2020).

Thus, the conserved structures that influence the syntaxin subcellular localization

differ between syntaxin proteins.

331 Conclusion

We presented a model describing the regulatory role of PbrSYP71 in ER

accumulation at the pollen tube subapical area, taking an important horticultural crop

pear as an example (Figure 9). PbrSYP71 protein anchors the ER via the TM domain

and bridges ER with F-actin through actin-binding domain (ABD) interaction. This

maintains ER accumulation at the subapical region of pollen tube, providing materials

for pollen tube growth. Insufficient PbrSYP71 protein, however, lead to the loss of

connection between ER and F-actin, resulting in no ER aggregation and evenly ER

distribution. Consequently, the reduction in secretory vesicles fails to provide

sufficient materials for pollen tube growth, causing abnormal pollen tube phenotypes.

The absence of track and collar structures for PbrSYP71 protein attachment,

following F-actin depolymerization, reduce ER accumulation and secretory vesicle

content. In conclusion, we elucidate the key role of *PbrSYP71* in the ER accumulation

during pollen tube growth, providing new insights for the role of syntaxin proteins in

exocytosis process during polarized cell growth.

Materials and methods

Plant materials

Pear trees (*P. bretschneideri* cv. "Dangshansuli") were grown in the Fruit

Experimental Field of Nanjing Agricultural University, Nanjing, China (32°N, 118°E).

Mature pollens from these trees were collected and stored at -20°C in a desiccator for

later use.

Identification and basic information analysis of *SYP* genes in pear

Protein sequences of 25 AtSYP were acquired from the TAIR

(https://www.arabidopsis.org/) and these sequences were used as queries to perform

BLASTP searches against the Pear Genome Database (http://pcwgdb.njau.edu.cn/)

(Hu *et al*., 2023) by TBtools software (China) with an e-value cut-off of $1.0 \times e^{-5}$

(Chen *et al*., 2023). Meanwhile, hidden Markov model profiles of the SYP domain

(PF05739) were submitted to HMMsearch software to search for candidate SYP

proteins in pear (Johnson *et al*., 2010). Redundant sequences from all obtained gene

sequences were removed. SMART (http://smart.embl-heidelberg.de/) and the Pfam

database (http://pfam.xfam.org/) were applied to detect SNARE functional domains in

obtained sequences (Letunic *et al*., 2021; Mistry *et al*., 2021). The relative molecular

weights and isoelectric points of PbrSYP proteins were determined by ExPASY

(http://web.expasy.org/compute_pi/) (Artimo *et al*., 2012). The transmembrane

structure and motifs of PbrSYP proteins were respectively predicted using CCTOP

(https://cctop.ttk.hu/) (Dobson *et al*., 2015) and MEME

(https://meme-suite.org/meme/) (Bailey *et al*., 2015). Actin-binding domain (ABD)

LIG_Actin_WH2_2 was predicted by the Eukaryotic Linear Motif (ELM) program

(Dinkel *et al*., 2016).

Phylogenetic analysis

Phylogenetic analysis of 43 pear *SYP* members was performed by MEGA7.0 software

(Kumar *et al*., 2016). Evolutionary analysis was conducted using 43 sequences of pear

SYP proteins and 25 sequences of *Arabidopsis* SYP proteins. ClusterW alignment of
all sequences was preferred, followed by the construction of a *SYP* evolutionary tree
using the Neighbor Joining, with internal branches being taken 1000 times.
Phylogenetic relationships of pear and *Arabidopsis* SYP7 subgroup proteins were
conducted using the Neighbor Joining method in MEGA7.

Pollen culture *in vitro*

The method of cultivating pollen *in vitro* can be found in our previous study (Tang *et*
al., 2023). Mature pollen was taken out from -20°C and slowly thawed to room
temperature, and evenly dispersed in a pollen medium (containing 0.5 mM calcium
nitrate $Ca(NO_3)_2$, 1.5 mM boric acid (H_3BO_3), 25 mM 2-(N-morpholino)
ethanesulfonic acid hydrate (MES) and 450 mM sucrose (sucrose) with pH 5.8, and
cultured at 25°C/120 rpm. Then, the pollen tubes were collected by centrifugation at
2,000 rpm for 5 min, and then stored at -80°C.

RNA extraction

387 Total RNA was extracted using a FastPure[®] Universal Plant Total RNA Isolation Kit
388 (Vazyme, China). The HiScript[®] II1st Strand cDNA Synthesis Kit (Vazyme, China)
389 was employed for reverse transcribing the RNA into cDNA.

Quantitative real-time PCR (RT-qPCR) assays

The quantitative specific primers of *SYP* gene family members in pear were designed
by the online Primer BLAST tool (https://www.ncbi.nlm.nih.gov/tools/primer-blast/)
(Ye *et al*., 2012). LC480 II light cycler (Roche, Germany) instrument was used for
qRT-PCR. The volume of each sample was 20 μL, which included 5 μL of specific
primers with a concentration of 10 μM, 2 μL of cDNA with a concentration of
100ng/μL, 3 μL of ddH_2O, and 10 μL of $2 \times$ SYBR mixed dye (Vazyme, China). The
reaction procedure for qPCR followed three steps, being that step 1 is incubating at
95℃ for 5 minutes, step 2 is incubating at 95 °C for 3 seconds, followed by 60 °C for
10 seconds and 72 °C for 10 seconds, for a total of 45 cycles, and step 3 is drawing a
dissolution curve to verify primer specificity. These data were analyzed using
LightCycler480 software. *PbrUBQ* (Pbr035452.1) was used as an internal reference in
pear pollen tube to normalize the expression levels of *SYP* genes. The relative

expression level of *SYP* was calculated according to the $2^{-\Delta\Delta Ct}$ method. The primers of

qRT-PCR assays were shown in Supplemental Table S3.

Analysis of *PbrSYP* genes expression based on transcriptome data

The expression levels of *PbrSYP* genes were analyzed based on our previous

transcriptomic data (http://pcwgdb.njau.edu.cn/) (Zhou *et al*., 2016; Hu *et al*., 2023) in

19 pear tissues including four seedling tissues (JSM, juvenile stem; CT, cotyledon;

JLF, juvenile leaf; and RT, root), six flower organ tissues (OA, ovary; SP, sepal; ST,

stamen; SY, style; PT, petal; and FLS, flower stalk), five fruit tissues (FTC, fruit core;

FS, flesh ;FTS, fruit stalk; SD, seed; and PR, pericarp), flower bud (FLB), stem (SM),

adult leaf (ALF), leaf bud (LFB). The transcripts per kilobase million (TPM) values

were used for measuring the *PbrSYP* gene expression level at different developmental

stages during pollen germination and pollen tube growth (MP, mature pollen grains;

HP, hydrated pollen grains; PT, growing pollen tubes; and SPT, stopped-growth pollen

tubes). The RPKM (reads per kilobase per million) values were used for measuring

the gene expression level. These data of expression levels of *PbrSYP* genes were

given in Supplemental Table S4 and Table S5.

Antisense oligodeoxynucleotide (as-ODN) assays

Antisense oligodeoxynucleotides (ODN) assays were performed as described

previously in reference to the previous study (Chen *et al*., 2018). First, the mixture of

ODN sequence and Lipofectamine 3000 (Thermo Fisher, USA) transfection reagent

was incubated at room temperature for 15 minutes. Then the mixture was added to the

pollen tube culture medium. The final concentration of ODN was 40 μM. Pollen was

cultured for 3 h with 25℃ and 120 rpm. The images of pollen germination and pollen

tube growth were obtained using an optical microscope, and the diameter and length

of pollen tubes were measured using ImageJ software (https://imagej.nih.gov/ij/). The

as-ODN Primer of PbrSYP71 gene was designed using online RNA fold website

Server (http://rna.tbi.univie.ac.at//cgi-bin/RNAWebSuite/RNAfold.cgi). The primers

of ODN were shown in Supplemental Table S3.

Detection of cellulose in pollen tube cell wall

Cellulose staining of pollen tube cell wall was conducted based on our previous study

(Li *et al.*, 2022). First, prepared pollen tubes were washed with phosphate buffered

saline (PBS) three times, and then fixed with 4% paraformaldehyde for 30 minutes.

After fixation, pollen tubes were transferred to PBS containing 0.01% Pontamine Fast

Scarlet 4B (S4B) (Sigma-Aldrich, USA), then incubated at room temperature for 5

minutes and washed with PBS three times. The dyed pollen tube was observed using a

confocal microscope LSM800 (Zeiss, Germany). The fluorescence was measured

using ZEN 3.1 software (Zeiss, Germany).

Subcellular localization assays

The subcellular localization assays were performed based on (Sparkes *et al.*, 2006).

Firstly, the CDS of *PbrSYP71*, *PbrSYP71-TM* (amino acid residue 236-265aa), and

PbrSYP71-N (amino acid residue 1-235aa) without stop codon were cloned into the

pCAMBIA1300-35S::GFP vector to construct 35S::PbrSYP71-GFP,

35S::PbrSYP71-TM-GFP and 35S::PbrSYP71-N-GFP vectors. The CDS of AtRTP1

without stop codon was cloned into the pCAMBIA1300-35S::mCherry vector to

construct 35S::AtRTP1-mCherry vector. Then, vectors were transformed into

epidermal cells of *N. benthamiana*. Meanwhile, the pCAMBIA1300-35S::GFP vector

was used as the control. Nucleus was marked by DAPI dye. Finally, the fluorescence

was observed using an LSM800 laser scanning confocal microscope (Zeiss, Germany).

The primers of subcellular localization assays were shown in Supplemental Table S3.

ER labeling

ER-Tracker Red (Beyotime, shanghai, China) – a specific red fluorescent probe for

ER – was used as a marker of the ER of eukaryotic cells (Wang *et al.*, 2017; Su *et al.*,

2021). ER of the pollen tube was labeled ER-Tracker Red as follows: pollen tubes

were transferred to pollen culture medium containing 20 μM ER-Tracker Red and

allowed to incubate for 30 min. The fluorescence of ER-Tracker Red was observed

using an LSM800 laser scanning confocal microscope (Zeiss, Germany).

Secretory vesicles labeling

Parton's method was used to measure secretory vesicles (Parton *et al.*, 2001). The

pollen tubes were transferred to pollen culture medium containing 8 μM FM4-64 and

allowed to incubate for 15 min. The fluorescence of FM4-64 was observed using an

LSM800 laser scanning confocal microscope (Zeiss, Germany).

Yeast two-hybrid (Y2H) assay

The *PbrSYP71* and *PbrActin1* CDS sequences without stop codon were cloned into

the pGBDT7 and pGADT7 vector to construct the PbrSYP71-BD vector and the

PbrSYP71-AD vector, respectively. Then, different combinations were transferred

into Y2HGold yeast cells. Finally, these yeast cells grew on selective medium lacking

Leu, Trp, Ade, and His (−L/−T/−A/−H) to detect possible interactions. Growth in the

medium lacking Leu, Trp, Ade, and His indicated protein-protein interactions. The

AD-T and BD-lam combination was used as a negative control, and the AD-T and

BD-53 combination was used as a positive control. The primers of yeast two-hybrid

assays were shown in Supplemental Table S3.

Luciferase complementation imaging (LCI) assays

Firstly, The *PbrActin1* CDS without stop codon was cloned into the

pCAMBIA1300-nLuc to construct PbrActin1-nLuc nLuc. The *PbrSYP71* and

PbrSYP71-ABD (amino acid residue 77-95 aa) CDS without stop codon was cloned

into the pCAMBIA1300-cLuc to construct PbrSYP71-cLuc and

PbrSYP71-ABD-cLuc, respectively. Then, different combinations were transformed

into leaf cells of *N. benthamiana.* Finally, the LUC activity in leaves of *N.*

benthamiana was detected using In Vivo Plant Imaging System (PIXIS 1024B,

Princeton Instrument, USA). The primers of LCI assays were shown in Supplemental

Table S3.

Movie production

485 The video of PbrSYP71-GFP movement was produced using the following method.

486 First, at least 35 consecutive fluorescence images were obtained using LSM800 laser

487 scanning confocal microscope (Zeiss, Germany), in which fluorescence images were

488 obtained every 14 s. Then, these images were played in chronological order (Frame

489 Per Second=5) using ZEN 2012 software (Zeiss, Germany).

Statistical analysis

GraphPad Prism 8.2.1 (GraphPad Software, USA) was used to plot the results

including their averages and the corresponding standard deviation. SPSS v22 (IBM,

USA) was used to determine statistical significance. Significant statistical differences

for two samples were analyzed using the Student's *t*-test (P < 0.01). Significant

statistically differences for multiple samples were analyzed using Duncan LSD test of

one-way ANOVA (P < 0.05).

Supplementary data

The following supplemental materials are available:

Fig. S1 Neighbor-Joining phylogeny of SYP family proteins from *Pyrus* and

Arabidopsis.

Fig. S2 Conservative domains analysis of proteins of SYP7 subgroup.

Fig. S3 Conserved motif analysis of SYP proteins

Fig. S4 Expression patterns of *PbrSYP* genes in different tissues of pear.

Fig. S5 The expression of *PbrSYP* genes during four development stages of pear

pollen.

Fig. S6 Intracellular localization of PbrSYP71-TM to the ER in tobacco leaf cells.

Fig. S7 The distribution of ER and vesicle in pear pollen tube.

Table S1 Detailed information on *SYP* gene family members in pear.

Table S2 The 1D structure of PbrSYP proteins.

Table S3 Primers used in this study

Table S4 Expression patterns of PbrSYP genes in different tissues of pear.

Table S5 The expression of PbrSYP genes during four development stages of pear

pollen.

Movie S1 The movement of PbrSYP71-GFP in tobacco leaf cells.

Movie S2 The movement of PbrSYP71-GFP treated with the actin depolymerizing

agent Cytochalasins B (CB) in tobacco leaf cells .

Movie S3 The movement of PbrSYP71-TM-GFP in tobacco leaf cells.

Movie S4 The movement of PbrSYP71-TM-GFP treated with the CB in tobacco leaf

cells.

Movie S5 The movement of PbrSYP71-N-GFP in tobacco leaf cells.

Movie S6 The movement of PbrSYP71-N-GFP treated with the CB in tobacco leaf

cells.

Acknowledgements

We thank Dr. Yuehua Ma from Central laboratory of College of Horticulture in

Nanjing Agricultural University for her technical assistance in the use of laser

confocal microscopy LSM800 (Zeiss, Germany).

Conflict of interest

The authors declare no competing interests.

References

Akita K, Kobayashi M, Sato M, Kutsuna N, Ueda T, Toyooka K, Nagata N, Hasezawa S,

Higaki T. Cell wall accumulation of fluorescent proteins derived from a trans-Golgi cisternal

membrane marker and paramural bodies in interdigitated *Arabidopsis* leaf epidermal cells.

Protoplasma. 2017:**254:**1367-377. https://doi.org/10.1007/s00709-016-0955-1

Artimo P, Jonnalagedda M, Arnold K, Baratin D, Csardi G, de Castro E, Duvaud S, Flegel V,

Fortier A, Gasteiger E, et al. ExPASy: SIB bioinformatics resource portal. Nucleic Acids Res.

2012:**40:**W597-603. https://doi.org/10.1093/nar/gks400

Bailey T, Johnson J, Grant C, Noble W. The MEME Suite. Nucleic Acids Res. 2015:**43:**W39-49.

https://doi.org/10.1093/nar/gkv416

Brandizzi F, Snapp E, Roberts A, Lippincott-Schwartz J, Hawes C. Membrane protein

transport between the endoplasmic reticulum and the Golgi in tobacco leaves is energy

dependent but cytoskeleton independent: evidence from selective photobleaching. Plant Cell.

2002:**14:**1293-1309. https://doi.org/10.1105/tpc.001586

Cai G, Parrotta L, Cresti M. Organelle trafficking, the cytoskeleton, and pollen tube growth. J

Integr Plant Biol. 2015:**57:**63-78. https://doi.org/10.1111/jipb.12289

Campanoni P, Blatt M. Membrane trafficking and polar growth in root hairs and pollen tubes. J

Exp Bot. 2007:**58:**65-74. https://doi.org/10.1093/jxb/erl059

Cao P, Renna L, Stefano G, Brandizzi F. SYP73 Anchors the ER to the Actin Cytoskeleton for

Maintenance of ER Integrity and Streaming in *Arabidopsis*. Curr Biol. 2016:**26:**3245-3254.

https://doi.org/10.1016/j.cub.2016.10.024

Chen C, Wu Y, Li J, Wang X, Zeng Z, Xu J, Liu Y, Feng J, Chen H, He Y, et al. TBtools-II: A

"one for all, all for one" bioinformatics platform for biological big-data mining. Mol Plant.

2023:**16:**1733-1742. https://doi.org/10.1016/j.molp.2023.09.010

Chen J, Wang P, de Graaf B, Zhang H, Jiao H, Tang C, Zhang S, Wu J. Phosphatidic Acid

Counteracts S-RNase Signaling in Pollen by Stabilizing the Actin Cytoskeleton. Plant Cell.

2018:**30:**1023-1039. https://doi.org/10.1105/tpc.18.00021

Cheung A, Wu H. Structural and functional compartmentalization in pollen tubes. J Exp Bot.

2007: **58:** 75-82. https://doi.org/10.1093/jxb/erl122

Dinkel H, Van Roey K, Michael S, Kumar M, Uyar B, Altenberg B, Milchevskaya V,

Schneider M, Kühn H, Behrendt A, et al. ELM 2016--data update and new functionality of

the eukaryotic linear motif resource. Nucleic Acids Res. 2016:**44:**D294-300.

https://doi.org/10.1093/nar/gkv1291

Dobson L, Reményi I, Tusnády G. CCTOP: a Consensus Constrained TOPology prediction web

server. Nucleic Acids Res. 2015:**43:**W408-412. https://doi.org/10.1093/nar/gkv451

Fan J, Yang X, Lu J, Chen L, Xu P. Role of H(abc) domain in membrane trafficking and

targeting of syntaxin 1A. Biochem Biophys Res Commun. 2007:**359:** 245-250.

https://doi.org/10.1016/j.bbrc.2007.05.065

Gao G, Banfield D. Multiple features within the syntaxin Sed5p mediate its Golgi localization.

Traffic. 2020**21:**274-296. https://doi.org/10.1111/tra.12720

Gupta G, Free S, Levina N, Keränen S, Heath I. Two divergent plasma membrane syntaxin-like

SNAREs, nsyn1 and nsyn2, contribute to hyphal tip growth and other developmental processes

in Neurospora crassa. Fungal Genet Biol. 2003:**40:**271-286.

https://doi.org/10.1016/s1087-1845(03)00109-9

Han J, Pluhackova K, Böckmann R. The Multifaceted Role of SNARE Proteins in Membrane

Fusion. Front Physiol. 2017:**8:**5. https://doi.org/10.3389/fphys.2017.00005

Hatsuzawa K, Tamura T, Hashimoto H, Hashimoto H, Yokoya S, Miura M, Nagaya H, Wada

I. Involvement of syntaxin 18, an endoplasmic reticulum (ER)-localized SNARE protein, in

ER-mediated phagocytosis. Mol Biol Cell. 2006:**17:** 3964-3977.

https://doi.org/10.1091/mbc.e05-12-1174

Hepler P, Winship L. The pollen tube clear zone: clues to the mechanism of polarized growth. J

Integr Plant Biol. 2015:**57:** 79-92. https://doi.org/10.1111/jipb.12315

Hirano T, Ebine K, Ueda T, Higaki T, Watanabe-Nakayama T, Konno H,

Takigawa-Imamura H, Sato M. The SYP123-VAMP727 SNARE complex delivers secondary

cell wall components for root hair shank hardening in *Arabidopsis*. Plant Cell.

2023:30;35(12)**:**4347-4365. https://doi.org/10.1093/plcell/koad240

Holthuis J, Nichols B, Pelham H. The syntaxin Tlg1p mediates trafficking of chitin synthase III

to polarized growth sites in yeast. Mol Biol Cell. 1998:**9:** 3383-3397.

https://doi.org/10.1091/mbc.9.12.3383

Hong SY, So J, Lee J, Min K, Son H, Park C, Yun SH, Lee YW. Functional analyses of two
syntaxin-like SNARE genes, GzSYN1 and GzSYN2, in the ascomycete Gibberella zeae. Fungal
Genet Biol. 2010:**47:** 364-372. https://doi.org/10.1016/j.fgb.2010.01.005

Hu J, Huang B, Yin H, Qi K, Jia Y, Xie Z, Gao Y, Li H, Li Q, Wang Z, et al. PearMODB: a
multiomics database for pear (*Pyrus*) genomics, genetics and breeding study. Database (Oxford).
2023:baad050. https://doi.org/10.1093/database/baad050

Ichikawa M, Iwano M, Sato M. Nuclear membrane localization during pollen development and
apex-focused polarity establishment of SYP124/125 during pollen germination in Arabidopsis
thaliana. Plant Reprod. 2015:**28:** 143-151. https://doi.org/10.1007/s00497-015-0265-3

Jahn R, Scheller R. SNAREs--engines for membrane fusion. Nature reviews. Nat Rev Mol Cell
Biol. 2006:**7:** 631-643. https://doi.org/10.1038/nrm2002

Johnson L, Eddy S, Portugaly E. Hidden Markov model speed heuristic and iterative HMM
search   procedure.   BMC   Bioinformatics.   2010:**11:**
431.https://doi.org/10.1186/1471-2105-11-431

Kasai K, Akagawa K. Roles of the cytoplasmic and transmembrane domains of syntaxins in
intracellular   localization   and   trafficking.   J   Cell   Sci.   2001:**114:**   3115-3124.
https://doi.org/10.1242/jcs.114.17.3115

Kawabata M, Matsuo H, Koito T, Murata M, Kubori T, Nagai H, Tagaya M, Arasaki K.
Legionella hijacks the host Golgi-to-ER retrograde pathway for the association of
Legionella-containing   vacuole   with   the   ER.   PLoS   Pathog.   2021:**17:**   e1009437.
https://doi.org/10.1371/journal.ppat.1009437

Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version
7.0   for   Bigger   Datasets.   Mol   Biol   Evol.   2016:**33:**   1870-1874.
https://doi.org/10.1093/molbev/msw054

Li B, Li Y, Liu F, Tan X, Rui Q, Tong Y, Qiao L, Gao R, Li G, et al. Overexpressed Tomosyn
Binds Syntaxins and Blocks Secretion during Pollen Development. Plant Physiol.
2019:**181:**1114-1126. https://doi.org/10.1104/pp.19.00965

Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in 2020.
Nucleic Acids Res. 2021:**49:** D458-D460. https://doi.org/10.1093/nar/gkaa937

Li X, Tang C, Li X, Zhu X, Cai Y, Wang P, Zhang S, Wu J. Cellulose accumulation mediated

by PbrCSLD5, a cellulose synthase-like protein, results in cessation of pollen tube growth in

Pyrus bretschneideri. Physiol Plant. 2022:**174:** e13700. https://doi.org/10.1111/ppl.13700

Lipka V, Kwon C, Panstruga R. SNARE-ware: the role of SNARE-domain proteins in plant

biology.   Annu   Rev   Cell   Dev   Biol.   2007:**23:**   147-174.

https://doi.org/10.1146/annurev.cellbio.23.090506.123529

Lovy-Wheeler A, Cárdenas L, Kunkel J, Hepler P. Differential organelle movement on the

actin cytoskeleton in lily pollen tubes. Cell Motil Cytoskeleton. 2007:**64:** 217-232.

https://doi.org/10.1002/cm.20181

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar G, Sonnhammer E, Tosatto S,

Paladin L, Raj S, Richardson L, et al. Pfam: The protein families database in 2021. Nucleic

Acids Res. 2021:**49:** D412-D419. https://doi.org/10.1093/nar/gkaa913

Pan Q, Cui B, Deng F, Quan J, Loake G, Shan W. RTP1 encodes a novel endoplasmic reticulum

(ER)-localized protein in Arabidopsis and negatively regulates resistance against biotrophic

pathogens. New Phytol. 2016:**209:** 1641-1654. https://doi.org/10.1111/nph.13707

Parton R, Fischer-Parton S, Watahiki M, Trewavas A. Dynamics of the apical vesicle

accumulation and the rate of growth are related in individual pollen tubes. J Cell Sci. 2001:**114:**

2685-2695. https://doi.org/10.1242/jcs.114.14.2685

Qi Z, Liu M, Dong Y, Zhu Q, Li L, Li B, Yang J, Li Y, Ru Y, Zhang H, et al. The syntaxin

protein (MoSyn8) mediates intracellular trafficking to regulate conidiogenesis and

pathogenicity of rice blast fungus. New Phytol. 2016:**209:** 1655-1667.

https://doi.org/10.1111/nph.13710

Ruan H, Li J, Wang T, Ren H. Secretory Vesicles Targeted to Plasma Membrane During Pollen

Germination and Tube Growth. Frontiers in cell and developmental biology. 2020:**8:** 615447.

https://doi.org/10.3389/fcell.2020.615447

Rui Q, Tan X, Liu F, Li Y, Liu X, Li B, Wang J, Yang H, Qiao L, Li T, et al. Syntaxin of

plants31 (SYP31) and SYP32 is essential for Golgi morphology maintenance and pollen

development. Plant Physiol. 2021:**186:** 330-343.https://doi.org/10.1093/plphys/kiab049

Sanderfoot A, Assaad F, Raikhel N. The *Arabidopsis* genome. An abundance of soluble

N-ethylmaleimide-sensitive factor adaptor protein receptors. Plant Physiol. 2000:**124:**

1558-1569. https://doi.org/10.1104/pp.124.4.1558

Schwarz D, Blower M. The endoplasmic reticulum: structure, function and response to cellular

signaling. Cell Mol Life Sci. 2016:**73:** 79-94. https://doi.org/10.1007/s00018-015-2052-6

Silva P, Ul-Rehman R, Rato C, Di Sansebastiano G, Malhó R. Asymmetric localization of

Arabidopsis SYP124 syntaxin at the pollen tube apical and sub-apical zones is involved in tip

growth. BMC Plant Biol. 2010:**10:**179. https://doi.org/10.1186/1471-2229-10-179

Spang A. Retrograde traffic from the Golgi to the endoplasmic reticulum. Cold Spring Harb

Perspect Biol. 2013:**5.** https://doi.org/10.1101/cshperspect.a013391

Spang A, Schekman R. Reconstitution of retrograde transport from the Golgi to the ER in vitro. J

Cell Biol. 1998:**143:**589-599. https://doi.org/10.1083/jcb.143.3.589

Sparkes I, Runions J, Kearns A, Hawes C. Rapid, transient expression of fluorescent fusion

proteins in tobacco plants and generation of stably transformed plants. Nat Protoc. 2006:**1:**

2019-2025. 10.1038/nprot.2006.286

Steer M, Steer J. Pollen tube tip growth. New Phytol. 1989:**111:** 323-358.

https://doi.org/10.1111/j.1469-8137.1989.tb00697.x

Su X, Yan X, Chen X, Guo M, Xia Y, Cao Y. Calcofluor white hypersensitive proteins contribute

to stress tolerance and pathogenicity in entomopathogenic fungus, Metarhizium acridum. Pest

Manag Sci 2021:**77:** 1915-1924. https://doi.org/10.1002/ps.6218

Tang C, Wang P, Zhu X, Qi K, Xie Z, Zhang H, Li X, Gao H, Gu T, Gu C, et al. Acetylation of

inorganic pyrophosphatase by S-RNase signaling induces pollen tube tip swelling by repressing

pectin methylesterase. Plant Cell. 2023:1;35(9)**:**3544-3565.

https://doi.org/10.1093/plcell/koad162

Teng F, Wang Y, Tang B. The syntaxins. Genome Biol. 2001:**2:** REVIEWS3012.

https://doi.org/10.1186/gb-2001-2-11-reviews3012

Wang H, Niu H, Zhai Y, Lu M BiPCharacterization of Genes from Pepper (*L.*) and the Role of

in Response to Endoplasmic Reticulum and Multiple Abiotic Stresses. Front Plant Sci. 2017:**8:**

1122. https://doi.org/10.3389/fpls.2017.01122

Wang M, Zhang H, Zhao X, Zhou J, Qin G, Liu Y, Kou X, Zhao Z, Wu T, Zhu J, et al.

SYNTAXIN OF PLANTS81 regulates root meristem activity and stem cell niche maintenance

via ROS signaling. Plant Physiol.2023:**191:**1365-1382. https://doi.org/10.1093/plphys/kiac530

Wang P, Hussey P NETWORKED 3B: a novel protein in the actin cytoskeleton-endoplasmic

reticulum interaction. J Exp Bot. 2017:**68:** 1441-1450. https://doi.org/10.1093/jxb/erx047

Watson R, Pessin J. Functional cooperation of two independent targeting domains in syntaxin 6

is required for its efficient localization in the trans-golgi network of 3T3L1 adipocytes. J Biol

Chem. 2000:**275:** 1261-1268. https://doi.org/10.1074/jbc.275.2.1261

Watson R, Pessin J. Transmembrane domain length determines intracellular membrane

compartment localization of syntaxins 3, 4, and 5. Am J Physiol Cell Physiol. 2001:**281:**

C215-223. https://doi.org/10.1152/ajpcell.2001.281.1.C215

Weimbs T, Low S, Chapin S, Mostov K, Bucher P, Hofmann K. A conserved domain is present

in different families of vesicular fusion proteins: a new superfamily. Proc Natl Acad Sci U S A.

1997:**94:** 3046-3051. https://doi.org/10.1073/pnas.94.7.3046

Westrate L, Lee J, Prinz W, Voeltz G. Form follows function: the importance of endoplasmic

reticulum shape. Annu Rev Biochem. 2015:**84:** 791-811.

https://doi.org/10.1146/annurev-biochem-072711-163501

Wu J, Wang Z, Shi Z, Zhang S, Ming R, Zhu S, Khan M, Tao S, Korban S, Wang H, et al.

The genome of the pear (*Pyrus bretschneideri* Rehd.). Genome Res. 2013:**23:** 396-408.

https://doi.org/10.1101/gr.144311.112

Yang X, Xu P, Xiao Y, Xiong X, Xu T. Domain requirement for the membrane trafficking and

targeting of syntaxin 1A. J Biol Chem. 2006:**281:** 15457-15463.

https://doi.org/10.1074/jbc.M513246200

Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden T. Primer-BLAST: a tool to

design target-specific primers for polymerase chain reaction. BMC Bioinformatics. 2012:**13:**

134. https://doi.org/10.1186/1471-2105-13-134

Zheng P, Obara C, Szczesna E, Nixon-Abell J, Mahalingan K, Roll-Mecak A,

Lippincott-Schwartz J, Blackstone C. ER proteins decipher the tubulin code to regulate

organelle distribution. Nature. 2022:**601:** 132-138. https://doi.org/10.1038/s41586-021-04204-9

Zhou H, Yin H, Chen J, Liu X, Gao Y, Wu J, Zhang S. Gene-expression profile of developing

pollen tube of *Pyrus bretschneideri*. Gene Expr Patterns. 2016:**20:** 11-21.

https://doi.org/10.1016/j.gep.2015.10.004

Zhou X, Zheng Y, Wang L, Li H, Guo Y, Li M, Sun M, Zhao P. SYP72 interacts with the

mechanosensitive channel MSL8 to protect pollen from hypoosmotic shock during hydration.

Nat Commun. 2022:**13:** 73. https://doi.org/10.1038/s41467-021-27757-9

Figure legends

Figure 1. PbrSYP71, a member of the plant SYP7 subgroup of SNARE, was

expressed in pear pollen and pollen tube. A) Sequence alignment of PbrSYP71,

PbrSYP72, PbrSYP73, PbrSYP74, PbrSYP75, AtSYP71, AtSYP72 and AtSYP73.

The black line represents the SNARE-Qc domain and transmembrane (TM) domain.

B) Phylogenetic relationship of pear and Arabidopsis SYP7 subgroup proteins. The

tree was generated using the Maximum Likelihood method in MEGA7. **C)** Expression

level of *PbrSYP71*, *PbrSYP73*, and *PbrSYP75* in pollen grain and pollen tube were

determined by RT-qPCR assay. Data are mean ± SD of 3 independent biological

replicates. Different letters indicate significant differences as determined by Duncan

LSD test (P < 0.05).

Figure 2. ***PbrSYP71* is essential for pear pollen tube growth. A)** Pear pollen treated

with as-ODN for 3 h. Bars = 50 μm. The experiments were repeated 3 times with

similar results, and representative pictures were shown. **B)** Expression level of

PbrSYP71 in pollen tube transfected with sense oligodeoxynucleotide of *PbrSYP71*

(s-ODN), and antisense oligodeoxynucleotide (as-ODN), were determined by

RT-qPCR assays. **C)** The germination rate of pollen in (A). **D)** The length of pollen

tube in (A). **E)** pollen tube width in (A), at least 10 pollen tubes were measured. **F)**

Cellulose content of pollen tube wall was indicated using S4B dye under s-ODN and

as-ODN treatments. Bars = 10 μm. The experiments were repeated 3 times with

similar results, and representative pictures were shown. **G)** Fluorescence intensity of

S4B at 20 μm distance from pollen tube tips along the white thread in (F). **H)**

Quantitative analysis of the S4B fluorescence intensity at the apex of the pollen tube

about 5 μm from the tip along black line in (F), at least 15 pollen tubes were measured.

For (B-D), data are mean ± SD of 3 independent biological replicates. For (B-E, and

H), differences were identified using Student's *t*-test, significant differences (P < 0.01)

between means are indicated by **.

Figure 3. Intracellular localization of PbrSYP71 to the endoplasmic reticulum

(ER) in tobacco leaf cells. A) Vector as control and PbrSYP71-GFP proteins were

transiently expressed in transformed tobacco leaf epidermal cells. Green color

represents the fluorescence intensity of GFP. The nucleus was stained by DAPI as a

marker, for which the fluorescence is purple. White fluorescence showed

colocalization of green and purple fluorescence. Bars = 20 μm. **B)** A locally enlarged

image further shows that PbrSYP71-GFP was located in the cytoplasm, with the black

arrow indicating the position of the cell wall. Bar = 5 μm. **C)** A locally enlarged

image further shows that the location of PbrSYP71-GFP around the nucleus, with the

black arrow indicating the position of nucleus. Bar = 5 μm. **D)** The fluorescence

intensity of PbrSYP71-GFP and DAPI on the section line in Figure C. Red represents

DAPI, green represents PbrSYP71. **E)** Intracellular localization of PbrSYP71-GFP

and ER marker (AtRTP1-mCherry). Red color represents the fluorescence intensity of

mCherry. Yellow fluorescence showed colocalization of green and red fluorescence.

Bars = 20 μm.

Figure 4. PbrSYP71 is essential for endoplasmic reticulum (ER) and vesicles

accumulation of pear pollen tube. A) Pear pollen tube stained by ER-tracker Red

treated with as-ODN for 3 h. Bars = 10 μm. **B)** Pear pollen tube stained by FM4-64

treated with as-ODN for 3 h. Bars = 10 μm. For (A and B), the experiments were

repeated 3 times with similar results, and representative pictures were shown. **C)**

Quantitative analysis of ER fluorescence intensity of pollen sub-apcial area at black

box in (A), 7 pollen tubes were measured. **D)** Fluorescence intensity of FM4-64 at 40

μm distance from pollen tube tips along the white thread in (B). **E)** Quantitative

analysis of FM4-64 fluorescence intensity of pollen apcial area at black box in (B), at

least 7 pollen tubes were measured. For (C and E), differences were identified using

Student's *t*-test, significant differences (P < 0.01) between means are indicated by **.

Figure 5. PbrSYP71 interacts with PbrActin1. A) Schematic diagram of PbrSYP71

protein demonstrating the actin-binding domain (ABD), a SNARE-Qc domain and a

transmembrane domain. **B)** Alignment of the predicted actin-binding domains in

PbrSYP71 protein and AtSYP73 protein. **C)** PbrSYP71 interacts with PbrActin1 in

yeast two-hybrid assays. PbrActin fused to the pGADT7 vector and that of PbrSYP71

fused to the pGBDT7 vector. Growth in the medium that lacked Leu, Trp, Ade, and

His indicated protein-protein interactions. **D)** LCI assay showing interaction between

PbrSYP71 and PbrActin1 in vivo. Fluorescence on the leaf surface indicated

protein-protein interactions. **E)** LCI assay showing interaction between

PbrSYP71-ABD and PbrActin1 in vivo.

Figure 6. The distribution of PbrSYP71 depended on actin cytoskeleton. A)

Schematic diagram of segments of PbrSYP71 protein demonstrating the SYP71,

SYP71-TM and SYP71-N. **B)** PbrSYP71-GFP presented bundle structure in tobacco

leaf epidermal cells. Bars = 10 μm. **C)** The bundle structure of PbrSYP71-GFP

collapsed to cisternae network when treated with actin cytoskeleton depolymerizing

agent Cytochalasins B (CB). Bars = 10 μm. **D)** PbrSYP71-TM-GFP presented

cisternae network. Bars = 10 μm. **E)** The cisternae structure of PbrSYP71-TM-GFP

proteins was enlarged when treated with CB. Bars = 10 μm. **F)** PbrSYP71-N-GFP

presented stronger bundle structure. Bars = 10 μm. **G)** The bundle structure of

PbrSYP71-N-GFP collapsed to circles with randomly distributing when treated with

CB. Bars = 10 μm. Green color represents the fluorescence intensity of GFP.

Figure 7. F-actin is essential for endoplasmic reticulum (ER) and vesicles

accumulation of pear pollen tube. A) Pear pollen tubes were stained by ER-tracker

Red and treated with actin cytoskeleton depolymerizing agent Cytochalasins B (CB).

Bars = 10 μm. **B)** Pear pollen tubes were stained by FM4-64 and treated with CB.

Bars = 10 μm. For (A and B), the experiments were repeated 3 times with similar

results, and representative pictures were shown. **C)** Quantitative analysis of ER

fluorescence intensity of pollen apcial area at black box in (A), at least 8 pollen tubes

were measured. **D)** Fluorescence intensity of FM4-64 at 40 μm distance from pollen

tube tips along the white thread in (B). **E)** Quantitative analysis of FM4-64

fluorescence intensity of pollen apcial area at black box in (B), at least 8 pollen tubes

were measured. For (C and E), differences were identified using Student's *t*-test,

significant differences (P < 0.01) between means are indicated by **.

Figure 8. F-actin is essential for pear pollen tube growth. A) Pear pollen tubes

were treated with actin cytoskeleton depolymerizing agent Cytochalasins B (CB) for 3

h. Bars=50 μm. The experiments were repeated 3 times with similar results, and

representative pictures were shown. **B)** The germination rate of pollen in (A). **C)** The

length of pollen tube in (A). **D)** pollen tube width in (A), 7 pollen tubes were

measured. For (B and C) data are mean ± SD of three independent biological

replicates. Differences were identified using Student's *t*-test, significant differences (P

< 0.01) between means are indicated by ** . **E)** Cellulose content of pollen tube wall

was indicated using S4B dye under CB treatments. Bars = 10 μm. The experiments

were repeated 3 times with similar results, and representative pictures were shown. **F)**

Fluorescence intensity of S4B at 20 μm distance from pollen tube tips along the white

thread in (E). **G)** Quantitative analysis of the S4B fluorescence intensity at the apex of

the pollen tube about 5 μm from the tip along black line in (E), at least 14 pollen tubes

were measured. For (B, C, and G), differences were identified using Student's *t*-test,

significant differences (P < 0.01) between means are indicated by **.

Figure 9. Hypothetical model of function of PbrSYP71 in endoplasmic reticulum

(ER) distribution. A) PbrSYP71 anchors the ER to actin filament (F-actin) to

maintain the accumulation of the ER at subapcial area in pear pollen tube. **B)** The

accumulation of the ER was disturbed when the *PbrSYP71* expression was knocked

down. **C)** The accumulation of the ER was disturbed when F-actin was

depolymerized.

Figure S1. Neighbor-Joining phylogeny of SYP family proteins from *Pyrus*

***bretschneideri* and *Arabidopsis thaliana*.** The phylogenetic tree was constructed by

MEGA 7.0 software using full-length protein sequences. A bootstrap analysis was

conducted with 1000 replicates. Different subgroups are distinguished by different

color backgrounds (SYP1-SYP8), respectively. AtSYPs and PbrSYPs from

Arabidopsis and *Pyrus*, respectively. The scale bar is 0.1.

Figure S2. Conservative domains analysis of proteins of SYP7 subgroup. A)

Schematic diagram of SYP7 subgroup proteins demonstrating a SNARE-Qc domain.

B) Predicted actin-binding domains (ABD) and transmembrane (TM) domain have

been identified in PbrSYP71-75.

Figure S3. Conserved motif analysis of SYP proteins. A) Phylogenetic tree of the

PbrSYP genes. **B)** Composition and position of 9 conserved motifs of pear PbrSYP

proteins indicated by different color blocks.

Figure S4. Expression patterns of *PbrSYP* genes in different tissues of

"Dangshansuli" pear. Expression patterns of *PbrSYP* genes in 19 different tissues,

including 4 seedling tissues (JSM, juvenile stem; CT, cotyledon; JLF, juvenile leaf;

RT, root), flower bud (FLB), stem (SM), adult leaf (ALF), leaf bud (LFB), 6 flower

organ tissues (OA, ovary, SP, sepal;; ST, stamen; SY, style; PT, petal; FLS, flower

stalk), 5 fruit tissues (FTC, fruit core; FS, flesh ;FTS, fruit stalk; SD, seed; PR,

pericarp). The expression levels were calculated as log2-transformed (TPM) values.

white, yellow, and red indicate low, moderate, and high levels, respectively.

Figure S5. The expression of *PbrSYP* genes during 4 development stages of pear

pollen, including mature pollen grains (MP), hydrated pollen grains (HP), growing

pollen tubes (PT) and stopped-growth pollen tubes (SPT). Color scale represents

log2-transformed (FPKM) values. The expression levels were calculated as

log2-transformed (RPKM) values. white, yellow, and red indicate low, moderate, and

high levels, respectively.

Figure S6. Intracellular localization of PbrSYP71-TM to the endoplasmic

reticulum (ER) in tobacco leaf cells. A) Schematic diagram of PbrSYP71. **B)**

Schematic   diagram   of   PbrSYP71-TM.   **C)**   Intracellular   localization   of

PbrSYP71-TM-GFP and ER marker (AtRTP1-mCherry). Red color represents the

fluorescence intensity of mCherry. Yellow fluorescence showed colocalization of

green and red fluorescence. Bars = 20 μm.

Figure S7. The distribution of the endoplasmic reticulum (ER) and vesicle in

pear Pollen tube. A) ER-Tracker Red staining distribution in a growing pollen tube

of pear. Bars = 10 μm. **B)** FM4-64 staining distribution in a growing pollen tube of

pear. Bars = 10 μm. **C)** Quantification of fluorescence intensity along the white line

through the bright image in (A). **D)** Quantification of fluorescence intensity along the

white line through the bright image in (B).

Figure 1. *PbrSYP71*, a member of the plant SYP7 subgroup of SNARE, was expressed in pear pollen and pollen tube. **A)** Sequence alignment of PbrSYP71, PbrSYP72, PbrSYP73, PbrSYP74, PbrSYP75, AtSYP71, AtSYP72 and AtSYP73. The black line represents the SNARE-Qc domain and transmembrane (TM) domain. **B)** Phylogenetic relationship of pear and Arabidopsis SYP7 subgroup proteins. The tree was generated using the Maximum Likelihood method in MEGA7. **C)** Expression level of *PbrSYP71*, *PbrSYP73*, and *PbrSYP75* in pollen grain and pollen tube were determined by RT-qPCR assay. Data are mean ± SD of 3 independent biological replicates. Different letters indicate significant differences as determined by Duncan LSD test (P < 0.05).

Figure 2. *PbrSYP71* **is essential for pear pollen tube growth. A)** Pear pollen treated with as-ODN for 3 h. Bars = 50 μm. The experiments were repeated 3 times with similar results, and representative pictures were shown. **B)** Expression level of *PbrSYP71* in pollen tube transfected with sense oligodeoxynucleotide of PbrSYP71 (s-ODN), and antisense oligodeoxynucleotide (as-ODN), were determined by RT-qPCR assays. **C)** The germination rate of pollen in (A). **D)** The length of pollen tube in (A). **E)** pollen tube width in (A), at least 10 pollen tubes were measured. **F)** Cellulose content of pollen tube wall was indicated using S4B dye under s-ODN and as-ODN treatments. Bars = 10 μm. The experiments were repeated 3 times with similar results, and representative pictures were shown. **G)** Fluorescence intensity of S4B at 20 μm distance from pollen tube tips along the white thread in (F). **H)** Quantitative analysis of the S4B fluorescence intensity at the apex of the pollen tube about 5 μm from the tip along black line in (F), at least 15 pollen tubes were measured. For (B-D), data are mean ± SD of 3 independent biological replicates. For (B-E, and H), differences were identified using Student's t-test, significant differences ($P < 0.01$) between means are indicated by **.

Figure 3. Intracellular localization of PbrSYP71 to the endoplasmic reticulum (ER) in tobacco leaf cells. A) Vector as control and PbrSYP71-GFP proteins were transiently expressed in transformed tobacco leaf epidermal cells. Green color represents the fluorescence intensity of GFP. The nucleus was stained by DAPI as a marker, for which the fluorescence is purple. White fluorescence showed colocalization of green and purple fluorescence. Bars = 20 μm. **B)** A locally enlarged image further shows that PbrSYP71-GFP was located in the cytoplasm, with the black arrow indicating the position of the cell wall. Bar = 5 μm. **C)** A locally enlarged image further shows that the location of PbrSYP71-GFP around the nucleus, with the black arrow indicating the position of nucleus. Bar = 5 μm. **D)** The fluorescence intensity of PbrSYP71-GFP and DAPI on the section line in Figure C. Red represents DAPI, green represents PbrSYP71. **E)** Intracellular localization of PbrSYP71-GFP and ER marker (AtRTP1-mCherry). Red color represents the fluorescence intensity of mCherry. Yellow fluorescence showed colocalization of green and red fluorescence. Bars = 20 μm.

Figure 4. *PbrSYP71* is essential for endoplasmic reticulum (ER) and vesicles accumulation of pear pollen tube. A) The ER was stained with ER-tracker Red dye after treating the pollen tube with as-ODN for 3 hours. Bars = 10 μm. **B)** The vesicles were stained with FM4-64 dye after treating the pollen tube with as-ODN for 3 hours. Bars = 10 μm. For (A and B), the experiments were repeated 3 times with similar results, and representative pictures were shown. **C)** Quantitative analysis of ER fluorescence intensity of pollen sub-apcial area at black box in (A), 7 pollen tubes were measured. **D)** Fluorescence intensity of FM4-64 at 40 μm distance from pollen tube tips along the white thread in (B). **E)** Quantitative analysis of FM4-64 fluorescence intensity of pollen apcial area at black box in (B), at least 7 pollen tubes were measured. For (C and E), differences were identified using Student's *t*-test, significant differences (P < 0.01) between means are indicated by **.

Figure 5. PbrSYP71 interacts with PbrActin1. A) Schematic diagram of PbrSYP71 protein demonstrating the actin-binding domain (ABD), a SNARE-Qc domain and a transmembrane domain (TM). **B)** Alignment of the predicted actin-binding domains in PbrSYP71 protein and AtSYP73 protein. **C)** PbrSYP71 interacts with PbrActin1 in yeast two-hybrid assays. PbrActin fused to the pGADT7 vector and that of PbrSYP71 fused to the pGBDT7 vector. Growth in the medium that lacked Leu, Trp, Ade, and His indicated protein-protein interactions. **D)** LCI assay showing interaction between PbrSYP71 and PbrActin1 in vivo. Fluorescence on the leaf surface indicated protein-protein interactions. **E)** LCI assay showing interaction between PbrSYP71-ABD and PbrActin1 in vivo.

Figure 6. The distribution of PbrSYP71 depended on actin cytoskeleton. A) Schematic diagram of segments of PbrS-YP71 protein demonstrating the SYP71, SYP71-TM and SYP71-N. **B)** PbrSYP71-GFP presented bundle structure in tobacco leaf epidermal cells. Bars = 10 μm. **C)** The bundle structure of PbrSYP71-GFP collapsed to cisternae network when treated with actin cytoskeleton depolymerizing agent Cytochalasins B (CB). Bars = 10 μm. **D)** PbrSYP71-TM-GFP presented cisternae network. Bars = 10 μm. **E)** The cisternae structure of PbrSYP71-TM-GFP proteins was enlarged when treated with CB. Bars = 10 μm. **F)** PbrSYP71-N-GFP presented stronger bundle structure. Bars = 10 μm. **G)** The bundle structure of PbrSYP71-N-GFP collapsed to circles with randomly distributing when treated with CB. Bars = 10 μm. Green color represents the fluorescence intensity of GFP.

Figure 7. F-actin is essential for endoplasmic reticulum (ER) and vesicles accumulation of pear pollen tube. A) The ER was stained with ER-tracker Red dye after treating the pollen tube with actin cytoskeleton depolymerizing agent Cytochalasins B (CB). Bars = 10 μm. **B)** The vesicles were stained with FM4-64 dye after treating the pollen tube with CB. Bars = 10 μm. For (A and B), the experiments were repeated 3 times with similar results, and representative pictures were shown. **C)** Quantitative analysis of ER fluorescence intensity of pollen apcial area at black box in (A), at least 8 pollen tubes were measured. **D)** Fluorescence intensity of FM4-64 at 40 μm distance from pollen tube tips along the white thread in (B). **E)** Quantitative analysis of FM4-64 fluorescence intensity of pollen apcial area at black box in (B), at least 8 pollen tubes were measured. For (C and E), differences were identified using Student's *t*-test, significant differences (P < 0.01) between means are indicated by **.

Figure 8. F-actin is essential for pear pollen tube growth. A) Pear pollen tubes were treated with actin cytoskeleton depolymerizing agent Cytochalasins B (CB) for 3 h. Bars=50 μm. The experiments were repeated 3 times with similar results, and representative pictures were shown. **B)** The germination rate of pollen in (A). **C)** The length of pollen tube in (A). **D)** pollen tube width in (A), 7 pollen tubes were measured. For (B and C) data are mean ± SD of three independent biological replicates. Differences were identified using Student's *t*-test, significant differences (P < 0.01) between means are indicated by ** . **E)** Cellulose content of pollen tube wall was indicated using S4B dye under CB treatments. Bars = 10 μm. The experiments were repeated 3 times with similar results, and representative pictures were shown. **F)** Fluorescence intensity of S4B at 20 μm distance from pollen tube tips along the white thread in (E). **G)** Quantitative analysis of the S4B fluorescence intensity at the apex of the pollen tube about 5 μm from the tip along black line in (E), at least 14 pollen tubes were measured. For (B, C, and G), differences were identified using Student's *t*-test, significant differences (P < 0.01) between means are indicated by **.

Figure 9. Hypothetical model of function of *PbrSYP71* in endoplasmic reticulum (ER) distribution. A) PbrSYP71 protein anchors the ER to actin filament (F-actin) to maintain the accumulation of the ER at subapcial area in pear pollen tube. **B)** The accumulation of the ER was disturbed when the *PbrSYP71* expression was knocked down. **C)** The accumulation of the ER was disturbed when F-actin was depolymerized.

Parsed Citations

Akita K, Kobayashi M, Sato M, Kutsuna N, Ueda T, Toyooka K, Nagata N, Hasezawa S, Higaki T. Cell wall accumulation of fluorescent proteins derived from a trans-Golgi cisternal membrane marker and paramural bodies in interdigitated Arabidopsis leaf epidermal cells. Protoplasma. 2017:254:1367-377. https://doi.org/10.1007/s00709-016-0955-1
Google Scholar: Author Only Title Only Author and Title

Artimo P, Jonnalagedda M, Arnold K, Baratin D, Csardi G, de Castro E, Duvaud S, Flegel V, Fortier A, Gasteiger E, et al. ExPASy: SIB bioinformatics resource portal. Nucleic Acids Res. 2012:40:W597-603. https://doi.org/10.1093/nar/gks400
Google Scholar: Author Only Title Only Author and Title

Bailey T, Johnson J, Grant C, Noble W. The MEME Suite. Nucleic Acids Res. 2015:43:W39-49. https://doi.org/10.1093/nar/gkv416
Google Scholar: Author Only Title Only Author and Title

Brandizzi F, Snapp E, Roberts A, Lippincott-Schwartz J, Hawes C. Membrane protein transport between the endoplasmic reticulum and the Golgi in tobacco leaves is energy dependent but cytoskeleton independent: evidence from selective photobleaching. Plant Cell. 2002:14:1293-1309. https://doi.org/10.1105/tpc.001586
Google Scholar: Author Only Title Only Author and Title

Cai G, Parrotta L, Cresti M. Organelle trafficking, the cytoskeleton, and pollen tube growth. J Integr Plant Biol. 2015:57:63-78. https://doi.org/10.1111/jipb.12289
Google Scholar: Author Only Title Only Author and Title

Campanoni P, Blatt M. Membrane trafficking and polar growth in root hairs and pollen tubes. J Exp Bot. 2007:58:65-74. https://doi.org/10.1093/jxb/erl059
Google Scholar: Author Only Title Only Author and Title

Cao P, Renna L, Stefano G, Brandizzi F. SYP73 Anchors the ER to the Actin Cytoskeleton for Maintenance of ER Integrity and Streaming in Arabidopsis. Curr Biol. 2016:26:3245-3254. https://doi.org/10.1016/j.cub.2016.10.024
Google Scholar: Author Only Title Only Author and Title

Chen C, Wu Y, Li J, Wang X, Zeng Z, Xu J, Liu Y, Feng J, Chen H, He Y, et al. TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data mining. Mol Plant. 2023:16:1733-1742. https://doi.org/10.1016/j.molp.2023.09.010
Google Scholar: Author Only Title Only Author and Title

Chen J, Wang P, de Graaf B, Zhang H, Jiao H, Tang C, Zhang S, Wu J. Phosphatidic Acid Counteracts S-RNase Signaling in Pollen by Stabilizing the Actin Cytoskeleton. Plant Cell. 2018:30:1023-1039. https://doi.org/10.1105/tpc.18.00021
Google Scholar: Author Only Title Only Author and Title

Cheung A, Wu H. Structural and functional compartmentalization in pollen tubes. J Exp Bot. 2007: 58: 75-82. https://doi.org/10.1093/jxb/erl122
Google Scholar: Author Only Title Only Author and Title

Dinkel H, Van Roey K, Michael S, Kumar M, Uyar B, Altenberg B, Milchevskaya V, Schneider M, Kühn H, Behrendt A, et al. ELM 2016--data update and new functionality of the eukaryotic linear motif resource. Nucleic Acids Res. 2016:44:D294-300. https://doi.org/10.1093/nar/gkv1291
Google Scholar: Author Only Title Only Author and Title

Dobson L, Reményi I, Tusnády G. CCTOP: a Consensus Constrained TOPology prediction web server. Nucleic Acids Res. 2015:43:W408-412. https://doi.org/10.1093/nar/gkv451
Google Scholar: Author Only Title Only Author and Title

Fan J, Yang X, Lu J, Chen L, Xu P. Role of H(abc) domain in membrane trafficking and targeting of syntaxin 1A. Biochem Biophys Res Commun. 2007:359: 245-250. https://doi.org/10.1016/j.bbrc.2007.05.065
Google Scholar: Author Only Title Only Author and Title

Gao G, Banfield D. Multiple features within the syntaxin Sed5p mediate its Golgi localization. Traffic. 202021:274-296. https://doi.org/10.1111/tra.12720
Google Scholar: Author Only Title Only Author and Title

Gupta G, Free S, Levina N, Keränen S, Heath I. Two divergent plasma membrane syntaxin-like SNAREs, nsyn1 and nsyn2, contribute to hyphal tip growth and other developmental processes in Neurospora crassa. Fungal Genet Biol. 2003:40:271-286. https://doi.org/10.1016/s1087-1845(03)00109-9
Google Scholar: Author Only Title Only Author and Title

Han J, Pluhackova K, Böckmann R. The Multifaceted Role of SNARE Proteins in Membrane Fusion. Front Physiol. 2017:8:5. https://doi.org/10.3389/fphys.2017.00005
Google Scholar: Author Only Title Only Author and Title

Hatsuzawa K, Tamura T, Hashimoto H, Hashimoto H, Yokoya S, Miura M, Nagaya H, Wada I. Involvement of syntaxin 18, an

endoplasmic reticulum (ER)-localized SNARE protein, in ER-mediated phagocytosis. Mol Biol Cell. 2006:17: 3964-3977. https://doi.org/10.1091/mbc.e05-12-1174

Google Scholar: Author Only Title Only Author and Title

Hepler P, Winship L. The pollen tube clear zone: clues to the mechanism of polarized growth. J Integr Plant Biol. 2015:57: 79-92. https://doi.org/10.1111/jipb.12315

Google Scholar: Author Only Title Only Author and Title

Hirano T, Ebine K, Ueda T, Higaki T, Watanabe-Nakayama T, Konno H, Takigawa-Imamura H, Sato M. The SYP123-VAMP727 SNARE complex delivers secondary cell wall components for root hair shank hardening in Arabidopsis. Plant Cell. 2023:30;35(12):4347-4365. https://doi.org/10.1093/plcell/koad240

Google Scholar: Author Only Title Only Author and Title

Holthuis J, Nichols B, Pelham H. The syntaxin Tlg1p mediates trafficking of chitin synthase III to polarized growth sites in yeast. Mol Biol Cell. 1998:9: 3383-3397. https://doi.org/10.1091/mbc.9.12.3383

Google Scholar: Author Only Title Only Author and Title

Hong SY, So J, Lee J, Min K, Son H, Park C, Yun SH, Lee YW. Functional analyses of two syntaxin-like SNARE genes, GzSYN1 and GzSYN2, in the ascomycete Gibberella zeae. Fungal Genet Biol. 2010:47: 364-372. https://doi.org/10.1016/j.fgb.2010.01.005

Google Scholar: Author Only Title Only Author and Title

Hu J, Huang B, Yin H, Qi K, Jia Y, Xie Z, Gao Y, Li H, Li Q, Wang Z, et al. PearMODB: a multiomics database for pear (Pyrus) genomics, genetics and breeding study. Database (Oxford). 2023:baad050. https://doi.org/10.1093/database/baad050

Google Scholar: Author Only Title Only Author and Title

Ichikawa M, Iwano M, Sato M. Nuclear membrane localization during pollen development and apex-focused polarity establishment of SYP124/125 during pollen germination in Arabidopsis thaliana. Plant Reprod. 2015:28: 143-151. https://doi.org/10.1007/s00497-015-0265-3

Google Scholar: Author Only Title Only Author and Title

Jahn R, Scheller R. SNAREs--engines for membrane fusion. Nature reviews. Nat Rev Mol Cell Biol. 2006:7: 631-643. https://doi.org/10.1038/nrm2002

Google Scholar: Author Only Title Only Author and Title

Johnson L, Eddy S, Portugaly E. Hidden Markov model speed heuristic and iterative HMM search procedure. BMC Bioinformatics. 2010:11: 431.https://doi.org/10.1186/1471-2105-11-431

Google Scholar: Author Only Title Only Author and Title

Kasai K, Akagawa K. Roles of the cytoplasmic and transmembrane domains of syntaxins in intracellular localization and trafficking. J Cell Sci. 2001:114: 3115-3124. https://doi.org/10.1242/jcs.114.17.3115

Google Scholar: Author Only Title Only Author and Title

Kawabata M, Matsuo H, Koito T, Murata M, Kubori T, Nagai H, Tagaya M, Arasaki K. Legionella hijacks the host Golgi-to-ER retrograde pathway for the association of Legionella-containing vacuole with the ER. PLoS Pathog. 2021:17: e1009437. https://doi.org/10.1371/journal.ppat.1009437

Google Scholar: Author Only Title Only Author and Title

Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. Mol Biol Evol. 2016:33: 1870-1874. https://doi.org/10.1093/molbev/msw054

Google Scholar: Author Only Title Only Author and Title

Li B, Li Y, Liu F, Tan X, Rui Q, Tong Y, Qiao L, Gao R, Li G, et al. Overexpressed Tomosyn Binds Syntaxins and Blocks Secretion during Pollen Development. Plant Physiol. 2019:181:1114-1126. https://doi.org/10.1104/pp.19.00965

Google Scholar: Author Only Title Only Author and Title

Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in 2020. Nucleic Acids Res. 2021:49: D458-D460. https://doi.org/10.1093/nar/gkaa937

Google Scholar: Author Only Title Only Author and Title

Li X, Tang C, Li X, Zhu X, Cai Y, Wang P, Zhang S, Wu J. Cellulose accumulation mediated by PbrCSLD5, a cellulose synthase-like protein, results in cessation of pollen tube growth in Pyrus bretschneideri. Physiol Plant. 2022:174: e13700. https://doi.org/10.1111/ppl.13700

Google Scholar: Author Only Title Only Author and Title

Lipka V, Kwon C, Panstruga R. SNARE-ware: the role of SNARE-domain proteins in plant biology. Annu Rev Cell Dev Biol. 2007:23: 147-174. https://doi.org/10.1146/annurev.cellbio.23.090506.123529

Google Scholar: Author Only Title Only Author and Title

Lovy-Wheeler A, Cárdenas L, Kunkel J, Hepler P. Differential organelle movement on the actin cytoskeleton in lily pollen tubes. Cell Motil Cytoskeleton. 2007:64: 217-232. https://doi.org/10.1002/cm.20181

Google Scholar: Author Only Title Only Author and Title

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar G, Sonnhammer E, Tosatto S, Paladin L, Raj S, Richardson L, et al. Pfam: The protein families database in 2021. Nucleic Acids Res. 2021:49: D412-D419. https://doi.org/10.1093/nar/gkaa913
Google Scholar: Author Only Title Only Author and Title

Pan Q, Cui B, Deng F, Quan J, Loake G, Shan W. RTP1 encodes a novel endoplasmic reticulum (ER)-localized protein in Arabidopsis and negatively regulates resistance against biotrophic pathogens. New Phytol. 2016:209: 1641-1654. https://doi.org/10.1111/nph.13707
Google Scholar: Author Only Title Only Author and Title

Parton R, Fischer-Parton S, Watahiki M, Trewavas A. Dynamics of the apical vesicle accumulation and the rate of growth are related in individual pollen tubes. J Cell Sci. 2001:114: 2685-2695. https://doi.org/10.1242/jcs.114.14.2685
Google Scholar: Author Only Title Only Author and Title

Qi Z, Liu M, Dong Y, Zhu Q, Li L, Li B, Yang J, Li Y, Ru Y, Zhang H, et al. The syntaxin protein (MoSyn8) mediates intracellular trafficking to regulate conidiogenesis and pathogenicity of rice blast fungus. New Phytol. 2016:209: 1655-1667. https://doi.org/10.1111/nph.13710
Google Scholar: Author Only Title Only Author and Title

Ruan H, Li J, Wang T, Ren H. Secretory Vesicles Targeted to Plasma Membrane During Pollen Germination and Tube Growth. Frontiers in cell and developmental biology. 2020:8: 615447. https://doi.org/10.3389/fcell.2020.615447
Google Scholar: Author Only Title Only Author and Title

Rui Q, Tan X, Liu F, Li Y, Liu X, Li B, Wang J, Yang H, Qiao L, Li T, et al. Syntaxin of plants31 (SYP31) and SYP32 is essential for Golgi morphology maintenance and pollen development. Plant Physiol. 2021:186: 330-343.https://doi.org/10.1093/plphys/kiab049
Google Scholar: Author Only Title Only Author and Title

Sanderfoot A, Assaad F, Raikhel N. The Arabidopsis genome. An abundance of soluble N-ethylmaleimide-sensitive factor adaptor protein receptors. Plant Physiol. 2000:124: 1558-1569. https://doi.org/10.1104/pp.124.4.1558
Google Scholar: Author Only Title Only Author and Title

Schwarz D, Blower M. The endoplasmic reticulum: structure, function and response to cellular signaling. Cell Mol Life Sci. 2016:73: 79-94. https://doi.org/10.1007/s00018-015-2052-6
Google Scholar: Author Only Title Only Author and Title

Silva P, Ul-Rehman R, Rato C, Di Sansebastiano G, Malhó R. Asymmetric localization of Arabidopsis SYP124 syntaxin at the pollen tube apical and sub-apical zones is involved in tip growth. BMC Plant Biol. 2010:10:179. https://doi.org/10.1186/1471-2229-10-179
Google Scholar: Author Only Title Only Author and Title

Spang A. Retrograde traffic from the Golgi to the endoplasmic reticulum. Cold Spring Harb Perspect Biol. 2013:5. https://doi.org/10.1101/cshperspect.a013391
Google Scholar: Author Only Title Only Author and Title

Spang A, Schekman R. Reconstitution of retrograde transport from the Golgi to the ER in vitro. J Cell Biol. 1998:143:589-599. https://doi.org/10.1083/jcb.143.3.589
Google Scholar: Author Only Title Only Author and Title

Sparkes I, Runions J, Kearns A, Hawes C. Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. Nat Protoc. 2006:1: 2019-2025. 10.1038/nprot.2006.286
Google Scholar: Author Only Title Only Author and Title

Steer M, Steer J. Pollen tube tip growth. New Phytol. 1989:111: 323-358. https://doi.org/10.1111/j.1469-8137.1989.tb00697.x
Google Scholar: Author Only Title Only Author and Title

Su X, Yan X, Chen X, Guo M, Xia Y, Cao Y. Calcofluor white hypersensitive proteins contribute to stress tolerance and pathogenicity in entomopathogenic fungus, Metarhizium acridum. Pest Manag Sci 2021:77: 1915-1924. https://doi.org/10.1002/ps.6218
Google Scholar: Author Only Title Only Author and Title

Tang C, Wang P, Zhu X, Qi K, Xie Z, Zhang H, Li X, Gao H, Gu T, Gu C, et al. Acetylation of inorganic pyrophosphatase by S-RNase signaling induces pollen tube tip swelling by repressing pectin methylesterase. Plant Cell. 2023:1;35(9):3544-3565. https://doi.org/10.1093/plcell/koad162
Google Scholar: Author Only Title Only Author and Title

Teng F, Wang Y, Tang B. The syntaxins. Genome Biol. 2001:2: REVIEWS3012. https://doi.org/10.1186/gb-2001-2-11-reviews3012
Google Scholar: Author Only Title Only Author and Title

Wang H, Niu H, Zhai Y, Lu M BiPCharacterization of Genes from Pepper (L.) and the Role of in Response to Endoplasmic Reticulum and Multiple Abiotic Stresses. Front Plant Sci. 2017:8: 1122. https://doi.org/10.3389/fpls.2017.01122
Google Scholar: Author Only Title Only Author and Title

Wang M, Zhang H, Zhao X, Zhou J, Qin G, Liu Y, Kou X, Zhao Z, Wu T, Zhu J, et al. SYNTAXIN OF PLANTS81 regulates root meristem activity and stem cell niche maintenance via ROS signaling. Plant Physiol. 2023:191:1365-1382. https://doi.org/10.1093/plphys/kiac530

Google Scholar: Author Only Title Only Author and Title

Wang P, Hussey P NETWORKED 3B: a novel protein in the actin cytoskeleton-endoplasmic reticulum interaction. J Exp Bot. 2017:68: 1441-1450. https://doi.org/10.1093/jxb/erx047

Google Scholar: Author Only Title Only Author and Title

Watson R, Pessin J. Functional cooperation of two independent targeting domains in syntaxin 6 is required for its efficient localization in the trans-golgi network of 3T3L1 adipocytes. J Biol Chem. 2000:275: 1261-1268. https://doi.org/10.1074/jbc.275.2.1261

Google Scholar: Author Only Title Only Author and Title

Watson R, Pessin J. Transmembrane domain length determines intracellular membrane compartment localization of syntaxins 3, 4, and 5. Am J Physiol Cell Physiol. 2001:281: C215-223. https://doi.org/10.1152/ajpcell.2001.281.1.C215

Google Scholar: Author Only Title Only Author and Title

Weimbs T, Low S, Chapin S, Mostov K, Bucher P, Hofmann K. A conserved domain is present in different families of vesicular fusion proteins: a new superfamily. Proc Natl Acad Sci U S A. 1997:94: 3046-3051. https://doi.org/10.1073/pnas.94.7.3046

Google Scholar: Author Only Title Only Author and Title

Westrate L, Lee J, Prinz W, Voeltz G. Form follows function: the importance of endoplasmic reticulum shape. Annu Rev Biochem. 2015:84: 791-811. https://doi.org/10.1146/annurev-biochem-072711-163501

Google Scholar: Author Only Title Only Author and Title

Wu J, Wang Z, Shi Z, Zhang S, Ming R, Zhu S, Khan M, Tao S, Korban S, Wang H, et al. The genome of the pear (Pyrus bretschneideri Rehd.). Genome Res. 2013:23: 396-408. https://doi.org/10.1101/gr.144311.112

Google Scholar: Author Only Title Only Author and Title

Yang X, Xu P, Xiao Y, Xiong X, Xu T. Domain requirement for the membrane trafficking and targeting of syntaxin 1A. J Biol Chem. 2006:281: 15457-15463. https://doi.org/10.1074/jbc.M513246200

Google Scholar: Author Only Title Only Author and Title

Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden T. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. BMC Bioinformatics. 2012:13: 134. https://doi.org/10.1186/1471-2105-13-134

Google Scholar: Author Only Title Only Author and Title

Zheng P, Obara C, Szczesna E, Nixon-Abell J, Mahalingan K, Roll-Mecak A, Lippincott-Schwartz J, Blackstone C. ER proteins decipher the tubulin code to regulate organelle distribution. Nature. 2022:601: 132-138. https://doi.org/10.1038/s41586-021-04204-9

Google Scholar: Author Only Title Only Author and Title

Zhou H, Yin H, Chen J, Liu X, Gao Y, Wu J, Zhang S. Gene-expression profile of developing pollen tube of Pyrus bretschneideri. Gene Expr Patterns. 2016:20: 11-21. https://doi.org/10.1016/j.gep.2015.10.004

Google Scholar: Author Only Title Only Author and Title

Zhou X, Zheng Y, Wang L, Li H, Guo Y, Li M, Sun M, Zhao P. SYP72 interacts with the mechanosensitive channel MSL8 to protect pollen from hypoosmotic shock during hydration. Nat Commun. 2022:13: 73. https://doi.org/10.1038/s41467-021-27757-9

Google Scholar: Author Only Title Only Author and Title