Neutralizing gut-derived lipopolysaccharide as a novel therapeutic strategy for

severe leptospirosis

Xufeng Xie[a], Xi Chen[a], Shilei Zhang[a], Jiuxi Liu[a], Wenlong Zhang[a*], Yongguo Cao[a,b*]

[a]Department of Clinical Veterinary Medicine, College of Veterinary Medicine, Jilin

University, Changchun, 130062, People's Republic of China; [b]Key Laboratory for

Zoonosis Research, Ministry of Education, College of Veterinary Medicine, Jilin

University, Changchun, People's Republic of China

* Corresponding author. E-mail addresses: zwenlong123@126.com and

ygcao82@jlu.edu.cn

Abstract

Leptospirosis is an emerging infectious disease caused by pathogenic *Leptospira*

spp. Humans and some mammals can develop severe forms of leptospirosis

accompanied by a dysregulated inflammatory response, which often results in death.

The gut microbiota has been increasingly recognized as a vital element in systemic

health. However, the precise role of the gut microbiota in severe leptospirosis is still

unknown. Here, we aimed to explore the function and potential mechanisms of the gut

microbiota in a hamster model of severe leptospirosis. Our study showed that

leptospires were able to multiply in the intestine, cause pathological injury, and induce

intestinal and systemic inflammatory responses. 16S rRNA gene sequencing analysis

revealed that *Leptospira* infection changed the composition of the gut microbiota of

hamsters with an expansion of Proteobacteria. In addition, gut barrier permeability

was increased after infection, as reflected by a decrease in the expression of tight

junctions. Translocated Proteobacteria were found in the intestinal epithelium of

moribund hamsters, as determined by fluorescence in situ hybridization, with elevated

LPS levels in the serum. Moreover, gut microbiota depletion reduced the survival

time, increased the leptospiral load, and promoted the expression of proinflammatory

cytokines after *Leptospira* infection. Intriguingly, fecal filtration and serum from

moribund hamsters both increased the transcription of *TNF-α*, *IL-1β*, *IL-10*, and *TLR4*

in macrophages compared with those from uninfected hamsters. These stimulating

activities were inhibited by LPS neutralization using polymyxin B. Based on our

findings, we identified an LPS neutralization therapy that significantly improved the

survival rates in severe leptospirosis when used in combination with antibiotic therapy

or polyclonal antibody therapy. In conclusion, our study not only uncovers the role of

the gut microbiota in severe leptospirosis but also provides a therapeutic strategy for

severe leptospirosis.

Introduction

Leptospirosis is a worldwide emerging zoonotic disease caused by *Leptospira* spp.

(1). *Leptospira interrogans* impacts approximately one million people resulting 60000

deaths every year (1). In human leptospirosis, the clinical manifestations can range

from mild illness to multiorgan failure, characterized by jaundice, kidney, and liver

failure, and even death (2, 3). Currently, one important challenge is that leptospirosis

is often misdiagnosed as other febrile diseases, and once the optimal treatment period

is missed, leptospirosis quickly develops into the severe form, which often ends in

death (4). However, antibiotic treatment is often useless in severe leptospirosis and

even aggravates it, called the Jarisch-Herxheimer Reaction (5). Thus, it is imperative

to explore precision medicine strategies for the treatment of severe leptospirosis.

90% of infected humans may resolve spontaneously while 10% of patients are

designated as susceptible hosts (6). Intestinal bleeding is a common but neglected

symptom of severe leptospirosis in humans and hamster (7). A healthy gut

environment, shaped by the presence of a healthy functional microbial ecosystem, is

fundamental to instruct the immune system towards homeostasis. The intestine

harbours a densely populated resident microbial community, which consists of

bacteria, viruses, and fungi, called microbiota (8). Gut dysbiosis has been linked to

increased susceptibility to disseminated bacterial infections and sepsis in humans. In

addition, studies of patients in intensive care units (ICUs) have found that dysbiosis

increases the risk of nosocomial infections, sepsis, and mortality (9). Our previous

study proved that the gut microbiota contributed to the defence against *Leptospira*

infection in mice, a resistant model of leptospirosis, without obvious intestinal lesions

or increased intestinal permeability (10). Focusing on the role of the gut microbiota in

leptospirosis may help explain the differences in susceptibility to *Leptospira* infection

in humans and develop targeted therapy for acute leptospirosis. However, whether a

relationship exists between the gut microbiota and severe leptospirosis is still unclear.

In this study, we used the hamster model of leptospirosis to explore the functional

role of the gut microbiota in acute leptospirosis. We provided evidence that

Leptospira infection disrupted intestinal homeostasis, and targeting gut-derived LPS

improved the survival rates of hamsters with severe leptospirosis. Our study provides

a basis for the development of novel therapeutic strategies to control severe

leptospirosis.

Materials and methods

Ethics statement

A previous study reported that sex influenced host susceptibility to leptospirosis (11)

and that sex may affect the composition of the gut microbiota (12). To reduce

variability in infection outcomes (13), most infection experiments were performed in

female Syrian golden hamsters *(Mesocricetus auratus)*. For experiments utilizing

neutralizing LPS therapy combined with antibiotic or antibody therapy for severe

leptospirosis, both sexes were used. Specific pathogen-free female Syrian hamsters

were maintained on standard rodent chow with water supplied ad libitum during the

experimental period. All animal experiments were conducted according to the

regulations of the Administration of Affairs Concerning Experimental Animals in

China. The protocol was approved by the Institutional Animal Care and Use

Committee of Jilin University (20170318).

Bacterial strains and animals

Pathogenic *L. interrogans* serovar Lai strain Lai (56601) was grown in liquid Ellinghausen-McCullough-Johnson-Harris (EMJH) medium at 29 °C without agitation. A Petroff-Hauser chamber was used to count leptospires. Six-week-old female hamsters were provided by Liaoning Changsheng Biotechnology Co. Ltd.

Experimental infections

Part I hamster infection

To explore the effect of challenge route on leptospiral colonization in the intestine and the survival rate, six-week-old female hamsters were injected intraperitoneally or subcutaneously with 10^7 leptospires. Hamsters were euthanized at 0 h, 6 h, 48 h, 96 h post infection (p.i.), and articulo mortis stage (AM). The ileums and colons were collected aseptically. The intestinal contents were excluded from the intestine with PBS, and the tissues were stored at -80 °C for leptospiral load analysis. For the survival assay, hamsters were monitored daily for signs of illness including ruffled fur, gait difficulty, loss of appetite, or weight loss of ≥10% of the animal's maximum weight, which was considered moribund (4). Hamsters were euthanized when they were moribund.

To explore the effect of *Leptospira* infection on the gut microbiota, six-week-old female hamsters were injected intraperitoneally with a lethal dose of 10^7 leptospires. Fresh fecal pellets were collected aseptically at 0 days, 2 days, and AM p.i., immediately frozen in liquid nitrogen, and stored at −80 °C. Colons were also collected aseptically and the intestinal contents were excluded from the intestine with PBS. Then, tissues were stored at -80 °C for gene expression measurement.

To examine the role of the gut microbiota on acute leptospirosis, the gut microbiota depletion and fecal microbiota transplantation (FMT: a therapeutic procedure aimed at restoring a normal intestinal microbiota by application of fecal microorganisms from

a healthy subject into the gastrointestinal tract of a dysbiotic subject (14)) were

performed as described previously with some modifications (15). Briefly,

six-week-old female hamsters were administered antibiotics (ampicillin, 100 mg/kg;

metronidazole, 100 mg/kg; neomycin sulfate, 100 mg/kg (Sigma-Aldrich, USA) and

vancomycin, 50 mg/kg (BiochemPartner, China) intragastrically once daily for ten

consecutive days (16). Fresh fecal pellets were collected from uninfected female

hamsters and then resuspended in sterile normal PBS (1 fecal pellet in 1 mL of PBS).

The pellets were immediately homogenized and the homogenate was centrifuged at

100 rpm, 4 °C for 5 min. 200 μL of the supernatant was given to the gut

microbiota-depleted hamster by oral gavage for 5 consecutive days after antibiotic

treatment was stopped. Then, hamsters were intraperitoneally infected with 10^6 *L.*

interrogans. Hamsters were monitored daily for 21 days to record the survival rate.

Hamsters were euthanized when they were moribund. Blood was collected to

determine leptospiral load and the gene expression of inflammatory cytokines at 6

days p.i..

To explore the effect of LPS neutralization on severe leptospirosis, six-week-old

male or female hamsters were inoculated intraperitoneally with 10^7 leptospires.

Treatments were started immediately after detection of the first death regardless of the

assigned group (17). Group 1: saline control (400 μL/hamster, i.p.); Group 2:

Polymyxin B (PMB) (1 mg/kg, i.p., Solarbio, China); Group 3: Antibody against

Leptospira (Ab) (16 mg/kg, subcutaneous injection); Group 4: Doxycycline (Dox) (5

mg/kg, i.p., Solarbio, China); Group 5: PMB (1 mg/kg, i.p.) and Ab (16 mg/kg,

subcutaneous injection); Group 6: PMB (1 mg/kg, i.p.) and Dox (5 mg/kg, i.p.).

Hamsters were treated twice a day for 3 consecutive days. Hamsters were monitored

daily for 21 days. Hamsters were euthanized when they were moribund. Blood from

female hamsters was collected to determine the leptospiral load and gene expression

of inflammatory cytokines at 6 days p.i..

Part II rabbit infection

The purification of polyclonal antibody (Ab) was performed as previously

described (18). Briefly, six eight-month-old female New Zealand White rabbits were

injected intravenously into the marginal ear vein with a dose of $2 \pm 4\times10^8$ live

leptospires/mL in PBS according to the following schedule: day 1, 1 mL; day 6, 2 mL;

day 11, 4 mL; and day 16 and 21, 6 mL each. One week after the last injection, the

rabbits were anaesthetized and then exsanguinated through cardiocentesis. The rabbits

were then euthanized. During the immunization period, the rabbits did not exhibit

clinical signs of leptospirosis. Antisera were collected by centrifuging blood at 3,000

g. The polyclonal Ab IgG was purified from the antisera using the caprylic acid

ammonium sulfate precipitation method from the antisera. The concentration of the

IgG-polyclonal Ab was determined with the BCATM Protein Assay Kit (Thermo,

USA) and adjusted to a final concentration of 20 mg/mL with phosphate buffered

saline (PBS). The Ab was stored at -20 °C until used for treatment. The protocol was

approved by the Committee on the Ethics of Animal Experiments of the First Norman

Bethune Hospital of Jilin University, China [(2013) clinical trial (2013-121)] (18).

Bacterial isolation and culture

For isolation of the translocated microbiota, six-week-old female hamsters were

intraperitoneally infected with 10^7 leptospires. Hamsters were anaesthetized by

intraperitoneal injection of a mix of ketamine/xylazine (200 and 10 mg/kg,

respectively). Blood from the uninfected group and AM group by cardiac puncture

were plated on LB or MacConkey agar plates aerobically or anaerobically at 37 °C for

36 h.

In vivo **barrier permeability**

Six-week-old female hamsters were intraperitoneally infected with 10^7 leptospires.

The permeability of the intestine was determined at 0 days, 2 days, and AM p.i..

Hamsters were starved for 6 h before gavage with FITC-dextran (4 kDa, 400 mg/kg

body/hamster). After 2 h, hamsters were anaesthetized by intraperitoneal injection of a

mix of ketamine/xylazine (200 and 10 mg/kg, respectively) and blood was collected

and centrifuged at 3000 rpm and 4 °C for 10 min. The samples were analyzed with a

fluorescence spectrophotometer (Synergy II plate reader with Gen5 software; BioTek

Instruments, Winooski, VT) (19).

Minimum inhibitory concentration (MIC) determination

The MIC of PMB and Dox against *L. interrogans* serovar Lai strain Lai 56601 was

determined according to a previous method (20). Briefly, leptospires in the

exponential-phase were deposited at a final concentration of 2×10^6 leptospires/mL in

each well of 96 well plates with serial two-fold dilutions of PMB and Dox ranging

from to 0.0156 μg/L in EMJH. The final volume was 200 μL in each well. The plates

were incubated for 3 days at 30 °C, and then 20 μL of Alamar Blue (Invitrogen,

Thermo Fisher Scientific) was added to each well. Then, the samples were incubated

at 30 °C for 2 days. Each strain–drug combination was tested in duplicate, and

positive (bacteria and no antibiotic added) and negative (no bacteria added) controls

were included in each plate. The results were recorded by a microplate reader

(MultiSkan FC, Thermo Fisher Scientific).

Limulus amebocyte lysate assay

Blood from uninfected hamsters and AM hamsters was collected and centrifuged at

3000 rpm for 10 min. Then, serum was diluted (1:10) and detected by the Limulus

amebocyte lysate assay according to the manufacturer's instructions (LAL

Chromogenic Endotoxin Quantitation Kit; Xiamen Biotechnology Co., Ltd., China). *E.*

coli LPS at 1 EU/mL and *E. coli* LPS at 10 EU/mL (Sigma-Aldrich, USA) were used

as positive controls.

Fluorescence in situ hybridization (FISH)

Six-week-old female hamsters were injected intraperitoneally with 10^7 leptospires.

Hamsters were euthanized at 0 days, 2 days, and AM p.i.. Colons were collected

aseptically. The intestinal contents were excluded from the intestine with PBS and a

segment of colon was fixed in 4% paraformaldehyde solution overnight, washed and

passed through 15% and 30% sucrose solutions. The sample was then embedded in

optimal cutting temperature compound (O.C.T., Tissue-Tek) and cryo-sectioned into 5

μm longitudinal sections (Leica). Slides were equilibrated in hybridization buffer (0.9

M NaCl, 20 mM Tris-HCl, 0.01% sodium dodecyl sulfate, 10% formamide, pH 7.5)

and incubated in Cy3-lablled 10 ng/μl FISH probe (Comate Bioscience Co.,Ltd.,

China) for 14 h at 42 °C in a humidified chamber. The probe information is listed in

Table S1. Slides were then incubated for 20 min in wash buffer (0.9 M NaCl, 20 mM

Tris-HCl, pH 7.5) preheated to 42 °C and washed three times. Samples were then

incubated in the dark with 10 μg/ml DAPI (Servicebio Technology Co., Ltd., China)

in PBS for 10 min at room temperature, washed three times with PBS and mounted

with Vectashield mounting medium (Vector Labs). Images were acquired on a Nikon

A1 confocal microscope.

FISH and immunofluorescence (IF) double labelling (FISH/IF)

After the preparation and digestion of paraffin-embedded sections and before

hybridization, the probe hybridization solution was added at a concentration of 1 μM

and the sections were incubated in a humidified chamber to hybridize with the

Cy3-lablled 10 ng/μL FISH probe (Comate Bioscience Co.,Ltd., China) for 14 h at

42 °C. Then, the sections were washed, blocked, and incubated with rabbit anti-*L.*

interrogans serovar Lai strain Lai 56601 (1:500) as the primary antibody overnight at

4 °C. The next day, the samples were washed with PBS three times for 5 min. After

washing, the samples were blocked with 5% BSA and incubated at room temperature

for 1 h with the secondary antibody. Samples were then incubated in the dark with 10

μg/ml DAPI (Servicebio Technology Co., Ltd., China) in PBS for 10 min at room

temperature, washed three times with PBS and mounted with Vectashield mounting

medium (Vector Labs) (21, 22). Images were acquired on a Nikon A1 confocal

microscope.

Isolation of primary macrophages

Primary macrophages were isolated as previously described (10). Six-week-old

female hamsters were injected intraperitoneally with 2 mL of 3% thioglycolate.

Hamsters were euthanized after 4 days, and macrophages were lavage by sterile PBS

and enriched by plating the lavage cells on tissue culture plates in RPMI 1640

supplemented with 10% FCS and 1% penicillin and streptomycin at 37 °C for 2 h.

Then, the plate was washed 3 times with sterile PBS to remove nonadherent cells.

Culture of primary macrophages with fecal filtration or serum

Fecal filtration or serum were collected from uninfected hamsters and AM hamsters.

Fecal pellets were homogenized using PBS (1 mL per 0.1 g of fecal matter). The

homogenate was centrifuged at 100 rpm and 4 °C for 5 min and filtered using 0.22

μm syringe filters. The quantity of the fecal filtration or serum were adjusted to 5% or

20% of the total volume to stimulate macrophages *in vitro*. After incubation at 37 °C

for 4 h, RNA was extracted by TRIzol.

To determine the role of LPS in the induction of cytokine release, fecal filtration or

serum were pre-incubated with LPS neutralizing reagent polymyxin B (PMB;

Solarbio, China) for 2 h at 37 °C resulting in a final concentration of 50 μg/mL PMB

after addition to the macrophages. After 4 h of incubation at 37 °C, RNA was

extracted with TRIzol.

Real-time and reverse transcription-qPCR (RT-qPCR)

Total RNA of the colon samples was extracted by TRIzol (Invitrogen, USA). RNA

was reverse transcribed into cDNA by using random primers from a TransScript

One-Step gDNA Removal kit and cDNA Synthesis SuperMix (TransGen Biotech,

China). The primers used in this study are listed in Table S1. The PCR conditions

were as follows: 50 °C for 2 min, 95 °C for 10 min, followed by 45 cycles of

amplification at 95 °C for 15 s and 60 °C for 60 s (23). qPCR was conducted by an

Applied Bioscience 7500 thermocycler and FastStart Universal SYBR Green Master

Mix (Roche Applied Science, Germany). The expression of target genes was

normalized to that of GAPDH using the $2^{-\Delta\Delta CT}$ method.

Bacterial load and qPCR assay

The leptospiral burdens in organs were determined by quantitative PCR using an

Applied Bioscience 7500 thermocycler and FastStart Universal SYBR green Master

(Roche Applied Science, Germany). The specimens (0.09 to 0.15 g) of tissues were

homogenized with PBS (w/v, 1/10). The homogenate was centrifuged at 2000 rpm

and 4 °C for 5 min. The supernatant was transferred into a new tube. Then, it was

centrifuged at 12000 rpm and 4 °C for 5 min. The pellets were extracted using the

TIANamp Bacteria DNA kit (Tiangen, China) according to the manufacturer's

instructions (24). The concentration of DNA was measured by spectrometry. The

genomic DNA of a counted number of *L. interrogans* was used as a calibrator. The

primers used in this study were listed in Table S1. The leptospiral load was presented

as the number of genome equivalents per μg of tissue DNA (25).

Histopathological examination

Colons from the uninfected group and AM group were collected and fixed in

buffered 4% formaldehyde. Hematoxylin and eosin (H&E) staining was performed on

4–5 μm paraffin-embedded sections. Blinded samples were then scored by two

pathologists. Each section was evaluated for inflammation, epithelial hyperplasia,

erosion and ulceration, and the extent of lesion as listed in Table S2 (26).

Microbiota assay

Fresh fecal pellets were collected under sterile conditions, immediately frozen in liquid nitrogen, and then stored at −80 °C. Total genomic DNA from the samples was extracted using the CTAB/SDS method. DNA concentration and purity were monitored on 1% agarose gels. Then, 16S rRNA genes of distinct regions were amplified using specific primers with barcodes (27F, AGAGTTTGATCMTGGCTCAG; 1492R, ACCTTGTTACGACTT). All PCRs were carried out with TransStart FastPfu DNA Polymerase (TransGen Biotech). PCR products were purified using the Qiagen Gel Extraction Kit (Qiagen, Germany). Sequencing libraries were generated using the SMRTbellTM Template Prep Kit (PacBio) following the manufacturer's recommendations. The library quality was assessed on the Qubit 2.0 Fluorometer (Thermo Scientific) and FEMTO Pulse system. Finally, the library was sequenced on the PacBio Sequel platform. Raw sequences were initially processed through the PacBio SMRT portal. Sequences were filtered for a minimum of 3 passes, and a minimum predicted accuracy of 90% (minfullpass = 3, minPredictedAccuacy = 0.9). The predicted accuracy was 90%, which was defined as the threshold below which CCS was considered noise. The files generated by the PacBio platform were then used for amplicon size trimming to remove sequences outside the expected amplicon size (minLength 1340 bp, maxLength 1640 bp). The reads were assigned to samples based on their unique barcode and trimmed by removing the barcode and primer sequence. The reads were compared with the reference database using the UCHIME algorithm to detect chimeric sequences, and then the chimeric sequences were removed to obtain clean reads. Alpha diversity (Chao1 and Shannon) was calculated with QIIME (Version 1.9.1) and displayed with R software (Version 2.15.3). Principal coordinate analysis (PCoA) was performed to obtain principal coordinates and visualize the complex, multidimensional data. A distance matrix of weighted UniFrac among samples obtained before was transformed to a new set of orthogonal axes, by which the maximum variation factor was

demonstrated by the first principal coordinate, and the second maximum variation

factor was demonstrated by the second principal coordinate. PCoA analysis results

were displayed by the WGCNA package, stat packages and ggplot2 package in R

software (Version 2.15.3).

Statistical analysis

Survival differences between the study groups were compared by using the

Kaplan-Meier log-rank test. All values are expressed as the mean ± SEM. Differences

between mean values of normally distributed data were analyzed using the Wilcoxon

rank-sum test. The results were considered statistically significant at $p < 0.05$.

Results

L. interrogans **proliferated in the intestine, impaired the intestinal structure, and**

induced proinflammatory response

Recently, Inamasu *et al*. reported that subcutaneous infection with leptospires led to

intestinal dissemination (27). We wondered whether different challenge routes

affected the dissemination of *Leptospira* into the intestine. Thus, we compared the

common intraperitoneal challenge with subcutaneous challenge in a hamster model of

leptospirosis. Our study indicated that subcutaneous challenge and intraperitoneal

challenge both led to leptospiral proliferation in the ileum and colon (Supplementary

Fig. S1A and Fig. 1B). There were approximately 7 times more leptospires in the

colon than in the ileums at AM p.i. after intraperitoneal challenge (Fig. 1B). Hamsters

infected by intraperitoneal challenge had shortened survival time compared with

hamsters infected by subcutaneous challenge (Supplementary Fig. S1B). *L.*

interrogans infection disrupted the intestinal structure, caused bleeding, and led to the

infiltration of inflammatory cells (Fig. 1C). In addition, *L. interrogans* infection

promoted the inflammatory response in the colons as determined by the upregulated

gene expression of *lipocalin 2* (*Lcn2*)/*neutrophil gelatinase-associated lipocalin*

(*NGAL*), a marker of intestinal inflammation (28), *TNF-α*, *IL-1β*, *Nos2*, and *TLR4* (a

well-known receptor of LPS that is vital against *Leptospira* infection (29)) and the

downregulated gene expression of *IL-10* (Fig. 1D-I). We wondered whether

proinflammatory gene expression was also related to other TLRs (30). The results

showed that the gene expression of *TLR2* and *TLR9* was significantly increased, while

TLR3, *TLR5*, and *TLR7* showed no obvious difference in the AM group compared

with the D0 group (Supplementary Fig. S2A-E). It is well-known that *Leptospira*

infection can cause leptospiremia and cytokine storms at late stages (6, 31). Our

results also proved that the gene expression of *TNF-α*, *IL-1β*, and *IL-10* were

increased in the blood of the AM group compared with that of D0 group

(Supplementary Fig. S2F-H). These results indicated that *L. interrogans* was able to

proliferate in the intestine and impair the intestinal structure, along with inducing

intestinal inflammation and systemic inflammation.

The disruption of the gut microbiota after *L. interrogans* infection

To obtain insight into the composition of the gut microbiota during leptospirosis,

we analyzed the feces from *Leptospira*-infected hamsters by 16S rRNA gene

sequencing at the indicated time points (Fig. 2A). The results showed that the

bacterial richness (Observed species and Ace) and the diversity (Simpson index and

Shannon index) of the gut microbiota were decreased at 2 days p.i., while they were

increased at AM p.i. (Fig. 2B-E). Then, we pooled all groups together to perform

principal coordinate analysis (PCoA). The results showed that the composition of the

gut microbiota in the AM group was significantly different from that in the D0 group

(Fig. 2G). There was a large degree of dispersion among different individuals 2 days

p.i. (Fig. 2F), and the composition of the gut microbiota in the D2 group was also

different from that of the other two groups (Supplementary Fig. S3A-B). To identify

bacterial taxa that were common in the D0 group or the AM group, we performed

linear discriminant analysis effect size (LEfSe) analysis. The results showed that the

abundance of Proteobacteria was increased in the AM group (Fig. 2I and 2K), while

the abundances of *Lactobacillus* and *Allobaculum* were decreased compared with

those in the D0 group (Fig. 2I, 2L, and 2M). The composition of the bacterial taxa in

the D2 group was also various compared with that of other two groups, especially the

abundance of *Lactobacillus* was also decreased in the AM group compared with that

of the D2 group (Supplementary Fig. S3C-D). Although the relative abundance of

Firmicutes seemed to gradually decrease after *Leptospira* infection (Fig. 2H), the ratio

of Firmicutes to Bacteroidetes showed no difference in the D0 and AM groups (Fig.

2J). These results demonstrated that *L. interrogans* infection changed the composition

of the gut microbiota of hamsters.

L. interrogans infection increased intestinal permeability by disrupting tight

junctions

The intestinal barrier is essential for maintaining intestinal homeostasis (32).

Epithelial tight junctions define the paracellular permeability of the intestinal barrier,

which is related to some tight junction proteins, such as Claudins, Occludin, and Zona

Occludens 1 (Zo-1) (33). The epithelium separates the organism from the external

environment and defines individual compartments within tissues. At some sites, the

epithelial cells form a nearly complete barrier, the disruption of which is catastrophic

(33). Most studies have relied on a probe, FITC-4 kDa dextran, which has a

hydrodynamic diameter of 28 Å, to probe intestinal permeability (34-36). As shown in

Fig. 1B, we detected leptospiral proliferation in the gut. We wondered whether

Leptospira infection could influence intestinal permeability.   The results showed that

the intestinal permeability of hamsters was increased significantly after infection (Fig.

3A). Then, we further investigated the gene expression of tight junction proteins and

Mucin-2 in hamsters. The results showed that the gene expression of *Claudin-2* was

increased, while the gene expression of *Claudin-3*, *JAMA*, *ZO-1* and *Mucin-2* was

decreased in the AM group compared with the D0 group (Fig. 3B-3F).

L. interrogans infection caused the translocation of Proteobacteria

Damaged intestinal structure may increase the translocation of pathobionts (37, 38) and our 16S rRNA sequencing showed an increase in Proteobacteria in the AM group. We hypothesized that the pathogenic Proteobacteria could translocate into the intestinal epithelium or bloodstream after infection. First, we found that the intestinal epithelium of the AM group had more bacteria than that of the D0 and D2 groups, as determined by the EUB 338 probe, which was supposed to hybridize with all bacteria (Fig. 4). Then, we examined the intestines of the D0 group and the AM group by FISH assays with the GAM42a probe, which was supposed to hybridize γ-Proteobacteria, and IF assays with the anti-*L. interrogans* serovar Lai strain Lai 56601 antiserum. The results showed that the intestinal epithelium of the AM group indeed had γ-Proteobacteria that was not colocalized with leptospires (Supplementary Fig. S4A). Then, we tested whether the gut microbiota could translocate into the blood. However, no bacteria were recovered from the blood by aerobic or anaerobic culture on LB and MacConkey agar plates (Supplementary Fig. S4B).

Gut microbiota depletion exacerbated leptospirosis

To determine the role of the gut microbiota in hamsters during *L. interrogans* infection, we treated hamsters with broad-spectrum antibiotics to deplete the gut microbiota, as shown in Fig. 5A. 16S rRNA sequencing analysis indicated that the richness of the gut microbiota decreased after antibiotic treatment (Fig. 5B-C). In addition, the diversity of the gut microbiota in the Abx-treated hamsters was lower than that of the control group, and this change in diversity was reversed by FMT (Fig. 5D-E). Antibiotic treatment changed the composition of the gut microbiota in hamsters (Fig. 5F, Supplementary Fig. S5A-D), leading to the expansion of Proteobacteria and a decrease in Bacteroidota (Fig. 5F-G). The survival time of Abx-treated hamsters was significantly reduced after infection with 10^6 leptospires

(Fig. 5I). In addition, we also measured leptospiral load and the gene expression of

inflammatory cytokines in the blood. The results showed that the leptospiral load of

the Abx group was significantly increased compared with that of the control group,

and this increase was partially reversed by FMT (Fig. 5J). Abx treatment slightly

increased the gene expression of *TNF-α* and *IL-1β* compared with those of the

untreated group. Consistent with this result, the Abx-treated infection group exhibited

higher expression of *TNF-α* and *IL-1β* than the untreated infection group, which was

also partially reversed by FMT (Fig. 5K-L). Although *Leptospira* infection

significantly increased the gene expression of *IL-10* compared with that of the

untreated group, there was no significant difference among the control infection group,

the Abx-treated infection group, and the FMT infection group (Fig. 5M). These results

indicated that gut microbiota homeostasis helped fight *Leptospira* infection in

hamsters.

Gut-derived LPS in the intestine and serum induced inflammation

Gut dysbiosis is a major determinant of endotoxaemia via dysfunction of the

intestinal barrier scaffold, which is a prerequisite for LPS translocation into systemic

circulation (39). A previous study demonstrated that gut-derived LPS was a central

mediator in alcoholic steatohepatitis (40), white adipose tissue inflammation (41), and

peripheral inflammation (42). Cytokine storms are characteristic of severe

leptospirosis (6). Our results showed that the proportion of Proteobacteria and

intestinal permeability were increased in the AM group; thus, we hypothesized that

gut-derived LPS promoted the inflammatory response in severe leptospirosis. We

found that the LPS level was higher in the serum of the AM group than in that of the

D0 group (Supplementary Fig. S6B). The LPS level of in the Abx-treated AM group

was lower than that in the PBS-treated AM group (Supplementary Fig. S6B). To

further gain insight into the role of gut-derived LPS in inflammation, we cocultured

macrophages with different concentrations of fecal filtration or serum and measured

the changes in gene expression (Fig. 6A). The results showed that fecal filtration from

uninfected hamsters and AM hamsters both upregulated the gene expression of *TNF-α*,

IL-1β, *IL-10*, and *TLR4* (Fig. 6B-E). The serum of uninfected hamsters significantly

increased the transcription of *TNF-α* (Fig. 6F), while the serum of AM hamsters only

slightly upregulated the gene expression of *TNF-α* (Fig. 6F). Serum from uninfected

hamsters and AM hamsters promoted the transcription of IL-1β (Fig. 6G). However,

only the serum of AM hamsters significantly upregulated the gene expression of *IL-10*

and *TLR4* (Fig. 6H-I). The identification of TLR4, a well-known receptor of LPS,

prompted us to examine whether the proinflammatory activity of fecal filtration and

serum of AM hamsters was mediated by LPS. A previous study demonstrated that

cytokine responses to heat-killed leptospires were not inhibited by PMB (29). Our

results showed that LPS neutralization with PMB inhibited gene expression of *TNF-α*,

IL-1β, *IL-10*, and *TLR4* in macrophages after stimulation by fecal filtration or serum

from AM hamsters (Fig. J-Q). These results indicated that gut-derived LPS might be

an essential factor that causes inflammatory storm in hamsters with severe

leptospirosis.

LPS neutralization synergized with antibiotic therapy or polyclonal antibody

therapy improved the survival rate in severe leptospirosis

Based on the above results, we proposed that in addition to leptospires, gut-derived

LPS might also play a pathogenic role in severe leptospirosis and induce an

uncontrolled inflammatory response, in which case neutralizing LPS could be

effective in treating severe leptospirosis. Therefore, we explored the effect of LPS

neutralization therapy compared with antibiotic therapy and our published rabbit

polyclonal antibody therapy in severe leptospirosis (18). As shown in Fig. 7A,

moribund hamsters were treated twice a day for 3 consecutive days. LPS

neutralization therapy, antibiotic therapy, and rabbit polyclonal antibody therapy did

not significantly improve the conditions of female hamsters with severe leptospirosis.

However, LPS neutralization combined with antibiotic therapy or rabbit polyclonal

antibody therapy significantly increased the survival rate up to 50% or 66.7% (Fig.

7B). To further explore the factors leading to differences in survival, we determined

the leptospiral load and gene expression of inflammatory cytokines in the blood at 6

days p.i.. The results showed that antibiotic therapy and rabbit polyclonal antibody

therapy decreased the leptospiral load in the blood, while LPS neutralization therapy

did not reduce the leptospiral load in the blood (Fig. 7C). Although antibiotic therapy

reduced the leptospiral load, it significantly increased the gene expression of

proinflammatory cytokines (*TNF-α* and *IL-1β*) compared with that of the control

group (Fig. 7D-E). LPS neutralization therapy combined with *Leptospira*-killing

therapy decreased the gene expression of inflammatory cytokines (*TNF-α*, *IL-1β* and

IL-10) compared with that of the control group (Fig. 7D-F). Finally, we examined the

effectiveness of LPS neutralization therapy in male hamsters with severe leptospirosis.

The results showed that LPS neutralization synergized with antibiotic therapy or

polyclonal antibody therapy to improve the conditions of male hamsters with severe

leptospirosis (Supplementary Fig. S8B). Our results indicated that LPS neutralization

could be a novel therapy for severe leptospirosis.

Discussion

Recently, Inamasu *et al*. investigated the pathological changes and distribution of

leptospires in the gut of hamsters after subcutaneous infection (27). Their study

highlighted that not only the urine of infected animals but also the feces could be a

source of infection. However, the exact role of the gut microenvironment in the

pathogenesis of leptospirosis is still unknown. In our study, we found that leptospires

could proliferate in the ileum and colon following intraperitoneal challenge or

subcutaneous challenge, which was consistent with the results of Inamasu's study (27).

Hamsters infected intraperitoneally exhibited more rapid death than hamsters infected

subcutaneously. A previous study reported that the route of infection affected the

kinetics of hematogenous dissemination, kidney colonization, and inflammation (43).

In addition, the subcutaneous route induced a slower-progressing infection than the

intraperitoneal route in a leptospirosis rat model (44). However, there was a higher

leptospiral load in the colons than in the ileums of hamsters infected by the

intraperitoneal route. The capacity of *L. interrogans* to breach the immune defence at

each port of entry determined when *Leptospira* gained access to the blood. *Leptospira*

infection destroyed the intestinal structure and induced intestinal inflammation in the

colon. However, Inamasu *et al*. reported that *Leptospira* was not involved in the

pathogenesis of the colon, suggesting that the jejunum and ileum are the main sites of

lesions. Differences in bacterial strains and the route of infection might lead to

variations in results. Infection via the intraperitoneal and subcutaneous routes bypass

the epidermal defence system. Early damage to the intestinal structure and

environment may be one of the causes of rapid death following infection by

intraperitoneal injection. The proliferation of leptospires in the intestine makes feces a

potential excretion route. Our previous study demonstrated that leptospires were

rapidly cleared in the intestine and that *Leptospira* infection did not increase the

intestinal permeability of mice (10). A recent study reported that pathogenic

Leptospira spp. were highly present in human and gorilla stool samples in the Congo

but were absent in other African nonhuman primates, therefore suggesting a possible

gorilla-humans transfer (45). Our 16S rRNA sequencing results also showed that the

relative abundance of Spirochetes was increased in the AM group compared with that

of the D0 group, which might be related to the excretion of leptospires in the feces.

However, no leptospires were detected in the feces from *Leptospira*-infected rats (44).

Serovar specificity and species differences may determine the excretion route of

leptospires and the disruption of intestinal homeostasis may influence the progression

of leptospirosis.

Our 16S rRNA sequencing results showed that the species richness and diversity of

the gut microbiota in the AM group were even higher than those in the D2 group. The

species richness and diversity of the gut microbiota decreased first and then increased

with disease progression. This phenomenon was also reported in melioidosis and

HIV-1 infection (46, 47). Lankelma *et al.* explained that several "big players" in the

gut microbiota were eliminated by the host immune system during severe systemic

bacterial infection, giving way to the proliferation of other bacteria (46). Specific

microbiota community composition, rather than species richness, drives

microbiota-mediated resistance against *Leptospira* infection. Becattini *et al*.

demonstrated that microbiota composition and the type of immune stimulus impacted

bacterial responses and that the intestinal metabolome was rapidly reshaped by host

immune activation (48). We hypothesized that the early colonization and proliferation

of leptospires in the intestine rapidly activated intestinal immunity leading to the

expansion of specific microbes in the AM group. The composition of the gut

microbiota in the D2 group was discrete, as shown by PCoA. This difference might be

related to the disease progression. In addition, we found that the levels of

Lactobacillus and *Allobaculum* were higher in the D0 group than in the AM group at

the genus level. Meanwhile, the level of Proteobacteria increased in the AM group.

Our study proved that *Leptospira* infection increased the gene expression of *TLR2*,

TLR4, *TLR9*, and some proinflammatory cytokines, which created a proinflammatory

environment in the intestine. A previous study demonstrated that intestinal

inflammation increased the bioavailability of respiratory electron acceptors for

Enterobacteriaceae, which promoted the expansion of pathogenic Enterobacteriaceae

(49). Therefore, it is plausible that *Leptospira* infection triggered intestinal

inflammation, which promoted the expansion of Proteobacteria by increasing the

bioavailability of respiratory electron acceptors for pathogenic Proteobacteria.

The intestinal barrier is critical for maintaining homeostasis in the host. The apical

junctional complex (AJC) encircles pairs of neighbouring epithelial and endothelial

cells to create an adhesive network and maintain barrier integrity (50). Our results

showed that *Leptospira* infection significantly disrupted the intestinal permeability of

hamsters at the late stage of infection. Previous studies have demonstrated that

Leptospira infection reduced the signalling between cell-cell junction proteins and

caused the mislocalisation of AJC proteins, such as occludin and Zo-1 (51). A recent

study reported that *L. interrogans* could disassemble AJCs in renal proximal tubule

epithelial cells and transmigrate through the paracellular route for dissemination in the

host (51). Thus, we hypothesize that *L. interrogans* was also able to transmigrate

through the intestinal epithelial barrier, and feces excretion route of leptospires was

possible in susceptible hosts during leptospirosis. Our results showed that the gene

expression of *Claudin-2*, *Claudin-3*, *Claudin-4*, *Zo-1* and *Mucin-2* were

compensatively upregulated in the D2 group compared with those in the D0 group.

However, the gene expression of *Claudin-3*, *JAMA*, *Zo-1* and *Mucin-2* was

downregulated in the AM group compared with that in the D0 group. In contrast, the

gene expression of *Claudin-2* was higher in the AM group that in the D0 group, which

might explain why moribund hamsters had diarrhoea (34).

Gut microbiota-depleted hamsters exhibited reduced survival time, increased

leptospiral load, and upregulated gene expression of proinflammatory cytokines

compared with untreated hamsters after *Leptospira* infection, while FMT partially

reversed these effects. Our previous study demonstrated that the leptospiral load was

increased in the organs of gut microbiota-depleted mice and that gut microbiota

depletion diminished the bactericidal activity of macrophages during *Leptospira*

infection (10). These results indicated that the homeostasis of the gut microbiota is

essential for both susceptible and resistant hosts to combat *Leptospira* infection. 16S

rRNA sequencing analysis revealed that antibiotic treatment decreased the diversity of

the gut microbiota, and FMT partially reversed these changes. Antibiotic treatment

increased the relative abundance of proinflammatory bacteria, such as

γ-Proteobacteria, while it reduced the relative abundance of anti-inflammatory

bacteria, such as *Allobaculum*. The alteration of the gut microbiota might skew the

host to an inflammatory condition and reduce antileptospiral capacity. FMT aims to

replace an unfavourable resident gut microbiota with a favourable microbiota from a

healthy donor. FMT has demonstrated clear efficacy in the treatment of recurrent

Clostridioides difficile infection (rCDI), inflammatory bowel disease, and

neurological disorders (14). The efficiency of FMT determines the phenotypes of

subjects in the infection model. Prolonging the duration of FMT might help recover

the gut microbiota homeostasis and the capacity to combat anti-*Leptospira* infection.

Our study showed the translocation of Proteobacteria to the intestinal epithelium.

Although no bacteria were found in the blood of the AM group, the level of LPS was

increased in the PBS-treated AM group compared with that of the uninfected group,

and it was decreased in the Abx-treated AM group compared with that of the

PBS-treated AM group. This result indicated that gut-derived LPS was involved in

severe leptospirosis. Gut dysbiosis is a primary determinant of low-grade

endotoxaemia mediated by dysfunction of the intestinal barrier scaffold, which is a

prerequisite for LPS translocation into the systemic circulation (39). Importantly, mice

(resistant to leptospirosis) can tolerate levels of LPS endotoxin 250 higher than the

level that humans can tolerate (52). In addition, classical gram-negative LPS had a

1000-fold higher potency than leptospiral LPS (53). Fecal filtration from the AM

group and the uninfected group both increased the gene expression of *TLR4*, while

only serum from the AM group increased the gene expression of *TLR4*. TLR4

expression was inhibited by PMB treatment. In severe leptospirosis, excessive

immune activation is a hallmark of disease (6). A previous study demonstrated that the

cytokine response to heat-killed whole leptospires was not inhibited by PMB in

macrophages (29). Leptospiral LPS has been described as atypical in terms of

structure, and silver-stained electrophoretic profiles of LPS from pathogenic

leptospiral strains were shown to be different from those of *E. coli* LPS (54).

Although it is still difficult to distinguish the specific roles of gut-derived LPS and

leptospiral LPS in promoting inflammation, we hypothesize that gut-derived LPS may

also be an essential driver of inflammatory reactions in severe leptospirosis. In a rat

cirrhosis model, gut decontamination with a 2-week course of antibiotics led to a

redistributed microbiota, reduced proinflammatory activation of mucosal immune

cells and diminished gut bacterial translocation (37). Another study showed that

endogenous endotoxins caused a pan enteric inflammatory ileus after colonic surgery

and that gut decontamination alleviated small intestinal muscularis inflammation (55).

Our results showed that doxycycline treatment significantly reduced the leptospiral

load in the blood in severe leptospirosis; however, the gene expression of

proinflammatory cytokines was upregulated, which was consistent with the

Jarisch-Herxheimer Reaction (5). Recently, Li *et al*. proposed an all-in-one drug

delivery strategy equipped with bactericidal, LPS neutralization, and detoxification

activities to treat sepsis-associated infections and hyperinflammation (56). Our study

also highlighted the important role of endogenous gut-derived LPS induced by

Leptospira infection. Thus, the neutralization of gut-derived LPS and antileptospiral

therapy are both essential in the treatment of severe leptospirosis; thus, these

strategies provide a novel avenue for the treatment of human acute leptospirosis in the

future.

Our study has potentially important clinical implications for the care of critically ill

patients with severe leptospirosis. In human infections, gastrointestinal symptoms

such as abdominal pain, vomiting, and diarrhea have been frequently observed, which

has been linked to an increased risk of organ dysfunction, and even death (2, 57). Our

findings implicate impairment of the intestinal barrier by *Leptospira* infection as a

mechanism underlying the association between gut dysbiosis and disseminated

gut-derived LPS in the ICU. More importantly, from a translational perspective, our

findings suggest that the increased risk of death in severe leptospirosis can be reduced

by LPS neutralization combined with anti-*Leptospira* therapy. In fact, gut microbiota

dysbiosis in COVID-19 patients or HIV infection is also associated with microbial

translocation, LPS biosynthesis, and an expansion of Proteobacteria (58, 59). The

application of LPS antagonist therapies, whether microbial or pharmaceutical, may be

a viable precision-medicine strategy to increase the survival rate of patients with

severe leptospirosis or other gut-damaging infections by protecting against the spread

of LPS in the bloodstream.

Acknowledgements

We thank Dr. Xiaokui Guo (Shanghai Jiao Tong University, Shanghai, China) for

providing *Leptospira interrogans* serovar Lai strain Lai (56601).

Author contributions

WZ and YC conceived ideas, and XX and WZ designed experiments. XX, JL, ZN and

XC performed the mouse experiments. JL and SZ conducted the bacteriology

experiments. XX and JL analyzed data. WZ and YC performed bioinformatics

analysis. XX and WZ wrote the original draft. YC supervised the study and reviewed

the final version of the text.

Funding

This work was supported by the National Natural Science Foundation of China (No.

32172872 and 32302879), the Fundamental Research Funds for the Central

Universities, China Postdoctoral Science Foundation funded project (BX20220131),

and Basic and Applied Basic Research Foundation of Guangdong Province

(2022A1515110966).

Competing interests

The authors declare no competing interests.

Data availability statement

Data are available in a public, open access repository. The bacterial 16S rDNA

sequences were deposited to the sequence read archive. The accession number for the

16S sequencing data reported in this paper are PRJNA772361 and PRJNA1036417.

Reference

1. Coburn J, Picardeau M, Woods CW, Veldman T, Haake DA. Pathogenesis
insights from an ancient and ubiquitous spirochete. PLoS Pathog.
2021;17(10):e1009836.

2. Haake DA, Levett PN. Leptospirosis in humans. Curr Top Microbiol Immunol.
2015;387:65-97.

3. Maia MAC, Bettin EB, Barbosa LN, de Oliveira NR, Bunde TT, Pedra ACK, et al.
Challenges for the development of a universal vaccine against leptospirosis revealed
by the evaluation of 22 vaccine candidates. Front Cell Infect Microbiol.
2022;12:940966.

4. Coutinho ML, Matsunaga J, Wang LC, de la Pena Moctezuma A, Lewis MS,
Babbitt JT, et al. Kinetics of *Leptospira interrogans* infection in hamsters after
intradermal and subcutaneous challenge. PLoS Negl Trop Dis. 2014;8(11):e3307.

5. Guerrier G, D'Ortenzio E. The Jarisch-Herxheimer reaction in leptospirosis: a
systematic review. PLoS One. 2013;8(3):e59266.

6. Cagliero J, Villanueva S, Matsui M. Leptospirosis Pathophysiology: Into the
Storm of Cytokines. Front Cell Infect Microbiol. 2018;8:204.

7. Alventosa Mateu C, Plana Campos L, Larrey Ruíz L, Acedo Mayordomo R,
Sanchís Artero L, Peño Muñoz L, et al. [Gastrointestinal bleeding and acute hepatic
failure by leptospirosis: an entity that should not be forgotten]. Rev Gastroenterol
Peru. 2017;37(1):96-9.

8. Kamada N, Chen GY, Inohara N, Nunez G. Control of pathogens and pathobionts
by the gut microbiota. Nat Immunol. 2013;14(7):685-90.

9. McDonald B, Zucoloto AZ, Yu IL, Burkhard R, Brown K, Geuking MB, et al.
Programing of an Intravascular Immune Firewall by the Gut Microbiota Protects
against Pathogen Dissemination during Infection. Cell Host Microbe.
2020;28(5):660-8 e4.

10. Xie X, Liu J, Chen X, Zhang S, Tang R, Wu X, et al. Gut microbiota involved in
leptospiral infections. ISME J. 2022;16(3):764-73.

11. Gomes CK, Guedes M, Potula HH, Dellagostin OA, Gomes-Solecki M. Sex
Matters: Male Hamsters Are More Susceptible to Lethal Infection with Lower Doses
of Pathogenic *Leptospira* than Female Hamsters. Infect Immun. 2018;86(10).

12. Vemuri R, Sylvia KE, Klein SL, Forster SC, Plebanski M, Eri R, et al. The
microgenderome revealed: sex differences in bidirectional interactions between the
microbiota, hormones, immunity and disease susceptibility. Semin Immunopathol.
2019;41(2):265-75.

13. Tawk C, Lim B, Bencivenga-Barry NA, Lees HJ, Ramos RJF, Cross J, et al.
Infection leaves a genetic and functional mark on the gut population of a commensal
bacterium. Cell Host Microbe. 2023;31(5):811-26 e6.
14. Schmidt TSB, Li SS, Maistrenko OM, Akanni W, Coelho LP, Dolai S, et al.
Drivers and determinants of strain dynamics following fecal microbiota
transplantation. Nat Med. 2022;28(9):1902-12.
15. Sun L, Pang Y, Wang X, Wu Q, Liu H, Liu B, et al. Ablation of gut microbiota
alleviates obesity-induced hepatic steatosis and glucose intolerance by modulating
bile acid metabolism in hamsters. Acta Pharm Sin B. 2019;9(4):702-10.
16. Zhai L, Huang C, Ning Z, Zhang Y, Zhuang M, Yang W, et al. Ruminococcus
gnavus plays a pathogenic role in diarrhea-predominant irritable bowel syndrome by
increasing serotonin biosynthesis. Cell Host Microbe. 2023;31(1):33-44 e5.
17. Soares LM, Macedo JO, de Azevedo EC, Santos CS, Sampaio Mde Q, dos Santos
AC, et al. Immunomodulatory treatment with thalidomide in experimental
leptospirosis in Golden Syrian hamsters (Mesocricetus auratus). Trans R Soc Trop
Med Hyg. 2014;108(2):105-11.
18. Jin X, Zhang W, Ding Z, Wang H, Wu D, Xie X, et al. Efficacy of the Rabbit
Polyclonal Anti-leptospira Antibody against Homotype or Heterotype *Leptospira*
Infection in Hamster. PLoS Negl Trop Dis. 2016;10(12):e0005191.
19. Spalinger MR, Schwarzfischer M, Hering L, Shawki A, Sayoc A, Santos A, et al.
Loss of *PTPN22* abrogates the beneficial effect of cohousing-mediated fecal
microbiota transfer in murine colitis. Mucosal Immunol. 2019;12(6):1336-47.
20. Murray CK, Hospenthal DR. Broth microdilution susceptibility testing for
Leptospira spp. Antimicrob Agents Chemother. 2004;48(5):1548-52.
21. Xu J, Wang J, Chen M, Chao B, He J, Bai Y, et al. miR-101-5p suppresses
trophoblast cell migration and invasion via modulating the DUSP6-ERK1/2 axis in
preeclampsia. J Assist Reprod Genet. 2023;40(7):1597-610.
22. Wu J, Xiao J, Bai M, Shi C, Xin Y, Zhao W, et al. Single-Cell RNA Sequencing
Reveals Unique Alterations in the Immune Panorama and Treg Subpopulations in
Mice during the Late Stages of *Echinococcus granulosus* Infection. Infect Immun.
2023;91(5):e0002923.
23. Cao Y, Xie X, Zhang W, Wu D, Tu C. Low-dose Norfloxacin-treated leptospires
induce less IL-1beta release in J774A.1cells following discrepant leptospiral gene
expression. Microb Pathog. 2018;119:125-30.
24. Zhang W, Xie X, Wang J, Song N, Lv T, Wu D, et al. Increased inflammation
with crude *E. coli* LPS protects against acute leptospirosis in hamsters. Emerg
Microbes Infect. 2020;9(1):140-7.
25. Zhang W, Zhang N, Xie X, Guo J, Jin X, Xue F, et al. Toll-Like Receptor 2
Agonist Pam3CSK4 Alleviates the Pathology of Leptospirosis in Hamster. Infect
Immun. 2016;84(12):3350-7.
26. Sanchez KK, Chen GY, Schieber AMP, Redford SE, Shokhirev MN, Leblanc M,
et al. Cooperative Metabolic Adaptations in the Host Can Favor Asymptomatic

Infection and Select for Attenuated Virulence in an Enteric Pathogen. Cell.
2018;175(1):146-+.

27. Inamasu Y, Nikaido Y, Miyahara S, Maruoka T, Takigawa T, Ogawa M, et al.
Dissemination of *Leptospira* into the intestinal tract resulting in fecal excretion in a
hamster model of subcutaneous infection with *Leptospira interrogans*. Microb Pathog.
2022;165:105481.

28. Kitamoto S, Nagao-Kitamoto H, Jiao Y, Gillilland MG, 3rd, Hayashi A, Imai J, et
al. The Intermucosal Connection between the Mouth and Gut in Commensal
Pathobiont-Driven Colitis. Cell. 2020;182(2):447-62 e14.

29. Viriyakosol S, Matthias MA, Swancutt MA, Kirkland TN, Vinetz JM. Toll-like
receptor 4 protects against lethal *Leptospira interrogans* serovar icterohaemorrhagiae
infection and contributes to in vivo control of leptospiral burden. Infect Immun.
2006;74(2):887-95.

30. Aldape MJ, Bryant AE, Katahira EJ, Hajjar AM, Finegold SM, Ma Y, et al. Innate
immune recognition of, and response to, *Clostridium sordellii*. Anaerobe.
2010;16(2):125-30.

31. Limothai U, Lumlertgul N, Sirivongrangson P, Kulvichit W, Tachaboon S,
Dinhuzen J, et al. The role of leptospiremia and specific immune response in severe
leptospirosis. Sci Rep. 2021;11(1):14630.

32. Turner JR. Intestinal mucosal barrier function in health and disease. Nat Rev
Immunol. 2009;9(11):799-809.

33. Horowitz A, Chanez-Paredes SD, Haest X, Turner JR. Paracellular permeability
and tight junction regulation in gut health and disease. Nat Rev Gastroenterol Hepatol.
2023;20(7):417-32.

34. Tsai PY, Zhang B, He WQ, Zha JM, Odenwald MA, Singh G, et al. IL-22
Upregulates Epithelial Claudin-2 to Drive Diarrhea and Enteric Pathogen Clearance.
Cell Host Microbe. 2017;21(6):671-81 e4.

35. Zeng Y, Wu R, Wang F, Li S, Li L, Li Y, et al. Liberation of daidzein by gut
microbial beta-galactosidase suppresses acetaminophen-induced hepatotoxicity in
mice. Cell Host Microbe. 2023;31(5):766-80 e7.

36. Yang Y, Nguyen M, Khetrapal V, Sonnert ND, Martin AL, Chen H, et al.
Within-host evolution of a gut pathobiont facilitates liver translocation. Nature.
2022;607(7919):563-70.

37. Munoz L, Borrero MJ, Ubeda M, Conde E, Del Campo R, Rodriguez-Serrano M,
et al. Intestinal Immune Dysregulation Driven by Dysbiosis Promotes Barrier
Disruption and Bacterial Translocation in Rats With Cirrhosis. Hepatology.
2019;70(3):925-38.

38. Bertocchi A, Carloni S, Ravenda PS, Bertalot G, Spadoni I, Lo Cascio A, et al.
Gut vascular barrier impairment leads to intestinal bacteria dissemination and
colorectal cancer metastasis to liver. Cancer Cell. 2021.

39. Violi F, Cammisotto V, Bartimoccia S, Pignatelli P, Carnevale R, Nocella C.
Gut-derived low-grade endotoxaemia, atherothrombosis and cardiovascular disease.
Nat Rev Cardiol. 2022.
40. Szabo G. Gut-liver axis in alcoholic liver disease. Gastroenterology.
2015;148(1):30-6.
41. Caesar R, Reigstad CS, Backhed HK, Reinhardt C, Ketonen M, Lunden GO, et al.
Gut-derived lipopolysaccharide augments adipose macrophage accumulation but is
not essential for impaired glucose or insulin tolerance in mice. Gut.
2012;61(12):1701-7.
42. Lappi J, Kolehmainen M, Mykkanen H, Poutanen K. Do large intestinal events
explain the protective effects of whole grain foods against type 2 diabetes? Crit Rev
Food Sci Nutr. 2013;53(6):631-40.
43. Maruoka T, Nikaido Y, Miyahara S, Katafuchi E, Inamasu Y, Ogawa M, et al.
Correlation between renal distribution of leptospires during the acute phase and
chronic renal dysfunction in a hamster model of infection with *Leptospira interrogans*.
PLoS Negl Trop Dis. 2021;15(6):e0009410.
44. Zilber AL, Belli P, Grezel D, Artois M, Kodjo A, Djelouadji Z. Comparison of
Mucosal, Subcutaneous and Intraperitoneal Routes of Rat *Leptospira* Infection. PLoS
Negl Trop Dis. 2016;10(3):e0004569.
45. Medkour H, Amona I, Akiana J, Laidoudi Y, Davoust B, Bitam I, et al. Bacterial
Infections in Humans and Nonhuman Primates from Africa: Expanding the
Knowledge. Yale J Biol Med. 2021;94(2):227-48.
46. Lankelma JM, Birnie E, Weehuizen TAF, Scicluna BP, Belzer C, Houtkooper RH,
et al. The gut microbiota as a modulator of innate immunity during melioidosis. PLoS
Negl Trop Dis. 2017;11(4):e0005548.
47. Lozupone CA, Li M, Campbell TB, Flores SC, Linderman D, Gebert MJ, et al.
Alterations in the gut microbiota associated with HIV-1 infection. Cell Host Microbe.
2013;14(3):329-39.
48. Becattini S, Sorbara MT, Kim SG, Littmann EL, Dong Q, Walsh G, et al. Rapid
transcriptional and metabolic adaptation of intestinal microbes to host immune
activation. Cell Host Microbe. 2021;29(3):378-93 e5.
49. Byndloss MX, Olsan EE, Rivera-Chavez F, Tiffany CR, Cevallos SA, Lokken KL,
et al. Microbiota-activated PPAR-gamma signaling inhibits dysbiotic
Enterobacteriaceae expansion. Science. 2017;357(6351):570-5.
50. Rusu AD, Georgiou M. The multifarious regulation of the apical junctional
complex. Open Biol. 2020;10(2):190278.
51. Sebastian I, Okura N, Humbel BM, Xu J, Hermawan I, Matsuura C, et al.
Disassembly of the apical junctional complex during the transmigration of Leptospira
interrogans across polarized renal proximal tubule epithelial cells. Cell Microbiol.
2021;23(9):e13343.

52. Copeland S, Warren HS, Lowry SF, Calvano SE, Remick D, Inflammation, et al.
Acute inflammatory response to endotoxin in mice and humans. Clin Diagn Lab
Immunol. 2005;12(1):60-7.
53. Werts C, Tapping RI, Mathison JC, Chuang TH, Kravchenko V, Saint Girons I, et
al. Leptospiral lipopolysaccharide activates cells through a TLR2-dependent
mechanism. Nat Immunol. 2001;2(4):346-52.
54. Bonhomme D, Santecchia I, Vernel-Pauillac F, Caroff M, Germon P, Murray G, et
al. Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated
antimicrobial responses through O antigen and associated lipoproteins. PLoS Pathog.
2020;16(8):e1008639.
55. Turler A, Schnurr C, Nakao A, Togel S, Moore BA, Murase N, et al. Endogenous
endotoxin participates in causing a panenteric inflammatory ileus after colonic surgery.
Ann Surg. 2007;245(5):734-44.
56. Li X, Qu S, Song X, Wu C, Shen J, Zhu K. In Situ Neutralization and
Detoxification of LPS to Attenuate Hyperinflammation. Adv Sci (Weinh).
2023;10(26):e2302950.
57. Jimenez JIS, Marroquin JLH, Richards GA, Amin P. Leptospirosis: Report from
the task force on tropical diseases by the World Federation of Societies of Intensive
and Critical Care Medicine. J Crit Care. 2018;43:361-5.
58. Bernard-Raichon L, Venzon M, Klein J, Axelrad JE, Zhang C, Sullivan AP, et al.
Gut microbiome dysbiosis in antibiotic-treated COVID-19 patients is associated with
microbial translocation and bacteremia. Nat Commun. 2022;13(1):5926.
59. Zhang Y, Xie Z, Zhou J, Li Y, Ning C, Su Q, et al. The altered metabolites
contributed by dysbiosis of gut microbiota are associated with microbial translocation
and immune activation during HIV infection. Front Immunol. 2022;13:1020822.

Figure legends

Fig. 1. *L. interrogans* proliferated in the intestine, destroyed the intestinal

structure, and increased intestinal inflammation. (A) Flow diagram of the

experiment. Six-week-old female hamsters were injected intraperitoneally with 10^7

leptospires. Hamsters were euthanized at 0 h, 6 h, 48 h, 96 h, and AM p.i.. The ileums

and colons were collected aseptically for further analysis. (B) Leptospiral burdens in

the ileums and colons of *Leptospira*-infected hamsters were determined by qPCR (n =

6). (C) Left panel, histopathology of colons of D0 hamsters (scale bar, 50 μm; n = 4)

and AM hamsters (scale bar, 50 μm; n = 4). Representative photographs are presented.

Right panel, histopathology scores of colons. D0, uninfected hamster. AM, articulo

mortis. (D-I) The gene expression of *Lcn2* (D), *TNF-α* (E), *IL-1β* (F), *IL-10* (G), *TLR4*

(H), and *Nos2* (I) was analyzed by RT-qPCR. The mRNA levels of genes in the colons

of the D0 group (n = 3) and the AM group (n = 3) were normalized to the expression

of the housekeeping gene *GAPDH*. Each infection experiment was repeated three

times. Data are shown as the mean ± SEM and analyzed by the Wilcoxon rank-sum

test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Fig. 2. *L. interrogans* infection changed the composition of the gut microbiota. (A)

Flow diagram of the experiment. Six-week-old female hamsters were injected

intraperitoneally with 10^7 leptospires. Fecal pellets were collected aseptically for 16S

rRNA gene sequencing at 0 days, 2 days, and AM p.i.. D0, uninfected hamster. AM,

articulo mortis. (B-C) Observed species (B) and Ace (C) in the feces of hamsters.

Observed species and Ace indicate species richness. (D-E) Simpson index (D) and

Shannon index (E) in the feces of hamsters. Simpson and Shannon index indicate

species diversity. (F) Principal coordinates analysis (PCoA) of fecal samples based on

16S rRNA gene sequencing using weighted UniFrac. (G) PCoA of fecal samples

based on 16S rRNA gene sequencing using weighted UniFrac. (H) Relative

abundance of the top 10 phyla in the feces of hamster. (I) Linear discriminant analysis

(LDA) of effect size (LEfSe) between D0 and AM hamsters (LDA score > 3).

Cambridge blue bars indicate taxa enrichment in D0 hamsters, and pink bars indicate

taxa enrichment in AM hamsters. (J) The ratio of Firmicutes to Bacteroidetes in the

D0 and AM. (K-M) The relative abundance of Proteobacteria (K), *Lactobacillus* (L),

and *Allobaculum* (M) in the D0 and AM groups. (D0, n = 10; D2, n = 10; AM, n = 8).

Data are shown as the mean ± SEM and analyzed by the Wilcoxon rank-sum test. $*p$

< 0.05, $**p < 0.01$.

Fig. 3. *L. interrogans* infection disrupted intestinal tight junctions. Six-week-old

female hamsters were injected intraperitoneally with 10^7 leptospires. Blood was

collected to determine intestinal permeability at 0 days, 2 days, and AM p.i.. Colons

were collected aseptically to measure gene expression at 0, 2, and AM p.i.. (A) The

intestinal permeability of *Leptospira*-infected hamsters was analyzed with a

fluorescence spectrophotometer at indicated time (D0, n = 4; D2, n = 6; AM, n = 6).

AM, articulo mortis. (B-G) The relative gene expression of *ZO-1* (B), *Claudin-3* (C),

JAMA (D), *Claudin-2* (E), and *Mucin-2* (F) in *Leptospira*-infected hamsters (n = 6-10

per group) was determined by qPCR. D0, uninfected hamster. AM, articulo mortis.

Each infection experiment was repeated three times. Data are shown as the mean ±

SEM and analyzed by the Wilcoxon rank-sum test. *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001.

Fig. 4. Translocated microbiota in the intestinal epithelium following *L.*

***interrogans* infection.** Six-week-old female hamsters were injected intraperitoneally

with 10^7 leptospires. Hamsters were euthanized at 0 days, 2 days, and AM p.i.. AM,

articulo mortis. Colons were collected aseptically. The intestinal contents were

excluded from the intestine with PBS and a segment of colon was fixed in 4%

paraformaldehyde solution overnight and analyzed by FISH with an EUB 338 probe

(red). (A) Th results are representative photographs of three groups. (B) Number of

positive bacteria per field in the three groups. n = 6 per group. D0, uninfected hamster.

AM, articulo mortis. Each infection experiment was repeated three times. Data are

shown as the mean ± SEM and analyzed by the Wilcoxon rank-sum test. ***$p <$

0.001.

Fig. 5. Microbiota depletion exacerbated leptospirosis. (A) Flow diagram of the

experiment. Hamsters were administered with Abx or water intragastrically once daily

for ten consecutive days. Then, fecal pellets were collected aseptically for 16S rRNA

gene sequencing. (B-C) Observed species (B) and Chao1 (C) in the feces of hamsters.

Observed species and Chao1 indicate species richness. (D-E) Simpson index (D) and

Shannon index (E) in the feces of hamsters. Simpson and Shannon index indicate

species diversity. (F) Relative abundance of the top 10 phyla in the feces of hamsters.

(G) The relative abundance of Bacteroidetes, Firmicutes, and Proteobacteria in the

Con, Abx, and FMT groups. n = 4 per group. (H) Hamsters were treated with Abx as

described above. After a 2-day washout period, hamsters were intraperitoneally

infected with 10^6 *L. interrogans*. Then, the survival rate of the hamsters was recorded.

n = 6 per group. (J) Leptospiral load in the blood of the Con, Abx, and FMT groups. n

= 6 per group. (K-M) Gene expression of *TNF-α* (K), *IL-1β* (L), and *IL-10* (M) in the

blood of the three groups. n = 5 per group. Each infection experiment was repeated

three times. Data are shown as the mean ± SEM and analyzed by the Wilcoxon

rank-sum test. Survival differences between the study groups were compared by using

the Kaplan-Meier log-rank test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. ns, not

significant.

Fig. 6. *Leptopsira* infection induced intestinal and systemic inflammation that

was partially inhibited by LPS neutralization. (A) Flow diagram of the experiment.

Fecal filtration or serum of uninfected hamsters and AM hamsters was directly added

into dishes or pretreated with PMB at 37 °C for 2 h, and then cocultured with

macrophages for 4 h. RNA was extracted for gene expression analysis. (B-E) The

gene expression of *TNF-α* (B), *IL-1β* (C), *IL-10* (D), and *TLR4* (E) of fecal filtration

of uninfected hamsters (n = 3) and AM hamsters (n = 3) were analyzed by RT-qPCR.

(F-I) The gene expression of *TNF-α* (F), *IL-1β* (G), *IL-10* (H) and *TLR4* (I) of serum

of uninfected hamsters (n = 3) and AM hamsters (n = 3) were analyzed by RT-qPCR.

(J-M) The effect of LPS neutralization on the gene expression of *TNF-α* (J), *IL-1β* (K),

IL-10 (L) and *TLR4* (M) in the fecal filtration of AM hamsters (n = 3) were analyzed

by RT-qPCR. (N-Q) The effect of LPS neutralization on the gene expression of *TNF-α*

(N), *IL-1β* (O), *IL-10* (P) and *TLR4* (Q) of in the serum of AM hamsters (n = 3) was

analyzed by RT-qPCR. The mRNA levels in untreated controls were set as 1.0. Each

infection experiment was repeated three times. Data are shown as the mean ± SEM

and analyzed by the Wilcoxon rank-sum test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Fig. 7. LPS neutralization improved the survival rate of female hamsters with

severe leptospirosis. (A) Flow diagram of the experiment. Six-week-old female

hamsters were injected intraperitoneally with 10^7 leptospires. Group 1: saline control;

Group 2: Polymyxin B (PMB) (1 mg/kg, i.p.); Group 3: Antibody against *Leptospira*

(Abl) (16 mg/kg, subcutaneous injection); Group 4: Doxycycline (Dox) (5 mg/kg,

i.p.); Group 5: TLR4 antibody (25 μg/hamster, i.p.); Group 6: PMB (1 mg/kg, i.p.)

and Abl (16 mg/kg, subcutaneous injection); Group 7: PMB (1 mg/kg, i.p.) and Dox

(5 mg/kg, i.p.); Group 8: TLR4 antibody (25 μg/hamster, i.p.) and Dox (5 mg/kg, i.p.).;

Hamsters were treated twice a day for 3 consecutive days. Hamsters were monitored

daily for 21 days. (B) Survival rate of female hamsters of severe leptospirosis after

different treatments. n = 6 per group. (C) Leptospiral load in the blood of different

groups. n = 5 per group. (D-F) Gene expression of *TNF-α* (D, *IL-1β* (E), and *IL-10* (F)

in the blood of different groups. The mRNA levels of cytokines in uninfected controls

were set as 1.0. n = 5 per group. Each infection experiment was repeated three times.

Survival differences between the study groups were compared by using the

Kaplan-Meier log-rank test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. ns, not significant.

A

	EUB 338	DAPI	Merge	Enlarge

D0

D2

AM

100 μm (repeated in each panel)

B

Number of positive bacteria/field

-50   0   50   100   150

D0

D2

AM

***