Modeling and Correction of Protein Conformational Disease in iPSC-derived Neurons through Personalized Base Editing

Colin T Konishi[1,2], Nancy Mulaiese[,2], Tanvi Butola[3], Qinkun Zhang[1,2], Dana Kagan[1,2], Qiaoyan Yang[1,2], Mariel Pressler[1,4], Brooke G Dirvin[1,2], Orrin Devinsky[4,†], Jayeeta Basu[3†], Chengzu Long[1,2,3,4,†]

[1] NYU Cardiovascular Research Center, NYU Grossman School of Medicine, New York, NY 100016,

[2] Leon H. Charney Division of Cardiology, NYU Grossman School of Medicine, New York, NY 100016,

[3] Department of Neuroscience and Physiology, NYU Grossman School of Medicine, New York, NY 100016,

[4] Department of Neurology, NYU Grossman School of Medicine, New York, NY 100016.

† Corresponding authors:

E-mail: chengzu.long@nyumc.org (C.L.), jayeeta.basu@nyulangone.org (J.B.) and od4@nyu.edu (O.D.)

Short title: Base editing correction in conformational disease

Abstract

Altered protein conformation can cause incurable neurodegenerative disorders. Mutations in

SERPINI1, the gene encoding neuroserpin, can alter protein conformation resulting in cytotoxic

aggregation leading to neuronal death. Familial encephalopathy with neuroserpin inclusion

bodies (FENIB) is a rare autosomal dominant progressive myoclonic epilepsy that progresses to

dementia and premature death. We developed HEK293T and induced pluripotent stem cell

(iPSC) models of FENIB, harboring a patient-specific pathogenic *SERPINI1* variant or stably

overexpressing mutant neuroserpin fused to GFP (MUT NS-GFP). Here, we utilized a

personalized adenine base editor (ABE)-mediated approach to correct the pathogenic variant

efficiently and precisely to restore neuronal dendritic morphology. ABE-treated MUT NS-GFP

cells demonstrated reduced inclusion size and number. Using an inducible MUT NS-GFP

neuron system, we identified early prevention of toxic protein expression allowed aggregate

clearance, while late prevention halted further aggregation. To address several challenges for

clinical applications of gene correction, we developed a neuron-specific engineered virus-like

particle to optimize neuronal ABE delivery, resulting in higher correction efficiency. Our findings

provide a targeted strategy which may treat FENIB and potentially other neurodegenerative

diseases due to altered protein conformation such as Alzheimer's and Huntington's diseases.

Introduction

Altered protein conformation can cause progressive and incurable neurodegenerative disorders such as Alzheimer's and Huntington's diseases.[1,2] Neuroserpin (NS) is a neuron-specific serine protease inhibitor involved in axon growth and synapse formation.[3,4] Some mutations in the *SERPINI1* gene alter neuroserpin protein stability, leading to protein aggregation within the endoplasmic reticulum (ER), causing ER and oxidative stress, neuronal dysfunction, and cell death.[5–7] Familial encephalopathy with neuroserpin inclusion bodies (FENIB) is a rare autosomal dominant progressive myoclonic epilepsy, causing seizures, dementia, motor impairments, and premature death.[8–10] The variant, *SERPINI1* c.1175 G>A (p.G392E) causes one of the most severe forms of FENIB, characterized by absence seizures, myoclonic and tonic-clonic seizures, and progressive dementia beginning in late childhood to early adolescence.[11,12] There are no disease-modifying therapies.[13,14] Many clinical and preclinical strategies for the treatment of conformational diseases, including Alzheimer's and Huntington's diseases, aim to reduce or eliminate expression of precursor forms of aggregation-prone proteins.[15,16] These strategies employ RNAi, antisense oligonucleotides, small molecular modulators, or genome editors such as zinc-finger nucleases, transcription activator-like effector nucleases (TALEN), and clustered regularly interspace palindromic repeat (CRISPR)/CRISPR-associated sequence (Cas) nucleases to target pathogenic forms of RNA and DNA.[17–20] All approaches face their own challenges.

CRISPR/Cas genome editors are sequence-specific RNA-guided nucleases that generate double-strand breaks (DSB) in target DNA.[21–23] DSB repair through non-homologous end joining facilitates gene disruption due to the formation of insertions and deletions, while homology directed repair (HDR) produces sequence-specific modifications by introducing an exogenous repair template.[24,25] Cas9 can edit stable cell lines and induced pluripotent stem cells (iPSCs) *in vitro* and neurons, cardiomyocytes, hepatocytes, and other cell types *in vivo*.[26–28] Advances in

iPSC technology and CRISPR/Cas genome editing have enabled the generation of isogenic

cells harboring patient-specific disease-causing variants. These cells can be differentiated into

various cell fates to model disease.[29–31] However, HDR has been demonstrated to be primarily

active during S and G2 phases of the cell cycle which limits its therapeutic applications in post-

mitotic cells.[32–34] Adenine base editing (ABE), a CRISPR/Cas9 modality, uses an evolved

transfer RNA deaminase (tadA) fused to Cas9 nickase (nCas9) to act on single-stranded DNA,

converting adenine (A) to guanine (G) without a DSB. ABEs can generate precise A>G

transition mutations and have the potential to correct the pathogenic *SERPINI1* c.1175 G>A

mutation.[35]

Therapeutic delivery of gene editors remains a common challenge for all gene therapies.[36]

Adenovirus, adeno-associated virus (AAV), and lentivirus are highly efficient viral vectors but

have limited scope due to immunogenicity and genomic packaging capacity inherent to using

viruses.[37,38] Engineered virus-like particles (eVLPs) are an extracellular vesicle formed of viral

proteins that can deliver cargo fused to retroviral Gag proteins but do not contain viral genomes.

eVLPs target tissues with far fewer immunogenic effects and are expressed for a shorter

duration than AAVs and lentivirus, thereby reducing off-target edits.[38–40] Further, eVLPs can

target cell-types by modulating surface glycoproteins.[39] However, the efficiency of eVLP-

delivered ABEs to correct mutations in cortical neurons remains unknown, as neurons use

nonclassical nuclear import, with lower expression of KPNA2 and KPNA7, which are recognized

by proteins with classical nuclear localization signals.

In this proof-of-concept study, we generated a stable overexpression model of FENIB in human

embryonic kidney 293T (HEK293T) cells, and for the first time, endogenous knock-in and

overexpression models of FENIB in iPSC-derived neurons. We used ABEs to target and correct

the pathogenic variant. Correction of the variant restored dendritic morphology in KI neurons,

and clearance of MUT NS-GFP aggregates in HEK293T cells. Using our inducible system, early

treatment of MUT NS-GFP cells led to aggregate clearance, while later treatment prevented

aggregate progression. We also tested eVLPs with neuron-specific nuclear localization signals

(NLS) and identified improved neuronal editing. eVLP-mediated ABE editing appears to be a

safe and efficient treatment for FENIB and potentially other diseases of protein conformation.

Results

CRISPR/Cas9 mediated HDR KI of patient-specific *SERPINI1* variant.

The autosomal dominant *SERPINI1* c.1175 G>A (p. G392E) variant causes neuroserpin to form

cytotoxic aggregates (**Fig. 1A, B**).[9] To investigate ABE correction as a means to treat this

variant, we first generated a *SERPINI1* c.1175 G>A cell line using CRISPR/Cas9 mediated HDR

to knock-in the variant in HEK293T cells. We tested SpCas9 and SpCas9-NG editors to target

SERPINI1 exon 9 (**Fig. 1B**).[41] Single guide RNAs (gRNA) were cloned into plasmids encoding

SpCas9-2A-GFP (g1) or SpCas9-NG-2A-GFP (g2). g1 and g2 cleaved *SERPINI1* proximal to

c.1175, and a single stranded oligodeoxynucleotide donor (ssODN) was used for templated

HDR-mediated knock-in. g1 used a 5'-GGG-3' PAM located at *SERPINI1* c.1173-c.1175. KI of

SERPINI1 c.1175 G>A alters the PAM sequence to 5'-GG<u>A</u>-3', preventing recutting by SpCas9,

to increase KI efficiency by preventing recurrent cleavage**.** g2 used a 5'-TG-3' PAM to generate

a DSB immediately downstream of the KI sequence to optimize HDR-mediated KI efficiency.[42]

First, HEK293T cells were transfected using lipofectamine 2000 with g1 or g2 plasmids without

ssODN (**Fig. S1A, B**). 72-hours post-transfection, cells were sorted by GFP expression, and

editing efficiency was measured by PCR-based T7 endonuclease I (T7E1) assay, Sanger

sequencing, and Synthego Inference of CRISPR Edits (Synthego ICE) analysis tool.[43,44] Both g1

and g2 demonstrated high editing activity on target sites. Next, we nucleofected HEK293T cells

with g1 or g2 plasmids and *SERPINI1* c.1175 G>A ssODN for templated KI using the Lonza

nucleofection SF system (**Fig. S1C**). 72-hours post-nucleofection, we sorted GFP-positive cells

into bulk populations or single cells to assess editing efficiencies and selected clones harboring

the patient-specific *SERPINI1* c.1175 G>A variant. ICE identified modest HDR efficiencies by g1

(16.5%) and g2 (11.5%). g2 produced far fewer nonspecific indels (50%) compared to g1 (80%).

The similar KI efficiencies but improved nonspecific indel profile of g2 prompted us to focus on

g2 for HEK293T modeling.

Next, we nucleofected iPSCs with either g1 or g2 plasmid and *SERPINI1* c.1175 G>A ssODN

for templated KI. Both gRNAs efficiently generated cells harboring the variant sequence (g1:

24%, g2: 21%), consistent with our HEK293T cell editing (**Fig. S1D**). All identified FENIB

patients are heterozygous for *SERPINI1* variants. [45] However, we sought to assess the impact

of gene dosage and selected both heterozygous *SERPINI1* c.1175 G>A/+ (heterozygous KI)

and homozygous *SERPINI1* c.1175 G>A/G>A (homozygous KI) clones. Patient-specific iPSCs

and isogenic control wildtype clones were clonally expanded for future experiments.

Stable overexpression of GFP fused neuroserpin in human cells.

Previous studies have used immortalized cell lines, and mouse primary neural progenitors and

neurons overexpressing neuroserpin protein to model FENIB.[5,14,46,47] To more accurately

recapitulate patient disease, we implemented overexpression of mutant neuroserpin in human

HEK293T cells and iPSCs in addition to our KI cells. We designed a lentivirus vector expressing

wild-type (WT) or mutant (MUT) neuroserpin protein fused to enhanced green fluorescent

protein (NS-GFP) under constitutive cytomegalovirus (CMV) or doxycycline-inducible

tetracycline response element (TRE) promoters (**Fig. 1C**). We transduced HEK293T cells with

TRE-NS-GFP and reverse tetracycline-controlled transactivator (rtTA) lentiviruses. One day

post-transduction, we induced NS-GFP expression by adding doxycycline to the culturing

media. Using a combination of imaging native GFP fluorescence and immunocytochemical

labeling, we assessed NS-GFP expression over the span of one week. One day after NS-GFP

induction (d1), cells expressed NS-GFP with no observable differences between WT and MUT

NS-GFP expressing cells. Both types of cells had diffuse NS-GFP fluorescence which

colocalized with immunofluorescent neuroserpin staining. By day 7 (d7), MUT NS-GFP cells

harbored aggregates in the form of concentrated puncta of GFP expression (**Fig. 1C**).

Fluorescent GFP puncta colocalized with the anti-neuroserpin immunofluorescent staining. WT

NS-GFP fluorescence remained cytoplasmic and diffuse throughout the duration of our

experiments, with no detectable puncta at any timepoint.

Protein aggregation in MUT NS-GFP cells was confirmed using denaturing sodium dodecyl-

sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and nondenaturing native-PAGE

immunoblotting (**Fig. 1D and Fig. S1E**). Immunoblot of SDS-PAGE using anti-neuroserpin

antibody revealed a band of ~85kDa in the WT and MUT NS-GFP lanes; the combined

molecular weights of neuroserpin (55kDa) and GFP (30kDa) fusion protein (**Fig. 1D**). Native-

PAGE immunoblots of WT and MUT NS-GFP confirmed observations made during fluorescent

microscopy imaging. Native-PAGE immunoblots of WT NS-GFP revealed a single dominant

band indicative of NS-GFP monomers. MUT NS-GFP samples had a NS-GFP monomer band in

addition to banding and smearing at higher molecular weights, reflecting mutant NS-GFP

aggregate polymers. Next, we validated our results by resolving SDS-PAGE and native-PAGE

using anti-GFP antibodies (**Fig. S1E**). Immunoblot of SDS-PAGE using anti-GFP antibodies

revealed a ~85kDa band in both WT and MUT NS-GFP lanes. Validating results obtained with

the anti-neuroserpin antibody, native-PAGE immunoblot of WT NS-GFP and MUT NS-GFP

revealed a shared NS-GFP monomer band while MUT NS-GFP had additional high molecular

weight banding and smearing.

Neuroserpin aggregation in the endoplasmic reticulum (ER) causes ER stress mediated through

the unfolded protein response (UPR) and ER overload response (EROR).[7,48,49] We collected

RNA from WT and MUT NS-GFP HEK293T cells, cells stably expressing GFP, and control cells

without any form of overexpression. We assessed ER stress through quantitative PCR (qPCR)

of UPR response genes *BiP* and spliced *XBP1* (**Fig. S1F**). When normalized to control

HEK293T cells, cells overexpressing GFP show no difference in fold expression of either gene.

We find that both WT and MUT NS-GFP cells revealed a significant increase in both *BiP* (WT

NS-GFP: 4.1, MUT NS-GFP: 7.8, p<0.0001) and *sXBP1* (WT NS-GFP: 2.8, MUT NS-GFP: 3.9,

p<0.0001), with the greatest increase in MUT NS-GFP cells when normalized to *18s* transcript

expression. Previous studies of cell metabolic activity have found that overexpression of WT NS

reduces cellular viability[14]. In all, our immunofluorescent microscopy, native-PAGE immunoblot,

and transcript analyses demonstrated stable overexpression of NS-GFP results in FENIB

neuroserpin aggregation in human cells.

Generation of patient-specific iPSC-derived FENIB neurons through *Neurogenin2*

induction.

To evaluate our iPSC *in vitro* FENIB phenotype model, we differentiated iPSCs into excitatory

cortical neurons by overexpressing *Neurogenin2* (Ngn2) (**Fig. 1E and Fig. S2A**).[31] We

assessed induction efficiency through immunocytochemical labeling of iPSCs prior to induction

and cells 21 days (d21) post-induction. iPSCs showed standard iPSC morphology and

pluripotency marker expression when labeled with stage-specific embryonic antigen 4 (SSEA4)

and SRY-Box transcription factor 2 (SOX2). iPSC-derived neurons (iNs) displayed standard

neuronal morphology and neuronal expression of neuron-specific microtubule associated

protein-2 (MAP2), and synapse protein synapsin-1 (SYN1) (**Fig. 1E, Fig. S2B, and Fig. S2E**).

At day 35 (d35), iNs also expressed neurofilament heavy (NFH) and neuronal nuclear protein

(NeuN), characteristic of mature neurons (**Fig. S2C**).

Next, we sought to confirm the neural identity of iNs by comparing intrasample differential

transcript expression in d35 iNs against previously published iN datasets (**Fig. S2D**).[31,50] We

collected RNA from three independent inductions per genotype and performed polyA selected

bulk RNA-seq to target mRNAs. Transcriptional profiles of d35 neurons revealed an enrichment

of genes confirming neuronal identity. In a comparison to the Zhang 2013 dataset, we identified

pan-neuronal (*DCX*, *MAP2*, *TUBB3*), glutamate-related (*SLC17A6*), and post-synaptic (*DLG4*)

genes within our top five hits. Similarly, comparing against the Lindhout 2020 dataset, we see

further enrichment for axon (*GAP43*, *BASP1*), and neuron (*STMN1*, *ENO2*, *SYP*) genes.

Further, we confirm that our iNs express *SERPINI1* ($\log_2$(TPM+1): wild-type: 7.14, heterozygous

KI: 7.30, homozygous KI: 7.21, n = 3). Despite expression of *SERPINI1*, we were unable to

detect NS aggregation in KI iNs throughout the duration of our studies. Our

immunocytochemical and transcriptomic analyses together confirm the neural identity of Ngn2

induced neurons.

iPSC-derived neuron models of FENIB demonstrate typical cortical neuron

electrophysiology.

We assessed the electrophysiological function of iPSC-derived neurons by analyzing evoked

action potentials (AP) using whole-cell patch-clamp recordings in current clamp mode (**Fig.**

S3A-G). While single APs were elicited in all neurons by day 14 (d14), with a burst of firing in a

subset of tested cells, further maturation of iNs until day 42 (d42) resulted in consistent

prototypical APs across all cells recorded. Hence, we assessed d42-44 iNs for intrinsic

excitability. No significant differences were observed between wild-type, heterozygous KI, and

homozygous KI neurons in AP rheobase (WT: 29.7 ± 2.1 pA, n = 54, HET.: 25.4 ± 1.6 pA, n =

46, HOM: 28.1 ± 2.0 pA, n = 47, p>0.05), full-width at half-maxima (WT: 2.0 ± 0.1 ms, n = 52,

HET: 2.0 ± 0.1 ms, n = 46, HOM: 2.0 ± 0.1 ms, n = 47, p>0.05), amplitude (WT: 84.7 ± 1.2 mV, n

= 52, HET: 82.5 ± 1.5 mV, n = 46, HOM: 84.6 ± 1.2 mV, n = 47, p>0.05), firing threshold (WT: -

46.2 ± 1.1 mV, n = 52, HET: -48.5 ± 1.2 mV, n = 46, HOM:-47.9 ± 1.3 mV, n = 47 p>0.05), cell-

capacitance (WT: 15.7 ± 0.7 pF, n = 53, HET: 15.5 ± 0.6 pF, n = 46, HOM: 16.5 ± 0.7 pF, n = 49,

p>0.05), and resting membrane potential (WT: -49.9 ± 1.3 mV, n = 54, HET: -49.5 ± 1.3 mV, n =

45, HOM: -49.9 ± 1.6 mV, n = 50, p>0.05). Our evoked AP findings were consistent with *in vitro*

cortical neurons, validating our human neuronal model of FENIB.[51–53]

Characterizing neuroserpin inclusion bodies in NS-GFP overexpression iPSC-derived neuron FENIB model.

We hypothesized that preventing mutant neuroserpin expression would facilitate clearance of pre-existing aggregated protein in MUT NS-GFP iNs (**Fig. 2A**).[54] To test this, we transduced iPSCs with either doxycycline inducible WT or MUT NS-GFP lentivirus in addition to Ngn2 and rtTA lentivirus for inducing neuronal fate. Triply transduced cells were selected for following doxycycline induction using puromycin and blasticidin, with resistance conferred from Ngn2-T2A-PURO and NS-GFP-T2A-BSD lentivirus respectively.

NS-GFP expression was induced in iNs, with GFP signal observed one day after induction, similar to our HEK293T cell models. By d4, NS-GFP aggregates began forming in MUT NS-GFP cells. WT NS-GFP iNs behaved similarly to WT NS-GFP HEK293T cells, with NS-GFP signal detected as diffuse GFP. We sought to test whether early termination of mutant protein expression would facilitate the clearance of mutant protein aggregates. We induced expression of NS-GFP for 6 days. Beginning on d7, we separated cells into continued expression (+DOX) or discontinued expression through doxycycline withdrawal (W/D) until day 10 (**Fig. 2A**). We imaged d7 prior to separation, d10 +DOX, and d10 W/D cells using immunofluorescent confocal microscopy to identify aggregate formation and clearance in iNs (**Fig. 2B-D**). iNs were labeled with antibodies specific for NeuN and neuroserpin and expressed either WT NS-GFP, MUT NS-GFP, or a no overexpression control (**Fig. 2B**). Images were acquired as Z-stacks to capture fluorescent signal throughout the thickness of the iN culture. We then generated 2D reconstructions of the iNs using the "maximal projection" method in ImageJ to visualize the maximum signal of a given pixel in a 2D picture. d7 and d10 +DOX MUT NS-GFP cells harbored extensive NS-GFP aggregates in the form of GFP puncta which colocalized with neuroserpin immunofluorescent labeling. d7 and d10 +DOX WT NS-GFP cells had diffuse NS-GFP fluorescence which colocalized with neuroserpin immunofluorescent labeling. d10 W/D MUT

NS-GFP cells revealed an almost complete clearance of NS-GFP aggregates. Similarly, d10

W/D WT NS-GFP cells had almost no detectable NS-GFP signal, which was verified by the loss

of neuroserpin immunofluorescent labeling. Aggregated MUT NS-GFP signal produced distinct

GFP puncta which are brighter than diffuse WT NS-GFP. Using ImageJ image analysis

software, we set a fluorescence intensity threshold for GFP expression to establish an unbiased

method to distinguish aggregated NS-GFP from diffuse NS-GFP signal. Next, we used the

ImageJ particle analysis tool to quantify NS-GFP aggregate count and area **(Fig. S4A)**. These

values were normalized to cells labeled with neuronal marker, NeuN, and 4'6-diamidino-2-

phenylindole (DAPI) to obtain NS-GFP aggregate count per neuron and NS-GFP aggregate

area per neuron. By day 7, MUT NS-GFP cells harbored extensive NS-GFP aggregates with no

aggregates in either WT NS-GFP or no overexpression control (ctrl) counterpart cells (d7 MUT

NS-GFP: 4.22 aggregates/neuron, d7 WT NS-GFP: 0 aggregates/neuron, ctrl: 0

aggregates/neuron, p<0.0001). Following doxycycline withdrawal, d10 W/D MUT NS-GFP cells

had significantly reduced aggregate count compared to +DOX counterparts (d10 +DOX MUT

NS-GFP: 3.32 aggregates/neuron, W/D MUT NS-GFP: 0.56 aggregates/neuron, p<0.0001). d10

W/D MUT NS-GFP values were statistically non-significant when compared to WT NS-GFP and

no overexpression control aggregate counts (d10 +DOX ctrl: 0 aggregates/neuron, d10 W/D ctrl:

0 aggregates/neuron, d10 +DOX WT NS-GFP: 0.06 aggregates/neuron, W/D WT NS-GFP: 0

aggregates/neuron) (**Fig. 2C**). Aggregate clearance was also reflected in a reduction in

aggregate area per neuron (d7 MUT NS-GFP: 4.31 μm^2/neuron, d7 WT NS-GFP: 0 μm^2/neuron,

d7 ctrl: 0 μm^2/neuron, p<0.0001). After doxycycline withdrawal, d10 W/D MUT NS-GFP cells

had a statistically significant reduction in aggregate burden compared to continuously induced

d10 +DOX MUT NS-GFP cells (d10 +DOX MUT NS-GFP: 3.70 μm^2/neuron, d10 W/D MUT NS-

GFP: 0.20 μm^2/neuron, p<0.0001) (**Fig. 2D**). When compared to d10 WT NS-GFP and d10 no

overexpression control counterpart cells, d10 W/D MUT NS-GFP aggregate area per neuron

values were statistically non-significant (d10 +DOX ctrl: 0 μm^2/neuron, d10 W/D ctrl: 0

μm^2/neuron, d10 +DOX WT NS-GFP: 0.01 μm^2/neuron, d10 W/D WT NS-GFP: 0 μm^2/neuron).

Based off these findings, early termination of mutant NS expression is sufficient for the

clearance of aggregated protein.

Next, we assessed if prolonged aggregate accumulation could be cleared in more mature

neurons (**Fig. 3A-D**). We induced aggregation until day 21 (d21), then split samples and used

confocal microscopy to visualize and assess per neuron aggregation count and area at d21, day

28 (d28) (+DOX and W/D), and day 35 (d35) (+DOX and W/D) (**Fig. 3A**). Neurons were labeled

with antibodies specific for NeuN and expressed MUT NS-GFP. (**Fig. 3B**). Using the ImageJ

particle analysis method described above, we found that aggregate count remained similar from

d21 to d28. W/D d28 iNs trended towards fewer aggregates, however, aggregate count between

d21, +DOX d28, and W/D d28 iNs were not statistically significant (d21: 0.82

aggregates/neuron, +DOX d28: 0.71 aggregates/neuron, W/D d28: 0.62 aggregates/neuron,

p>0.05) (**Fig. 3C**). When we terminated expression at d21 and assessed clearance in d35

neurons, W/D d35 iNs showed a significant decrease in aggregate count compared to +DOX

counterparts (+DOX d35: 1.24 aggregates/neuron, W/D d35: 0.75 aggregates/neuron, p<0.01).

W/D d35 aggregates per neuron was similar to d21 and d28 neurons (p>0.05). Next, we

assessed aggregate area per neuron. W/D d35 iNs had significantly reduced aggregate area

per neuron compared to +DOX iNs (+DOX d35: 3.2 μm^2/neuron, W/D d35: 1.1 μm^2/neuron,

p<0.0001) (**Fig. 3D**). The difference in aggregate area per neuron identified between +DOX d35

iNs and W/D d35 iNs was not present when comparing W/D d35 iNs to d21 or either +DOX or

W/D d28 iNs (d21: 1.427 μm^2/neuron, +DOX d28: 1.16 μm^2/neuron, W/D d28: 0.89 μm^2/neuron,

p>0.05). Thus, preventing MUT NS-GFP expression at day 21, and allowing cells to resolve

toxic protein aggregation until day 35 was insufficient to clear aggregates. However, by stopping

expression of the mutant protein, we were successful in preventing the formation of new

aggregates. These doxycycline inducible NS-GFP studies demonstrate that early cessation of

protein aggregation allowed for complete aggregate clearance. By contrast, late-stage cessation

prevented further aggregate accumulation.

iPSC-derived neurons overexpressing MUT NS-GFP show changes in membrane
291
excitability.

We next assessed neuronal excitability and physiological function using whole cell patch clamp

in current clamp mode to evoke APs in NS-GFP iNs (**Fig. S5A-G**). Based on the maturation of

neurons at day 42 (**Fig. S3**), NS-GFP iNs at d42 were assessed for intrinsic properties of

excitability including rheobase, full-width at half-maxima, amplitude, firing threshold, cell

capacitance, and resting membrane potential. +DOX MUT NS-GFP iNs had increased rheobase

compared to the WT NS-GFP counterparts (+DOX MUT NS-GFP: 73.1 ± 14.3 pA, n = 8, +DOX

WT MUT NS-GFP: 37.5 ± 5.9, n = 8, p<0.05) **(Fig. S5B)**. This suggests a decrease in

membrane excitability consistent with other reports of protein aggregation based

neurodegenerative disease models. [55,56] W/D d42 MUT NS-GFP (51.4 ± 5.1 pA, n = 7) neurons

had similar rheobase as WT NS-GFP (+DOX WT NS-GFP: 37.5 ± 5.9 pA, n = 8, and W/D WT

NS-GFP: 48.6 ± 7.0 pA, n = 7, p>0.05) and control (+DOX CTRL: 52.5 ± 4.5 pA, n = 8, and W/D

CTRL: 63.3 ± 8.2, n = 9, p>0.05) neurons indicating that alteration to rheobase was rescued

following termination of MUT NS-GFP expression. AP full-width at half-maxima (+DOX CTRL:

1.6 ± 0.2 ms, n = 8, +DOX WT NS-GFP: 1.6 ± 0.1 ms, n = 8, +DOX MUT NS-GFP: 1.4 ± 0.1 ms,

n = 8, W/D CTRL: 1.8 ± 0.2 ms, n = 9, W/D WT NS-GFP: 1.7 ± 0.2 ms, n = 7, W/D MUT NS-

GFP: 1.6 ± 0.1 ms, n = 7, p>0.05), amplitude (+DOX CTRL: 87.0 ± 2.8 mV, n = 8, +DOX WT

NS-GFP: 85.3 ± 3.3 mV, n = 8, +DOX MUT NS-GFP: 84.3 ± 2.0 mV, n = 8, W/D CTRL: 91.3 ±

2.2 mV, n = 9, W/D WT NS-GFP: 85.6 ± 0.8 mV, n = 7, W/D MUT NS-GFP: 88.0 ± 3.6 mV, n = 7,

p>0.05), firing threshold (+DOX CTRL: -49.0 ± 2.7 mV, n = 8, +DOX WT NS-GFP: -52.3 ± 1.6

mV, n = 8, +DOX MUT NS-GFP: -44.3.3 ± 2.9 mV, n = 8, W/D CTRL: -49.1 ± 2.7 mV, n = 9, W/D

WT NS-GFP: -46.2 ± 2.8 mV, n = 7, W/D MUT NS-GFP: -48.8 ± 3.1, n = 7, p>0.05), cell

capacitance (+DOX CTRL: 30.2 ± 3.9 pF, n = 8, +DOX WT NS-GFP: 24.4 ± 1.6 pF, n = 8, +DOX

MUT NS-GFP: 26.1 ± 2.9 pF, n = 7, W/D CTRL: 27.2 ± 3.2 pF, n = 9, W/D WT NS-GFP: 24.0 ±

2.7 pF, n = 8, W/D MUT NS-GFP: 29.8 ± 4.2 pF, n = 6, p>0.05), and resting membrane potential

(+DOX CTRL: -54.1 ± 2.1 mV, n = 11, +DOX WT NS-GFP: -54.2 ± 4.7 mV, n = 6, +DOX MUT

NS-GFP: -56.6 ± 2.9, n = 7, W/D CTRL: -59.5 ± 2.5 mV, n = 8, W/D WT NS-GFP: -57.6 ± 5.0

mV, n = 8, W/D MUT NS-GFP: -61.9 ± 3.3 mV, n = 9, p>0.05) were all similar, with no

statistically significant differences identified.

Based on our finding of dysfunctional membrane excitability due to an increase in rheobase in

+DOX MUT NS-GFP iNs, we plotted an I-O curve to assess firing rate in +DOX and W/D iNs

(**Fig. S5H and Fig. S5I**). When focused in on the minimum current required to increase the

firing rate, we identified a minor trending but statistically non-significant difference between the

initial increase in firing rate of d42 +DOX MUT NS-GFP iNs (40 pA) compared to WT NS-GFP

(20 pA) and CTRL (30 pA) counterparts. Comparisons of d42 W/D iNs revealed no statistically

significant differences. Taken together, our results demonstrate that MUT NS-GFP iNs have

dysfunctional membrane excitability, which is rescued once MUT NS-GFP expression is

terminated.

ABE-mediated correction of *SERPINI1* c.1175 G>A variant.

After identifying that termination of mutant neuroserpin expression through withdrawal of

doxycycline in inducible MUT NS-GFP neurons led to aggregate clearance or prevented further

aggregation, we sought to generate an ABE approach to target and correct the pathogenic

variant, *SERPINI1* c.1175 G>A, back to the wild-type sequence. We identified a shortened 5'-

TG-3' PAM sequence downstream of the patient variant, recognized by ABEs with an SpCas9-

NG C-terminal domain, which positioned the variant within the ABE editing window at the A5

position (**Fig. 4A**). Rationally designed and evolved ABE variants increase editing efficiency and

reduce off-target effects.[57–59] We tested four ABEs to identify an optimal ABE for targeting the

SERPINI1 c.1175 locus. We subcloned the SpCas9-NG C-terminal region into four next-

generation ABEs resulting in NG-ABEmax, NG-SECURE-miniABEmax-V82G (NG-V82G), NG-

SECURE-miniABEmax-K20AR21A (NG-K20AR21A), and NG-ABE8e.[58,60,61] We found a

candidate bystander mutation within the editing window at the A6 position. We hypothesized

that this bystander mutation would be tolerated due to its location at the codon wobble position.

Combination target and bystander editing encodes the wild-type residue (G, glycine), while

bystander-only editing would not alter the pathogenic residue (E, glutamic acid).

We cloned a mutant-specific sgRNA into the plasmid, pBSU6-sgRNA-GFP (g3), which was

nucleofected along with a plasmid encoding ABE using the Lonza P3 nucleofection system in

both heterozygous and homozygous KI iPSCs (**Fig. 4B**). We quantified ABE correction using

EditR Sanger sequencing trace analysis and deep sequencing of the target locus compared to

unedited control cells.[62] EditR analysis of heterozygous KI iPSCs revealed the greatest

correction efficiency with NG-ABE8e (82% A:T to G:C conversion efficiency) and NG-ABEmax

(49%), and modest editing by NG-V82G (10%) and NG-K20AR21A (5%) at the A5 position.

Analysis of homozygous KI iPSCs largely mirrored editing in heterozygous KI iPSCs (NG-

ABE8e: 86%, NG-ABEmax: 48% NG-V82G: 16%, and NG-K20AR21A: 13% A:T to G:C

conversion efficiency). Individuals harboring the *SERPINI1* c.1175 G>A variant are

heterozygous as FENIB variants are often de novo mutations.[10–12] We performed deep

sequencing on heterozygous KI iPSCs treated with the two most efficient ABEs, NG-ABEmax

and NG-ABE8e (**Fig. 4C**). Deep sequencing reads revealed codon specific editing outcomes

with NG-ABEmax encoding 75% wild-type glycine (GGA or GGG) and NG-ABE8e encoding

92% glycine. Further analysis of glycine encoding reads revealed that NG-ABEmax activity

resulted in more reversion to the wild-type codon (GGA: 62%, GGG: 13%) while NG-ABE8e had

more codons harboring the permissive bystander mutation (GGA: 35%, GGG: 57%).

We treated MUT NS-GFP iPSCs with plasmids encoding g3 and NG-ABE8e to assess if NG-

ABE8e editing could correct MUT NS-GFP iPSCs (**Fig. 4D**). EditR base calling analysis of

Sanger sequencing traces revealed highly efficient editing in both homozygous KI (90%) and

MUT NS-GFP (65%) iPSCs. Thus, NG-ABE8e with g3 can efficiently correct the variant within

the endogenous genomic context in heterozygous and homozygous KI cells, and as the stably

incorporated MUT NS-GFP model of FENIB.

ABE correction of MUT NS-GFP in HEK293T cells allows aggregate clearance.

After identifying a highly efficient method of base editing the pathogenic variant back to the wild-

type *SERPINI1* sequence, we evaluated ABE treatment on NS-GFP aggregate clearance. Using

lentivirus transduction, we stably incorporated either WT or MUT NS-GFP constitutively

expressed under the CMV promoter into HEK293T cells. 72-hours after transduction, we sorted

GFP positive cells into single cells to generate biological replicates. Cells were clonally

expanded and expressed WT NS-GFP or MUT NS-GFP control for 3 weeks to induce extensive

NS-GFP aggregation. MUT NS-GFP HEK293T clones were transfected with NG-ABE8e-T2A-

tagRFP and g3 plasmids co-expressing tagRFP (ABE + g3), or NG-ABE8e-T2A-tagRFP only

(ABE only) as control. 72-hours post-transfection, GFP+/RFP+ double positive cells were either

bulk or single cell sorted to assess DNA correction and NS-GFP aggregate clearance. EditR

analysis of bulk sorted cells revealed highly efficient correction of the variant back to the wild-

type sequence in ABE + g3 MUT NS-GFP cells (71% A:T to G:C conversion efficiency) (**Fig.**

5A). We sought to visualize NS-GFP expression and aggregation using confocal microscopy in

ABE + g3 MUT NS-GFP cells, and ABE only MUT NS-GFP, MUT NS-GFP pretreatment, and

WT NS-GFP control cells (**Fig. 5B**). Native NS-GFP fluorescence was imaged alongside

immunocytochemical labeling of neuroserpin using an antibody which recognizes all

neuroserpin forms (1A10, all NS) and an antibody specific for aggregated neuroserpin polymers

(purified polyclonal antibody, MUT poly). [63] We used Cellpose to count DAPI stained nuclei and

determined total cells per imaging field to assess for culturing fitness differences. We did not

identify any statistically significant changes (WT NS-GFP: 224 cells, pretreatment MUT NS-

GFP: 235 cells, ABE only MUT NS-GFP: 233 cells, ABE + g3 MUT NS-GFP: 190 cells, p>0.05)

(**Fig. 5C**). Immunoblotting whole cell lysate from ABE + g3 MUT NS-GFP, or ABE only MUT NS-

GFP and WT NS-GFP controls revealed similar levels of NS-GFP protein expression when

probed with anti-neuroserpin antibody (**Fig. S6A**). Therefore, any differences observed in

aggregated and diffuse NS-GFP expression was caused by protein aggregation, as opposed to

total protein expression.

Next, we quantified the effects of ABE + g3 treatment of MUT NS-GFP cells on NS-GFP

aggregation (**Fig. 5D and 5E**). ABE + g3 MUT NS-GFP cells had statistically significant reduced

aggregates per cell values compared to pretreatment MUT NS-GFP and ABE only MUT NS-

GFP cells (pretreatment MUT NS-GFP: 3.98 aggregates/cell, p<0.01, ABE only MUT NS-GFP:

5.77 aggregates/cell, p<0.0001, ABE + g3 MUT NS-GFP: 1.22 aggregates/cell) (**Fig. 5D**). ABE +

g3 treatment was statistically non-significant when compared to WT NS-GFP control cells (WT

NS-GFP: 0 aggregates/cell). ABE + g3 MUT NS-GFP cells also had a statistically significant

reduction in NS-GFP aggregate area per cell compared to pretreatment MUT NS-GFP and ABE

only MUT NS-GFP cells (pretreatment MUT NS-GFP: 3.61 μm^2/cell, p<0.05, ABE only MUT NS-

GFP: 6.23 μm^2/cell,p<0.0001, ABE + g3 MUT NS-GFP: 1.02 μm^2/cell) (**Fig. 5E**). ABE + g3

treatment was sufficient to restore aggregate area per cell back to WT NS-GFP (WT NS-GFP: 0

μm^2/cell, p>0.05).

Next, we analyzed ABE + g3 treated single clones from two MUT NS-GFP HEK293T clones,

Clone A and Clone B (**Fig. 5F and Fig. S6B**). Sanger sequencing of ABE + g3 treated single

clones revealed that all clones (n = 26 total, Clone A = 6, Clone B = 20) were either corrected to

the wild-type sequence (c.1175 G, Clone A = 3/6, Clone B = 16/20), or remained unedited

(c.1175 G>A, Clone A=3/6, Clone B = 4/20), with no partially edited cells. Quantification of NG-

ABE editing in single clones reflected editing efficiencies previously identified in bulk sorted cells

(single clones: 72%, bulk sorted cells: 71%) (**Fig. 5F**). Following Sanger sequencing, we used

epifluorescence microscopy to detect NS-GFP aggregates in ABE + g3 treated single clones

(**Fig. S6B**). All corrected clones with the wild-type (c.1175 G) sequence had diffuse NS-GFP

expression, while all unedited clones harbored NS-GFP aggregates. Based on our results, we

conclude that ABEs can correct the FENIB mutation and allow aggregate clearance following

extensive NS-GFP aggregation. Further, our clonal assay demonstrates that ABE correction is

sufficient for total aggregate clearance in single cells.

Impaired dendritic arborization in heterozygous and homozygous KI iPSC-derived

neurons is partially restored with ABE treatment.

Neuroserpin is expressed during late stage neuronal maturation with roles in dendritic

morphology, synaptic maturation, and neuronal differentiation.[64–66] Cultured rat hippocampal

neurons overexpressing neuroserpin have increased dendritic arborization and altered spine

shapes.[65] *In vivo* analysis of neuroserpin-deficient mice revealed a decrease in spine-

synapses.[67] We evaluated whether dendritic morphology was affected in heterozygous and

homozygous KI neurons, and whether ABE corrected neurons showed restored morphology

(Fig. S7A). We transfected d35 WT, heterozygous KI, homozygous KI, and ABE corrected

neurons with GFP expressing plasmids for 30 minutes to sparsely label single cells. 24-hours

post-transfection, we fixed coverslips for staining with anti-GFP antibodies to increase signal in

imaging for Sholl analysis. Using ImageJ, we set a fluorescence intensity threshold against GFP

expression to image the morphology of single neurons. Then using Simple Neurite Tracer

(SNT), we assessed the number of dendritic intersections at 10 μm radial distances from the

neuron soma **(Fig. 6A).**[68,69]

We first sought to assess differences between FENIB KI iNs at d35 **(Fig. 6B-D)**. We observed

no significant differences in dendritic intersections between WT, heterozygous KI, homozygous

KI, and ABE corrected neurons at d35 for any radial distance assessed. Intersection numbers

peaked at 20 µm for WT (9.6 dendrites), heterozygous KI (9 dendrites), and ABE corrected (7.3

dendrites) and at 30 µm for homozygous KI (8.6 dendrites) (**Fig. 6B**). Using the number of

intersections at radial distances, we calculated area under the curve (AUC) to determine overall

dendritic arborization (**Fig. 6C**). Although WT and ABE corrected neurons appeared to have

slightly greater AUC than both KI groups, statistical testing reveal no significant difference (WT:

819 arbitrary units (AU), heterozygous KI: 752 AU, homozygous KI: 620 AU, ABE corrected: 853

AU, $p>0.05$). Similarly, we quantified total dendritic length and found no statistically significant

differences (WT: 1022.2 µm, heterozygous KI: 938.4 µm, homozygous KI: 864.4 µm, ABE

corrected: 1177.5 µm, $p>0.05$) (**Fig. 6D**). We next sought to test differences in number of

intersections at 10 µm increments from the soma across all groups using a two-way ANOVA

with Bonferroni correction. We identified statistically significant differences in dendrite length in

ABE corrected neurons compared to heterozygous KI and homozygous KI neurons at 70-130

µm radial distances from the soma (70 µm: ABE corrected: 6.2 intersections, homozygous KI:

3.6 intersections, $p<0.05$, 80 µm: ABE corrected: 5.8 intersections, homozygous KI: 2.9

intersections, $p<0.01$, 90 µm: ABE corrected: 5.9 intersections, homozygous KI: 2.7

intersections, $p<0.01$, heterozygous KI: 3.5, $p<0.05$, 100 µm: ABE corrected: 6.6 intersections,

homozygous KI: 2.3, $p<0.0001$, heterozygous KI: 3.6, $p<0.01$, 110 µm: ABE corrected: 5.6

intersection, homozygous KI: 2.3 intersections, $p<0.01$, 120 µm: ABE corrected: 5.1

intersections, homozygous KI: 2.2 intersections, $p<0.01$, 130 µm: ABE corrected: 5.1

intersections, homozygous KI: 2 intersections, $p<0.01$). WT and ABE corrected neurons

remained largely similar, with one statistically significant difference identified at 100 µm (ABE

corrected: 6.6 intersections, WT: 3.6 intersections, $p<0.05$).

As neuroserpin functions in late-stage neuronal maturation, we allowed further neuronal

development and assessed dendritic morphology again at d42 (**Fig. 6E-G)**. Intersection number

peaked at 30 μm for WT (11.6 dendrites), homozygous KI (7.9 dendrites), and ABE corrected

neurons (11.8 dendrites), and at 20 μm for heterozygous KI (8.4 dendrites) (**Fig. 6E**). When we

compared AUC for d42 neurons, we identified a significant impairment in overall dendritic

arborization in heterozygous and homozygous KI neurons compared to WT (WT: 1293 AU,

heterozygous KI: 822.5 AU, homozygous KI: 887.8 AU, $p < 0.01$ for WT vs. heterozygous KI,

$p < 0.05$ for WT vs. homozygous KI) (**Fig. 6F**). Comparison of heterozygous and homozygous KI

to ABE corrected neurons was trending but statistically non-significantly (ABE corrected: 1174

AU, $p > 0.05$). When we compared AUC between WT and ABE corrected neurons, we identified

no differences in overall dendritic arborization ($p > 0.05$), suggesting that ABE correction results

in neurons with dendritic arborization phenotype in-between WT and KI variant neurons. We

identified a similar impairment when we assessed total dendritic length in d42 neurons (**Fig.**

6G). Heterozygous and homozygous KI neurons had reduced total dendritic length compared to

WT control neurons (WT: 1704.7 μm, heterozygous KI: 1086.4 μm, homozygous KI μm: 1093.8

μm, $p < 0.05$). Once again, ABE corrected neurons were statistically non-significant from either

WT or KI neurons (ABE corrected: 1536 μm, $p > 0.05$). Next, we ran a two-way ANOVA with

Bonferroni correction to test differences in number of intersections at 10 μm radial distances

from the soma across WT, heterozygous KI, homozygous KI, and ABE corrected iNs. Similar to

d35, we identified statistically significant differences between both WT and heterozygous and

homozygous KI neurons, and ABE corrected and heterozygous and homozygous KI neurons.

These differences occur at 20-40 and 60-90 μm radial distances from soma for WT neurons (20

μm: WT: 11.4 intersections, homozygous KI: 7.0 intersections, $p < 0.01$, 30 μm: WT: 11.6

intersections, heterozygous KI: 7.8 intersections, $p < 0.05$, homozygous KI: 7.9 intersections,

$p < 0.05$, 40 μm: WT: 10.7 intersections, homozygous KI: 7.4 intersections, $p < 0.05$, 60 μm: WT:

10.5 intersections, heterozygous KI: 6.5 intersections, $p < 0.01$, homozygous KI: 7.1

intersections, $p < 0.05$, 70 μm: WT: 10.2 intersections, heterozygous KI: 6.1 intersections, $p < 0.01$,

homozygous KI: 5.9 intersections, $p < 0.01$, 80 μm: WT: 9.1 intersections, heterozygous KI: 5.1

intersections, p<0.01, homozygous KI: 5.4 intersections, p<0.05, 90 µm: WT: 8.8 intersections,

heterozygous KI: 4.3 intersections, p<0.01, homozygous KI: 5.3 intersections, p<0.05) and 20-

40 µm radial distances for ABE corrected neurons (20 µm: ABE corrected: 10.4 intersections,

homozygous KI: 7.0 intersections, p<0.05, 30 µm: ABE corrected: 11.8 intersections,

heterozygous KI: 7.8 intersections, p<0.001, homozygous KI: 7.9 intersections, p<0.01, 40 µm:

ABE corrected: 11.1 intersections, heterozygous KI: 7.9 intersections, p<0.05, homozygous KI:

7.4 intersections, p<0.05). These findings support a developmental impairment arising from the

mutant neuroserpin activity in addition to the toxic gain-of-function of aggregating mutant

neuroserpin.

eVLP-mediated delivery of ABEs can efficiently correct the FENIB variant.

Delivery of gene editors to target tissues is an outstanding challenge for therapeutic genome

editing.[36,70] We decided to test eVLPs to target neurons due to the high cell-targeting efficiency

and subsequent editing, and simplicity in reconfiguring eVLP constructs. We produced eVLPs

by transfecting HEK293T cells with plasmids encoding VSV-G glycoprotein, MMLV-gag-pol, and

MMLV fused NG-ABE8e ribonucleoprotein, henceforth called the eVLP plasmids (**Fig. 7A**). We

first tested eVLP editing on target loci with documented high editing; *HEK2* and *HEK3* (**Fig. 7B**

and 7C).[39] To generate eVLPs, we transfected HEK293T cells with the eVLP plasmids, and

collected the eVLP containing media after 72 hours.[39] We concentrated the supernatant 100-fold

using PEG-IT, a polyethylene glycol (PEG) based reagent. Next, we treated HEK293T cells with

100-fold concentrated eVLPs and sequenced the target loci 72 hours after treatment. We

observed robust editing at both the *HEK2* and *HEK3* loci, with the greatest editing at the A7

position (88%) and A6 position (64%), respectively.

After confirming the capacity of eVLPs to deliver ABE and edit target loci in HEK293T cells, we

sought to test eVLP editing in *SERPINI1* KI iPSCs. We treated homozygous KI iPSCs with eVLP

containing supernatant in Stemflex iPSC media at ratios of 1:10, 1:3, and 1:1 (**Fig. S7B**). After

72 hours, we collected DNA from the treated cells and Sanger sequenced the samples at the

SERPINI1 c.1175 G>A locus. Without concentration or selection, we observed >10 % editing at

the A5 and A6 positions in our 1:1 ratio of eVLP containing supernatant to Stemflex media when

using EditR Sanger sequencing analysis. We next generated concentrated eVLPs and treated

homozygous KI iPSCs and sequenced the target locus 72 hours after treatment without

selection (**Fig. 7D**). eVLP editing using EditR revealed a dosage effect, with the 30 µl high dose

editing resulting in 22.5% correction to the wild-type sequence at the A5 position and 25%

correction at the A6 position.

We assessed neural editing by treating homozygous KI neurons with optimized eVLP

conditions. Concentrated eVLPs were added with neurotrophic factors, ARAC, and doxycycline

at d7 of neural induction. Treated neurons were collected without selection and Sanger

sequenced at the target locus after 72 hours (**Fig. S7C**). We observed a dosage effect in our

homozygous KI iNs. However, our highest dose, 30ul, only reached 2% at the A5 and 1.5%

editing at the A6 position, necessitating further optimization.

We tested a modified rabies virus glycoprotein, FuG-B2, to increase eVLP uptake through

enhancing neuronal specificity and selectivity.[39,71] Homozygous KI neurons treated with

concentrated eVLPs enveloped with FuG-B2 glycoprotein increased editing to 1% at the A5 and

3% at the A6 positions (**Fig. S7D**). To optimize neuron specific nuclear localization, we

generated NG-ABE8e flanked by Nipah virus W protein (NiV W), methyl CpG binding protein 2

(MeCP2), and heterogenous nuclear ribonucleoprotein D (hnRNP D) NLSs.[72] We produced

each NLS variant in eVLPs with both VSV-G and FuG-B2 glycoproteins. For VSV-G eVLPs, NiV

W NLS (1.7%) resulted in the highest correction at the A5 position, followed by MeCP2 NLS

(0.6%), BP/SV40 NLS (0.5%), and hnRNPD NLS (0%) (**Fig. S7E**). FuG-B2 eVLP results were

similar with NiV W NLS (3.2%) resulting in the highest correction efficiency, followed by

BP/SV40 NLS (2.9%), MeCP2 NLS (0.8%), and hnRNPD NLS (0%) (**Fig. S7F**). The difference

in eVLP editing efficiencies between HEK293T cells and iPSCs, and iPSC-derived neurons

suggested an issue with eVLP uptake.

We hypothesized that eVLP-cell interactions were insufficient for eVLP uptake and decided to

test the addition of polybrene (**Fig. 7E**). Polybrene is a cationic polymer which facilitates

receptor-independent virus adsorption.[73] Both 5 µg/ml and 10 µg/ml polybrene increased eVLP

editing by 10-fold in VSV-G eVLPs. eVLP editing in homozygous KI neurons with VSV-G

glycoprotein with 5 µg/ml polybrene had the highest editing efficiency at the A5 position (28%)

while 10 µg/ml polybrene followed closely (18%).

We assessed eVLPs with neuron specific NLS with 5 µg/ml polybrene to determine optimal

neuron editing. We treated homozygous KI neurons with VSV-G glycoprotein eVLPs and either

BP/SV40 or neuron-specific NLS (**Fig. 7F**). Homozygous KI neurons treated with eVLPs with

NiV W NLS (38.5%, p<0.0001) had the highest editing efficiency, significantly greater than

BP/SV40 (20.9%). We also found that MeCP2 NLS (29.7%, p<0.01) had statistically increased

editing at the A5 position. Editing between hnRNP D NLS and BP/SV40 was statistically non-

significant (18.8% p=0.76). In addition to differences in editing efficiency, we found that neuron

specific NLS eVLPs had different editing profiles than standard BP/SV40. There was low but

detectable editing at all A positions along the gRNA. BP/SV40 had the broadest editing window

with A5 (20.9%) A6 (13.7%). The neuronal NLS editors had a higher ratio of desired A5 to

bystander A6 editing, with NiV W 38.5%:10.9%, MeCP2 29.7%:7.2%, and hnRNP D at

18.8%:4.7%.

We next assessed eVLP editing of MUT NS-GFP neurons (**Fig. S7G**). VSV-G with BP/SV40

NLS (13.4%) had greater editing efficiency at the A5 position. Unlike homozygous KI neurons,

eVLP editing of the MUT NS-GFP loci by NiV W (4.8% p<0.0001), MeCP2 (0.4%, p<0.0001),

and hnRNP D (0.16% p<0.0001) were statistically less efficient than BP/SV40. Thus, neuron

specific NLS eVLPs increase editing efficiency in KI neurons and increased the desired A5 to

bystander A6 editing ratio, but not in the MUT NS-GFP neurons.

Discussion

We aimed to generate patient-specific models of protein conformational disease to assess

whether ABEs can efficiently correct the pathogenic variant, and if preventing novel mutant

protein expression would facilitate clearance of neuroserpin aggregates. Our results support

that CRISPR/Cas-mediated genome editing systems can model and eliminate aggregates or

prevent aggregation progression in this rare patient-specific variant of brain disease using

HEK293T cells, iPSCs, and neuron models.

We hypothesized that disease modifying treatments for FENIB would require resolving NS

aggregation, similar to potential therapies for Alzheimer's and Huntington's diseases which

target aggregate-prone protein precursors.[16,19] Doxycycline withdrawal experiments

demonstrated that early prevention of mutant protein expression facilitates complete clearance

of NS-GFP aggregates. Whether clearance of aggregates resulted from reduced aggregate

burden or because neurons were younger remains unclear. The average number of aggregates

per cell and size of aggregates were similar in d10 neurons and d21 neurons, suggesting that

the difference in aggregate clearance reflected neuron age, not aggregate burden. In more

mature neurons, we were unable to achieve complete clearance, but prevented further NS-GFP

aggregation. Coupling prevention of novel protein production with methods to promote protein

turnover may facilitate aggregate clearance. Embelin, a small molecule derived from the

Japanese Ardisia (*Ardisia japonica*) herb, prevented *in vitro* NS aggregation and dissolved

aggregates.[74,75] ABE correction of the pathogenic variant to prevent new aggregation combined

with embelin disruption of existing aggregates is a potential effective therapy.

Research of conformational neurodegenerative diseases is often limited by experimental

timescale where aggregate burden requires aging over years or decades for accumulation and

neurotoxicity.[76,77] Overexpression of NS overcomes low levels and reduced overall duration of

endogenous NS protein expression, and has been used to successfully study FENIB and NS

aggregation both *in vitro* and *in vivo*.[5,14,46,49,63,63,78–81] However, overexpression of NS-GFP to

study aggregation obscures the potential phenotypes which may arise given patients'

heterozygosity. We complemented NS-GFP with heterozygous and homozygous KI neurons

and identified a novel developmental deficit in dendritic arborization and total dendritic length.

Dendritic spines are directly involved in receiving excitatory synaptic inputs in excitatory cortical

neurons. Impaired dendritic spines are a known abnormality in epilepsy, but it remains unclear

whether this is a cause or result of epilepsy.[82] Dendritic pathologies have been also identified in

neurological disorders with epilepsy including Rett and Down syndrome.[83,84] Approximately 90%

of patients with Rett syndrome have epilepsy. Knockdown of endogenous *Mecp2*, the gene

implicated in Rett syndrome, using shRNAs reduced the number of dendritic nodes, total

dendritic length, and dendritic intersections in a neuronal model.[85] Further analysis of dendritic

spines in FENIB will be illustrative to assess whether heterozygous and homozygous KI

neurons suffer impairments to spine morphology in addition to dendritic arbor.

Whole-cell patch clamp electrophysiology revealed no differences between WT and

heterozygous or homozygous KI neurons, suggesting that partial or complete loss of wild-type

NS protein does not directly cause seizures. Because our model generates cortical excitatory

neurons, it remains possible that other neuronal subtypes, specifically inhibitory neurons, could

be affected in FENIB. Additionally, our *in vitro* cultures may not sufficiently model the

interactions between NS and tissue-type plasminogen activator (tPA).[86–88] tPA enhances brain

N-methyl-D-aspartate receptor (NMDAR) activity through a mechanism that has yet to be

completely understood.[89,90] NS protects against NMDAR excitotoxicity *in vivo* and *in vitro* by

limiting tPA's pro-excitotoxic effects.[91] Likely, further *in vivo* modeling of aged FENIB neurons will

be required to assess whether the NS-tPA and NMDAR interactions are responsible for the

epileptic phenotype.

Delivery of genome editors larger than 5kb remains a challenge. This is due to the genomic

packaging limitations of AAVs. eVLPs with neuron-specific NLSs improved editing efficiency of

in vitro cortical excitatory neurons with the additional benefit of narrowing the on-target:

bystander editing ratio. The delivery method of genome editors affects the duration of Cas9

expression. Plasmids-based methods results in days, AAV-based methods in months, and

lentivirus-based methods in lifelong expression. eVLPs deliver genome editors as

ribonucleoprotein, resulting in the turnover of most protein after 24 hours. [92] We hypothesize

that due to the minimal window of expression, eVLP mediated ABE delivery requires faster

nuclear import observed when using neuron-specific NLSs. [72] For this reason, we envision that

neuron-specific NLS's will improve editing with both AAV and lipid-nanoparticle RNPs, as both

require NLSs for nuclear entry.

In summary, our CRISPR/Cas genome editors effectively modeled and corrected the severe

patient-specific FENIB variant. Our modeling strategy uses complementary cellular systems to

investigate partial or complete loss of wild-type NS and toxic gain-of-function present in FENIB.

Heterozygous and homozygous KI neurons revealed developmental deficits resulting in

impaired dendritic morphology. Early termination of MUT NS-GFP expression can completely

clear aggregates, while late termination prevented further NS-GFP aggregation. Treatment of

MUT NS-GFP cells with ABE facilitated NS-GFP clearance. Finally, we increased editing

efficiency in neurons using NiV W NLS peptide to shuttle NG-ABE8e into neuronal nuclei.

Materials and Methods

iPSC culture

Induced pluripotent stem cells (iPSC) were cultured in Stemflex media (Gibco; Grand Island,

NY) supplemented with 1% antibiotic-antimycotic (Gibco; Grand Island, NY) with daily media

changes. Cells were split and passaged at 70% confluency, approximately every 3-4 days,

using 0.5mM EDTA (Invitrogen; Waltham, MA) in DPBS (Gibco; Grand Island, NY) or Accutase

(Innovative Cell Technologies; San Diego, CA) and plated onto geltrex (Gibco; Grand Island,

NY) coated tissue culture treated wells. When passing, iPSCs were plated in Stemflex media

with 1% antibiotic-antimycotic supplemented with 10uM ROCK-inhibitor (LC Laboratories;

Woburn, MA). Cell media was changed with Stemflex media without ROCK-inhibitor the

following day. Cells were maintained at 37°C with 5.0% CO_2.

Neural induction

iPSC-derived neurons were generated through doxycycline (Sigma Aldrich; Saint Louis, MO)

induced expression of Ngn2.[31] Briefly, iPSCs were detached and passaged onto geltrex coated

plates, and transduced with lentivirus expressing ubiquitin-C promoter expressed reverse

tetracycline transactivator, or tetracycline response element On promoter expressed mouse

Neurogenin2-T2A-puromycin (*Ngn2*) resistance in Stemflex media with 10uM ROCK-inhibitor.

On day 1, culturing media was changed to iN media consisting of Neurobasal medium (Gibco;

Grand Island, NY) with 1X B-27 (Gibco; Grand Island, NY) and further supplemented with 10 µM

ROCK-inhibitor and doxycycline to induce *Ngn2* expression. Culturing media was replaced

every day with iN media until day 6, with doxycycline induction from days 1-7, and puromycin

(Invivogen; San Diego, CA) selection from days 2-5. On day 6, cells were detached and

replated onto glia isolated from p3 mouse pups in iN media supplemented with 5% FBS

(GeminiBio; West Sacramento, CA) on either geltrex coated glass coverslips for

immunocytochemical labeling and patch clamp electrophysiology or plastic tissue culture plates

for RNA and protein processing. Beginning d7, culturing media was replaced every 2-3 days

with iN media supplemented with BDNF (STEMCELL Technologies; Vancouver, Canada) GDNF

(STEMCELL Technologies, Vancouver, Canada), NT-3 (STEMCELL Technologies; Vancouver,

Canada), and ARAC (Selleckchem; Houston, TX). Cells were maintained at 37°C with 5.0%

CO_2.

HEK293T cell culture

HEK293T cells were cultured in media consisting of DMEM (Gibco; Grand Island, NY)

supplemented with 10% fetal bovine serum (FBS) and 1% antibiotic-antimycotic with media

changes every 2-3 days. Cells were passaged every 5-7 days or when 70% confluent using

0.25% trypsin/EDTA (Gibco; Grand Island, NY). Cells were maintained at 37°C with 5.0% CO_2.

Plasmids

The plasmid pSpCas9(BB)-2A-GFP (PX458) was a gift from Feng Zhang (#48138, Addgene),

the plasmid pTet-O-Ngn2-puro was a gift from Marius Wernig (#52047, Addgene), the plasmid

FUW-M2rtTA was a gift from Rudolf Jaenisch (#20342, Addgene), the plasmids pCMV_ABEmax

(#112095, Addgene), ABE8e (#138489, Addgene), and pCMV-MMLVgag-3xNES-ABE8e-NG

(#181754, Addgene) were gifts from David Liu, the plasmids SECURE miniABEmax-V82G

(pJUL1828) (#131313, Addgene) and SECURE miniABEmax-K20A/R21A (pJUL1774)

(#131312, Addgene) were gifts from Keith Joung, the plasmid pBS-CMV-gagpol was a gift from

Patrick Salmon (#35614, Addgene), the plasmid pCMV-VSV-G was a gift from Bob Weinberg

(#8454, Addgene), the plasmid pCAGGS-FuG-B2 was a gift from Kazuto Kobayashi (#67513,

Addgene), the plasmids cTNT-BLAST-T2A-eGFP-RP, psPAX2, and pMD2G were gifts from Dr.

Lei Bu, and the plasmid pBSU6_FE_Scaffold_RSV_GFP was a gift from Dr. Dirk Grimm.

The plasmid SpCas9-NG was generated by introducing the mutations L1111R, D1135V,

G1218R, E1219F, A1322R, R1335V, T1337R into the pSpCas9(BB)-2A-GFP (PX458) plasmid

using primer directed mutagenesis and Gibson assembly. [41] gRNAs targeting *SERPINI1* c.1175

G>A were cloned into the plasmid pBSU6_FE_Scaffold_RSV_GFP. The C-terminal portion of

SpCas9-NG fused to 2A-GFP expressing fragment was amplified from the SpCas9-NG plasmid

and cloned into the ABEmax, ABE8e, V82G, and K20A/R21A ABE plasmids using Gibson

assembly. pTet-O-WT-NS-GFP-BSD and pTet-O-MUT-NS-GFP-BSD plasmids were cloned into

pTet-O-Ngn2-puro plasmids. Wild-type (WT) and mutant (MUT) *SERPINI1* cDNA were amplified

from cDNA synthesized from wild-type iPSC-derived neurons and homozygous KI iPSC-derived

neurons expressing *SERPINI1* transcripts, respectively. The GFP expressing fragment was

amplified from pSpCas9(BB)-2A-GFP. A 3X glycine-glycine-glycine-serine linker sequence was

added between the *SERPINI1* and GFP sequences by primer directed mutagenesis. The 2A-

BSD expressing fragment was amplified from cTNT-BLAST-T2A-eGFP-RP. The *SERPINI1*,

GFP, and 2A-BSD fragments were cloned into the pTet-O-Ngn2-puro plasmid using Gibson

assembly. CMV-MMLVgag-3xNES-NiVW-ABE8e-NG, pCMV-MMLVgag-3xNES-MeCP2-ABE8e-

NG, and pCMV-MMLVgag-3xNES-hnRNPD-ABE8e-NG plasmids were generated by replacing

the SV40 and BP NLS sequences in the pCMV-MMLVgag-3xNES-ABE8e-NG plasmid with

neuron-specific NLS sequences using primer directed mutagenesis and Gibson assembly. All

Gibson assembly primers are listed in Table 1, all plasmid sequences are listed in Table 2, all

plasmid sequencing primers are listed in Table 3.

Plasmids were transformed into DH5A (Invitrogen; Waltham, MA) or Stbl3 (Invitrogen; Waltham,

MA) bacteria, and amplified using Nucleobond Xtra Maxi plasmid purification kit (Macherey-

Nagel; Allentown, PA). Plasmid sequences were verified using Sanger sequencing or Plasmid-

EZ (Genewiz; Waltham, MA).

Transfections

Transfections were performed using nucleofections, lipofectamine, or polyethylenimine.

Nucleofections used 4D nucleofector system with P3 or SF reagent (Lonza; Morris Township,

NJ) for iPSCs and HEK293T cells respectively, per manufacturer's guidelines. For generating

the knock-in model, cells were nucleofected with 5 µg px458 or NG-px458 plasmid with locus

specific gRNA and 5 µl of 10 µM ssODN (Integrated DNA technologies; Coralville, IA).

Lipofection was performed using lipofectamine 2000 (Invitrogen; Waltham, MA) as reverse

transfections, per manufacturer's guidelines. PEI (Sigma Aldrich; Saint Louis, MO) transfections

were performed as reverse transfections, using a 1:5 ratio of plasmid to PEI in Opti-MEM

(Gibco; Grand Island, NY) All gRNA sequences are listed in Table 4.

eVLP and lentivirus preparation

Lentivirus and eVLP vectors were produced by reverse transfection of 1.2×10^7 HEK293T cells

using PEI in Opti-MEM. To generate lentivirus, cells were reverse transfected with psPAX2 (4.7

µg), pMD2.G (1.6 µg), and expression plasmids (6.3 µg) per 10 cm dish. To generate eVLPs,

cells were reverse transfected with VSV-G or FuG-B2 (400 ng), MMLVgag-pro-pol (3,375 ng),

MMLVgag-3XNES-NG-ABE8e or neuron-specific NLS variants (1,125 ng), and sgRNA (4,400

ng) plasmids per 10 cm dish. 24-hours post-transfection, culturing media was replaced with

DMEM+/+. 72-hours post-transfection, vector containing media was harvested and centrifuged

for 10 min at 500 RCF to pellet cells. Vector containing supernatant was filtered using a 0.45 µm

PVDF syringe filter (Corning; Corning, NY). Vector supernatant was concentrated 100-fold using

Lenti-X concentrator (Takara Bio; Shiga, Japan) or PEG-IT virus precipitation solution (System

Biosciences; Palo Alto, CA) per manufacturer's instructions in cold 1 X DPBS. Concentrated

vectors were either immediately used or stored at -80°C avoiding multiple freeze-thaws.

Flow cytometry

HEK293T cells and iPSCs were dissociated into a single cell suspension using 0.25%

trypsin/EDTA or Accutase, respectively, and sorted 72-hours post-transfection using either a

MoFlo-XDP or SY3200 cell sorter. Cells were gated via doublet discrimination, followed by

live/dead viability based on DAPI staining. Cells were sorted by GFP or GFP and RFP signal

based on the experiment. Cells were sorted using a 100 μm nozzle. To generate stable clones,

cells were single-cell sorted into 96-well plates into DMEM +/+ for HEK293T cells or Stemflex

supplemented with 1X CloneR (STEMCELL Technologies; Vancouver, Canada) for iPSCs.

DNA extraction

DNA extraction was performed on a minimum of 10,000 cells. Cells were processed with

DirectPCR lysis-reagent cell (Viagen; Los Angeles, CA) to extract crude DNA. Crude DNA was

directly used for PCR.

PCR

GoTaq PCR (Promega; Madison, WI) was used to generate amplicons using sequence-specific

primers (Integrated DNA technologies; Coralville, IA) in the Mastercycler nexus X2 thermal

cycler (Eppendorf; Hamburg, Germany). Primer sequences can be found in Table 5.

T7E1, Synthego ICE, and EditR

T7 endonuclease I (T7E1) assay was performed by reannealing raw PCR product from GoTaq

PCR amplification followed by incubation of reannealed product with T7E1 for digestion of

mismatched paired strands. Synthego ICE analysis was performed by comparing Sanger

sequencing .ab1 files per application instructions.[44] Briefly, .ab1 Sanger sequencing traces from

edited samples are provided along with unedited control traces, sgRNA sequences, and HDR KI

template sequences. These files are used to compare the edited sequence against the unedited

control and determine the percentage of signal corresponding to the intended KI sequence,

nonspecific indels, and unedited wild-type sequence. EditR analysis was performed by

calculating the difference in base calls at the target locus between ABE treated and unedited

control cells as determined by EditR software (1.0.10).[62] Sanger sequencing trace files were

uploaded in combination with gRNA sequence, with P-value cutoff at the default value, 0.01. 5'

start and 3' ends were determined empirically based off Sanger sequencing trace quality for

each locus tested as follows: *SERPINI1* 5' start: 50bp, 3' end: 280 bp, *HEK2* 5' cut start: 50 bp,

3' end: 280 bp, *HEK3* 5' start: 50 bp, 3' end: 550 bp.

Immunocytochemistry

Cells were plated on geltrex coated 12 mm No.1 coverslips (Chemglass Life Sciences;

Vineland, NJ) in a 24-well plate. At experimental endpoints, cells were washed with DPBS to

remove residual media, then fixed in 4% PFA (Electron Microscopy Sciences; Hatfield, PA) in

PBS at room temperature. Samples were subsequently permeabilized using 0.1% triton-x 100

(Sigma Aldrich; Saint Louis, MO) in PBS, washed three times with PBS, then blocked using 10%

either donkey serum (Abcam; Waltham, MA), 0.25% triton-x 100 in PBS, and washed three

times with PBS. Cells were incubated with primary antibodies overnight at 4°C. The following

antibodies were used for immunocytochemistry: Rabbit anti-neurofilament 200 (1:1000, Sigma

Aldrich, N4142-25ul), rabbit anti-neuroserpin (1:200, Abcam, ab33077), rabbit anti-synapsin 1

(1:1000, Sigma Aldrich, AB1543), mouse anti-NeuN (1:500, Abcam, ab104224), mouse anti-

MAP2 (1:500, Sigma Aldrich, MAB3418), rabbit anti-synapsin 1 (1:1000, Invitrogen, A-6442),

mouse anti-SSEA4 (1:250, Invitrogen, MC-813-70), rabbit anti-SOX2 (1:250, Invitrogen, PA1-

094), mouse anti-GFP (1:1000, MBL Life Science, M048-3), rabbit anti-neuroserpin (1:1000, a

gift from Dr. M. Elena Miranda Banos, purified polyclonal antibody) and mouse anti-neuroserpin

(1:1000, a gift from Dr. M. Elena Miranda Banos, 1A10). [63] The following day, cells were washed

three times with PBS and incubated with secondary antibodies in 10% donkey serum in PBS for

1 hour at room temperature. The following secondary antibodies were used: donkey anti-mouse

IgG (H+L) highly cross-adsorbed secondary antibody Alexa Fluor 488 (1:250, Invitrogen, A-

21202), donkey anti-mouse IgG (H+L) highly cross-adsorbed secondary antibody Alexa Fluor

594 (1:250, Invitrogen, A-21203), donkey anti-mouse IgG (H+L) highly cross-adsorbed

secondary antibody Alexa Fluor 647 (1:250, Invitrogen, A-31571), donkey anti-rabbit IgG (H+L)

highly cross-adsorbed secondary antibody Alexa Fluor 488 (1:250, Invitrogen, A-21206), donkey

anti-rabbit IgG (H+L) highly cross-adsorbed secondary antibody Alexa Fluor 594 (1:250,

Invitrogen, A-21207). Nuclei were stained with DAPI (Invitrogen; Waltham, MA), then stained

slides were mounted with fluorescence mounting medium (Dako; Glostrup, Denmark). Cells

were imaged with a Zeiss-880 inverted confocal microscope with a magnification of 40X N.A.

1.1, 1 Airy unit pinhole, half-maximal optimized Z-stack increments with 2x2 tiled images.

Western Blot

Cells were lysed using a lysis buffer containing 20 mM Tris-HCl, pH 7.5, 150 mM NaCl,

cOmplete mini EDTA-free protease inhibitor cocktail (Roche; Indianapolis, IN), PhosSTOP

phosphatase inhibitor cocktail (Roche; Indianapolis, IN), 1% Triton-X 100, in PBS. Samples

were incubated on ice for 30 minutes and vortexed every 10 minutes. Samples were centrifuged

at 20,000 RCF for 30 minutes at 4°C to remove insoluble material.

Protein concentration was determined using the Pierce BCA kit (Thermo Fisher Scientific;

Waltham, MA). 25 µg of whole cell lysate was loaded into a 7.5% mini-protein precast protein

gel (Bio-Rad; Hercules, CA). Lysate for SDS-PAGE was prepared in 4X sample buffer with 2-

mercaptoethanol (Sigma Aldrich; Saint Louis, MO) and heated at 98°C for 5 minutes. Lysate for

native-PAGE was prepared with BlueJuice Gel loading buffer (Thermo Fisher Scientific;

Waltham, MA). Gels were run at 180V for 1 hour at 4°C in either SDS-PAGE (25 mM Tris, 192

mM glycine, 1% SDS in diH2O) or native-PAGE running buffer (25 mM Tris, 192 mM glycine in

diH2O) to separate proteins. Gels were electrophoretically transferred onto Immun-Blot PVDF

membrane (Bio-Rad; Hercules, CA) overnight in transfer buffer (25 mM Tris, 190 mM glycine,

0.1% SDS, 20% methanol in diH2O) at a constant 20mA. Membranes were washed with TBST

(20 mM Tris, 150 mM NaCl, 0.1% tween 20 in diH2O) then blocked in 5% nonfat dry milk in TBST. Membranes were rotationally incubated with diluted primary antibodies in 5% nonfat dry milk in TBST at 4°C overnight. The following antibodies were used for immunoblotting: rabbit anti-neuroserpin (1:200, Abcam, ab33077), mouse anti-vinculin (1:1000, Sigma Aldrich, V9131-2ML), mouse anti-GFP (1:1000, MBL Life Science, M048-3). The following day, membranes were washed with TBST and incubated for 1 hour at room temperature with either goat anti-mouse IgG (H+L) secondary antibody conjugated to HRP (1:5000, Invitrogen, 31430) or goat anti-rabbit IgG secondary antibody conjugated to HRP (1:5000, Invitrogen, 31460) in TBST. Membranes were washed three times with TBST, and a final wash with TBS before applying Western Blotting Luminol Reagent (Santa Cruz Biotechnology; Dallas, TX) and imaging with a Bio Rad ChemiDoc imaging system.

RNA extraction and cDNA synthesis

RNA was extracted using RNeasy mini kit (Qiagen; Hilden, Germany) with on-column DNase digestion (Qiagen; Hilden, Germany) per the manufacturer's guidelines. cDNA was synthesized from RNA using the SuperScript III First-Strand cDNA Synthesis Kit (Thermo Fisher Scientific; Waltham, MA).

Quantitative PCR

Quantitative PCR was performed on cDNA synthesized as mentioned above using ssoAdvanced Universal SYBR Green Supermix reagent (Bio-Rad; Hercules, CA). For each assay, 500 nM of forward and reverse primers, and 1X final concentration of SYBR green master mix was used. Assays were performed using a StepOne Real-time PCR system (Applied Biosystems; Waltham, MA) using the following settings: 30 seconds at 95°C, followed by 40 cycles of 10 seconds at 95°C and 60 seconds at 60°C, followed by a melt curve analysis of 40

cycles at 65°C -95°C increasing in 0.5°C increments at a rate of 5 seconds per step. All qPCR

primers are listed in table 6.

Next generation sequencing

RNA was extracted using RNeasy mini kit as mentioned above. Automated polyA selected

stranded RNA-seq library prep was performed by the New York University Langone Health

Genome Technology Center. Libraries were sequenced using an Illumina NovaSeq6000 next

generation sequencer (Illumina Inc; San Diego, CA) as paired end runs. After quality control,

samples were mapped using BBTools software, Sequence Expression Analyzer (Seal) (Joint

Genome Institute; Berkeley, CA) against both mouse and human reference genomes to exclude

reads from mouse glia.[93] Reads were mapped using Spliced Transcripts Aligned to a Reference

(STAR), and counted with featureCounts.[94,95] Read counts were normalized as transcripts per

million (TPM) and plotted as $\log_2$(TPM+1).

Aggregate analysis and detection

Aggregate analysis was performed using ImageJ FIJI software. Confocal microscopy acquired

images were projected as maximum projections of Z-stack images to create a two-dimensional

image of aggregates. The NS-GFP channel was separated and thresholded for the brightest

fluorescence expression. All images from one experiment were thresholded using the same

values. Aggregates were counted using ImageJ "analyze particles" function. Particles of less

than 0.1 microns were excluded to avoid including single pixel values in the analysis. The total

number of particles in an image was used to determine aggregate count. The total area of the

particles in an image was used to determine the aggregate area.

NS-GFP aggregate detection for single clone assay was performed using the Olympus CKX53

epifluorescence microscope (Evident Corporation, Tokyo, Japan). GFP fluorescence and

brightfield images were acquired using cellSens imaging software (Evident Corporation, Tokyo,

Japan). All images were acquired under the same gain and exposure settings. Images were

processed using ImageJ FIJI software. The NS-GFP channel was thresholded for the brightest

fluorescence to detect NS-GFP aggregates. Images were categorized as containing aggregates

if any particle greater than 0.1 micron was detected.

Cell count

Confocal microscopy acquired images were used to identified and count cells using Cellpose

v2.2.2 software.[96] Images were initially processed in ImageJ to isolate the DAPI stained nuclei

channel, and project using the sum projection option to produce an image of all cells within the

field-of-view. The nuclei channels were processed in Cellpose set to nuclei mode. Cell diameter

was automatically determined using the calibration function. Flow_threshold was set to 0.4,

cellprob_threshold was set to 0.0, and stitch_threshold was set to 0.0. Image saturation was set

to automatic. The resulting cell count was used to determine the total number of cells per field.

To determine NeuN and DAPI double positive cells, images of NeuN and DAPI stained cells

were processed using ImageJ. The NeuN channel was isolated and automatically thresholded

to generate a binary mask. The mask was processed using the "Fill Holes" function, to generate

a mask representing whole NeuN positive nuclei. Next, isolated DAPI stained nuclei channels

were processed with corresponding NeuN masks using the image calculator function "multiply"

to produce a binary image with only NeuN/DAPI double positive nuclei remaining. This image

was processed in Cellpose, as mentioned above, to produce the number of NeuN/DAPI double

positive cells for normalization.

Sholl analysis

iPSC-derived neurons were cultured in geltrex coated 12 mm No.1 coverslips. Neurons were

transfected with pmaxGFP[TM] (Lonza; Morris Township, NJ) using Lipofectamine 2000 to

sparsely label neurons. 24-hours post-transfection, cells were fixed with 4% PFA in PBS. Cells

were stained with mouse anti-GFP (1:1000, Invitrogen, G10362) overnight at 4°C. The following

day, cells were labeled with donkey anti-rabbit IgG (H+L) cross-adsorbed secondary antibody

DyLight™ 488 (1:250, Invitrogen, SA5-10038). Coverslips were mounted on slides using Dako

fluorescent mounting media.

Neurite morphology analyses on transfected iPSC-derived neurons were performed on

randomly sampled neurons. Images of neurons were acquired using the Zeiss-880 confocal at

63X magnification with 1 Airy unit pinhole, Z-stack increments of 4.28 µm, and laser gain

between 700-800 for all channels. Imaging parameters were kept consistent between

inductions. Images were processed using NIH image software (ImageJ) as a maximum

projection of 2x2 tiled images. Neurites were manually traced using NIH ImageJ with the SNT

plugin and Sholl analysis was performed using Image J with the Sholl analysis plugin. To ensure

accuracy, neurons with overlapping neurites were excluded from analysis.

Electrophysiology

Whole-cell patch clamp recording was performed on iPSC-derived neurons 42-44 days after

neural induction. Glass pipettes were filled with a solution containing the following (in mM): 135

KmeSO$_3$, 5 KCl, 0.1 EGTA, 10 HEPES, 2 NaCl, 5 MgATP, 0.4 Na$_2$GTP, 10 Na$_2$PhosphoCreatine

(pH 7.3). Cells were perfused with artificial cerebrospinal fluid (ACSF) containing (in mM)- 125

NaCl, 22.5 D-glucose, 25 NaHCO$_3$, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 3 C$_3$H$_3$NaO$_3$, 1 C$_6$H$_8$O$_6$, 2 CaCl, 1

MgCl, saturated with 95% O$_2$, 5% CO$_2$ (pH 7.3). The ACSF was continuously flowing at a rate of

2-3 ml/min at room temperature. Recording electrodes were pulled from borosilicate glass

pipettes (Sutter Instrument; Novato, CA) and had a tip resistance of 3 – 5 MΩ when filled with

internal solution. All recordings were performed using a MultiClamp 700B amplifier interfaced

with a PC through a Digidata 1550 (Molecular Devices; Sunnyvale, CA). Data was acquired with

pClamp 10.7 software (Molecular Devices; Sunnyvale, CA) and analyzed with Clampfit 10.7

(Molecular Devices; Sunnyvale, CA) and Prism (Graphpad; LaJolla; CA).

eVLP delivery of ABE

eVLP delivery of ABE was performed by adding concentrated eVLP to HEK293T, iPSC, or iNs.

For HEK293T cells, cells were dissociated into a single cell suspension with 0.25%

trypsin/EDTA. 40,000 cells were plated per well of a 48-well plates. 25 µl of 100X concentrated

eVLPs were added as a reverse transduction and media was changed after 24 hours. For

iPSCs, cells were gently dissociated into a single cell suspension with Accutase treatment and

40,000 cells were plated on geltrex coated 48-well plates. The indicated volumes of eVLPs were

treated as a reverse transduction and directly added into culturing media in either

unconcentrated or concentrated form. Culturing media was changed after 24 hours. All cells

were collected for analysis without selection at 72 hours. For iNs, cells were plated as normal

onto glia coated cultureware. The indicated volumes of concentrated eVLPs were added as part

of a standard media change with the addition of polybrene (Sigma Aldrich; Saint Louis, MO).

Media was changed the following day with complete iN media with neurotrophic factors. Cells

were collected for analysis without selection after 72 hours.

Statistics

All statistical analyses were computed using Prism as described. Graphs are presented as

means ± standard error of the mean (SEM), or box and whisker plots showing 10-90th percentile

range, with sample number indicated per experiment. Unpaired Student's t tests, one-way

ANOVA with multiple comparisons tests, and two-way ANOVA with multiple comparisons tests

with Bonferroni corrections were performed to determine statistical significance at 95%

confidence interval.

Data availability statement

For original data, please contact corresponding authors chengzu.long@nyulangone.org,

jayeeta.basu@nyulangone.org, and od4@nyu.edu.

Acknowledgements

We would like to acknowledge members of the Long and Basu labs for comments and technical
assistance. We would like to thank members of the NYU Langone core facilities including
Cytometry and Cell Sorting Laboratory, Genome Technology Center, High Performance
Computing Core, Small Instrument Fleet, and Microscopy Laboratory for their technical
assistance. This work was supported by grants from the Departmental Start-Up Grant, NYU
Langone Health, Kids Connect Charitable Fund, NIH (T32GM136542), NYU Clinical and
Translational Science Institute (CTSI) Pilot Project (UL1TR001445, NIH/NCATS
(JB/CL/OD/TB)), and Cancer Center Support Grant (P30CA016086). American Epilepsy Society
Junior Investigator Award (JB), Parekh Center for Interdisciplinary Neurology (PCIN) Pilot
Research Program 2023 (JB and OD), FACES (Finding a Cure for Epilepsy and Seizures), NYU
Langone Health (CL/JB/OD/TB), NIH NINDS BRAIN INITIATIVE 1R01NS109994 (JB), NIH
NINDS 1R01NS109362-01 (JB).

Author contributions

C.T.K. and C.L. conceptualized the project. C.T.K., T.B., O.D., J.B., and C.L. acquired funding.
C.T.K. and N.M. performed experiments and analyzed results. T.B. and Q.Y. offered training,
guidance, and oversaw experiments. V.Z., D.K., M.P. provided technical support. C.T.K. and
C.L. wrote the manuscript. C.T.K., N.M., T.B., O.D., J.B., and C.L. reviewed and edited the
manuscript.

Declaration of interests

The authors declare no competing interests.

Keywords

Protein conformational disease, protein aggregation, iPSC-derived neurons, CRISPR/Cas9,
adenine base editing, engineered virus-like particles

References

1. Chiti, F., and Dobson, C.M. (2006). Protein Misfolding, Functional Amyloid, and Human
Disease.   Annual   Review   of   Biochemistry   *75*,   333–366.
https://doi.org/10.1146/annurev.biochem.75.101304.123901.

2. Soto, C., and Pritzkow, S. (2018). Protein misfolding, aggregation, and conformational
strains   in   neurodegenerative   diseases.   Nat   Neurosci   *21*,   1332–1340.
https://doi.org/10.1038/s41593-018-0235-9.

3. Stoeckli, E.T., Lemkin, P.F., Kuhn, T.B., Ruegg, M.A., Heller, M., and Sonderegger, P.
(1989). Identification of proteins secreted from axons of embryonic dorsal-root-ganglia
neurons. European Journal of Biochemistry *180*, 249–258. https://doi.org/10.1111/j.1432-
1033.1989.tb14640.x.

4. Osterwalder, T., Contartese, J., Stoeckli, E.T., Kuhn, T.B., and Sonderegger, P. (1996).
Neuroserpin, an axonally secreted serine protease inhibitor. The EMBO Journal *15*, 2944–
2953. https://doi.org/10.1002/j.1460-2075.1996.tb00657.x.

5. Miranda, E., Römisch, K., and Lomas, D.A. (2004). Mutants of Neuroserpin That Cause
Dementia Accumulate as Polymers within the Endoplasmic Reticulum*. Journal of Biological
Chemistry *279*, 28283–28291. https://doi.org/10.1074/jbc.M313166200.

6. Belorgey, D., Crowther, D.C., Mahadeva, R., and Lomas, D.A. (2002). Mutant Neuroserpin
(S49P) That Causes Familial Encephalopathy with Neuroserpin Inclusion Bodies Is a Poor
Proteinase Inhibitor and Readily Forms Polymers in Vitro *. Journal of Biological Chemistry
277, 17367–17373. https://doi.org/10.1074/jbc.M200680200.

7. López-González, I., Pérez-Mediavilla, A., Zamarbide, M., Carmona, M., Torrejón Escribano,
B., Glatzel, M., Galliciotti, G., and Ferrer, I. (2016). Limited Unfolded Protein Response and
Inflammation   in   Neuroserpinopathy.   J   Neuropathol   Exp   Neurol   *75*,   121–133.
https://doi.org/10.1093/jnen/nlv011.

8. Davis, R.L., Holohan, P.D., Shrimpton, A.E., Tatum, A.H., Daucher, J., Collins, G.H., Todd,
R., Bradshaw, C., Kent, P., Feiglin, D., et al. (1999). Familial Encephalopathy with
Neuroserpin Inclusion Bodies. The American Journal of Pathology *155*, 1901–1913.
https://doi.org/10.1016/S0002-9440(10)65510-1.

9. Davis, R.L., Shrimpton, A.E., Carrell, R.W., Lomas, D.A., Gerhard, L., Baumann, B.,
Lawrence, D.A., Yepes, M., Kim, T.S., Ghetti, B., et al. (2002). Association between
conformational mutations in neuroserpin and onset and severity of dementia. The Lancet
359, 2242–2247. https://doi.org/10.1016/S0140-6736(02)09293-0.

10. Kara, B., Sarıkaya, C.E., Bayrak, Y.E., Güneş, A.S., Güngör, M., and Yeşil, G. (2020).
Early-onset rapidly progressive myoclonic epilepsy associated with G392R likely pathogenic
variant in SERPINI1. Seizure *80*, 181–182. https://doi.org/10.1016/j.seizure.2020.06.022.

11. Ranza, E., Garcia-Tarodo, S., Varvagiannis, K., Guipponi, M., Lobrinus, J.A., Bottani, A.,
Kern, I., Kurian, M., Pittet, M.-P., Antonarakis, S.E., et al. (2017). SERPINI1 pathogenic

variants: An emerging cause of childhood-onset progressive myoclonic epilepsy. American
Journal of Medical Genetics Part A *173*, 2456–2460. https://doi.org/10.1002/ajmg.a.38317.

12. Yang, X., Fang, Z., Yan, L., He, X., Luo, H., Han, Z., Gui, J., Cheng, M., and Jiang, L.
(2022). Role of SERPINI1 pathogenic variants in familial encephalopathy with neuroserpin
inclusion bodies: A case report and literature review. Seizure *103*, 137–147.
https://doi.org/10.1016/j.seizure.2022.11.008.

13. Knupp, K., and Wirrell, E. (2014). Progressive myoclonic epilepsies: It takes a village to
make a diagnosis. Neurology *82*, 378–379.
https://doi.org/10.1212/WNL.0000000000000091.

14. Guadagno, N.A., Moriconi, C., Licursi, V., D'Acunto, E., Nisi, P.S., Carucci, N., De Jaco, A.,
Cacci, E., Negri, R., Lupo, G., et al. (2017). Neuroserpin polymers cause oxidative stress in
a neuronal model of the dementia FENIB. Neurobiology of Disease *103*, 32–44.
https://doi.org/10.1016/j.nbd.2017.03.010.

15. Levey, A.I. (2021). Progress with Treatments for Alzheimer's Disease. New England Journal
of Medicine *384*, 1762–1763. https://doi.org/10.1056/NEJMe2103722.

16. Tabrizi, S.J., Flower, M.D., Ross, C.A., and Wild, E.J. (2020). Huntington disease: new
insights into molecular pathogenesis and therapeutic opportunities. Nat Rev Neurol *16*,
529–546. https://doi.org/10.1038/s41582-020-0389-4.

17. Evers, M.M., Miniarikova, J., Juhas, S., Vallès, A., Bohuslavova, B., Juhasova, J.,
Skalnikova, H.K., Vodicka, P., Valekova, I., Brouwers, C., et al. (2018). AAV5-miHTT Gene
Therapy Demonstrates Broad Distribution and Strong Human Mutant Huntingtin Lowering in
a Huntington's Disease Minipig Model. Molecular Therapy *26*, 2163–2177.
https://doi.org/10.1016/j.ymthe.2018.06.021.

18. Tabrizi, S.J., Leavitt, B.R., Landwehrmeyer, G.B., Wild, E.J., Saft, C., Barker, R.A., Blair,
N.F., Craufurd, D., Priller, J., Rickards, H., et al. (2019). Targeting Huntingtin Expression in
Patients with Huntington's Disease. N Engl J Med *380*, 2307–2316.
https://doi.org/10.1056/NEJMoa1900907.

19. Cummings, J., Zhou, Y., Lee, G., Zhong, K., Fonseca, J., and Cheng, F. (2023). Alzheimer's
disease drug development pipeline: 2023. Alzheimer's & Dementia: Translational Research
& Clinical Interventions *9*, e12385. https://doi.org/10.1002/trc2.12385.

20. Wild, E.J., and Tabrizi, S.J. (2017). Therapies targeting DNA and RNA in Huntington's
disease. The Lancet Neurology *16*, 837–847. https://doi.org/10.1016/S1474-
4422(17)30280-6.

21. Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., and Church,
G.M. (2013). RNA-Guided Human Genome Engineering via Cas9. Science *339*, 823–826.
https://doi.org/10.1126/science.1232033.

22. Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W.,
Marraffini, L.A., et al. (2013). Multiplex Genome Engineering Using CRISPR/Cas Systems.
Science *339*, 819–823. https://doi.org/10.1126/science.1231143.

23. Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012). A
Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity.
Science *337*, 816–821. https://doi.org/10.1126/science.1225829.

24. Chapman, J.R., Taylor, M.R.G., and Boulton, S.J. (2012). Playing the End Game: DNA
Double-Strand Break Repair Pathway Choice. Molecular Cell *47*, 497–510.
https://doi.org/10.1016/j.molcel.2012.07.029.

25. Urnov, F.D., Rebar, E.J., Holmes, M.C., Zhang, H.S., and Gregory, P.D. (2010). Genome
editing with engineered zinc finger nucleases. Nat Rev Genet *11*, 636–646.
https://doi.org/10.1038/nrg2842.

26. Long, C., Amoasii, L., Mireault, A.A., McAnally, J.R., Li, H., Sanchez-Ortiz, E.,
Bhattacharyya, S., Shelton, J.M., Bassel-Duby, R., and Olson, E.N. (2016). Postnatal
genome editing partially restores dystrophin expression in a mouse model of muscular
dystrophy. Science *351*, 400–403. https://doi.org/10.1126/science.aad5725.

27. Park, H., Oh, J., Shim, G., Cho, B., Chang, Y., Kim, S., Baek, S., Kim, H., Shin, J., Choi, H.,
et al. (2019). In vivo neuronal gene editing via CRISPR–Cas9 amphiphilic nanocomplexes
alleviates deficits in mouse models of Alzheimer's disease. Nat Neurosci *22*, 524–528.
https://doi.org/10.1038/s41593-019-0352-0.

28. Yin, H., Xue, W., Chen, S., Bogorad, R.L., Benedetti, E., Grompe, M., Koteliansky, V.,
Sharp, P.A., Jacks, T., and Anderson, D.G. (2014). Genome editing with Cas9 in adult mice
corrects a disease mutation and phenotype. Nat Biotechnol *32*, 551–553.
https://doi.org/10.1038/nbt.2884.

29. Chang, C.-J., Kotini, A., Teruya-Feldstein, J., Abdel-Wahab, O., Bradley, R., and
Papapetrou, E.P. (2016). Isogenic iPSC Models of SRSF2-Mutant Myelodysplastic
Syndrome Capture Disease Phenotypes, Splicing Defects and Drug Responses. Blood *128*,
962–962. https://doi.org/10.1182/blood.V128.22.962.962.

30. Xie, Y., Ng, N.N., Safrina, O.S., Ramos, C.M., Ess, K.C., Schwartz, P.H., Smith, M.A., and
O'Dowd, D.K. (2020). Comparisons of dual isogenic human iPSC pairs identify functional
alterations directly caused by an epilepsy associated SCN1A mutation. Neurobiology of
Disease *134*, 104627. https://doi.org/10.1016/j.nbd.2019.104627.

31. Zhang, Y., Pak, C., Han, Y., Ahlenius, H., Zhang, Z., Chanda, S., Marro, S., Patzke, C.,
Acuna, C., Covy, J., et al. (2013). Rapid Single-Step Induction of Functional Neurons from
Human Pluripotent Stem Cells. Neuron *78*, 785–798.
https://doi.org/10.1016/j.neuron.2013.05.029.

32. Mao, Z., Bozzella, M., Seluanov, A., and Gorbunova, V. (2008). DNA repair by
nonhomologous end joining and homologous recombination during cell cycle in human
cells. Cell Cycle *7*, 2902–2906. https://doi.org/10.4161/cc.7.18.6679.

33. Saleh-Gohari, N., and Helleday, T. (2004). Conservative homologous recombination
preferentially repairs DNA double-strand breaks in the S phase of the cell cycle in human
cells. Nucleic Acids Research *32*, 3683–3688. https://doi.org/10.1093/nar/gkh703.

34. Lin, S., Staahl, B.T., Alla, R.K., and Doudna, J.A. (2014). Enhanced homology-directed
human genome engineering by controlled timing of CRISPR/Cas9 delivery. eLife *3*, e04766.
https://doi.org/10.7554/eLife.04766.

35. Gaudelli, N.M., Komor, A.C., Rees, H.A., Packer, M.S., Badran, A.H., Bryson, D.I., and Liu,
D.R. (2017). Programmable base editing of A•T to G•C in genomic DNA without DNA
cleavage. Nature *551*, 464–471. https://doi.org/10.1038/nature24644.

36. Raguram, A., Banskota, S., and Liu, D.R. (2022). Therapeutic in vivo delivery of gene
editing agents. Cell *185*, 2806–2827. https://doi.org/10.1016/j.cell.2022.03.045.

37. Bulcha, J.T., Wang, Y., Ma, H., Tai, P.W.L., and Gao, G. (2021). Viral vector platforms
within the gene therapy landscape. Sig Transduct Target Ther *6*, 1–24.
https://doi.org/10.1038/s41392-021-00487-6.

38. Lyu, P., Wang, L., and Lu, B. (2020). Virus-Like Particle Mediated CRISPR/Cas9 Delivery
for Efficient and Safe Genome Editing. Life (Basel) *10*, 366.
https://doi.org/10.3390/life10120366.

39. Banskota, S., Raguram, A., Suh, S., Du, S.W., Davis, J.R., Choi, E.H., Wang, X., Nielsen,
S.C., Newby, G.A., Randolph, P.B., et al. (2022). Engineered virus-like particles for efficient
in vivo delivery of therapeutic proteins. Cell *185*, 250-265.e16.
https://doi.org/10.1016/j.cell.2021.12.021.

40. Choi, J.G., Dang, Y., Abraham, S., Ma, H., Zhang, J., Guo, H., Cai, Y., Mikkelsen, J.G., Wu,
H., Shankar, P., et al. (2016). Lentivirus pre-packed with Cas9 protein for safer gene editing.
Gene Ther *23*, 627–633. https://doi.org/10.1038/gt.2016.27.

41. Nishimasu, H., Shi, X., Ishiguro, S., Gao, L., Hirano, S., Okazaki, S., Noda, T., Abudayyeh,
O.O., Gootenberg, J.S., Mori, H., et al. (2018). Engineered CRISPR-Cas9 nuclease with
expanded targeting space. Science *361*, 1259–1262.
https://doi.org/10.1126/science.aas9129.

42. Richardson, C.D., Ray, G.J., DeWitt, M.A., Curie, G.L., and Corn, J.E. (2016). Enhancing
homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using
asymmetric donor DNA. Nature Biotechnology *34*, 339–344.
https://doi.org/10.1038/nbt.3481.

43. Synthego Performance Analysis, ICE analysis. 2019. v3.0. Sythego.

44. Conant, D., Hsiau, T., Rossi, N., Oki, J., Maures, T., Waite, K., Yang, J., Joshi, S., Kelso, R.,
Holden, K., et al. (2022). Inference of CRISPR Edits from Sanger Trace Data. The CRISPR
Journal *5*, 123–130. https://doi.org/10.1089/crispr.2021.0113.

45. Fra, A., D'Acunto, E., Laffranchi, M., and Miranda, E. (2018). Cellular Models for the
Serpinopathies. In Serpins: Methods and Protocols Methods in Molecular Biology., A.
Lucas, ed. (Springer), pp. 109–121. https://doi.org/10.1007/978-1-4939-8645-3_7.

46. D'Acunto, E., Gianfrancesco, L., Serangeli, I., D'Orsi, M., Sabato, V., Guadagno, N.A.,
Bhosale, G., Caristi, S., Failla, A.V., De Jaco, A., et al. (2022). Polymerogenic neuroserpin
causes mitochondrial alterations and activates NFκB but not the UPR in a neuronal model of

neurodegeneration FENIB. Cell. Mol. Life Sci. *79*, 437. https://doi.org/10.1007/s00018-022-
04463-3.

47. Schipanski, A., Oberhauser, F., Neumann, M., Lange, S., Szalay, B., Krasemann, S., van
Leeuwen, F.W., Galliciotti, G., and Glatzel, M. (2014). The lectin OS-9 delivers mutant
neuroserpin to endoplasmic reticulum associated degradation in familial encephalopathy
with neuroserpin inclusion bodies. Neurobiology of Aging *35*, 2394–2403.
https://doi.org/10.1016/j.neurobiolaging.2014.04.002.

48. Schipanski, A., Lange, S., Segref, A., Gutschmidt, A., Lomas, D.A., Miranda, E., Schweizer,
M., Hoppe, T., and Glatzel, M. (2013). A Novel Interaction Between Aging and ER Overload
in a Protein Conformational Dementia. Genetics *193*, 865–876.
https://doi.org/10.1534/genetics.112.149088.

49. Davies, M.J., Miranda, E., Roussel, B.D., Kaufman, R.J., Marciniak, S.J., and Lomas, D.A.
(2009). Neuroserpin Polymers Activate NF-κB by a Calcium Signaling Pathway That Is
Independent of the Unfolded Protein Response. J. Biol. Chem. *284*, 18202–18209.
https://doi.org/10.1074/jbc.M109.010744.

50. Lindhout, F.W., Kooistra, R., Portegies, S., Herstel, L.J., Stucchi, R., Snoek, B.L., Altelaar,
A.M., MacGillavry, H.D., Wierenga, C.J., and Hoogenraad, C.C. (2020). Quantitative
mapping of transcriptome and proteome dynamics during polarization of human iPSC-
derived neurons. eLife *9*, e58124. https://doi.org/10.7554/eLife.58124.

51. Gunhanlar, N., Shpak, G., van der Kroeg, M., Gouty-Colomer, L.A., Munshi, S.T.,
Lendemeijer, B., Ghazvini, M., Dupont, C., Hoogendijk, W.J.G., Gribnau, J., et al. (2018). A
simplified protocol for differentiation of electrophysiologically mature neuronal networks from
human induced pluripotent stem cells. Mol Psychiatry *23*, 1336–1344.
https://doi.org/10.1038/mp.2017.56.

52. Espuny-Camacho, I., Michelsen, K.A., Gall, D., Linaro, D., Hasche, A., Bonnefont, J., Bali,
C., Orduz, D., Bilheu, A., Herpoel, A., et al. (2013). Pyramidal Neurons Derived from Human
Pluripotent Stem Cells Integrate Efficiently into Mouse Brain Circuits In Vivo. Neuron *77*,
440–456. https://doi.org/10.1016/j.neuron.2012.12.011.

53. Simkin, D., Marshall, K.A., Vanoye, C.G., Desai, R.R., Bustos, B.I., Piyevsky, B.N., Ortega,
J.A., Forrest, M., Robertson, G.L., Penzes, P., et al. (2021). Dyshomeostatic modulation of
Ca2+-activated K+ channels in a human neuronal model of KCNQ2 encephalopathy. eLife
10, e64434. https://doi.org/10.7554/eLife.64434.

54. Budrikis, Z., Costantini, G., La Porta, C.A.M., and Zapperi, S. (2014). Protein accumulation
in the endoplasmic reticulum as a non-equilibrium phase transition. Nat Commun *5*, 3620.
https://doi.org/10.1038/ncomms4620.

55. Kopach, O., Esteras, N., Wray, S., Abramov, A.Y., and Rusakov, D.A. (2021). Genetically
engineered MAPT 10+16 mutation causes pathophysiological excitability of human iPSC-
derived neurons related to 4R tau-induced dementia. Cell Death Dis *12*, 1–12.
https://doi.org/10.1038/s41419-021-04007-w.

56. Virdi, G.S., Choi, M.L., Evans, J.R., Yao, Z., Athauda, D., Strohbuecker, S., Nirujogi, R.S.,
Wernick, A.I., Pelegrina-Hidalgo, N., Leighton, C., et al. (2022). Protein aggregation and

calcium dysregulation are hallmarks of familial Parkinson's disease in midbrain dopaminergic neurons. npj Parkinsons Dis. *8*, 1–22. https://doi.org/10.1038/s41531-022-00423-7.

57. Gehrke, J.M., Cervantes, O., Clement, M.K., Wu, Y., Zeng, J., Bauer, D.E., Pinello, L., and Joung, J.K. (2018). An APOBEC3A-Cas9 base editor with minimized bystander and off-target activities. Nature Biotechnology *36*, 977–982. https://doi.org/10.1038/nbt.4199.

58. Grünewald, J., Zhou, R., Iyer, S., Lareau, C.A., Garcia, S.P., Aryee, M.J., and Joung, J.K. (2019). CRISPR DNA base editors with reduced RNA off-target and self-editing activities. Nature Biotechnology *37*, 1041–1048. https://doi.org/10.1038/s41587-019-0236-6.

59. Rees, H.A., Wilson, C., Doman, J.L., and Liu, D.R. (2019). Analysis and minimization of cellular RNA editing by DNA adenine base editors. Science Advances *5*, eaax5717. https://doi.org/10.1126/sciadv.aax5717.

60. Koblan, L.W., Doman, J.L., Wilson, C., Levy, J.M., Tay, T., Newby, G.A., Maianti, J.P., Raguram, A., and Liu, D.R. (2018). Improving cytidine and adenine base editors by expression optimization and ancestral reconstruction. Nature Biotechnology *36*, 843–846. https://doi.org/10.1038/nbt.4172.

61. Richter, M.F., Zhao, K.T., Eton, E., Lapinaite, A., Newby, G.A., Thuronyi, B.W., Wilson, C., Koblan, L.W., Zeng, J., Bauer, D.E., et al. (2020). Phage-assisted evolution of an adenine base editor with improved Cas domain compatibility and activity. Nature Biotechnology *38*, 883–891. https://doi.org/10.1038/s41587-020-0453-z.

62. Kluesner, M.G., Nedveck, D.A., Lahr, W.S., Garbe, J.R., Abrahante, J.E., Webber, B.R., and Moriarity, B.S. (2018). EditR: A Method to Quantify Base Editing from Sanger Sequencing. CRISPR J *1*, 239–250. https://doi.org/10.1089/crispr.2018.0014.

63. Miranda, E., MacLeod, I., Davies, M.J., Pérez, J., Römisch, K., Crowther, D.C., and Lomas, D.A. (2008). The intracellular accumulation of polymeric neuroserpin explains the severity of the dementia FENIB. Human Molecular Genetics *17*, 1527–1539. https://doi.org/10.1093/hmg/ddn041.

64. Hermann, M., Reumann, R., Schostak, K., Kement, D., Gelderblom, M., Bernreuther, C., Frischknecht, R., Schipanski, A., Marik, S., Krasemann, S., et al. (2020). Deficits in developmental neurogenesis and dendritic spine maturation in mice lacking the serine protease inhibitor neuroserpin. Molecular and Cellular Neuroscience *102*, 103420. https://doi.org/10.1016/j.mcn.2019.103420.

65. Borges, V.M., Lee, T.W., Christie, D.L., and Birch, N.P. (2010). Neuroserpin regulates the density of dendritic protrusions and dendritic spine shape in cultured hippocampal neurons. Journal of Neuroscience Research *88*, 2610–2617. https://doi.org/10.1002/jnr.22428.

66. Kement, D., Reumann, R., Schostak, K., Voß, H., Douceau, S., Dottermusch, M., Schweizer, M., Schlüter, H., Vivien, D., Glatzel, M., et al. (2021). Neuroserpin Is Strongly Expressed in the Developing and Adult Mouse Neocortex but Its Absence Does Not Perturb Cortical Lamination and Synaptic Proteome. Front Neuroanat *15*, 627896. https://doi.org/10.3389/fnana.2021.627896.

67. Reumann, R., Vierk, R., Zhou, L., Gries, F., Kraus, V., Mienert, J., Romswinkel, E., Morellini,
F., Ferrer, I., Nicolini, C., et al. (2017). The serine protease inhibitor neuroserpin is required
for normal synaptic plasticity and regulates learning and social behavior. Learn. Mem. *24*,
650–659. https://doi.org/10.1101/lm.045864.117.

68. Arshadi, C., Günther, U., Eddison, M., Harrington, K.I.S., and Ferreira, T.A. (2021). SNT: a
unifying toolbox for quantification of neuronal anatomy. Nat Methods *18*, 374–377.
https://doi.org/10.1038/s41592-021-01105-7.

69. Ferreira, T.A., Blackman, A.V., Oyrer, J., Jayabal, S., Chung, A.J., Watt, A.J., Sjöström,
P.J., and van Meyel, D.J. (2014). Neuronal morphometry directly from bitmap images. Nat
Methods *11*, 982–984. https://doi.org/10.1038/nmeth.3125.

70. Wang, D., Zhang, F., and Gao, G. (2020). CRISPR-Based Therapeutic Genome Editing:
Strategies and In Vivo Delivery by AAV Vectors. Cell *181*, 136–150.
https://doi.org/10.1016/j.cell.2020.03.023.

71. Kato, S., Kobayashi, K., Inoue, K., Kuramochi, M., Okada, T., Yaginuma, H., Morimoto, K.,
Shimada, T., Takada, M., and Kobayashi, K. (2011). A Lentiviral Strategy for Highly Efficient
Retrograde Gene Transfer by Pseudotyping with Fusion Envelope Glycoprotein. Human
Gene Therapy *22*, 197–206. https://doi.org/10.1089/hum.2009.179.

72. Karasev, M.M., Baloban, M., Verkhusha, V.V., and Shcherbakova, D.M. (2022). Nuclear
Localization Signals for Optimization of Genetically Encoded Tools in Neurons. Frontiers in
Cell and Developmental Biology *10*.

73. Davis, H.E., Morgan, J.R., and Yarmush, M.L. (2002). Polybrene increases retrovirus gene
transfer efficiency by enhancing receptor-independent virus adsorption on target cell
membranes. Biophysical Chemistry *97*, 159–172. https://doi.org/10.1016/S0301-
4622(02)00057-1.

74. Visentin, C., Musso, L., Broggini, L., Bonato, F., Russo, R., Moriconi, C., Bolognesi, M.,
Miranda, E., Dallavalle, S., Passarella, D., et al. (2020). Embelin as Lead Compound for
New Neuroserpin Polymerization Inhibitors. Life *10*, 111.
https://doi.org/10.3390/life10070111.

75. Saga, G., Sessa, F., Barbiroli, A., Santambrogio, C., Russo, R., Sala, M., Raccosta, S.,
Martorana, V., Caccia, S., Noto, R., et al. (2016). Embelin binds to human neuroserpin and
impairs its polymerisation. Sci Rep *6*, 18769. https://doi.org/10.1038/srep18769.

76. Penney, J., Ralvenius, W.T., and Tsai, L.-H. (2020). Modeling Alzheimer's disease with
iPSC-derived brain cells. Mol Psychiatry *25*, 148–167. https://doi.org/10.1038/s41380-019-
0468-3.

77. LaFerla, F.M., and Green, K.N. (2012). Animal Models of Alzheimer Disease. Cold Spring
Harb Perspect Med *2*, a006320. https://doi.org/10.1101/cshperspect.a006320.

78. Kroeger, H., Miranda, E., MacLeod, I., Pérez, J., Crowther, D.C., Marciniak, S.J., and
Lomas, D.A. (2009). Endoplasmic Reticulum-associated Degradation (ERAD) and
Autophagy Cooperate to Degrade Polymerogenic Mutant Serpins. J. Biol. Chem. *284*,
22793–22802. https://doi.org/10.1074/jbc.M109.027102.

79. Roussel, B.D., Newton, T.M., Malzer, E., Simecek, N., Haq, I., Thomas, S.E., Burr, M.L.,
Lehner, P.J., Crowther, D.C., Marciniak, S.J., et al. (2013). Sterol metabolism regulates
neuroserpin polymer degradation in the absence of the unfolded protein response in the
dementia FENIB. Human Molecular Genetics *22*, 4616. https://doi.org/10.1093/hmg/ddt310.

80. Galliciotti, G., Glatzel, M., Kinter, J., Kozlov, S.V., Cinelli, P., Rülicke, T., and Sonderegger,
P. (2007). Accumulation of Mutant Neuroserpin Precedes Development of Clinical
Symptoms in Familial Encephalopathy with Neuroserpin Inclusion Bodies. The American
Journal of Pathology *170*, 1305–1313. https://doi.org/10.2353/ajpath.2007.060910.

81. Takasawa, A., Kato, I., Takasawa, K., Ishii, Y., Yoshida, T., Shehata, M.H., Kawaguchi, H.,
Mohafez, O.M.M., Sasahara, M., and Hiraga, K. (2008). Mutation-, Aging-, and Gene
Dosage-dependent Accumulation of Neuroserpin (G392E) in Endoplasmic Reticula and
Lysosomes of Neurons in Transgenic Mice. J. Biol. Chem. *283*, 35606–35613.
https://doi.org/10.1074/jbc.M804125200.

82. Wong, M., and Guo, D. (2013). Dendritic spine pathology in epilepsy: Cause or
consequence?   Neuroscience   *251*,   141–150.
https://doi.org/10.1016/j.neuroscience.2012.03.048.

83. Xu, X., Miller, E.C., and Pozzo-Miller, L. (2014). Dendritic spine dysgenesis in Rett
syndrome. Frontiers in Neuroanatomy *8*.

84. Torres, M.D., Garcia, O., Tang, C., and Busciglio, J. (2018). Dendritic spine pathology and
thrombospondin-1 deficits in Down syndrome. Free Radical Biology and Medicine *114*, 10–
14. https://doi.org/10.1016/j.freeradbiomed.2017.09.025.

85. Chapleau, C.A., Calfa, G.D., Lane, M.C., Albertson, A.J., Larimore, J.L., Kudo, S.,
Armstrong, D.L., Percy, A.K., and Pozzo-Miller, L. (2009). Dendritic spine pathologies in
hippocampal pyramidal neurons from Rett syndrome brain and after expression of Rett-
associated   MECP2   mutations.   Neurobiology   of   Disease   *35*,   219–233.
https://doi.org/10.1016/j.nbd.2009.05.001.

86. Hastings, G.A., Coleman, T.A., Haudenschild, C.C., Stefansson, S., Smith, E.P., Barthlow,
R., Cherry, S., Sandkvist, M., and Lawrence, D.A. (1997). Neuroserpin, a Brain-associated
Inhibitor of Tissue Plasminogen Activator Is Localized Primarily in Neurons IMPLICATIONS
FOR THE REGULATION OF MOTOR LEARNING AND NEURONAL SURVIVAL. J. Biol.
Chem. *272*, 33062–33067. https://doi.org/10.1074/jbc.272.52.33062.

87. Krueger, S.R., Ghisu, G.-P., Cinelli, P., Gschwend, T.P., Osterwalder, T., Wolfer, D.P., and
Sonderegger, P. (1997). Expression of Neuroserpin, an Inhibitor of Tissue Plasminogen
Activator, in the Developing and Adult Nervous System of the Mouse. J. Neurosci. *17*,
8984–8996. https://doi.org/10.1523/JNEUROSCI.17-23-08984.1997.

88. Osterwalder, T., Cinelli, P., Baici, A., Pennella, A., Krueger, S.R., Schrimpf, S.P., Meins, M.,
and Sonderegger, P. (1998). The Axonally Secreted Serine Proteinase Inhibitor,
Neuroserpin, Inhibits Plasminogen Activators and Plasmin but Not Thrombin. J. Biol. Chem.
273, 2312–2321. https://doi.org/10.1074/jbc.273.4.2312.

89. Samson, A.L., Nevin, S.T., Croucher, D., Niego, B., Daniel, P.B., Weiss, T.W., Moreno, E.,
Monard, D., Lawrence, D.A., and Medcalf, R.L. (2008). Tissue-type plasminogen activator

requires a co-receptor to enhance NMDA receptor function. J Neurochem *107*, 1091–1101.
https://doi.org/10.1111/j.1471-4159.2008.05687.x.

90. Nicole, O., Docagne, F., Ali, C., Margaill, I., Carmeliet, P., MacKenzie, E.T., Vivien, D., and
Buisson, A. (2001). The proteolytic activity of tissue-plasminogen activator enhances NMDA
receptor-mediated signaling. Nat Med *7*, 59–64. https://doi.org/10.1038/83358.

91. Lebeurrier, N., Liot, G., Lopez-Atalaya, J.P., Orset, C., Fernandez-Monreal, M.,
Sonderegger, P., Ali, C., and Vivien, D. (2005). The brain-specific tissue-type plasminogen
activator inhibitor, neuroserpin, protects neurons against excitotoxicity both in vitro and in
vivo. Molecular and Cellular Neuroscience *30*, 552–558.
https://doi.org/10.1016/j.mcn.2005.09.005.

92. Kim, S., Kim, D., Cho, S.W., Kim, J., and Kim, J.-S. (2014). Highly efficient RNA-guided
genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. Genome
Res. *24*, 1012–1019. https://doi.org/10.1101/gr.171322.113.

93. BBMap (2023). SourceForge. https://sourceforge.net/projects/bbmap/.

94. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

95. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.
https://doi.org/10.1093/bioinformatics/btt656.

96. Pachitariu, M., and Stringer, C. (2022). Cellpose 2.0: how to train your own model. Nat
Methods *19*, 1634–1641. https://doi.org/10.1038/s41592-022-01663-4.

1285

Figure 1 Generation of FENIB models.

(A) FENIB disease progression. Mutant neuroserpin protein aggregates into neuroserpin inclusion bodies. **(B)** *SERPINI1* gene locus. Top: patient-specific *SERPINI1* c.1175 G>A (p.G392E) mutation located in exon 9 (red arrowhead). Middle: gRNAs for targeting wild-type *SERPINI1* (g1 in dark red and g2 in blue). PAM (grey) and DNA cleavage sites indicated with matching arrowheads. Bottom: knock-in (KI) single-stranded oligodeoxynucleotide (ssODN) sequence with patient-specific c.1175 mutation in red (red arrowhead). **(C)** Wild-type (WT) or mutant (MUT) neuroserpin fused to GFP (NS-GFP) lentivirus under the tetracycline response element (TRE) or cytomegalovirus (CMV) promoter used to generate stably expressed NS-GFP cells. Representative confocal microscopy images of HEK293T cells expressing WT NS-GFP or MUT NS-GFP under the TRE promoter. Samples imaged at day 1 (d1) and day 7 (d7) post-doxycycline induction and blasticidin selection. HEK293T cells express fluorescent NS-GFP (green) and are labeled with antibody specific for neuroserpin (all NS) (red). Scale bar, 25 μm. **(D)** Immunoblot of cell lysate from HEK293T cells expressing WT NS-GFP, MUT NS-GFP, or control. Samples were resolved using anti-neuroserpin antibody in SDS-PAGE (top), anti-vinculin antibody in SDS-PAGE as control (middle), or anti-neuroserpin antibody in native-PAGE (bottom) to detect polymers. **(E)** iPSCs transduced with lentivirus expressing rtTA and Ngn2. Ngn2 expression induced for 7 days using doxycycline with puromycin selection for 5 days. Cells replated onto mouse glia and cultured in neurobasal media containing neurotrophic factors. Representative confocal microscopy images of iPSCs and d21 neurons expressing stage-specific proteins. iPSCs are labeled with antibodies specific for stage-specific embryonic antigen 4 (SSEA4) (green) and SRY-Box transcription factor 2 (SOX2) (red). Neurons are labeled with antibodies for neuron specific microtubule associated protein 2 (MAP2) (green) and synapse protein synapsin I (SYN1) (red). Scale bar, 50 μm.

Figure 2 Early prevention of MUT NS-GFP expression is sufficient for aggregate

clearance.

(A) Administration and withdrawal of doxycycline for the aggregation and clearance of mutant

NS-GFP. **(B-D)** Early prevention of MUT NS-GFP expression in inducible NS-GFP neurons.

+Doxycycline (+DOX) cells received continuous doxycycline and expressed NS-GFP through

day 10. Withdrawal (W/D) cells received doxycycline until day 7, at which point doxycycline was

withdrawn, allowing aggregate clearance until day 10, at which point all cells were processed for

imaging on day 10. Each sample is represented as a dot, indicative of an imaging field of a

biological replicate. Each field has ~100-200 cells. The graphs depict mean ± SEM of n = 6

biological replicates. **(B)** Representative confocal microscopy images of NS-GFP aggregation in

day 7 and 10 FENIB neurons expressing WT NS-GFP, MUT NS-GFP, or control. Cells are

labeled with antibodies specific for all neuroserpin forms (all NS) (red), NeuN (white), DAPI

(blue), and express NS-GFP (green). Scale bar, 25 µm. **(C)** Quantification of NS-GFP aggregate

count normalized to DAPI+/NeuN+ cells. (****p<0.0001) All comparisons left unmarked are

statistically non-significant. **(D)** Quantification of NS-GFP aggregate area normalized to

DAPI+/NeuN+ cells. (****p<0.001) All comparisons left unmarked are statistically non-significant**.**

Figure 3 Late prevention of MUT NS-GFP expression prevents aggregation progression.

(A) Administration and withdrawal of doxycycline for the aggregation and clearance of mutant

NS-GFP. **(B-D)** Late prevention of MUT NS-GFP expression in inducible NS-GFP neurons. MUT

NS-GFP aggregation in iPSC-derived neurons at indicated timepoints**.** Day 21, day 28, and day

35 neurons had continuous expression of MUT NS-GFP. Day 28 and day 35 withdrawal neurons

had MUT NS-GFP expression withdrawn at day 21. Each sample is represented as a dot,

indicative of an imaging field. Each field has ~100-200 cells. The graphs depict mean ± SEM of

n = 12-19 biological replicates (day 21 n = 12, day 28 dox and w/d n = 9, day 35 dox n = 18, day

35 w/d n = 19). **(B)** Representative confocal microscopy image of MUT NS-GFP neurons

labeled with NeuN (red) and DAPI (blue), harboring NS-GFP aggregates (green). Scale bar, 25

μm. **(C)** Quantification of NS-GFP aggregate count. NS-GFP aggregate count is normalized to

DAPI+/NeuN+ cells. (*p<0.05, **p<0.01). All comparisons left unmarked are statistically non-

significant. **(D)** Quantification of NS-GFP aggregate area normalized to DAPI+/NeuN+ cells.

(**p<0.001, ***p<0.001, ****p<0.0001). All comparisons left unmarked are statistically non-

significant.

Figure 4 Correction of FENIB mutation in iPSC models using NG-ABE8e.

(A) Patient-specific FENIB mutation *SERPINI1* c.1175 G>A targeted with adenine base editor

(ABE). ABE editing window (green) using gRNA g3 (dark red) with NG PAM (grey). Patient-

specific c.1175 G>A mutation is located at the A5 position of the gRNA sequence. **(B)**

Quantification of ABE editing in heterozygous KI iPSCs (top) and homozygous KI iPSCs

(bottom) using Sanger sequencing and EditR. The graphs depict mean ± SEM of n = 3

replicates. All adenine positions within the gRNA sequence are listed, with the c.1175 variant at

position A5 (red arrowhead). **(C)** Quantification of deep sequencing reads of heterozygous KI

iPSCs treated with NG-ABEmax or NG-ABE8e. Arrows represent editing outcomes from the

variant GAA codon. Wild-type GGG codon (grey) encoding glycine (G; black), patient-specific

variant GAA codon (red) encoding glutamic acid (E; red), wild-type synonymous GGG codon

(blue) encoding glycine (G; black), and bystander only GAG codon (purple) encoding glutamic

acid (E; red) are depicted above the graph. **(D)** Quantification of ABE editing using NG-ABE8e

in homozygous KI and MUT NS-GFP model of FENIB. The graphs depict mean ± SEM of n = 3

replicates.

Figure 5 Correction of MUT NS-GFP in HEK293Ts allows for aggregate clearance.

(A) Quantification of NG-ABE8e editing in MUT NS-GFP HEK293T cells using Sanger

sequencing and EditR. gRNA sequence outlines targetable adenine bases, with the variant

located at A5. The graph depicts mean ± SEM of n = 3 replicates. **(B)** Representative confocal

microscopy images of ABE + g3 treated MUT NS-GFP, ABE only sham MUT NS-GFP control,

pretreatment MUT NS-GFP control, or WT NS-GFP control HEK293T cells. Cells are labeled

with antibodies specific for all neuroserpin forms (all NS) (grey), mutant neuroserpin polymers

(MUT poly.) (red), DAPI (blue) and are expressing NS-GFP (green). Scale bar, 25 µm. **(C)**

Quantification of cells per field. All comparisons are statistically non-significant. **(D-E)**

Analysis of NS-GFP aggregate clearance following ABE + g3 treatment in confocal microscopy

imaged coverslips**. (D)** Quantification of NS-GFP aggregate count normalized to cell count

(****$p<0.0001$, **$p<0.01$). All comparisons left unmarked are statistically non-significant. **(E)**

Quantification of NS-GFP aggregate area normalized to cell count (****$p<0.0001$, ***$p<0.001$,

*$p<0.05$). All comparisons left unmarked are statistically non-significant. Each sample is

represented as a dot, indicative of an imaging field. Each field has ~200-400 cells. The graphs

depict mean ± SEM of n = 3 biological replicates and n = 6 images per replicate. Pre =

pretreatment. **(F)** Quantification of NG-ABE8e editing efficiency in MUT NS-GFP single clones

(n = 26) following ABE + g3 treatment compared to bulk sorted cells using Sanger sequencing

and EditR.

Figure 6 Dendritic outgrowth is impaired in FENIB KI neurons and is partially restored

following ABE treatment.

(A) Representative confocal microscopy images of d42 neurons labeled with GFP (green) and

DAPI (blue), reconstructed neuron trace used for Sholl analysis, and representative Sholl

intersection schematic of inset marked by dashed red box. Sholl intersection schematic depicts

10 µm steps from soma shown as concentric rings (red) with intersections marked with open

dots. Scale bar, 50 µm. **(B-D)** Sholl analysis performed on day 35 FENIB KI neurons. The

graphs depict mean ± SEM of n = 9-18 biological replicates (WT = 9, HET. KI = 10, HOM. KI =

10, CORR = 18). **(B)** Quantification of the number of dendritic intersections from 0-200 µm

radial distances from the soma of day 35 FENIB KI neurons. (* are used to depict statistical

significance between WT and comparison groups. # are used to depict statistical significance

between CORR and comparison groups. *p<0.05, #p<0.05, ##p<0.01, ####p<0.0001). All

comparisons left unmarked are statistically non-significant. **(C)** Quantification of the area under

the curve for day 35 FENIB KI neurons. All comparisons left unmarked are statistically non-

significant. **(D)** Quantification of total dendrite length for day 35 FENIB KI neurons. All

comparisons left unmarked are statistically non-significant. **(E-G)** Sholl analysis performed on

day 42 FENIB KI neurons. The graphs depict mean ± SEM of n = 15-17 biological replicates

(WT = 15, HET. KI = 16, HOM. KI = 16, CORR = 17). **(E)** Quantification of the number of

intersections from 0-200 µm radial distances from soma of day 42 FENIB KI neurons. (* are

used to depict statistical significance between WT and comparison groups. # are used to depict

statistical significance between CORR and comparison groups. *p<0.05, **p<0.01, #p<0.05,

##p<0.01). All comparisons left unmarked are statistically non-significant. **(F)** Quantification of

area under the curve for day 42 FENIB KI neurons. (*p<0.05, **p<0.01). All comparisons left

unmarked are statistically non-significant. **(G)** Quantification of total dendrite length for fay 42

FENIB KI neurons. (*p<0.05). All comparisons left unmarked are statistically non-significant. WT

= wild-type, HET. KI = heterozygous knock-in, HOM. KI = homozygous knock-in, CORR = ABE

corrected.

Figure 7 eVLP editing in iPSC and iPSC-derived neurons.

(A) MMLV-gag fused NG-ABE8e for eVLPs (top), structure of ABE containing eVLP (middle),

and depiction of editing and sequencing (bottom). MMLV-gag (green) fused NG-ABE8e (pink)

with nuclear localization signals (brown), nuclear localization signal (yellow), and protease

cleavage site (blue). eVLPs generated using MMLV-gag-pol (green and blue), VSV-G or FuG-B2

glycoprotein (orange), and MMLV-gag-NG-ABE8e (green and pink) (middle). Diagram of eVLP

treated cells and subsequent editing at target locus. **(B)** Quantification of eVLP-ABE editing at

the HEK2 locus in HEK293T cells. The graphs depict mean ± SEM of n = 3 biological replicates.

(C) Quantification of eVLP-ABE editing at the HEK3 locus in HEK293T cells. The graphs depict

mean ± SEM of n = 3 biological replicates. **(D)** Quantification of eVLP-ABE correction of FENIB

mutation in *SERPINI1* c.1175 homozygous KI iPSCs. The graphs depict mean ± SEM of n = 3

biological replicates. **(E)** Quantification of eVLP ABE editing in day 7 iPSC-derived neurons

using 5 µg/ml polybrene (PB) with BP/SV40 NLS (grey) or NiV NLS (red) or 10 µg/ml PB with

BP/SV40 NLS (blue) or NiVW NLS (green). The graphs depict mean ± SEM of n = 3 biological

replicates. **(F)** Quantification of eVLP-ABE correction of FENIB mutation in d7 iPSC-derived

neurons using neuron-specific nuclear localization signals NiV W (red), MeCP2 (blue), hnRNP D

(green) or BP/SV40 (grey) control. The graphs depict mean ± SEM of n = 3 biological replicates.

(****$p<0.0001$, **$p<0.001$.) All comparisons left unmarked are statistically non-significant.

Figure 1

Figure 2

A

| | d0 | d21 | d28 | d35 | Condition |

+Doxycycline (DOX)

Withdrawal (W/D)

B

+Doxycycline | Withdrawal

d21 | d28 | d35 | d28 | d35

Composite

NS-GFP

NeuN

C

NS-GFP aggregate count (normalized to neuron count)

2.5
2.0
1.5
1.0
0.5
0.0

*
**
**

DOX: +   +   W/D   +   W/D
      d21   d28   d35

D

NS-GFP aggregate area (μm²) (normalized to neuron count)

**
**
***
****

DOX: +   +   W/D   +   W/D
      d21   d28   d35

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7