Comparative analysis of retroviral Gag-host cell interactions: focus on the nuclear interactome

Gregory S. Lambert[1a], Breanna L. Rice[1a], Rebecca J. Kaddis Maldonado[1,2], Jordan Chang[1], and Leslie J. Parent [1,2]*

[1] Department of Medicine, Penn State College of Medicine, 500 University Drive, Hershey, PA 17033, USA; glambert@pennstatehealth.psu.edu; breanna.l.rice@gmail.com; rkaddis@pennstatehealth.psu.edu; jchang3@pennstatehealth.psu.edu

[2] Department of Microbiology and Immunology, Penn State College of Medicine, 500 University Drive, Hershey, PA 17033, USA

[a] These authors contributed equally to the manuscript.
* Correspondence: lparent@psu.edu

Abstract

Retroviruses exploit a variety of host proteins to assemble and release virions from infected cells. To date, most studies that examined possible interacting partners of retroviral Gag proteins focused on host proteins that localize primarily to the cytoplasm or plasma membrane. Given the recent findings that several full-length Gag proteins localize to the nucleus, identifying the Gag-nuclear interactome has high potential for novel findings that reveal previously unknown host processes. In this study, we systematically compared nuclear factors identified in published HIV-1 proteomic studies which had used a variety of experimental approaches. In addition, to contribute to this body of knowledge, we report results from a mass spectrometry approach using affinity-tagged (His_6) HIV-1 and RSV Gag proteins mixed with nuclear extracts. Taken together, the previous studies—as well as our own—identified potential binding partners of HIV-1 and RSV Gag involved in several nuclear processes, including transcription, splicing, RNA modification, and chromatin remodeling. Although a subset of host proteins interacted with both Gag proteins, there were also unique host proteins belonging to each interactome dataset. To validate one of the novel findings, we demonstrated the interaction of RSV Gag with a member of the Mediator complex, Med26, which is required for RNA polymerase II-mediated transcription. These results provide a strong premise for future functional studies to investigate roles for these nuclear host factors that may have shared functions in the biology of both retroviruses, as well as functions specific to RSV and HIV-1, given their distinctive hosts and molecular pathology.

Keywords: retroviruses; Rous sarcoma virus; HIV-1; mass spectrometry; proteomics

1. Introduction

Retroviral replication depends upon the selection and packaging of unspliced viral RNA (USvRNA) as the genome by the Gag polyprotein, the major structural protein shared by retroviruses. Historically, it was thought that the Gag protein performed this function in the cytoplasm before trafficking to the plasma membrane where budding of virions occurs. However, there is now compelling evidence that a population of retroviral Gag proteins enter the nucleus where they may initiate selection of genomic RNA (gRNA). The strongest evidence for this paradigm shift is based on studies of the Rous sarcoma virus (RSV) Gag protein, which depends on transient nucleocytoplasmic trafficking facilitated by host transport proteins to ensure efficient gRNA packaging [1-6]. This requirement for nuclear trafficking was demonstrated by a mutant of Gag that bypassed the nucleus, resulting in decreased gRNA packaging which was increased with restoration of nuclear localization [7]. RSV Gag co-opts the host karyopherins importin α/β, importin-11, and transportin-3 (TNPO3) to enter the nucleus [1-3,6], and nuclear egress is mediated by binding of a hydrophobic nuclear export signal (NES) in the p10 domain to the CRM1-RanGTP nuclear export complex [3-5]. A more recent study demonstrated that RSV Gag binds to newly synthesized USvRNA in discrete ribonucleoprotein complexes (RNPs) in the nucleus, and these Gag-vRNA RNPs have been observed trafficking across the nuclear envelope into the cytoplasm [8].

A population of human immunodeficiency virus type-1 (HIV-1) Gag, like that of RSV Gag, also undergoes nuclear localization [9-13]. HIV-1 Gag forms focal RNP complexes with nascent USvRNA in the nucleus, and traffics to the major viral RNA transcription site in T cells reactivated from latency [9]. Nuclear localization of HIV-1 Gag occurs in a concentration-independent manner shortly after Gag synthesis begins, and Gag colocalizes with transcriptionally-active euchromatin near the nuclear periphery [10]. The function(s) of these nuclear RNPs has yet to be thoroughly investigated, although these data demonstrate that both RSV and HIV-1 Gag proteins traffic to transcription sites and associate with their cognate USvRNAs. Although one possibility is that Gag-USvRNA binding in the nucleus could initiate the genomic RNA packaging process, nuclear localization of Gag could also influence other

important cellular processes, such as regulation of viral or host transcription, RNA modification

or processing, splicing, chromatin remodeling, or RNA export.

In addition to RSV and HIV-1, the Gag proteins of other retroviruses, including murine

leukemia virus (MLV), prototype foamy virus (PFV), feline immunodeficiency virus (FIV),

mouse mammary tumor virus (MMTV), and Mason-Pfizer monkey virus (MPMV) also have

been shown to localize to the nucleus [11,14-27]. For example, the PFV Gag protein is in-

volved in proviral integration through its interaction with chromatin [27]. The finding that nu-

clear trafficking of Gag is a feature conserved among many retroviruses raises the likelihood

that nuclear-localized Gag proteins participate in functions important for virus replication.

There are also Gag cleavage products that undergo nuclear localization, including the

nucleocapsid (NC) proteins of HIV-1, RSV, MLV, and MMTV, which localize to the nucleolus

[11,19,22,28], and the p12 protein of MLV, which binds to chromatin and influences proviral

integration [23,29].

It is well known that retroviruses exploit a variety of host pathways during replication, but

previous investigation of host factors that bind to Gag have focused on factors localized to the

cytoplasm and plasma membrane. However, the nuclear localization of retroviral Gag proteins

raises important questions concerning their functions, which can be informed by identifying

nuclear host partners. To gain further insight into what nuclear processes Gag could be in-

fluencing, we comprehensively analyzed and systematically compared six previously pub-

lished HIV-1 proteomic studies performed by other laboratories, which used various experi-

mental approaches to identify novel host proteins that interact with HIV-1 Gag. A variety of

techniques are represented in this analysis, including affinity purifications of GFP-tagged Gag,

tandem affinity purification of Gag, and BirA* Gag complexes [30-35]. To complement those

datasets, we performed affinity-tagged purification of both RSV and HIV-1 Gag, and identified

nuclear interacting partners using mass spectrometry. To further explore one of the novel hits,

we utilized immunoprecipitation and quantitative imaging approaches to validate the interac-

tion of RSV Gag with Mediator complex subunit 26 (Med26), a critical component of the

transcriptional Mediator complex, which is exploited by other viruses and endogenous

retroelements [36-43]. Together, published studies combined with our results suggest that

Gag proteins may interface with host nuclear factors to facilitate genomic RNA selection

and/or influence cellular processes, including gene expression, RNA processing, splicing,

nucleic acid metabolism, and/or chromatin modification.

2. Materials and Methods

Cells, Plasmids, and Purified Proteins

DF1 chicken embryo fibroblast cells, HeLa human cervical cancer cells, and QT6 quail

fibroblast cells were maintained as described [11,44,45]. The RSV Gag expression constructs

pGag.ΔPR (referred to herein as RSV Gag), pGag.L219A.ΔPR (referred to herein as RSV

Gag.L219A), pGag.ΔNC, and pGag.ΔPR-GFP (referred to herein as RSV Gag-GFP) [4,11,46]

and plasmids encoding for *Escherichia coli* (*E. coli*) expression of His-tagged RSV Gag

(pET28.TEV-Gag.3h) and HIV Gag (pET28a.WT.HIV.Gag.Δp6) were previously described

[47,48].

Subcellular Fractionation

QT6 cells were transfected with untagged RSV Gag constructs using the calcium phos-

phate method [49]. Sixteen hours later, the medium was changed to fresh primary growth

medium (PGM) and the cells were allowed to recover for 24 hours. All subsequent steps were

performed on ice or at 4°C with cold buffers unless otherwise stated. Cells were fractionated

using the method described in [50] with some minor modifications, as below. Cells were re-

moved from the plates using trypsin and then washed in cold PBS. The cell pellet was re-

suspended in sucrose buffer (10 mM HEPES pH 7.9, 10 mM KCl, 2 mM magnesium acetate, 3

mM $CaCl_2$, 340 mM sucrose, 1 mM DTT, 100 µg/ml phenylmethanesulfonyl fluoride (PMSF), 1

µg/ml pepstatin, and Roche Complete Protease Inhibitor Cocktail) and incubated on ice for 10

minutes. IGEPAL Nonidet P-40 was added to the final concentration of 0.5% and cells were

vortexed on high for 15 seconds, and then spun for 10 minutes at 3,500 x g at 4°C. The su-

pernatant (cytoplasm fraction) was collected, and the pelleted nuclei were resuspended in

nucleoplasm extraction buffer (50 mM HEPES pH 7.9, 150 mM potassium acetate, 1.5 mM MgCl$_2$, 0.1% IGEPAL Nonidet P-40, 1 mM DTT, 100 µg/ml PMSF, 1 µg/ml pepstatin, and Roche Complete Protease Inhibitor Cocktail) and transferred to a Dounce homogenizer and homogenized with 20 slow strokes. The homogenates were checked under a light microscope for completion of nuclear lysis, then transferred to a new tube and rotated at 4°C for 20 minutes. The lysates were spun at 16,000 x g for 10 minutes at 4°C. The supernatant (nucleoplasmic fraction) was collected and the remaining chromatin-containing pellet was resuspended in nuclease incubation buffer (50 mM HEPES pH 7.9, 10 mM NaCl, 1.5 mM MgCl$_2$, 1 mM DTT, 100 µg/ml PMSF, 1 µg/ml pepstatin, and Roche Complete Protease In-hibitor Cocktail) with 100 U/ml of OmniCleave nuclease (Epicentre) for 10 minutes at 37°C. NaCl was added to a final concentration of 150 mM and the lysates were incubated on ice for 20 minutes and spun for 10 minutes at 16,000 x g at 4°C. The supernatant (low-salt chromatin fraction) was collected and the pellet was resuspended in chromatin extraction buffer (50 mM HEPES pH 7.9, 500 mM NaCl, 1.5 mM MgCl$_2$, 0.1% Triton X-100, 1 mM DTT, 100 µg/ml PMSF, 1 µg/ml pepstatin, and Roche Complete Protease Inhibitor Cocktail), incubated for 20 minutes on ice, spun for 10 minutes at 16,000 x g at 4°C, and the supernatant (high-salt chromatin fraction) was collected.

Western Blot Analysis of Subcellular Fractions

Proteins from the subcellular fractions were analyzed via SDS-PAGE. Aliquots of the fractions were heated to 90°C in 4X SDS-PAGE sample buffer (250 mM Tris-HCl, pH 6.8, 40% glycerol, 0.4% bromophenol blue, 8% SDS, and 8% β-mercaptoethanol) for 10 minutes prior to loading on a 10% SDS-PAGE gel and analyzed by western blot. Proteins were detected using antibodies against RSV Gag [51], Calnexin (Enzo Life Sciences ADI-SPA-865), Med4 (Abcam ab129170), RCC1 (Abcam ab54600), Histone H2B (Abcam ab52484), GAPDH (UBP Bio Y1040), and the appropriate HRP-conjugated secondary antibodies (Invitrogen).

Signal densities of the protein bands on the antibody-stained membranes were analyzed using Bio-Rad Image Lab Software on a ChemiDoc MP system. Rectangles were drawn

around each band, as well as a blank background region, using the volume tools feature to quantify the signal intensity of each band. The background subtraction method was set to local, and the blank region that was highlighted by a rectangle was labelled as the background volume. The volumes report table was exported to Microsoft Excel. For each band corresponding to the Gag signal, the adjusted volumes for each fraction were added together to calculate the total adjusted volume. Then the percentages of each fraction were calculated by subtracting the fraction's adjusted volume from the total adjusted volume. Averages and standard deviations were calculated for each fraction for each Gag protein from three separate experiments.

Purified RSV Gag and HIV Gag Pulldowns

Lysate Preparation: DF1 and HeLa cells were fractionated using the NE-PER Nuclear and Cytoplasmic Extraction kit (ThermoFisher Scientific). All steps and buffers used were performed on ice or at 4°C unless otherwise stated. Cells were lysed in CERI buffer containing the Complete Protease Inhibitor Cocktail (Roche). Cells were vortexed on the highest setting for 15 seconds and incubated on ice for 10 minutes. Ice-cold CERII buffer was added and cells were vortexed on high for 5 seconds then centrifuged for 5 minutes at 16,000 x *g* in a microcentrifuge. The supernatant was collected (cytoplasmic fraction), and the pelleted nuclei were resuspended in ice-cold NER buffer with protease inhibitor cocktail added. The nuclei were vortexed on high for 15 seconds and incubated on ice for 10 minutes, then vortexed for 15 seconds every 10 minutes for a total of 40 minutes. The lysed nuclei were centrifuged at 16,000 x *g* for 10 minutes. The supernatant (nuclear fractionation) was diluted to 14 ml with Buffer A (25 mM Tris-HCl pH 8.0, 200 mM NaCl, 2 mM 2-Mercaptoethanol (BME), and protease inhibitor cocktail). The nuclear fraction was concentrated to ~1 ml in a 3 kD MWCO Amicon column and then was diluted to 14 ml and concentrated once more to ~1.2 ml.

Nickel Affinity Purifications: Three reactions were performed using 6 μg of RSV H6.Gag.3h or HIV-1 WT.Gag.Δp6.H6, and a no protein control for DF1 and HeLa nuclear lysates, respectively. The proteins and no protein control were incubated with pre-washed

nickel beads for 1 hour at 4°C with rotation. The beads were then washed three times in Wash

Buffer (300 mM NaCl, 50 mM NaH_2PO_4, pH 8.0), followed by incubation with 500 μg of nuclear

extract for 2 hours at 4°C with rotation. The beads were washed again three times in Wash

Buffer, and bound proteins were eluted from the beads using Wash Buffer + 300 mM imid-

azole for 15 minutes while rotating at 4°C. The eluates were buffer exchanged into water using

Zeba Spin Desalting Columns (ThermoFisher Scientific) and 20 ug of each sample was used

for mass spectrometry analysis.

Sample Preparation for Mass Spectrometry: The samples were prepared and processed

at the Mass Spectrometry and Proteomics Core Research Facility at Penn State College of

Medicine using an ABSciex 5600 TripleTOF. In a final volume of 100 μl, the samples were

incubated in 50 mM NH_4HCO_3, pH 8.0, 10% v/v acetronitrile, and 0.1 μg trypsin for at least 3

hours at 48°C. To evaporate off the NH_4HCO_3 and acetronitrile, samples were dried down

using a SpeedVac, and then resuspended in 200 μl H_2O with vortexing. The drying was re-

peated 3X total, but the final resuspension volume was 10 μl. To each sample, a $1/9^{th}$ volume

of 1% formic acid was added.

Mass Spectrometry

The following mass spectrometry workflows were performed two separate times and data

from both instances were combined to create the set of interactors presented herein.

2D-LC Separations: SCX (strong cation-exchange) separations were performed on a

passivated Waters 600E HPLC system, using a 4.6 X 250 mm PolySULFOETHYL

Aspartamide column (PolyLC, Columbia, MD) at a flow rate of 1 ml/min. The gradient was

100% Buffer A (10 mM ammonium formate, pH 2.7, in 20% acetonitrile/80% water) (0-22

minutes following sample injection), 0%→40% Buffer B (666 mM ammonium formate, pH 2.7,

in 20% acetonitrile/80% water) (16-48 min), 40%→100% Buffer B (48-49 min), isocratic 100%

Buffer B (49-56 min), then at 56 min switched back to 100% Buffer A to re-equilibrate for the

next injection. One milliliter fractions were collected and were dried down then resuspended in

9 µl of 2% (v/v) acetonitrile, 0.1% (v/v) formic acid, and were filtered prior to reverse phase

C18 nanoflow-LC separation.

Mass Spectrometry Analysis: Each SCX fraction was analyzed following a calibration run

using trypsin-digested β-Gal as a calibrant, then a blank run using the ABSciex 5600

TripleTOF. MS Spectra were then acquired from each sample using the newly updated default

calibration, using a 60-minute gradient from an Eksigent NanoLC-Ultra-2D Plus and Eksigent

cHiPLC Nanoflex through a 200 µm x 0.5 mm Chrom XP C18-CL 3 µm 120 Å Trap Column

and elution through a 75 µm x 15 cm Chrom XP C18-CL 3 µm 120 Å Nano cHiPLC Column.

Protein Identification and Analysis: Protein identification and quantitation were performed

using the Paragon algorithm as implemented in ProteinPilot 5.0 software (ProteinPilot 5.0,

which contains the Paragon Algorithm 5.0.0.0, build 4632 from ABI/MDS-Sciex) [52]. Spectra

were searched against *Homo sapien* or *Gallus gallus* RefSeq subsets (plus 389 common

contaminants) of the NCBInr database concatenated with a reversed "decoy" version of itself.

For the ProteinPilot analyses, the preset Thorough Identification Search settings were used,

and identifications needed to have a ProteinPilot Unused Score > 1.3 (>95% confidence in-

terval) to be accepted. In addition, the only protein identifications (IDs) accepted were required

to have a "Local False Discovery Rate" estimation of no higher than 5%, as calculated from

the slope of the accumulated Decoy database hits by the PSPEP (Proteomics System Per-

formance Evaluation Pipeline) [53]. Proteins that were labelled as contaminants or reversed

were removed from the analysis.

Analysis of Proteomics

The Database for Annotation, Visualization, and Integrated Discovery (DAVID, version 6.8)

[54,55] was used to assign each protein to its cellular compartment(s) and biological process

categories. Proteins were organized by their gene name for entries into DAVID and the *Homo*

sapiens species database was used. Data presented in the tables were generated using the

Gene Ontology GOTERM_BP_ALL to categorize proteins by their biological function, and

GOTERM_CC_ALL to first identify the proteins present in the nucleus. Categories with a

p-value of ≤ 0.05, as determined by modified Fisher's Exact Test, were considered statistically overrepresented, and any redundant categories (same p-value and proteins) were removed.

The Bioinformatics and Evolutionary Genomics online comparison tool was used to generate the Venn diagram (http://bioinformatics.psb.ugent.be/webtools/Venn/). Ingenuity Pathway Analysis (IPA) (QIAGEN Inc., https://www.qiagenbioinformatics.com/products/ingenuitypathway-analysis) was performed to categorize the functions of the identified proteins. Core Analysis was performed on the gene IDs that could be mapped by IPA, as some gene IDs were not recognized by IPA, on each separate proteomic list. Under the Core Analysis, Expression analysis was selected; direct and indirect relationships were examined. No endogenous chemicals were included in the analysis. The filters that were used included: all molecule types and data sources; confidence = experimentally observed; species = human only; no tissues or cell lines or mutations were included. Only examined categories associated with molecular and cellular functions, as out-lined by [56]. Additionally, protein interaction maps were generated using STRING consortium (https://string-db.org/) to visualize clusters of protein-protein interactions.

Generation of RC.V8-Infected QT6 Nuclear Lysates

QT6 cells were infected for 4 hours with cell culture medium obtained from a separate culture of QT6 cells transfected with pRC.V8. Cells were fractionated using the method de-scribed in [50] with minor modifications, as described below. Cells were trypsinized, pelleted at low speed, and washed in cold PBS. The cell pellet was resuspended in lysis buffer (10 mM HEPES pH 7.9, 10 mM KCl, 0.1 mM EDTA, 0.3% Nonidet P-40, 1 mM DTT, 100 µg/ml phenylmethanesulfonyl fluoride (PMSF), 1 µg/ml pepstatin, 100U/ml Omnicleave (Epicentre), and Roche Complete Protease Inhibitor Cocktail) and incubated on ice for 5 minutes. Cells were then spun for 5 minutes at 3,000 rpm at 4°C to pellet nuclei, and the supernatant (cyto-plasmic fraction) was collected. The pelleted nuclei were washed once with lysis buffer, then resuspended in nuclear extract buffer (20 mM HEPES pH 7.9, 10 mM NaCl, 1 mM DTT, 100 µg/ml PMSF, 1 µg/ml pepstatin, 100U/ml Omnicleave (Epicentre), and Roche Complete

Protease Inhibitor Cocktail) and incubated at 37°C in a water bath for 10 minutes. Nuclear lysate was placed on ice, and solution was brought to 400 mM NaCl and 1 mM EDTA, followed by vortexing on high for 15 seconds and 20 minutes of rotation at 4°C. Debris was pelleted at 13,000 rpm for 10 minutes at 4°C, and supernatant was transferred to a fresh tube (nuclear fraction). Protein concentration in lysates was determined by Bradford assay.

RSV Gag-Med26 Co-immunoprecipitation

RC.V8-infected nuclear lysates (500 µg) were pre-incubated for 2 hours with α-RSV-CA antibody (mouse α-RSV CA.A11, gift from Neil Christensen, Penn State College of Medicine) in low salt NET2 buffer (50 mM Tris pH 7.4, 150 mM NaCl, 0.05% Nonidet P-40; [57]) at 4°C with rotation. With 1 hour remaining, Pierce™ Protein G Magnetic Beads (60 µl of a 50% slurry per reaction) were washed 4 times with high salt NET2 buffer (50 mM Tris pH 7.4, 400 mM NaCl, 0.05% Nonidet P-40; [57]), then blocked with 5% w/v BSA in low salt NET2 buffer at 4°C with rotation. At the end of the 2 hours, blocking buffer was removed and beads were re-suspended in low salt NET2 buffer. An equal amount (~30 µl) was added to each reaction, and tubes were rotated at 4°C overnight.

After overnight incubation, buffer was removed and beads were washed 4 times with high salt NET2 buffer. Bound proteins were then eluted by boiling beads in 50 µl of 1X SDS-PAGE buffer for 10 minutes at 100°C. Beads were pelleted at 13,000 rpm for 5 minutes, and su-pernatant was taken for analysis.

Samples were run on 10% SDS-PAGE gels, transferred to PVDF, blocked for 30 minutes with 5% Milk/0.1% TBS-Tween, and then incubated with primary antibody (rabbit α-Med26, Proteintech, 21043-1-AP) in 0.5% Milk/0.1% TBS-Tween at 4°C with rocking overnight. Membranes were washed 3 times for 5 minutes with 0.1% TBS-Tween, then incubated with secondary antibody (goat α-rabbit-HRP, Sigma A0545) for 1 hour at room temperature. Washes were repeated, and membranes incubated with ECL 2 for 5 minutes. Western blots were imaged using a BioRad ChemiDoc MP imager. Blots were stripped and reprobed for

RSV Gag using rabbit α-RSV-W [gift from John Wills, Penn State College of Medicine [51] and

secondary antibody (goat α-rabbit-HRP, Sigma A0545)].

Confocal Imaging:

QT6 cells were plated at a density of 0.5 x 10^6 cells/well in 6-well tissue culture dishes

containing #1.5 coverslips and were allowed to settle overnight. The following afternoon, wells

were transfected with 500 ng of RSV Gag-GFP [11] and 125 ng of FLAG-tagged Med26 (a gift

from   Joan   Conaway   and   Ronald   Conaway   [Addgene   plasmid   #15367;

http://n2t.net/addgene:15367; RRID:Addgene_15367)[58]] expression vectors. The following

morning, cells were washed 2X quickly with warm PBS, and fixed in 3.7% formaldehyde in

PBS for 10 minutes at RT. Slides were then washed 3X with PBS (5 minutes per wash),

permeabilized for 15 minutes in 0.25% Triton X-100/PBS, washed 3X, and blocked for 30

minutes at RT in 10% BSA/PBS. Relevant slides were incubated for 2 hours with mouse

α-FLAG-M2 antibody (Sigma, F1804) in 3% BSA/PBS in a humid chamber at 37°C followed by

3X washes. Secondary antibody staining with donkey α-mouse-AlexaFluor647 (AF647) was

carried out for 2 hours in 3% BSA/PBS in a humid chamber at 37°C, followed by 3X washes.

Slides were DAPI stained for 1 minute at RT, mounted with ProLong™ Diamond (Life Tech-

nologies), and cured for 24 hours. Slides were imaged on a Leica AOBS SP8 confocal mi-

croscope with a 63x/1.4 oil objective, with pinhole set to 1 airy unit and frame averaging set to

four. FLAG-Med26-AF647 was excited using a white light laser (WLL) tuned to 647 nm at 2%

laser power, and emission was detected via hybrid detector. RSV Gag-GFP was excited by

WLL tuned to 488 nm at 3% laser power and detected via hybrid detector. DAPI was excited

using a 405 nm diode laser at 8% power and detected using a photomultiplier tube.

Colocalization between RSV Gag and Med26 was assessed using Imaris image analysis

software (Oxford Instruments). Briefly, images were Gaussian filtered and surfaces were

generated around cell nuclei using the DAPI channel as reference. Masks were created for

RSV Gag and Med26 signal within nuclear surfaces, and colocalization of this signal was

assessed using the Imaris colocalization tool. Manders' Overlap Coefficients were exported

from Imaris and data was assessed for outliers by Grubbs' test with an α = 0.05 using

GraphPad Prism (GraphPad Software, Inc.). Statistical analysis and generation of **Fig 4C** was

also done with this software. A total of 17 M1 (Med26∩Gag) and 18 M2 (Gag∩Med26) indi-

vidual data points were plotted. Representative images (**Fig 4B**) and video (**Video S1**) were

created using Imaris software.

3. Results

3.1. Subcellular localization

Prior experiments using microscopy revealed that both RSV and HIV-1 Gag proteins

localize to the perichromatin compartment of the nucleus where they associate with their

cognate USvRNAs at active transcription sites [3,9,11]. In addition, HIV-1 Gag associates

preferentially with marks associated with transcriptionally-active euchromatin compared to

heterochromatin [10]. To further define the nuclear subcompartments where RSV Gag is lo-

calized, cells expressing wild-type or mutant Gag proteins were separated into cytoplasmic,

nucleoplasmic, and chromatin-associated protein fractions (**Fig 1**). The first chromatin fraction

was extracted using a NaCl concentration of 150 mM, which removes proteins that are loosely

associated with chromatin (i.e., the euchromatin fraction). In the second chromatin fraction, a

higher salt concentration was used (500 mM NaCl) along with detergent, to remove proteins

that are more tightly bound to chromatin [50] (i.e., the heterochromatin fraction).

The signals of the protein bands were quantified to yield the relative ratio of Gag protein in

each fraction, demonstrating that RSV Gag was present in the cytoplasm (42.5 ± 2.6%) and

the nucleoplasm (39.1 ± 1.6%) of cells (**Fig 1A**). Interestingly, RSV Gag was also present in

both chromatin-associated protein fractions (euchromatin, 12.6 ± 4.5%; heterochromatin, 5.8

± 1.6%). Examination of the nuclear-restricted mutant Gag.L219A, which contains a point

mutation in the p10 NES, demonstrated decreased signal in the cytoplasm (20.3 ± 0.8%) and

increased signal in both the nucleoplasm (46.7 ± 6.2%) and heterochromatin-associated

fractions (16.7 ± 2.8%), compared to wild type Gag. Analysis of mutant Gag.ΔNC, which lacks

the NC domain required for genomic RNA packaging, demonstrated an increase in cyto-

plasmic Gag compared to wild type (66.8 ± 6.6%), with little to no Gag.ΔNC in either of the chromatin fractions. Controls for the fractions included Calnexin (endoplasmic reticulum/cytoplasm), Mediator subunit 4 (Med4; nucleoplasm and euchromatin), and histone 2B (chromatin fractions). We have observed similar localization of HIV-1 Gag to all four subcellular compartments, as reported [9].

3.2. Affinity purification and proteomic analysis

We next set out to identify potential nuclear interacting partners of RSV and HIV-1 Gag to provide clues about their possible role(s) in the nucleus. Recombinant His-tagged RSV and HIV-1 Gag proteins purified from *E. coli* were incubated with nuclear lysates from DF1 chicken fibroblast cells or HeLa cells, respectively. A beads-only control was also performed using DF1 lysates and HeLa lysates incubated with nickel beads. The affinity purifications were performed twice for both Gag proteins, as well as the beads-only controls. Proteins identified using mass spectrometry were combined into a single list for each Gag protein and analyzed using DAVID and Ingenuity Pathway Analysis, as described in Materials and Methods. Proteins identified from the beads-only purifications were removed from the lists of RSV and HIV-1 affinity-tagged purifications. Proteins that had an unused score, as defined by ProteinPilot, of less than 1.3 were removed, along with common contaminants (see Supplemental files S1, S2, S3, and S4 for raw data).

DAVID was used to assign each protein from the two final Gag proteome lists to their cellular compartment(s) and biological process categories [16,19]. We used the functional annotation tool to determine the most relevant gene ontology (GO) terms associated with the proteomics lists. The order of the GO terms that were enriched in the Gag proteome lists are dependent upon the p-values for each GO term. The smaller the p-value, the more that particular GO term was enriched. **Table S1** shows the results of using the DAVID analysis software to analyze the list of proteins obtained from the RSV Gag affinity purification using DF1 nuclear lysates, and the top 10 biological processes' GO terms are displayed. Only nuclear proteins, as determined by DAVID, were analyzed to determine the enriched biological

functions, and **Table S2** shows the top 10 results. Included in the results were generic GO terms such as: GO:0034641~cellular nitrogen compound metabolic process, GO:1901360~organic cyclic compound metabolic process, and GO:0006807~nitrogen compound metabolic process. There were also more specific GO terms that were enriched, including GO:0010467~gene expression and GO:0006396~RNA processing.

Next, the potential binding partners of HIV-1 Gag isolated from HeLa nuclear lysates were examined. The terms GO:0006396~RNA processing and GO:0010467~gene expression were among the top 10 GO biological process terms identified after DAVID analysis (**Table S3**). When only nuclear proteins were examined, the top 10 GO terms again included GO:0010467~gene expression and GO:0006396~RNA processing (**Table S4**).

The proteins identified for each RSV and HIV-1 Gag affinity purification were compared, and we found there were 57 proteins that overlapped from 317 total proteins from the RSV list and 436 total proteins from the HIV-1 list (**Fig 2**). **Table 1** contains the names and functions of the 57 proteins in common between the RSV and HIV-1 Gag. The functions of the 57 proteins are varied, encompassing cell cycle progression, RNA methylation, RNA processing and splicing, cytoskeletal regulation, mRNA nuclear export, transcription, DNA repair, and chromatin modification.

Table 1. Names and functions of the 57 cellular proteins identified in both the RSV and HIV-1 Gag protein interactomes.

Symbol	Description	Function (summarized from Genecards or NCBI)
ABLIM1	Actin Binding LIM Protein 1	A LIM zinc-binding domain-containing protein that binds to actin filaments. Mediates interactions between actin and cytoplasmic targets.
ACTN1	Actinin Alpha 1	A bundling F-actin cross-linking protein thought to anchor actin to a variety of intracellular structures.
ATXN2	Ataxin 2	Involved in EGFR trafficking, acting as negative regulator of endocytic EGFR internalization at the plasma membrane.

		modulates mTOR signals, modifying ribosomal translation and mitochondrial function.
BCL7B	BAF Chromatin Remodeling Complex Subunit BCL7B	Positive regulator of apoptosis. Negatively regulates the expression of Wnt signaling components CTNNB1 and HMGA1. Involved in cell cycle progression, maintenance of the nuclear structure and stem cell differentiation.
CBLL1	Cbl Proto-Oncogene Like 1	E3 ubiquitin-protein ligase that mediates ubiquitination of several tyrosine-phosphorylated Src substrates. Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that has a role in the efficiency of mRNA splicing and RNA processing.
CDC42BPB	CDC42 Binding Protein Kinase Beta	Serine/threonine-protein kinase which is an important downstream effector of CDC42 and plays a role in the regulation of cytoskeleton reorganization and cell migration.
CHTOP	Chromatin Target of PRMT1	Plays an important role in the ligand-dependent activation of estrogen receptor target genes. Binds to 5-hydroxymethylcytosine (5hmC) and associates with the methylosome complex. The CHTOP-methylosome complex associated with 5hmC is recruited to selective sites on the chromosome, where it methylates H4R3 and activates the transcription of genes involved in glioblastomagenesis. Required for effective mRNA nuclear export and is a component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA.
CYFIP1	Cytoplasmic FMR1 Interacting Protein 1	Component of the CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression. Regulates formation of membrane ruffles and lamellipodia. Plays a role in axon outgrowth. Part of the WAVE complex that regulates actin filament reorganization via its interaction with the

		Arp2/3 complex.
EIF4G2	Eukaryotic Translation Initiation Factor 4 Gamma 2	Plays a role in the switch from cap-dependent to IRES-mediated translation during mitosis, apoptosis and viral infection.
FAM120A	Family with Sequence Similarity 120A	Critical component of the oxidative stress-induced survival signaling. Activates src family kinases and acts as a scaffolding protein enabling src family kinases to phosphorylate and activate PI3-kinase. Binds RNA and promotes the secretion of IGF-II.
FOXP1	Forkhead Box P1	Essential transcriptional regulator of B-cell development. Involved in regulation of cardiac muscle cell proliferation. Involved in the columnar organization of spinal motor neurons. Represses transcription of various pro-apoptotic genes and cooperates with NF-kappa B-signaling in promoting B-cell expansion by inhibition of caspase-dependent apoptosis. Involved in endothelial cell proliferation, tube formation and migration indicative for a role in angiogenesis. Involved in transcriptional regulation in embryonic stem cells (ESCs). Stimulates expression of transcription factors that are required for pluripotency and decreases expression of differentiation-associated genes.
HNRNPH3	heterogeneous nuclear ribonucleoprotein H3	Involved in the splicing process and participates in early heat shock-induced splicing arrest.
IGF2BP3	Insulin Like Growth Factor 2 MRNA Binding Protein 3	RNA-binding factor that may recruit target transcripts to cytoplasmic protein-RNA complexes (mRNPs) for mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation.
ING2	Inhibitor of Growth Family Member 2	Seems to be involved in p53/TP53 activation and p53/TP53-dependent apoptotic pathways, probably by enhancing acetylation of p53/TP53. Component of a mSin3A-like corepressor complex, which is probably involved in

		deacetylation of nucleosomal histones.
IQGAP1	IQ Motif Containing GTPase Activating Protein 1	Plays a crucial role in regulating the dynamics and assembly of the actin cytoskeleton.
KDM2A	Lysine Demethylase 2A	Histone demethylase that specifically demethylates 'Lys-36' of histone H3, thereby playing a central role in histone code. Required to maintain the heterochromatic state. Associates with centromeres and represses transcription of small non-coding RNAs that are encoded by the clusters of satellite repeats at the centromere. Required to sustain centromeric integrity and genomic stability, particularly during mitosis.
KIAA1033 (WASHC4)	WASH Complex Subunit 4	Acts as a component of the WASH core complex that functions as a nucleation-promoting factor (NPF) at the surface of endosomes, where it recruits and activates the Arp2/3 complex to induce actin polymerization, playing a key role in the fission of tubules that serve as transport intermediates during endosome sorting.
KLC1	Kinesin Light Chain 1	Kinesin is a microtubule-associated force-producing protein that may play a role in organelle transport. The light chain may function in coupling of cargo to the heavy chain or in the modulation of its ATPase activity.
KLC2	Kinesin Light Chain 2	See above.
LEO1	LEO1 Homolog, Paf1/RNA Polymerase II Complex Component	Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'- phosphorylated forms and is involved in transcriptional elongation.
MAP2	Microtubule Associated Protein 2	The exact function of MAP2 is unknown but MAPs may stabilize the microtubules against depolymerization.
MAPRE2	Microtubule Associated Protein	May be involved in microtubule

	RP/EB Family Member 2	polymerization, and spindle function by stabilizing microtubules and anchoring them at centrosomes.
MED26	Mediator Complex Subunit 26	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery.
NDUFA9	NADH:Ubiquinone Oxidoreductase Subunit A9	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I). Complex I functions in the transfer of electrons from NADH to the respiratory chain.
OGT	O-Linked N-Acetylglucosamine (GlcNAc) Transferase	Catalyzes the transfer of a single N-acetylglucosamine from UDP-GlcNAc to a serine or threonine residue in cytoplasmic and nuclear proteins resulting in their modification with a beta-linked N-acetylglucosamine (O-GlcNAc). Glycosylates a large and diverse number of proteins and can regulate their cellular processes via cross-talk between glycosylation and phosphorylation or by affecting proteolytic processing. Component of a THAP1/THAP3-HCFC1-OGT complex that is required for the regulation of the transcriptional activity of RRM1. Plays a key role in chromatin structure by mediating O-GlcNAcylation of 'Ser-112' of histone H2B: recruited to CpG-rich transcription start sites of active genes via its interaction with TET proteins.
ORC5	Origin Recognition Complex Subunit 5	Component of the origin recognition complex (ORC) that binds origins of replication. ORC is required to assemble the pre-replication complex necessary to initiate DNA replication.
PDCD11	Programmed Cell Death 11	Essential for the generation of mature 18S rRNA. Directly interacts with U3 snoRNA. Involved in the biogenesis of rRNA.
PDLIM5	PDZ And LIM Domain 5	May play an important role in the heart development by scaffolding PKC to the Z-disk region. May play a role in the regulation of cardiomyocyte expansion.

		Contributes to the regulation of dendritic spine morphogenesis in neurons.
PLOD2	Procollagen-Lysine,2-Oxoglutarate 5-Dioxygenase 2	Forms hydroxylysine residues in -Xaa-Lys-Gly- sequences in collagens. These hydroxylysines serve as sites of attachment for carbohydrate units and are essential for the stability of the intermolecular collagen cross-links.
POLR2B	RNA Polymerase II Subunit B	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Second largest component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Proposed to contribute to the polymerase catalytic activity and forms the polymerase active center together with the largest subunit.
PPIB	Peptidylprolyl Isomerase B	PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding.
PRPF40A	Pre-MRNA Processing Factor 40 Homolog A	Binds to WASL/N-WASP and suppresses its translocation from the nucleus to the cytoplasm, thereby inhibiting its cytoplasmic function. Plays a role in the regulation of cell morphology and cytoskeletal organization.
PTBP1	Polypyrimidine Tract Binding Protein 1	Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. May promote RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA.
RAD21	RAD21 Cohesin Complex Component	A member of the cohesin complex, involved in sister chromatid cohesion from the time of DNA replication in S phase to their segregation in mitosis, a function that is essential for proper chromosome segregation, post-replicative DNA repair, and the prevention of inappropriate recombination between repetitive regions.
RBM33	RNA Binding Motif Protein 33	*Could not find a clear function

		associated with the protein.
RFC1	Replication Factor C Subunit 1	The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins PCNA and activator 1. This subunit binds to the primer-template junction. Can bind single- or double-stranded DNA. 5' phosphate residue is required for binding of the N-terminal DNA-binding domain to duplex DNA, suggesting a role in recognition of non-primer template DNA structures during replication and/or repair.
RFC5	Replication Factor C Subunit 5	See above.
SAFB	Scaffold Attachment Factor B	Binds to scaffold/matrix attachment region (S/MAR) DNA and forms a molecular assembly point to allow the formation of a 'transcriptosomal' complex (consisting of SR proteins and RNA polymerase II) coupling transcription and RNA processing. Can function as an estrogen receptor corepressor and can also bind to the HSP27 promoter and decrease its transcription. Can inhibit cell proliferation.
SAFB2	Scaffold Attachment Factor B2	See above with S/MAR DNA. Is involved in cell cycle regulation, apoptosis, differentiation, the stress response, and regulation of immune genes.
SCAF8	SR-Related CTD Associated Factor 8	Anti-terminator protein required to prevent early mRNA termination during transcription. Mechanistically, associates with the phosphorylated C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit (POLR2A), and subsequently binds nascent RNA upstream of early polyadenylation sites to prevent premature mRNA transcript cleavage and polyadenylation. Independently of SCAF4, also acts as a positive regulator of transcript elongation.
SEPT7	Septin 7	Filament-forming cytoskeletal GTPase. Required for normal organization of the actin cytoskeleton. Required for normal progress through mitosis.

SERBP1	SERPINE1 MRNA Binding Protein 1	May play a role in the regulation of mRNA stability. Seems to play a role in PML-nuclear bodies formation
SLC25A3	Solute Carrier Family 25 Member 3	Transport of phosphate groups from the cytosol to the mitochondrial matrix.
SMAD2	SMAD Family Member 2	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases.
SMC1A	Structural Maintenance of Chromosomes 1A	Involved in chromosome cohesion during cell cycle and in DNA repair. Involved in DNA repair via its interaction with BRCA1 and its related phosphorylation by ATM, or via its phosphorylation by ATR.
SMC4	Structural Maintenance of Chromosomes 4	Central component of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condense chromosomes.
SURF4	Surfeit 4	May play a role in the maintenance of the architecture of the endoplasmic reticulum-Golgi intermediate compartment and of the Golgi.
THOC3	THO Complex 3	Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not with unspliced pre-mRNA.
TJP1	Tight Junction Protein 1	TJP1, TJP2, and TJP3 are closely related scaffolding proteins that link tight junction (TJ) transmembrane proteins such as claudins, junctional adhesion molecules, and occludin to the actin cytoskeleton.
TPM1	Tropomyosin 1	Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.
TPM3	Tropomyosin 3	Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in

		association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.
TRA2B	Transformer 2 Beta Homolog	Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing. Can either activate or suppress exon inclusion. Binds to pre-mRNA.
TRIP12	Thyroid Hormone Receptor Interactor 12	E3 ubiquitin-protein ligase involved in ubiquitin fusion degradation pathway and regulation of DNA repair. Acts as a key regulator of DNA damage.
U2AF1	U2 Small Nuclear RNA Auxiliary Factor 1	Plays a critical role in both constitutive and enhancer-dependent splicing by mediating protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. Recruits U2 snRNP to the branch point. Directly mediates interactions between U2AF2 and proteins bound to the enhancers.
VWA9 (INTS14)	Integrator Complex Subunit 14	Probable component of the Integrator (INT) complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing.
ZEB1	Zinc Finger E-Box Binding Homeobox 1	Acts as a transcriptional repressor.
ZFC3H1	Zinc Finger C3H1-Type Containing	Subunit of the trimeric poly(A) tail exosome targeting (PAXT) complex, a complex that directs a subset of long and polyadenylated poly(A) RNAs for exosomal degradation.

400
401

3.3. Comparative proteomic analysis of previously published HIV-1 Gag mass spectrometry experiments.

Mass spectrometry experiments focusing on HIV-1 Gag cellular binding partners have been previously reported by other laboratories; however, these investigators did not focus on the nuclear proteins identified in these experiments. Engeland *et al.,* 2011 [31] performed five

independent affinity-tagged purification experiments to identify cellular proteins that interact with HIV-1 Gag. The techniques used consisted of a tandem affinity purification (TAP) tag for a C-terminally tagged Gag, GFP-TRAP A beads, and GFP microbeads for Gags with GFP fused either internally to the MA domain or the C-terminus of Gag. Each of these Gag constructs were transfected into 293T cells and whole cell lysates were used for the affinity pulldowns. The authors found 31 proteins that were identified in at least 3 of the experiments, and of these 31 proteins, 24 localize to the nucleus. Using DAVID to analyze these 24 proteins, GO:0006396~RNA processing and GO:0010467~gene expression were the top two biological functions identified (**Table S5**). In 2014, Engeland and colleagues [30] examined the HIV-1 Gag interactome again and found 944 proteins that met their inclusion criteria. Using DAVID to analyze these 944 proteins, 186 were found to be nuclear. The authors performed a GO enrichment analysis using DAVID to determine the most enriched biological processes and found nuclear processes including GO:0006396~RNA processing, GO:0008380~RNA splicing, and GO:0006334~nucleosome assembly. We performed a GO analysis using solely the 186 nuclear proteins identified and found similar results (**Table S6**).

In Jäger *et al.,* 2012 [32], the authors identified host proteins that interact with each of the HIV-1 polyproteins, processed proteins, and accessory proteins, in a systematic and quantitative manner. A purification strategy was used that consisted of appending two Strep tags and three FLAG tags at the C-terminal end of each HIV-1 protein, which were expressed separately in HEK293 or Jurkat cells. When examining their raw data, we found that 1,134 unique proteins were identified from affinity purifications using full-length Gag plus the proteolytic cleavage products of Gag (MA, CA, NC, and p6). Of these interacting proteins, 180 were identified as being nuclear by DAVID analysis. Limiting our GO analysis using DAVID to these nuclear proteins, GO:0006396~RNA processing was the top biological function term identified, followed by protein targeting functions and RNA metabolism (**Table S7**).

Ritchie *et al.,* 2015 [35] provided further information on the possible protein interactors of HIV-1 Gag using the *E. coli* biotin ligase BirA* to permit identification of protein partners in close proximity to the bait protein [59-61]. The BirA* tag was inserted within the MA domain of

Gag, and the Gag-BirA* construct was expressed in cells. The authors found 53 proteins after their exclusion criteria eliminated nonspecific interactors, and from these 53 proteins, we identified 17 nuclear proteins by DAVID analysis. **Table S8** shows DAVID analysis of these 17 proteins, identifying the top biological categories of GO:0098609~cell-cell adhesion, GO:0090304~nucleic acid metabolic process, and GO:0010608~posttranscriptional regulation of gene expression.

Le Sage *et al.,* 2015 [33] also identified potential host factors that interact with HIV-1 Gag. Similar to Ritchie *et al.*, they used the BirA* tagging system, except the tag was placed at the N-terminus of Gag. They found a total of 42 proteins, of which 19 were nuclear. When these 19 nuclear proteins alone were analyzed by DAVID, the top hits included protein targeting and RNA processing and metabolism (**Table S9**).

Li *et al.,* 2016 [34] examined binding partners of the MA domain of HIV-1 Gag by inserting a Strep tag at the C-terminus of MA and collecting MA-interacting complexes after HIV-1 infection. There were 97 proteins identified that met their inclusion criteria and were not present in the lysate-only control. Of these proteins, 63 were nuclear. When only the nuclear proteins were further analyzed by DAVID, the top categories were ER targeting and viral transcription/gene expression (**Table S10**).

There is one other publication that performed a tandem immunoprecipitation to identify HIV-1 Gag binding partners [62] that was not included in the comparative analysis presented here due to the small number of proteins identified. Gao *et al.,* identified 12 individual proteins and each of these were present in at least one of the previously discussed publications or found in the experiment we presented earlier in this report. They identified numerous ribosomal proteins, HNRNP proteins, and other nuclear proteins including histone proteins, EF1-α, nucleolin, B23, Nopp34, and SNRPD3.

To provide a comprehensive view of the set of proteins identified in multiple studies, proteins identified in the previously published laboratories' reports were compared to the list of proteins found in our HIV-1 Gag affinity purifications. **Table 2** shows the list of proteins identified in our HIV-1 Gag pulldown as well as at least one of the publications discussed above.

463 **Table 2** also indicates whether each protein was previously demonstrated to have a role in

464 HIV-1 replication based on the HIV-1 human protein interaction dataset [63-65]. A total of 129

465 proteins identified in our mass spectrometry analysis were also present in at least one of the

466 prior publications.

Table 2. Proteins identified in the present HIV-1 Gag affinity purification and found in at least one of the previously published HIV-1 Gag proteomic publications.

Symbol	Description	Cell Compartment[+]	Involved in HIV replication[#]	No. of publications	Ritchie	Engeland (2014)	Engeland (2011)	Li	Le Sage	Jäger
MTDH	metadherin	N/ER	Y	5	*	*	*		*	*
NAT10	N-acetyltransferase 10; RNA cytidine acetyltransferase	N	Y	4		*	*		*	*
ZC3HAV1	zinc finger CCCH-type containing, antiviral 1	N/C	Y	4	*	*	*			*
ABCF1	ATP binding cassette subfamily F member 1	C/N	Y	3		*	*			*
CCT8	chaperonin containing TCP1 subunit 8; T-complex protein 1 subunit theta	Ck/C	Y	3	*			*		*
CDC5L	cell division cycle 5 like	C/N	Y	3		*	*			*
HNRNPM	heterogeneous nuclear ribonucleoprotein M	N	Y	3	*	*				*
MAP4	microtubule associated protein 4	Ck	Y	3	*	*				
MOV10	RISC complex RNA helicase	N/C	Y	3		*	*			*
NUFIP2	FMR1 interacting protein 2; Nuclear fragile X mental retardation-interacting protein 2	N/C	N	3	*	*				*
PLRG1	pleiotropic regulator 1	N	N	3		*			*	*
PRPF3	pre-mRNA processing factor 3; U4/U6 small nuclear ribonucleoprotein Prp3	N	N	3		*	*			*
PRPF4	pre-mRNA processing factor 4; U4/U6	N	N	3		*	*			*

	small nuclear ribonucleoprotein Prp4								
RBM14	RNA binding motif protein 14	N/C	Y	3		*	*		*
RPL37A	60S ribosomal protein L37a	C/S/N	Y	3		*		*	*
SRPK1	SRSF protein kinase 1	N/C/ER	Y	3		*	*		*
SRSF3	serine/arginine rich splicing factor 3	N/C	Y	3		*	*		*
SRSF7	serine/arginine rich splicing factor 7	N/C	Y	3		*	*		*
TMPO	Thymopoietin; Lamina-associated polypeptide 2, isoform alpha	N	Y	3	*	*			*
TOP1	DNA topoisomerase I	N	Y	3		*	*		*
ALYREF	Aly/REF export factor; THO complex subunit 4	N/C	Y	2		*			*
ATAD3B	ATPase family, AAA domain containing 3B	M	Y	2		*			*
C7orf50	chromosome 7 open reading frame 50	N	N	2		*			*
CHTOP	chromatin target of PRMT1	N	N	2		*			*
DAP3	death associated protein 3; 28S ribosomal protein S29, mitochondrial	M	Y	2		*			*
DDX24	DEAD-box helicase 24	N/Me	Y	2		*			*
DDX27	DEAD-box helicase 27	N	Y	2		*			*
DDX47	DEAD-box helicase 47	N	Y	2		*			*
DHX30	DExH-box helicase 30	M/C	Y	2		*			*
DIMT1	DIM1 dimethyladenosine transferase 1 homolog	N	Y	2		*			*

DKC1	dyskerin pseudouridine synthase 1; H/ACA ribonucleoprotein complex subunit	N/C	Y	2	*	*
EIF2S1	eukaryotic translation initiation factor 2 subunit alpha	C	Y	2	*	*
ELAVL1	ELAV like RNA binding protein 1	N/C	Y	2	*	*
EXOSC2	exosome component 2	N/C	Y	2	*	*
FAM120A	family with sequence similarity 120A; Constitutive coactivator of PPAR-gamma-like protein 1	PM/C	N	2	*	*
FBL	Fibrillarin; rRNA 2'-O-methyltransferase	N	N	2	*	*
FXR1	FMR1 autosomal homolog 1; Fragile X mental retardation syndrome-related protein 1	C	Y	2	*	*
G3BP2	G3BP stress granule assembly factor 2; Ras GTPase-activating protein-binding protein 2	C	Y	2	*	*
GRSF1	G-rich RNA sequence binding factor 1	M	Y	2	*	*
GRWD1	glutamate rich WD repeat containing 1	N	N	2	*	*
GTF3C2	general transcription factor IIIC subunit 2	N	Y	2	*	*
H1FX	H1 histone family member X	N	Y	2	*	*
HIST1H1E	histone cluster 1 H1 family member e	N	Y	2	*	*
HNRNPDL	heterogeneous nuclear ribonucleoprotein D like	N/C	Y	2	*	*
HNRNPH3	heterogeneous nuclear ribonucleoprotein H3	N	Y	2	*	*

Symbol	Name	Localization				
HNRNPR	heterogeneous nuclear ribonucleoprotein R	ER/N/C	Y	2		*	*
HNRNPUL1	heterogeneous nuclear ribonucleoprotein U like 1	N	Y	2		*	*
IARS	isoleucyl-tRNA synthetase	C	Y	2		*	*
IGF2BP1	insulin like growth factor 2 mRNA binding protein 1	N/C	Y	2		*	*
IGF2BP3	insulin like growth factor 2 mRNA binding protein 3	N/C	Y	2		*	*
ILF2	interleukin enhancer binding factor 2	N/C	Y	2		*	*
ILF3	interleukin enhancer binding factor 3	N/C	Y	2		*	*
LSM12	LSM12 homolog	C/N	Y	2		*	*
MKI67	marker of proliferation Ki-67	N	Y	2	*		*
MRPS9	mitochondrial ribosomal protein S9	M	Y	2		*	*
MRPS23	mitochondrial ribosomal protein S23	M	Y	2		*	*
MRPS27	mitochondrial ribosomal protein S27	M/C	N	2		*	*
MRPS31	mitochondrial ribosomal protein S31	M	N	2		*	*
MRPS34	mitochondrial ribosomal protein S34	M	N	2		*	*
NOP2	NOP2 nucleolar protein; Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase	N	N	2		*	*
NOP16	NOP16 nucleolar protein	N	N	2		*	*
PES1	pescadillo ribosomal biogenesis factor 1	N	N	2		*	*
PDCD11	programmed cell death 11	N	Y	2		*	*

Symbol	Description							
PRRC2C	proline rich coiled-coil 2C	C	Y	2		*	*	
PTCD3	pentatricopeptide repeat domain 3	M	Y	2		*		*
PURA	purine rich element binding protein A	N	Y	2		*		*
QARS	glutaminyl-tRNA synthetase	C	Y	2		*		*
RBM10	RNA binding motif protein 10	N	Y	2		*		*
RFC1	replication factor C subunit 1	N	Y	2		*		*
RFC5	replication factor C subunit 5	N	Y	2		*		*
RPL5	60S ribosomal protein L5	C/N	Y	2		*		*
RPL15	60S ribosomal protein L15	Me	N	2		*		
RPS24	40S ribosomal protein S24	C/ER/N/Me	Y	2		*		*
RPS28	40S ribosomal protein S28	C/ER	Y	2		*		*
RSL1D1	ribosomal L1 domain containing 1	N	Y	2		*		*
SERBP1	SERPINE1 mRNA binding protein 1	N/C	N	2	*			*
SNRPA	U1 small nuclear ribonucleoprotein polypeptide A	N	Y	2		*		*
SPTAN1	spectrin alpha, non-erythrocytic 1	Ck	Y	2		*		*
STRBP	spermatid perinuclear RNA binding protein	C	Y	2		*		*
TRA2B	transformer 2 beta homolog	N	Y	2		*		*
UPF1	RNA helicase and ATPase	N/C	Y	2		*		*
WDR3	WD repeat domain 3	N	Y	2		*		*
XRN1	5'-3' exoribonuclease 1	C	Y	2	*			*

Gene	Description						
ABLIM1	actin binding LIM protein 1	Ck/C	N	1		*	
ACTN4	actinin alpha 4	Ck/C/N	Y	1			*
AFG3L2	AFG3 like matrix AAA peptidase subunit 2	M	Y	1			*
AHNAK	Neuroblast differentiation-associated protein	N	Y	1	*		
ANKRD17	ankyrin repeat domain 17	N/C	Y	1			*
ASUN (INTS13)	asunder, spermatogenesis regulator; integrator complex subunit 13	N/C	N	1		*	
BAG2	BCL2 associated athanogene 2	Ck/C	N	1			*
BAZ1A	bromodomain adjacent to zinc finger domain 1A	N	Y	1			*
BAZ1B	bromodomain adjacent to zinc finger domain 1B	N	Y	1			*
BSG	basigin	PM/ER	Y	1			*
BUB3	mitotic checkpoint protein	N	N	1			*
BYSL	bystin like	N/C	N	1			*
C1QBP	complement C1q binding protein	N/S/PM/M/C	Y	1			*
CCAR2	cell cycle and apoptosis regulator 2	N/C	Y	1			*
CCDC86	coiled-coil domain containing 86	N	N	1			*
CCT4	chaperonin containing TCP1 subunit 4	Ck/C	Y	1			*
CCT5	chaperonin containing TCP1 subunit 5	Ck/C	Y	1			*
CCT6A	chaperonin containing TCP1 subunit 6A	C	Y	1			

Gene	Description	Localization					
CCT7	chaperonin containing TCP1 subunit 7	C	Y	1			*
CD3EAP	CD3e molecule associated protein; DNA-directed RNA polymerase I subunit RPA34	N	N	1			*
CDK12	cyclin dependent kinase 12	N	N	1			*
CLTC	clathrin heavy chain	Ck/C	Y	1			*
CORO1C	coronin 1C	PM/Ck/Me	N	1			*
CTNNB1	catenin beta 1	Ck/PM/N/C	Y	1	*		
DDX20	DEAD-box helicase 20	N/C	Y	1	*		
DDX31	DEAD-box helicase 31	N	N	1			*
DDX46	DEAD-box helicase 46	N/Me	Y	1			*
DNAJA3	DnaJ heat shock protein family (Hsp40) member A3	C/M/PM	N	1		*	
DNMT1	DNA methyltransferase 1	N	Y	1			*
ECPAS (KIAA0368)	Ecm29 Proteasome Adapter and Scaffold	ER/N/Ck	N	1	*		
EEF1B2	eukaryotic translation elongation factor 1 beta 2	C	N	1			*
EXOSC4	exosome component 4	C/N	N	1			*
FAU	ubiquitin like and ribosomal protein S30 fusion	S/N/C	Y	1			*
FIP1L1	factor interacting with PAPOLA and CPSF1	N	Y	1			*
GAPDH	glyceraldehyde-3-phosphate dehydrogenase	C/Ck/N/Me	Y	1			*

Gene	Description	Localization	Y/N	Num			
GLYR1	glyoxylate reductase 1 homolog	N	N	1			*
GPATCH4	G-patch domain containing 4	N/C	Y	1	*		
HIST1H1A	histone cluster 1 H1 family member a	N	Y	1	*		
HIST2H2BF	histone cluster 2 H2B family member f	N	Y	1	*		
HNRNPUL2	heterogeneous nuclear ribonucleoprotein U like 2	N	N	1			*
HP1BP3	heterochromatin protein 1 binding protein 3	N	N	1	*		
INIP	INTS3 and NABP interacting protein; SOSS complex subunit C	N	N	1	*		
INTS9	integrator complex subunit 9	N	N	1		*	
ITPR1	inositol 1,4,5-trisphosphate receptor type 1	ER/S	Y	1	*		
KIF4A	Chromosome-associated kinesin family member 4A	Ck/C/N	Y	1	*		
KIF23	kinesin family member 23	N	Y	1			*
KLC2	kinesin light chain 2	Ck/Me	N	1	*		
KRT2	keratin 2	Ck/C/S/N/PM	Y	1			*
KRT16	keratin 16	Ck/C/E/N	Y	1			*
KRT19	keratin 19	Ck/C/E/PM	Y	1			*
LARP4	La ribonucleoprotein domain family member 4	C	Y	1			*
LARP7	La ribonucleoprotein domain family member 7	N	Y	1			*

Gene	Description						
MAPK1	mitogen-activated protein kinase 1	Ck/C/N	Y	1			*
MBD2	methyl-CpG binding domain protein 2	N	Y	1			*
MED26	mediator complex subunit 26	N	Y	1		*	
MEPCE	7SK snRNA methylphosphate capping enzyme	N	Y	1			*
MINA (RIOX2)	MYC induced nuclear antigen; Ribosomal oxygenase 2	N	N	1			*
MRPL10	39S mitochondrial ribosomal protein L10	M	Y	1			*
MRPL53	Mitochondrial 39S ribosomal protein L53	M	Y	1			*
MRPS14	mitochondrial 28S ribosomal protein S14	M	N	1			*
MRPS22	mitochondrial 28S ribosomal protein S22	M	N	1		*	
NOC4L	nucleolar complex associated 4 homolog	N	N	1			*
PABPN1	poly(A) binding protein nuclear 1	N/C	Y	1			*
PAXBP1	PAX3 and PAX7 binding protein 1	N	Y	1			*
PBRM1	polybromo 1	N	Y	1			*
PCBP1	poly(rC) binding protein 1	C/N	Y	1			*
PPIH	peptidylprolyl isomerase H	N/C	Y	1			*
PRRC2B	proline rich coiled-coil 2B	N/C	N	1	*		
PSMC3	proteasome 26S subunit, ATPase 3	N/C	Y	1		*	

Gene	Description	Loc							
PTBP1	polypyrimidine tract binding protein 1	N	Y	1					*
PTRF (CAVIN1)	polymerase I and transcript release factor; Caveolae-associated protein 1	C/N/ER/M/PM	N	1					*
PUM1	pumilio RNA binding family member 1	C	N	1					*
RBM7	RNA binding motif protein 7	N	Y	1			*		
RBM26	RNA binding motif protein 26	N	Y	1	*				
RCC1	regulator of chromosome condensation 1	N/C	Y	1		*			
RNF213	E3 ubiquitin-protein ligase; ring finger protein 213	C	Y	1		*			
RRP9	U3 small nucleolar RNA-interacting protein 2	N	Y	1					*
RSBN1L	round spermatid basic protein 1 like; Lysine-specific demethylase	N	Y	1				*	
SDF2L1	stromal cell derived factor 2 like 1	ER	Y	1					*
SEC24B	SEC24 homolog B, COPII coat complex component	C/ER	N	1					*
SLC25A3	solute carrier family 25, member 3	M	Y	1					*
SLC25A4	solute carrier family 25 member 4; ADP/ATP translocase 1	M/Me	Y	1					*
SLC25A11	solute carrier family 25 member 11; Mitochondrial 2-oxoglutarate/malate carrier protein	M	Y	1					*
SMARCA4	SWI/SNF related, matrix associated, actin dependent regulator of chromatin; Transcription activator BRG1	N	Y	1					*

Symbol	Description					
SMARCC2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin subfamily c member 2	N	Y	1	*	
SMC4	structural maintenance of chromosomes 4	C/N	Y	1	*	
SMN2	survival of motor neuron 2, centromeric	N/C	N	1	*	
SNRPD2	small nuclear ribonucleoprotein D2 polypeptide	C/N	Y	1		*
SPIN1	spindlin 1	N	Y	1		*
SPOUT1	SPOUT domain containing methyltransferase 1	Ck	Y	1		*
SRP9	signal recognition particle 9	C	Y	1		*
SUPT16H	SPT16 homolog, facilitates chromatin remodeling subunit	N	Y	1	*	
TAB1	TGF-beta activated kinase 1 (MAP3K7) binding protein 1	C/Me/N	N	1		*
TCEB1 (ELOC)	transcription elongation factor B subunit 1; Elongin-C	N	Y	1		*
TCP1	t-complex 1	C/Ck	Y	1		*
TPM1	tropomyosin 1 (alpha)	Ck	Y	1		*
TPM3	tropomyosin 3	Ck	Y	1		*
TRIP12	thyroid hormone receptor interactor 12; E3 ubiquitin-protein ligase	N	N	1		*
TRRAP	transformation/transcription domain associated protein	N	Y	1	*	

Gene	Description	Localization						
TUBB6	tubulin beta 6 class V	Ck	Y	1		*		
U2AF1	U2 small nuclear RNA auxiliary factor 1; Splicing factor U2AF 35 kDa	N	Y	1				*
UBAP2L	ubiquitin associated protein 2 like	C	Y	1	*			
URB1 (NPA1)	Nucleolar pre-ribosomal-associated protein 1	N	Y	1				*
UTP20	UTP20, small subunit processsome component	N	N	1				*
WDR5	WD repeat domain 5	N	Y	1				*
WDR82	WD repeat domain 82	N	Y	1		*		
ZC3H4	zinc finger CCCH-type containing 4	C/N	N	1				*

69 + - C = cytoplasm, N = nucleus, M = mitochondria, PM = plasma membrane, Me = membrane, S = secreted, ER = endoplasmic reticulum, Ck = cytoskeleton;

70 determined from UniProtKB (uniprot.org), or genecards.org if localization not available on Uniprot.org; order determines abundance.

71 # - Summarized from HIV-1, human protein interaction dataset [63-65].

These overlapping proteins were analyzed using Ingenuity Pathway Analysis (IPA) (**Fig**

3), and the interacting proteins found in this report and at least one of the previously published

lists were categorized based on their functions. **Figure 3** shows the distribution of the proteins

among 13 basic categories identified by Lippé [56]. Proteins may belong to more than one

functional category, which results in the number of proteins present in each category ex-

ceeding the total number of proteins. To address this issue, the number of proteins in each

category was divided by the total number of proteins to yield a percentage for each of the

categories represented in the protein list analyzed. **Figure 3** displays the relative values of the

Ingenuity Pathway Analysis rather than the raw data.

3.4. Closer examination of Gag interactions with host proteins involved in transcription and

splicing.

Numerous Gag-interacting proteins were found to be involved in RNA polymerase II

transcription (GO term: 0006366 transcription from RNA polymerase II promoter) and splicing

(GO term: 0008380 RNA splicing). We were interested in these two GO terms because our

previous work demonstrated that RSV and HIV-1 Gag localize to the perichromatin space and

associate with newly-transcribed USvRNA [8,9], suggesting that they may interface with

transcription and splicing processes. In addition, in Rice *et al.*, we demonstrated that RSV

Gag.L219A colocalizes with the splicing factors SF2 and SC35, and has similar mobility and

dynamics as proteins residing in splicing speckles [66]. Maldonado *et al.* subsequently

showed that RSV Gag forms discrete nuclear foci and interacts with USvRNA at active tran-

scription sites [8]. Taken together with the proteomic data presented here, it is reasonable to

propose that RSV Gag interacts with cellular proteins involved in transcription and

co-transcriptional processes such as splicing and RNA processing, at or near transcription

sites. **Tables 3** and **4** list the proteins involved in transcription and splicing, respectively, for

the RSV pulldowns. Similarly, Tuffy *et al.* and Chang *et al.* [9,10] both demonstrated that HIV-1

Gag localizes to transcriptionally active regions in HeLa cells and T cells reactivated from

latency. Given these findings, it is feasible that HIV-1 Gag interacts with host nuclear factors

involved in transcription, RNA processing, and chromatin remodeling. **Tables 5** and **6** list the proteins in these categories identified in our HIV-1 pulldowns and indicate whether each protein was identified in any of the other publications. Of note, proteins involved in processes such as splicing are included in the DAVID category of transcription interactome [67,68] because transcription and splicing have been shown to be linked and occur simultaneously [69-72].

3.5 Validation of RSV Gag-Med26 Interaction

We were intrigued by the finding that our proteomic analyses of interactors with RSV and HIV-1 Gag proteins identified several components of the Mediator complex (RSV: Med6, Med13L, Med22, Med24, Med26, Med30; HIV-1: Med9, Med13, Med15, Med21, Med23, Med26, Med28)(**Tables 3, 5, S1-S4**), a coactivator involved in the regulated transcription of nearly all RNAPII genes [58,73-76]. The presence of multiple Mediator proteins in our dataset raised the likelihood that Gag may interact with this multiprotein complex. Furthermore, Mediator proteins have been shown to be exploited by other viruses and endogenous retroelements [36-43]. Interestingly, Med26 and Med30 are both metazoan-specific Mediator proteins, implying that Gag may display selectivity for metazoan-specific Mediator complexes over those with protozoan orthologs. Med26 and Med30 both also have critical roles in transcription, with Med26 responsible for the recruitment of important elongation factors to sites of transcription, and Med30 providing stabilization of the intact Mediator core [77,78].

We previously showed that nuclear localization of Gag contributes to efficient USvRNA selection for packaging [7]. One possible model is that interaction of Gag with Mediator proteins could tether Gag at active transcription sites, increasing the chance that it would find and associate with nascent USvRNA. To examine the interaction of RSV Gag and Med26, we utilized confocal microscopy to assess colocalization between transfected RSV Gag and Med26 in QT6 cells and observed that these two proteins did in fact colocalize (**Fig 4A-B, Video S1**) (M1, Med26∩RSV Gag: 0.0364 ± 0.004; M2, RSV Gag∩Med26: 0.471 ± 0.067). Next, immunoprecipitation of endogenous RSV Gag from RC.V8-infected QT6 cell nuclear lysates was undertaken, and samples were resolved via SDS-PAGE and subjected to western

blotting, as described in the Methods section (**Fig 4C**). In line with our proteomic results, we

observed that Med26 and RSV Gag co-immunoprecipitated (**Fig 4C, Eluate lane**). The mo-

lecular mechanisms underlying this interaction and the possible involvement of additional

Mediator proteins will be pursued in future studies.

Table 3. Names and functions of the proteins identified in the RSV proteomics list under

GO:0006366~ transcription from RNA polymerase II promoter.

Symbol	Description	Function related to Transcription (mostly summarized from Genecards)
ARID4A	AT-rich interactive domain-containing protein 4A	DNA-binding protein which modulates activity of several transcription factors. May function as part of an mSin3A repressor complex.
ATXN1L	Ataxin-1-like	Chromatin-binding factor that repress Notch signaling in the absence of Notch intracellular domain by acting as a CBF1 corepressor. Binds to the HEY promoter and might assist, along with NCOR2, RBPJ-mediated repression.
CCNK	Cyclin-K	Regulatory subunit of cyclin-dependent kinases that mediates activation of target kinases. Plays a role in transcriptional regulation via its role in regulating the phosphorylation of the C-terminal domain (CTD) of the large subunit of RNA polymerase II (POLR2A).
CDK13	Cyclin Dependent Kinase 13	Hyperphosphorylates the C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit RPB1, thereby acting as a key regulator of transcription elongation.
CHD7	Chromodomain Helicase DNA Binding Protein 7	May interact with CTCF (see below). Is a member of the chromodomain helicase DNA-binding family of ATP-dependent chromatin remodeling enzymes [79].
CHTOP	Chromatin Target of PRMT1	Required for effective mRNA nuclear export and is a component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced

		pre-mRNA.
CNOT7	CCR4-NOT transcription complex subunit 7	Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation.
CTCF	CCCTC-Binding Factor	Chromatin binding factor that binds to DNA sequence specific sites. Involved in transcriptional regulation by binding to chromatin insulators and preventing interaction between promoter and nearby enhancers and silencers. Inversely, binding to target sites is prevented by CpG methylation. Plays an important role in chromatin remodeling preventing spreading of CpG methylation and maintaining methylation-free zones.
CUX1	Cut Like Homeobox 1	Transcription factor involved in the control of neuronal differentiation in the brain. It probably has a broad role in mammalian development as a repressor of developmentally regulated gene expression. May act by preventing binding of positively-activating CCAAT factors to promoters.
EIF4A3	Eukaryotic Translation Initiation Factor 4A3	Involved in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*
ETV6	ETS Variant Transcription Factor 6	Transcriptional repressor; binds to the DNA sequence 5'-CCGGAAGT-3'.
FOXK2	Forkhead Box K2	Transcriptional regulator involved in different processes such as glucose metabolism, aerobic glycolysis and autophagy. Recognizes and binds the forkhead DNA sequence motif (5'-GTAAACA-3') and can both act as a transcription activator or repressor, depending on the context.
FOXP1	Forkhead Box P1	Transcriptional repressor.
HCFC2	Host Cell Factor C2	Component of the MLL1/MLL complex, a multiprotein complex that mediates both methylation of 'Lys-4' of histone H3 (H3K4me) complex, a specific tag for epigenetic transcriptional activation.

HNRNPK	heterogeneous nuclear ribonucleoprotein H3	Plays an important role in p53/TP53 response to DNA damage, acting at the level of both transcription activation and repression.
INTS3	Integrator complex subunit 3	Component of the Integrator (INT) complex. The Integrator complex is involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A).
INTS5	Integrator Complex Subunit 5	Component of the Integrator (INT) complex (see above).
JUN	Jun Proto-Oncogene, AP-1 Transcription Factor Subunit	Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'.
LEF1	Lymphoid Enhancer Binding Factor 1	Transcription factor that binds DNA in a sequence-specific manner. Participates in the Wnt signaling pathway.
LEO1	LEO1 Homolog, Paf1/RNA Polymerase II Complex Component	Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation.
MAGOH	Mago Homolog, Exon Junction Complex Subunit	Involved in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*
MBD3	Methyl-CpG-binding domain protein 3	Acts as transcriptional repressor and plays a role in gene silencing.
MED6	Mediator of RNA polymerase II transcription subunit 6	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional

		preinitiation complex with RNA polymerase II and the general transcription factors.
MED24	Mediator of RNA polymerase II transcription subunit 24	Component of the Mediator complex (see above).
MED26	Mediator of RNA polymerase II transcription subunit 26	Component of the Mediator complex (see above).
MED30	Mediator of RNA polymerase II transcription subunit 30	Component of the Mediator complex (see above).
NOC2L	NOC2 Like Nucleolar Associated Transcriptional Repressor	Acts as an inhibitor of histone acetyltransferase activity; prevents acetylation of all core histones by the EP300/p300 histone acetyltransferase at p53/TP53-regulated target promoters in a histone deacetylases (HDAC)-independent manner. Acts as a transcription corepressor of p53/TP53- and TP63-mediated transactivation of the p21/CDKN1A promoter.
OGT	UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit	Component of a THAP1/THAP3-HCFC1-OGT complex that is required for the regulation of the transcriptional activity of RRM1. Plays a key role in chromatin structure by mediating O-GlcNAcylation of 'Ser-112' of histone H2B: recruited to CpG-rich transcription start sites of active genes via its interaction with TET proteins (TET1, TET2 or TET3). As part of the NSL complex indirectly involved in acetylation of nucleosomal histone H4 on several lysine residues.
PLK1	Polo Like Kinase 1	Phosphorylates FOXM1, a key mitotic transcription regulator, leading to enhance FOXM1 transcriptional activity. Phosphorylates the transactivation domain of the transcription factor p73/TP73, leading to inhibit p73/TP73-mediated transcriptional activation and pro-apoptotic functions.
POLR2B	RNA Polymerase II Subunit B	Second largest component of RNA polymerase II. Proposed to contribute to the polymerase catalytic activity and forms the polymerase active center together with the largest subunit.
RAD21	RAD21 Cohesin Complex Component	In interphase, cohesins may function in the control of gene expression by binding to numerous sites within the genome. May

		control RUNX1 gene expression. Binds to and represses APOB gene promoter.
RFC1	Replication Factor C Subunit 1	Could play a role in DNA transcription regulation as well as DNA replication and/or repair. Can bind single- or double-stranded DNA. Interacts with C-terminus of PCNA. 5' phosphate residue is required for binding of the N-terminal DNA-binding domain to duplex DNA, suggesting a role in recognition of non-primer template DNA structures during replication and/or repair.
RPRD1B	Regulation of nuclear pre-mRNA domain-containing protein 1B	Interacts with phosphorylated C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit POLR2A, and participates in dephosphorylation of the CTD by RPAP2. Prevents RNA polymerase II from reading through the 3' end termination site and may allow it to be recruited back to the promoter through promotion of the formation of a chromatin loop.
SAFB	Scaffold attachment factor B1	Binds to scaffold/matrix attachment region (S/MAR) DNA and forms a molecular assembly point to allow the formation of a 'transcriptosomal' complex (consisting of SR proteins and RNA polymerase II) coupling transcription and RNA processing.
SAP30	Sin3A Associated Protein 30	Involved in the functional recruitment of the Sin3-histone deacetylase complex (HDAC) to a specific subset of N-CoR corepressor complexes. Active in deacetylating core histone octamers (when in a complex) but inactive in deacetylating nucleosomal histones.
SMAD2	SMAD Family Member 2	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. The SMAD2/SMAD4 complex functions to activate transcription.
SP3	Sp3 Transcription Factor	Transcriptional factor that can act as an activator or repressor depending on isoform and/or post-translational modifications.
SRSF2	Serine and Arginine Rich Splicing	A unique SR protein that activates transcription in a position-dependent

(SC35)	Factor 2	manner. Binds to promoter-associated small RNAs to mediate transcription pause release [80].
SRSF11	Serine and Arginine Rich Splicing Factor 11	Involved in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*
STRN3	Striatin-3	Binding partner of glucocorticoid receptor (GR) that interferes with GR's ligand-dependent transactivation capacity.
TAF2	TATA-Box Binding Protein Associated Factor 2	Transcription factor TFIID is one of the general factors required for accurate and regulated initiation by RNA polymerase II. TFIID is a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. It requires core promoter-specific cofactors for productive transcription stimulation. TAF2 stabilizes TFIID binding to core promoter.
TAF7	TATA-Box Binding Protein Associated Factor 7	Functions as a component of the DNA-binding general transcription factor complex TFIID.
TAF8	TATA-Box Binding Protein Associated Factor 8	Functions as a component of the DNA-binding general transcription factor complex TFIID. Mediates both basal and activator-dependent transcription. Required for the integration of TAF10 in the TAF complex.
TAF13	TATA-Box Binding Protein Associated Factor 13	Functions as a component of the DNA-binding general transcription factor complex TFIID.
TCEB3 **(ELOA)**	Elongin A	Also known as elongin, is a general transcription elongation factor that increases the RNA polymerase II transcription elongation past template-encoded arresting sites. Subunit A is transcriptionally active and its transcription activity is strongly enhanced by binding to the dimeric complex of the SIII regulatory subunits B and C (elongin BC complex).
TCF3	Transcription Factor 3	Transcriptional regulator.
TCF20	Transcription Factor 20	Transcriptional activator that binds to the regulatory region of MMP3 and thereby controls stromelysin expression. It stimulates the activity of various

Symbol	Description	Function
		transcriptional activators such as JUN, SP1, PAX6 and ETS1, suggesting a function as a coactivator.
THOC3	THO complex 3	Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not unspliced pre-mRNA.
TTF2	Transcription Termination Factor 2	DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template.
U2AF1	Splicing factor U2AF 35 kDa subunit	Involved in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*
VWA9 (INTS14)	von Willebrand factor A domain containing 9 (Integrator Complex Subunit 14)	Probable component of the Integrator (INT) complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing.
XPO1	Exportin-1	Mediates the nuclear export of cellular proteins (cargos) bearing a leucine-rich nuclear export signal (NES) and of RNAs.
ZEB1	Zinc Finger E-Box Binding Homeobox 1	Acts as a transcriptional repressor.
ZMYND8	Protein kinase C-binding protein 1 (Zinc finger MYND domain-containing protein 8)	May act as a transcriptional corepressor for KDM5D.

536 **Table 4.** Names and functions of the proteins identified in the RSV proteomics list under

537 GO:0008380~RNA splicing.

Symbol	Description	Function related to Splicing (mostly summarized from Genecards)
AQR	Aquarius intron-binding spliceosomal factor	Intron-binding spliceosomal protein required to link pre-mRNA splicing and snoRNP (small nucleolar ribonucleoprotein) biogenesis
CACTIN	Cactin, Spliceosome C Complex Subunit	Essential for efficient splicing of many precursor mRNAs (pre-mRNAs). Interacts with the spliceosome-associated factors DHX8 and SRRM2 [81].
CDK13	Cyclin Dependent	Required for RNA splicing, probably by phosphorylating

	Kinase 13	SRSF1/SF2.
EIF4A3	Eukaryotic Translation Initiation Factor 4A3	Involved in pre-mRNA splicing as component of the spliceosome. Core component of the splicing-dependent multiprotein exon junction complex (EJC) deposited at splice junctions on mRNAs.
FUS	FUS RNA Binding Protein	Binds to nascent pre-mRNAs and acts as a molecular mediator between RNA polymerase II and U1 small nuclear ribonucleoprotein thereby coupling transcription and splicing.
HNRNPH1	Heterogeneous Nuclear Ribonucleoprotein H1	Mediates pre-mRNA alternative splicing regulation.
HNRNPH3	heterogeneous nuclear ribonucleoprotein H3	Involved in the splicing process and participates in early heat shock-induced splicing arrest.
HNRNPK	Heterogeneous Nuclear Ribonucleoprotein K	Member of the heterogeneous nuclear ribonucleoproteins (hnRNPs) family. The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport.
ISY1	ISY1 Splicing Factor Homolog	Component of the spliceosome C complex required for the selective processing of microRNAs during embryonic stem cell differentiation.
LUC7L3	LUC7-like 3 pre-mRNA splicing factor; human homolog of yeast U1 snRNP-associated factor	Acts as a bridge between the pre-mRNA and the U1 snRNP during splicing [82].
MAGOH	Mago Homolog, Exon Junction Complex Subunit	Required for pre-mRNA splicing as component of the spliceosome. Plays a redundant role with MAGOHB as core component of the exon junction complex (EJC) and in the nonsense-mediated decay (NMD) pathway.
NONO	Non-POU Domain Containing Octamer Binding	Involved in pre-mRNA splicing, probably as a heterodimer with SFPQ. Interacts with U5 snRNA.
POLR2B	RNA Polymerase II Subunit B	Second largest component of RNA polymerase II.
PRPF8	pre-mRNA	A core component of precatalytic, catalytic and

	processing factor 8	postcatalytic spliceosomal complexes, both of the predominant U2-type spliceosome and the minor U12-type spliceosome. Functions as a scaffold that mediates the ordered assembly of spliceosomal proteins and snRNAs. Required for the assembly of the U4/U6-U5 tri-snRNP complex.
PRPF39	Pre-MRNA Processing Factor 39	Involved in pre-mRNA splicing.
PRPF40A	PRP40 pre-mRNA processing factor 40 homolog A	May be involved in pre-mRNA splicing through interactions with SF1 (splicing factor 1) [83].
PTBP1	polypyrimidine tract binding protein 1	Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. May promote RNA looping when bound to two separate polypyrimidine tracts of introns in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA.
SCAF8	SR-Related CTD Associated Factor 8	Identified in a complex with CDC5L and other spliceosomal proteins.
SF3A3	splicing factor 3a subunit 3	Involved in pre-mRNA splicing as a component of the splicing factor SF3A complex that contributes to the assembly of the 17S U2 snRNP, and the subsequent assembly of the pre-spliceosome 'E' complex and the pre-catalytic spliceosome 'A' complex. Involved in pre-mRNA splicing as a component of pre-catalytic spliceosome 'B' complexes.
SMNDC1	Survival Motor Neuron Domain Containing 1	Necessary for spliceosome assembly.
SNRNP40	small nuclear ribonucleoprotein U5 subunit 40	Required for pre-mRNA splicing as a component of the U5 small nuclear ribonucleoprotein (snRNP) complex and the U4/U6-U5 tri-snRNP complex.
SNRNP200	small nuclear ribonucleoprotein U5 subunit 200	Plays role in pre-mRNA splicing as core component of precatalytic, catalytic and postcatalytic spliceosomal complexes. Involved in spliceosome assembly, activation and disassembly. Catalyzes the ATP-dependent unwinding of U4/U6 RNA duplexes, an essential step in the assembly of a catalytically active spliceosome.
SRSF2	Serine and Arginine Rich Splicing Factor 2	Required for formation of the earliest ATP-dependent splicing complex and interacts with spliceosomal components bound to both the 5'- and 3'-splice sites during spliceosome assembly. It also is required for ATP-dependent interactions of both U1 and U2 snRNPs with pre-mRNA. Interacts with other spliceosomal components, via the RS domains, to form a bridge

		between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF.
SRSF11	Serine and Arginine Rich Splicing Factor 11	May function in pre-mRNA splicing. Contains an arginine/serine-rich region similar to segments found in pre-mRNA splicing factors.
SUGP1 (SF4)	SURP and G-patch domain containing 1	Plays a role in pre-mRNA splicing. Interacts with U2AF2 [84,85]
THOC3	THO complex 3	Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not unspliced pre-mRNA.
TRA2B	Transformer 2 Beta Homolog	Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing. Can either activate or suppress exon inclusion. Alters pre-mRNA splicing patterns by antagonizing the effects of splicing regulators.
TTF2	Transcription Termination Factor 2	Interacts with cell division cycle 5-like, associates with human splicing complexes, and plays a role in pre-mRNA splicing.
U2AF1	U2 small nuclear RNA auxiliary factor 1	Plays a critical role in both constitutive and enhancer-dependent splicing by mediating protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. Recruits U2 snRNP to the branch point. Directly mediates interactions between U2AF2 and proteins bound to the enhancers and thus may function as a bridge between U2AF2 and the enhancer complex to recruit it to the adjacent intron.
USP39	Ubiquitin Specific Peptidase 39	Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP.
WTAP	Wilms tumor 1 associated protein	Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing. Acts as a mRNA splicing regulator.
ZCCHC8	Zinc Finger CCHC-Type Containing 8	An accessory factor to the nuclear RNA exosome complex. This complex is also thought to recruit the exosome to degrade intronic RNAs via its interaction with both the exosome and the spliceosome.
ZNF326	Zinc Finger Protein 326	Core component of the DBIRD complex, a multiprotein complex that acts at the interface between core mRNP particles and RNA polymerase II (RNAPII) and integrates transcript elongation with the regulation of alternative

	splicing.

538 *PPP1R9B was identified but not included in this list as a clear function in splicing was not

539 identified.

541 **Table 5.** Names and functions of the proteins identified in the HIV-1 proteomics list under

542 GO:0006366~ transcription from RNA polymerase II promoter.

Symbol	Description	Function related to Transcription (mostly summarized from Genecards)	Frequency (# of publications)
ALYREF	Aly/REF Export Factor	Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm. Involved in transcription elongation and genome stability. Acts as chaperone and promotes the dimerization of transcription factors containing basic leucine zipper (bZIP) domains and thereby promotes transcriptional activation.	2
ARID4A	AT-rich interactive domain-containing protein 4A	DNA-binding protein which modulates activity of several transcription factors. May function as part of an mSin3A repressor complex.	0
ASH2L	ASH2 Like, Histone Lysine Methyltransferase Complex Subunit	Component or associated component of some histone methyltransferase complexes which regulates transcription through recruitment of those complexes to gene promoters.	0

ASUN (INTS13)	asunder, spermatogenesis regulator	A probable component of the Integrator (INT) complex – involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and 3' processing. The INT complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.	1
BRMS1	breast cancer metastasis suppressor 1	Transcriptional repressor. Down-regulates NF-κB transcriptional activation. Promotes HDAC1 binding to promoter regions.	0
C1QBP	Complement component 1 Q subcomponent-binding protein, mitochondrial	May be involved in regulation of FOXC1 transcriptional activity and NFY/CCAAT-binding factor complex-mediated transcription.	1
CDK12	cyclin dependent kinase 12	Cyclin-dependent kinase that phosphorylates the C-terminal domain (CTD) of the large subunit of RNA polymerase II (POLR2A), thereby acting as a key regulator of transcription elongation. Regulates the expression of genes involved in DNA repair and is required for the maintenance of genomic stability.	1
CDKN2A	cyclin dependent kinase inhibitor 2A (Tumor suppressor ARF)	Has an isoform (ARF) that binds to BCL6 and down-regulates BCL6-induced transcriptional repression. Binds to E2F1 and MYC and blocks their transcriptional activator activity but has no effect on MYC transcriptional repression. Binds to TOP1/TOPOI and stimulates its activity. This complex binds to rRNA gene promoters and may play a role in rRNA transcription and/or maturation.	0
CGGBP1	CGG Triplet Repeat Binding Protein 1	Binds to nonmethylated 5'-d(CGG)(n)-3' trinucleotide repeats in the FMR1 promoter. May play a role in regulating FMR1 promoter.	0

CHTOP	Chromatin Target of PRMT1	Required for effective mRNA nuclear export and is a component of the TREX complex (see above).	2
CIC	capicua transcriptional repressor (protein capicua homolog)	Transcriptional repressor which plays a role in development of the central nervous system (CNS).	0
CLP1	Cleavage Factor Polyribonucleotide Kinase Subunit 1	Member of the tRNA splicing complex. May be involved in the termination of RNA polymerase II transcription (Reactome).*	0
CPSF3	cleavage and polyadenylation specific factor 3	Member of the cleavage and polyadenylation specificity factor (CPSF) complex. May be involved in the termination of RNA polymerase II transcription (Reactome).*	0
CPSF3L (INTS11)	cleavage and polyadenylation specific factor 3-like (Integrator Complex Subunit 11)	Catalytic component of the Integrator (INT) complex (see above).	0
CPSF7	Cleavage And Polyadenylation Specific Factor 7	Member of the CPSF complex. May be involved in the termination of RNA polymerase II transcription (Reactome).*	0
CREBBP	CREB binding protein	Acetylates histones, giving a specific tag for transcriptional activation.	0
CTNNB1	Catenin beta 1	In the presence of Wnt ligand, CTNNB1 accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes.	1
DDX20	DEAD-Box Helicase 20	Represses Egr2-mediated transcriptional activation [86].	1
DNAJA3	DnaJ heat shock protein family (Hsp40) member A3	Can modulate IFN-gamma-mediated transcriptional activity.	1
DNMT1	DNA methyltransferase	Mediates transcriptional repression by direct binding to HDAC2. In association with	1

		DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9.	
EHMT2	Euchromatic Histone Lysine Methyltransferase 2	Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones.	0
FIP1L1	Factor Interacting With PAPOLA And CPSF1	Member of the CPSF complex. May be involved in RNA polymerase II transcription termination (Reactome)*	1
FOXP1	forkhead box P1	Transcriptional repressor.	0
GATAD2A	GATA zinc finger domain containing 2A	Transcriptional repressor. Enhances MBD2-mediated repression. Efficient repression requires the presence of GATAD2B.	0
GTF2H1	general transcription factor IIH subunit 1	Component of the general transcription and DNA repair factor IIH (TFIIH) core complex, which is involved in general and transcription-coupled nucleotide excision repair (NER) of damaged DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. In transcription, TFIIH has an essential role in transcription initiation.	0
GTF2H3	general transcription factor IIH subunit 3	See above.	0
GTF2H4	general transcription factor IIH subunit 4	See above.	0
HDAC3	histone deacetylase 3	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation	0

		gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events.	
HIST1H1E	H1.4 Linker Histone, Cluster Member	Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation.	2
HIST2H3A	H3 Clustered Histone 15	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.	0
ILK	Integrin Linked Kinase	The kinase activity regulates key signaling pathways, leading to the stimulation of downstream effector kinases and transcription factors [87].	0
INO80	INO80 Complex ATPase Subunit	ATPase component of the chromatin remodeling INO80 complex which is involved in transcriptional regulation, DNA replication and DNA repair.	0
INTS9	integrator complex subunit 9	Component of the Integrator (INT) complex (see above).	1
LEO1	LEO1 homolog, Paf1/RNA polymerase II complex component	Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is	0

implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation.

MBD2	Methyl-CpG Binding Domain Protein 2	Recruits histone deacetylases and DNA methyltransferases. Acts as transcriptional repressor and plays a role in gene silencing.	0
MECP2	Methyl-CpG Binding Protein 2	Mediates transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A.	0
MED13	mediator complex subunit 13	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.	0
MED15	mediator complex subunit 15	Component of the Mediator complex (see above). Required for cholesterol-dependent gene regulation.	0
MED21	mediator complex subunit 21	Component of the Mediator complex (see above).	0
MED23	Mediator Complex Subunit 23	Required for transcriptional activation subsequent to the assembly of the pre-initiation complex. Component of the Mediator complex (see above).	0
MED26	Mediator Complex	Component of the Mediator	1

	Subunit 26	complex (see above).	
MEIS2	Meis Homeobox 2	Involved in transcriptional regulation.	1
MINA (RIOX2)	MYC induced nuclear antigen (Ribosomal Oxygenase 2)	Oxygenase that can act as both a histone lysine demethylase and a ribosomal histidine hydroxylase. Is involved in the demethylation of trimethylated 'Lys-9' on histone H3 (H3K9me3), leading to an increase in ribosomal RNA expression.	1
MTDH	Metadherin	Downregulates SLC1A2/EAAT2 promoter activity when expressed ectopically. Activates the nuclear factor kappa-B (NF-kappa-B) transcription factor.	4
NCOR1	Nuclear Receptor Corepressor 1	Mediates transcriptional repression by certain nuclear receptors. Part of a complex which promotes histone deacetylation and the formation of repressive chromatin structures which may impede the access of basal transcription factors.	0
NFYA	Nuclear Transcription Factor Y Subunit Alpha	Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors.	0
OGT	O-linked N-acetylglucosamine (GlcNAc) transferase	Component of a THAP1/THAP3-HCFC1-OGT complex that is required for the regulation of the transcriptional activity of RRM1. Plays a key role in chromatin structure by mediating O-GlcNAcylation of 'Ser-112' of histone H2B: recruited to CpG-rich transcription start sites of active genes via its interaction with TET proteins (TET1, TET2 or TET3). As part of the NSL complex indirectly involved in acetylation of nucleosomal histone H4 on	0

		several lysine residues.	
PABPN1	poly(A) binding protein nuclear 1	Cooperates with SKIP to synergistically activate E-box-mediated transcription through MYOD1 and may regulate the expression of muscle-specific genes.	1
PAXBP1	PAX3 And PAX7 Binding Protein 1	Adapter protein linking the transcription factors PAX3 and PAX7 to the histone methylation machinery. Associates with a histone methyltransferase complex that specifically mediates dimethylation and trimethylation of 'Lys-4' of histone H3.	1
PBX1	PBX homeobox 1	Acts as a transcriptional activator of PF4 in complex with MEIS1. PDX1:PBX1b:MEIS2b complex is involved in the transcriptional activation of the ELA1 enhancer. Probably in complex with MEIS2, is involved in transcriptional regulation by KLF4. Acts as a transcriptional activator of NKX2-5 and a transcriptional repressor of CDKN2B.	0
PCBP1	poly(rC) binding protein 1	Involved in the transcriptional activation of BRCA1 promoter [88], eukaryotic translation initiation factor 4E [89], and the mouse mu opioid receptor [90].	1
PCF11	PCF11 Cleavage and Polyadenylation Factor Subunit	The encoded protein is necessary for efficient Pol II transcription termination and may be involved in degradation of the 3' product of polyA site cleavage.	0
POLDIP3	DNA polymerase delta interacting protein 3	May be involved in RNA polymerase II transcription termination (Reactome)	0
POLR2B	RNA Polymerase II Subunit B	Second largest component of RNA polymerase II. Proposed to contribute to the polymerase catalytic activity and forms the polymerase active center together with the largest subunit.	0
POLR2E	RNA polymerase II	Common component of RNA	0

	subunit E	polymerases I, II and III. In Pol II, POLR2E/RPB5 is part of the lower jaw surrounding the central large cleft and thought to grab the incoming DNA template.	
PSMC3	proteasome 26S subunit, ATPase 3	One of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex.	1
RAD21	RAD21 cohesin complex component	In interphase, cohesins may function in the control of gene expression by binding to numerous sites within the genome. May control RUNX1 gene expression. Binds to and represses APOB gene promoter.	0
RBM10	RNA binding motif protein 10	Component of a large chromatin remodeling complex, composed of MYSM1, PCAF, RBM10 and KIF11/TRIP5.	2
RBM14	RNA Binding Motif Protein 14	Isoform 1 may function as a nuclear receptor coactivator, enhancing transcription through other coactivators such as NCOA6 and CITED1. Isoform 2, functions as a transcriptional repressor, modulating transcriptional activities of coactivators including isoform 1, NCOA6 and CITED1.	3
RFC1	replication factor C subunit 1	Could play a role in DNA transcription regulation as well as DNA replication and/or repair. Interacts with C-terminus of PCNA.	2
RNMT	RNA guanine-7 methyltransferase	Interacts with elongating form of polymerase II and RNGTT	0
SAFB	Scaffold Attachment Factor B	Binds to scaffold/matrix attachment region (S/MAR) DNA and forms a molecular assembly point to allow the formation of a 'transcriptosomal' complex (consisting of SR proteins and	0

		RNA polymerase II) coupling transcription and RNA processing.	
SETD2	SET Domain Containing 2, Histone Lysine Methyltransferase	Histone methyltransferase that specifically trimethylates 'Lys-36' of histone H3 (H3K36me3), a specific tag for epigenetic transcriptional activation. Plays a role in chromatin structure modulation during elongation by coordinating recruitment of the FACT complex and by interacting with hyperphosphorylated POLR2A.	0
SMAD2	SMAD family member 2	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. The SMAD2/SMAD4 complex functions to activate transcription.	0
SMAD5	SMAD Family Member 5	Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase.	0
SMARCA4	SWI/SNF Related, Matrix Associated, Actin Dependent Regulator Of Chromatin, Subfamily A, Member 4	Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner.	1
SMARCB1	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily b, member 1	Core component of the BAF (hSWI/SNF) complex. This ATP-dependent chromatin-remodeling complex. Involved in activation of CSF1 promoter.	0
SMARCC2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin	Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of	1

	subfamily c member 2	DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes (see above).	
SQSTM1	sequestosome 1	Regulates activation of the NF-kB signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-kB in response to upstream signals.	0
SRRM1	serine and arginine repetitive matrix 1	Has a role in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*	0
SRSF3 (SRp20)	serine and arginine rich splicing factor 3	Has a role in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*	3
SRSF7 (9G8)	serine and arginine rich splicing factor 7	Has a role in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*	3
SUDS3	SDS3 Homolog, SIN3A Corepressor Complex Component	Regulatory protein which represses transcription and augments histone deacetylase activity of HDAC1. May have a potential role in tumor suppressor pathways through regulation of apoptosis. May function in the assembly and/or enzymatic activity of the mSin3A corepressor complex or in mediating interactions between the complex and other regulatory complexes.	0
SUPT16H	SPT16 Homolog, Facilitates Chromatin Remodeling Subunit	Component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the	1

		nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II.	
SUZ12	SUZ12 Polycomb Repressive Complex 2 Subunit	Polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene.	0
TARDBP	TAR DNA Binding Protein	Transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription.	0
TCEB1 (ELOC)	transcription elongation factor B subunit 1	A general transcription elongation factor that increases the RNA polymerase II transcription elongation past template-encoded arresting sites.	1
TGFB1I1	transforming growth factor beta 1 induced transcript 1	In the nucleus, functions as a nuclear receptor coactivator regulating glucocorticoid, androgen, mineralocorticoid and progesterone receptor transcriptional activity.	0
THOC3	THO complex 3	Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not unspliced pre-mRNA.	0
TRIM33	tripartite motif containing 33	May act as a transcriptional repressor. Inhibits the transcriptional response to TGF-beta/BMP signaling cascade.	0
TRPS1	Transcriptional Repressor GATA Binding 1	Transcriptional repressor.	0
U2AF1	U2 small nuclear RNA auxiliary factor 1	Has a role in splicing. May be involved in RNA polymerase II transcription termination (Reactome).*	1

UBE2I	Ubiquitin Conjugating Enzyme E2 I	Interacts with various transcription factors such as TFAP2A, TFAP2B, TFAP2C, AR, ETS1 and SOX4.	0
VWA9 (INTS14)	von Willebrand factor A domain containing 9	Probable component of the Integrator (INT) complex (see above).	0
WDR5	WD repeat domain 5	Contributes to histone modification.	1
WDR61	WD repeat domain 61	Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II (see above). Required for Hox gene transcription.	0
ZBTB7B	Zinc Finger And BTB Domain Containing 7B	Transcription regulator that acts as a key regulator of lineage commitment of immature T-cell precursors. Transcriptional repressor of the collagen COL1A1 and COL1A2 genes. May also function as a repressor of fibronectin and possibly other extracellular matrix genes.	0
ZEB1	Zinc Finger E-Box Binding Homeobox 1	Acts as a transcriptional repressor.	0

544 **Table 6.** Names and function of the proteins identified in the HIV-1 proteomics list under the

545 GO:0008380~RNA splicing.

Symbol	Description	Function related to RNA Splicing (mostly summarized from Genecards)	Frequency (# of publications)
ALYREF	Aly/REF Export Factor	Involved in nuclear export of spliced and unspliced mRNAs through specific binding to the m5C by being a component of the TREX complex. The TREX complex functions to couple different steps during mRNA biogenesis, including mRNA transcription, processing, decay, and nuclear export.	2

C1QBP	Complement component 1 Q subcomponent-binding protein, mitochondrial	Involved in regulation of RNA splicing by inhibiting the RNA-binding capacity of SRSF1 and its phosphorylation. Required for the nuclear translocation of splicing factor U2AF1L4.	1
CCAR2	cell cycle and apoptosis regulator 2	A core component of the DBIRD complex, a multiprotein complex that acts at the interface between core messenger ribonucleoprotein (mRNP) particles and RNA polymerase II and integrates transcript elongation with the regulation of alternative splicing. The DBIRD complex affects local transcript elongation rates and alternative splicing of a large set of exons embedded in (A + T)-rich DNA regions.	1
CDC5L	cell division cycle 5 like	Plays role in precursor mRNA (pre-mRNA) splicing as core component of precatalytic, catalytic and postcatalytic spliceosomal complexes; the PRP19-CDC5L complex.	3
CDK12	cyclin dependent kinase 12	Required for RNA splicing, possibly by phosphorylating SRSF1/SF2.	1
CLP1	Cleavage Factor Polyribonucleotide Kinase Subunit 1	Component of the pre-mRNA cleavage complex II (CF-II), which seems to be required for mRNA 3'-end formation. Component of the tRNA splicing endonuclease complex and is implicated in tRNA, mRNA, and siRNA maturation.	0
CPSF3	cleavage and polyadenylation specific factor 3	Component of the cleavage and polyadenylation specificity factor (CPSF) complex that play a key role in pre-mRNA 3'-end formation and functions as mRNA 3'-end-processing endonuclease. Also involved in the histone 3'-end pre-mRNA processing through U7 small nuclear ribonucleoprotein (snRNP) interactions.	0

CPSF4	cleavage and polyadenylation specific factor 4	Component of the CPSF complex (see above). Binds RNA polymers with a preference for poly(U).	0
CPSF7	Cleavage and Polyadenylation Specific Factor 7	Component of the cleavage factor Im (CFIm) complex that functions as an activator of the pre-mRNA 3'-end cleavage and polyadenylation processing. Directly activates the mRNA 3'-processing machinery and binds to polyadenylation signals in RNA substrates. Interacts with the splicing factor U2 snRNP Auxiliary Factor (U2AF) 65 to link the splicing and polyadenylation complexes.	0
DDX20	DEAD-Box Helicase 20	DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. Directly interacts with survival of motor neurons (SMN) complex and may play a catalytic role in the function of the SMN complex on spliceosome RNPs. May also play a role in the metabolism of small nucleolar ribonucleoprotein (snoRNPs).	1
DDX46	DEAD-Box Helicase 46	DEAD box protein family. Plays an essential role in splicing, either prior to, or during splicing A complex formation.	1
DDX47	DEAD-box helicase 47	DEAD box protein family. May have a role in rRNA processing and mRNA splicing. Associates with pre-rRNA precursors.	2
DNAJC8	DnaJ Heat Shock Protein Family (Hsp40) Member C8	Interacts with SRPK1 (see below).	0
ELAVL1 (HuR)	ELAV like RNA binding protein 1	RNA-binding protein that binds to poly-U elements and AU-rich elements in the 3'-UTR region of mRNAs increasing their stability.	2
FAM98B	family with sequence similarity 98 member B	Component of the tRNA-splicing ligase complex [91,92].	0

FIP1L1	Factor Interacting with PAPOLA And CPSF1	Component of the CPSF complex. Contributes to poly(A) site recognition and stimulates poly(A) addition. Binds to U-rich RNA sequence elements surrounding the poly(A) site. May act to tether poly(A) polymerase to the CPSF complex.	1
HNRNPH3	heterogeneous nuclear ribonucleoprotein H3	Involved in the splicing process and participates in early heat shock-induced splicing arrest.	2
HNRNPM	heterogeneous nuclear ribonucleoprotein M	Member of the heterogeneous nuclear ribonucleoproteins (hnRNPs) family. The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport.	3
HNRNPR	Heterogeneous Nuclear Ribonucleoprotein R	Component of hnRNPs (see above). A member of the spliceosome C complex.	2
HNRNPUL1	heterogeneous nuclear ribonucleoprotein U like 1	Plays also a role in mRNA processing and transport. Binds avidly to poly(G) and poly(C) RNA homopolymers *in vitro*.	2
LSM3	LSM3 homolog, U6 small nuclear RNA and mRNA degradation associated	Plays a role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex that is involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex).	0
PABPN1	poly(A) binding protein nuclear 1	Involved in the 3'-end formation of pre-mRNAs by the addition of a poly(A) tail of 200-250 nt to the upstream cleavage product. Promotes intron excision in the context of 3'-end polyadenylation but not when bound to internal A-tracts [93].	1
PCBP1	poly(rC) binding protein 1	Interaction with SMAD3 regulates the alternative splicing of CD44 [94].	1

PCF11	PCF11 Cleavage and Polyadenylation Factor Subunit	Component of pre-mRNA cleavage complex II.	0
PLRG1	pleiotropic regulator 1	Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing.	3
POLR2B	RNA Polymerase II Subunit B	Second largest component of RNA polymerase II	0
POLR2E	RNA polymerase II subunit E	Common component of RNA polymerases I, II and III.	0
PPIH	peptidylprolyl isomerase H	Participates in pre-mRNA splicing. May play a role in the assembly of the U4/U5/U6 tri-snRNP complex of the spliceosome.	1
PRPF3	pre-mRNA processing factor 3	Plays role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex in spliceosome complex. Spliceosome B complex.	4
PRPF4	pre-mRNA processing factor 4	Plays role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex in spliceosome complex. Spliceosome B complex.	4
PRPF38B	pre-mRNA processing factor 38B	May be required for pre-mRNA splicing.	0
PRPF40A	Pre-MRNA Processing Factor 40 Homolog A	May be involved in pre-mRNA splicing. with SF1 (splicing factor 1) [83].	0
PTBP1	polypyrimidine tract binding protein 1	Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. May promote RNA looping when bound to two separate polypyrimidine tracts of introns in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA.	1
PUF60	poly(U) binding splicing factor 60	DNA- and RNA-binding protein, involved in several nuclear processes such as pre-mRNA splicing, apoptosis and	0

transcription regulation.
Promotes splicing of an intron
with weak 3'-splice site and
pyrimidine tract in a cooperative
manner with U2AF2. Modulates
alternative splicing of several
mRNAs.

RAVER2	ribonucleoprotein, PTB binding 2	Interactions with PTB may be required for role in splicing [95,96].	0
RBM7	RNA Binding Motif Protein 7	Interacts with splicing factors, SAP145 and SRp20 [97], as well as U4/U6.U5 tri-snRNP-associated proteins [98].	1
RBM10	RNA binding motif protein 10	Associates with hnRNP proteins and may be involved in regulating alternative splicing.	2
SCAF8	SR-Related CTD Associated Factor 8	Identified in a complex with CDC5L and other spliceosomal proteins.	0
SMN2	survival of motor neuron 2, centromeric	The SMN complex plays a catalyst role in the assembly of snRNPs of the spliceosome. Ensures the correct splicing of U12 intron-containing genes. May also play a role in the metabolism of snoRNPs.	1
SNRPA	small nuclear ribonucleoprotein polypeptide A	Component of the spliceosomal U1 snRNP, which is essential for recognition of the pre-mRNA 5' splice-site and the subsequent assembly of the spliceosome. In a snRNP-free form (SF-A) may be involved in coupled pre-mRNA splicing and polyadenylation process.	2
SNRPD2	small nuclear ribonucleoprotein D2 polypeptide	Plays role in pre-mRNA splicing as core component of the SMN-Sm complex that mediates spliceosomal snRNP assembly and as component of the spliceosomal U1, U2, U4 and U5 snRNPs. Component of both the pre-catalytic spliceosome B complex and activated spliceosome C complexes. A component of the minor U12	1

		spliceosome.	
SRPK1	SRSF Protein Kinase 1	Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing. Plays a central role in the regulatory network for splicing, controlling the intranuclear distribution of splicing factors in interphase cells and the reorganization of nuclear speckles during mitosis.	1
SRRM1	serine and arginine repetitive matrix 1	Part of pre- and post-splicing multiprotein mRNP complexes. Promotes constitutive and exonic splicing enhancer (ESE)-dependent splicing activation by bridging together sequence-specific and basal snRNP factors of the spliceosome.	0
SRSF3	serine and arginine rich splicing factor 3	Splicing factor that specifically promotes exon-inclusion during alternative splicing.	3
SRSF7	serine and arginine rich splicing factor 7	Required for pre-mRNA splicing.	3
SRSF12	Serine And Arginine Rich Splicing Factor 12	Splicing factor that seems to antagonize SR proteins in pre-mRNA splicing regulation.	0
TARDBP	TAR DNA Binding Protein	RNA-binding protein that is involved in various steps of RNA biogenesis and processing. Regulates the splicing of many non-coding and protein-coding RNAs. Regulates mRNA stability.	0
THOC3	THO complex 3z	Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates	0

		with spliced mRNA and not unspliced pre-mRNA.	
TRA2B	Transformer 2 Beta Homolog	Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing. Can either activate or suppress exon inclusion. Alters pre-mRNA splicing patterns by antagonizing the effects of splicing regulators.	2
U2AF1	U2 small nuclear RNA auxiliary factor 1	Plays a critical role in both constitutive and enhancer-dependent splicing by mediating protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. Recruits U2 snRNP to the branch point. Directly mediates interactions between U2AF2 and proteins bound to the enhancers and thus may function as a bridge between U2AF2 and the enhancer complex to recruit it to the adjacent intron.	1

4. Discussion

Several different laboratories have observed that the retroviral Gag proteins of HIV-1,

RSV, MMTV, MLV, FIV, PFV, and MPMV undergo nuclear localization [9-23,25,26,99-102].

As described in detail in this report, six previously published proteomic studies searching for

binding partners of HIV-1 Gag identified many nuclear proteins [30-35]. As we were specifi-

cally interested in the nuclear interactomes of RSV and HIV-1 Gag, we took a different ap-

proach than previous groups, using nuclear lysates incubated with recombinant Gag proteins

to perform affinity purification of complexes followed by mass spectrometry for our proteomic

analysis. Despite these differences in methodology, a set of overlapping factors were identi-

fied for HIV-1 Gag that included a large number of proteins involved in nuclear processes such

as transcription/gene expression, RNA processing, splicing, and chromatin remodeling.

Comparison of the potential binding partners of RSV and HIV-1 Gag indicated that 57 proteins

were found to be in common (**Fig 2**). Whether these factors have similar functions in RSV or

HIV-1 replication remains to be examined. When the HIV-1 Gag interactomes identified by other laboratories were compared to the interacting proteins identified by us in this report, 190 common proteins were found by at least two independent laboratory groups. Further experimentation will be needed to validate each of these factors to determine whether they play important roles in retrovirus replication or pathogenesis **(Fig 5)**.

Our past and present cell fractionation experiments demonstrated that, in addition to being present in the nucleoplasm, both the RSV and HIV-1 Gag proteins can be extracted from euchromatin and heterochromatin fractions, complementing our recently published report indicating that HIV-1 Gag localizes with euchromatin marks at the nuclear periphery [9,10]. These results suggest that HIV-1 Gag may be specifically targeted to a chromatin-associated compartment through interactions with host nuclear binding partners. Indeed, many of the proteins identified by us and others are chromatin-associated, raising the importance of further exploring the signals that target chromatin-associated regions, examining Gag interactions with chromatin factors, and elucidating possible roles for Gag in chromatin-related functions. To better understand how these chromatin-associated factors interact, and to identify which histone-associated protein networks appear most closely associated with HIV-1 Gag, we employed the use of the STRING Consortium database (**Fig 6**). Of those identified, histones H1 (8 proteins), H2A (14 proteins), and H2B (11 proteins) displayed the strongest evidence for association. Histones H3 and H4 were also identified, but each only displayed a single association. Additionally, four histone deacetylases (HDACs)—including the transcription-repressing class II HDACs 4 and 6—and seven members of the SWI/SNF chromatin remodeling complex (e.g. SMARCC2, SMARCA5, DEK, BAZ1A, MAZ1B, ARID1B, and ARID2) were identified [103]. SWI/SNF complexes are highly enriched at transcriptional enhancer regions, where they modulate accessibility to promote gene activations [104]. Interestingly, SWI/SNF complexes have been shown to play a role in HIV-1 infection [105-107], and also interact with histones H2A, H2B, and H4. Given the spatial organization of the nucleosome, it is possible that HIV-1 Gag may be interacting with factors residing near histones

H1, H2A, H2B, and H4, to locate viral transcription sites, influence transcription of viral or

cellular genes, or facilitate chromatin remodeling to expose the integrated provirus.

It will be important to investigate whether these interactions facilitate viral replication steps

occurring in the nucleus, such as gene expression, viral RNA processing and splicing,

unspliced viral RNA binding by Gag, and/or nuclear export of viral RNA complexes, or whether

Gag influences cellular processes for the benefit of the virus, for example by modulating ex-

pression of cellular genes involved in the immune response or altering splicing of host genes

to change their localization or function. **Figure 5** highlights the various nuclear processes that

HIV-1 Gag could be involved in based on the proteins identified in the HIV-1 proteomic stud-

ies.

Our published data indicating that RSV and HIV-1 Gag each interact specifically with their

cognate USvRNAs to form discrete foci to form viral RNP complexes in the interchromatin

space raise the possibility that interaction with host factors facilitates co-transcriptional retro-

viral genome selection [8,9]. In both viruses, these findings were confirmed in infected cells,

where Gag was targeted to the USvRNA transcription site, presumably at the site of proviral

integration. Among other transcription-related proteins, in this study we identified multiple

members of the Mediator complex, which has been shown to be hijacked by other viruses and

retroelements, and we demonstrated that RSV Gag colocalized and co-immunoprecipitated

with Med26 (**Fig 4 and Video S1**) [36-43]. Quantitative analysis of our imaging experiments

indicated that nearly half of the nuclear Gag signal was colocalized with Med26. Visual in-

spection of the colocalization channel revealed that much of this colocalization was at Gag foci

in the perichromatin space (**Fig 4A**). Our previous work demonstrating that RSV and HIV-1

Gag proteins can form biomolecular condensates (BMCs) [108,109], combined with evidence

for the existence of Mediator-containing transcriptional condensates [110,111], argues that

Gag may be utilizing condensate-driven compartmentalization to increase its chances of

encountering nascent viral RNA. Further studies will address this intriguing possibility.

Additionally, we identified two components of the WMM complex, CBLL1 (Cbl Pro-

to-Oncogene Like 1; present in RSV and HIV-1 datasets) and WTAP (Wilms tumor 1 associ-

ated protein; RSV dataset only), which mediates m6-methyladenosine (m6A) methylation of RNAs. This RNA modification affects RNA splicing and processing, and full length HIV-1 RNA containing m6A has displayed a bias towards serving as template RNA for the translation of viral proteins, as opposed to being packaged as gRNA [102]. Interaction with components of the WMM complex could simply serve to localize Gag to sites where viral RNA may exist, but it is intriguing to consider the possibility that Gag antagonizes the function of this complex to maintain a pool of genomic RNA.

For RSV, we have observed movement of viral RNPs across the nuclear membrane, suggesting that these complexes may ultimately be packaged into assembling virions at the plasma membrane [8]. We also previously found that RSV Gag colocalizes with splicing factors SF2 and SC35 [66] in splicing speckles, which are located in the perichromatin space, raising the possibility that Gag may influence splicing of viral and/or cellular RNAs in a co-transcriptional manner. It would be of interest to test the hypothesis that RSV Gag suppresses splicing at viral intron/exon junctions to promote retention of full-length viral RNA for use as the viral genome.

Tables 3-6 list the proteins involved in transcription and splicing identified in the RSV and HIV-1 Gag interactomes. We noted that there were Gag-interacting proteins identified at different stages of transcription, including initiation, elongation, and termination. Interestingly, PolR2B, the second largest subunit of RNA polymerase II (RNAPII), was identified in both the RSV and HIV-1 interactome datasets, raising the possibility that Gag interacts with RNAPII itself [112,113]. Each dataset also contained a member of the Elongin complex (TCEB1, or Elongin C, for HIV-1; TCEB3, or Elongin A, for RSV), which forms a complex with PolR2B to promote elongation [114]. Numerous members of the Integrator complex were also identified in the datasets (VWA9, RSV and HIV-1; INTS3 and INTS5, RSV only; ASUN, INTS5, INTS9, and CPSF3L, HIV-1 only). Integrator regulates transcription by binding to the C-terminal tail of PolR2A (the largest subunit of RNAPII) during pausing, which results in cleavage of newly synthesized RNA and termination of transcription [115-117]. Other factors that modulate transcription identified were LEO1 and SCAF8 (both RSV and HIV-1), CDK13 and ZNF326

(RSV only), and ALYREF, CCAR2, CDK12, SETD2, and SUPT16H (HIV-1 only). One intriguing idea is that Gag could alter transcription elongation rates or regulate pausing of nascent viral RNA synthesis to promote folding of the psi region, recruit essential host RNA binding factors, modulate or suppress termination or splicing of viral RNA, or alter other co-transcriptional processes that could have downstream effects on the fate of viral RNA.

The data presented here, along with our analysis of previously published HIV-1 Gag interactomes, indicate that many potential host protein partners of RSV and HIV-1 Gag reside in the nucleus in association with chromatin. As both RSV and HIV-1 Gag have nuclear populations that form vRNPs in the perichromatin region, the interaction of Gag proteins with chromatin-associated nuclear factors merits further investigation. Novel roles for Gag proteins of RSV, HIV-1, and other retroviruses may be uncovered, shedding light on previously unknown aspects of the replication cycle that could be targeted by antiviral therapies.

Supplementary Materials:

Spreadsheet S1. RSV Gag Dataset 1.xlsx;

Spreadsheet S2. RSV Gag Dataset 2.xlsx;

Spreadsheet S3. HIV Gag Dataset 1.xlsx;

Spreadsheet S4. HIV Gag Dataset 2.xlsx

Table S1. Top 10 DAVID biological processes for proteins isolated from the RSV Gag affinity purifications from DF1 nuclear lysates;

Table S2. Top 10 nuclear enriched DAVID biological processes for proteins isolated from the RSV Gag affinity purifications from DF1 nuclear lysates;

Table S3. Top 10 DAVID biological processes for proteins isolated from the purified HIV Gag affinity purifications from HeLa nuclear lysates;

Table S4. Top 10 nuclear enriched DAVID biological processes for proteins isolated from the purified HIV Gag affinity purifications from HeLa nuclear lysates;

Table S5. Top 10 DAVID biological processes of nuclear proteins identified in Engeland *et al.*, 2011;

Table S6. Top 10 DAVID biological processes of nuclear proteins identified in Engeland *et al.*, 2014;

Table S7. Top 10 DAVID biological processes of nuclear proteins identified in Jäger *et al.*, 2012;

Table S8. Top 10 DAVID biological processes of nuclear proteins identified in Ritchie *et al.*, 2015;

Table S9. Top 10 DAVID biological processes of nuclear proteins identified in Le Sage *et al.*, 2015

Table S10. Top 10 DAVID biological processes of nuclear proteins identified in Li *et al.*, 2016;

Video S1. RSV Gag and Mediator complex subunit 26 (Med26) colocalize in transfected QT6 cells

Funding: This project was funded in part by NIH awards P50GM103297 (LJP), R01CA076534 (LJP), R01 GM139392 (LJP), R21/R33 DA053689 (LJP), T32 CA60395 (BLR), and F31 CA196292 (BLR). This project was also supported by the Penn State College of Medicine's Comprehensive Health Studies Program. This content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgments: We would like to acknowledge the expert assistance of Anne Stanley and Bruce Stanley in the PSU College of Medicine Mass Spectrometry and Proteomics Core, for their assistance with the mass spectrometry experiments and analysis. We greatly appreciate and acknowledge John Flanagan in the Department of Biochemistry and Molecular Biology at Penn State College of Medicine for purification of the recombinant RSV and HIV-1 Gag proteins. We acknowledge Bradley Winters at the Penn State College of Medicine for creating the illustration in Figure 5. We thank members of the Parent laboratory, Eunice Chen, Malgorzata Sudol, and Kevin Tuffy, for insightful discussions and suggestions.

The Mass Spectrometry and Proteomics Core (RRID:SCR_017831) and Advanced Light Microscopy core (RRID:SCR_022526) services and instruments used in this project were funded, in part, by the Pennsylvania State University College of Medicine via the Office of the Vice Dean of Research and Graduate Students and the Pennsylvania Department of Health using Tobacco Settlement Funds (CURE). The content is solely the responsibility of the authors and does not necessarily represent the official views of the University or College of Medicine. The Pennsylvania Department of Health specifically disclaims responsibility for any analyses, interpretations or conclusions.

Conflicts of Interest: The authors declare no conflict of interest.

Figure Legends

Figure 1. Schematic diagram and western blot with quantitation. (**A**) RSV Gag constructs. (**B**) Subcellular fractionations were performed to separate the cytoplasm and nucleoplasm from the chromatin fractions, which were further separated using differential NaCl concentrations. The 150 mM chromatin fraction (Chr 150 mM) contains proteins associated with open chromatin (euchromatin). The 500 mM chromatin fraction (Chr 500 mM) contains proteins that are associated with condensed chromatin (heterochromatin). Wild type RSV Gag and Gag.L219A were detected in all of the fractions at different ratios. Gag.ΔNC was primarily detected in the cytoplasm with very little in the chromatin fractions. (**C**) Band densities were determined for each Gag construct for each fraction and are displayed as mean ± standard error of the mean (SEM) for three biological replicates. To assess fraction purity, cellular proteins were detected using antibodies against calnexin (cytoplasm), Med4 (nucleoplasm and euchromatin), and Histone H2B (euchromatin and heterochromatin). The position of molecular weight markers, in kilodaltons, are indicated on the right.

Figure 2. The number of proteins identified in both the RSV and HIV-1 Gag affinity tagged purifications. 57 proteins were found to be in common between RSV and HIV-1 Gag.

Figure 3. The relative representation of the IPA categories present in each protein list. (**A**) The proteins identified in our HIV-1 Gag pulldown that were also found in at least one of the previously published reports were analyzed for molecular and cellular functions. The color key on the right is the same for each pie graph. Protein functions identified in this publication (Rice) (**B**), Engeland 2011 (**C**), Engeland 2014 (**D**), Le Sage (**E**), Li (**F**), Ritchie (**G**), and Jager (**H**).

Figure 4. RSV Gag colocalized and co-immunoprecipitated with Med26. (**A**) Transfected RSV Gag-GFP (red) and FLAG-Med26 (green) colocalize (white) within QT6 cells. Image representative of average colocalization, as quantified in panel (**B**). Nuclei (blue) are outlined by a dotted white line, and regions boxed in the main images are enlarged below. White arrows are included to guide the eye. Scale bars = 2 µm. (**B**) Manders' Overlap Coefficient values for image set represented by (**A**). Individual values are shown in addition mean ± SEM, n ≥ 17; ****, $p<0.0001$ by unpaired two-tailed t-test. (**C**) 500 µg of RC.V8-infected QT6 nuclear lysates were incubated with an α-RSV Gag antibody (mouse α-RSV CA.A11, gift from Neil Christensen, Penn State College of Medicine), followed by antibody capture on Pierce™ Protein G Magnetic Beads. After extensive washing, proteins were eluted from beads by boiling in 1X SDS-PAGE sample buffer and run on a 10% SDS-PAGE gel, transferred to PVDF, and Western blotted first for Med26 (top) followed by RSV Gag (bottom). The position of molecular weight markers, in kilodaltons, are indicated on the left. FT, flow through; Beads, lysate only; Eluate, lysate plus antibody. Images representative of three independent experiments.

Figure 5. HIV-1 interactome pathway analysis. This diagram illustrates the HIV-1 Gag-interacting nuclear host factors discussed in this study. Factors shaded blue were uniquely identified in the present study (newly identified). Proteins shaded in purple were identified in this publication as well as at least one other published report (previously identi-

fied). Gray shading was used to show host protein complexes that are involved with Gag-interacting factors in chromatin remodeling, gene expression, nuclear export, and splicing. (**A**) HIV-1 Gag interacting factors that promote an open chromatin structure (euchromatin state) are on the left, whereas proteins involved in condensing chromatin are shown on the right. (**B**) Proteins involved in regulation of gene expression are depicted. Factors that promote transcription initiation are indicated by arrows and green plus signs. Factors that suppress or inhibit transcription are demarcated by red blocking lines. (**C**) The two nucleoporin proteins NUP98 and NUP188 were identified and are involved in trafficking between the nucleus and the cytoplasm through the nuclear pore complex. (**D**) Proteins that localize to the spliceosome and are involved in RNA splicing are shown.

Figure 6. STRING protein network map of HIV-1 Gag interacting host chromatin proteins. Protein lists were generated from mass spectrometry experiments of Gag interacting proteins. Gene lists were then categorized into gene ontology (GO) terms to identify those which were chromatin associated. This refined list was input into STRING Consortium v12.0 (https://string-db.org/) to generate a protein-protein interaction map. A total of 129 proteins were queried and the following physical protein-protein interactions were made. Among the 129 proteins queried, 117 proteins are displayed after a minimum required interaction score of 0.4 was applied. Proteins that do not have any interacting partners are shown at the bottom left corner of the map. Lines are generated based on known interactions (purple lines), predicted gene fusions (red lines), predicted gene neighborhoods (green lines), and predicted gene co-occurrences (blue lines) from the literature.

References

1.   Butterfield-Gerson, K.L.; Scheifele, L.Z.; Ryan, E.P.; Hopper, A.K.; Parent, L.J. Importin-β Family Members Mediate Alpharetrovirus Gag Nuclear Entry via Interactions with Matrix and Nucleocapsid. *J Virol* **2006**, *80*, 1798-1806, doi:10.1128/jvi.80.4.1798-1806.2006.

2.   Gudleski, N.; Flanagan, J.M.; Ryan, E.P.; Bewly, M.C.; Parent, L.J. Directionality of nucleocytoplasmic transport of the retroviral Gag protein depends on sequential

binding of karyopherins and viral RNA. *Proceedings of the National Academy of Sciences* **2010**, *107*, 9358-9363.

3. Scheifele, L.Z.; Garbitt, R.A.; Rhoads, J.D.; Parent, L.J. Nuclear entry and CRM1-dependent nuclear export of the Rous sarcoma virus Gag polyprotein. *Proceedings of the National Academy of Sciences* **2002**, *99*, 3944-3949, doi:10.1073/pnas.062652199.

4. Scheifele, L.Z.; Ryan, E.P.; Parent, L.J. Detailed Mapping of the Nuclear Export Signal in the Rous Sarcoma Virus Gag Protein. *J Virol* **2005**, *79*, 8732-8741, doi:10.1128/jvi.79.14.8732-8741.2005.

5. Scheifele, L.Z.; Kenney, S.P.; Cairns, T.M.; Craven, R.C.; Parent, L.J. Overlapping roles of the Rous sarcoma virus Gag p10 domain in nuclear export and virion core morphology. *J Virol* **2007**, *81*, 10718-10728, doi:10.1128/JVI.01061-07.

6. Rice, B.L.; Stake, M.S.; Parent, L.J. TNPO3-Mediated Nuclear Entry of the Rous Sarcoma Virus Gag Protein Is Independent of the Cargo-Binding Domain. *J Virol* **2020**, *94*, doi:10.1128/JVI.00640-20.

7. Garbitt-Hirst, R.; Kenney, S.P.; Parent, L.J. Genetic evidence for a connection between Rous sarcoma virus Gag nuclear trafficking and genomic RNA packaging. *J.Virol.* **2009**, *83*, 6790-6797.

8. Maldonado, R.J.K.; Rice, B.; Chen, E.C.; Tuffy, K.M.; Chiari, E.F.; Fahrbach, K.M.; Hope, T.J.; Parent, L.J. Visualizing Association of the Retroviral Gag Protein with Unspliced Viral RNA in the Nucleus. *mBio* **2020**, *11*, e00524-00520, doi:10.1128/mBio.00524-20.

9. Tuffy, K.M.; Maldonado, R.J.K.; Chang, J.; Rosenfeld, P.; Cochrane, A.; Parent, L.J. HIV-1 Gag Forms Ribonucleoprotein Complexes with Unspliced Viral RNA at Transcription Sites. *Viruses* **2020**, *12*.

10. Chang, J.; Parent, L.J. HIV-1 Gag colocalizes with euchromatin histone marks at the nuclear periphery. J Virol 2023 Vol. 97 Issue 12 Pages e0117923; 37991367 PMCID: PMC10734548 DOI: 10.1128/jvi.01179-23.

11. Lochmann, T.L.; Bann, D.V.; Ryan, E.P.; Beyer, A.R.; Mao, A.; Cochrane, A.; Parent, L.J. NC-Mediated Nucleolar Localization of Retroviral Gag Proteins. *Virus research* **2013**, *171*, 304-318, doi:10.1016/j.virusres.2012.09.011.

12. Gallay, P.; Swingler, S.; Song, J.; Bushman, F.; Trono, D. HIV nuclear import is governed by the phosphotyrosine-mediated binding of matrix to the core domain of integrase. *Cell* **1995**, *83*, 569-576.

13. Zhang, J.; Crumpacker, C.S. Human immunodeficiency virus type 1 nucleocapsid protein nuclear localization mediates early viral mRNA expression. *J Virol* **2002**, *76*, 10444-10454.

14. Kemler, I.; Meehan, A.; Poeschla, E.M. Live-Cell Coimaging of the Genomic RNAs and Gag Proteins of Two Lentiviruses. *J Virol* **2010**, *84*, 6352-6366, doi:10.1128/jvi.00363-10.

15. Kemler, I.; Saenz, D.; Poeschla, E. Feline Immunodeficiency Virus Gag Is a Nuclear Shuttling Protein. *J Virol* **2012**, *86*, 8402-8411, doi:10.1128/jvi.00692-12.

825   16.   Bohl, C.R.; Brown, S.M.; Weldon, R.A. The pp24 phosphoprotein of Mason-Pfizer
826   monkey virus contributes to viral genome packaging. *Retrovirology* **2005**, *2*, 68,
827   doi:10.1186/1742-4690-2-68.

828   17.   Weldon, R.A.; Sarkar, P.; Brown, S.M.; Weldon, S.K. Mason–Pfizer monkey virus Gag
829   proteins interact with the human sumo conjugating enzyme, hUbc9. *Virology* **2003**,
830   *314*, 62-73, doi:https://doi.org/10.1016/S0042-6822(03)00348-9.

831   18.   Baluyot, M.F.; Grosse, S.A.; Lyddon, T.D.; Janaka, S.K.; Johnson, M.C.
832   CRM1-dependent trafficking of retroviral Gag proteins revisited. *J Virol* **2012**, *86*,
833   4696-4700.

834   19.   Beyer, A.R.; Bann, D.V.; Rice, B.; Pultz, I.S.; Kane, M.; Goff, S.P.; Golovkina, T.V.;
835   Parent, L.J. Nucleolar Trafficking of the Mouse Mammary Tumor Virus Gag Protein
836   Induced by Interaction with Ribosomal Protein L9. *J Virol* **2013**, *87*, 1069-1082,
837   doi:10.1128/jvi.02463-12.

838   20.   Elis, E.; Ehrlich, M.; Prizan-Ravid, A.; Laham-Karam, N.; Bacharach, E. p12 Tethers
839   the Murine Leukemia Virus Pre-integration Complex to Mitotic Chromosomes. *PLOS
840   Pathogens* **2012**, *8*, e1003103, doi:10.1371/journal.ppat.1003103.

841   21.   Nash, M.A.; Meyer, M.K.; Decker, G.L.; Arlinghaus, R.B. A subset of Pr65Gag is
842   nucleus associated in murine leukemia virus-infected cells. *J Virol* **1993**, *67*,
843   1350-1356.

844   22.   Risco, C.; Menendez-Arias, L.; Copeland, T.D.; Pinto da Silva, P.; Oroszlan, S.
845   Intracellular transport of the murine leukemia virus during acute infection of NIH 3T3
846   cells: nuclear import of nucleocapsid protein and integrase. *Journal of Cell Science*
847   **1995**, *108*, 3039.

848   23.   Schneider, W.M.; Brzezinski, J.D.; Aiyer, S.; Malani, N.; Gyuricza, M.; Bushman, F.D.;
849   Roth, M.J. Viral DNA tethering domains complement replication-defective mutations in
850   the p12 protein of MuLV Gag. *Proceedings of the National Academy of Sciences* **2013**,
851   *110*, 9487-9492, doi:10.1073/pnas.1221736110.

852   24.   Müllers, E.; Stirnnagel, K.; Kaulfuss, S.; Lindemann, D. Prototype foamy virus Gag
853   nuclear localization: a novel pathway among retroviruses. *J Virol* **2011**, *85*, 9276-9285,
854   doi:10.1128/JVI.00663-11.

855   25.   Renault, N.; Tobaly-Tapiero, J.; Paris, J.; Giron, M.-L.; Coiffic, A.; Roingeard, P.; Saïb,
856   A. A nuclear export signal within the structural Gag protein is required for prototype
857   foamy virus replication. *Retrovirology* **2011**, *8*, 6, doi:10.1186/1742-4690-8-6.

858   26.   Schliephake, A.W.; Rethwilm, A. Nuclear localization of foamy virus Gag precursor
859   protein. *J Virol* **1994**, *68*, 4946-4954.

860   27.   Tobaly-Tapiero, J.; Bittoun, P.; Lehmann-Che, J.; Delelis, O.; Giron, M.L.; de Thé, H.;
861   Saïb, A. Chromatin tethering of incoming foamy virus by the structural Gag protein.
862   *Traffic* **2008**, *9*, 1717-1727, doi:10.1111/j.1600-0854.2008.00792.x.

863   28.   Yu, K.L.; Lee, S.H.; Lee, E.S.; You, J.C. HIV-1 nucleocapsid protein localizes
864   efficiently to the nucleus and nucleolus. *Virology* **2016**, *492*, 204-212,
865   doi:10.1016/j.virol.2016.03.002.

29. Brzezinski, J.D.; Modi, A.; Liu, M.; Roth, M.J. Repression of the Chromatin-Tethering Domain of Murine Leukemia Virus p12. *J Virol* **2016**, *90*, 11197-11207, doi:10.1128/JVI.01084-16.

30. Engeland, C.E.; Brown, N.P.; Börner, K.; Schümann, M.; Krause, E.; Kaderali, L.; Müller, G.A.; Kräusslich, H.-G. Proteome analysis of the HIV-1 Gag interactome. *Virology* **2014**, *460-461*, 194-206, doi:https://doi.org/10.1016/j.virol.2014.04.038.

31. Engeland, C.E.; Oberwinkler, H.; Schümann, M.; Krause, E.; Müller, G.A.; Kräusslich, H.-G. The Cellular Protein Lyric Interacts with HIV-1 Gag. *J Virol* **2011**, *85*, 13322-13332, doi:10.1128/jvi.00174-11.

32. Jäger, S.; Cimermancic, P.; Gulbahce, N.; Johnson, J.R.; McGovern, K.E.; Clarke, S.C.; Shales, M.; Mercenne, G.; Pache, L.; Li, K.; et al. Global landscape of HIV–human protein complexes. *Nature* **2011**, *481*, 365, doi:10.1038/nature10719

https://www.nature.com/articles/nature10719#supplementary-information.

33. Le Sage, V.; Cinti, A.; Valiente-Echeverría, F.; Mouland, A.J. Proteomic analysis of HIV-1 Gag interacting partners using proximity-dependent biotinylation. *Virology Journal* **2015**, *12*, 138, doi:10.1186/s12985-015-0365-6.

34. Li, Y.; Frederick, K.M.; Haverland, N.A.; Ciborowski, P.; Belshan, M. Investigation of the HIV-1 matrix interactome during virus replication. *Proteomics – Clinical Applications* **2016**, *10*, 156-163, doi:10.1002/prca.201400189.

35. Ritchie, C.; Cylinder, I.; Platt, E.J.; Barklis, E. Analysis of HIV-1 Gag Protein Interactions via Biotin Ligase Tagging. *J Virol* **2015**, *89*, 3988-4001, doi:10.1128/jvi.03584-14.

36. Kornberg, R.D. Mediator and the mechanism of transcriptional activation. *Trends Biochem Sci* **2005**, *30*, 235-239, doi:10.1016/j.tibs.2005.03.011.

37. Boyer, T.G.; Martin, M.E.; Lees, E.; Ricciardi, R.P.; Berk, A.J. Mammalian Srb/Mediator complex is targeted by adenovirus E1A protein. *Nature* **1999**, *399*, 276-279, doi:10.1038/20466.

38. Yang, M.; Hay, J.; Ruyechan, W.T. Varicella-zoster virus IE62 protein utilizes the human mediator complex in promoter activation. *J Virol* **2008**, *82*, 12154-12163, doi:10.1128/JVI.01693-08.

39. Lester, J.T.; DeLuca, N.A. Herpes simplex virus 1 ICP4 forms complexes with TFIID and mediator in virus-infected cells. *J Virol* **2011**, *85*, 5733-5744, doi:10.1128/JVI.00385-11.

40. Vijayalingam, S.; Chinnadurai, G. Adenovirus L-E1A activates transcription through mediator complex-dependent recruitment of the super elongation complex. *J Virol* **2013**, *87*, 3425-3434, doi:10.1128/JVI.03046-12.

41. Pei, J.; Beri, N.R.; Zou, A.J.; Hubel, P.; Dorando, H.K.; Bergant, V.; Andrews, R.D.; Pan, J.; Andrews, J.M.; Sheehan, K.C.F.; et al. Nuclear-localized human respiratory syncytial virus NS1 protein modulates host gene transcription. *Cell Rep* **2021**, *37*, 109803, doi:10.1016/j.celrep.2021.109803.

42. Rovnak, J.; Quackenbush, S.L. Exploitation of the Mediator complex by viruses. *PLoS Pathog* **2022**, *18*, e1010422, doi:10.1371/journal.ppat.1010422.

43. Asimi, V.; Sampath Kumar, A.; Niskanen, H.; Riemenschneider, C.; Hetzel, S.; Naderi, J.; Fasching, N.; Popitsch, N.; Du, M.; Kretzmer, H.; et al. Hijacking of transcriptional condensates by endogenous retroviruses. *Nat Genet* **2022**, doi:10.1038/s41588-022-01132-w.

44. Himly, M.; Foster, D.N.; Bottoli, I.; Iacovoni, J.S.; Vogt, P.K. The DF-1 Chicken Fibroblast Cell Line: Transformation Induced by Diverse Oncogenes and Cell Death Resulting from Infection by Avian Leukosis Viruses. *Virology* **1998**, *248*, 295-304, doi:https://doi.org/10.1006/viro.1998.9290.

45. Craven, R.C.; Leure-duPree, A.E.; Weldon, R.A.; Wills, J.W. Genetic analysis of the major homology region of the Rous sarcoma virus Gag protein. *J Virol* **1995**, *69*, 4213-4227.

46. Kenney, S.P.; Lochmann, T.L.; Schmid, C.L.; Parent, L.J. Intermolecular Interactions between Retroviral Gag Proteins in the Nucleus. *J Virol* **2008**, *82*, 683-691, doi:10.1128/jvi.02049-07.

47. Bewley, M.C.; Reinhart, L.; Stake, M.S.; Nadaraia-Hoke, S.; Parent, L.J.; Flanagan, J.M. A non-cleavable hexahistidine affinity tag at the carboxyl-terminus of the HIV-1 Pr55Gag polyprotein alters nucleic acid binding properties. *Protein Expression and Purification* **2017**, *130*, 137-145, doi:https://doi.org/10.1016/j.pep.2016.10.001.

48. Rye-McCurdy, T.D.; Nadaraia-Hoke, S.; Gudleski-O'Regan, N.; Flanagan, J.M.; Parent, L.J.; Musier-Forsyth, K. Mechanistic Differences between Nucleic Acid Chaperone Activities of the Gag Proteins of Rous Sarcoma Virus and Human Immunodeficiency Virus Type 1 Are Attributed to the MA Domain. *J Virol* **2014**, *88*, 7852-7861, doi:10.1128/jvi.00736-14.

49. Fujiwara, T.; Oda, K.; Yokota, S.; Takatsuki, A.; Ikehara, Y. Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *Journal of Biological Chemistry* **1988**, *263*, 18545-18552.

50. Chase, G.P.; Rameix-Welti, M.-A.; Zvirbliene, A.; Zvirblis, G.; Götz, V.; Wolff, T.; Naffakh, N.; Schwemmle, M. Influenza Virus Ribonucleoprotein Complexes Gain Preferential Access to Cellular Export Machinery through Chromatin Targeting. *PLOS Pathogens* **2011**, *7*, e1002187, doi:10.1371/journal.ppat.1002187.

51. Weldon, R.A.; Erdie, C.R.; Oliver, M.G.; Wills, J.W. Incorporation of chimeric Gag protein into retroviral particles. *J Virol* **1990**, *64*, 4169-4179.

52. Shilov, I.V.; Seymour, S.L.; Patel, A.A.; Loboda, A.; Tang, W.H.; Keating, S.P.; Hunter, C.L.; Nuwaysir, L.M.; Schaeffer, D.A. The Paragon Algorithm, a Next Generation Search Engine That Uses Sequence Temperature Values and Feature Probabilities to Identify Peptides from Tandem Mass Spectra. *Molecular & Cellular Proteomics* **2007**, *6*, 1638-1655, doi:10.1074/mcp.T600050-MCP200.

53. Tang, W.H.; Shilov, I.V.; Seymour, S.L. Nonlinear Fitting Method for Determining Local False Discovery Rates from Decoy Database Searches. *Journal of Proteome Research* **2008**, *7*, 3661-3667, doi:10.1021/pr070492f.

54. Huang, D.W.; Sherman, B.T.; Lempicki, R.A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* **2008**, *4*, 44, doi:10.1038/nprot.2008.211

https://www.nature.com/articles/nprot.2008.211#supplementary-information.

55. Huang, D.W.; Sherman, B.T.; Lempicki, R.A. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research* **2009**, *37*, 1-13, doi:10.1093/nar/gkn923.

56. Lippé, R. Deciphering Novel Host–Herpesvirus Interactions by Virion Proteomics. *Frontiers in Microbiology* **2012**, *3*, doi:10.3389/fmicb.2012.00181.

57. Nojima, T.; Gomes, T.; Grosso, A.R.F.; Kimura, H.; Dye, M.J.; Dhir, S.; Carmo-Fonseca, M.; Proudfoot, N.J. Mammalian NET-Seq Reveals Genome-wide Nascent Transcription Coupled to RNA Processing. *Cell* **2015**, *161*, 526-540, doi:10.1016/j.cell.2015.03.027.

58. Sato, S.; Tomomori-Sato, C.; Parmely, T.J.; Florens, L.; Zybailov, B.; Swanson, S.K.; Banks, C.A.; Jin, J.; Cai, Y.; Washburn, M.P.; et al. A set of consensus mammalian mediator subunits identified by multidimensional protein identification technology. *Mol Cell* **2004**, *14*, 685-691, doi:10.1016/j.molcel.2004.05.006.

59. Kim, D.I.; KC, B.; Zhu, W.; Motamedchaboki, K.; Doye, V.; Roux, K.J. Probing nuclear pore complex architecture with proximity-dependent biotinylation. *Proceedings of the National Academy of Sciences* **2014**, 201406459, doi:10.1073/pnas.1406459111.

60. Mechold, U.; Gilbert, C.; Ogryzko, V. Codon optimization of the BirA enzyme gene leads to higher expression and an improved efficiency of biotinylation of target proteins in mammalian cells. *J Biotechnol* **2005**, *116*, 245-249.

61. Roux, K.J.; Kim, D.I.; Raida, M.; Burke, B. A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. *Journal of Cell Biology* **2012**, *196*, 801-810, doi:10.1083/jcb.201112098.

62. Gao, W.; Li, M.; Zhang, J. Tandem immunoprecipitation approach to identify HIV-1 Gag associated host factors. *J Virol Methods* **2014**, *203*, 116-119.

63. Fu, W.; Sanders-Beer, B.E.; Katz, K.S.; Maglott, D.R.; Pruitt, K.D.; Ptak, R.G. Human immunodeficiency virus type 1, human protein interaction database at NCBI. *Nucleic Acids Res* **2009**, *37*, 15.

64. Ptak, R.G.; Fu, W.; Sanders-Beer, B.E.; Dickerson, J.E.; Pinney, J.W.; Robertson, D.L.; Rozanov, M.N.; Katz, K.S.; Maglott, D.R.; Pruitt, K.D.; et al. Cataloguing the HIV type 1 human protein interaction network. *AIDS Res Hum Retroviruses* **2008**, *24*, 1497-1502.

65. Pinney, J.W.; Dickerson, J.E.; Fu, W.; Sanders-Beer, B.E.; Ptak, R.G.; Robertson, D.L. HIV-host interactions: a map of viral perturbation of the host system. *Aids* **2009**, *23*, 549-554.

66. Rice, B.; Kaddis, R.; Stake, M.; Lochmann, T.; Parent, L. Interplay between the alpharetroviral Gag protein and SR Proteins SF2 and SC35 in the nucleus. *Frontiers in Microbiology* **2015**, *6*, doi:10.3389/fmicb.2015.00925.

67. Jassal, B.; Matthews, L.; Viteri, G.; Gong, C.; Lorente, P.; Fabregat, A.; Sidiropoulos, K.; Cook, J.; Gillespie, M.; Haw, R.; et al. The reactome pathway knowledgebase. *Nucleic Acids Res* **2020**, *48*, D498-D503.

68. Wu, G.; Haw, R. Functional Interaction Network Construction and Analysis for Disease Discovery. *Methods Mol Biol* **2017**, 6783-6784_6711.

69. Kornblihtt, A.R.; De La Mata, M.; Fededa, J.P.; Muñoz, M.J.; Nogués, G. Multiple links between transcription and splicing. *RNA* **2004**, *10*, 1489-1498, doi:10.1261/rna.7100104.

70. Brody, Y.; Shav-Tal, Y. Transcription and splicing. *Transcription* **2011**, *2*, 216-220, doi:10.4161/trns.2.5.17273.

71. Herzel, L.; Ottoz, D.S.M.; Alpert, T.; Neugebauer, K.M. Splicing and transcription touch base: co-transcriptional spliceosome assembly and function. *Nature Reviews Molecular Cell Biology* **2017**, *18*, 637-650, doi:10.1038/nrm.2017.63.

72. Tellier, M.; Maudlin, I.; Murphy, S. Transcription and splicing: A two-way street. *WIREs RNA* **2020**, *11*, e1593, doi:10.1002/wrna.1593.

73. Soutourina, J. Transcription regulation by the Mediator complex. *Nat Rev Mol Cell Biol* **2018**, *19*, 262-274, doi:10.1038/nrm.2017.115.

74. Conaway, R.C.; Conaway, J.W. Function and regulation of the Mediator complex. *Curr Opin Genet Dev* **2011**, *21*, 225-230, doi:10.1016/j.gde.2011.01.013.

75. Conaway, R.C.; Conaway, J.W. The Mediator complex and transcription elongation. *Biochim Biophys Acta* **2013**, *1829*, 69-75, doi:10.1016/j.bbagrm.2012.08.017.

76. Ansari, S.A.; Morse, R.H. Mechanisms of Mediator complex action in transcriptional activation. *Cell Mol Life Sci* **2013**, *70*, 2743-2756, doi:10.1007/s00018-013-1265-9.

77. Takahashi, H.; Parmely, T.J.; Sato, S.; Tomomori-Sato, C.; Banks, C.A.; Kong, S.E.; Szutorisz, H.; Swanson, S.K.; Martin-Brown, S.; Washburn, M.P.; et al. Human mediator subunit MED26 functions as a docking site for transcription elongation factors. *Cell* **2011**, *146*, 92-104, doi:10.1016/j.cell.2011.06.005.

78. Tan, C.; Zhu, S.; Chen, Z.; Liu, C.; Li, Y.E.; Zhu, M.; Zhang, Z.; Zhang, L.; Gu, Y.; Liang, Z.; et al. Mediator complex proximal Tail subunit MED30 is critical for Mediator core stability and cardiomyocyte transcriptional network. *PLoS Genet* **2021**, *17*, e1009785, doi:10.1371/journal.pgen.1009785.

79. Basson, M.A.; van Ravenswaaij-Arts, C. Functional Insights into Chromatin Remodelling from Studies on CHARGE Syndrome. *Trends in Genetics* **2015**, *31*, 600-611, doi:https://doi.org/10.1016/j.tig.2015.05.009.

80. Mo, S.; Ji, X.; Fu, X.-D. Unique role of SRSF2 in transcription activation and diverse functions of the SR and hnRNP proteins in gene expression regulation. *Transcription* **2013**, *4*, 251-259, doi:10.4161/trns.26932.

81. Zanini, I.M.Y.; Soneson, C.; Lorenzi, L.E.; Azzalin, C.M. Human cactin interacts with DHX8 and SRRM2 to assure efficient pre-mRNA splicing and sister chromatid cohesion. *Journal of Cell Science* **2017**, *130*, 767, doi:10.1242/jcs.194068.

82. Fortes, P.; Bilbao-Cortés, D.; Fornerod, M.; Rigaut, G.; Raymond, W.; Séraphin, B.; Mattaj, I.W. Luc7p, a novel yeast U1 snRNP protein with a role in 5′ splice site recognition. *Genes & Development* **1999**, *13*, 2425-2438.

83. Crisci, A.; Raleff, F.; Bagdiul, I.; Raabe, M.; Urlaub, H.; Rain, J.-C.; Krämer, A. Mammalian splicing factor SF1 interacts with SURP domains of U2 snRNP-associated proteins. *Nucleic Acids Research* **2015**, *43*, 10456-10473, doi:10.1093/nar/gkv952.

84. Hegele, A.; Kamburov, A.; Grossmann, A.; Sourlis, C.; Wowro, S.; Weimann, M.; Will, C.L.; Pena, V.; Luhrmann, R.; Stelzl, U. Dynamic protein-protein interaction wiring of the human spliceosome. *Mol Cell* **2012**, *45*, 567-580.

85. Wang, J.; Huo, K.; Ma, L.; Tang, L.; Li, D.; Huang, X.; Yuan, Y.; Li, C.; Wang, W.; Guan, W.; et al. Toward an understanding of the protein interaction network of the human liver. *Mol Syst Biol* **2011**, *7*, 67.

86. Gillian, A.L.; Svaren, J. The Ddx20/DP103 Dead Box Protein Represses Transcriptional Activation by Egr2/Krox-20. *Journal of Biological Chemistry* **2004**, *279*, 9056-9063, doi:10.1074/jbc.M309308200.

87. Wu , C.; Dedhar , S. Integrin-linked kinase (ILK) and its interactors : a new paradigm for the coupling of extracellular matrix to actin cytoskeleton and signaling complexes. *Journal of Cell Biology* **2001**, *155*, 505-510, doi:10.1083/jcb.200108077.

88. Thakur, S.; Nakamura, T.; Calin, G.; Russo, A.; Tamburrino, J.F.; Shimizu, M.; Baldassarre, G.; Battista, S.; Fusco, A.; Wassell, R.P.; et al. Regulation of BRCA1 Transcription by Specific Single-Stranded DNA Binding Factors. *Molecular and Cellular Biology* **2003**, *23*, 3774, doi:10.1128/mcb.23.11.3774-3787.2003.

89. Lynch, M.; Chen, L.; Ravitz, M.J.; Mehtani, S.; Korenblat, K.; Pazin, M.J.; Schmidt, E.V. hnRNP K Binds a Core Polypyrimidine Element in the Eukaryotic Translation Initiation Factor 4E (eIF4E) Promoter, and Its Regulation of eIF4E Contributes to Neoplastic Transformation. *Molecular and Cellular Biology* **2005**, *25*, 6436, doi:10.1128/mcb.25.15.6436-6453.2005.

90. Choi, H.S.; Song, K.Y.; Hwang, C.K.; Kim, C.S.; Law, P.-Y.; Wei, L.-N.; Loh, H.H. A proteomic approach for identification of single-strand DNA-binding proteins involved in transcriptional regulation of mouse mu-opioid receptor gene. *Molecular & Cellular Proteomics* **2008**, doi:10.1074/mcp.M800052-MCP200.

91. Popow, J.; Englert, M.; Weitzer, S.; Schleiffer, A.; Mierzwa, B.; Mechtler, K.; Trowitzsch, S.; Will, C.L.; Lührmann, R.; Söll, D.; et al. HSPC117 Is the Essential Subunit of a Human tRNA Splicing Ligase Complex. *Science* **2011**, *331*, 760-764, doi:10.1126/science.1197847.

92. Popow, J.; Jurkin, J.; Schleiffer, A.; Martinez, J. Analysis of orthologous groups reveals archease and DDX1 as tRNA splicing factors. *Nature* **2014**, *511*, 104-107.

93. Muniz, L.; Davidson, L.; West, S. Poly(A) Polymerase and the Nuclear Poly(A) Binding Protein, PABPN1, Coordinate the Splicing and Degradation of a Subset of Human Pre-mRNAs. *Molecular and Cellular Biology* **2015**, *35*, 2218, doi:10.1128/mcb.00123-15.

1070   94.   Tripathi, V.; Sixt, K.M.; Gao, S.; Xu, X.; Huang, J.; Weigert, R.; Zhou, M.; Zhang, Y.E.
1071   Direct Regulation of Alternative Splicing by SMAD3 through PCBP1 Is Essential to the
1072   Tumor-Promoting Role of TGF-beta. *Mol Cell* **2016**, *64*, 549-564.

1073   95.   Kleinhenz, B.; Fabienke, M.; Swiniarski, S.; Wittenmayer, N.; Kirsch, J.; Jockusch,
1074   B.M.; Arnold, H.H.; Illenberger, S. Raver2, a new member of the hnRNP family. *FEBS*
1075   *Lett* **2005**, *579*, 4254-4258.

1076   96.   Henneberg, B.; Swiniarski, S.; Sabine, B.; Illenberger, S. A conserved peptide motif in
1077   Raver2 mediates its interaction with the polypyrimidine tract-binding protein. *Exp Cell*
1078   *Res* **2010**, *316*, 966-979.

1079   97.   Guo, T.B.; Boros, L.G.; Chan, K.C.; Hikim, A.P.S.; Hudson, A.P.; Swerdloff, R.S.;
1080   Mitchell, A.P.; Salameh, W.A. Spermatogenetic Expression of RNA-Binding Motif
1081   Protein 7, a Protein That Interacts With Splicing Factors. *Journal of Andrology* **2003**,
1082   *24*, 204-214, doi:10.1002/j.1939-4640.2003.tb02664.x.

1083   98.   Nag, A.; Steitz, J.A. Tri-snRNP-associated proteins interact with subunits of the
1084   TRAMP and nuclear exosome complexes, linking RNA decay and pre-mRNA splicing.
1085   *RNA Biology* **2012**, *9*, 334-342, doi:10.4161/rna.19431.

1086   99.   Lesbats, P.; Serrao, E.; Maskell, D.P.; Pye, V.E.; O'Reilly, N.; Lindemann, D.;
1087   Engelman, A.N.; Cherepanov, P. Structural basis for spumavirus Gag tethering to
1088   chromatin. *Proceedings of the National Academy of Sciences* **2017**, *114*, 5509.

1089   100.   Müllers, E.; Stirnnagel, K.; Kaulfuss, S.; Lindemann, D. Prototype Foamy Virus Gag
1090   Nuclear Localization: a Novel Pathway among Retroviruses. *J Virol* **2011**, *85*,
1091   9276-9285, doi:10.1128/jvi.00663-11.

1092   101.   Tobaly-Tapiero, J.; Bittoun, P.; Lehmann-Che, J.; Delelis, O.; Giron, M.L.; Thé, H.D.;
1093   Saïb, A. Chromatin Tethering of Incoming Foamy Virus by the Structural Gag Protein.
1094   *Traffic* **2008**, *9*, 1717-1727, doi:doi:10.1111/j.1600-0854.2008.00792.x.

1095   102.   Pereira-Montecinos, C.; Toro-Ascuy, D.; Ananías-Sáez, C.; Gaete-Argel, A.;
1096   Rojas-Fuentes, C.; Riquelme-Barrios, S.; Rojas-Araya, B.; García-de-Gracia, F.;
1097   Aguilera-Cortés, P.; Chnaiderman, J.; et al. Epitranscriptomic regulation of HIV-1
1098   full-length RNA packaging. *Nucleic Acids Res* **2022**, *50*, 2302-2318,
1099   doi:10.1093/nar/gkac062.

1100   103.   Thiagalingam, S.; Cheng, K.H.; Lee, H.J.; Mineva, N.; Thiagalingam, A.; Ponte, J.F.
1101   Histone deacetylases: unique players in shaping the epigenetic histone code. *Ann N Y*
1102   *Acad Sci* **2003**, *983*, 84-100, doi:10.1111/j.1749-6632.2003.tb05964.x.

1103   104.   Mittal, P.; Roberts, C.W.M. The SWI/SNF complex in cancer - biology, biomarkers and
1104   therapy. *Nat Rev Clin Oncol* **2020**, *17*, 435-448, doi:10.1038/s41571-020-0357-3.

1105   105.   La Porte, A.; Cano, J.; Wu, X.; Mitra, D.; Kalpana, G.V. An Essential Role of
1106   INI1/hSNF5 Chromatin Remodeling Protein in HIV-1 Posttranscriptional Events and
1107   Gag/Gag-Pol Stability. *J Virol* **2016**, *90*, 9889-9904, doi:10.1128/JVI.00323-16.

1108   106.   Lesbats, P.; Botbol, Y.; Chevereau, G.; Vaillant, C.; Calmels, C.; Arneodo, A.;
1109   Andreola, M.L.; Lavigne, M.; Parissi, V. Functional coupling between HIV-1 integrase
1110   and the SWI/SNF chromatin remodeling complex for efficient in vitro integration into
1111   stable   nucleosomes.   *PLoS   Pathog*   **2011**,   *7*,   e1001280,
1112   doi:10.1371/journal.ppat.1001280.

107. Mahmoudi, T.; Parra, M.; Vries, R.G.; Kauder, S.E.; Verrijzer, C.P.; Ott, M.; Verdin, E. The SWI/SNF chromatin-remodeling complex is a cofactor for Tat transactivation of the HIV promoter. *J Biol Chem* **2006**, *281*, 19960-19968, doi:10.1074/jbc.M603336200.

108. Kaddis Maldonado, R.; Lambert, G.S.; Rice, B.L.; Sudol, M.; Flanagan, J.M.; Parent, L.J. The Rous sarcoma virus Gag Polyprotein Forms Biomolecular Condensates Driven by Intrinsically-disordered Regions. *J Mol Biol* **2023**, *435*, 168182, doi:10.1016/j.jmb.2023.168182.

109. Monette, A.; Niu, M.; Maldonado, R.K.; Chang, J.; Lambert, G.S.; Flanagan, J.M.; Cochrane, A.; Parent, L.J.; Mouland, A.J. Influence of HIV-1 Genomic RNA on the Formation of Gag Biomolecular Condensates. *J Mol Biol* **2023**, *435*, 168190, doi:10.1016/j.jmb.2023.168190.

110. Sabari, B.R.; Dall'Agnese, A.; Boija, A.; Klein, I.A.; Coffey, E.L.; Shrinivas, K.; Abraham, B.J.; Hannett, N.M.; Zamudio, A.V.; Manteiga, J.C.; et al. Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **2018**, *361*, doi:10.1126/science.aar3958.

111. Sabari, B.R.; Dall'Agnese, A.; Young, R.A. Biomolecular Condensates in the Nucleus. *Trends Biochem Sci* **2020**, *45*, 961-977, doi:10.1016/j.tibs.2020.06.007.

112. Cramer, P.; Bushnell, D.A.; Kornberg, R.D. Structural basis of transcription: RNA polymerase II at 2.8 angstrom resolution. *Science* **2001**, *292*, 1863-1876, doi:10.1126/science.1059493.

113. Acker, J.; Wintzerith, M.; Vigneron, M.; Kédinger, C. Primary structure of the second largest subunit of human RNA polymerase II (or B). *J Mol Biol* **1992**, *226*, 1295-1299, doi:10.1016/0022-2836(92)91071-v.

114. Chen, Y.; Kokic, G.; Dienemann, C.; Dybkov, O.; Urlaub, H.; Cramer, P. Structure of the transcribing RNA polymerase II-Elongin complex. *Nat Struct Mol Biol* **2023**, *30*, 1925-1935, doi:10.1038/s41594-023-01138-w.

115. Fianu, I.; Chen, Y.; Dienemann, C.; Dybkov, O.; Linden, A.; Urlaub, H.; Cramer, P. Structural basis of Integrator-mediated transcription regulation. *Science* **2021**, *374*, 883-887, doi:10.1126/science.abk0154.

116. Welsh, S.A.; Gardini, A. Genomic regulation of transcription and RNA processing by the multitasking Integrator complex. *Nat Rev Mol Cell Biol* **2023**, *24*, 204-220, doi:10.1038/s41580-022-00534-2.

117. Wagner, E.J.; Tong, L.; Adelman, K. Integrator is a global promoter-proximal termination complex. *Mol Cell* **2023**, *83*, 416-427, doi:10.1016/j.molcel.2022.11.012.

A.

B.

C.

	Cytoplasm	Nucleoplasm	Chr 150 mM	Chr 500 mM
RSV Gag	42.5 ± 2.6%	39.1 ± 1.6%	12.6 ± 4.5%	5.8 ± 1.6%
RSV Gag.L219A	20.3 ± 0.8%	46.7 ± 6.2%	16.4 ± 5.2%	16.7 ± 2.8%
RSV Gag.ΔNC	66.8 ± 6.6%	32.1 ± 5.9%	1.1 ± 0.7%	0.0 ± 0.0%

1149

1150

1151

1152

A) Common Among All

- ■ Cell Assembly & Organization
- ■ Cell Cycle
- ■ Cell Death & Survival
- ■ Cell Proliferation
- ■ Cell Signaling
- ■ Development & Function
- ■ DNA Repl., Recomb, & Repair
- ■ Gene Expression
- ■ Metabolism
- ■ Other
- ■ Protein Synthesis, Modification & Repair
- ■ RNA Modification & Repair
- ■ Transport/Movement

B) Rice

C) Engeland 2011 *et al*

D) Engeland 2014 *et al*

E) LeSage *et al*

F) Li *et al*

G) Ritchie *et al*

H) Jager *et al*

1153

1154

A.

| | RSV Gag-GFP | FLAG-Med26 | Merge | Colocalization |

B.

Manders' Overlap Coefficient

M2 (Gag ∩ Med26)

M1 (Med26 ∩ Gag)

****

0.0   0.2   0.4   0.6   0.8   1.0

C.

IP: α-RSV Gag

| Input | FT | Beads | Eluate | WB: |

250
130
100
70

Med26

250
130
100
70

Gag-Pol

Gag

1155

1156

1157

1158

1159