Eukaryotic genomic data uncover an extensive host range of mirusviruses

Hongda Zhao, Lingjie Meng, Hiroyuki Hikida, Hiroyuki Ogata*

Institute for Chemical Research, Kyoto University, Uji, Kyoto, Japan

* Corresponding author: Hiroyuki Ogata (ogata@kuicr.kyoto-u.ac.jp)

Highlights

• Mirusvirus signals detected in genomic data from eight eukaryotic supergroups.

• Habits of putative mirusvirus hosts not limited to marine environments.

• Major capsid sequences from these assemblies show new mirusviral lineages.

• Three circular mirusvirus genomes were identified.

Summary

A recent metagenomic study has revealed a novel group of viruses designated

mirusviruses, which are proposed to form an evolutionary link between two

double-stranded DNA virus realms, *Varidnaviria* and *Duplodnaviria*. Metagenomic

data suggest that these viruses infect marine microeukaryotes, but their host range

remains largely unknown. In this study, we investigated the presence of mirusvirus

marker genes in 1,901 eukaryotic assemblies, mainly derived from unicellular

eukaryotes, to identify potential hosts of mirusviruses. Mirusvirus marker sequences

were identified in 1,348 assemblies spanning 284 eukaryotic genera across eight

supergroups. The habitats of the putative mirusvirus hosts included not only marine

but also other diverse environments. Among the major capsid protein (MCP) sequence

signals in the genome assemblies, we identified 85 sequences that showed high

sequence and structural similarities to reference mirusvirus MCPs. A phylogenetic

analysis of these sequences revealed their distant evolutionary relationships with the

seven previously reported mirusvirus clades. Most of the scaffolds with these MCP

sequences encoded multiple mirusvirus homologs, underscoring the impact of

mirusviral infection on the evolution of the host genome. We also identified three

circular mirusviral genomes within the genomic data of *Schizochytrium* sp. and

Ostreobium quekettii. Overall, mirusviruses probably infect a wide spectrum of

eukaryotes and are more diverse than previously reported.

Keywords: virus–host relationship, mirusvirus, major capsid protein, protist,

endogenous virus

Introduction

Viruses pervade diverse environments across the Earth and play crucial

ecological and evolutionary roles.[1–3] A recent metagenomic analysis identified a

previously unrecognized but diverse group of double-stranded (ds) DNA viruses,

designated mirusviruses, that are abundant, widespread, and active in the global

marine ecosystem.[4] Based on their virion morphogenesis genes (e.g., HK97-fold

MCP), mirusviruses are proposed to form the phylum '*Mirusviricota*' in the realm *Duplodnaviria*, which includes herpesviruses and caudoviruses. However, a large number of genes encoded by mirusviruses, including informational genes, are more closely related to homologs in nucleocytoviruses. Nucleocytoviruses belong to another dsDNA viral realm, *Varidnaviria*, and are known to play important roles in marine ecosystems. The genomic similarities between mirusviruses and nucleocytoviruses suggest an evolutionary interplay between the two distinct groups of viruses and their similar habitats.

Viral groups have different host ranges. Nucleocytoviruses infect diverse eukaryotes, including protists, green algae, and animals.[5] Within *Duplodnavira*, caudoviruses infect a broad spectrum of prokaryotes,[6,7] whereas herpesviruses have a relatively narrow host range, limited to animals.[8,9] Until the discovery of mirusviruses, no duplodnaviruses were known to infect early-branching eukaryotes, such as protists. Mirusviruses were the first group of duplodnaviruses suggested to infect unicellular eukaryotes. This suggestion was based on the horizontal gene transfer of heliorhodopsin genes between mirusviruses and green algae, and the fact that mirusvirus signals were detected in the metagenomes and metatranscriptomes derived from size fractions corresponding to unicellular planktonic eukaryotes.[4] Therefore, the discovery of mirusviruses was considered to fill the host gap in the duplodnaviruses (between prokaryotes and animals). However, the evidence of mirusvirus hosts was limited and the taxonomic breadth of their hosts remained poorly investigated.

Two recent studies investigating the viral signals in eukaryotic genomes revealed

some potential hosts of mirusviruses. Specifically, endogenized mirusviral genomes

have been detected in the thraustochytrids species *Aurantiochytrium limacinum* and

Hondaea fermentalgiania,[10] and the green algal species *Cymbomonas*

tetramitiformis.[11] Notably, an additional circular mirusvirus-like genome (probably in

the form of an episome) was identified in the *A. limacinum* genome assembly,

suggesting an episomal form of mirusvirus in the host cells.[10] In general, the viral

sequences within eukaryotic assemblies can be categorized into three types:

transferred genes, integrated viral genome fragments, and free viral genomes.[12–14] The

previously reported endogenized mirusviral sequences and the circular genome

correspond to the latter two types, respectively. These three types of evidence strongly

suggest that eukaryotes containing viral signals have acted as the hosts of viruses.

In the present study, we systematically screened 1,901 genome assemblies from a

diverse group of predominantly unicellular eukaryotes for mirusvirus signals. Of the

318 eukaryotic genera analyzed, 284 contained mirusvirus signals. The mirusviral

MCP sequences identified in this study formed distinct phylogenetic clades, separate

from previously reported mirusviral MCP sequences derived from marine

environments. Moreover, three circular mirusviral genomes encoding a nearly

complete set of marker genes were identified. This study suggests that the host range

of the mirusviruses is broad, and that the mirusviruses display previously

unrecognized diversity.

Results

Mirusviral marker sequences detected in a wide range of eukaryotes

The dataset for this study was collected from GenBank and comprised 1,901 eukaryotic genomic assemblies. These assemblies spanned over 318 minor lineages (mostly at the genus level, and are hereafter referred to as 'genera' for simplicity) and 16 major lineages of eukaryotic organisms (mostly at the phylum level or higher; Supplementary Table S1, Data S1), and cover eight of the nine eukaryotic supergroups.[15] In these assemblies, we identified nearly 118 million open reading frames (ORFs). Of these protein sequences, 6,659 showed significant sequence similarities (E-value $< 10^{-5}$) to the hidden Markov models (HMMs) of five selected mirusviral marker sequences: MCP, capsid triplex subunit 1 (Triplex1), capsid triplex subunit 2 (Triplex2), capsid portal protein (Portal), and capsid maturation protease (Maturation). To ensure their specificity to the mirusviruses, we aligned these sequences with other duplodnavirus (caudovirus and herpesvirus) sequences and HMMs in the PFAM database, and excluded possible false positives. In this way, we identified 6,042 marker sequences (541 MCP, 1,202 Portal, 509 Triplex1, 625 Triplex2, and 3,165 Maturation) that are specifically related to mirusvirus counterparts in 1,348 eukaryotic genome assemblies (71% of the analyzed assemblies).

The 1,348 eukaryotic genome assemblies containing viral marker sequences

included 284 genera spanning 15 major lineages and eight eukaryotic supergroups

(Data S2). Specifically, MCP was detected in 98 genera, Portal in 171 genera,

Triplex1 in 114 genera, Triplex2 in 135 genera, and Maturation in 249 genera. Of

these 284 genera, we selected 90 genera that showed strong signals for mirusviral

marker sequences based on the criterion that the assemblies in the genus contained

four or five distinct marker sequences (Fig. 1). Of the 6,042 marker sequences

identified, 4,365 belonged to these 90 genera. These genera were spread across 11

major lineages and seven supergroups of eukaryotes.

Fig. 1 Eukaryotic genera showing four or five different mirusvirus markers in

We investigated the sequence similarities between all 6,042 mirusviral marker sequences detected in the eukaryotic assemblies and reference marker sequences encoded in previously described mirusvirus genomes derived from marine metagenomes.[4] Most of the marker sequences shared low sequence similarity scores with the reference markers, and the median bit-score calculated with hmmscan ranged from 10 to 15 for all five marker genes (Fig. 2A). The median lengths of the marker sequences in the eukaryotic assemblies were smaller than those of the reference markers and ranged from 100 to 200 amino acids (Fig. 2B). These features suggest that many of the marker sequences detected were decaying nonfunctional genes (Fig. 2C). Nevertheless, there were also marker sequences of normal length but that shared low bit scores with reference sequences, implying the existence of diversified viral lineages.

Fig. 2 Lengths of detected marker sequences and their similarities to reference sequences. (A) Boxplots of bit-scores against five reference marker genes. **(B)** Boxplots of the length distributions of the detected marker sequences. Yellow stars represent the median lengths of the reference mirusvirus orthogroups. **(A–B)** Red diamonds represent the outlier points. **(C)** Relationships between length and bit-score for each marker.

MCPs from eukaryotic assemblies form clades distinct from those of known mirusviruses

We next investigated the evolutionary relationships between the mirusvirus signals in the eukaryotic genome assemblies and previously described mirusviruses in

marine metagenomes, focusing on MCP. This protein is the most important viral

structural component and is critical for viral taxonomic classification.[1] Furthermore,

MCP sequences are reported to represent an evolutionary path for the mirusviruses,

similar to informational genes.[4] To minimize the interference degraded genes impose

on phylogenetic analyses, we specifically filtered MCP marker sequences suitable for

evolutionary analyses as follows. First, from the initial 541 MCP sequences identified

in eukaryotic assemblies, we selected 262 sequences suitable for structural prediction.

The selection was based on the absence of ambiguous amino acids (due to sequencing

quality) and a length of 200–1,500 amino acids. We predicted the three-dimensional

(3D) structures of these 262 MCPs and then compared their structural similarities with

the marine mirusviral MCPs. Of the 262 structural models, 170 showed a template

modeling score (TM-score) of > 0.5 (a criterion for the potential same fold[16,17]) with

at least one reference mirusviral MCP, supporting their structural similarities.

We further refined our selection of MCPs from eukaryotic assemblies for

phylogenetic analysis based on the conservation of functional domains. The

HK97-fold is an important shared feature of duplodnaviral MCP structures. There are

multiple conserved elements in the HK97-fold, including the axial domain

(A-domain), peripheral domain (P-domain), extended loop (E-loop), and N-terminal

arm (N-arm).[18] The floor domain, which includes the last three elements, exists in all

HK97 MCPs. We used the E-loop, a long two-stranded β-sheet hairpin, as an indicator

of the presence of the floor domain. Of the 170 sequences with high structural

similarities to reference mirusviral MCPs, 150 were found to contain the two-stranded β-sheet hairpin with a β-strand longer than 10 amino acids on each side.

Finally, we compared the structures of these 150 MCP sequences with those of other representative duplodnaviral MCPs (from caudoviruses and herpesviruses). Of these 150 MCP sequences, 85 showed higher structural similarities to mirusviruses than to other viruses (Supplementary Fig. S1). Therefore, we considered these 85 MCP sequences as the most appropriate set of sequences for the subsequent phylogenetic analysis. These 85 MCP sequences were derived from 15 assemblies (14 species), including organisms from Alveolata, Amoebozoa, Chlorophyta, Cryptophyta, Rhizaria, and Stramenopiles (Table 1). Most of the homologs showed a bit-score of > 100 and a TM-score to the reference mirusvirus MCPs of > 0.7 (Supplementary Fig. S2). However, homologs from Rhizaria generally displayed low bit-scores, but a wide range of TM-scores to the reference mirusviral MCPs.

Table 1: Species and assemblies containing the 85 MCP homologs.

Species	Major lineage	Assembly	Location of isolation	Environment	Clade
Cymbomonas tetramitiformis	Chlorophyta	GCA_001247695.1	English Channel	Sea	E01
Aurantiochytrium sp.	Stramenopiles	GCA_001462505.1	Madeira	Sea	E02
Aurantiochytrium sp.	Stramenopiles	GCA_003116975.1	Hiroshima	Sea	E02
Schizochytrium sp.	Stramenopiles	GCA_004764695.1	missing	Sea	E02
Parietichytrium sp.	Stramenopiles	GCA_012862575.1	Okinawa	Sea	E02
Hondaea fermentalgiana	Stramenopiles	GCA_014084085.1	Mayotte	Mangrove	E02

Aurantiochytrium acetophilum	Stramenopiles	GCA_004332575.1	Biscayne Bay	Mangrove	E02
Ostreobium quekettii	Chlorophyta	GCA_905146915.1	Catanduanes Island	Sea	E02
Euplotes weissei	Alveolata	GCA_021440005.1	Qingdao	Sea	E02
Cryptophyta sp.	Cryptophyta	GCA_026770585.1	Northern Baffin Bay	Sea	E03
Uncultured Cryptomonadales	Cryptophyta	GCA_947538865.1	Římov Reservoir	Fresh water	E03
Guillardia theta	Cryptophyta	GCA_000315625.1	Connecticut	Sea	E03
Acanthamoeba pearcei	Amoebozoa	GCA_000826505.1	missing	missing	E04
Paulinella micropora	Rhizaria	GCA_009731375.1	Ibaraki	Fresh water	E05
Paulinella micropora	Rhizaria	GCA_019918135.1	Chungnam	Fresh water	E05

After all filters were applied, we reconstructed a phylogenetic tree comprising

these 85 MCP homologs, together with 79 reference mirusviral MCPs derived from

marine metagenomes (Fig. 3). The topology of the subtree of the reference mirusviral

MCPs (clades M01 to M07) was generally consistent with a previous report.[4] The

MCP homologs detected in eukaryotic assemblies were not grouped within the

reference mirusvirus clades. Instead, they formed five distinct clades, E01 to E05,

which were distantly related to the reference MCPs. The representative sequences for

the individual clades (i.e., the longest homolog from each clade) displayed predicted

3D structures similar to those of the reference MCP, including the floor domain (Fig.

3). Notably, clades E01–E03 had an additional conserved antiparallel β-strand

adjacent to the E-Loop.

The newly identified clades corresponded to distinct eukaryotic lineages, except

Clade E02. Clade E01 consisted of nine homologs from a single genomic assembly of

the green algal species *C. tetramitiformis*.[11] Clade E02 was the only clade consisting

of homologs (n = 13) from different organisms, including *Euplotes weissei*

(Alveolata), *Ostreobium quekettii* (Chlorophyta), and six thraustochytrids

(Stramenopiles) members.[10] Clade E03 consisted of 18 homologs from three

Cryptophyta assemblies. Clade E04 contained a single homolog from an assembly of

Acanthamoeba pearcei. Clade E05 consisted of 44 homologs from two different

strains of *Paulinella micropora* (Rhizaria).

Fig. 3 Maximum-likelihood phylogenetic tree of MCPs. Different clades are

indicated with different colors: blue represents the reference mirusviral MCPs.

Predicted protein structures of the longest homolog within each clade are displayed

around the tree. β-sheets are colored orange. Above the structure, the clade number is

210 Existence of viral regions and three circular viral genomes

We also investigated the genomic context around the 85 most highly conserved

MCP homologs. These homologs were found in 83 contigs or scaffolds (hereafter

referred to as 'scaffolds' for simplicity). The lengths of these scaffolds ranged from

1.2 kilobases to 19.7 megabases. We analyzed all the predicted ORFs on these

scaffolds and found that most of the scaffolds encoded many homologs of mirusviral

genes, except the very short scaffolds (Fig. 4). Most of the scaffolds displayed a

higher predicted ORF density than the average level within the same assembly

(Supplementary Fig. S3).

Fig. 4 Genomic maps of the 83 scaffolds. Scaffolds were divided into three groups according to their lengths. The predicted ORFs with mirusvirus orthogroup best hits are colored red. The predicted ORFs with PFAM best hits are colored blue. The tip of the triangle represents the direction of the ORF.

From database records at the National Center for Biotechnology Information (NCBI), we found that CM015707.1 and CM015708.1 are circular contigs from *Schizochytrium* sp. (Stramenopiles), with lengths of 289 kilobases and 275 kilobases, respectively. When we examined the remaining 81 scaffolds, we found that CAJHUC010000281.1 (from *Ostreobium quekettii*, Chlorophyta) is also likely to be a circular contig (325 kilobases) (Supplementary Fig. S4). These circular contigs

encode 3–5 mirusvirus markers (of the five selected for this study) and additional

functionally important proteins, such as terminases, DNA/RNA polymerases, and

heliorhodopsins (Fig. 5). Notably, the three circular genomes lack RNA polymerase

subunit B, even though RNA polymerase subunit B is the most commonly detected

gene and is conserved in 89% of the reported mirusviral genomes. In contrast, the

circular genomes encoded two copies of RNA polymerase subunit A. These circular

contigs belonged to Clade E02. A family B DNA polymerase (PolB)-based

phylogenetic analysis confirmed the close relationships between these three circular

genomes (Supplementary Fig. S5). No split genes were detected in these circular

contigs, indicating that most genes were intact and not pseudogenized.

Fig. 5 Three circular mirusviral genomes recovered from eukaryotic assemblies.

The five markers used in this study are colored brown. Functionally annotated genes

are colored orange. Other predicted genes are colored yellowish.

Discussion

     Mirusviruses are reported to be a deep-branching diverse group of dsDNA viruses,

but previous studies have provided only limited information on their potential hosts in

marine environemnts.[4,10,11] In the present study, we detected mirusviral maker

sequence signals in a large number of eukaryotic (mostly unicellular) genome assemblies (1,348 assemblies; 71% of those analyzed). These eukaryotes included nearly 90% of the eukaryotic genera analyzed and 15 of the 16 major lineages analyzed (Data S2). These potential mirusvirus hosts included not only organisms living in marine environments (e.g., *Aurantiochytrium*, *Euplotes*) but also those living in other environments (e.g., freshwater: *Paulinella, Yamagishiella*; soil: *Dictyostelium*, *Physarum*; parasites of animals or plants: *Trypanosoma*, *Phytophthora*). Therefore, mirusviruses probably infect a broad spectrum of eukaryotes in various types of habitat.

Many of the mirusviral marker sequences identified in eukaryotic assemblies showed low sequence similarity scores to the reference mirusviral sequences (Fig. 2A). This sequence divergence from the reference sequences is probably attributable to the accumulation of mutations (e.g., pseudogenization[19]) after their ancient insertion into the host genomes. This is supported by the existence of abnormally short ORFs in the mirusviral marker sequences (Fig. 2B). Despite the presence of decaying genes, 90 genera displayed a comprehensive set of mirusviral marker genes (i.e., four or five markers). Such a high level of marker gene conservation may reflect the relatively recent insertion of the mirusviral genomes into the eukaryotic genomes (Fig. 1).

Our analysis also revealed a wide range of sequence similarity scores, even for ORFs in the normal size range (Fig. 2C). This suggests that factors beyond

pseudogenization have contributed to the low sequence similarities between the newly

detected sequences and the reference sequences. We analyzed the phylogeny of 85

selected MCP sequences that showed strong similarities (both in sequence and

structure) to reference mirusviral MCPs and were thus considered to represent

relatively recent insertions (Supplementary Fig. S2). These homologs were only

distantly related to the reference mirusviral MCPs and formed five distinct clades (Fig.

3). Therefore, these MCP marker sequences probably originated from viral lineages

that are distinct from the previously described mirusviral lineages.

Current research into mirusviruses is based on genomic data from environmental

samples or eukaryotes, because no mirusvirus has yet been isolated. Despite their

widespread presence in marine environments, the uncertainty regarding their original

hosts poses challenges for isolation studies. In this study, we identified organisms

showing strong and fresh mirusvirus signals. Notably, a three-stranded antiparallel

β-sheet was observed in clade E01, E02, and E03 MCP homologs (Fig. 3). This

structure consists of two β-strands in the E-loop at the N-terminus and an additional

β-strand at the C-terminus,[1] and its presence indicates the high structural integrity of

the MCP. A previous study reported a circular mirusviral genome in a thraustochytrids

species.[10] We detected three circular mirusviral genomes, without evidence of

pseudogenization, in another thraustochytrids species and a green algal species (Fig.

5). The circular status of the genomes suggests that these mirusviruses either

latently[10,20] or persistently[21] infect their hosts. Eukaryotes harboring circular

mirusviral genomes or high-integrity MCP sequences (E01–E03 in Table 1) represent

promising candidates for the isolation of mirusviruses because their infections are

probably recent or on-going events.

Viral genes are known to contribute to the critical evolution of their hosts.[22,23] The

invasion of eukaryotic host genomes by DNA viral genomes is widespread among

unicellular eukaryotes.[13] In particular, large dsDNA viruses have been reported to

form giant viral endogenous elements within their host genomes.[24,25] The detection of

mirusvirus signals in various eukaryotes indicates that mirusviruses are not an

exception to this phenomenon (Data S1, S2). We identified multiple mirusvirus

homologs in the scaffolds that harbored mirusviral MCPs (Fig. 4), suggesting that the

genomic regions are giant endogenous viral elements originating from mirusvirus

genomes. It has been suggested that giant viral endogenous elements confer unique

capabilities upon their hosts (usually unicellular eukaryotes) by providing various

genes of viral origin,[24] in a similar way to endogenous viral elements in higher

eukaryotes.[26,27] Given the extensive presence of mirusvirus-like genomic regions in

eukaryotic genomes, mirusviruses may have contributed to the genomic innovation of

their hosts.[28]

Methods

Unicellular eukaryotic genome assembly data

We compiled the unicellular eukaryotic genome assembly data by retrieving all

the relevant eukaryotic genome assemblies from the GenBank database (as of June

2023). Assemblies from Fungi (NCBI Taxonomy ID 4751), Metazoa (NCBI

Taxonomy ID 33208), and Streptophyta (NCBI Taxonomy ID 35493) were excluded.

A total of 1,901 assemblies were collected for subsequent analysis. Organisms that

were not classified at the genus level were treated as distinct genera. To identify

mirusvirus-originating sequences within these assemblies, gene calling was performed

on all compiled data with the Prodigal v2.6.3 software, with the parameter '-p meta'.[29]

Predicted sequences shorter than 20 amino acids or longer than 4,000 amino acids

were discarded.

Creation of mirusvirus marker protein models

The reference metagenome-assembled mirusvirus genomes were obtained from a

previous study.[4] Gene calling was performed with Prodigal, and orthologous groups

were then generated with Orthofinder v2.5.2.[30] The core gene orthologous groups

were annotated with a BLASTp search against the RefSeq database with Diamond

v2.1.8,[31] and subsequent manual curation. For the five virion module protein markers,

redundant sequences with > 90% sequence identity were removed from the five

orthologous groups with cd-hit v4.8.1 and the parameter '-c 0.9'.[32] The HMMs of the

five markers were built with HMMER v3.3.2. [32]

Detection mirusvirus marker sequences in eukaryotic assemblies

To detect mirusviral marker sequences within the eukaryotic assemblies, we used

HMMs corresponding to five marker genes. We screened our predicted protein

database, retaining hits with an E-value $< 10^{-5}$ as determined with HMMER. To

ensure the specificity of our results for mirusviruses, we established additional

controls to exclude similar sequences from other viruses. First, we downloaded

caudovirus and herpesvirus protein sequences from NCBI Virus (as of October 26,

2023). We then curated a dataset of homologs for the selected marker genes (for

caudoviruses: MCP, Portal, Maturation; for herpesviruses: MCP, Triplex1, Triplex2,

Portal, Maturation) using keyword filtering based on the NCBI annotations. Sequence

redundancy was removed with cd-hit, with a cut-off of 90% sequence identity. We

then constructed HMMs for each of the caudovirus and herpesvirus proteins and used

these models to examine our mirusvirus hits. Any protein sequences from the

eukaryotic assemblies that demonstrated a lower E-value when matched with the

caudovirus or herpesvirus protein models than when compared with the mirusvirus

models were excluded from further analysis. We also used hmmscan to scan the

remaining mirusvirus hits against HMMs in the PFAM database (as of November 7,

2023),[33] and discarded hits with lower E-values against any PFAM HMM.

3D structural analyses of MCP homologs

Protein 3D structural predictions were made with AlphaFold v2.3.2 (-t

2023-11-14).[34] For structural comparisons with known HK97-fold MCPs, we also

predicted the 3D structures of reference MCP sequences from mirusviruses,

herpesviruses, and caudoviruses. Sequences shorter than 200 amino acids or with >

70% sequence identity to other sequences in the same viral group were removed from

the reference MCP sequences. In total, 37 herpesvirus MCP models and 252

caudovirus MCP models were referenced. Structural similarities between proteins

were calculated with Foldseek v6-29e2557 with the program 'easy-search'.[35] Protein

structures were visualized with ChimeraX v1.7.[36]

Phylogenetic analysis

Multiple-sequence alignments of MCP and PolB sequences longer than 200

amino acids were generated with Clustal-Omega v1.2.4.[37] We trimmed the alignments

with trimAl v1.4.1, with the parameter '-gt 0.1'.[38] Maximum-likelihood phylogenetic

trees were constructed with IQ-TREE v2.2.2.6,[39] with 1,000 ultrafast bootstrap

replicates. Models were selected with ModelFinder.[40] The phylogenetic trees were

visualized with iTOL v6.8.1.[41] For the phylogenetic tree of PolB, we collected all

mirusviral PolB homologous sequences on those 83 scaffolds and also used

eukaryotic reference sequences from a previous study.[42]

Annotation of contigs and scaffolds

We identified mirusvirus orthologous groups that were shared by more than 10

reference mirusviral metagenome-assembled genomes. Using the HMMs of these

orthologous groups as the queries, we scanned all 83 scaffolds (E-value $< 10^{-3}$). We

also used PFAM models as queries to scan these scaffolds (E-value $< 10^{-3}$). For the

predicted genes that showed similarities to both mirusvirus orthologous groups and

PFAM models, the annotation result was based on the hit with the lower E-value. To

determine whether these scaffolds were likely to be circular, we performed a

similarity comparison of different parts of the same sequence with NCBI web-based

versions of BLASTn and DigAlign to determine whether the extremities of the

sequence showed the same sequence (https://www.genome.jp/digalign/).[43] For the

three circular mirusvirus contigs, we re-predicted their ORFs with the default

parameters of Prodigal. The genomic maps of the circular contigs were generated with

the Python Package 'DNA features viewer',[44] and the annotations were based on the

best hits of the predicted ORFs to the mirusvirus orthogroups (E-value $< 10^{-3}$). To

investigate pseudogenization, we used Diamond BLASTp to query all ORFs derived

from the circular genomes against the RefSeq database, using the parameter '-k 3 -e

10'. We looked for adjacent ORFs showing similarity to different parts of the same

reference sequence as potential signals of pseudogenization.

Acknowledgments

We thank Junyi Wu and Ruixuan Zhang for valuable discussions. This work was

supported, in part, by JSPS/KAKENHI (nos 22H00384 and 19H05667).

Computational time was provided by the Supercomputer System, Institute for

Chemical Research, Kyoto University (Kyoto, Japan). Molecular graphics and

analyses were performed with UCSF ChimeraX, developed by the Resource for

Biocomputing, Visualization, and Informatics at the University of California, San

Francisco (CA, USA), with support from National Institutes of Health

(R01-GM129325) and the Office of Cyber Infrastructure and Computational Biology,

National Institute of Allergy and Infectious Diseases (MD, USA). We thank Edanz

(https://jp.edanz.com/ac) for editing a draft of this manuscript.

Author Contributions

HZ designed the work, performed all the analyses, and wrote the initial draft of

the manuscript. LM contributed to the data preparation and assisted with the analyses.

HH and HO supervised HZ and contributed to the finalization of the manuscript. All

authors have read and approved the final version of the manuscript.

Declaration of Interests

The authors declare no competing interests.

Data and Code Availability

• The annotations, predicted ORFs, HMM files, alignments, predicted structure

files, and tree files are available through GenomeNet

(https://www.genome.jp/ftp/db/community/Mirusvirus_host_range/).

• This paper does not report original code.

• Any additional information required to reanalyze the data reported in this paper is

available from the authors upon request.

References

1.   Krupovic, M., and Koonin, E.V. (2017). Multiple origins of viral capsid proteins from cellular ancestors.

Proceedings of the National Academy of Sciences *114*, E2401–E2410. 10.1073/pnas.1621061114.

419 2. Forterre, P. (2006). The origin of viruses and their possible roles in major evolutionary transitions. Virus
420 Research *117*, 5–16. 10.1016/j.virusres.2006.01.010.

421 3. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2015). Origins and evolution of viruses of eukaryotes: The
422 ultimate modularity. Virology *479–480*, 2–25. 10.1016/j.virol.2015.02.039.

423 4. Gaïa, M., Meng, L., Pelletier, E., Forterre, P., Vanni, C., Fernandez-Guerra, A., Jaillon, O., Wincker, P.,
424 Ogata, H., Krupovic, M., et al. (2023). Mirusviruses link herpesviruses to giant viruses. Nature *616*, 783–789.
425 10.1038/s41586-023-05962-4.

426 5. Schulz, F., Abergel, C., and Woyke, T. (2022). Giant virus biology and diversity in the era of
427 genome-resolved metagenomics. Nat Rev Microbiol, 1–16. 10.1038/s41579-022-00754-5.

428 6. Pagaling, E., Haigh, R.D., Grant, W.D., Cowan, D.A., Jones, B.E., Ma, Y., Ventosa, A., and Heaphy, S.
429 (2007). Sequence analysis of an Archaeal virus isolated from a hypersaline lake in Inner Mongolia, China. BMC
430 Genomics *8*, 410. 10.1186/1471-2164-8-410.

431 7. Piel, D., Bruto, M., Labreuche, Y., Blanquart, F., Goudenège, D., Barcia-Cruz, R., Chenivesse, S., Le Panse,
432 S., James, A., Dubert, J., et al. (2022). Phage–host coevolution in natural populations. Nat Microbiol *7*, 1075–1086.
433 10.1038/s41564-022-01157-1.

434 8. Bandín, I., and Dopazo, C.P. (2011). Host range, host specificity and hypothesized host shift events among
435 viruses of lower vertebrates. Veterinary Research *42*, 67. 10.1186/1297-9716-42-67.

436 9. Rosani, U., Gaia, M., Delmont, T.O., and Krupovic, M. (2023). Tracing the invertebrate herpesviruses in the
437 global sequence datasets. Frontiers in Marine Science *10*.

438 10. Collier, J.L., Rest, J.S., Gallot-Lavallée, L., Lavington, E., Kuo, A., Jenkins, J., Plott, C., Pangilinan, J.,
439 Daum, C., Grigoriev, I.V., et al. (2023). The protist Aurantiochytrium has universal subtelomeric rDNAs and is a
440 host for mirusviruses. Current Biology. 10.1016/j.cub.2023.10.009.

441 11. Gyaltshen, Y., Rozenberg, A., Paasch, A., Burns, J.A., Warring, S., Larson, R.T., Maurer-Alcalá, X.X.,
442 Dacks, J., Narechania, A., and Kim, E. (2023). Long-Read-Based Genome Assembly Reveals Numerous
443 Endogenous Viral Elements in the Green Algal Bacterivore Cymbomonas tetramitiformis. Genome Biol Evol *15*,
444 evad194. 10.1093/gbe/evad194.

445 12. Irwin, N.A.T., Pittis, A.A., Richards, T.A., and Keeling, P.J. (2022). Systematic evaluation of horizontal
446 gene transfer between eukaryotes and viruses. Nat Microbiol *7*, 327–336. 10.1038/s41564-021-01026-3.

447 13. Bellas, C., Hackl, T., Plakolb, M.-S., Koslová, A., Fischer, M.G., and Sommaruga, R. (2023). Large-scale
448 invasion of unicellular eukaryotic genomes by integrating DNA viruses. Proceedings of the National Academy of
449 Sciences *120*, e2300465120. 10.1073/pnas.2300465120.

450 14. Zhao, H., Zhang, R., Wu, J., Meng, L., Okazaki, Y., Hikida, H., and Ogata, H. (2023). A 1.5-Mb continuous
451 endogenous viral region in the arbuscular mycorrhizal fungus Rhizophagus irregularis. Virus Evolution *9*, vead064.
452 10.1093/ve/vead064.

453 15. Burki, F., Roger, A.J., Brown, M.W., and Simpson, A.G.B. (2020). The New Tree of Eukaryotes. Trends in
454 Ecology & Evolution *35*, 43–55. 10.1016/j.tree.2019.08.008.

455 16. Zhang, Y., and Skolnick, J. (2005). TM-align: a protein structure alignment algorithm based on the TM-score.
456 Nucleic Acids Res *33*, 2302–2309. 10.1093/nar/gki524.

457 17. Xu, J., and Zhang, Y. (2010). How significant is a protein structure similarity with TM-score = 0.5?
458 Bioinformatics *26*, 889–895. 10.1093/bioinformatics/btq066.

459 18. Duda, R.L., and Teschke, C.M. (2019). The amazing HK97 fold: versatile results of modest differences. Curr
460 Opin Virol *36*, 9–16. 10.1016/j.coviro.2019.02.001.

19. Katzourakis, A., and Gifford, R.J. (2010). Endogenous Viral Elements in Animal Genomes. PLoS Genet *6*,

e1001191. 10.1371/journal.pgen.1001191.

20. Cohen, J.I. (2020). Herpesvirus latency. J Clin Invest *130*, 3361–3369. 10.1172/JCI136225.

21. Blanc-Mathieu, R., Dahle, H., Hofgaard, A., Brandt, D., Ban, H., Kalinowski, J., Ogata, H., and Sandaa,

R.-A. (2021). A Persistent Giant Algal Virus, with a Unique Morphology, Encodes an Unprecedented Number of

Genes Involved in Energy Metabolism. Journal of Virology *95*, e02446-20. 10.1128/JVI.02446-20.

22. Ochman, H., Lawrence, J.G., and Groisman, E.A. (2000). Lateral gene transfer and the nature of bacterial

innovation. Nature *405*, 299–304. 10.1038/35012500.

23. Koonin, E.V., and Krupovic, M. (2018). The depths of virus exaptation. Current Opinion in Virology *31*, 1–8.

10.1016/j.coviro.2018.07.011.

24. Moniruzzaman, M., Weinheimer, A.R., Martinez-Gutierrez, C.A., and Aylward, F.O. (2020). Widespread

endogenization of giant viruses shapes genomes of green algae. Nature *588*, 141–145.

10.1038/s41586-020-2924-2.

25. Moniruzzaman, M., Erazo-Garcia, M.P., and Aylward, F.O. (2022). Endogenous giant viruses contribute to

intraspecies genomic variability in the model green alga Chlamydomonas reinhardtii. Virus Evolution *8*, veac102.

10.1093/ve/veac102.

26. Fujino, K., Horie, M., Honda, T., Merriman, D.K., and Tomonaga, K. (2014). Inhibition of Borna disease

virus replication by an endogenous bornavirus-like element in the ground squirrel genome. Proceedings of the

National Academy of Sciences *111*, 13175–13180. 10.1073/pnas.1407046111.

27. Suzuki, Y., Frangeul, L., Dickson, L.B., Blanc, H., Verdier, Y., Vinh, J., Lambrechts, L., and Saleh, M.-C.

(2017). Uncovering the Repertoire of Endogenous Flaviviral Elements in Aedes Mosquito Genomes. Journal of

Virology *91*, 10.1128/jvi.00571-17. 10.1128/jvi.00571-17.

28. Aylward, F.O. (2023). Microbiology: The curious case of the mysterious mirusvirus. Current Biology *33*,

R1234–R1235. 10.1016/j.cub.2023.10.037.

29. Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010). Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics *11*, 119.

10.1186/1471-2105-11-119.

30. Emms, D.M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics.

Genome Biology *20*, 238. 10.1186/s13059-019-1832-y.

31. Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment using DIAMOND. Nat

Methods *12*, 59–60. 10.1038/nmeth.3176.

32. Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLOS Computational Biology *7*, e1002195.

10.1371/journal.pcbi.1002195.

33. Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger, A., Hetherington, K.,

Holm, L., Mistry, J., et al. (2014). Pfam: the protein families database. Nucleic Acids Research *42*, D222.

10.1093/nar/gkt1223.

34. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R.,

Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*,

583–589. 10.1038/s41586-021-03819-2.

35. van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M., Söding, J., and

Steinegger, M. (2023). Fast and accurate protein structure search with Foldseek. Nat Biotechnol, 1–4.

10.1038/s41587-023-01773-0.

36.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H., and Ferrin, T.E. (2023).

UCSF ChimeraX: Tools for structure building and analysis. Protein Science *32*, e4792. 10.1002/pro.4792.

37.   Sievers, F., and Higgins, D.G. (2018). Clustal Omega for making accurate alignments of many protein

sequences. Protein Sci *27*, 135–145. 10.1002/pro.3290.

38.   Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: a tool for automated alignment

trimming in large-scale phylogenetic analyses. Bioinformatics *25*, 1972–1973. 10.1093/bioinformatics/btp348.

39.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and Lanfear,

R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era.

Molecular Biology and Evolution *37*, 1530–1534. 10.1093/molbev/msaa015.

40.   Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and Jermiin, L.S. (2017). ModelFinder:

fast model selection for accurate phylogenetic estimates. Nat Methods *14*, 587–589. 10.1038/nmeth.4285.

41.   Letunic, I., and Bork, P. (2007). Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display

and annotation. Bioinformatics *23*, 127–128. 10.1093/bioinformatics/btl529.

42.   Kazlauskas, D., Krupovic, M., Guglielmini, J., Forterre, P., and Venclovas, Č. (2020). Diversity and

evolution of B-family DNA polymerases. Nucleic Acids Res *48*, 10142–10156. 10.1093/nar/gkaa760.

43.   Johnson, M., Zaretskaya, I., Raytselis, Y., Merezhuk, Y., McGinnis, S., and Madden, T.L. (2008). NCBI

BLAST: a better web interface. Nucleic Acids Research *36*, W5–W9. 10.1093/nar/gkn201.

44.   Zulkower, V., and Rosser, S. (2020). DNA Features Viewer: a sequence annotation formatting and plotting

library for Python. Bioinformatics *36*, 4350–4352. 10.1093/bioinformatics/btaa213.

E02 BGKB01000168.1_100
L = 921 aa, TM-score = 0.7926

E01 LGRX01004602.1_6
L = 1478 aa, TM-score = 0.8462

Referenced
Mirusviruses

E02

M06 & M07

E01

M05

E03

M01 & M03

E03 JH993101.1_6
L = 1258 aa, TM-score = 0.8065

E04

M02

E04 CDFJ01186769.1_15
L = 1039 aa, TM-score = 0.8786

Mirusvirus reference
M03_0104_TARA_00027_24
L = 1105

E05 BJOX01001092.1_29
L = 1286 aa, TM-score = 0.5635

M01 & M04

E05

Tree scale: 1

CM015708.1
275050 bp
(Schizochytrium sp.)*

Primase
Phosphotransferase
Translesion related gene
Phosphotransferase
Ribonucleotide Reductase
Transcription factor
Portal
Transcription factor 2
Maturation
Terminase
Holliday Junction Resolvase
Phosphotransferase
PolB
MCP
RNA polymerase subunit A
RNA polymerase subunit A
Triplex2

CM015707.1
289439 bp
(Schizochytrium sp.)*

Holliday Junction Resolvase
Phosphotransferase
Phosphotransferase
PolB
Phosphotransferase
Ribonucleotide Reductase
Ribonucleotide Reductase I
RNA polymerase subunit A
Phosphotransferase
Maturation
Triplex2
RNA polymerase subunit A
Triplex1
MCP
ATPase
Portal
Translesion related gene
Primase
Transcription factor
Heliorhodopsin
Transcription factor 2
Terminase
Holliday Junction Resolvase

Heliorhodopsin
Translesion related gene
Transcription factor
Terminase
Transcription factor
Portal
MCP
Primase
Endonuclease
ATPase
Triplex2
Transcription factor 2
RNA polymerase subunit A
Transcription factor

CAJHUC010000281.1
324836 bp
(Ostreobium quekettii)*

Holliday Junction Resolvase
Ribonucleotide Reductase I
Ribonucleotide Reductase
RNA polymerase subunit A
Phosphotransferase
Heliorhodopsin
Endonuclease
PolB

■ **Predicted genes**

■ **Mirusvirus markers**

■ **Annotated genes**