Programmable DNA shell scaffolds for directional membrane budding

Michael T. Pinner[1,2], Hendrik Dietz[1,2]*

[1] Laboratory for Biomolecular Nanotechnology, Department of Biosciences, School of Natural Sciences, Technical University of Munich; Garching bei München, 85748, Germany.
[2] Munich Institute of Biomedical Engineering, Technical University of Munich; Garching bei München, 85748, Germany.
* Corresponding author. Email: dietz@tum.de

Abstract: In the pursuit of replicating biological processes at the nanoscale, controlling cellular membrane dynamics has emerged as a key area of interest. Here, we report a system mimicking virus assembly to control directional membrane budding. We employ three-dimensional DNA origami techniques to construct cholesterol-modified triangles that self-assemble into polyhedral shells on lipid vesicles, resulting in gradual curvature induction, bud formation and spontaneous neck scission. Strategic positioning of cholesterols on the triangle surface provides control over the directionality of bud growth and yields daughter vesicles with DNA endo- or exoskeletons reminiscent of clathrin-coated vesicles. This process occurs with rapid kinetics and across various lipid compositions. When combined into a two-step process, nested bivesicular objects with DNA shells encapsulated between lipid vesicles could be produced. Our work replicates key aspects of natural endocytic and exocytic pathways, opening new avenues for exploring membrane mechanics and applications in targeted drug delivery and synthetic biology.

Main text:

Membrane budding is a ubiquitous biological phenomenon in which parent lipid membranes release daughter vesicles for the transport of materials across the membrane. In biological systems, membrane curvature changes are often induced by curved proteins such as clathrin. With its distinctive triskelion shape, clathrin acts as a molecular scaffold by binding to membrane-bound receptor-adaptor protein complexes, followed by self-assembly into spherical cage-like structures that induce curvature in the cell membrane[1–3]. Recent studies confirmed clathrin's membrane-bending properties in vitro[4] and in cells[5]. We hypothesised that any material capable of self-assembly into curved shapes could act as a molecular scaffold for membrane budding, with DNA origami emerging as a promising candidate. Herein, a several kilobases long, single-stranded 'scaffold' strand is mixed with an array of shorter oligonucleotides and exposed to a thermal annealing ramp to drive the folding of the DNA origami structure[6–8]. Previous efforts in the field related to lipid membranes have successfully tethered membrane-spanning DNA nanopores of various complexity to lipid bilayers using hydrophobic moieties such as cholesterol[9], ethyl-phosphorothioate[10] or tetraphenylporphyrin[11]. Notably, cholesterol-labelled DNA origami objects tethered to lipid membranes retain diffusive mobility influenced by factors like linker oligonucleotide length[12], paving the way for the 2D assembly of monomeric structures into multimeric lattices[13]. Prior studies have also shown that both curved[14–16] and planar[14] origami structures can induce tubulation in giant unilamellar lipid vesicles. On-membrane polymerisation of origami

46 structures likewise resulted in the deformation of the underlying membrane reminiscent of
47 clathrin networks involved in cellular budding processes[13,17–20].
48
Building upon this foundational work, we have developed an artificial, DNA-origami-based
membrane budding system with spontaneous neck scission. At its core are three-
dimensional membrane-interacting triangular subunits constructed from DNA with bevelled
edges[21]. Strategically placed shape-complementary protrusions and recesses at the triangle
edges direct the self-assembly of these triangular subunits into icosahedral shells through
adhesive base pair stacking interactions[22]. Interactions with lipid membranes of giant vesicles
(GVs) were promoted using cholesterol moieties placed on the origami structure. We
hypothesised that membrane deformation processes may be sufficient for the spontaneous
scission of the bud neck without the need for active constriction by GTPases like dynamin[23].
Indeed, the self-assembly of membrane-bound triangles induced local curvature and finally
led to the formation and eventual scission of buds. The positioning of cholesterol determined
whether the bud-formation occurs inward or outward from the source vesicle, leading to
DNA-shell-coated vesicles (DCV) and vesicle-coated DNA shells (VCD), respectively.
62
DNA origami triangles as molecular scaffolds
We adapted a previously developed programmable icosahedral shell canvas, in which twenty
equilateral DNA origami triangles form a closed icosahedron triggered by elevated
concentrations of magnesium chloride for membrane-supported assembly (Supplementary
Figure 1 & Supplementary Tables 1-24)[21]. To provide the necessary hydrophobic interactions
required for association with lipid membranes, we incorporated multiple cholesterol-bearing
oligonucleotides (chol-oligos) at the shell-inward-facing surface of the constituent triangular
DNA origami subunits (Fig. 1a & Supplementary Figure 2). We hypothesised that the 2D
diffusion of triangles on a fluid membrane can support the assembly of icosahedral shells
whilst pulling membrane material along, resulting in increasingly deformed membrane buds
growing away from the parent membrane until the bud neck is cut and a DNA-shell-coated
vesicle (DCV) is released into solution.
75
We screened cholesterol density and positioning to balance the solubility of the triangles with
membrane affinity (Supplementary Figure 3a & b) and then validated their assembly into the
expected icosahedral shells in the absence of membranes (Supplementary Figure 3c).
Introducing cholesterol resulted in a shift from the previously observed relatively uniform
icosahedral shells to a mixture containing both octahedral and icosahedral shells. This
variation was controllable: by limiting the reach of each cholesterol moiety, we could largely
revert the assembly to predominantly icosahedral structures. The addition of lipid membranes
into the system introduced another variable, again leading to the occurrence of octahedral
structures. This tendency varied depending on the amount of cholesterol per triangle and the
presence of lipid vesicles, detailed in Supplementary Figures 4, 5 & 6a. We believe this effect
arises from a complex interplay of forces, including crowding of membrane-bound triangles,
altered or biased relative alignment of triangles (possibly influenced by local curvature
induction), and elastic deformation within the shells caused by less-than-ideal angles
between the triangle subunits. Our shells share this structural ambiguity with clathrin, which
likewise assembles into differently shaped cages without negatively impacting its role as a
molecular scaffold, and we thus considered both shell types suitable for our objectives[3].

We prepared giant vesicles (GVs) composed entirely of DOPC and fluorescently labelled
DOPE to serve as model membranes in our budding assays[24]. Aliquots of GVs and chol-
decorated triangles were then mixed to anchor the nanostructures to the vesicles (Fig. 1b,
detailed in Supplementary Figure 7). The inclusion of NaCl in the reaction mixture minimised
non-specific binding and ensured that the triangles attached to the vesicles in the correct
orientation as defined by the positioning of their cholesterol moieties[16]. We then triggered the
assembly of membrane-bound triangles by increasing the $MgCl_2$ concentration and
incubating the solution at 37 °C for up to multiple days. Negative stain transmission electron
microscopy (TEM) imaging revealed abundant DCVs, characterised by dark DNA shells fully
encapsulating bright lipid vesicles (Fig. 1c, top right; Supplementary Figure 8). Occasionally,
the shell-enclosed vesicles appeared deformed or cracked, which we attribute to staining
and drying artefacts (Supplementary Figure 9). To also study DCVs in their native state, we
acquired micrographs of vitrified samples by cryogenic transmission electron microscopy
(cryoEM, Fig. 1c, bottom right; Supplementary Figure 10a-c). This data generally agrees with
the conclusions drawn from negative stain TEM data, but the enclosed vesicles seen by
cryoEM rarely had any defects except for occasionally encapsulating even smaller vesicles
themselves. In the cryo-EM micrographs, the vesicle typically fills up the available space
inside the origami shell, with the membrane in direct contact with the enclosed DNA shell.
111

As the size difference between DCVs and GVs spans multiple orders of magnitude, direct
observation of DCV formation is challenging. To overcome this problem, we instead used
large vesicles (LVs) of approx. 200 nm in our budding assays. By additionally using isotonic
$MgCl_2$ solution and shortening the time of shell assembly before vitrification, we could resolve
intermediate states of DCV formation by cryoEM (Fig. 1d & Supplementary Figure 10d). Whilst
membrane patches free of DNA triangle subunits appear undisturbed, those with triangles on
them had deformations that matched the curvature of the triangle assemblies. Notably, some
vesicles carry multiple buds with clearly visible necks. This image data strongly supports the
notion that the assembly of membrane-bound triangles induces curvature and that budding
is the mechanism behind DCV formation.
122

Characterisation of DCVs
We analysed DCV formation efficiency by agarose gel electrophoresis using fluorescently
labelled DNA origami and GVs (Supplementary Figure 11). Lipid material colocalised
exclusively with the higher-order shell assemblies and was virtually absent at the level of
monomeric triangles and intermediates. We also note that cholesterol-free triangles, variants
incapable of shell formation, and triangles added to GVs after several hours of membrane-
free assembly all failed to produce DCVs efficiently (Supplementary Figures 6b, 12 & 13).
These findings underscore the critical role of on-membrane triangle assembly in DCV
formation.
132

To evaluate whether the lipid vesicles were fully enclosed by well-ordered polyhedral DNA
shells, we recorded single-axis tilt series to generate tomograms, allowing us to study DCVs
in 3D (Fig. 2a). The representative tomograms confirm full engulfment of the vesicle by closed
shells featuring the symmetry properties expected by design. Occasionally, we found DCVs
where the DNA origami shell had assembly "scars" consisting of one to several missing

triangles (Fig. 2b). Considering that the increasingly high curvature at the bud neck can
become a steric barrier to completion of the DNA origami shell, we believe the occurrence of
scars to be a signature of the proposed budding process. In this scenario, spontaneous
membrane scission outcompetes the completion of the DNA origami shell.

The addition of concentrated $MgCl_2$ solution as a trigger for shell assembly inevitably
increases the osmolality of the sample buffer and may thus alter the biophysical properties
of the membrane by deflating the vesicles. Indeed, hyperosmotic stress has previously been
exploited to alter vesicle shape by membrane-bound DNA origami[16]. We studied the influence
of membrane tension on DCV yield by mixing triangle-covered GVs with increasing amounts
of glycine as an osmolyte. Interestingly, DCV yields correlated strongly with the osmotic
environment. In the hypotonic regime, we observed moderate to low yields, with the lowest
tonicity tested producing only a very faint band by agarose gel electrophoresis (Fig. 2c,
Supplementary Figure 14 & Supplementary Table 25). Conversely, as we transitioned into the
hypertonic regime, we noted a marked and progressive increase in DCV yields, with the most
pronounced yield achieved at the highest buffer osmolality. These findings reveal a direct
relationship between membrane tension and DCV yields, reinforcing the notion that DCVs
emerge from intricate membrane remodelling processes.

Our system's sensitivity to membrane tension draws a parallel to clathrin-mediated
endocytosis, which depends on actin engagement for budding under increased membrane
tension[25]. Another point of similarity is that the assembly of both our DNA triangles and
clathrin itself is sufficient to deform model membranes in vitro[4]. However, vesicle deformation
has also been demonstrated by attaching DNA origami structures to lipid vesicles using
cholesterol as an anchor, not requiring the assembly of subunits[14]. Having demonstrated the
necessity of triangle assembly for DCV formation (Supplementary Figure 12), we next
investigated the roles of cholesterol insertion and outer leaflet expansion in inducing
curvature and budding. To evaluate this, we prepared GVs using a lipid mixture containing
3% biotinylated DOPE and replaced chol-oligos with biotinylated oligonucleotides linked to
NeutrAvidin (b-NAv) as membrane anchors. This allowed DNA triangles to attach to the
membrane via NeutrAvidin-biotin interactions without inserting foreign lipids into the outer
leaflet. Remarkably, this approach still resulted in DCV formation and adding excess chol-
oligos with an orthogonal sequence did not improve yields (Fig. 2d & Supplementary Figure
15). These findings show that cholesterol insertion is not required for DCV formation,
supporting the idea that membrane buds are primarily shaped by shell assembly.

In our study, we primarily used vesicles composed of DOPC, supplemented with a small
fraction of fluorescently labelled DOPE species. The high fluidity and homogeneity of DOPC
membranes provide idealised conditions that do not necessarily reflect the properties of
biological membranes. We next investigated the versatility of our budding system using GVs
with diverse lipid compositions. These included single-lipid species compositions (low-
melting DOPC, and high-melting DPPC and DSPC), binary mixtures (70% DOPC with either
30% cholesterol, DPPC, or DSPC), and four polar extracts of natural origin (derived from soy,
brain, yeast, and *E. coli*). With the exception of GVs composed entirely of high-melting lipids,
DCVs successfully formed from all tested compositions with varying efficiencies (Fig. 2e &
Supplementary Figure 16). The successful formation of DCVs from binary mixtures containing

high-melting lipids suggests that the overall membrane fluidity, rather than the mere presence
of high-melting lipids, is a critical factor in determining budding success. This is supported
by the fact that low-melting DOPC constituted the majority (70%) of these mixtures, rendering
the bilayer overall fluid.
Another important factor for DCV formation is neck scission. Constriction of the bud neck
may be driven by spontaneous and induced curvature at the neck due to a high triangle
density on the membrane and adhesion forces[26,27]. Scission can also be helped by line
tension in phase-separated membranes, where the phase boundary represents a linear
defect along which fission is facilitated[28]. Indeed, when comparing DCV yields obtained from
pure   DOPC   membranes   versus   phase-separated   membranes   composed   of
DOPC/DPPC/chol or DOPC/DSPC/chol (40:40:20 mol%), we found that the phase-separated
membranes generally yielded more DCVs (Fig. 2f & Supplementary Figure 17).
196

We next analysed the kinetics of DCV formation by starting the assembly reaction and
collecting aliquots at various time points. The budding process was rapid, with a substantial
number of DCVs already present at the 0-minute time point (Fig. 2g & Supplementary Figure
18). We attribute the rapid kinetics to the membrane confinement of the origami triangles,
which generates elevated local concentrations and restricts mobility to two dimensions.
Consistent with earlier observations, the majority of DCVs formed initially exhibited an
octahedral geometry. Over time, however, a growing fraction of icosahedral DCVs was
detected. This progression mirrors the behaviour observed in vesicle-free control assemblies,
where DNA origami triangles rapidly assemble into intermediates resembling octahedra in the
agarose gel, which subsequently mature into icosahedral shells. However, the maturation of
octahedral DCVs into icosahedral ones appears restricted. We believe that size constraints
imposed by the vesicle inside the DCV stabilises its configuration and limits the expansion of
the inner cavity necessary to form larger icosahedral species.
210

Inward budding
Having studied the formation and characterisation of DCVs, we next investigated the
directional reversal of this process. DCV budding mimics endocytic processes with the
important difference that in our system the triangular subunits approach the parent vesicle
from the exterior. By shifting the placement of cholesterols on the triangle surface from the
bottom (shell-inward) to the top (shell-outward) face, we hypothesised that we could reverse
the budding directionality in a process reminiscent of exocytosis. As the triangles would still
bind to the vesicles from the outside, we expect the formation and release of vesicle-coated
DNA shells (VCDs) into the lumen of the parent vesicle (Fig. 3a), effectively creating an
endosome-like compartment with a DNA origami endoskeleton. We prepared large vesicles
(LVs) by extruding GVs through 200 nm-sized pore filters to facilitate cryoEM imaging, mixed
them with triangles carrying one cholesterol moiety per side and triggered shell assembly.
Direct imaging with cryoEM (Fig. 3b) revealed VCDs as lipid vesicles tightly wrapped around
icosahedral shells located within their respective parent vesicles. Occasionally, we also
observed free VCDs, presumably set free by the bursting of the parent vesicle, and the
presence of holes in the DNA shell similar to those observed in DCVs.
To capture the dynamics of the suspected budding process, we prepared vitrified samples
with shortened assembly times and analysed them by cryoEM. The micrographs we obtained
revealed the gradual deformation of the parent vesicle in response to the assembly of

membrane-bound triangles, leading to the formation of inward-growing buds (Fig. 3c). In
contrast to fully formed VCDs, these particles remain connected to the parent membrane,
sometimes just by a narrow bud neck.

We finally explored the creation of bivesicular shell structures by combining outward and
inward budding reactions in a two-step process using triangles with sequence-orthogonal
linker handles on both faces (Fig. 4a). First, we produced DCVs by hybridising chol-oligos to
the triangles' bottom face and mixing the triangles with GVs as described earlier. After
isolating the DCVs by centrifugation, we prepared LVs and coated them with chol-oligos
complementary to the linkers on the DCV's outer face. Mixing the DCV concentrate with these
LVs and incubating at 37 °C resulted in the formation of vesicle-coated DCVs (VCDCVs), as
confirmed by cryoEM (Fig. 4b). Like VCDs, VCDCVs were typically enclosed within their
parent vesicles as a result of the proposed budding mechanism. These particles may be
considered simple examples of synthetically created organelles featuring a cytoplasm-like
compartment (the inner vesicle) and a periplasm-like compartment (the void between the
outer envelope membrane and the inner lipid membrane). The icosahedral DNA shell acts as
a stabilising mechanical skeleton, offering engineering options for including additional
molecular functionalities.

Discussion
In this study, we developed a DNA origami-based membrane budding system that
recapitulates key aspects of clathrin-mediated endocytosis (CME) without relying on
components of cellular budding machineries. Like clathrin, the DNA origami triangles function
as molecular scaffolds whose self-assembly into cage-like structures induces membrane
deformation and ultimately drives the formation of vesicles. Free energy is supplied by the
self-assembly process and relayed onto the bilayer by either cholesterol or biotin-NeutrAvidin
linkages, mimicking the way clathrin associates with membranes via adaptor proteins. Using
average stacking energies reported for DNA nanostructures (-2.97 k_BT, T = 300 K), we
estimate the free energy of icosahedral assembly (30 edges with 16 base-stacking
interactions each) at approx. -1400 k_BT, and -570 k_BT for octahedra[29]. Our estimates exceed
the energetic cost of bending DOPC membranes into spherical vesicles (460 k_BT), derived
from Helfrich's theory (E = $8\pi k_C$, where k_C = 18.3 k_BT at 298 K), and support our experimental
observations[30,31].

Both clathrin cages and DNA origami shells demonstrate structural flexibility, forming various
geometries while maintaining their scaffolding function[3]. The triangles noticeably shifted from
icosahedral to octahedral shells when assembled on the lipid membrane with influencing
factors including lipid composition and membrane tether type. We suspect electrostatic
interactions between the membrane and the origami, as well as molecular crowding effects,
as key drivers behind this phenomenon. However, a detailed analysis lies beyond the scope
of this study.
Our system demonstrates robust performance across a variety of lipid compositions,
including natural lipid extracts, with rapid kinetics highlighting its versatility and potential for
broader applications. Nonetheless, we identified limitations under specific conditions. DCV
formation was impaired in membranes composed entirely of high-melting lipids (DPPC,
DSPC) and yields were reduced under hypotonic conditions. These constraints can be

explained by the Helfrich Hamiltonian, which relates the energetic cost of membrane
deformation to the mechanical properties of the bilayer[30]. Higher bending rigidities of gel-
phase membranes and increased membrane tension under hypotonic conditions impose
energetic penalties directly impacting DCV yields[32]. While clathrin underlies the same
limitations, biological membranes make use of the cytoskeleton to enable CME even under
high membrane tension[25]. Future work could explore active neck constriction to overcome
the current limitations of our system.
283
In summary, we present a versatile platform for engineering vesicles of controlled size
featuring addressable endo- or exoskeletons. This approach can be extended to encapsulate
molecules of interest during the budding process, potentially leading to molecular payload
applications. Using nested bivesicular structures, it becomes possible to design scenarios
where two different molecular payloads are delivered simultaneously but separately. This
work may thus pave the way for innovative applications in targeted drug delivery, vaccine
development, and synthetic biology. The potential to manipulate membrane dynamics at the
nanoscale also opens new horizons in synthesising artificial cells and organelles, offering a
valuable tool for researchers in their quest to harness the power of nanotechnology for
practical and therapeutic purposes.
294

References
1. Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nat. Rev. Mol.*
Cell Biol. **19**, 313–326 (2018).
2. Crowther, R. A., Pinch, J. T. & Pearse, B. M. F. On the structure of coated vesicles. *J.*
Mol. Biol. **103**, 785–798 (1976).
3. Morris, K. L. *et al.* Cryo-EM of multiple cage architectures reveals a universal mode of
clathrin self-assembly. *Nat. Struct. Mol. Biol.* **26**, 890–898 (2019).
4. Dannhauser, P. N. & Ungewickell, E. J. Reconstitution of clathrin-coated bud and vesicle
formation with minimal components. *Nat. Cell Biol.* **14**, 634–639 (2012).
5. Cail, R. C., Shirazinejad, C. R. & Drubin, D. G. Induced nanoscale membrane curvature
bypasses the essential endocytic function of clathrin. *J. Cell Biol.* **221**, e202109013 (2022).
6. Rothemund, P. W. K. Folding DNA to create nanoscale shapes and patterns. *Nature* **440**,
297–302 (2006).
7. Douglas, S. M. *et al.* Self-assembly of DNA into nanoscale three-dimensional shapes.
Nature **459**, 414–418 (2009).
8. Dey, S. *et al.* DNA origami. *Nat. Rev. Methods Primer* **1**, 1–24 (2021).
9. Langecker, M. *et al.* Synthetic Lipid Membrane Channels Formed by Designed DNA
Nanostructures. *Science* **338**, 932–936 (2012).
10. Burns, J. R., Stulz, E. & Howorka, S. Self-Assembled DNA Nanopores That Span Lipid
Bilayers. *Nano Lett.* **13**, 2351–2356 (2013).
11. Burns, J. R. *et al.* Lipid-Bilayer-Spanning DNA Nanopores with a Bifunctional Porphyrin
Anchor. *Angew. Chem. Int. Ed.* **52**, 12069–12072 (2013).
12. Khmelinskaia, A., Mücksch, J., Petrov, E. P., Franquelim, H. G. & Schwille, P. Control of
Membrane Binding and Diffusion of Cholesteryl-Modified DNA Origami Nanostructures
by DNA Spacers. *Langmuir* **34**, 14921–14931 (2018).

13. Journot, C. M. A., Ramakrishna, V., Wallace, M. I. & Turberfield, A. J. Modifying
Membrane Morphology and Interactions with DNA Origami Clathrin-Mimic Networks.
ACS Nano **13**, 9973–9979 (2019).

14. Grome, M. W., Zhang, Z., Pincet, F. & Lin, C. Vesicle Tubulation with Self-Assembling
DNA Nanosprings. *Angew. Chem. Int. Ed.* **57**, 5330–5334 (2018).

15. Liu, L., Xiong, Q., Xie, C., Pincet, F. & Lin, C. Actuating tension-loaded DNA clamps drives
membrane tubulation. *Sci. Adv.* **8**, eadd1830 (2022).

16. Franquelim, H. G., Khmelinskaia, A., Sobczak, J.-P., Dietz, H. & Schwille, P. Membrane
sculpting by curved DNA origami scaffolds. *Nat. Commun.* **9**, 811 (2018).

17. Yang, J. *et al.* Modulating Lipid Membrane Morphology by Dynamic DNA Origami
Networks. *Nano Lett.* **23**, 6330–6336 (2023).

18. Baumann, K. N. *et al.* Coating and Stabilization of Liposomes by Clathrin-Inspired DNA
Self-Assembly. *ACS Nano* **14**, 2316–2323 (2020).

19. Kocabey, S. *et al.* Membrane-Assisted Growth of DNA Origami Nanostructure Arrays.
ACS Nano **9**, 3530–3539 (2015).

20. Zhan, P. *et al.* 3D DNA origami pincers that multitask on giant unilamellar vesicles. *Sci.*
Adv. **10**, eadn8903 (2024).

21. Sigl, C. *et al.* Programmable icosahedral shell system for virus trapping. *Nat. Mater.* **20**,
1281–1289 (2021).

22. Gerling, T., Wagenbauer, K. F., Neuner, A. M. & Dietz, H. Dynamic DNA devices and
assemblies formed by shape-complementary, non–base pairing 3D components.
Science **347**, 1446–1452 (2015).

23. Sweitzer, S. M. & Hinshaw, J. E. Dynamin Undergoes a GTP-Dependent Conformational
Change Causing Vesiculation. *Cell* **93**, 1021–1029 (1998).

24. Weinberger, A. *et al.* Gel-Assisted Formation of Giant Unilamellar Vesicles. *Biophys. J.*
105, 154–164 (2013).

25. Boulant, S., Kural, C., Zeeh, J.-C., Ubelmann, F. & Kirchhausen, T. Actin dynamics
counteract membrane tension during clathrin-mediated endocytosis. *Nat. Cell Biol.* **13**,
1124 (2011).

26. Steinkühler, J. *et al.* Controlled division of cell-sized vesicles by low densities of
membrane-bound proteins. *Nat. Commun.* **11**, 905 (2020).

27. Agudo-Canalejo, J. & Lipowsky, R. Stabilization of membrane necks by adhesive
particles, substrate surfaces, and constriction forces. *Soft Matter* **12**, 8155–8166 (2016).

28. Jülicher, F. & Lipowsky, R. Domain-induced budding of vesicles. *Phys. Rev. Lett.* **70**,
2964–2967 (1993).

29. Kilchherr, F. *et al.* Single-molecule dissection of stacking forces in DNA. *Science* **353**,
aaf5508 (2016).

30. Helfrich, W. Elastic properties of lipid bilayers: theory and possible experiments. *Z.*
Naturforschung Teil C Biochem. Biophys. Biol. Virol. **28**, 693–703 (1973).

31. Doktorova, M., Harries, D. & Khelashvili, G. Determination of bending rigidity and tilt
modulus of lipid membranes from real-space fluctuation analysis of molecular dynamics
simulations. *Phys. Chem. Chem. Phys. PCCP* **19**, 16806–16818 (2017).

32. Solmaz, M. E. *et al.* Optical stretching as a tool to investigate the mechanical properties
of lipid bilayers. *RSC Adv.* **3**, 10.1039/C3RA42510J (2013).

Fig. 1 | DNA-shell-coated vesicles form by membrane budding. a, Schematic of a DNA origami triangle-based membrane budding system. The triangular structures are decorated with cholesterol moieties to tether them to the membrane of a lipid vesicle. Upon increasing the MgCl$_2$ concentration, the triangles self-assemble into polyhedral shells, pulling the membrane underneath along to form a bud. Following spontaneous neck scission, a DNA-shell-coated vesicle (DCV) is released. **b,** Pseudocoloured micrographs of DNA origami triangle attachment (grey) onto giant vesicles (GVs, magenta). Triangles will adsorb onto GVs even if they are not decorated with cholesterol (top left & top right panels). However, adding sodium chloride suppresses this non-specific adsorption (bottom left panel), and membrane binding will only occur if cholesterol is attached to the triangle (bottom right panel). Scale bars: 10 μm. **c,** Transmission electron micrographs of empty DNA shells and DCVs. In negative stain TEM (top panels), lipid vesicles within DCVs appear as white rings inside the DNA shell. In cryogenic electron microscopy (cryoEM, bottom panels), the vesicle is visible as a dark ring inside the DNA shell. Scale bars: 100 nm. **d,** CryoEM micrographs of large vesicle (LV) deformation upon assembly of membrane-bound DNA origami triangles. Bud curvature increases gradually and closely follows the intrinsic curvature of the assembling shells. Scale bars: 100 nm.

384
385
Fig. 2 | Characterisation of DNA-shell-coated vesicles. a, Tomogram of an icosahedral
DCV depicted as summed up z-slices of the top, central and bottom segments. The topmost
segment appears as a tilted, incomplete pentamer. The central segment reveals a lipid vesicle
trapped within the origami shell. The bottommost segment is a pentameric shell cap. **b,** DCVs
occasionally appear incomplete with visible 'scars' of missing triangles. The bud neck
prevents closure of the DNA shell by steric hindrance, resulting in a gap in the origami shell
that remains visible after scission of the neck. Left panel: negative stain TEM; Right panel:
cryoEM. Scale bars: 100 nm. **c,** Agarose gel of DCVs prepared under varying osmotic
conditions. Samples in the top gel contain triangles (grey, EtBr) and GVs (magenta, DOPE-
Atto643), whereas the bottom gel contains triangles only and serves as assembly control.
Glycine and MgCl₂ were added at increasing concentrations to vary sample tonicity and
promote triangle assembly. Budding efficiency strongly correlates with tonicity, working best
under hypertonic conditions and coming to a near halt at the most hypotonic condition tested.
The unhindered assembly of triangles in absence of vesicles indicates that DCV formation is
strongly influenced by changes in membrane tension. **d,** Left: Agarose gel of DCVs prepared
by using biotin-NeutrAvidin (b-NAv) interactions to connect origami triangles (grey, Atto643)
and GVs (magenta, DOPE-Atto488). The addition of orthogonal chol-oligos did not influence
DCV yields. Right: TEM micrographs of DCVs obtained by this approach. Scale bar: 100 nm.
e, Agarose gel of DCVs obtained from GVs of varying lipid composition. Only GVs composed
entirely of high-melting lipids prevented DCV formation. Grey: DNA (EtBr); Magenta: Lipids
(DOPE-Atto643); **f,** Left: Agarose gel comparing DCVs obtained from GVs composed of
DOPC or phase-separated ternary lipid mixtures. Phase-separated vesicles yielded more

DCVs, suggesting an influence of phase boundaries on the budding mechanism. Grey: DNA
(EtBr); Magenta: Lipids (LissRhod-PE); Right: Micrographs of exemplary phase-separated
vesicles showing the lipid-ordered (cyan) and -disordered (red) phases. Scale bars: 20 µm. **g,**
DCV formation kinetics (top) compared against vesicle-free assembly kinetics. DCV budding
occurs instantly and yields mostly octahedral species, but at longer incubation times a
notable fraction of icosahedral DCVs is formed. Grey: DNA (EtBr); Magenta: Lipids (DOPE-
Atto643);
415

416
417
Fig. 3 | Vesicle-coated DNA shells. a, Illustration of vesicle-coated DNA shell (VCD)
formation by budding into its parent vesicle. The budding direction is reversed with respect
to DCVs by shifting the cholesterol linker handles from the shell-inner face to the shell-outer
face of the triangle subunits. **b,** cryoEM micrographs of VCDs within their parent vesicles.
Scale bars: 100 nm. **c,** cryoEM micrographs of inward-oriented membrane buds at different
stages. Scale bars: 100 nm.
424

top

bottom

● single-stranded linker handle

● cholesterol

inner vesicle
DNA shell
outer vesicle
parent vesicle

425
426
Fig. 4 | Multivesicular DNA shells. a, Illustration of a bivesicular DNA shell (vesicle-coated
DCV; VCDCV) contained within its parent vesicle, produced in a 2-step budding assay using
triangles with orthogonal linker handles on their top and bottom faces. In the first step, the
bottom face linkers are hybridised to cholesterol-modified DNA strands for DCV formation by
outward budding. Then, DCVs are mixed with LVs carrying single-stranded linker handles
complementary to those on the shell-outer face of the triangles, resulting in inward budding
and VCDCV formation. As a result, VCDCVs have four layers: An inner vesicle stemming from
the first step (orange), the DNA origami shell (white), an outer vesicle (pink) coupled to the
shell by complementary linker handles, and the parent vesicle (purple) with its DNA linker
handles (green blobs). **b,** cryoEM micrographs of VCDCVs. Images were Gaussian blurred
and levelled for clarity. Scale bars: 100 nm.
438

Methods

Materials
All chemicals were purchased from Sigma Aldrich and Carl Roth unless stated otherwise. 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC), polar lipid extracts (soy, brain, yeast, e. coli), TopFluor cholesterol and Lissamine Rhodamine B-1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) were obtained from Avanti Research. Atto488- and Atto643-modified DOPE was obtained from Atto-Tec. NeutrAvidin was purchased from Thermo Scientific. Regular DNA oligonucleotides were purchased from IDT, and modified oligonucleotides were obtained from Biomers. Scaffold DNA was produced biotechnologically in-house. Buffer osmolalities were measured using an Osmomat 3000 freeze point osmometer (Gonotec). For buffer compositions, refer to Supplementary Table 26.

DNA origami folding, purification & quantification
DNA origami triangles were folded as described previously[21]. Briefly, circular scaffold strands of bacteriophage origin (sc8064, 50 nM; Supplementary Table 1) and staple oligonucleotides (200 nM per staple; Supplementary Tables 2-23) were mixed in folding buffer and exposed to a thermal annealing ramp (65 °C, 15'; 58-53 °C, 1 h/°C; stored at 20 °C) in a Tetrad2 (Bio-Rad) thermocycler. For some experiments, structures were fluorescently labelled by including Atto643 labelled oligonucleotides in the folding reaction. The folded product was purified by agarose gel electrophoresis by excising the leading band, crushing the gel pieces and centrifuging them in 0.45 µm Costar Spin-X spin columns (Corning) for 10 min at 8 krcf. When needed, the purified structures were washed and concentrated by ultrafiltration (Amicon Ultra 0.5 mL, 100 kDa cut-off, Millipore), exchanging the buffer for sodium buffer. Origami solutions were quantified using a NanoDrop8000 spectrophotometer (Thermo Scientific).

Vesicle preparation and quantification
Giant vesicles were prepared by gentle swelling of dry lipid films on polyvinyl alcohol gels as described previously[24], with slight modifications. A petri dish (⌀ 10 cm) was plasma cleaned in a glow discharge device, and then 1800 µl of a 5% solution of polyvinyl alcohol (Mowiol 28-99, Sigma Aldrich) in ddH$_2$O was evenly distributed across its surface. The dish was incubated for at least 30 min at 50 °C until a dry gel formed. Then, 162 µl of the lipid mixture (typically DOPC with additional 0.5-1% fluorescently labelled DOPE if visualisation by fluorescence was desired, or 0.4% TopFluor cholesterol and LissRhod-PE each for visualisation of lipid phases. Total lipid concentration: 2.54 mM) in chloroform was evenly spread out across the surface using a Drigalski spatula until the chloroform evaporated. Remaining traces of solvent were removed by exposing the dried lipid cake to a vacuum for at least 15 min. Next, 4-5 mL of sodium buffer was added to the petri dish and the solution was left in a humid chamber in the dark at RT for at least 1 h. GVs were detached by gently tapping against the dish and pipetting up and down with a cut pipette tip. The GV suspension was stored at 4 °C in the dark.

When GVs were to be washed and concentrated, the dry lipid film was swollen in caesium buffer or sucrose buffer and, after harvesting, transferred into a falcon tube. The tube was filled up with sodium buffer (isosmotic to the swelling buffers) and centrifuged for at least 30 min at 300 rcf. The supernatant was discarded, and the GV pellet was washed 2-3 more times by adding 900 µl sodium buffer and centrifuging the suspension for 7 min at 300 rcf.

Smaller-sized vesicles were prepared using a mini-extruder (Avanti Research) and polycarbonate membranes with the desired pore size (typically 200 nm; Whatman). GV preparations were pushed back and forth through the membrane 21 times.

491
492 Lipids in vesicle preparations were quantified using a colourimetric phospholipid
493 quantification assay kit (Sigma Aldrich CS0001) following the manufacturer's instructions.
494 The absorbance was measured at 570 nm on a Clariostar plus microplate reader (BMG
495 Labtech).
496
Vesicles used in the lipid mixture screens (Fig. 2 e,f) were swollen in sucrose buffer at 60 °C
to account for differences in lipid charge and melting temperature and washed as described
above. Lipid quantity was estimated by measuring the lipid content in sample A (DOPC
vesicles) using the colourimetric assay mentioned above. As this assay can only detect
choline-containing lipids, the lipid quantity in the remaining samples was estimated based on
the fluorescence of Atto643-DOPE species included at equal molar percentages in all lipid
mixtures. For this, vesicle preparations were diluted 1:10 and 10 µl aliquots were mixed with
90 µl Triton-X100 (2%) solution in a black 96 well plate. The plate was incubated at 60 °C for
1 h to disintegrate the vesicles and the fluorescence of Atto643-DOPE was measured in a
plate reader. The lipid quantity in the other vesicle preparations was finally calculated by
normalising the fluorescence readouts in respect to that of sample A, and multiplying the
obtained ratio by the colourimetric quantification result for sample A. Images of phase-
separated vesicles were obtained using a ThermoFisher EVOS M7000 imaging system.
510
Vesicle binding study
Microscope slides and coverslips were washed with ddH$_2$O and incubated in denatured BSA
blocking buffer (10 mM TRIS, 150 mM NaCl, 50 µM BSA, pH 8) overnight as described
previously[33]. Next, the glassware was thoroughly washed with ddH$_2$O to remove residual
blocking buffer and dried at 50 °C for 1 h. 6 µl of GVs (99.95% DOPC, 0.05% DOPE-Atto488;
swollen in imaging buffer A) was mixed with origami triangles (+/- cholesterol modified
strands; fluorescently labelled with Atto643 modified oligonucleotides; final origami
concentration: 7 nM) and topped up to 40 µl with imaging buffer B. 300 mM NaCl was added
to samples testing the effect of elevated sodium on the binding behaviour of origami. The
samples were pipetted into an imaging chamber formed by sandwiching a silicon isolator
(Grace Bio-Labs) between passivated coverslips and microscope slides. The samples were
imaged using a Leica TCS SP5 confocal microscope with a Leica HCX PL APO CS 63x/1.40-
0.60 oil immersion objective. Intensity profiles were obtained using Fiji[34].
524
Budding assays
Vesicles were prepared as described above using 99.0 – 100% DOPC with the remainder
being fluorescently labelled DOPE if visualisation by fluorescence was desired. Origami
triangles were hybridised to cholesterol-modified oligonucleotides by adding them at a 1.5 x
excess with respect to the total amount of compatible linker handles on the origami. The
hybridisation reaction was incubated at RT for at least 30 min. For outward budding reactions,
24 µl of untreated GV suspension was mixed with origami triangles with linker handles on the
shell-inward face at a final concentration of approx. 2 nM origami. In experiments where lipid
quantities in GV preparations were known, 1000 pmol lipid of GVs prepared in caesium buffer,
or 350 pmol of GVs prepared in sucrose buffer was mixed with the origami (4.5 µl at 15 nM).
The sample was topped up to 30 µl with sodium buffer, and the MgCl$_2$ concentration was
adjusted to 65 mM by adding 1 M MgCl$_2$ solution (hypertonic) or to 60 mM by adding isotonic
assembly buffer. Budding reactions were incubated at 37 °C for up to 3 d.
538
539 For the tonicity screen, undiluted, chol-hybridised triangles were mixed with GVs and diluted
540 using water and glycine buffer. The samples were incubated at RT for 2 h for triangles to bind
541 to the vesicles and for vesicles to adjust to the altered osmotic conditions. Finally, MgCl$_2$
542 solution was added to trigger assembly and samples were incubated at 37 °C overnight. For
543 detailed sample compositions, refer to Supplementary Table 25.

544
545 The budding kinetics were obtained by mixing samples with gel loading dye (50% Ficoll400,
546 20 mM $MgCl_2$, Orange G) and freezing them in liquid nitrogen at the indicated time points.
547 Once all time points were taken, the samples were thawed, mixed and used for agarose gel
548 electrophoresis.
549
550 For inward budding reactions, origami triangles (3 µl at 20 nM) with linker handles on the
551 shell-outward face were mixed with LVs (1500 pmol lipids; approx. 200 nm diameter) and
552 incubated for up to 3 d.
553
For bivesicular DNA shells (VCDCVs), triangles with orthogonal linker handles on the top and
bottom face were used, and inward and outward budding reactions were conducted
successively. First, DCVs were produced in an outward budding assay (1200 µl in total) using
cholesterol-modified oligonucleotides hybridised to the bottom face linker handles. The
sample was then centrifuged at 300 rcf for 5-8 min, and the pellet containing residual GVs
was discarded. The supernatant was then ultracentrifuged at 55 krpm (avg. 108 krcf) for 30
min in a Beckman Coulter Optima MAX-TL ultracentrifuge equipped with a TLA-110 fixed
angle rotor. 80-90% of the supernatant was removed, and the pellet was resuspended in the
remaining volume.
Meanwhile, LVs (200 nm diameter) were mixed with a second chol-oligo complementary to
the linker handles on the top face of the triangles. These oligonucleotides were added at a
50x excess with respect to the total amount of complementary linker handles on the origami,
and the solution was incubated at RT for 15-30 min. Then, the magnesium concentration in
the LV solution was adjusted to 60 mM $MgCl_2$ by adding isotonic assembly buffer. Finally,
LVs (1500 pmol lipids) were mixed with the DCV concentrate (to a final concentration of 1-5
nM DNA), and the samples were incubated at 37 °C for 1-3 d.
570
Budding using biotin-NeutrAvidin
For budding reactions using biotin-NeutrAvidin interactions to anchor triangles to the
membranes, GVs containing 3% biotinylated DOPE, 1% fluorescently labelled DOPE and 96%
DOPC were prepared. Origami triangles were folded with linker handles (Supplementary
Table 20) complementary to biotinylated oligonucleotides (Supplementary Tables 3) and
linker handles for purification using magnetic beads (Supplementary Table 4). Biotinylated
oligonucleotides were already included in the folding reaction to remove unbound strands
and excess staples by gel extraction. The triangles were then mixed with a 500-fold excess
of NeutrAvidin and incubated for 1 h at 37 °C on an orbital shaker. Next, the sample was
transferred into a tube previously passivated with denatured BSA blocking buffer and mixed
with magnetic beads (Dynabeads M-270 Streptavidin, Invitrogen) coated with dual-
biotinylated DNA linker handles (Supplementary Table 4) complementary to the linkers on the
origami. The sample was incubated for 1 h at RT on a rotary shaker, and the beads now
carrying the origami triangles were washed 3-4 times with sodium buffer. Finally, an invader
strand complementary to the full length of the purification linker handles (Supplementary
Table 4) was added in great excess, and the solution was incubated at 37 °C for 1 h to
uncouple the origami from the magnetic beads by strand displacement. The NeutrAvidin
bearing triangles were then mixed with biotinylated GVs, and the $MgCl_2$ concentration was
adjusted to 60 mM by adding isotonic assembly buffer. The sample was incubated at 37 °C
for 2.5 d.
591
Negative stain transmission electron microscopy
Sample solutions were incubated on glow-discharged copper grids (400 mesh) with a carbon
support film (Science Services) for 4-8 minutes, depending on the sample concentration. The
solution was then blotted off using filter paper (Whatman). Then, the grid was washed once
using stain solution (2% aqueous uranyl formate + 25 mM NaOH), followed by incubating

another stain droplet on the grid for 30 s. Excess stain was blotted off using filter paper, and
the grid was air-dried before imaging.
599
Grids were typically imaged at x26-30k magnification using SerialEM with an FEI Tecnai T12
electron microscope operated at 120kV and a Tietz TEMCAM-F416 camera. Image contrast
was auto-levelled using Fiji.
For tomography, one-directional tilt series were acquired from -50° to +50° in steps of 2°
using SerialEM. The resulting image stacks were imported into Etomo (IMOD[35]), and images
were aligned without fiducials by cross-correlation with cumulative correlation switched on.
Tomograms were generated by filtered back-projection (gaussian filter cut-off: 0.35; fall-off:
0.035;). For visualisation, the z-stacks thus generated were imported into Fiji and z-projected
by summing up z-slices corresponding to the top, middle and bottom sections of the
respective particle of interest.
610
Cryogenic electron microscopy
Sample grids for cryoEM were prepared using an automated vitrification system (FEI Vitrobot
Mark V). First, grids (Quantifoil R1.2/1.3, Cu 200 mesh, 100 holey carbon films, 2 nm carbon
support film; for images of DCVs: Ted Pella, Cu 400 mesh, lacey carbon, <3 nm carbon
support film) were glow-discharged and then inserted into the Vitrobot. The conditions inside
the device's environmental chamber were kept at either 4 °C or 22 °C and 100% relative
humidity. Samples were typically incubated on the grids inside the chamber for 5 min and
plunge-frozen in liquid ethane (blot force: -1, 0 or 25; blot time: 2-3 s; blot total: 1; drain time:
0 s;). The grids were imaged with a spherical-aberration (Cs)-corrected Titan Krios G2
electron microscope (Thermo Fisher) operated at 300kV and equipped with a Falcon III 4k
direct electron detector (Thermo Fisher). Samples were imaged at x29k magnification and
defocus values between -1 and -4 µm. Images were auto-levelled in Fiji.
623
Agarose gel electrophoresis and image processing
Agarose gels were cast by dissolving UltraPure agarose (Invitrogen) in running buffer (0.5 x
TBE, Carl Roth) supplemented with $MgCl_2$ and optionally ethidium bromide (0.025%, Carl
Roth). Samples were run for 2 h at 90 V and then scanned using a Typhoon FLA 9500 laser
scanner (GE Healthcare) at a pixel size of 50 µm. Gels containing only monomeric origami
structures were run with 5.5 mM $MgCl_2$ in the gel and running buffer and an agarose
concentration of 1.5%. To resolve and preserve assembled shell structures, the gel and
running buffer were supplemented with 20 mM $MgCl_2$ and the agarose concentration was
reduced to 0.8%. Due to the elevated salt concentration in these gels, the gel boxes were
cooled in a water-ice bath, and the running buffer was exchanged every 45 min to replenish
lost magnesium. For visualisation, brightness/contrast of the area underneath the gel pockets
was auto-levelled in Adobe Photoshop and Fiji, intentionally oversaturating the gel pockets
in the process.
Gels containing both fluorescent DNA and lipids were scanned using appropriate excitation
wavelengths and emission filters. A merged-colour image was created in Fiji, setting DNA
signal grey and lipid signal magenta.
640
References

33. Park, J. H. *et al.* Temperature-Induced Denaturation of BSA Protein Molecules for
Improved Surface Passivation Coatings. *ACS Appl. Mater. Interfaces* **10**, 32047–32057
(2018).
34. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).

35. Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer Visualization of Three-
Dimensional Image Data Using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996).

Acknowledgements
This manuscript has been edited in part with the assistance of the large language models
GPT3.5 (openAI) and Perplexity AI. We thank Andreas Bausch and Friedrich Simmel for
technical assistance, Björn Högberg for discussions, and Volodymyr Mykhailiuk for help with
illustrations. We acknowledge financial support received from the European Research
Council Advanced Grant awarded to H.D. (grant agreement 101018465) and the Deutsche
Forschungsgemeinschaft for the Gottfried Wilhelm Leibniz Program grants provided to H.D.
This project has received funding from the European Union's Horizon 2020 research and
innovation programme under the Marie Skłodowska-Curie grant agreement No 765703.

659
660   **Author contributions:** The research was performed by M.T.P., supervised by H.D. The
661   manuscript was written by M.T.P. and H.D.
662
663   **Competing interests:** The authors declare no competing interests.
664
665   **Data availability:** All data are available in the main text or supplementary materials.
666
667   **Supplementary Materials**
668   Supplementary Figures 1 to 18
669   Supplementary Tables 1 to 26
670

672
673
674 **Supplementary Figure 1 | Triangle design map.** Exemplary linker handles extending from
675 the origami surface are shown in red (shell-outer face) and green (shell-inner face). The
676 three main domains represent the three sides comprising the triangle. Protrusions used for
677 triangle assembly are highlighted in orange, the shape-complementary recesses are visible
678 as holes in the design.
679

origami surface

chol. bearing oligo

lipid bilayer

680
681
682 **Supplementary Figure 2 | Attachment of origami triangles to vesicles via cholesterol-**
683 **bearing oligonucleotides.** Linker handles extending from the origami surface are
684 hybridised to cholesterol-bearing oligonucleotides with complementary sequences.
685 Cholesterol acts as a lipid membrane anchor to tether the triangles to lipid vesicles.
686

687
688
Supplementary Figure 3 | The influence of cholesterol position and orientation on
origami triangles. a, Linker handle positions on three different triangle variants, and
scheme of distal and proximal cholesterol configurations. The upper, white helix represents
the triangle surface (side view), the pink strand represents the cholesterol-bearing oligo
(chol-oligo). Cholesterol may be oriented facing away (distal) or towards (proximal) the
origami surface. **b,** The overall reach of chol-oligos determines the migration behaviour of
origami triangles. When cholesterols are positioned closer to the outer edge and the
stacking contacts in a distal configuration (A_D), bands in an agarose gel appear shifted and
smeared compared to the same triangles not hybridised to chol-oligos. This effect could be
reduced by changing the orientation of the cholesterol to a proximal configuration with
respect to the origami, such that the reach of the cholesterol is minimised (A_P). When chol-
oligos are positioned closer to the centre of the structure, and thus further away from the
stacking contacts, the migration distance of these triangles is comparable to that of sample
A_P, albeit with more smearing. Combining both variants to double the cholesterol count per
triangle from 9 to 18 moieties yielded bands appearing as combinations of the individual
variants. **c,** The reach of cholesterol influences the assembly of cholesterol-decorated
triangles. An assembly control (AC) of triangles lacking protruding linker handles assembles
mostly into icosahedral species, as do triangles with protruding linker handles not
hybridised to chol-oligos. When chol-oligos are introduced, distal cholesterol (A_D, B_D)
promotes the formation of primarily octahedral species, whereas proximal cholesterol (A_P)
does not noticeably influence the assembly behaviour. Reducing the reach of individual
cholesterols by positioning them closer to the centre of the triangles and choosing proximal
configurations reduces unwanted interactions.
712

715 **Supplementary Figure 4 | The influence of cholesterol count per triangle in assembly**
716 **reactions containing GVs.** Whilst optimising cholesterol positioning on the triangle can
717 reduce unexpected assembly behaviour, introducing lipid vesicles yet again shifts the
718 assembly towards octahedra, the degree of which increases with the number of
719 cholesterols per triangle.
720

721
722
Supplementary Figure 5 | GVs shift the assembly of cholesterol-bearing triangles towards octahedral shells. The assembly behaviour of cholesterol-bearing triangles, tagged with Atto643 for visualisation, depends on the presence or absence of GVs. Without GVs, triangles mainly assemble into icosahedral shells as per their design. With GVs, however, most shells detected by agarose gel electrophoresis are octahedral. The abundance of smaller, intermediate species between closed shells and monomeric triangles increases with the number of cholesterols per triangle and mostly represents aggregated triangles.

732
733
Supplementary Figure 6 | GVs only alter the assembly behaviour of chol-bearing
triangles. a, Agarose gel of DNA shells made visible by Atto643-bearing
oligonucleotides.Triangles assemble predominantly into icosahedral shells even if either
GVs or chol-bearing oligonucleotides are present in the sample. However, if both GVs and
chol-bearing oligonucleotides are present in a sample at the same time, the assembly of
triangles is shifted towards octahedral species. **b,** The gel in a rescanned for visualisation of
DOPE-Atto488 species included in the vesicle membranes. Only the sample containing
chol-bearing triangles and GVs produced lipid bands coinciding with the DNA shell bands
in a, therefore indicating that the octahedral shells found in a are mostly octahedral DNA-
shell-coated vesicles. (M: triangle monomers)
744

745
746
Supplementary Figure 7 | Sodium chloride prevents non-specific adsorption of DNA
origami onto giant vesicles. Cross-sectional intensity profiles and channel-separated
confocal microscopy images of giant vesicles mixed with DNA origami triangles.
Conditions: +/- cholesterol-modified oligonucleotides (chol) and +/- 300 mM NaCl, as
indicated. At low-salt conditions, triangles adsorb onto lipid vesicles even if not hybridised
to chol-oligos. By adding 300 mM NaCl, non-specific origami-vesicle association is
suppressed without interfering with cholesterol-mediated association. Scale bars: 10 µm
754

755
756

Supplementary Figure 8 | DCVs obtained from triangles with 3, 6 or 9 cholesterols. Full
fields-of-view of DCV samples imaged using negative stain TEM. DCVs are encircled green
and empty shells are encircled red. Particles with ambiguous appearance, DCVs with
irregular shapes (e.g. particles larger than octa- and icosahedral particles), and mono- or
oligomeric triangles have not been marked. The micrographs were obtained under the same
conditions from triangles carrying either **a,** 3, **b,** 6, or **c,** 9 cholesterol moieties. Under
optimised conditions, DCVs make up the largest particle fraction. Scale bars: 100 nm.

765
766
Supplementary Figure 9 | Appearance of DCVs in TEM micrographs. a, Uranium stain
for negative stain TEM (ns-TEM) increases the contrast of biological samples, allowing for
easy differentiation between different shell species. Empty shells accumulate stain within
their cavity, rendering it black, while lipid vesicles within DCVs displaces the stain,
rendering the cavity white. **b,** Staining and subsequent drying of sample grids commonly
alters the appearance of lipid vesicles. Consequently, the vesicles within DCVs commonly
appear distorted, incomplete or broken in ns-TEM. **c,** Cryogenic electron microscopy
(cryoEM) allows imaging of DCVs in their native state without sample staining or drying
steps. DCVs can be distinguished from empty shells by a dark ring lining the inner cavity of
the DCV, representing the lipid vesicle. The artefacts shown in b are absent in cryoEM
micrographs. All scale bars: 100 nm
778

a empty octahedral shells

b octahedral DCVs

c octahedral & icosahedral DCVs

d octahedral budding

Supplementary Figure 10 | CryoEM images of octahedral DCVs. a, Appearance of
empty octahedral shells. **b,** Octahedral DCVs. Differences in appearance are due to
different orientations of the particles. **c,** Size comparison between an octahedral (left) and
icosahedral (right) DCV. As icosahedral particles are composed of 20 triangles (versus 8 in
octahedra), the inner cavity and the overall size of the particle are both larger than in
octahedral species. **d,** Budding of two octahedral DCVs. To the left of the two triangle-
covered buds is an empty octahedral shell. All scale bars: 100 nm.

Atto643   Atto488 composite
(DNA)   (DOPE)

788
789
Supplementary Figure 11 | Identification of DCVs in agarose gel electrophoresis by
colocalisation analysis. The relative efficiency of DCV formation can be analysed by
agarose gel electrophoresis using fluorescently labelled DNA oligos attached to the origami
structure or ethidium bromide, and a small fraction of fluorescently labelled DOPE species
in the lipid mixture comprising the vesicles. Whereas multiple bands can typically be
detected in the DNA channel, representing triangles in monomeric, oligomeric or fully
assembled states, lipid signal was only seen at the height of assembled shells. The absence
of lipid signal colocalised with monomeric triangles suggests that lipids predominantly enter
the gel as part of a DCV. The gel pockets contain origami-coated GVs, which were
oversaturated to better visualise the much fainter bands below.
800

801
802
Supplementary Figure 12 | Triangle assembly is the driving force behind DCV
formation. The importance of shell assembly for DCV formation was studied by incubating
triangles at various MgCl$_2$ concentrations at (65 mM) or below (10-40 mM) the typical
threshold concentration for shell assembly. For visualisation by agarose gel electrophoresis,
the triangles were hybridised to Atto643 bearing oligonucleotides, and the GVs contained a
fraction of Atto488 labelled DOPE. Two triangle species were prepared: A non-passivated
version with blunt-ended stacking contacts and a passivated version where staple
oligonucleotides at the base stacking contacts were extended by additional unpaired
thymidines. Whereas the non-passivated versions formed DCVs, as seen by the bands in
the lipid channel (lower panel) colocalising with the shells in the DNA channel (upper panel),
only faint bands below the typical height of DNA shells were found in samples containing
the passivated version. TEM analysis (right panels) revealed that while the passivated
triangle samples contained mostly monomeric triangles, there was a small number of
aggregated triangles held together by small vesicles, likely representing the faint bands. The
lack of DCVs in samples containing passivated triangles highlights the importance of shell
assembly for DCV formation. Scale bars: 100 nm.
819

Lipid channel (DOPE-Atto643)

820
821
Supplementary Figure 13 | Shell preassembly reduces DCV yields. Cholesterol-bearing
origami triangles were incubated without GVs for 4 h at 37 °C in either 5 mM (monomer
conditions) or 60 mM (assembly conditions) MgCl$_2$ to study the effect partially assembled
shells have on DCV yields. Afterwards, GVs (containing a small fraction of DOPE-Atto643)
were added, and the MgCl$_2$ concentration in the 5 mM controls was increased to 60 mM.
Aliquots taken and frozen at t = 0 d were compared to samples incubated at 37 °C for 2.5 d
using agarose gel electrophoresis, and DCV yields were analysed by comparing lipid
bands. Whereas t = 0 d aliquots looked alike in samples pre-incubated in either 5 mM or 60
mM MgCl$_2$, we finally obtained stronger lipid signals in samples initially incubated at 5 mM.
We deduce that DCVs form best when monomeric triangles are assembled directly on the
membrane. Partially assembled shells may fail to bind GVs efficiently due to steric
hindrance, resulting in reduced DCV yields.
834

835
836
Supplementary Figure 14 | Budding assays at varying tonicities (channel-separated).
Agarose gel of budding assays ran at different tonicities and visualised using two
fluorophores. The top image shows signal obtained from DNA origami stained by ethidium
bromide, while the bands in the bottom image stem from Atto643-labelled DOPE species
included in the lipid mixture of the GVs. Budding efficiency strongly correlates with tonicity,
working best under hypertonic conditions and coming to a near halt at the most hypotonic
condition tested. Refer to Figure 2c for a merged-colour image.
844

Supplementary Figure 15 | Membrane anchor screen (channel-separated). a, Agarose gel of DCVs prepared by using biotin-NeutrAvidin (b-NAv) interactions to connect origami triangles (Atto643, top image) and GVs (DOPE-Atto488, bottom image) containing a fraction of biotinylated DOPE. Linker handles on the triangles were complementary to b-NAv-oligos, but not to chol-oligos. Lipid material remains in the pockets unless triangles are mixed with b-NAv-oligos and GVs. The bands in the lipid channel colocalise with polyhedral shells but not with monomeric triangles, confirming the formation of DCVs. The addition of orthogonal chol-oligos did not improve yields, showing that the insertion of cholesterols into the bilayer is not a requirement. Gel pockets were oversaturated for better band visualisation. Refer to Figure 2d for a merged-colour image. **b,** Agarose gel of DCVs prepared under the same conditions as in a, using triangles with linker handles complementary to chol-oligos.

Supplementary Figure 16 | Budding assay using giant vesicles of varying lipid composition (channel-separated). Agarose gel of DCVs obtained from GVs of various lipid compositions. All compositions tested allowed budding of DCVs at varying efficiencies, except for GVs composed entirely of high-melting lipids. Top image: DNA (EtBr); Bottom image: Lipids (DOPE-Atto643); Refer to Figure 2e for a merged-colour image.

Supplementary Figure 17 | Budding assay from phase-separated vesicles (channel-separated). Agarose gel comparing DCVs obtained from GVs composed of DOPC or phase-separated ternary lipid mixtures. Phase-separated vesicles yielded more DCVs, suggesting an influence of phase boundaries on the budding mechanism. Top images: DNA (EtBr); Bottom images: Lipids (LissRhod-PE); Refer to Figure 2f for a merged-colour image.

873
874
875 **Supplementary Figure 18 | Budding kinetics (channel-separated).** DCV budding occurs
876 instantly and yields mostly octahedral species, but at longer incubation times a notable
877 fraction of icosahedral DCVs is formed. Top image: DNA (EtBr); Bottom image: Lipids
878 (LissRhod-PE); Refer to Figure 2g for a merged-colour image.

sc8064 sequence

```
GGCAATGACCTGATAGCCTTTGTAGATCTCTCAAAAATAGCTACCCTCTCCGGCATTAATTTATCAGCTAGAACGGTTGAATATCATATTGATGGTGATTTGACTGTCTCCG
GCCTTTCTCACCCTTTTGAATCTTTACCTACACATTACTCAGGCATTGCATTTAAAATATATGAGGGTTCTAAAAATTTTTATCCTTGCGTTGAAATAAAGGCTTCTCCCGC
AAAAGTATTACAGGGTCATAATGTTTTTGGTACAACCGATTTAGCTTTATGCTCTGAGGCTTTATTGCTTAATTTTGCTAATTCTTTGCCTTGCCTGTATGATTTATTGGAT
GTTAATGCTACTACTATTAGTAGAATTGATGCCACCTTTTCAGCTCGCGCCCCAAATGAAAATATAGCTAAACAGGTTATTGACCATTTGCGAAATGTATCTAATGGTCAAA
CTAAATCTACTCGTTCGCAGAATTGGGAATCAACTGTTATATGGAATGAAACTTCCAGACACCGTACTTTAGTTGCATATTTAAAACATGTTGAGCTACAGCATTATATTCA
GCAATTAAGCTCTAAGCCATCCGCAAAAATGACCTCTTATCAAAAGGAGCAATTAAAGGTACTCTCTAATCCTGACCTGTTGGAGTTTGCTTCCGGTCTGGTTCGCTTTGAA
GCTCGAATTAAAACGCGATATTTGAAGTCTTTCGGGCTTCCTCTTAATCTTTTTGATGCAATCCGCTTTGCTTCTGACTATAATAGTCAGGGTAAAGACCTGATTTTTGATT
TATGGTCATTCTCGTTTTCTGAACTGTTTAAAGCATTTGAGGGGGATTCAATGAATATTTATGACGATTCCGCAGTATTGGACGCTATCCAGTCTAAACATTTTACTATTAC
CCCCTCTGGCAAAACTTCTTTTGCAAAAGCCTCTCGCTATTTTGGTTTTTATCGTCGTCTGGTAAACGAGGGTTATGATAGTGTTGCTCTTACTATGCCTCGTAATTCCTTT
TGGCGTTATGTATCTGCATTAGTTGAATGTGGTATTCCTAAATCTCAACTGATGAATCTTTCTACCTGTAATAATGTTGTTCCGTTAGTTCGTTTTATTAACGTAGATTTTT
CTTCCCAACGTCCTGACTGGTATAATGAGCCAGTTCTTAAAATCGCATAAGGTAATTCACAATGATTAAAGTTGAAATTAAACCATCTCAAGCCCAATTTACTACTCGTTCT
GGTGTTTCTCGTCAGGGCAAGCCTTATTCACTGAATGAGCAGCTTTGTTACGTTGATTTGGGTAATGAATATCCGGTTCTTGTCAAGATTACTCTTGATGAAGGTCAGCCAG
CCTATGCGCCTGGTCTGTACACCGTTCATCTGTCCTCTTTCAAAGTTGGTCAGTTCGGTTCCCTTATGATTGACCGTCTGCGCCTCGTTCCGGCTAAGTAACATGGAGCAGG
TCGCGGATTTCGACACAATTTATCAGGCGATGATACAAATCTCCGTTGTACTTTGTTTCGCGCTTGGTATAATCGCTGGGGGTCAAAGATGAGTGTTTTAGTGTATTCTTTT
GCCTCTTTCGTTTTAGGTTGGTGCCTTCGTAGTGGCATTACGTATTTTACCCGTTTAATGGAAACTTCCTCATGAAAAAGTCTTTAGTCCTCAAAGCCTCTGTAGCCGTTGC
TACCCTCGTTCCGATGCTGTCTTTCGCTGCTGAGGGTGACGATCCCGCAAAAGCGGCCTTTAACTCCCTGCAAGCCTCAGCGACCGAATATATCGGTTATGCGTGGGCGATG
GTTGTTGTCATTGTCGGCGCAACTATCGGTATCAAGCTGTTTAAGAAATTCACCTCGAAAGCAAGCTGATAAACCGATACAATTAAAGGCTCCTTTTGGAGCCTTTTTTTTG
GAGATTTTCAACGTGAAAAAATTATTATTCGCAATTCCTTTAGTTGTTCCTTTCTATTCTCACTCCGCTGAAACTGTTGAAAGTTGTTTAGCAAAATCCCATACAGAAAATT
CATTTACTAACGTCTGGAAAGACGACAAAACTTTAGATCGTTACGCTAACTATGAGGGCTGTCTGTGGAATGCTACAGGCGTTGTAGTTTGTACTGGTGACGAAACTCAGTG
TTACGGTACATGGGTTCCTATTGGGCTTGCTATCCCTGAAAATGAGGGTGGTGGCTCTGAGGGTGGCGGTTCTGAGGGTGGCGGTTCTGAGGGTGGCGGTACTAAACCTCCT
GAGTACGGTGATACACCTATTCCGGGCTATACTTATATCAACCCTCTCGACGGCACTTATCCGCCTGGTACTGAGCAAAACCCCGCTAATCCTAATCCTTCTCTTGAGGAGT
CTCAGCCTCTTAATACTTTCATGTTTCAGAATAATAGGTTCCGAAATAGGCAGGGGGCATTAACTGTTTATACGGGCACTGTTACTCAAGGCACTGACCCCGTTAAAACTTA
TTACCAGTACACTCCTGTATCATCAAAAGCCATGTATGACGCTTACTGGAACGGTAAATTCAGAGACTGCGCTTTCCATTCTGGCTTTAATGAGGATTTATTTGTTTGTGAA
TATCAAGGCCAATCGTCTGACCTGCCTCAACCTCCTGTCAATGCTGGCGGCGGCTCTGGTGGTGGTTCTGGTGGCGGCTCTGAGGGTGGTGGCTCTGAGGGTGGCGGTTCTG
AGGGTGGCGGCTCTGAGGGAGGCGGTTCCGGTGGTGGCTCTGGTTCCGGTGATTTTGATTATGAAAAGATGGCAAACGCTAATAAGGGGGCTATGACCGAAAATGCCGATGA
AAACGCGCTACAGTCTGACGCTAAAGGCAAACTTGATTCTGTCGCTACTGATTACGGTGCTGCTATCGATGGTTTCATTGGTGACGTTTCCGGCCTTGCTAATGGTAATGGT
GCTACTGGTGATTTTGCTGGCTCTAATTCCCAAATGGCTCAAGTCGGTGACGGTGATAATTCACCTTTAATGAATAATTTCCGTCAATATTTACCTTCCCTCCCTCAATCGG
TTGAATGTCGCCCTTTTGTCTTTGGCGCTGGTAAACCATATGAATTTTCTATTGATTGTGACAAAATAAACTTATTCCGTGGTGTCTTTGCGTTTCTTTTATATGTTGCCAC
CTTTATGTATGTATTTTCTACGTTTGCTAACATACTGCGTAATAAGGAGTCTTAATCATGCCAGTTCTTTTGGGTATTCCGTTATTATTGCGTTTCCTCGGTTTCCTTCTGG
TAACTTTGTTCGGCTATCTGCTTACTTTTCTTAAAAAGGGCTTCGGTAAGATAGCTATTGCTATTTCATTGTTTCTTGCTCTTATTATTGGGCTTAACTCAATTCTTGTGGG
TTATCTCTCTGATATTAGCGCTCAATTACCCTCTGACTTTGTTCAGGGTGTTCAGTTAATTCTCCCGTCTAATGCGCTTCCCTGTTTTTATGTTATTCTCTCTGTAAAGGCT
GCTATTTTCATTTTTGACGTTAAACAAAAAATCGTTTCTTATTTGGATTGGGATAAATAATATGGCTGTTTATTTTGTAACTGGCAAATTAGGCTCTGGAAAGACGCTCGTT
AGCGTTGGTAAGATTCAGGATAAAATTGTAGCTGGGTGCAAAATAGCAACTAATCTTGATTTAAGGCTTCAAAACCTCCCGCAAGTCGGGAGGTTCGCTAAAACGCCTCGCG
TTCTTAGAATACCGGATAAGCCTTCTATATCTGATTTGCTTGCTATTGGGCGCGGTAATGATTCCTACGATGAAAATAAAAACGGCTTGCTTGTTCTCGATGAGTGCGGTAC
TTGGTTTAATACCCGTTCTTGGAATGATAAGGAAAGACAGCCGATTATTGATTGGTTTCTACATGCTCGTAAATTAGGATGGGATATTATTTTTCTTGTTCAGGACTTATCT
ATTGTTGATAAACAGGCGCGTTCTGCATTAGCTGAACATGTTGTTTATTGTCGTCGTCTGGACAGAATTACTTTACCTTTTGTCGGTACTTTATATTCTCTTATTACTGGCT
CGAAAATGCCTCTGCCTAAATTACATGTTGGCGTTGTTAAATATGGCGATTCTCAATTAAGCCCTACTGTTGAGCGTTGGCTTTATACTGGTAAGAATTTGTATAACGCATA
TGATACTAAACAGGCTTTTTCTAGTAATTATGATTCCGGTGTTTATTCTTATTTAACGCCTTATTTATCACACGGTCGGTATTTCAAACCATTAAATTTAGGTCAGAAGATG
AAATTAACTAAAATATATTTGAAAAAGTTTTCTCGCGTTCTTTGTCTTGCGATTGGATTTGCATCAGCATTTACATATAGTTATATAACCCAACCTAAGCCGGAGGTTAAAA
AGGTAGTCTCTCAGACCTATGATTTTGATAAATTCACTATTGACTCTTCTCAGCGTCTTAATCTAAGCTATCGCTATGTTTTCAAGGATTCTAAGGGAAAATTAATTAATAG
CGACGATTTACAGAAGCAAGGTTATTCACTCACATATATTGATTTATGTACTGTTTCCATTAAAAAAGGTAATTCAAATGAAATTGTTAAATGTAATTAATTTTGTTTTCTT
GATGTTTGTTTCATCATCTTCTTTTGCTCAGGTAATTGAAATGAATAATTCGCCTCTGCGCGATTTTGTAACTTGGTATTCAAAGCAATCAGGCGAATCCGTTATTGTTTCT
CCCGATGTAAAAGGTACTGTTACTGTATATCATCTGACGTTAAACCTGAAAATCTACGCAATTTCTTTATTTCTGTTTTACGTGCAAATAATTTTGATATGGTAGGTTCTA
ACCCTTCCATTATTCAGAAGTATAATCCAAACAATCAGGATTATATTGATGAATTGCCATCATCTGATAATCAGGAATATGATGATAATTCCGCTCCTTCTGGTGGTTTCTT
TGTTCCGCAAAATGATAATGTTACTCAAACTTTTAAAATTAATAACGTTCGGGCAAAGGATTTAATACGAGTTGTCGAATTGTTTGTAAAGTCTAATACTTCTAAATCCTCA
AATGTATTATCTATTGACGGCTCTAATCTATTAGTTGTTAGTGCTCCTAAAGATATTTTAGATAACCTTCCTCAATTCCTTTCAACTGTTGATTTGCCAACTGACCAGATAT
TGATTGAGGGTTTGATATTTGAGGTTCAGCAAGGTGATGCTTTAGATTTTTCATTTGCTGCTGGCTCTCAGCGTGGCACTGTTGCAGGCGGTGTTAATACTGACCGCCTCAC
CTCTGTTTTATCTTCTGCTGGTGGTTCGTTCGGTATTTTTAATGGCGATGTTTTAGGGCTATCAGTTCGCGCATTAAAGACTAATAGCCATTCAAAAATATTGTCTGTGCCA
CGTATTCTTACGCTTTCAGGTCAGAAGGGTTCTATCTCTGTTGGCCAGAATGTCCCTTTTATTACTGGTCGTGTGACTGGTGAATCTGCCAATGTAAATAATCCATTTCAGA
CGATTGAGCGTCAAAATGTAGGTATTTCCATGAGCGTTTTTCCTGTTGCAATGGCTGGCGGTAATATTGTTCTGGATATTACCAGCAAGGCCGATAGTTTGAGTTCTTCTAC
TCAGGCAAGTGATGTTATTACTAATCAAAGAAGTATTGCTACAACGGTTAATTTGCGTGATGGACAGACTCTTTTACTCGGTGGCCTCACTGATTATAAAAACACTTCTCAG
GATTCTGGCGTACCGTTCCTGTCTAAAATCCCTTTAATCGGCCTCCTGTTTAGCTCCCGCTCTGATTCTAACGAGGAAAGCACGTTATACGTGCTCGTCAAAGCAACCATAG
TACGCGCCCTGTAGCGGCGCATTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTCGCTTTCTTCCCTTCCT
TTCTCGCCACGTTCGCCGGCTTTCCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAAACTTGATTTGGGTGA
TGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAACCCT
ATCTCGGGCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTCGGAACCACCATCAAACAGGATTTTCGCCTGCTGGGGCAAACCAGCGTGGACCGCTTGCTGCAACTCTC
TCAGGGCCAGGCGGTGAAGGGCAATCAGCTGTTGCCCGTCTCACTGGTGAAAAGAAAAACCACCCTGGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTCA
TTAATGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCAGGCTTTACACTTTATGCTT
CCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATAACAATTTCACACAGGAAACAGCTATGACCATGATTACGAATTCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAA
GGCTCACGGACGCGAAGAACAGGCACGCGTGCTGGCAGAAACCCCCGGTATGACCGTGAAAACGGCCCGCCGCATTCTGGCCGCAGCACCATGGGATCTCCAACGCGTTGGATGCAGGCGCGCAGT
GACACTGCGCTGGATCGTCTGATGCAGGGGGCACCGGCACCGCTGGCTGCAGGTAACCCGGCATCTGATGCCGTTAACGATTTGCTGAACACACCAGTGTAAGGGATGTTTA
TGACGAGCAAAGAAACCTTTACCCATTACCAGCCGCAGGGCAACAGTGACCCGGCTCATACCGCAACCGCGCCCGGCGGATTGAGTGCGAAAGCGCCTGCAATGACCCCGCT
TACAAGTCCGGCACGTTCCGTTATGAGGATGTGCTCTGGCCGGAGGCTGCCAGCGACGAGACGAAAAAACGGACCGCGTTTGCCGGAACGGCAATCAGCATCGTTTAACTTT
ACCCTTCATCACTAAAGGCCGCCTGTGCGGCTTTTTTTACGGGATTTTTTTATGTCGATGTACACAACCGCCCAACTGCTGGCGGCAAATGAGCAGAAATTTAAGTTTGATC
CGCTGTTTCTGCGTCTCTTTTTCCGTGAGAGCTATCCCTTCACCACGGAGAAAGTCTATCTCTCACAAATTCCGGGACTGGTAAACATGGCGCTGTACGTTTCGCCGATTGT
TTCCGGTGAGGTTATCCGTTCCCGTGGCGGCTCCACCTCTGAAAGCTTGGCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAAAACCCTGGCGTTACCCAACTTAATCGC
CTTGCAGCACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGAATGGCGCTTTGCCTGGTTTC
CGGCACCAGAAGCGGTGCCGGAAAGCTGGCTGGAGTGCGATCTTCCTGAGGCCGATACTGTCGTCGTCCCCTCAAACTGGCAGATGCACGGTTACGATGCGCCCATCTACAC
CAACGTGACCTATCCCATTACGGTCAATCCGCCGTTTGTTCCCACGGAGAATCCGACGGGTTGTTACTCGCTCACATTTAATGTTGATGAAAGCTGGCTACAGGAAGGCCAG
ACGCGAATTATTTTTGATGGCGTTCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAATGCGAATTTTAACAAAATATTAACGTTTACAATTTAAATATTTGCTTA
TACAATCTTCCTGTTTTTGGGGCTTTTCTGATTATCAACCGGGGTACATATGATTGACATGCTAGTTTTACGATTACCGTTCATCGATTCTCTTGTTTGCTCCAGACTCTCA
```

Supplementary Table 1 | Sequence of scaffold sc8064.

experiment	reference figure	core mix	internal linker mix	external linker mix	internal functionalised strands	external functionalised strands
Vesicle deformation & buds in cryoEM; octahedral DCVs	Fig. 1D S. Fig. 10	C1	I1	E1	CHOL1	*no linkers*
Budding by tonicity	Fig. 2C & S. Fig. 14	X6-20 / E6_1 C1	I1	E1	CHOL1	*no linkers*
Budding by membrane anchor (b-NAv)	Fig. 2D & S. Fig. 15A	C1	I3	E2	BIOTIN	FLUO1 & bead purification linkers
Budding by lipid mixture	Fig. 2E & S. Fig. 16	X6-20 / E6_1 C1	I1	E1	CHOL1	*no linkers*
Budding from phase-separated vesicles	Fig. 2F & S. Fig. 17	X6_20 / E6_8 C1	I1	E3	CHOL1	*FLUO2*
Budding kinetics	Fig. 2G & S. Fig. 18	X6_20/E6_1 C1	I1	E1	CHOL1	*no linkers*
Inward budding	Fig. 3B, C	C1	I4	E4	*no linkers*	CHOL1
Bivesicular budding	Fig. 4B	C1	I1	E5	CHOL1	CHOL2 & FLUO1*
Cholesterol position screen	S. Fig. 3	C1	I5 or I6	E1	CHOL2 or CHOL3	*no linkers*
Cholesterol count screen	S. Fig. 4	C1	I7 or I8 or I9	E6	CHOL4	FLUO1
Influence of GVs on triangle assembly	S. Fig. 5	C1	I10 or I11 or I12	E6	CHOL4	FLUO1
Influence of GVs on triangle assembly 2	S. Fig. 6	X6_2/E6_2 C1	I10	E6	CHOL4	FLUO1
DCVs from triangles with 3/6/9 cholesterols	S. Fig. 8	X6_2/3/4 C1	I10 or I11 or I12	E6	CHOL4	FLUO1
DNA-lipid colocalisation in agarose gel electrophoresis	S. Fig. 11	C1	I10	E6	CHOL4	FLUO1
Assembly as the driving force for DCV budding	S. Fig. 12	C1 or C2	I2	E3	CHOL2	FLUO2
Preassembled shells	S. Fig. 13	C1	I1	E1	CHOL1	*no linkers*
Budding by membrane anchor (chol.)	S. Fig. 15B	C1	I2	E2	CHOL2	FLUO1 & bead purification linkers

881
Supplementary Table 2 | Triangle variants used in this study. For different experiments
(see 'experiment' and 'reference figure' columns), slight variations of the triangular origami
structure have been used differing mostly in the sequence, number and positioning of its
linker handles. For simplicity, the final staple mixture was divided into three sub-mixtures
comprising a constant part (core mix), staples with linker handle extensions on the shell-
inner face (internal linker mix), and staples with linker handle extensions on the shell-outer
face (external linker mix). For foldings, these mixtures were mixed according to their staple
count (equal staple concentrations in the final mixture). The corresponding complementary
functionalised oligonucleotides (see Supplementary Table 3 for sequences), if any, for both
faces are listed in additional columns. Note that cholesterol-bearing strands were not
included in folding reactions but added just before mixing with vesicle solutions (see

Materials & Methods section). Refer to the following supplementary tables for sequences.
*Linker handles for FLUO1 were left single-stranded (no fluorescence labelling was
required).

functionalised strands	sequence	modification
CHOL1	CTCCACCTCTCACTCTTT	3'-(cholesterol-TEG)
CHOL2	ATCATCTCACCTTTT	3'-(cholesterol-TEG)
CHOL3	TTTTCCACTCTACTA	5'-(cholesterol-TEG)
CHOL4	ATCATCTCACCTTTT	3'-(cholesterol-prolinol)
FLUO1	CCAAGCTACAGCCTGAAA	3'-(Atto643)
FLUO2	GCCAGTCCGCCTCGCAAA	3'-(Atto643)
BIOTIN	TACGCCGGAAGCAGC	3'-(biotin-TEG)
purification linker handle	see Supplementary Table 4	

897   **Supplementary Table 3 | List of functionalised strands.**

functionalised strands	sequence	modification
CHOL1	CTCCACCTCTCACTCTTT	
CHOL2	ATCATCTCACCTTTT	
CHOL3	TTTTCCACTCTACTA	
CHOL4	ATCATCTCACCTTTT	
FLUO1	CCAAGCTACAGCCTGAAA	
FLUO2	GCCAGTCCGCCTCGCAAA	
BIOTIN	TACGCCGGAAGCAGC	

Strand	sequence	modification
Bead handle	TTTCTATCACCTCCA	5'-(dual biotin)
Origami handle	--TTTGATGTGTAGAGTTGGAGGTGATAG	
Invader strand	CCTTCAATCTTACTATCACCTCCAACTCTACACATC	

899 **Supplementary Table 4 | List of sequences for magnetic bead purification of DNA**
900 **origami.** Bead handles are coupled to Dynabeads M-270 Streptavidin. The stated
901 sequence of the origami handle is part of a staple with the full sequences listed in
902 Supplementary Table 20. Complementary sequence snippets are highlighted. The invader
903 strands can optionally be removed from the origami by its reverse complement.

Bead handle	TTTCTATCACCTCCA	5'-(dual biotin)
Origami handle	--TTTGATGTGTAGAGTTGGAGGTGATAG	
Invader strand	CCTTCAATCTTACTATCACCTCCAACTCTACACATC	

No.	C1 sequences (non-passivated triangles)
01	GGAAACCACGGTGCGGGCCTCTTCAGCCCAATGTATAAGCAAAAGCCC
02	GCTATTACTAAAATTCGCATTGCTTTAAACAGTT
03	CGCCACCCTCAGAACCGGAATAGCGCAATAATAGGACTGTAGCGCGT
04	AATACCCAACGGAATACAAGATAGCAGCACCG
05	AGTGTACCCCGGTTGCTGCTCCAATTTTCAGGGATAGCA
06	TGTTACTTAGCCGCATTAGACGGAAGACACCAAAGAACT
07	TTAATGCCAGAATCGAGAATATAAACGGTGTACCAACTTTGAAAGAGG
08	GCAGACGGTCAATCATTCCGCGACATAATCAGAAATATTTCTCAGAG
09	AAAATAGCAAAGTCAGAGGAGAAAACTTTTTC
10	AGCCCGAGGAAGATTAGGAACCCATAAATTGTGTCGAAAAAGGGAAC
11	GCCGAACAAAGTTACCAGTAAGACTCAGAGAGATACTGAACA
12	GCCTTTAGCGTCACAAAGGATAAAAATTTTTAGAACCCT
13	TCAAGTTTGGCATGATAAGGAAACAGAGCAAGTTTAGTACACGTTGT
14	AATAAAGCAAATTGTAAACGCGGCCACCCTC
15	TTTTGTCAGGATATCAGGTCATTGCCTGAGAACCCAGCT
16	CCAGTAGCACCGTCACCGACTTGAAACGCTAAATAACATA
17	TGGGTTATATGGAAACAGTACATACCAAGCTTTCAGAGGTGGAGCCG
18	TGTCAATCATATTTAAACGAGGCACAATTTTTTAGTTGC
19	AGAACGCGAGGCGTTTTAGCGAACGGCAGAGG
20	GTCTGGAGAACTAGCAGACAGTCATGTAATACTTTTGAGG
21	CTTGCGGGAGGTTTTGAAGCCTTAAAAAGCCTAAAACAGGGAAGCGG
22	TATTTTGCACCTTCATCAAGAGTAGCGCATAGGCTGGCTG
23	CAAAAATGCATTTTCGTTTGAAATACCGACCGGCTTAGGTATTGAGCG
24	CATATATTAATCAGTAGCGAGGAACCGCCT
25	ACCCCCAGCGATTATATTACCCAAATTCATCAACAGATGA
26	AGCCTTTATTTCAACGCAAATTAAGCTAAAGATT
27	TTTAAATGCAATGCCTGAGTAATGTGTCGGGAGA
28	ATATAAAGTATTTTTCGACAAAAGTTCATTTTTCTTCTGACCTAAA
29	TTAAAGGTGCCAAAGACAAAAGGGCTTTTTACATTCAACCGATTGA
30	AGGTAAATATTGTTTTTCGGAAATTATTCATGCCAGTT
31	TTATTTATCCCCCAATAGCAAGCAAATCAGATTTTTTTAGAAGGCTTATCCG
32	GCTAATGCAGAACGCGCATCAAGAAAACACAAGACTTTTTAAGAACGCGGTA
33	CCACGGGATTTCATTTGATTTTTTTACCTTTTTTAATAACTAT
34	TTTCATCGGCATTTTTTTTCGGACCAGAGCCTTTTTCCACCCAGAA
35	AAATATCAATAAGAAACGATTTTTTTTTTTGTTTAACGTACAAAATA
36	AGCCGCCGCAGGTCAGACGATTGGCTTTTTTTGACATTAGCAAGGCCGGA
37	AACGTAGAAAATTTTTACATACATAAAGGCAGTAT
38	ATGTAAATTTTTTCTGATGCAAATCCAATGTTATTTTAGTTAAT
39	GCTCATGGAGCACCAGCAGAAGATAAAACAGAGGTGACAAATCAACA
40	GTAAAGTAATTCTGTCTTACCGCGCAATCCAACATTCCAA
41	TCACCCAAGAGAACAAGCAAGCCGAAGTACCGCACTCATC
42	AATGCGCGAAGAATACGTGGCACAGCACTAAAGCGCGTAA
43	GCCAGCGGGAGGTGTCCAGCATCAATCCCGTAAAAAAAGCCGCACAG
44	TCGGCGAACTCCGTGGTGAAGGGAAAGATGATAACCTATTGTTAATGC
45	TGATTGCTTTGAATAGTACCTTTCCGTTGTAGCCACAGTGCCACGCT
46	CCCTCAGTTTTCATACCCTCAGAAAATAACAGTGCCCGCCTGAACAATTACC
47	TGAGCAAAAGTAGCTCTCACTCCTCAAGAGAAAGTAAGCAGATA
48	AACCACCACCAGGCCACCA
49	AGAAAAATAATCCGCGCTTAATGGGAAAAACATTAGTAA
50	ATTTACGAGCATGTAGAGATAAGTTATAAACAATTCTGAACACCGGA
51	GCTGGCAAGTTGCTTTGACTGCACTCTGTGGT
52	GCTAGGGCAGCTAACTGGTTTCTGCCAGCACGTACCTGCACACGTATA
53	TCAGTATTTAACATCACCATCACGTTAAAGGGGATTCGCCGTTGTGT
54	AAAAACCGTCTATCAGGGCGATGGCCGAGCTCCCACCACACCCGATC
55	TCCTCATTAGGAGTGTGGGAGGGGAGAACGGGTGGCTGTCTTTCCTTAT
56	TGGCTTTTAACGGGGTAGGTTGAGACCCTCAGAACCGCCA
57	TAAAGAACGTGGACTCCAACGTCAAATGAGTG
58	CAGTGCCTTGAGTACCCATCCTATTTTGGGGGTGAACCA
59	CTACATTTTTTGTTCCAGTAAGCGTCATACAATCAAGTT
60	ATCAAAATACATGAAAGTATATATCATTTCA
61	AACACCGCCTGCAATCAGCCGCCACCAG
62	GTTAGCATATTTCGGGAAACAAACCTGTTTATCAACAATAAACCAAT
63	CCGAGTAAAAGAGTCTGTCTTGCCTGTCCTCGTTCAGGGCGC
64	GCCTTGCTGGTTTTTTATATCCAGAACAAACTCAA
65	GCGCGGGGAGAGGCGGGAATGCCAACGGCTGTCACTTTTTGCGCGCCTGGAG
66	GCGGCCTTCCCACGCAACTTTTTAGCTTACGGCTGTGCCGGTG
67	CCCCCTGCTTTTTTCAGACGATCCAGCGGCCGGCCAGAATGCGG
68	GCTGCAACAAGGGAAGAAAGTTTTTGAAAGGAGCGGGCCTTGACGG
69	GGAATTGGTCAATAGATAATACATTTTTTGAGGGCCCTAAAACATCGCC
70	CGCTCACTGCTTTTTCGCTTTCCACGGGTTTTTCGTTTTCACGGTC
71	CGAACGTGGCGAGATAGGGTTGAGTGTTGTTCTTTTTAGTTTGGAACAAGAG
72	GAGAGCCAGCAGCTTTTTAATGTATCTGGTCTTTTTGTTGGGGCGG
73	TCTGACCTATTGGCAGATTCACCAGTTTTTCACACGACCAGTAATA
74	ATTCTGGCCAACTTTTTGAGATAGAACCCTATTTAGAG
75	AACTCCAAAGGTGTTTAGCTATATTTTCATTTGGGGCGTTGTACCAA
76	CAGTTCCTGATTATCTTTTTCTTCGTGGTGCTGGTCTGG
77	CGAGAAACACCAGAACGAGTAGTACACCAACC
78	ACTAATGCTAAAACGACGATATATTCGGTCGCAAAAGGCTGAGGCTTT
79	AGTAACATGGAGCGGATTAGAGCCAGGAAGGTTATCTAAA
80	AATAAAGACGTGGGAATGCCGGACTTGATGTTTAGACTG

81	CAGAAAACGAGAATGACCTTGCATCAAAGTTTTGACCAGACG
82	CAAACCCTCTGAATAATGGAGACAACGTCAG
83	GTGGCATCAATTCACCTCAAATATATATCTTT
84	CCTTATGCGGTGGTTCCGAAATCGGAAAATCCTGTTTGAT
85	ACTATCGATCCTGATAACCGCAATTTGCGTATTGGGCGCCGCCTGGC
86	CTGAAAAGAAAGCGGAATAAATCAGATAGCGTGAGTAACAAGTTTGA
87	AGATACATAACCAAAATAGACAGCTTGATACC
88	GCTGATTGCCCTTCACCAGGGTGGAGATGATGTTATACTT
89	TCCTCATATACCATATCAAAAATCAGTGAGGCCA
90	AACGAGTATGTCTGGAAGTTTCATGGACGTTGACGAGGCA
91	GTCGGGAAACCTGTCGAATAGCCCGAGAAAGGCCACGCTG
92	CCAAAAGGCACAGACAGCCCTCATTCGGCCTCAGGAAGATCGCACTC
93	AGGATTAGACATTATTAATTTTAAAAGTTTGGATTTTAA
94	CTTGAGATGGTTTAATTTCAACTTCGAAAGACTAGTAAGAGCAAGAC
95	CGATGCTGCTCCGGCCAGAGCACATCAGCAGCTGTTTGGAGCAATTCA
96	ATTATCATCATAGTCGGGCAACAGAACTGGCGTGAATTA
97	TTCACCAGTGACACTATCATAACCGGAAGCATCAGAAGC
98	TTACAAACTGCGGAACAAAGGGACGTTTGCCCAGTTGCAGCAAGCGGT
99	AGCATTAACATCCTTTTTATAACATAAAGCTTTTTTAATCGGCGAG
100	ATGACCCAATCACCATCAATATGATTTTTTTTCATTGACCATTAGATACA
101	AAGACTTCAAATTTTTATCGCGTTTTAAAAGAGGA
102	CACTAAAACATTTTTTCATCTTTGGACATTTTTCAACCATCGCCCA
103	TGCAACTAGTCATTTTTGCGGATGGTTTTTTTAGAGCTTAATTGCT
104	GATAGAGAGTTGAGATTTAGTTTTTAATACCACATTCAACGGAACA
105	ATTGTATCTTTTTGTTTATCAGCTTGCTCGGTTGCGCCGACAAT
106	TGCTGTAGCTCATTTTTCATGTTTTAAATAACGAACTA
107	TTTGTATCATCGCCTGATGTACCGTAACAGAGGTGTTTTTATTTCTTAACGA
108	CAGCCAGCAAACTACAACTTTTTCCTGTAGCATTCAGCCTTTA
109	AGGTAGAAAGATCAACGTAACAAAGCTGCTCATTTTTTCAGTGAATAAGGCT
110	ACAAAGTACAATTCCTGAGTTT
111	CGCCACCCTCAGAAGCAGTTGGGTAACGCCAGGGTTTTCCAATAGTGAATT
112	TATCAAAATCATAGGTCTGAGAATTGAGTTACCA
113	TAATTACTAGAAATCAAGAATCCTGAATCTTACCGCCATTTGCAATCAAT
114	ATCTTGACAAGAACCGGATATTCACCCAAGCGCGAA
115	GTATAAAGCCAACAGAATAAACACCGGAATCA
116	GTTTAGTAGTTAAATAGCTCAACAAGAATCCTTGAAAACATAGCG
117	ATAAGGCTCAACATCGCCATATGCGTTATACAAATTCTTACCA
118	CCATATTTAACAAGTAATTTACTCCCGATTTCCAGAGCCTAATT
119	AGAGACTATGACTGAAGACGATAACCTTGCTTCTGTAAATCGTC
120	GCTATTAATTAATAACCTCCGTGTGATAACCCTGAACAGCCTTTA
121	ATAGCTTAGATTAGAAGAGTCCAGTCACG
122	TATGTGAGCCTTTTTTTTCCCTTGTAGGGCTTAATTGAGAATCG
123	CTGGTGTGTTCAGCAAATCAGCGGGAGCTAATATCTTCTTTG
124	CCAAGTTACAAAAGAAATTTCTGCTCATTTGCCGCCAGCACATCCCTTACA
125	ATCCCCGGGTACCCACTACTCGAGGTGCCGTAAAGACAATATTGACGCTC
126	GAAATTGTTATCCAGCCTCCTCACAGTTGAGG
127	TTTTTATTTTCATCGTAGGAATCACAGACGACGAC
128	AATAAACAACATCGAAATTAAT
129	GAAGCATAAAGTGGGGTGCCTAAGGGCGCTAAAGGGAGCCCCCG
130	GGTTGCGGTATGATGCCGGGTCGTGCCTGGTACTATGGTGTAGCG
131	GAATTCGTCGTCCGTGGCTCACAAACTGTTGCCCTGCGGCTGGTA
132	ATGGGTAAAGGTTGTCATAAAGTTGGGCG
133	GTTAACGGCGCGCTCTCTTTTTTTCGCACTCAATCCGCCGGGCGC
134	ATTGCAGGCATCAGAGCCGGGTCTTCCACACAACATACGAGCCG
135	TTCTTCGAATCCTGTAAAGCATGGTCATAGCTGTTTCCTGTGT
136	TTTGAGGACTAAACCGCTTTTGCGGGATCGTC
137	GAATTGCGAATAATAATTTGGTAATAGTAAATAGTATTATAG
138	GTAGATGGGCGCATCGTAACCGTGAACAACTAAAG
139	ACCCTCAGCAGTAATCATTTCATTATACCAGTCATCCATATAAGAGTACC
140	GCAAAATCCCTTATAAATCAAAAGTGCCAGCTGCA
141	TTAATGAATCGCAGAGCACCGT
142	AACAGTTTCAGCGTAGAAAGGCATCTGCC
143	TTTCACGTTAAAGAAGAGTGAGTTTTGTCGTCTTTCCAGACGTT
144	AGCATCGGGTTAAAGGGACTTTTTGGATTTTGCTAAACAACTTTC
145	CGGGTAAAATACGTACGAAGGAATTGGGAATCTACGTTAATAAA
146	GCAGGGAAACCCACTAATGGAGGGTAGCAACGGCTACAGAGGC
147	TAACGATCTGAAAATTCTGTATGCATGAGGAAGTTTCCATTAAA
148	AGTAAATGAATTTCTCCAAAATGAGGCTTACGATAAAAACGCCAA

Supplementary Table 5 | List of staples in core mix 1 (C1). This staple mixture was used
to fold non-passivated triangles capable of self-assembly into shells by shape-
complementary base-stacking interactions.

No.	C2 sequences (passivated triangles)
01	GGAAACCACGGTGCGGGCCTCTTCAGCCCAATGTATAAGCAAAAGCCC
02	GCTATTACTAAAATTCGCATTGCTTTAAACAGTT
03	CGCCACCCTCAGAACCGGAATAGCGCAATAATAGGACTGTAGCGCGT
04	AATACCCAACGGAATACAAGATAGCAGCACCG
05	AGTGTACCCCGGTTGCTGCTCCAATTTTCAGGGATAGCA
06	TGTTACTTAGCCGCATTAGACGGAAGACACCAAAGAACT
07	TTAATGCCAGAATCGAGAATATAAACGGTGTACCAACTTTGAAAGAGG
08	GCAGACGGTCAATCATTCCGCGACATAATCAGAAATATTTCTCAGAG
09	AAAATAGCAAAGTCAGAGGAGAAAACTTTTTC
10	AGCCCGAGGAAGATTAGGAACCCATAAATTGTGTCGAAAAAGGGAAC
11	GCCGAACAAAGTTACCAGTAAGACTCAGAGAGATACTGAACA
12	GCCTTTAGCGTCACAAAGGATAAAAATTTTTAGAACCCT
13	TCAAGTTTGGCATGATAAGGAAACAGAGCAAGTTTAGTACACGTTGT
14	AATAAAGCAAATTGTAAACGCGGCCACCCTC
15	TTTTGTCAGGATATCAGGTCATTGCCTGAGAACCCAGCT
16	CCAGTAGCACCGTCACCGACTTGAAACGCTAAATAACATA
17	TGGGTTATATGGAAACAGTACATACCAAGCTTTCAGAGGTGGAGCCG
18	TGTCAATCATATTTAAACGAGGCACAATTTTTTAGTTGC
19	AGAACGCGAGGCGTTTTAGCGAACGGCAGAGG
20	GTCTGGAGAACTAGCAGACAGTCATGTAATACTTTTGAGG
21	CTTGCGGGAGGTTTTGAAGCCTTAAAAAGCCTAAAACAGGGAAGCGG
22	TATTTTGCACCTTCATCAAGAGTAGCGCATAGGCTGGCTG
23	CAAAAATGCATTTTCGTTTGAAATACCGACCGGCTTAGGTATTGAGCG
24	CATATATTAATCAGTAGCGAGGAACCGCCT
25	ACCCCCAGCGATTATATTACCCAAATTCATCAACAGATGA
26	AGCCTTTATTTCAACGCAAATTAAGCTAAAGATT
27	TTTAAATGCAATGCCTGAGTAATGTGTCGGGAGA
28	ATATAAAGTATTTTTCGACAAAAGTTCATTTTTCTTCTGACCTAAA
29	TTAAAGGTGCCAAAGACAAAAGGGCTTTTTACATTCAACCGATTGA
30	AGGTAAATATTGTTTTTCGGAAATTATTCATGCCAGTT
31	TTATTTATCCCCCAATAGCAAGCAAATCAGATTTTTTTAGAAGGCTTATCCG
32	GCTAATGCAGAACGCGCATCAAGAAAACACAAGACTTTTTAAGAACGCGGTA
33	CCACGGGATTTCATTTGATTTTTTTACCTTTTTTAATAACTAT
34	TTTCATCGGCATTTTTTTTCGGACCAGAGCCTTTTTCCACCCAGAA
35	AAATATCAATAAGAAACGATTTTTTTTTTGTTTAACGTACAAAATA
36	AGCCGCCGCAGGTCAGACGATTGGCTTTTTTTGACATTAGCAAGGCCGGA
37	AACGTAGAAAATTTTTACATACATAAAGGCAGTAT
38	ATGTAAATTTTTTCTGATGCAAATCCAATGTTATTTTAGTTAAT
39	GCTCATGGAGCACCAGCAGAAGATAAAACAGAGGTGACAAATCAACA
40	GTAAAGTAATTCTGTCTTACCGCGCAATCCAACATTCCAA
41	TCACCCAAGAGAACAAGCAAGCCGAAGTACCGCACTCATC
42	AATGCGCGAAGAATACGTGGCACAGCACTAAAGCGCGTAA
43	GCCAGCGGGAGGTGTCCAGCATCAATCCCGTAAAAAAAGCCGCACAG
44	TCGGCGAACTCCGTGGTGAAGGGAAAGATGATAACCTATTGTTAATGC
45	TGATTGCTTTGAATAGTACCTTTCCGTTGTAGCCACAGTGCCACGCT
46	CCCTCAGTTTTCATACCCTCAGAAAATAACAGTGCCCGCCTGAACAATTACC
47	TGAGCAAAAGTAGCTCTCACTCCTCAAGAGAAAGTAAGCAGATA
48	AACCACCACCAGGCCACCA
49	AGAAAAATAATCCGCGCTTAATGGGAAAAACATTAGTAA
50	ATTTACGAGCATGTAGAGATAAGTTATAAACAATTCTGAACACCGGA
51	GCTGGCAAGTTGCTTTGACTGCACTCTGTGGT
52	GCTAGGGCAGCTAACTGGTTTCTGCCAGCACGTACCTGCACACGTATA
53	TCAGTATTTAACATCACCATCACGTTAAAGGGGATTCGCCGTTGTGT
54	AAAAACCGTCTATCAGGGCGATGGCCGAGCTCCCACCACACCCGATC
55	TCCTCATTAGGAGTGTGGGAGGGAGAACGGGTGGCTGTCTTTCCTTAT
56	TGGCTTTTAACGGGGTAGGTTGAGACCCTCAGAACCGCCA
57	TAAAGAACGTGGACTCCAACGTCAAATGAGTG
58	CAGTGCCTTGAGTACCCATCCTATTTTGGGGGTGAACCA
59	CTACATTTTTTGTTCCAGTAAGCGTCATACAATCAAGTT
60	ATCAAAATACATGAAAGTATATATCATTTCA
61	AACACCGCCTGCAATCAGCCGCCACCAG
62	GTTAGCATATTTCGGGAAACAAACCTGTTTATCAACAATAAACCAAT
63	CCGAGTAAAAGAGTCTGTCTTGCCTGTCCTCGTTCAGGGCGC
64	GCCTTGCTGGTTTTTTATATCCAGAACAAACTCAA
65	GCGCGGGGAGAGGCGGGAATGCCAACGGCTGTCACTTTTTGCGCGCCTGGAG
66	GCGGCCTTCCCACGCAACTTTTTAGCTTACGGCTGTGCCGGTG
67	CCCCCTGCTTTTTTCAGACGATCCAGCGGCCGGCCAGAATGCGG
68	GCTGCAACAAGGGAAGAAAGTTTTTGAAAGGAGCGGGCCTTGACGG
69	GGAATTGGTCAATAGATAATACATTTTTTGAGGGCCCTAAAACATCGCC
70	CGCTCACTGCTTTTTCGCTTTCCACGGGTTTTTCGTTTTCACGGTC
71	CGAACGTGGCGAGATAGGGTTGAGTGTTGTTCTTTTTAGTTTGGAACAAGAG
72	GAGAGCCAGCAGCTTTTTAATGTATCTGGTCTTTTTGTTGGGGCGG
73	TCTGACCTATTGGCAGATTCACCAGTTTTTCACACGACCAGTAATA
74	ATTCTGGCCAACTTTTTGAGATAGAACCCTATTTAGAG
75	AACTCCAAAGGTGTTTAGCTATATTTTCATTTGGGGCGTTGTACCAA
76	CAGTTCCTGATTATCTTTTTCTTCGTGGTGCTGGTCTGG
77	CGAGAAACACCAGAACGAGTAGTACACCAACC
78	ACTAATGCTAAAACGACGATATATTCGGTCGCAAAAGGCTGAGGCTTT
79	AGTAACATGGAGCGGATTAGAGCCAGGAAGGTTATCTAAA
80	AATAAAGACGTGGGAATGCCGGACTTGATGTTTAGACTG

81	CAGAAAACGAGAATGACCTTGCATCAAAGTTTTGACCAGACG
82	CAAACCCTCTGAATAATGGAGACAACGTCAG
83	GTGGCATCAATTCACCTCAAATATATATCTTT
84	CCTTATGCGGTGGTTCCGAAATCGGAAAATCCTGTTTGAT
85	ACTATCGATCCTGATAACCGCAATTTGCGTATTGGGCGCCGCCTGGC
86	CTGAAAAGAAAGCGGAATAAATCAGATAGCGTGAGTAACAAGTTTGA
87	AGATACATAACCAAAATAGACAGCTTGATACC
88	GCTGATTGCCCTTCACCAGGGTGGAGATGATGTTATACTT
89	TCCTCATATACCATATCAAAAATCAGTGAGGCCA
90	AACGAGTATGTCTGGAAGTTTCATGGACGTTGACGAGGCA
91	GTCGGGAAACCTGTCGAATAGCCCGAGAAAGGCCACGCTG
92	CCAAAAGGCACAGACAGCCCTCATTCGGCCTCAGGAAGATCGCACTC
93	AGGATTAGACATTATTAATTTTAAAAGTTTGGATTTTAA
94	CTTGAGATGGTTTAATTTCAACTTCGAAAGACTAGTAAGAGCAAGAC
95	CGATGCTGCTCCGGCCAGAGCACATCAGCAGCTGTTTGGAGCAATTCA
96	ATTATCATCATAGTCGGGCAACAGAACTGGCGTGAATTA
97	TTCACCAGTGACACTATCATAACCGGAAGCATCAGAAGC
98	TTACAAACTGCGGAACAAAGGGACGTTTGCCCAGTTGCAGCAAGCGGT
99	AGCATTAACATCCTTTTTATAACATAAAGCTTTTTTAATCGGCGAG
100	ATGACCCAATCACCATCAATATGATTTTTTTTCATTGACCATTAGATACA
101	AAGACTTCAAATTTTTATCGCGTTTTAAAAGAGGA
102	CACTAAAACATTTTTTCATCTTTGGACATTTTTCAACCATCGCCCA
103	TGCAACTAGTCATTTTTGCGGATGGTTTTTTTAGAGCTTAATTGCT
104	GATAGAGAGTTGAGATTTAGTTTTTAATACCACATTCAACGGAACA
105	ATTGTATCTTTTTGTTTATCAGCTTGCTCGGTTGCGCCGACAAT
106	TGCTGTAGCTCATTTTTCATGTTTTAAATAACGAACTA
107	TTTGTATCATCGCCTGATGTACCGTAACAGAGGTGTTTTTATTTCTTAACGA
108	CAGCCAGCAAACTACAACTTTTTTCCTGTAGCATTCAGCCTTTA
109	AGGTAGAAAGATCAACGTAACAAAGCTGCTCATTTTTTTCAGTGAATAAGGCT
110	ATCTTGACAAGAACCGGATATTCACCAAGCGCGAATTTTT
111	TTTTTTATCAAAATCATAGGTCTGAGAATTGAGTTACCA
112	CGCCACCCTCAGAAGCAGTTGGGTAACGCCAGGGTTTTCCAATAGTGAATTTTTT
113	TTTTTCGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
114	TTTTTACAAAGTACAATTCCTGAGTTTTTTTT
115	TTTTTTAATTACTAGAAATCAAGAATCCTGAATCTTACCGCCATTTGCAATCAAT
116	TTTTTCCATATTTAACAAGTAATTTACTCCCGATTTCCAGAGCCTAATT
117	ATAAGGCTCAACATCGCCATATGCGTTATACAAATTCTTACCATTTTT
118	TTTTTGCTATTAATTAATAACCTCCGTGTGATAACCCTGAACAGCCTTTA
119	TATGTGAGCCTTTTTTTTCCCTTGTAGGGCTTAATTGAGAATCGTTTTT
120	TTTTTATAGCTTAGATTAGAAGAGTCCAGTCACG
121	AGAGACTATGACTGAAGACGATAACCTTGCTTCTGTAAATCGTCTTTTT
122	GTTTAGTAGTTAAAATAGCTCAACAAGAATCCTTGAAAACATAGCGTTTTT
123	TTTTTCTGGTGTGTTCAGCAAATCAGCGGGAGCTAATATCTTCTTTG
124	TTTTTTACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
125	TTTTTGAAATTGTTATCCAGCCTCCTCACAGTTGAGGTTTTT
126	CCAAGTTACAAAAGAAATTTCTGCTCATTTGCCGCCAGCACATCCCTTACATTTTT
127	TTTTTAATAAACAACATCGAAATTAATTTTTT
128	TTTTTATCCCCGGGTACCCACTACTCGAGGTGCCGTAAAGACAATATTGACGCTC
129	TTCTTCGAATCCTGTAAAGCATGGTCATAGCTGTTTCCTGTGTTTTTT
130	ATTGCAGGCATCAGAGCCGGGTCTTCCACACAACATACGAGCCGTTTTT
131	TTTTTGGTTGCGGTATGATGCCGGGTCGTGCCTGGTACTATGGTGTAGCG
132	GAATTCGTCGTCCGTGCGTCACAAACTGTTGCCCTGCGGCTGGTATTTTT
133	GTTAACGGCGCGCTCTCTTTTTTCGCACTCAATCCGCCGGGCGCTTTTT
134	TTTTTGAAGCATAAAGTGGGGTGCCTAAGGGCGCTAAAGGGAGCCCCCG
135	TTTTTATGGGTAAAGGTTGTCATAAAGTTGGGCG
136	TTTTTGAATTGCGAATAATAATTTGGTAATAGTAAATAGTATTATAG
137	TTTTTTTAATGAATCGCAGAGCACCGTTTTT
138	GCAAAATCCCTTATAAATCAAAAGTGCCAGCTGCATTTTT
139	TTTTTACCCTCAGCAGTAATCATTTCATTATACCAGTCATCCATATAAGAGTACC
140	TTTTTTTTGAGGACTAAACCGCTTTTGCGGGATCGTCTTTTT
141	GTAGATGGGCGCATCGTAACCGTGAACAACTAAAGTTTTT
142	GCAGGGAAACCCACTAATGGAGGGTAGCAACGGCTACAGAGGCTTTTT
143	TTTTTAGTAAATGAATTTCTCCAAAATGAGGCTTACGATAAAAACGCCAA
144	TTTTTAACAGTTTCAGCGTAGAAAGGCATCTGCC
145	TTTTTCGGGTAAAATACGTACGAAGGAATTGGGAATCTACGTTAATAAA
146	TAACGATCTGAAAATTCTGTATGCATGAGGAAGTTTCCATTAAATTTTT
147	TTTCACGTTAAAGAAGAGTGAGTTTTGTCGTCTTTCCAGACGTTTTTTT
148	AGCATCGGGTTAAAGGGACTTTTTGGATTTTGCTAAACAACTTTCTTTTT

Supplementary Table 6 | List of staples in core mix 2 (C2). This staple mixture was used
to fold passivated triangles incapable of self-assembly into shells by shape-complementary
base-stacking interactions due to protruding dT nucleotides at the stacking contacts.

No.	I1 sequences (internal mix 1)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
04	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
05	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
06	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAA
07	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
08	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
09	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
10	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
11	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
12	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
13	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
14	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTT
15	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
16	CATTCGCCATGTCATAAATATTCAGCCCAATACT
17	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
18	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
19	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
20	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
21	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
22	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
23	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
24	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
25	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC

Supplementary Table 7 | List of staples in internal linker mix 1 (I1).

No.	I2 sequences (internal mix 2)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA
04	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
05	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
06	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
07	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
08	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
09	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
10	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
11	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
12	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
13	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
14	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
15	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
16	CATTCGCCATGTCATAAATATTCAGCCCAATACT
17	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
18	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
19	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
20	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
21	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
22	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
23	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
24	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
25	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
26	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAATTTTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
27	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
28	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCATTTTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT

915 **Supplementary Table 8 | List of staples in internal linker mix 2 (I2).** Highlighted
916 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I3 sequences (internal mix 3)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA
04	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
05	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
06	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
07	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
08	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
09	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
10	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
11	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
12	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
13	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
14	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
15	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
16	CATTCGCCATGTCATAAATATTCAGCCCAATACT
17	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
18	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
19	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
20	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
21	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
22	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
23	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
24	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
25	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
26	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTTAATTTTTTTTTTTTTTTTTTTTTT**GCTGCTTCCGGCGTA**
27	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT**GCTGCTTCCGGCGTA**
28	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCATTTTTTTTTTTTTTTTTTTTTTT**GCTGCTTCCGGCGTA**

918 **Supplementary Table 9 | List of staples in internal linker mix 3 (I3).** Highlighted
919 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I4 sequences (internal mix 4)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA
04	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
05	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
06	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
07	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAA
08	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
09	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
10	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
11	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
12	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
13	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
14	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
15	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
16	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTT
17	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
18	CATTCGCCATGTCATAAATATTCAGCCCAATACT
19	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
20	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
21	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
22	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
23	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
24	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
25	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
26	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
27	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
28	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC

920
921 **Supplementary Table 10 | List of staples in internal linker mix 4 (I4).**

No.	I5 sequences (internal mix 5)
01	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA
02	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
03	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAA
04	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
05	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
06	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTT
07	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
08	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
09	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
10	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
11	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
12	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
13	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
14	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCGCCATGTTTAC
15	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTCCGGCAAACGC
16	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGCCATTCGCCAT
17	GGGGGATGAATAAATTTTTGTTAAATCAAAA
18	CGTCACCAGTACTTTCCGGCACCGCTT
19	GCAGAAACGAAAGGATTAGGATTATGCCGTC
20	TACATTTAACAAACGGATAACCTCACC
21	CGTAATGGATATTATTTGCACGTAAGGTTTA
22	CGGTGGTGCCATTAGTGATGAAGGGTA
23	**AAAAGGTGAGATGAT**TTTAAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
24	**AAAAGGTGAGATGAT**TTTACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
25	**AAAAGGTGAGATGAT**TTTTGGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
26	**AAAAGGTGAGATGAT**TTTTATAGCTATCTTACAGGAAACAATG
27	**AAAAGGTGAGATGAT**TTTTGTACGCCAGAATCACGATTTTAGA
28	**AAAAGGTGAGATGAT**TTTGTCATAAATATTCAGCCCAATACT
29	**AAAAGGTGAGATGAT**TTTCTGGTGCCTGCTGCAAGGCGATTATGGCGAAA
30	**AAAAGGTGAGATGAT**TTTGGAAACAAAGCGGATCAAACTTAAAAAGAGAC
31	**AAAAGGTGAGATGAT**TTTAAGTTAAAGATAGGTCACGTTGGTGGATTGAC

923 **Supplementary Table 11 | List of staples in internal linker mix 5 (I5).** Highlighted
924 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I6 sequences (internal mix 6)
01	CCATGTTTACATAGCTATCTTACAGGAAACAATG
02	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
03	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
04	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
05	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
06	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
07	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
08	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
09	CATTCGCCATGTCATAAATATTCAGCCCAATACT
10	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
11	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
12	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
13	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
14	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
15	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
16	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
17	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
18	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
19	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
20	GTCTTTACTGATACTAATAGTAGT
21	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA**TAGTAGAGTGGAAAA**
22	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT**TAGTAGAGTGGAAAA**
23	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG**TAGTAGAGTGGAAAA**
24	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA**TAGTAGAGTGGAAAA**
25	ACCCGTCGGATTCTCAGCCAGCT**TAGTAGAGTGGAAAA**
26	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA**TAGTAGAGTGGAAAA**
27	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTTAA**TAGTAGAGTGGAAAA**
28	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTT**TAGTAGAGTGGAAAA**
29	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA**TAGTAGAGTGGAAAA**

926 **Supplementary Table 12 | List of staples in internal linker mix 6 (I6).** Highlighted
927 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I7 sequences (internal mix 7)
01	CCATGTTTACATAGCTATCTTACAGGAAACAATG
02	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
03	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
04	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
05	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAA
06	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
07	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
08	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
09	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
10	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
11	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
12	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTT
13	CATTCGCCATGTCATAAATATTCAGCCCAATACT
14	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
15	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
16	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
17	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
18	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
19	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
20	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
21	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
22	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
23	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
24	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
25	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
26	GTCTTTACTGATACTAATAGTAGT
27	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCATTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
28	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGGTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
29	ACCCGTCGGATTCTCAGCCAGCTTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**

929 **Supplementary Table 13 | List of staples in internal linker mix 7 (I7).** Highlighted
930 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I8 sequences (internal mix 8)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
03	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
04	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
05	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCGCCATGTTTAC
06	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTCCGGCAAACGC
07	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGCCATTCGCCAT
08	GGGGGATGAATAAATTTTTGTTAAATCAAAA
09	CGTCACCAGTACTTTCCGGCACCGCTT
10	GCAGAAACGAAAGGATTAGGATTATGCCGTC
11	TACATTTAACAAACGGATAACCTCACC
12	CGTAATGGATATTATTTGCACGTAAGGTTTA
13	CGGTGGTGCCATTAGTGATGAAGGGTA
14	GTCTTTACTGATACTAATAGTAGT
15	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCATTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
16	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
17	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGGTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
18	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCATTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
19	ACCCGTCGGATTCTCAGCCAGCTTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
20	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGATTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
21	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAATTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
22	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTTTTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
23	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCATTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
24	**AAAAGGTGAGATGAT**TTTAAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
25	**AAAAGGTGAGATGAT**TTTACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
26	**AAAAGGTGAGATGAT**TTTGGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
27	**AAAAGGTGAGATGAT**TTTATAGCTATCTTACAGGAAACAATG
28	**AAAAGGTGAGATGAT**TTTGTACGCCAGAATCACGATTTTAGA
29	**AAAAGGTGAGATGAT**TTTGTCATAAATATTCAGCCCAATACT
30	**AAAAGGTGAGATGAT**TTTCTGGTGCCTGCTGCAAGGCGATTATGGCGAAA
31	**AAAAGGTGAGATGAT**TTTGGAAACAAAGCGGATCAAACTTAAAAAGAGAC
32	**AAAAGGTGAGATGAT**TTTAAGTTAAAGATAGGTCACGTTGGTGGATTGAC

931
932 **Supplementary Table 14 | List of staples in internal linker mix 8 (I8).** Highlighted
933 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I9 sequences (internal mix 9)
01	CCATGTTTACATAGCTATCTTACAGGAAACAATG
02	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
03	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
04	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
05	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTTAA
06	CAGTCCCGGAATTTGTTTTTTGAGAGATAGACTTTACGTACAGCG
07	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
08	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
09	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
10	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
11	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
12	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
13	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTT
14	CATTCGCCATGTCATAAATATTCAGCCCAATACT
15	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
16	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
17	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
18	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
19	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
20	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
21	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
22	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
23	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
24	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
25	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
26	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
27	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
28	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCATTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**

934
935 **Supplementary Table 15 | List of staples in internal linker mix 9 (I9).** Highlighted
936 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I10 sequences (internal mix 10)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCA
04	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
05	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
06	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTT
07	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
08	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
09	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
10	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
11	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGG
12	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCA
13	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
14	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
15	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
16	CATTCGCCATGTCATAAATATTCAGCCCAATACT
17	ACCCGTCGGATTCTCAGCCAGCTGTCTTTACTGATACTAATAGTAGT
18	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGA
19	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
20	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
21	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
22	TCAGGCTGCGCAACTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
23	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
24	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
25	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
26	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTTAATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
27	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
28	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT

938 **Supplementary Table 16 | List of staples in internal linker mix 10 (I10).** Highlighted
939 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I11 sequences (internal mix 11)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
04	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
05	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAA
06	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
07	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
08	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
09	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
10	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
11	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
12	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTT
13	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
14	CATTCGCCATGTCATAAATATTCAGCCCAATACT
15	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
16	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
17	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
18	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
19	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCA
20	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
21	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
22	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
23	GTCTTTACTGATACTAATAGTAGT
24	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCATTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
25	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
26	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGGTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
27	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCATTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
28	ACCCGTCGGATTCTCAGCCAGCTTTTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**
29	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGATTTTTTTTTTTTTTT**AAAAGGTGAGATGAT**

941 **Supplementary Table 17 | List of staples in internal linker mix 11 (I11).** Highlighted
942 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	I12 sequences (internal mix 12)
01	TGATATAAGTATAGCCAACCAATACAAAGAATTAATTAATATTTTGT
02	CCATGTTTACATAGCTATCTTACAGGAAACAATG
03	AAAACGACGGCCAGTGAATCAATAAAATAGCACTAATATC
04	GGCGATTATGGCGAAAGGGGGATGAATAAATTTTTGTTAAATCAAAA
05	CAGTCCCGGAATTTGTTTTTGAGAGATAGACTTTACGTACAGCG
06	AAACTTAAAAAGAGACGCAGAAACGAAAGGATTAGGATTATGCCGTC
07	ACATCGACATAAAAAAGCGGGGTCCAGGAACGACGTGCTT
08	CGGCAAACGCGTACGCCAGAATCACGATTTTAGA
09	TGAATATACAGTAACAGTACCAGGTAGCGTTTGCATAAGAGGCTGAG
10	GGTCCGTTTTTTCGTTTTTTTCGTCGCTGGCAGCATTGCCGTTC
11	CACCTTGCCCTAGGGTTAGAACCACGGAACGCAAACGGC
12	CATTCGCCATGTCATAAATATTCAGCCCAATACT
13	AAAATCAGTTCATCAACGTCTGGCCTTCCTGTAATTGCGT
14	ACGTTGGTGGATTGACCGTAATGGATATTATTTGCACGTAAGGTTTA
15	GGGGACGACGACAGTAAGTTAGCGGCGGAATCTGCAAAAG
16	TCAGGCTGCGCAACTTTTTTTTTGGGAAGGGCGATGGCAAAGCGC
17	CGTCACCAGTACTTTCCGGCACCGCTTCTGGTGCCTGCTGCAA
18	TACATTTAACAAACGGATAACCTCACCGGAAACAAAGCGGATC
19	CGGTGGTGCCATTAGTGATGAAGGGTAAAGTTAAAGATAGGTC
20	GTCTTTACTGATACTAATAGTAGT
21	GCCACCGCCACCCTCAGAGAGCCCAATAATACGAGGAAAGTGTATCATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
22	ATCATACAGGCAAGGGGAACGCCATCAGCTCATTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
23	ACGTCGCGCAGAGGCGAATACAGGAGGCCGACAAATTAATACATCGGTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
24	TCATAGCCCCCTTATCGGATAAGGCGGGGTTTTGCTCATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
25	ACCCGTCGGATTCTCAGCCAGCTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
26	CAATCAAAAAAATCTAAAGCATAGATTTTCAAACAGATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
27	GAGAGGGTCCGTACTCAGGCGAAGCCCTTTTTTTTTAATTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
28	ACGTCAGAGAGAAACAATACTGAGAAGTGTTTTTTTTTTTTTTTTTTTTTTTTTAAAAGGTGAGATGAT
29	ATAATTCGCATTAAATGTGATTGAATCCCTTTTTCTCATTTTTTTTTTTTTTTTAAAAGGTGAGATGAT

944 **Supplementary Table 18 | List of staples in internal linker mix 12 (I12).** Highlighted
945 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	E1 sequences (external mix 1)
01	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTC
02	CAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
03	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG
04	AATCGTAACAAACAAGGGAGAGGGTAGCTATTTTTGAGAAGGCCGGA
05	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA
06	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTAGAGCCAG
07	TTACCAGCGAATTATCACCATTACTATTCACAATTTTTCAAATAAA
08	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG
09	TAGTCTTTATTAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
10	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC
11	TAAGTTTTGATGATACAAAGCCAGAATGGAAAGCGCAGTTTGACAGG
12	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGAATGGCTAT
13	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACC
14	TTATTTACGAAAGCGTAACTGATAATTTAGAAGTTTTTATTAGACT
15	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC
16	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAA
17	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC
18	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTGATTCCCA
19	ATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
20	CACCAGAATATCATTTAATTCGACAACTCGTATTAAATCCTAATAGA
21	GATAAGAGAAGTACGGGATTTAGTACCGTTCTATTTTTCTGATAAA

947 **Supplementary Table 19 | List of staples in external linker mix 1 (E1).**

No.	E2 sequences (external mix 2)
01	CAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
02	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTAGAGCCAG
03	TAGTCTTTATTAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
04	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGAATGGCTAT
05	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTGATTCCCA
06	ATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
07	AATCGTAACAAACAAGGGAGAGG
08	TTACCAGCGAATTATCACCATTAC
09	TAAGTTTTGATGATACAAAGCCA
10	TTATTTACGAAAGCGTAACTGATA
11	CACCAGAATATCATTTAATTCGA
12	GATAAGAGAAGTACGGGATTTAGT
13	**TTTCAGGCTGTAGCTTGG**TTTGTAGCTATTTTTGAGAAGGCCGGA
14	**TTTCAGGCTGTAGCTTGG**TTTTATTCACAATTTTTCAAATAAA
15	**TTTCAGGCTGTAGCTTGG**TTTGAATGGAAAGCGCAGTTTGACAGG
16	**TTTCAGGCTGTAGCTTGG**TTTATTTAGAAGTTTTTATTAGACT
17	**TTTCAGGCTGTAGCTTGG**TTTCAACTCGTATTAAATCCTAATAGA
18	**TTTCAGGCTGTAGCTTGG**TTTACCGTTCTATTTTTCTGATAAA
19	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG**TTTCAGGCTGTAGCTTGG**TTT
20	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA**TTTCAGGCTGTAGCTTGG**TTT
21	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC**TTTCAGGCTGTAGCTTGG**TTT
22	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG**TTTCAGGCTGTAGCTTGG**TTT
23	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC**TTTCAGGCTGTAGCTTGG**TTT
24	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC**TTTCAGGCTGTAGCTTGG**TTT
25	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTCTTT**GATGTGTAGAGT**TGGAGGTGATAG
26	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACCTTT**GATGTGTAGAGT**TGGAGGTGATAG
27	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAATTT**GATGTGTAGAGT**TGGAGGTGATAG

949 **Supplementary Table 20 | List of staples in external linker mix 2 (E2).** Highlighted
950 sequences are complementary to functionalised oligonucleotides (Supplementary Tables 3
951 & 4).

No.	E3 sequences (external mix 3)
01	CAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
02	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTAGAGCCAG
03	TAGTCTTTATTAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
04	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGAATGGCTAT
05	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTGATTCCCA
06	ATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
07	AATCGTAACAAACAAGGGAGAGG
08	TTACCAGCGAATTATCACCATTAC
09	TAAGTTTTGATGATACAAAGCCA
10	TTATTTACGAAAGCGTAACTGATA
11	CACCAGAATATCATTTAATTCGA
12	GATAAGAGAAGTACGGGATTTAGT
13	**TTTGCGAGGCGGACTGGC**TTTGTAGCTATTTTTGAGAAGGCCGGA
14	**TTTGCGAGGCGGACTGGC**TTTTATTCACAATTTTTCAAATAAA
15	**TTTGCGAGGCGGACTGGC**TTTGAATGGAAAGCGCAGTTTGACAGG
16	**TTTGCGAGGCGGACTGGC**TTTATTTAGAAGTTTTTATTAGACT
17	**TTTGCGAGGCGGACTGGC**TTTCAACTCGTATTAAATCCTAATAGA
18	**TTTGCGAGGCGGACTGGC**TTTACCGTTCTATTTTTCTGATAAA
19	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG**TTTGCGAGGCGGACTGGC**TTT
20	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTC**TTTGCGAGGCGGACTGGC**TTT
21	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA**TTTGCGAGGCGGACTGGC**TTT
22	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC**TTTGCGAGGCGGACTGGC**TTT
23	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACC**TTTGCGAGGCGGACTGGC**TTT
24	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG**TTTGCGAGGCGGACTGGC**TTT
25	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC**TTTGCGAGGCGGACTGGC**TTT
26	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAA**TTTGCGAGGCGGACTGGC**TTT
27	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC**TTTGCGAGGCGGACTGGC**TTT

953 **Supplementary Table 21 | List of staples in external linker mix 3 (E3).** Highlighted
954 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	E4 sequences (external mix 4)
01	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTC
02	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG
03	AATCGTAACAAACAAGGGAGAGGGTAGCTATTTTTGAGAAGGCCGGA
04	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA
05	TTACCAGCGAATTATCACCATTACTATTCACAATTTTTCAAATAAA
06	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG
07	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC
08	TAAGTTTTGATGATACAAAGCCAGAATGGAAAGCGCAGTTTGACAGG
09	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACC
10	TTATTTACGAAAGCGTAACTGATAATTTAGAAGTTTTTATTAGACT
11	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC
12	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAA
13	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC
14	CACCAGAATATCATTTAATTCGACAACTCGTATTAAATCCTAATAGA
15	GATAAGAGAAGTACGGGATTTAGTACCGTTCTATTTTTCTGATAAA
16	AGAGCCAGCAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
17	ATGGCTATTAGTCTTTATTAAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
18	GATTCCCAATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
19	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTTTTTTTTTTTTTTTTTTTTTTT**AAAGAGTGAGAGGTGGAG**
20	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGATTTTTTTTTTTTTTTTTTTTTTTT**AAAGAGTGAGAGGTGGAG**
21	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTTTTTTTTTTTTTTTTTTTTTTTTTT**AAAGAGTGAGAGGTGGAG**

956 **Supplementary Table 22 | List of staples in external linker mix 4 (E4).** Highlighted
957 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	E5 sequences (external mix 5)
01	CAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
02	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTAGAGCCAG
03	TAGTCTTTATTAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
04	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGAATGGCTAT
05	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTGATTCCCA
06	ATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
07	AATCGTAACAAACAAGGGAGAGG
08	TTACCAGCGAATTATCACCATTAC
09	TAAGTTTTGATGATACAAAGCCA
10	TTATTTACGAAAGCGTAACTGATA
11	CACCAGAATATCATTTAATTCGA
12	GATAAGAGAAGTACGGGATTTAGT
13	**AAAAGGTGAGATGAT**TTTGTAGCTATTTTTGAGAAGGCCGGA
14	**AAAAGGTGAGATGAT**TTTTATTCACAATTTTTCAAATAAA
15	**AAAAGGTGAGATGAT**TTTGAATGGAAAGCGCAGTTTGACAGG
16	**AAAAGGTGAGATGAT**TTTATTTAGAAGTTTTTATTAGACT
17	**AAAAGGTGAGATGAT**TTTCAACTCGTATTAAATCCTAATAGA
18	**AAAAGGTGAGATGAT**TTTTACCGTTCTATTTTTCTGATAAA
19	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG**TTTCAGGCTGTAGCTTGG**TTT
20	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTC**TTTCAGGCTGTAGCTTGG**TTT
21	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA**TTTCAGGCTGTAGCTTGG**TTT
22	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC**TTTCAGGCTGTAGCTTGG**TTT
23	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACC**TTTCAGGCTGTAGCTTGG**TTT
24	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG**TTTCAGGCTGTAGCTTGG**TTT
25	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC**TTTCAGGCTGTAGCTTGG**TTT
26	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAA**TTTCAGGCTGTAGCTTGG**TTT
27	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC**TTTCAGGCTGTAGCTTGG**TTT

959 **Supplementary Table 23 | List of staples in external linker mix 5 (E5).** Highlighted
960 sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

No.	E6 sequences (external mix 6)
01	CTTATTACGTGGCAACATATAAAACATATGGTCAGAGAGACGAGCGTC
02	CAAAATCAAACGTCACAGAAAATTGAAACGCAGAGAATTAAACCCACA
03	AAACATTCAAAAACATGAACGGTCGAACTGACAGACCAG
04	TTTAATGGAGCCAGTAATAAGAGAGTATTCTAAACAGCCATA
05	CAATGAAACCATCAAGGGTGAGAAGATCTACAAAGGCATTAGAGCCAG
06	ATACCGGGCACATTAATTGCGTTGTCCACTATGGAAAGCCGG
07	TAGTCTTTATTAAAAAAATCGTCTTTGCAACACGCCGCTAAGAATCAG
08	CCCTCAGCCCCTGCCACTGGTAACAATAATCATTAAACC
09	TACCGAACGAACCCGCCGCCAGCACTCTGAATTTACCTGAATGGCTAT
10	AGTAGAAGATATTACCGCCAGCCAGAAATGGAGTCACGCTTCGGAACC
11	CGCATAACAAGAGGCAAAAGAATATGCCCTGAACATTATTAC
12	AAAAGATTTTCGAGCTTCAAAGCGCTCCTTTTAAGGAATTGGAAGAAA
13	GTTGAAATCAATATAAAAGAAACCCTGAGAGCAGCAGGC
14	ATGGTCAATAACCAGCACTAACAACTTTGCCCGAACGGTTGATTCCCA
15	ATTCTGCGTTTCGCAATTTAATTGAACCAGACCCTCGTTTCCAGAGGG
16	AATCGTAACAAACAAGGGAGAGG
17	TTACCAGCGAATTATCACCATTAC
18	TAAGTTTTGATGATACAAAGCCA
19	TTATTTACGAAAGCGTAACTGATA
20	CACCAGAATATCATTTAATTCGA
21	GATAAGAGAAGTACGGGATTTAGT
22	**TTTCAGGCTGTAGCTTGG**TTTGTAGCTATTTTTGAGAAGGCCGGA
23	**TTTCAGGCTGTAGCTTGG**TTTTATTCACAATTTTTCAAATAAA
24	**TTTCAGGCTGTAGCTTGG**TTTGAATGGAAAGCGCAGTTTGACAGG
25	**TTTCAGGCTGTAGCTTGG**TTTATTTAGAAGTTTTTATTAGACT
26	**TTTCAGGCTGTAGCTTGG**TTTCAACTCGTATTAAATCCTAATAGA
27	**TTTCAGGCTGTAGCTTGG**TTTACCGTTCTATTTTTCTGATAAA

Supplementary Table 24 | List of staples in external linker mix 6 (E6). Highlighted
sequences are complementary to functionalised oligonucleotides (Supplementary Table 3).

Osmolality difference	Glycine (mM)	MgCl$_2$ (mM)	NaCl (mM)	Total volume (µl)
-78%	0	25	26.3	30
-58%	100	30	26.3	30
-42%	200	35	26.3	30
-22%	300	45	26.3	30
-4	400	50	26.3	30
+12%	500	55	26.3	30
+33%	600	65	26.3	30
+54%	700	75	26.3	30

966 **Supplementary Table 25 | Tonicity screen sample compositions.** Composition of the
967 samples shown in Fig. 2c. The magnesium optimum for triangle assembly increased with
968 increasing glycine concentrations. Concentrations above or below the optima presented in
969 this table result in incomplete or impaired assembly.

	TRIS (mM)	EDTA (mM)	MgCl$_2$ (mM)	NaCl (mM)	CsCl (mM)	Sucrose (mM)	Glucose (mM)	Glycine (mM)	pH
Folding Buffer	5	1	20	5					8.0
Sodium Buffer	5	1	5	305					8.0
Caesium Buffer	5	1	5	5	320				8.0
Sucrose Buffer	10	1				470			8.0
Imaging Buffer A	5	1	5	5		280			8.0
Imaging Buffer B	5	1	5	5			280		8.0
Isotonic assembly buffer	5	1	210	5					8.0
Glycine buffer	5	1	5					1000	6.8

972 **Supplementary Table 26 | Buffer compositions**

973