Programmable DNA shell scaffolds for directional membrane budding

Michael T. Pinner[1,2] and Hendrik Dietz[1,2]

[1] Laboratory for Biomolecular Nanotechnology, Department of Biosciences, School of Natural Sciences, Technical University of Munich, Am Coulombwall 4a, 85748 Garching, Germany.
[2] Munich Institute of Biomedical Engineering, Technical University of Munich, Boltzmannstraße 11, 85748 Garching, Germany.

Abstract

In the pursuit of understanding and replicating biological processes at the nanoscale, controlling cellular membrane dynamics has emerged as a key area of interest. Here, we report a system capable of controlling directional membrane budding through a mechanism mimicking virus assembly on a lipid membrane. We employed three-dimensional DNA origami techniques to construct icosahedral shells that self-assemble and interact with the lipid membranes of giant vesicles via cholesterol moieties. The membrane-constrained assembly of shells results in the programmable formation of DNA-shell-coated vesicles or vesicle-coated DNA shells, effectively demonstrating membrane curvature induction and spontaneous neck scission. We further show that these processes can be combined into a 2-step budding reaction, giving rise to nested bivesicular objects with DNA shells encapsulated between two lipid vesicles. Our findings replicate key aspects of natural endocytic and exocytic pathways, and open new avenues for the exploration of membrane mechanics and for applications in targeted drug delivery and synthetic biology.

Introduction

Membrane budding is a ubiquitous biological phenomenon in which parent lipid membranes release daughter vesicles for the transport of materials across the membrane. It plays a pivotal role in the lifecycle of many viruses, particularly enveloped viruses, and is intimately tied to virus infection and virus release processes from host cells[1]. Understanding and

potentially controlling budding processes may contribute to comprehending viral
pathogenesis and developing antiviral strategies. Further, in clathrin-mediated endocytosis
(CME), clathrin, with its distinctive triskelion shape, remodels lipid membranes by binding to
membrane-bound receptor-adaptor protein complexes, followed by self-assembly into
spherical cage-like structures that induce curvature in the cell membrane[2–4]. This process
culminates in the formation of a clathrin-coated vesicle[2]. Artificial membrane budding
systems are of particular interest as they allow studying the underlying mechanics in a
controlled environment, potentially offering a deeper understanding of cellular processes.
Other benefits may include the development of improved drug delivery systems or, in the
context of immunology, vaccine vectors by producing non-infectious but immunogenic
biopsies of pathogenic cells. They may also support the creation of synthetic cells or
organelles, contributing to the field of synthetic biology where there is growing interest in
building biological systems with novel functions.

Scaffolded DNA origami is a unique method to self-assemble defined 3D structures on the
nanoscale using DNA as a building material[5–7]. A several kilobases long, single-stranded
'scaffold' strand is mixed with an array of shorter oligonucleotides and exposed to a thermal
annealing ramp to drive the folding of the DNA origami structure. The programmability of
Watson-Crick base pairing interactions has allowed the production of nanostructures ranging
from simple bricks[6] to giga-Dalton-sized multimeric assemblies[8–11]. The relative ease of
producing DNA origami structures expanded the experimental toolkit on the nanoscale and
accelerated research towards synthetic biology applications, including efforts to interact with
and manipulate natural membranes.

Previous efforts in the field related to lipid membranes have successfully tethered membrane-
spanning DNA nanopores of various complexity to lipid bilayers using hydrophobic moieties
such as cholesterol[12], ethyl-phosphorothioate[13] or tetraphenylporphyrin[14]. Notably,
cholesterol-labelled DNA origami objects tethered to lipid membranes retain diffusive mobility
influenced by factors like linker oligo length[15], paving the way for the 2D assembly of
monomeric structures into multimeric lattices[16]. Prior studies have also shown that both
curved[17–19] and planar[17] origami structures can induce tubulation in giant unilamellar lipid
vesicles (GUVs). On-membrane polymerisation of origami structures into 2D lattices likewise
resulted in the deformation of the underlying membrane reminiscent of clathrin networks[16,20–
22].

In the biological systems that inspire our work here, membrane curvature changes are often
induced by curved proteins, including Clathrin and Bin/amphiphysin/Rvs (BAR) domain
proteins. These proteins remodel lipid bilayers through mechanisms like the insertion of
hydrophobic residues into one of the leaflets of the bilayer, causing asymmetrical surface
area expansion, imprinting the protein's curved shape onto the bilayer through strong binding,
and crowding effects[23,24]. Further, BAR domain proteins also act as curvature sensors and
recruit the GTPase dynamin to the bud neck, promoting scission[2,25]. Interestingly, the
amphipathic helix used by N-BAR proteins for membrane anchoring alone can constrict and
sever the bud neck[26]. This mechanism is echoed in biological entities like the influenza virus
M2 protein, which facilitates membrane scission through an amphipathic helix independent
of its ion channel function or the cellular budding machinery[27].
78
Building upon this foundational work, we have developed an artificial, DNA origami-based
membrane budding system with spontaneous neck scission. At its core are three-
dimensional membrane-interacting triangular subunits constructed from DNA with bevelled
edges[28]. Strategically placed shape-complementary protrusions and recesses at the
triangles-edges direct the self-assembly of these triangular subunits into icosahedral shells
through adhesive base pair stacking interactions[29]. Interactions with lipid membranes of giant
vesicles (GVs) were promoted using cholesterol moieties placed on specific triangular
surfaces. We hypothesised that membrane deformation processes may be sufficient for the
spontaneous scission of the bud neck without the need for active constriction by GTPases
like dynamin[25]. Indeed, the self-assembly of membrane-bound triangles induced local
curvature and finally led to the formation and eventual scission of buds. The positioning of
cholesterol determined whether the bud-formation occurs inward or outward from the source
vesicle, leading to DNA-shell-coated vesicles (DCV) and vesicle-coated DNA shells (VCD),
respectively.

Results

1. DNA-Shell-Coated Vesicles

We adapted a previously developed programmable icosahedral shell canvas, in which twenty
equilateral DNA origami triangles form a closed icosahedron triggered by elevated
concentrations of magnesium chloride for membrane-supported assembly[28]. To provide the
necessary hydrophobic interactions required for association with lipid membranes, we
incorporated multiple cholesterol-bearing oligonucleotides at the shell-inward-facing surface
of the constituent triangular DNA origami subunits (Fig. 1a & Supplementary Fig. 1). We

hypothesised that the 2D diffusion of triangles on a fluid membrane can support the assembly

of icosahedral shells whilst pulling membrane material along, resulting in increasingly

deformed membrane buds growing away from the parent membrane until the bud neck is cut

and a DNA-shell-coated vesicle (DCV) is released into solution.

We screened cholesterol density and positioning to balance the solubility of the triangles with

membrane affinity (Supplementary Fig. 2a) and then validated the assembly of cholesterol-

modified triangles into the expected icosahedral shells in the absence of membranes.

Introducing cholesterol moieties resulted in a shift from the previously observed relatively

uniform icosahedral shells to a mixture containing both octahedral and icosahedral shells.

This variation was controllable: by limiting the reach of each cholesterol moiety, as shown in

Supplementary Figure 2b, we could largely revert the assembly to predominantly icosahedral

structures. The addition of lipid membranes into the system introduced another variable,

again leading to the occurrence of octahedral structures. This tendency varied depending on

the amount of cholesterol per triangle, detailed in Supplementary Figures 3 and 4. We believe

this effect arises from a complex interplay of forces, including the attraction between

cholesterols and membranes, membrane deformation, and elastic deformation within the

shells caused by less-than-ideal angles between the triangle subunits. Nevertheless,

octahedral and icosahedral forms were both considered suitable for our objectives.

We chose giant vesicles (GVs) composed of 1,2-dioleoyl-sn-glycero-3-phosphocholine

(DOPC) with up to 1% of fluorescently labelled 1,2-dioleoyl-sn-glycero-3-

phosphoethanolamine (DOPE) as model membranes to study membrane deformation and

budding. Vesicles were grown by gentle swelling of dried lipid cakes on polyvinyl alcohol

(PVA) coated surfaces[30] in folding buffer containing 5 mM $MgCl_2$ and 300 mM NaCl or 320

mM CsCl. Triangles bearing linker handles were hybridised to chol-bearing antihandles at a

1.5-fold excess for at least 30 min at room temperature. Thereafter, aliquots of GVs and chol-

decorated triangles were mixed to anchor the nanostructures to the vesicles. GVs could be

readily coated with the thus-modified DNA triangle subunits (Fig. 1b). Non-specific binding

of triangle subunits to vesicles was suppressed by including NaCl in the reaction mixture,

ensuring triangles bind vesicles in the designed orientation as defined by the location of their

cholesterol moieties[19].

We triggered DNA shell assembly on membranes by increasing the $MgCl_2$ concentration and

incubating the solution at 37 °C for up to multiple days. Imaging the samples by negative

stain transmission electron microscopy (TEM) revealed DCVs appearing as dark DNA shells
that fully engulfed lipid vesicles appearing as bright rings in the images (Fig. 1c, top right).
Occasionally, the shell-enclosed vesicles appeared deformed or cracked, which we attribute
to staining and drying artefacts. To study DCVs in their native conditions, we also acquired
micrographs of vitrified samples by cryogenic transmission electron microscopy (cryoEM, Fig.
1c, bottom right). The data generally agrees with the conclusions drawn from negative stain
TEM data, but the enclosed vesicles seen by cryoEM rarely did have any defects except for
occasionally encapsulating even smaller vesicles themselves. In the cryo-EM micrographs,
the vesicle typically fills up the available space inside the origami shell, with the membrane
in direct contact with the enclosed DNA shell. These findings support budding triggered by
shell assembly on a parent membrane as the predominant mechanism of DCV formation as
opposed to capturing small unilamellar vesicles (SUVs). We also note that variants of the
triangle subunits that cannot form shells also did not produce DCVs, supporting the idea that
shell assembly on membranes is an essential driver for DCV formation (Supplementary Fig.
5).
151
As our setup involves budding off nanoscopic vesicles that are multiple orders of magnitude
smaller than their parent vesicles, direct observation of DCV formation from GVs is
challenging. However, when using large vesicles (LVs), direct observation of intermediate
steps in DCV formation is possible using cryoEM. To this end, we thoroughly washed GVs to
remove contaminating SUVs and then pushed them through a 200 nm pore filter to produce
LVs of roughly that size. Furthermore, we shortened the on-membrane incubation of origami
triangles to capture growing buds before they are released into the solution as DCVs. We
mixed vesicles with isosmotic $MgCl_2$ solutions to avoid artefacts by vesicle deflation. We thus
succeeded in acquiring micrographs that show various intermediate states of deformation of
the parent vesicle induced by the emerging DNA shells (Fig. 2a). Whilst membrane patches
free of DNA triangle subunits appear undisturbed, those with triangle objects on them had
deformations that matched the curvature of the triangle assemblies. Notably, some vesicles
have multiple buds with a clearly visible bud neck. This image data further supports the notion
that budding is the mechanism behind DCV formation from LVs and GVs. Occasionally, we
also found DCVs where the DNA origami shell had assembly "scars" consisting of one or two
missing triangles (Fig. 2b). We believe these scars to be another signature of budding,
considering that the increasingly high curvature at the bud neck can become a steric barrier
to completion of the DNA origami shell. In this scenario, spontaneous membrane scission

competes with the completion of the DNA origami shell, which may occasionally lead to such
assembly scars.

To evaluate whether the lipid vesicles were fully enclosed by well-ordered icosahedral or
octahedral DNA shells, we recorded single-axis tilt series to generate tomograms, allowing
us to study DCVs in 3D (Fig. 2c). The representative tomograms confirm full engulfment of
the vesicle by closed shells featuring the symmetry properties expected by design.

We next studied the influence of osmolality on DCV formation. The addition of concentrated
magnesium chloride solution as a trigger for icosahedral shell assembly inevitably increases
the osmolality of the sample buffer and may thus alter the biophysical properties of the
membrane by deflation of GVs. Indeed, hyperosmotic stress has previously been exploited
to alter vesicle shape by membrane-bound DNA origami[19]. We tested the influence of
osmolality in DCV formation by triggering DNA shell assembly using either hyperosmotic or
isosmotic $MgCl_2$ solutions (Fig. 2d). We measured the extent of DCV formation by agarose
gel electrophoresis using fluorescent labels on DNA origami components and a fraction of
the lipids constituting the GVs (Supplementary Fig. 6). Fluorescent lipid bands colocalised
only with the higher-order shell assemblies and were virtually absent at the level of
monomeric triangles. Direct comparison of DCVs formed under hyperosmotic and isosmotic
conditions reveals that osmotic pressure differences do not noticeably influence assembly
yields as measured by comparing the brightness of lipid bands relative to the DNA shell band
in the gels. Hyperosmotic deflation of GVs, therefore, is not required for DCVs to form.

In our study, we primarily used vesicles made of 99.0 – 99.5% DOPC, supplemented with a
small fraction of fluorescently labelled DOPE species. Our initial theory suggested that
introducing cholesterol into the outer layer of the vesicle membrane via DNA triangle tethering
might be crucial for DCV formation. To evaluate this, we introduced 3% biotinylated DOPE
into the lipid mixture and prepared GVs from it. Instead of attaching cholesterol-modified
oligonucleotides to the origami triangles, we used biotinylated oligonucleotides connected to
NeutrAvidin (b-NAv). NeutrAvidin's unoccupied binding sites could then capture the vesicles'
biotinylated lipid components, allowing the DNA triangles to attach to the membrane without
directly inserting cholesterol or other foreign anchors into the membrane leaflets. Interestingly,
we observed the formation of OCVs under both hyperosmotic and isosmotic conditions using
this approach, as shown in Figure 2e. Furthermore, excess cholesterol-modified
oligonucleotides with a different sequence did not influence this result. This indicates that the

integration of cholesterol into the membrane is not necessary for the formation of DCVs and
supports the notion that DCVs are driven by curvature that emerges from the assembly of
the DNA origami shells.
208
We also investigated the efficiency of DCV assembly as a function of the average parent
vesicle size. To this end, we passed GVs through filters with 400, 200, or 50 nm pores,
producing vesicles of specific sizes. We then used these vesicles in a controlled lipid-to-DNA
triangles stoichiometry budding assay to examine the efficiency of DCV formation, which we
measured by quantifying DNA and lipid band intensities of DCVs ran on an agarose gel. As
Figure 2f shows, there is a defined optimal lipid quantity for each vesicle size except for GVs.
Interestingly, whereas DCV yield curves for SUVs and LVs were quite similar, the curve for
GVs peaked and then plateaued over a broad lipid quantity range (see Figure 2g). To test
whether triangles also assemble in solution into partial shells (i.e. half icosahedra) and then
subsequently form a bud on GVs, we inverted the DCV preparation protocol, i.e. we first
preassembled shells in the absence of membranes and then added GV with a temporal offset
of several hours, which led to a substantial decrease in DCV yield (Supplementary Fig. 7).
This data further supports the notion that on-membrane assembly of shells is a key driver for
bud formation.
223
The DCVs prepared thus far formed in a process reminiscent of exocytosis, where vesicles
bud outwardly from a cell, but with the difference that the components forming the bud are
approaching from the exterior of the parent vesicle. We were also able to invert the
directionality toward mimicking endocytic processes. To this end, we placed the cholesterol
moieties that promote membrane association on the shell-outward face of the triangle
subunits. Shell assembly on the membrane should then lead to the formation and release of
a vesicle-coated DNA shell (VCD) into the lumen of the parent vesicle (Fig. 3a), effectively
creating an organelle- or endosome-like compartment. We extruded GVs through 200 nm-
sized pore filters and prepared triangles carrying one cholesterol moiety per side. Direct
imaging with cryoEM (Fig. 3b) revealed VCDs, i.e. lipid vesicles tightly wrapped around
icosahedral shells located within their parent vesicles. Occasionally, we also observed free
VCDs presumably set free by the bursting of the parent vesicle. As with DCVs, we
occasionally spotted scars on the VCDs where the DNA shell was visibly missing triangles.
237
238   Finally, we tested if bivesicular shell structures could be produced by combining both
239   outward and inward budding in a two-step process using triangle subunits with sequence-

orthogonal linker handles on their top and bottom faces (Fig. 3c). First, we hybridised cholesterol-modified DNA strands to the linker handles on the bottom face of the triangles and mixed the DNA triangles with GVs to produce DCVs as described earlier. We then centrifuged the sample at low speed to remove some of the remaining GVs, followed by ultracentrifugation of the supernatant to pellet any remaining vesicles, DNA components and DCVs. Most of the supernatant was discarded, and the pellet was resuspended in the remaining volume. We then prepared LVs by extruding GVs through 200 nm pore size filters and added cholesterol-modified oligonucleotides to this vesicle mixture to cover the membrane with single-stranded linker handles. The sequence of these strands is complementary to the single-stranded linker handles on the shell-outer face of the triangles but orthogonal to those on the shell-inner face used in the first step for outward budding. Finally, the DCV concentrate was mixed with the LVs covered with complementary linker handles, and the mixture was incubated at 37 °C. We analysed the sample using cryoEM and indeed found vesicle-coated DCVs (VCDCVs). Like VCDs, VCDCVs were typically inside of their respective parent vesicle. These particles may be considered examples of synthetically created organelles featuring a cytoplasm-like compartment (the inner vesicle) and a periplasm-like compartment (the void between the outer envelope membrane and the inner lipid membrane). The DNA icosahedral shell acts as a stabilising mechanical skeleton, offering engineering options for including additional molecular functionalities.

Discussion

A key difference between our system and cellular clathrin-mediated endocytosis is the absence of active neck scission. In cells, dynamin is needed for vesicle release[31], but in our system, DNA-shell-coated vesicles (DCVs) release spontaneously, albeit after a comparably long incubation time. Neck scission may currently represent the kinetic bottleneck in DCV formation. Different membrane anchors, such as the influenza virus protein M2, could improve the system's kinetics. Previously, cholesterol-labelled DNA objects of varying shapes have been described to cause vesicle tubulation, a phenomenon also observed with Clathrin's adaptor protein epsin[17–19,32]. We find that the DNA origami's membrane anchor type does not significantly affect bud formation. The assembly of DNA triangles into shells is the primary driver of bud formation, as these shells imprint their curvature onto the membrane, shaping the buds. We also found that DCV formation varies with vesicle size and lipid quantity, where high lipid quantities contained in GV reservoirs yielded good amounts of DCVs, whereas budding from smaller vesicles had reduced yields.

We could trigger inward budding by shifting the position of cholesterols from the inner face to the outer face of a triangle. When using DCVs instead of monomeric triangles, we could produce nested bivesicular structures carrying both an inner and an outer vesicle, demonstrating the system's modularity and level of control. However, further optimisation is needed to improve yields and reduce aggregation for efficient production of bivesicular DNA shells.

In conclusion, our DNA origami-based system offers an attractive approach to engineer vesicles of controlled size featuring addressable endo- or exoskeletons. By performing the budding on GVs containing molecules of interest, the budding system could be further developed toward molecular payload applications. Using nested bivesicular structures, it becomes possible to design scenarios where two different molecular payloads are delivered simultaneously but separately. This work may thus pave the way for innovative applications in targeted drug delivery, vaccine development, and synthetic biology. Future explorations could extend to fine-tuning the system's efficiency and exploring the breadth of its applications, particularly in medical and biotechnological fields. The potential to manipulate membrane dynamics at the nanoscale also opens new horizons in synthesising artificial cells and organelles, offering a valuable tool for researchers in their quest to harness the power of nanotechnology for practical and therapeutic purposes.

Acknowledgements

The authors acknowledge financial support received from the European Research Council Advanced Grant awarded to H.D. (grant agreement 101018465) and the Deutsche Forschungsgemeinschaft for the Gottfried Wilhelm Leibniz Program grants provided to H.D. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 765703. We thank Andreas Bausch for technical assistance, Björn Högberg for discussions and Volodymyr Mykhailiuk for help with illustrations.

Author contributions

The research was performed by M.P., supervised by H.D. The manuscript was written by M.P. and H.D.

Figures & Captions

Figure 1: DNA-shell-coated vesicle formation

a, Schematic of a DNA origami triangle-based membrane budding system. The triangular

structures are decorated with cholesterol moieties to tether them to the membrane of a lipid

vesicle. Upon increasing the $MgCl_2$ concentration, the triangles self-assemble into DNA shells,

pulling the membrane underneath along to form a bud. Eventually, the bud neck is cut, and

a DNA-shell-coated vesicle (DCV) is released. **b,** Attachment of DNA origami triangles (yellow)

onto giant vesicles (GVs, cyan). Triangles will adsorb onto GVs even if they are not decorated

with cholesterol (top left & right panel). However, adding sodium chloride suppresses this

non-specific adsorption (bottom left panel), and membrane binding will only occur if

cholesterol is attached to the triangle (bottom right panel). Scale bars: 50 µm. **c,** Transmission

electron micrographs of empty DNA shells and DCVs. In negative-stain TEM (top panels),

lipid vesicles within DCVs appear as white rings inside the DNA shell. In cryogenic electron

microscopy (cryoEM, bottom panels), the vesicle is visible as a dark ring inside the DNA shell.

Scale bars: 100 nm.

Figure 2: DNA-shell-coated vesicles form by membrane budding.

a, CryoEM micrographs of large vesicle (LV) deformation upon assembly of membrane-bound DNA origami triangles. Bud curvature closely follows the intrinsic curvature of the assembling. Scale bars: 100 nm. **b,** DCVs occasionally appear incomplete with visible 'scars' of missing triangles. The bud neck prevents closure of the DNA shell by steric hindrance, resulting in a gap in the origami shell that remains visible after scission of the neck. Left panel: negative-stain TEM; Right panel: cryoEM. Scale bars: 100 nm. **c,** Tomogram of an icosahedral DCV depicted as summed up z-slices of the top, central and bottom segments of the DCV. The topmost segment appears as a tilted, incomplete pentamer. The central segment reveals a lipid vesicle trapped within the origami shell. The bottommost segment is a pentameric shell cap. **d,** Agarose gel of DCVs prepared under hyperosmotic and isosmotic conditions. Lipid material (cyan, DOPE-Atto488) remains in the pockets unless cholesterol-hybridised triangles (yellow, Atto643) are mixed with GVs. Lipids form bands colocalising with octahedral and

icosahedral shells but not with monomeric triangles. Thus, lipid material is dragged into the

gel as DCVs. Gel pockets were oversaturated for better band visualisation. **e,** Agarose gel of

DCVs prepared by using biotin-NeutrAvidin interactions to connect origami triangles (yellow,

Atto643) and GVs (cyan, DOPE-Atto488) containing a fraction of biotinylated DOPE. **f,**

Relative lipid band intensities of DCVs prepared from vesicles of different sizes and quantities.

g, Relative lipid and DNA band intensities of DCVs obtained from different quantities of GVs.

Figure 3: Vesicle-coated DNA shells and multivesicular DNA shells

a, Illustration of vesicle-coated DNA shell (VCD) formation by budding into a parent vesicle. The budding direction is reversed with respect to DCVs by shifting the cholesterol linker handles from the shell-inner face to the shell-outer face of the triangle subunits. **b,** cryoEM micrographs of VCDs within their parent vesicle. Scale bars: 100 nm. **c,** Illustration of a bivesicular DNA shell (vesicle-coated DCV; VCDCV) contained in a parent vesicle, produced in a 2-step budding assay using triangles with orthogonal linker handles on their top and bottom faces. In the first step, the bottom face linkers are hybridised to cholesterol-modified DNA strands for DCV formation by outward budding. Then, DCVs are mixed with LVs carrying single-stranded linker handles complementary to those on the shell-outer face of the triangles, resulting in inward budding and VCDCV formation. As a result, VCDCVs have four layers: An inner vesicle stemming from the first step (orange), the DNA origami shell (white), an outer vesicle (pink) coupled to the shell by complementary linker handles, and the parent vesicle (purple) with its DNA linker handles (green blobs). **d,** cryoEM micrographs of VCDCVs. Images were Gaussian blurred and levelled for clarity. Scale bars: 100 nm.

Materials and Methods

Materials

All chemicals were purchased from Sigma Aldrich and Carl Roth unless stated otherwise. 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC) and 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(cap biotinyl) (biotinylated DOPE) were obtained from Avanti Polar Lipids, and Atto488 and Atto643 modified 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) was obtained from Atto-Tec. NeutrAvidin was purchased from Thermo Scientific. Regular DNA oligonucleotides were purchased from IDT, and modified oligonucleotides were obtained from Biomers. Scaffold DNA was produced biotechnologically in-house.

DNA origami folding, purification & quantification

DNA origami triangles were folded as described previously[28]. Briefly, circular scaffold strands of bacteriophage origin (p8064, 50 nM) and staple oligonucleotides (200 nM per staple) were mixed in folding buffer (5 mM Tris base, 1 mM EDTA, 5 mM NaCl, pH 8) containing 20 mM $MgCl_2$ and exposed to a thermal annealing ramp (65 °C, 15'; 58-53 °C, 1 h/°C; stored at 20 °C) in a Tetrad2 (Bio-Rad) thermocycler. For some experiments, structures were fluorescently labelled by including Atto643 labelled oligonucleotides in the folding reaction. The folded product was purified by agarose gel electrophoresis by excising the leading band, crushing the gel pieces and centrifuging them in 0.45 µm Costar Spin-X spin columns (Corning) for 10 min at 8 krcf. When needed, the purified structures were washed and concentrated by ultrafiltration (Amicon Ultra 0.5 mL, 100 kDa cut-off, Millipore), exchanging the buffer for folding buffer containing 5 mM $MgCl_2$ and 300 mM NaCl. Origami solutions were quantified using a NanoDrop8000 spectrophotometer (Thermo Scientific).

Vesicle preparation and quantification

Giant vesicles were prepared by gentle swelling of dry lipid films on polyvinyl alcohol gels as described previously[30], with slight modifications. A petri dish (∅ 10 cm) was plasma cleaned in a glow discharge device, and then 1800 µl of a 5% solution of polyvinyl alcohol (Mowiol 28-99, Sigma Aldrich) in ddH$_2$O was evenly distributed across its surface. The dish was incubated for at least 30 min at 50 °C until a dry gel formed. Then, 162 µl of the lipid mixture (DOPC with additional 0.5-1% fluorescently labelled DOPE (DOPE-Atto488 or DOPE-Atto643, supplied by Atto-Tec) if visualisation by fluorescence was desired; 2.54 mM total lipid concentration) in chloroform was evenly spread out across the surface using a Drigalski spatula until the chloroform evaporated. Remaining traces of solvent were removed by

exposing the dried lipid cake to a vacuum for at least 15 min. Next, 4-5 mL of folding buffer

(+5 mM $MgCl_2$ and 300 mM NaCl) was added to the petri dish and the solution was left in a

humid chamber in the dark at RT for at least 1 h. GVs were detached by gently tapping against

the dish and pipetting up and down with a cut pipette tip. The GV suspension was stored at

4 °C in the dark.

When GVs were to be washed and concentrated, the dried lipid film was swollen in folding

buffer (+5 mM $MgCl_2$ and 320 mM CsCl) and, after harvesting, transferred into a falcon tube.

The tube was filled up with isosmotic buffer (folding buffer containing 5 mM $MgCl_2$ and 300

mM NaCl) and centrifuged for at least 30 min at 300 rcf. The supernatant was discarded, and

the GV pellet was washed 2-3 more times by adding 900 µl buffer and centrifuging the

suspension for 7 min at 300 rcf.

Smaller-sized vesicles were prepared using a mini-extruder (Avanti polar lipids) and

polycarbonate membranes with the desired pore size (400, 200, 50 nm; Whatman). GV

preparations were pushed back and forth through the membrane 21 times.

Lipids in vesicle preparations were quantified using a colourimetric phospholipid

quantification assay kit (Sigma Aldrich) following the manufacturer's instructions. The

absorbance was measured at 570 nm on a Clariostar plus microplate reader (BMG Labtech).

The result was extrapolated based on the molar composition of the lipid mix used to account

for PE species, which are not detectable by this method.

Buffer osmolalites were measured using an Osmomat 3000 freeze point osmometer

(Gonotec).

Vesicle binding study

Microscope slides and coverslips were washed with ddH_2O and incubated in denatured BSA

blocking buffer (10 mM TRIS, 150 mM NaCl, 50 µM BSA, pH 8) overnight as described

previously[33]. Next, the glassware was thoroughly washed with ddH_2O to remove residual

blocking buffer and dried at 50 °C for 1 h. 6 µl of GVs (99.95% DOPC, 0.05% DOPE-Atto488;

swollen in folding buffer (+5 mM $MgCl_2$, 280 mM sucrose) was mixed with origami triangles

(+/- cholesterol modified strands; fluorescently labelled with Atto643 modified

oligonucleotides; final origami concentration: 7 nM) and topped up to 40 µl with folding buffer

(+5 mM $MgCl_2$, 280 mM glucose). 300 mM NaCl was added to samples testing the effect of

elevated sodium on the binding behaviour of origami. The samples were pipetted into an

imaging chamber formed by sandwiching a silicon isolator (Grace Bio-Labs) between

passivated coverslips and microscope slides. The samples were imaged using a Leica TCS

SP5 confocal microscope with a Leica HCX PL APO CS 63x/1.40-0.60 oil immersion

objective.

Budding assays

Vesicles were prepared as described above using 99.0 – 100% DOPC with the remainder

being fluorescently labelled DOPE if visualisation by fluorescence was desired. Origami

triangles were hybridised to cholesterol-modified oligonucleotides by adding them at a 1.5 x

excess with respect to the total amount of compatible linker handles on the origami. The

hybridisation reaction was incubated at RT for at least 30 min. For outward budding reactions,

24 µl of untreated GV suspension was mixed with origami triangles with linker handles on the

shell-inward face at a final concentration of approx. 2 nM origami. In experiments where lipid

quantities in GV preparations were known, 1000 pmol lipid of the GV suspension was mixed

with the origami (6 µl at 10 nM). The sample was topped up to 30 µl with folding buffer (+5

mM $MgCl_2$, 300 mM NaCl), and the $MgCl_2$ concentration was adjusted to 65 mM by adding

1 M $MgCl_2$ solution (hyperosmotic conditions) or to 60 mM by adding folding buffer (+210

mM $MgCl_2$; isosmotic conditions). Budding reactions were incubated at 37 °C for 1-3 d.

For inward budding reactions, samples were prepared in the same way with the difference

that origami triangles (3 µl at 20 nM) with linker handles on the shell-outward face were mixed

with LVs (1500 pmol lipids; approx. 200 nm diameter).

For bivesicular DNA shells (VCDCVs), triangles with orthogonal linker handles on the top and

bottom face were used, and inward and outward budding reactions were conducted

successively. First, DCVs were produced in an outward budding assay (1200 µl in total) using

cholesterol-modified oligonucleotides hybridised to the bottom face linker handles. The

sample was then centrifuged at 300 rcf for 5-8 min, and the pellet containing residual GVs

was discarded. The supernatant was then ultracentrifuged at 55 krpm (avg. 108 krcf) for 30

min in a Beckman Coulter Optima MAX-TL ultracentrifuge equipped with a TLA-110 fixed

angle rotor. 80-90% of the supernatant was removed, and the pellet was resuspended in the

remaining volume.

Meanwhile, LVs (200 nm diameter) were mixed with a second cholesterol-bearing

oligonucleotide complementary to the linker handles on the top face of the triangles. These

oligonucleotides were added at a 50x excess with respect to the total amount of
complementary linker handles on the origami, and the solution was incubated at RT for 15-
30 min. Then, the magnesium concentration in the LV solution was adjusted to 60 mM $MgCl_2$
by adding an isosmotic folding buffer containing 210 mM $MgCl_2$. Finally, LVs (1500 pmol
lipids) were mixed with the DCV concentrate (to a final concentration of 1-5 nM DNA), and
the samples were incubated at 37 °C for 1-3 d.

Budding using biotin-NeutrAvidin

For budding reactions using biotin-NeutrAvidin interactions to anchor triangles to the
membranes, GVs containing 3% biotinylated DOPE, 1% fluorescently labelled DOPE and 96%
DOPC were prepared. Origami triangles were folded with linker handles for biotinylated
oligonucleotides and linker handles for purification using magnetic beads. Biotinylated
oligonucleotides were already included in the folding reaction to remove unbound strands
and excess staples by gel extraction. The triangles were then mixed with a 500-fold excess
of NeutrAvidin and incubated for 1 h at 37 °C on an orbital shaker. Next, the sample was
transferred into a tube previously passivated with denatured BSA blocking buffer and mixed
with magnetic beads (Dynabeads M-270 Streptavidin, Invitrogen) coated with DNA strands
complementary to a section of the purification linker handles on the triangles. The sample
was incubated for 1 h at RT on a rotary shaker, and the beads now carrying the origami
triangles were washed 3-4 times with folding buffer containing 5 mM $MgCl_2$ and 300 mM
NaCl. Finally, an invader strand complementary to the full length of the purification linker
handles was added in great excess, and the solution was incubated at 37 °C for 1 h to
uncouple the origami from the magnetic beads by strand displacement. The NeutrAvidin
bearing triangles were then mixed with biotinylated GVs, and the $MgCl_2$ concentration was
adjusted to 65 mM by adding 1 M $MgCl_2$ solution (hyperosmotic conditions) or to 60 mM by
adding folding buffer (+210 mM $MgCl_2$; isosmotic conditions). The sample was incubated at
37 °C for 2.5 d.

Negative-staining transmission electron microscopy

Sample solutions were incubated on glow-discharged copper grids (400 mesh) with a carbon
support film (Science Services) for 4-8 minutes, depending on the sample concentration. The
solution was then blotted off using filter paper (Whatman). Then, the grid was washed once
using stain solution (2% aqueous uranyl formate + 25 mM NaOH), followed by incubating
another stain droplet on the grid for 30 s. Excess stain was blotted off using filter paper, and
the grid was air-dried before imaging.

Grids were typically imaged at x26-30k magnification using SerialEM with an FEI Tecnai T12

electron microscope operated at 120kV and a Tietz TEMCAM-F416 camera. Image contrast

was auto-levelled using Fiji[34].

For tomography, one-directional tilt series were acquired from -50° to +50° in steps of 2°

using SerialEM. The resulting image stacks were imported into Etomo (IMOD[35]), and images

were aligned without fiducials by cross-correlation with cumulative correlation switched on.

Tomograms were generated by filtered back-projection (gaussian filter cut-off: 0.35; fall-off:

0.035;). For visualisation, the z-stacks thus generated were imported into Fiji and z-projected

by summing up z-slices corresponding to the top, middle and bottom sections of the

respective particle of interest.

Cryogenic electron microscopy

Sample grids for cryoEM were prepared using an automated vitrification system (FEI Vitrobot

Mark V). First, grids (Quantifoil R1.2/1.3, Cu 200 mesh, 100 holey carbon films, 2 nm carbon

support film; for images of DCVs: Ted Pella, Cu 400 mesh, lacey carbon, <3 nm carbon

support film) were glow-discharged and then inserted into the Vitrobot. The conditions inside

the device's environmental chamber were kept at either 4 °C or 22 °C and 100% relative

humidity. Samples were typically incubated on the grids inside the chamber for 5 min and

plunge-frozen in liquid ethane (blot force: -1, 0 or 25; blot time: 2-3 s; blot total: 1; drain time:

0 s;). The grids were imaged with a spherical-aberration (Cs)-corrected Titan Krios G2

electron microscope (Thermo Fisher) operated at 300kV and equipped with a Falcon III 4k

direct electron detector (Thermo Fisher). Samples were imaged at x29k magnification and

defocus values between -1 and -4 μm. Images were auto-levelled in Fiji.

Agarose gel electrophoresis and gel quantification

Agarose gels were cast by dissolving UltraPure agarose (Invitrogen) in running buffer (0.5 x

TBE, Carl Roth) supplemented with $MgCl_2$ and optionally ethidium bromide (0.025%, Carl

Roth). Samples were run for 2 h at 90 V and then scanned using a Typhoon FLA 9500 laser

scanner (GE Healthcare) at a pixel size of 50 μm. Gels containing only monomeric origami

structures were run with 5.5 mM $MgCl_2$ in the gel and running buffer and an agarose

concentration of 1.5%. To resolve and preserve assembled shell structures, the gel and

running buffer were supplemented with 20 mM $MgCl_2$ and the agarose concentration was

reduced to 0.8%. Due to the elevated salt concentration in these gels, the gel boxes were

cooled in a water-ice bath, and the running buffer was exchanged every 45 min to replenish

lost magnesium. For visualisation, aggregates in the pockets were oversaturated, and the
bands' contrast underneath was auto-levelled.

Gel lanes were quantified using Fiji. Images were processed by removing outliers (radius: 5.0
pixels; pixel threshold: 200;), spanning a rectangular selection box across the region of
interest and plotting its intensity profile. Peak valleys were connected by straight lines, and
the area between the peaks and the added lines was measured. The resulting values were
normalised against the largest value in the same gel.

References

1.   Votteler, J. & Sundquist, W. I. Virus Budding and the ESCRT Pathway. *Cell Host Microbe* **14**, 232–241 (2013).

2.   Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nat. Rev. Mol. Cell Biol.* **19**, 313–326 (2018).

3.   Crowther, R. A., Pinch, J. T. & Pearse, B. M. F. On the structure of coated vesicles. *J. Mol. Biol.* **103**, 785–798 (1976).

4.   Morris, K. L. *et al.* Cryo-EM of multiple cage architectures reveals a universal mode of clathrin self-assembly. *Nat. Struct. Mol. Biol.* **26**, 890–898 (2019).

5.   Rothemund, P. W. K. Folding DNA to create nanoscale shapes and patterns. *Nature* **440**, 297–302 (2006).

6.   Douglas, S. M. *et al.* Self-assembly of DNA into nanoscale three-dimensional shapes. *Nature* **459**, 414–418 (2009).

7.   Dey, S. *et al.* DNA origami. *Nat. Rev. Methods Primer* **1**, 1–24 (2021).

8.   Wagenbauer, K. F., Sigl, C. & Dietz, H. Gigadalton-scale shape-programmable DNA assemblies. *Nature* **552**, 78–83 (2017).

9.   Wintersinger, C. M. *et al.* Multi-micron crisscross structures grown from DNA-origami slats. *Nat. Nanotechnol.* **18**, 281–289 (2023).

10.   Ong, L. L. *et al.* Programmable self-assembly of three-dimensional nanostructures from 10,000 unique components. *Nature* **552**, 72–77 (2017).

11.   Tikhomirov, G., Petersen, P. & Qian, L. Fractal assembly of micrometre-scale DNA origami arrays with arbitrary patterns. *Nature* **552**, 67–71 (2017).

12.   Langecker, M. *et al.* Synthetic Lipid Membrane Channels Formed by Designed DNA Nanostructures. *Science* **338**, 932–936 (2012).

13.   Burns, J. R., Stulz, E. & Howorka, S. Self-Assembled DNA Nanopores That Span Lipid Bilayers. *Nano Lett.* **13**, 2351–2356 (2013).

14.   Burns, J. R. *et al.* Lipid-Bilayer-Spanning DNA Nanopores with a Bifunctional Porphyrin Anchor. *Angew. Chem. Int. Ed.* **52**, 12069–12072 (2013).

15.   Khmelinskaia, A., Mücksch, J., Petrov, E. P., Franquelim, H. G. & Schwille, P. Control of Membrane Binding and Diffusion of Cholesteryl-Modified DNA Origami Nanostructures by DNA Spacers. *Langmuir* **34**, 14921–14931 (2018).

16.   Journot, C. M. A., Ramakrishna, V., Wallace, M. I. & Turberfield, A. J. Modifying Membrane Morphology and Interactions with DNA Origami Clathrin-Mimic Networks. *ACS Nano* **13**, 9973–9979 (2019).

17.   Grome, M. W., Zhang, Z., Pincet, F. & Lin, C. Vesicle Tubulation with Self-Assembling
DNA Nanosprings. *Angew. Chem. Int. Ed.* **57**, 5330–5334 (2018).

18.   Liu, L., Xiong, Q., Xie, C., Pincet, F. & Lin, C. Actuating tension-loaded DNA clamps
drives membrane tubulation. *Sci. Adv.* **8**, eadd1830 (2022).

19.   Franquelim, H. G., Khmelinskaia, A., Sobczak, J.-P., Dietz, H. & Schwille, P.
Membrane sculpting by curved DNA origami scaffolds. *Nat. Commun.* **9**, 811 (2018).

20.   Yang, J. *et al.* Modulating Lipid Membrane Morphology by Dynamic DNA Origami
Networks. *Nano Lett.* **23**, 6330–6336 (2023).

21.   Baumann, K. N. *et al.* Coating and Stabilization of Liposomes by Clathrin-Inspired
DNA Self-Assembly. *ACS Nano* **14**, 2316–2323 (2020).

22.   Kocabey, S. *et al.* Membrane-Assisted Growth of DNA Origami Nanostructure Arrays.
ACS Nano **9**, 3530–3539 (2015).

23.   Bassereau, P., Sorre, B. & Lévy, A. Bending lipid membranes: Experiments after W.
Helfrich's model. *Adv. Colloid Interface Sci.* **208**, 47–57 (2014).

24.   Simunovic, M., Evergren, E., Callan-Jones, A. & Bassereau, P. Curving Cells Inside
and Out: Roles of BAR Domain Proteins in Membrane Shaping and Its Cellular Implications.
Annu. Rev. Cell Dev. Biol. **35**, 111–129 (2019).

25.   Sweitzer, S. M. & Hinshaw, J. E. Dynamin Undergoes a GTP-Dependent
Conformational Change Causing Vesiculation. *Cell* **93**, 1021–1029 (1998).

26.   Boucrot, E. *et al.* Membrane Fission Is Promoted by Insertion of Amphipathic Helices
and Is Restricted by Crescent BAR Domains. *Cell* **149**, 124–136 (2012).

27.   Rossman, J. S., Jing, X., Leser, G. P. & Lamb, R. A. Influenza Virus M2 Protein
Mediates ESCRT-Independent Membrane Scission. *Cell* **142**, 902–913 (2010).

28.   Sigl, C. *et al.* Programmable icosahedral shell system for virus trapping. *Nat. Mater.*
20, 1281–1289 (2021).

29.   Gerling, T., Wagenbauer, K. F., Neuner, A. M. & Dietz, H. Dynamic DNA devices and
assemblies formed by shape-complementary, non–base pairing 3D components. *Science*
347, 1446–1452 (2015).

30.   Weinberger, A. *et al.* Gel-Assisted Formation of Giant Unilamellar Vesicles. *Biophys.*
J. **105**, 154–164 (2013).

31.   Dannhauser, P. N. & Ungewickell, E. J. Reconstitution of clathrin-coated bud and
vesicle formation with minimal components. *Nat. Cell Biol.* **14**, 634–639 (2012).

32.   Ford, M. G. J. *et al.* Curvature of clathrin-coated pits driven by epsin. *Nature* **419**,
361–366 (2002).

33.   Park, J. H. *et al.* Temperature-Induced Denaturation of BSA Protein Molecules for

Improved Surface Passivation Coatings. *ACS Appl. Mater. Interfaces* **10**, 32047–32057 (2018).

34.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

35.   Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer Visualization of Three-

Dimensional Image Data Using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996).

SUPPLEMENTARY INFORMATION

Supplementary Figure 1 | Attachment of origami triangles to vesicles via cholesterol-bearing oligonucleotides. Linker handles extending from the origami surface are hybridised to cholesterol-bearing oligonucleotides with complementary sequences. Cholesterol acts as a lipid membrane anchor to tether the triangles to lipid vesicles.

Supplementary Figure 2 | The influence of cholesterol position and orientation on origami triangles. (**a**) The overall reach of cholesterol-bearing oligonucleotides (chol-oligo) determines the migration behaviour of origami triangles. When cholesterols are positioned closer to the outer edge and the stacking contacts in a distal configuration (A_D), bands in an agarose gel appear shifted and smeared compared to the same triangles not hybridised to chol-oligos. This effect could be reduced by changing the orientation of the cholesterol to a proximal configuration with respect to the origami, such that the reach of the cholesterol is minimised (A_P). When chol-oligos are positioned closer to the centre of the structure, and thus further away from the stacking contacts, the migration distance of these triangles is comparable to that of sample A_P, albeit with more smearing. Combining both variants to double the cholesterol count per triangle from 9 to 18 moieties yielded bands appearing as combinations of the individual variants. (**b**) The reach of cholesterol influences the assembly of cholesterol-decorated triangles. An assembly control (AC) of triangles lacking protruding linker handles assembles mostly into icosahedral species, as do triangles with protruding linker handles not hybridised to chol-oligos. When chol-oligos are introduced, distal cholesterol (A_D) promotes the formation of primarily octahedral species, whereas proximal cholesterol (A_P) does not noticeably influence the assembly behaviour. Reducing the reach of

individual cholesterols either by positioning them closer to the centre of the triangles or by

choosing proximal configurations reduces unwanted interactions.

Supplementary Figure 3 | The influence of cholesterols per triangle in assembly reaction

containing GVs. Whilst optimising cholesterol positioning on the triangle can reduce

unexpected assembly behaviour, introducing lipid vesicles yet again shifts the assembly

towards octahedra, the degree of which increases with the number of cholesterols per

triangle.

Supplementary Figure 4 | GVs shift the assembly of cholesterol-bearing triangles

towards octahedral shells. The assembly behaviour of cholesterol-bearing triangles, tagged

with Atto643 for visualisation, depends on the presence or absence of GVs. Without GVs,

triangles mainly assemble into icosahedral shells as per their design. With GVs, however,

most shells detected by agarose gel electrophoresis are octahedral. The abundance of

smaller, intermediate species between closed shells and monomeric triangles increases with

the number of cholesterols per triangle and mostly represents aggregated triangles.

670
671

Supplementary Figure 5 | Triangle assembly is the driving force behind DCV formation.
The importance of shell assembly for DCV formation was studied by incubating triangles at various $MgCl_2$ concentrations at (65 mM) or below (10-40 mM) the typical threshold concentration for shell assembly. For visualisation by agarose gel electrophoresis, the triangles were hybridised to Atto643 bearing oligonucleotides, and the GVs contained a fraction of Atto488 labelled DOPE. Furthermore, two triangle species were prepared: A non-passivated version with blunt-ended stacking contacts and a passivated version where the staple oligonucleotides at the base stacking contacts were extended by introducing additional unpaired thymidines. Whereas the non-passivated version formed DCVs as seen by the bands in the lipid channel (lower panel) colocalising with the shells in the DNA channel (upper panel), only faint bands below the typical height of DNA shells were found in samples containing the passivated version. These results were compared by TEM, which further revealed that while the passivated triangle samples contained mostly monomeric triangles, there was a small number of aggregated triangles held together by small vesicles, likely representing the faint bands. Shell assembly is, therefore, a main factor behind DCV formation. Scale bars: 100 nm.

688
689
690
691

Supplementary Figure 6 | Analysis of DCV formation by colocalisation analysis of DNA and lipid bands. The relative efficiency of DCV formation could be analysed by agarose gel electrophoresis using fluorescently labelled DNA oligos attached to the origami structure or ethidium bromide, and a small fraction of fluorescently labelled DOPE species in the lipid mixture comprising the vesicles. Whereas multiple bands were detected in the DNA channel representing triangles in monomeric or assembled states, lipid signal was only seen at the height of assembled shells. The absence of lipid signal colocalised with monomeric triangles suggests that lipids predominantly enter the gel as part of a DCV. The gel pockets contain origami-coated GVs, which were oversaturated to better visualise the much fainter bands below.

Supplementary Figure 7 | Shell preassembly reduces DCV yields. Cholesterol-bearing origami triangles were incubated without GVs (containing a small fraction of DOPE-Atto643) for 4 h at 37 °C in either 5 mM (monomer conditions) or 60 mM (assembly conditions) $MgCl_2$ to study the effect partially assembled shells have on DCV yields. Afterwards, GVs were added, and the $MgCl_2$ concentration in the 5 mM controls was increased to 60 mM. Aliquots taken and frozen at t = 0 d were compared to samples incubated at 37 °C for 2.5 d using agarose gel electrophoresis, and DCV yields were analysed by comparing lipid bands. Whereas t = 0 d aliquots looked alike in samples pre-incubated in either 5 mM or 60 mM $MgCl_2$, we finally obtained stronger lipid signals in samples initially incubated at 5 mM. We deduce that DCVs form best when monomeric triangles are assembled directly on the membrane. Partially assembled shells may fail to bind GVs efficiently due to steric hindrance, resulting in reduced DCV yields.

719