Structural features of heteromeric channels composed of CALHM2 and

CALHM4 paralogs

Katarzyna Drożdżyk[1], Martina Peter[1] and Raimund Dutzler[1*]

[1]Department of Biochemistry, University of Zurich, Zurich, Switzerland

*For correspondence: dutzler@bioc.uzh.ch (RD)

Abstract

The CALHM proteins constitute a family of large pore channels that contains six closely related

paralogs in human. Two family members, CALHM1 and 3, have been associated with the release

of ATP during taste sensation. Both proteins form heteromeric channels that activate at positive

potential and decreased extracellular Ca^{2+} concentration. Although the structures of several family

members displayed large oligomeric organizations of different size, their function has in most

cases remained elusive. Our previous study has identified the paralogs CALHM2, 4 and 6 to be

highly expressed in the placenta and defined their structural properties as membrane proteins

exhibiting features of large pore channels with unknown activation properties (Drozdzyk et al.,

2020). Here we investigated whether these placental paralogs would form heteromers and

characterized heteromeric complexes consisting of CALHM2 and CALHM4 subunits using

specific binders as fiducial markers. Both proteins assemble with different stoichiometries with

the largest population containing CALHM2 as predominant component. In these oligomers, the

subunits segregate and reside in their preferred conformation found in homomeric channels. Our

study has thus revealed the properties that govern the formation of CALHM heteromers in a

process of potential relevance in a cellular context.

Introduction

Large pore channels encompass a heterogeneous group of membrane proteins in higher eukaryotes, which form large oligomeric assemblies that facilitate the diffusion of diverse substrates with poor selectivity, including metabolites and signaling molecules such as the nucleotide ATP (Syrjanen et al., 2021). The calcium homeostasis modulators (CALHM) constitute one family of large pore channels that contains six members in humans (Ma et al., 2016). Their role in physiology is currently best understood for the paralogs CALHM1 and 3. Both proteins are expressed in type II taste bud cells to contribute to the sensation of umami, sweet and bitter taste (Ma et al., 2018b; Taruno et al., 2013b). There, they are involved in the non-vesicular release of ATP as a consequence of membrane depolarization following the initial sensation of tastants by G-protein coupled receptors (Taruno et al., 2013a). ATP in turn acts as neurotransmitter to activate P2X-receptors on the postsynaptic side. CALHM proteins reside in a non-conducting state at resting potentials and are activated in response to membrane depolarization with a decrease in extracellular Ca^{2+} shifting the threshold of activation towards more hyperpolarizing voltages, thereby allowing conduction also at negative potentials (Ma et al., 2012). The open channel shows poor selectivity and is permeable to the multivalent anion ATP. Although these properties are shared by homomeric assemblies of CALHM1 in heterologous expression systems, their slow kinetics of activation would prohibit fast signaling and thus does not reflect the behavior observed in taste bud cells (Ma et al., 2012). To enhance its activation properties, CALHM1 assembles as heteromeric complex with its paralog CALHM3, which on its own does not form functional ion channels (Ma et al., 2018b). The physiological role of other CALHM paralogs is currently much less well understood, although the ubiquitously expressed CALHM2 was also associated with ATP

release (Liao et al., 2023; Ma et al., 2018a) and CALHM6 was recently described as channel in

immunological synapses of macrophages (Danielli et al., 2023).

Based on their assumed topology containing four membrane-spanning segments, the CALHM

family was originally predicted to share a close structural relationship with large pore channels of

the connexin, pannexin and LRRC8 families (Siebert et al., 2013). However, this relationship was

refuted in numerous structures of homomeric assemblies of different CALHM paralogs that have

recently been determined by cryo-electron microscopy (cryo-EM) from samples obtained from

heterologous expression in mammalian and insect cell culture (Choi et al., 2019; Demura et al.,

2020; Drozdzyk et al., 2020; Foskett, 2020; Liu et al., 2020; Ren et al., 2022; Ren et al., 2020;

Syrjanen et al., 2020; Yang et al., 2020). Although sharing the same number of membrane-

spanning segments, their arrangement is unique and differs from other large pore channels, with

an extended helix located on the C-terminus and forming the core of a cytoplasmic domain further

mediating subunit interactions. These structures show channels that assemble into higher

oligomers around an axis of symmetry that presumably defines the conduction path. These

assemblies are heterogeneous in nature, with paralogs frequently adopting different oligomeric

states in the same sample. The smallest assemblies were generally observed for CALHM1, where

structures from different species contained between seven and nine subunits (Demura et al., 2020;

Ren et al., 2022; Ren et al., 2020; Syrjänen et al., 2023; Syrjanen et al., 2020). The size of the

respective assemblies is generally larger for other CALHM paralogs, ranging from decamers to

tridecamers (Choi et al., 2019; Drozdzyk et al., 2020; Liu et al., 2020; Syrjanen et al., 2020). In

detergent solution, these channels do in certain cases pair to form gap-junction like organizations,

although there is currently no evidence for such transcellular interactions in a physiological

environment (Choi et al., 2019; Syrjanen et al., 2020). In oligomeric channels, the pore is lined by

the first transmembrane segment (TM1) of each subunit, which forms an inner, of two concentric

rings, with the outer ring being constituted by TM2-4, with TM2 and 4 of neighboring subunits

engaging in extended interactions (Choi et al., 2019; Drozdzyk et al., 2020; Liu et al., 2020;

Syrjanen et al., 2020). The loosely packed TM1 is found in distinct conformations in different

structures to shape the pore geometry and thus presumably alter conduction properties (Choi et al.,

2019; Drozdzyk et al., 2020).

We have previously studied the structural and functional properties of the CALHM paralogs 2, 4

and 6, whose transcripts were found to be abundant in placental epithelia in dependence of the

developmental stage of the organ (Drozdzyk et al., 2020). These placental paralogs all form large

oligomers, which in CALHM4 consists of similar sized populations of decameric and undecameric

assemblies in the same sample. Both oligomers were found to pair via contacts on their intracellular

side into large tubular structures. In this sample, both assemblies adopt a similar subunit

conformation, leading to large cylindrical pores that are constricted by a short N-terminal helix

preceding TM1, which is oriented parallel to the membrane plane and projects towards the pore

axis (Drozdzyk et al., 2020). In the same study, a similar pairing of channels containing between

11 and 12 subunits was also found for CALHM2, where the preferred orientation of the particles

within the sample has prevented a three-dimensional reconstruction at high resolution. In contrast,

no dimerization was detected in a CALHM6 preparation, where, although also existing as

undecamers, a decameric assembly was prevalent and thus better resolved. As consequence of a

conformational rearrangement of TM1, which has detached from its interaction with the rest of the

subunit, resulting in a higher mobility, the CALHM6 pore has adopted a conical shape with

decreasing diameter towards the intracellular side suggesting that it might reside in a different

functional state (Drozdzyk et al., 2020).

Despite the observed large assemblies, none of the three paralogs overexpressed in *X. laevis*

oocytes showed any activity in electrophysiology experiments under conditions where CALHM1

was conductive, regardless of their expression at the cell surface, suggesting that they may be

activated by different stimuli (Drozdzyk et al., 2020). This was unexpected in light of the distinct

conformations observed for CALHM4 and 6 and their large apparent pore diameter, which is

particularly pronounced in case of CALHM4. In this respect, it was remarkable to find continuous

density with bilayer-like features inside the pore of CALHM4, matching the hydrophobic character

of the pore lining, which might explain the lack of activity observed for this protein (Drozdzyk et

al., 2020). This density is not found in CALHM6, where the altered shape of the pore could not

accommodate a bilayer. The functional properties of placental CALHM paralogs thus remained

elusive.

Since the formation of heteromers composed of CALHM1 and 3 paralogs strongly altered the

functional properties of the respective homomers, we wondered whether the tree paralogs

expressed in the placenta would share a similar property and whether their heteromerization would

increase their activity under the investigated conditions. To clarify this question, we have here

investigated the ability of the overexpressed CALHM homologs 2, 4 and 6, to form heteromeric

assemblies. Robust interactions were identified between CALHM2 and 4, but to a much lower

extent between any of the two paralogs and CALHM6. The subsequent structural characterization

of CALHM2/4 heteromers, employing synthetic subunit-specific nanobodies (sybodies), showed

channels with a broad subunit distribution. Predominant species contain a predominance of

CALHM2 with two to four CALHM4 subunits, where both paralogs segregate in the heteromeric

assemblies. While subunits maintain their distinct structural features found in homomeric

channels, they also influence the conformations of contacting subunits.

Results

Formation of CALHM heteromers

The formation of heteromers is a hallmark of different eukaryotic ion channels, with their subunit composition shaping functional properties. Such heteromerization has also been found in the CALHM family of large-pore channels, where the paralogs CALHM1 and 3 oligomerize to form channels that are activated by membrane depolarization and the decrease of the extracellular Ca^{2+}-concentration (Ma et al., 2018b). To probe whether this property would also extend to other family members, we have initially investigated the ability of the three placental paralogs CALHM2, 4 and 6 to form heteromeric assemblies. To this end, we have co-transfected HEK293 cells with pairs of constructs coding for different paralogs and containing unique tags for affinity purification. After solubilization, specific subunits were captured via their attached tags and the sample was assayed for the co-purification of the second subunit by SDS-PAGE and Western blot. In that way, we have identified robust heteromerization of the paralogs CALHM2 and 4, and weak pairing of both paralogs with CALHM6, thus suggesting that specific heteromerization might be a property that is inherent to certain CALHM paralogs (Figure 1A).

After identifying the heteromerization of CALHM2 and 4, we have tested whether this process has altered the activation properties of these channels. To this end we have expressed either single or pairs of different CALHM paralogs in HEK293 cells and investigated their functional properties by patch-clamp electrophysiology. When recorded in the whole-cell configuration, we were able to reproduce the previously observed phenotype of CALHM1 to form channels that activate at positive voltages (Figure 1B, Figure 1-figure supplement 1A). The observed functional properties are enhanced upon co-expression of CALHM1 and 3 subunits, which results in the increase of

currents and activation kinetics compared to CALHM1, when recorded in the same manner (Figure

1C, Figure 1-figure supplement 1B). In contrast, neither CALHM2 nor CALHM4 show

appreciable currents under equivalent conditions, which conforms with a previous investigation

performed in *X. laevis* oocytes (Drozdzyk et al., 2020) (Figure 1D, Figure 1-figure supplement

1C). Finally, we have investigated whether their assembly as heteromers would enhance the

activity of CALHM2 and CALHM4 and thus proceeded with the functional characterization of

cells that were co-transfected with plasmids encoding each subunit. However, similar to the

respective homomers, we were unable to detect any current response at positive voltage, thus

emphasizing that these proteins share an activation mechanism that is distinct from CALHM1/3

(Figure 1E, F, Figure 1-figure supplement 1D, E).

Selection of paralog-specific binders

Following, we set out to characterize the structural properties of CALHM2/4 heteromers by cryo-

EM. This task is complicated by the unknown arrangement of subunits sharing similar structural

features and the potentially large compositional heterogeneity of heteromers, which might prohibit

classification and alignment of oligomeric complexes. We have thus initially engaged in the

generation of paralog-specific binders that can be used as fiducial markers to aid the identification

of subunits in heteromeric channels. As in previous projects, we have decided for synthetic

nanobodies (termed sybodies), which were selected from large construct libraries by different

display methods (Zimmermann et al., 2020). In independent efforts, we have selected binders

against homomeric assemblies of CALHM2 and CALHM4 (Figure 2-figure supplement 1) and

subsequently used them for a structural characterization of respective homomers to define their

binding epitopes and their effect on subunit conformations.

Features of the CALHM4/SbC4 complex

Our selection against CALHM4 was carried out with paired assemblies of channels containing

either ten or eleven subunits each, which is reflected in the low elution volume of the sample during

size exclusion chromatography (Figure 2-figure supplement 1B). Dimerization in CALHM4

proceeds at its intracellular region, resulting in large particles, which expose their extracellular

parts at both ends of a tubular assembly (Drozdzyk et al., 2020). This sample has allowed us to

isolate a specific binder, termed Sb^{CALHM4} or short SbC4, that forms a tight complex with the

CALHM4 channel pair, which remains intact during size exclusion chromatography as confirmed

by SDSPAGE (Figure 2-figure supplement 1A, B). Upon characterization by surface plasmon

resonance, we measured a binding affinity of about 100 nM (Figure 2-figure supplement 1C).

Conversely, we did not find any evidence of this protein to engage in strong interactions with

CALHM2 (Figure 2-figure supplement 1D). When investigated by cryo-EM, we observed two

populations of particles either exhibiting D10 or D11 symmetry (corresponding to dimers of

decameric and undecameric channels) that closely resemble equivalent structures of CALHM4

determined in previous studies (Drozdzyk et al., 2020) (Figure 2-figure supplement 2). Between

the two populations, the reconstruction of the density with D10 symmetry imposed was of higher

resolution and we thus restrict our detailed analysis to this structure (Figure 2A, B, Figure 2-figure

supplement 2). Subunits in the oligomeric assembly are arranged around an axis of symmetry,

which presumably defines the pore of a large channel (Figure 2B). Each subunit contains four

membrane-spanning helices, three of which (TM2-4) are tightly interacting to form an outer ring

with large diameter that separates the inside of the pore from its surrounding membrane

environment (Figure 2C, D). In this ring, TM2 and 4 of adjacent subunits engage in extended

interactions (Figure 2D). In contrast, TM1 forms an inner ring of helices lining the pore, with the

bent helical N-terminus (NH) running perpendicular to the membrane and projecting towards the

pore axis (Figure 2D). TM1 packs against TM3 of the outer ring but is not involved in direct

subunit interactions. Within each channel, we find residual density of low resolution, whose

distribution resembles the headgroup region of a lipid bilayer, which was thus previously ascribed

to a membrane assembled in the lumen of the pore (Drozdzyk et al., 2020) (Figure 2E). In this

cryo-EM map, we find density of sybodies binding at the extracellular side of each subunit forming

two rings of binders at both ends of the assembly (Figure 2A, B). Although not sufficiently well

resolved to define detailed interactions with certainty, neighboring sybodies appear not to clash

with epitopes being confined to a single subunit, which provides ideal properties for the

identification of CALHM4 subunits in heteromeric channels.

Features of the CALHM2/SbC2 complex

In case of CALHM2, our selection efforts have allowed us to identify a binder termed Sb^{CALHM2}

(short SbC2) forming tight complexes with homomeric assemblies (Figure 2-figure supplement

1A, E-G). This sybody remains bound to the channel during size exclusion chromatography, which

is manifested in the decreased elution volume of the complex and the presence of both proteins in

peak fractions as confirmed by SDS-PAGE (Figure 2-figure supplement 1E). As in case of SbC4,

the binding is subunit specific (Figure 2-figure supplement 1F, G). We have then proceeded with

the structural characterization of CALHM2/SbC2 complexes by cryo-EM and found two

populations of particles showing either C10 or C11 symmetry. In this data-set, only the

reconstruction of undecameric particles resulted in high resolution density that resembles

structures of CALHM2 determined in previous studies (Choi et al., 2019; Demura et al., 2020;

Syrjanen et al., 2020) (Figure 3A, B, Figure 3-figure supplement 1). In contrast to the CALHM4

structure, we did not detect any dimerization of channels (Figure 3-figure supplement 1). Another

pronounced difference of the CALHM2 structure concerns the conformation of TM1, which is

detached from its interaction with TM3 as found in CALHM4 and instead projects towards the

pore axis (Figure 3C). Due to its high mobility, the density of this helix is only defined at its C-

terminal end at the boundary to the extracellular loop connecting to TM2, whereas its remainder

towards the N-terminus is not resolved (Figure 3C). This arrangement of TM1 in an 'up' position

resembles the conformation observed in a previous structure of a complex of CALHM2 with its

putative inhibitor rubidium red, although its density in the inhibitor complex is better defined (Choi

et al., 2019). In contrast to CALHM4, where TM3 and TM4 of the same subunit tightly interact

throughout, in CALHM2 both helices have moved apart at their intracellular halve (Figure 3D). In

the structure, the resulting narrow fenestration connecting the channel interior and the membrane

is filled by a bound lipid (Figure 3D). Finally, unlike in the CALHM4 structure where TM1 resides

in a 'down' conformation, the lumen of the CALHM2 channel does not contain pronounced

bilayer-like residual density (Figure 3E). In the CALHM2 assembly, strong density of appropriate

size and shape at each of the subunits defines the location of SbC2 binding to the intracellular side

of the channel via its extended CDR3 loop (Figure 3A-E). On CALHM2, the epitope encompasses

residues located on the intracellular end of TM3, the region immediately following TM4 and part

of the C-terminal loop connecting the short helices CT2H and CT3H of the contacted subunit.

Additionally, the very C-terminus of the adjacent subunit (CT^{+1}, located in counterclockwise

position when viewed from the extracellular side), which contains six additional residues added to

the expression construct contributes to the interaction (Figure 3D). This C-terminus is unstructured

in absence of the sybody but contacts the region preceding CT3H on the neighboring CALHM2

subunit and the CDR3 region of SbC2 to become partly buried in the binding interface (Figure

3D). As a consequence, about 75% of the binding interface covering about 2'200 Å^2 of the

combined molecular surface is contributed by the CT^{+1}. Despite the described involvement of CT^{+1}

to SbC2 binding, sybodies bound to neighboring subunits do not overlap and bind remote from the

primary subunit interface and thus clearly label CALHM2 subunits engaging in homomeric

interactions in the +1 position.

Characterization of CALHM2/4 heteromers

Their high selectivity combined with the distinct location of their epitopes make the two sybodies

complementary tools for the identification of paralogs in heteromeric channels. We have thus used

both binders for the structure determination of channels consisting of CALHM2 and CALHM4

subunits (CALHM2/4). To this end, we have transfected HEK-293 cells with equimolar amounts

of DNA encoding constructs of the respective paralogs and isolated heteromeric channels by

tandem purification, sequentially capturing the different tags that are attached to either construct.

This strategy has allowed us to isolate channels containing both subunits, which constitute a small

fraction (<10%) of the total expressed protein, most of which assembles as homomers. We have

then added sybodies in stoichiometric excess and determined two datasets of CALHM2/4-sybody

complexes, one containing both binders and a second SbC4 only. Both datasets are of high quality

and provide insight into the organization of heteromeric channels with the sybodies defining the

distribution of CALHM2 and 4 subunits (Figure 4-figure supplements 1 and 2). In both datasets,

the position of CALHM4 subunits is marked by the binding of SbC4 to the extracellular part of

the protein, whereas the dataset containing both sybodies, SbC2 additionally labels CALHM2

subunits. Both samples show a heterogeneous distribution of complexes with variable ratio of the

two paralogs, which in all cases segregate into clusters to minimize heteromeric interactions

(Figure 4-figure supplements 1 and 2). A 3D classification shows particles of three distinct

oligomeric organizations. A population of proteins with D10 symmetry exhibits the characteristic

organization of CALHM4 channels, which pairwise interact on their intracellular side (Figure 4A,

Figure 4-figure supplement 2B). These channels consist predominantly of CALHM4 with

interspersed CALHM2 subunits (Figure 4A). In contrast, in two larger populations of undecameric

and dodecameric channels that do not dimerize, CALHM2 subunits predominate (Figure 4B, C).

A further classification resulted in a distribution of channels containing two to four consecutively

arranged CALHM4 subunits, thus suggesting that tight homotypic interactions between subunits

prevail in heteromeric channels (Figure 4B, C).

Features of heteromeric CALHM2/4 channels

Although our 3D classification has defined a broad distribution of different heteromeric assemblies

of CALHM2/4 channels, the population in each distinct assembly is too small to reach high

resolution. We have thus pooled the classes of all undecameric channels containing 2-4 copies of

CALHM4, and proceeded with further map refinement, which has allowed us to obtain maps

extending to global resolutions of 3.8 Å and 3.3 Å for the datasets of samples containing either

SbC4 only (CALHM2/4/SbC4), or both SbC4 and SbC2 (CALHM2/4/SbC2/SbC4), respectively

(Figure 5, Figure 4-figure supplements 1 and 2). The two cryo-EM densities provide congruent

views of the relative arrangement and conformational preferences of CALHM2 and 4 subunits in

heteromeric channels. In both datasets, the two positions uniformly occupied by CALHM4

(referred to as 1 and 2 when numbered in counterclockwise direction from the extracellular side)

are well defined and clearly discernable by SbC4 bound to its extracellular epitope (Figure 5).

Similarly, the seven following positions (3-9), occupied by CALHM2, are well defined, whereas

the two positions adjacent to the two CALHM4 subunits in clockwise direction (11 and 10), which

contain CALHM2 or 4 at different ratios are of somewhat lower resolution due to the averaging of

two different subunits. This feature is particularly pronounced in the dataset of

CALHM2/4/SbC2/SbC4, containing both sybodies, where position 11 bears closer resemblance to

CALHM4, as further evidenced by weak residual density of SbC4 (Fig. 5D). Conversely, features

in the neighboring position 10 are closer to CALHM2, as expected for an average of channel

populations containing two to four CALHM4 subunits (Figure 5D). In the

CALHM2/4/SbC2/SbC4 complex, most of the positions occupied by CALHM2 (*i.e.* positions 4-

8) are also distinguished by strong density of SbC2 binding to its intracellular epitope (Figure 5D).

However, since the C-terminus of the neighboring CALHM2 (CT^{+1}) contributes to binding, we

find weaker density in position 9 (Figure 5D). In contrast, the weaker density of SbC2 in position

3, which is fully occupied by CALHM2, is probably a consequence of the distinct conformation

of this particular subunit, which compromises sybody binding.

Conformational properties of single subunits in CALHM2/4 channels

Whereas in CALHM2/4 heteromers, the CALHM4 subunits reside in the preferential 'down'

conformation observed in homomeric channels, irrespective of their relative position in the

channel, the conformational properties of CALHM2 subunits are influenced by their environment

(Figure 6A). CALHM2 subunits that are distant from CALHM4 predominantly reside in the

familiar 'up' position of TM1 observed in in the structure of the CALHM2/SbC2 complex (Figure

6A). In contrast, CALHM2 in position 3 shows a distinct conformation, where TM1 has rearranged

to maximize its contacts with TM3, resembling the 'down' conformation of adjacent CALHM4

subunits (Figure 6A, B). As a consequence of this interaction, TM2 and TM3 have moved towards

the outer ring of helices constituted by TM2-4 to close the fenestration between TM3 and TM4,

which in the CALHM2/SbC2 complex was found to be occupied by a lipid (Figure 3D, Figure 6B,

C). These movements affect the local structure of the epitope recognized by SbC2 as manifested

in the weaker density observed in the CALHM2/4/SbC2/SbC4 complex, which indicates a

deterioration of the interaction (Figure 5D). The influence of CALHM4 on CALHM2

conformations is also observed in position 4, where we find a mix of both states, and in weak

density of the 'down' conformation in CALHM2 subunits located remote from CALHM4. The

distinct preference of subunit conformations is also manifested in the distribution of residual

density in the pore region, which is evident in low-pass filtered maps (Figure 6D). Here we find a

pronounced toroidal density surrounding subunits in a down-conformation, which contains

bilayer-like features observed in CALHM4 homomers and which is absent around CALHM2

subunits sharing a predominant up-conformation (Figures 2D, 3E and 6D). Together, our results

have defined the oligomeric organization of heteromeric channels composed of CALHM2 and 4

subunits and they show how their assembly impacts the conformational preference of subunits.

CALHM Subunit interfaces

The CALHM2/4 heteromer contains four distinct subunit interfaces that provide insight into the

structural basis for heteromerization. Whereas the homotypic interactions between pairs of

CALHM2 or CALHM4 subunits are essentially identical as in the respective homomeric channels,

the structure also reveals the well-defined interaction interface between CALHM4 and 2 subunits

(viewed from the extracellular side in counterclockwise direction) whereas the details of the 2/4

interface are blurred by the mix of subunits at the 10 and 11 positions (Figures 5 and 7). With

5,400 $Å^2$, the interface buried between two CALHM2 subunits is 40% larger compared to the

corresponding CALHM4 interface (burying 3960 $Å^2$), with latter being similar in size to the

heteromeric CALHM4/CALHM2 interface (3572 $Å^2$, Figure 7A-C). The larger surface buried

between CALHM2 subunits, which presumably translates in a stronger interaction, reflects the

observed prevalence of homomers upon coexpression of subunits. In CALHM2/4 heteromers it

accounts for the dominance of one paralog in different channel populations combined with the

observed segregation of subunits (Figure 4).

The subunit interfaces consist of two separated parts, one contributed by the membrane-inserted

helices TM2 and TM4, and the other by the cytoplasmic parts following TM4, comprising the

helix CTH and subsequent regions (Figure 7A-C). Within the membrane, contacts are similar and

dominated by hydrophobic residues several of which are conserved between both paralogs,

although polar contacts extend further to the extracellular region in CALHM2 (Figure 7A). In the

cytoplasmic domain, the interactions show pronounced differences with the contact area buried by

CALHM2 being considerably larger (3,003 $Å^2$) compared to CALHM4 (1,909 $Å^2$), which is a

consequence of the parts located C-terminally to CTH, which differ in both paralogs (Figure 7D,

E). Whereas interactions mediated by the conserved CTH are similar between CALHM2 and 4,

the following region in CALHM2 consists of three short helices (CTH', CT2H and CT3H), and

connecting loops which wrap around CTH to increase the interface and contribute to subunit

interactions (Figure 7D). In CALHM4, of these three short helices only CT3H is present which

engages in similar conserved interactions with the CTH as the equivalent region in CALHM2,

although with different subunits (Figure 7E). While, due to the altered location of CT2H, the helix

in CALHM2 primarily binds to an epitope located in the interacting subunit in the +1 position, in

CALHM4 the helix binds to the equivalent region in the same subunit. A generally similar pattern

of interactions is also found between subunits at the CALHM4/2 interface where the buried surface

area at the C-terminus is even smaller than in CALHM4/4 interfaces (1,671 Å^2, Figure 7F). In

summary, we find conserved complementary contacts between CALHM2 and CALHM4 subunits,

which promote the formation of heteromeric channels. However, the larger area that is buried in

interfaces relating equivalent subunits underlines the observed prevalence of homomeric

interactions.

Discussion

The formation of heteromers of close paralogs of a protein family is a common mechanism in

eukaryotic ion channels to confer distinct functional properties to an oligomeric assembly.

Heteromeric assemblies are found in pentameric ligand-gated ion channels, glutamate receptors

and volume-regulated anion channels of the LRRC8 family (Baumann et al., 2001; Greger and

Mayer, 2019; Voss et al., 2014). In the CALHM family of large pore channels, the formation of

heteromers has been demonstrated for the paralogs CALHM1 and CALHM3, where the pairing of

subunits results in a strong improvement of the activation properties compared to their homomeric

equivalents (Figure 1B, C) (Ma et al., 2018b). We reasoned that a similar process might also extend

to other family members and thus investigated the heteromerization of three paralogs that we have

previously shown to be expressed in placental epithelia but for which we were so far unable to

detect any function (Drozdzyk et al., 2020). Our experiments here have demonstrated the ability

of at least two paralogs, CALHM2 and 4, to form heteromeric assemblies (Figure 1A). Although

the existence of such mixed assemblies in a cellular context is still uncertain, our study has shed

light on the detailed features of the interaction of both subunits. In that way it has revealed

energetic relationships between homo- and heteromeric subunit pairing, providing insight that

might also be relevant for other ion channels.

The ability to form heteromers appears specific to certain members of the CALHM family, since

not all of the investigated combinations of paralogs were equally suited to interact (Figure 1A).

The general, ability of subunits to become part of the same assembly is encoded in their structure.

Based on the comparison of known structures, there is an apparent incompatibility between one

group consisting of CALHM1 and 3 and the second encompassing CALHM 2, 4, 5, and 6. This is

reflected in the distinct orientation of the C-terminal domain, which constitutes a large part of the

interaction interface (Figure 8A, B). In its orientation, the first helix of this domain termed CTH

is similar in CALHM2, 4, 5 and 6 and, although different degrees of pairing have been detected

between these subunits (Figure 1A), the energetic penalty to form heteromers is presumably small

(Figure 8A). In contrast, the conformation of the C-terminus of CALHM1 and presumably also

CALHM3 would likely interfere with their incorporation in a heteromeric assembly with any of

the other four subunits (Figure 8B). In this respect it is noteworthy that the same unit was

previously shown to be responsible for the smaller oligomeric assembly of CALHM1 as either

heptamer or octamer (Ren et al., 2022; Syrjanen et al., 2020). The fact that in case of the co-

expression of CALHM2 and CALHM4, the homomeric populations prevail, underlines the

energetic preference of homotypic interactions. Such prevalence was previously also observed

upon overexpression of different members of the unrelated LRRC8 family (Rutz et al., 2023),

where homomeric assemblies are believed not to be present in a cellular environment, pointing

towards a tightly regulated oligomerization process that might be disturbed upon overexpression.

However, in contrast to LRRC8A/C heteromers which, when expressed under equivalent

conditions, adopt a single oligomeric organization with a 2:1 ration of A to C subunits (Rutz et al.,

2023), we find a broad distribution of different assemblies in case of CALHM2/4 heteromers

(Figures 4 and 8C). For either protein family, the predominance of one of the paralogs and a

segregation of subunits within the oligomer provides additional evidence for the higher affinity of

homomeric interactions (Figure 8C).

Finally, we found a similar general prevalence of subunit conformations observed in homomers

also in heteromeric assemblies, except that in some cases the distribution of conformations is

influenced by their neighborhood. This is illustrated in the preference of CALHM2 subunits to

reside in a lifted 'up' conformation of TM1 if the subunit is located distant from CALHM4 subunits

whereas this conformation changes into a 'down' conformation in vicinity to CALHM4 subunits

(Figure 5 and 6A). The conformational preference of subunits is also extended to the shape of

residual density inside the pore, which exclusively around subunits in a 'down' conformation

shows the described bilayer-like features found in CALHM4 channels (Figure 6D and 8D). These

observations underline the general independence of subunits in assuming their conformation but

also indicate some coupling between neighbors. Thus, a change between conformations would not

necessarily have to proceed as a fully concerted transition.

Despite the observed diversity of subunit assemblies and pore conformations in CALHM2/4

heteromers, it is puzzling not to find detectable changes in their functional properties, as neither

of the constructs shows any sign of activity under conditions where CALHM1/3 heteromers

robustly conduct ions (Figure 1B-F, Figure 1-figure supplement 1), despite the fact that their larger

assembly should result in a wider pore diameter. The activation properties of these CALHM

paralogs have thus remained elusive and might require still unidentified interaction partners.

Alternatively, we also cannot exclude that these large oligomers might serve a function or substrate

preference that differs from classical large pore channels, which could be modified by the observed

formation of heteromers.

Methods

Expression constructs

CALHM coding sequences (synthesized by GeneScript) were inserted into a modified FX-cloning

compatible pcDNA1.3 vector (Drozdzyk et al., 2020; Geertsma, 2013; Geertsma and Dutzler,

2011). The expressed CALHM constructs contained at their C-terminus either a Venus, myc and

Streptavidin-Binding Peptide (Venus-myc-SBP) tag or a myc and His_6 (myc-His) tag. A 3C

protease cleavage sequence between the CALHM and tag sequences permitted removal of the c-

terminal fusion after purification. Sybody coding sequences were cloned into the FX-compatible

pSb_init vector, which contains a chloramphenicol-resistance gene and an arabinose-inducible

promotor (Zimmermann et al., 2020). The expressed sybody constructs contained an N-terminal

pelB leader sequence and a C-terminal His_6-tag. To obtain tag-free sybodies used in structural

characterization of heteromeric CALHM2/4 channels, sybody sequences were cloned into a vector

containing an N-terminal pelB signal sequence followed by a His_{10}-tag, maltose binding protein

and a 3C cleavage site (pBXNPHM3).

Expression in HEK 293 suspension cell culture

Human CALHM proteins were transiently expressed in suspension HEK293 $GnTI^-$ cells (ATCC

CRL-3022) following previously described protocols (Drozdzyk et al., 2020). HEK293 $GnTI^-$

cells were cultured in HyCell HyClone TransFx-H medium (Cytiva) supplemented with 1% fetal

bovine serum (FBS), 4 mM L-glutamine, 1.5% Poloxamer 188, and penicillin/streptomycin. The

cells were maintained at 37 °C in a humidified incubator at a 5% CO_2 atmosphere at 185 rpm. One

day before transfection, the cells were diluted to a density of 5 10^5 cells/ml and transfected with a

mixture of DNA and PEI. In co-transfections, a mixture of DNA constructs at equivalent molar

ratio was used. For 300 ml of cell culture, 0.5 mg of DNA and 1.2 mg of PEI MAX were each

suspended in 16 ml of DMEM medium. DNA solutions were mixed with the PEI solutions and

incubated for 15 minutes. To increase transfection efficiency, the mixture was supplemented with

4 mM valproic acid. The transfection mixtures were then added to cell cultures and expression was

carried out for 36 hours. After this period, the cells were pelleted by centrifugation at 500 g for 25

min, washed with PBS buffer and stored at -20 °C for further use.

CALHM protein purification

All protein purification steps were conducted at 4 °C. HEK293 cells expressing SBP-tagged

CALHM proteins for purification of homomeric channels or co-expressing SBP tagged CALHM2

and His_{10}-tagged CALHM4 for purification of heteromeric channels were harvested (typically

from 1 to 2 l of cell culture for purification of homomeric channels and from 9 l for purification of

heteromeric channels) and suspended in lysis buffer consisting of 25 mM HEPES (pH 7.6), 150

mM NaCl, 1% lauryl maltose-neopentyl glycol (LMNG), 0.5 mM $CaCl_2$, 2 mM $MgCl_2$, protease

inhibitors, RNase and DNase. After 1 hour, the lysate was clarified by centrifugation at 15,000 g

for 20 min. The clarified lysate was mixed with Strep-Tactin SuperFlow resin (1 ml of bed-resin

per 5 g of cell pellet) and incubated for 1 hour. Following, the resin was washed with 5 column

volumes (CV) of SEC buffer containing 10 mM HEPES (pH 7.6), 150 mM NaCl, 50 µM GDN, 2

mM CaCl₂. Bound proteins were eluted from the resin using 3 CV of SEC buffer supplemented

with 10 mM d-biotin. For the isolation of heteromeric CALHM2/4 channels, an extra affinity

chromatography step on Ni-NTA resin was performed after the initial purification on Strep-Tactin

resin. To this end, the eluate from the first purification step was incubated with NiNTA resin at a

1:6 v/v ratio and incubated for 1 hour under gentle agitation. Subsequently, the resin was washed

with 30 column volumes (CV) of SEC buffer, and bound proteins were eluted using the same

buffer supplemented with 300 mM imidazole. To remove fusion tags, eluates were incubated with

a 1:1 molar ratio of 3C protease for 30 min. Samples were then concentrated to 0.5 ml using an

Amicon Ultra centrifugal filter (100 kDa cutoff; Millipore), filtered through 0.22 µm filters and

subjected to size exclusion chromatography on a Superose 6 10/300 GL column equilibrated with

SEC buffer. Peak fractions were pooled, concentrated using a 100 kDa MWCO centrifugal filter

and immediately used for the preparation of cryo-EM grids.

Pulldown binding assays

For pull-downs of CALHM homologues, HEK293 GnTI⁻ cells were co-transfected using mixtures

of vectors encoding one CALHM paralog containing a fusion to Venus-myc-SBP and another

paralog containing a myc-His tag at equivalent molar ratios. All steps of the pulldown binding

assays were performed at 4 °C. Cells from a 300 ml culture were lysed in lysis buffer (25 mM

HEPES (pH 7.6), 150 mM NaCl, 1% LMNG, protease inhibitors, RNase, DNAse, 0.5 mM CaCl₂,

2 mM MgCl₂) for 1 hour and the lysates were cleared by centrifugation at 15,000 rcf for 10 min.

The supernatants were filtered through a 5 µm filter, mixed with 2 ml of Strep-Tactin SuperFlow

resin and incubated for 2 hours under gentle agitation. The resin was then washed with 5 CV of

wash buffer and bound proteins were eluted with elution buffer (wash buffer supplemented with

10 mM d-desthiobiotin, pH 7.6). For Western blot analysis, samples were loaded on a 4–20%

SDS-polyacrylamide gel and transferred to a polyvinylidene fluoride membrane by a semi-dry

blotting procedure. The membranes were first blocked at room temperature for 1 hour with 5%

non-fat milk in TBST buffer (50 mM Tris (pH 7.6), 150 mM NaCl, 0.075% Tween20) and then

incubated with anti-Myc primary antibodies overnight at 4 °C. The membranes were then washed

three times for 20 min with TBST and incubated with horse-radish peroxidase-conjugated goat

anti-mouse IgG antibody in 5% milk-TBST at room temperature for 1 hour. To remove unbound

primary antibodies, the membranes were washed with TBST buffer and subsequently blotted with

goat anti-rabbit-HRP conjugated secondary antibody for 2 hours at 4 °C. After blotting, the

membranes were washed with TBST buffer and developed with the Amersham ECL Prime

Western Blotting Detection kit (GE Healthcare).

Sybody selection

For sybody selection procedures, CALHM2 and CALHM4 proteins were chemically biotinylated

using amine-reactive EZ-Link™ NHS-PEG4-Biotin (ThermoFisher Scientific). For this purpose,

proteins were diluted to a concentration of 25 µM in SEC buffer and the coupling agent was added

at an 8-fold molar excess. Samples were incubated for 1 hour on ice and the reaction was quenched

by the addition of 5 mM Tris-HCl (pH 8.0). Excess biotin was removed from the biotinylated

protein on a PD-10 desalting column. The eluted proteins were concentrated, mixed with glycerol

to a final concentration of 10%, aliquoted, flash frozen in liquid N_2 and stored at -80 °C. Sybody

selection was performed as previously described (Zimmermann et al., 2020). Three synthetic

ribosome-display libraries, concave, loop , and convex libraries, and the plasmids, were kindly

provided by Prof. Markus Seeger. Sybodies were selected against chemically biotinylated

CALHM proteins. In brief, one round of ribosome display was performed, and the output RNA

was reverse transcribed and cloned into a phage-display compatible vector. The obtained plasmids

were transformed into *E. coli* SS320 cells to produce phages using a helper phage M13KO7. The

produced phages were then used in subsequent two rounds of phage display. Phagemids were

isolated from the output phages of the second round, and the sequences coding for sybodies were

subcloned into an pSB_init expression vector. The resulting plasmids were transformed into *E.*

coli MC1061, which were plated on agar plates. Single clones were expressed on a small scale and

used for binding tests by ELISA. Positive clones were sequenced.

Sybody expression and purification

DNA constructs of sybodies were transformed into *E. coli* MC1061 by a heat shock procedure and

grown in one liter of Terrific Broth medium supplemented with 25 µg/ml chloramphenicol at

37 °C. After initial incubation at 37 °C for 2 hours, the temperature was lowered to 22 °C and once

the cultures reached an OD_{600} of 0.5, L-arabinose was added to a final concentration of 0.02% to

induce expression. After 16-18 hours of expression, bacteria were harvested by centrifugation at

8,000 g for 20 min and pellets were either used immediately or stored at -80 °C until further use.

For purification, pellets were resuspended in TBS buffer (50 mM Tris (pH 7.6), 150 mM NaCl)

supplemented with 0.1 mg/ml lysozyme at a 1:30 w/v ratio. Bacteria were lysed by three freeze-

thaw cycles followed by sonication to release the periplasmic content. Lysates were centrifuged at

8,000 g for 30 min at 4 °C. The supernatants were collected and mixed with Ni-NTA resin and

incubated for 30 min. The resin was washed with TBS and sybodies were eluted with TBS

supplemented with 300 mM imidazole. For removal of the affinity tag, eluates were incubated

overnight with 3C protease while dialyzing with a MWCO of 5 kDa. The eluates were then

concentrated and filtered through a 0.22 μm centrifugal filter and subjected to size exclusion

chromatography on a Sepax SRT-10C SEC100 column equilibrated in TBS. Fractions of the

monomeric sybody peaks were collected and stored at -80 °C for further use.

Binding assays

Purified CALHM proteins were mixed with sybodies in a molar ratio of 1:2 to a final concentration

of 25 μM CALHM proteins. Prior to mixing, sybodies were supplemented with GDN to a final

concentration of 50 μM. Protein mixtures were applied to a Superose 6 5/150 Increase column

equilibrated with SEC buffer. Peak fractions corresponding to CALHM proteins were collected,

concentrated using Amicon Ultra 0.5 ml centrifugal filters (10 kDa cut-off, Ultracel) and analyzed

by SDS-PAGE. SPR measurements were performed at the Functional Genomic Centre Zurich

using a Biacore T200 instrument (GE Healthcare). The chemically biotinylated CALHM proteins

were immobilized on a streptavidin-coated sensor chip (XanTec), resulting in a maximum response

value of approximately 600 response units (RU). Measurements were performed at 10°C at a flow

rate of 30 μl/min. The buffer used during the experiments consisted of 10 mM HEPES at pH 7.6,

150 mM NaCl, 2 mM $CaCl_2$ and 0.006% GDN for CALHM2 or 0.01% LMNG for CALHM4.

Flow cell 1 was left empty to act as a reference cell for the measurements. Prior to measurements,

the system was equilibrated with the appropriate buffer for 2 hours. The sybodies were injected in

a single cycle kinetic measurement at concentrations of 7.8, 15.6, 31.25, 125 and 500 nM for

sybody SbC2 and 8, 24, 72, 216 and 648 nM for sybody SbC4. Binding data was analyzed using

BIAevaluation software (GE Healthcare) and fitted to a single-site binding model.

Whole-cell patch-clamp recordings

Adherent HEK293T cells were gently detached from their support and seeded in 10-cm dishes at

10% confluency. After a 3-hour incubation, cells were transfected with a mixture of 12 µg PEI and

3 µg of plasmids encoding CALHM constructs, which were C-terminally tagged with Venus-Myc-

SBP. For co-expression experiments, cells were transfected at a 1:1 ratio of the constructs at a total

amount of 3 µg of plasmids. Mock cells were transfected with a vector expressing Venus tag only.

Prior to transfection, DNA and PEI were separately incubated in 0.5 ml of DMEM medium. After

5 min, both solutions were mixed, incubated for 15 min and added to respective cell cultures.

Whole-cell currents were recorded 16-24 hours after transfection at room temperature (20-22 °C).

Recordings were conducted using borosilicate glass capillaries that were pulled and polished to

achieve a final resistance of 4-6 MΩ. Patches were held at -40 mV and the voltage was altered

between -100 mV and +100 mV in 10 mV increments for 500 ms by a step protocol. Cells were

perfused using a gravity-fed system. The solutions were as previously described (Syrjanen et al.,

2020). The pipette solution contained 147 mM NaCl, 10 mM EGTA, and 10 mM HEPES at pH

7.0, using NaOH for adjusting the pH. The bath solution contained 147 mM NaCl, 13 mM glucose,

10 mM HEPES (pH 7.3), using NaOH for adjusting the pH, 2 mM KCl, 2 mM $CaCl_2$, and 1 mM

$MgCl_2$. In Ca^{2+}-free experiments, the bath solution was similar but without addition of $CaCl_2$. Data

acquisition was performed using an Axopatch 200B amplifier and either Digidata 1440 or 1550

(Molecular Devices). Analog signals were digitized at a sampling rate of 10-20 kHz and filtered

at 5 kHz using the built-in 4-pole Bessel filter. Clampex 10.6 software (Molecular Devices) was

used for data acquisition and GraphPad Prism 8 for data analysis.

Cryo-EM sample preparation and data collection

Cryo-EM grids were prepared with freshly purified CALHM proteins. Prior to freezing, sybodies

were added to the CALHM samples at a 1:1.5 molar excess (based on the total amount of CALHM

subunits). In the sample containing heteromeric CALHM2/4 channels and both sybodies, SbC4

was added at a 1.5 and SbC2 at a 2.5 molar excess. The samples were frozen at different

concentrations of CALHM proteins (20 µM in the CALHM2/SbC2 sample, 47 µM in the

CALHM4/SbC4 sample, 38 µM in the CALHM2/4/SbC4 sample and 45 µM in the

CALHM2/4/SbC2/SbC4 sample). Samples were applied to glow-discharged holey carbon grids

(Quantifoil R1.2/1.3 Au 200 mesh). Excess liquid was removed in a controlled environment (4 °C

and 100% relative humidity) by blotting grids for 2–4 s. Grids were subsequently flash-frozen in

a liquid propane-ethane mix using a Vitrobot Mark IV (Thermo Fisher Scientific) and stored in

liquid nitrogen. Samples were imaged with a 300 kV Titan Krios G3i with a 100 µm objective

aperture and data were collected using a post-column BioQuantum energy filter with a 20 eV slit

and a K3 direct electron detector operating in super-resolution counted mode. Dose-fractionated

micrographs were recorded with a defocus range of -1 µm to -2.4 µm in an automated mode using

EPU 2.9. Datasets were recorded at a nominal magnification of 130,000 corresponding to a pixel

size of 0.651 Å per pixel (0.3255 Å per pixel in super-resolution mode).

Cryo-EM image processing

Micrographs from all datasets were pre-processed in a similar manner. Micrographs of the CALHM2-/SbC2, CALHM2/4/SbC4 and CALHM2/4/SbC2/SbC4 datasets were binned prior to processing resulting in a pixel size of 0.651Å. The raw movies were used to correct for beam-induced motion using patch motion correction in cryoSPARC v.3.2.0 and v.4.0.3, followed by patch CTF estimation (Punjani et al., 2017). For the CALHM4/SbC4 dataset the raw movies were used for correction of the beam-induced movement using a dose-weighting scheme in RELION's own implementation of the MotionCor2 algorithm available in version 3.0 (Zivanov et al., 2018) and the CTF parameters were estimated on summed movie frames using CTFFIND4.1 (Rohou and Grigorieff, 2015). Poor quality images with significant drift, ice contamination or poor CTF estimates were discarded.

The dataset of CALHM4/SbC4 was processed entirely in RELION-3.1 (Scheres, 2012). Particles were first selected by Laplacian-of-Gaussian based auto-picking and subjected to a first round of 2D classification to generate 2D average templates. Next, four 2D classes with protein-like features were used as references for a second template-based picking. The particles were extracted with a box size of 720 pixels and downscaled four times. A second 2D classification was performed and the particles from the five best classes were used for the generation of an initial 3D model, imposing C10 symmetry. The extracted particles from the second 2D classification round were subjected to a round of 3D classification with C10 symmetry imposed, using the initial model low pass filtered to 60 Å as a reference map. The best 3D class was selected and the particles belonging to this class were unbinned to their original pixel size and subjected to 3D refinement with D10 symmetry imposed. Additionally, CTF and aberration refinement and Bayesian polishing was applied (Rohou and Grigorieff, 2015; Zivanov et al., 2019). The map was sharpened with an

isotropic b-factor of -153 Å^2, resulting in a global resolution of 3.7 Å. The local resolution was

estimated using RELION (Zivanov et al., 2018).

The dataset of CALHM2/SbC2 was processed entirely in cryoSPARC v.3.2.0 and v.4.0.3 (Punjani

et al., 2017). Particles were initially picked with the blob picker and subjected to 2D classification

to generate 2D average templates for more accurate template-based particle picking. Particles were

selected and extracted with a box size of 512 pixels and downscaled twice. Selected particles were

subjected to four rounds of 2D classification to remove false positives and low-quality particles

and subjected to an *ab initio* reconstruction with three classes. The obtained classes included

decameric, undecameric and ambiguous assemblies of CALHM2. Particles belonging to classes

showing an undecameric assembly were further sorted by a round of heterogeneous refinement

using three *ab initio* reconstructions as references. The same procedure was used for particles

belonging to classes showing a decameric assembly, which yielded a low-quality reconstruction.

Therefore, further processing was restricted to the more promising undecameric assemblies. The

quality of the reconstruction was further improved by CTF global and local refinements and non-

uniform refinement with C11 symmetry imposed, yielding a map at 3.07 Å. The local resolution

was estimated with cryoSPARC. To improve the resolution of the CALHM-sybody interface

region, a local refinement with C11 symmetry imposed was performed in the presence of a tight

mask restricted to the region surrounding the sybody binding site, yielding a reconstruction of this

region at a global resolution of 2.8 Å.

The datasets of CALHM2/4 in complex with sybodies were processed in cryoSPARC v.4.0.3

(Punjani et al., 2017), unless indicated otherwise. Particles from the CALHM2/4 datasets

containing either both sybodies (CALHM2/4/SbC2/SbC4) or SbC4 alone (CALHM2/4/SbC4)

were initially picked with the blob picker and subjected to 2D classification to generate 2D

templates for more accurate template-based particle picking. Particles were selected and extracted

with a box size of 512 pixels, downscaled twice and subjected to multiple rounds of 2D

classification to remove false positives and poor-quality particles. Visual inspection of the 2D

classes indicated the presence of different assemblies. To separate these assemblies, particles were

exported to RELION (version 4.0) (Kimanius et al., 2021) and used to generate initial models.

Subsequently, particles were subjected to several rounds of 3D classification using the initial

models as references. In the first two rounds of classification, low quality particles were excluded

and only particles belonging to the high-quality classes were subjected to a final 3D classification

with 10 distinct classes. To obtain high resolution reconstructions of complexes, particles

belonging to 2D classes of channel dimers were excluded and only particles belonging to high

quality 2D classes with single channels were used to generate *ab initio* reconstructions in

cryoSPARC. This yielded decoy reconstructions and undecameric reconstructions. Particles of

undecameric reconstructions were used as input for heterogeneous refinements with three

references including a decoy class and two proper classes. This approach yielded reconstructions

of undecameric and dodecameric assemblies. The heterogeneous refinement was repeated four

times using particles belonging to the undecameric assembly from the previous refinement as input

and undecameric and dodecameric reconstructions as references. Resulting undecameric

reconstructions were further improved by local and global CTF, and non-uniform refinements in

C1. . These approaches yielded maps with global resolutions of 3.8Å and 3.3Å for the

CALHM2/4/SbC4 and CALHM2/4/SbC2/SbC4 complexes, respectively. The local resolution of

maps was estimated within cryoSPARC.

Model building and Refinement

Models were based on previously determined structures of CALHM2 (PDB:6UIW) (Choi et al.,

2019) and CALHM4 (PDB:6YTL) (Drozdzyk et al., 2020). Sybodies were initially modelled

based on high-resolution structures (PDB:1ZVH for SbC2 and PDB:3K1K for SbC4). Models

were first fitted into their corresponding densities using ChimeraX (Pettersen et al., 2021). The

cryo-EM density of CALHM2/SbC2 allowed us to assign residues 39-308 and 314-323 of

CALHM2 and 324-329 of the remaining sequence of the expression tag, and the entire model of

SbC2 whose CDRs interacting with CALHM2 were rebuilt into the cryo-EM density. The cryo-

EM density of CALHM4/SbC4 was of sufficiently high resolution to unambiguously assign

residues 5–82 and 94–276 of CALHM4. In contrast, the limited quality of the density of SbC4 did

not permit a complete interpretation of its atomic model and thus a truncated version lacking the

variable regions was placed into the density.

The model of the heteromeric CALHM2/4 in complex with sybodies SbC2 and SbC4 was built by

placing eight CALHM2 and three CALHM4 subunits into the cryo-EM density obtained from the

CALHM2/4/SbC2/SbC4 dataset at 3.3 Å. Coordinates of CALHM subunits and sybodies were

obtained from the structures of the homomeric CALHM2/SbC2 and CALHM4/SbC4 complexes.

Residues 19-46 of CALHM2 in the downward conformation were built de novo. The subunit

displaying a mix of CALHM2 and 4 (i.e. position 10) was interpreted by a CALHM2 subunit

where loop regions not defined in the density were removed. The structure of the

CALHM2/4/SbC4 complex at 3.8 Å was built based on the model of CALHM2/4/SbC2/SbC4 after

removal of the SbC2 subunits, except that it contains a truncated version of CALHM2 also in

position 11. Refinement and validation of all models was performed in PHENIX (Afonine et al.,

2018). Figures containing molecular structures and densities were prepared with DINO

(http://www.dino3d.org), PyMOL (DeLano, 2002) and ChimeraX (Pettersen et al., 2021).

Interaction interfaces were calculated with CNS_SOLVE (Brunger, 2007).

Acknowledgments This research was supported by a grant from the Swiss National Science

Foundation (No. 310030B_182828). Cryo-EM data was collected at the Center for Microscopy

and Image Analysis (ZMB) of the University of Zurich. We thank Markus Seeger for providing

access to the sybody libraries. Jens Sobek and Marton Liciczai are acknowledged for help with

surface plasmon resonance experiments recorded at the Functional Genomics Center of UZH/ETH

Zurich. All members of the Dutzler lab are acknowledged for their help at various stages of the

project.

Competing interests

The authors declare no competing interests.

Accession codes

The cryo-EM density maps have been deposited in the Electron Microscopy Data Bank under

following ID codes: EMD-19365 (CALHM4/SbC4), EMD-19362 (CALHM2/SbC2), EMD-

19363 (CALHM2/CALHM4/SbC4), EMD-19364 (CALHM2/CALHM4/SbC2/SbC4). The

coordinates of the corresponding atomic models of have been deposited in the Protein Data Bank

under ID codes 8RMN (CALHM4/SbC4), 8RMK (CALHM2/SbC2), 8RML

(CALHM2/CALHM4/SbC4), 8RMM (CALHM2/CALHM4/SbC2/SbC4).

References

Afonine, P.V., B.K. Poon, R.J. Read, O.V. Sobolev, T.C. Terwilliger, A. Urzhumtsev, and P.D. Adams. 2018. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol*. 74:531-544.

Baumann, S.W., R. Baur, and E. Sigel. 2001. Subunit arrangement of gamma-aminobutyric acid type A receptors. *J Biol Chem*. 276:36275-36280.

Brunger, A.T. 2007. Version 1.2 of the Crystallography and NMR system. *Nat Protoc*. 2:2728-2733.

Choi, W., N. Clemente, W. Sun, J. Du, and W. Lu. 2019. The structures and gating mechanism of human calcium homeostasis modulator 2. *Nature*. 576:163-167.

Danielli, S., Z. Ma, E. Pantazi, A. Kumar, B. Demarco, F.A. Fischer, U. Paudel, J. Weissenrieder, R.J. Lee, S. Joyce, J.K. Foskett, and J.S. Bezbradica. 2023. The ion channel CALHM6 controls bacterial infection-induced cellular cross-talk at the immunological synapse. *EMBO J*. 42:e111450.

DeLano, W.L. 2002. Pymol: An open-source molecular graphics tool. *CCP4 Newsletter On Protein Crystallography*. 40:82-92.

Demura, K., T. Kusakizako, W. Shihoya, M. Hiraizumi, K. Nomura, H. Shimada, K. Yamashita, T. Nishizawa, A. Taruno, and O. Nureki. 2020. Cryo-EM structures of calcium homeostasis modulator channels in diverse oligomeric assemblies. *Sci Adv*. 6:eaba8105.

Drozdzyk, K., M. Sawicka, M.I. Bahamonde-Santos, Z. Jonas, D. Deneka, C. Albrecht, and R. Dutzler. 2020. Cryo-EM structures and functional properties of CALHM channels of the human placenta. *Elife*. 9.

Foskett, J.K. 2020. Structures of CALHM channels revealed. *Nat Struct Mol Biol*. 27:227-228.

Geertsma, E.R. 2013. FX cloning: a versatile high-throughput cloning system for characterization

of enzyme variants. *Methods Mol Biol*. 978:133-148.

Geertsma, E.R., and R. Dutzler. 2011. A versatile and efficient high-throughput cloning tool for

structural biology. *Biochemistry*. 50:3272-3278.

Greger, I.H., and M.L. Mayer. 2019. Structural biology of glutamate receptor ion channels:

towards an understanding of mechanism. *Curr Opin Struct Biol*. 57:185-195.

Kimanius, D., L. Dong, G. Sharov, T. Nakane, and S.H.W. Scheres. 2021. New tools for automated

cryo-EM single-particle analysis in RELION-4.0. *Biochemical Journal*. 478:4169-4185.

Liao, Y., Y. Wang, Q.-Q. Tao, C. Yang, J. Wang, J. Cheng, J. Ma, Z.-Y. Wu, R.-Y. Pan, and Z.

Yuan. 2023. CALHM2 V136G polymorphism reduces astrocytic ATP release and is

associated with depressive symptoms and Alzheimer's disease risk. *Alzheimer's &*

Dementia. 19:4407-4420.

Liu, J., F. Wan, Q. Jin, X. Li, E.A. Bhat, J. Guo, M. Lei, F. Guan, J. Wu, and S. Ye. 2020. Cryo-

EM structures of human calcium homeostasis modulator 5. *Cell Discov*. 6:81.

Ma, J., X. Qi, C. Yang, R. Pan, S. Wang, J. Wu, L. Huang, H. Chen, J. Cheng, R. Wu, Y. Liao, L.

Mao, F.C. Wang, Z. Wu, J.X. An, Y. Wang, X. Zhang, C. Zhang, and Z. Yuan. 2018a.

Calhm2 governs astrocytic ATP releasing in the development of depression-like behaviors.

Mol Psychiatry. 23:883-891.

Ma, Z., A.P. Siebert, K.H. Cheung, R.J. Lee, B. Johnson, A.S. Cohen, V. Vingtdeux, P.

Marambaud, and J.K. Foskett. 2012. Calcium homeostasis modulator 1 (CALHM1) is the

pore-forming subunit of an ion channel that mediates extracellular Ca2+ regulation of

neuronal excitability. *Proc Natl Acad Sci U S A*. 109:E1963-1971.

Ma, Z., J.E. Tanis, A. Taruno, and J.K. Foskett. 2016. Calcium homeostasis modulator (CALHM)

748   ion channels. *Pflugers Arch*. 468:395-403.

749   Ma, Z., A. Taruno, M. Ohmoto, M. Jyotaki, J.C. Lim, H. Miyazaki, N. Niisato, Y. Marunaka, R.J.

750   Lee, H. Hoff, R. Payne, A. Demuro, I. Parker, C.H. Mitchell, J. Henao-Mejia, J.E. Tanis,

751   I. Matsumoto, M.G. Tordoff, and J.K. Foskett. 2018b. CALHM3 Is Essential for Rapid Ion

752   Channel-Mediated Purinergic Neurotransmission of GPCR-Mediated Tastes. *Neuron*.

753   98:547-561 e510.

754   Pettersen, E.F., T.D. Goddard, C.C. Huang, E.C. Meng, G.S. Couch, T.I. Croll, J.H. Morris, and

755   T.E. Ferrin. 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and

756   developers. *Protein Science*. 30:70-82.

757   Punjani, A., J.L. Rubinstein, D.J. Fleet, and M.A. Brubaker. 2017. cryoSPARC: algorithms for

758   rapid unsupervised cryo-EM structure determination. *Nature Methods*. 14:290-296.

759   Ren, Y., Y. Li, Y. Wang, T. Wen, X. Lu, S. Chang, X. Zhang, Y. Shen, and X. Yang. 2022. Cryo-

760   EM structure of the heptameric calcium homeostasis modulator 1 channel. *J Biol Chem*.

761   298:101838.

762   Ren, Y., T. Wen, Z. Xi, S. Li, J. Lu, X. Zhang, X. Yang, and Y. Shen. 2020. Cryo-EM structure

763   of the calcium homeostasis modulator 1 channel. *Sci Adv*. 6:eaba8161.

764   Rohou, A., and N. Grigorieff. 2015. CTFFIND4: Fast and accurate defocus estimation from

765   electron micrographs. *Journal of structural biology*. 192:216-221.

766   Rutz, S., D. Deneka, A. Dittmann, M. Sawicka, and R. Dutzler. 2023. Structure of a volume-

767   regulated heteromeric LRRC8A/C channel. *Nat Struct Mol Biol*. 30:52-61.

768   Scheres, S.H. 2012. RELION: implementation of a Bayesian approach to cryo-EM structure

769   determination. *Journal of structural biology*. 180:519-530.

Siebert, A.P., Z. Ma, J.D. Grevet, A. Demuro, I. Parker, and J.K. Foskett. 2013. Structural and

functional similarities of calcium homeostasis modulator 1 (CALHM1) ion channel with

connexins, pannexins, and innexins. *J Biol Chem*. 288:6140-6153.

Syrjanen, J., K. Michalski, T. Kawate, and H. Furukawa. 2021. On the molecular nature of large-

pore channels. *J Mol Biol*. 433:166994.

Syrjänen, J.L., M. Epstein, R. Gómez, and H. Furukawa. 2023. Structure of human CALHM1

reveals key locations for channel regulation and blockade by ruthenium red. *Nature*

Communications. 14:3821.

Syrjanen, J.L., K. Michalski, T.H. Chou, T. Grant, S. Rao, N. Simorowski, S.J. Tucker, N.

Grigorieff, and H. Furukawa. 2020. Structure and assembly of calcium homeostasis

modulator proteins. *Nat Struct Mol Biol*.

Taruno, A., I. Matsumoto, Z. Ma, P. Marambaud, and J.K. Foskett. 2013a. How do taste cells

lacking synapses mediate neurotransmission? CALHM1, a voltage-gated ATP channel.

Bioessays. 35:1111-1118.

Taruno, A., V. Vingtdeux, M. Ohmoto, Z. Ma, G. Dvoryanchikov, A. Li, L. Adrien, H. Zhao, S.

Leung, M. Abernethy, J. Koppel, P. Davies, M.M. Civan, N. Chaudhari, I. Matsumoto, G.

Hellekant, M.G. Tordoff, P. Marambaud, and J.K. Foskett. 2013b. CALHM1 ion channel

mediates purinergic neurotransmission of sweet, bitter and umami tastes. *Nature*. 495:223-

226.

Voss, F.K., F. Ullrich, J. Munch, K. Lazarow, D. Lutter, N. Mah, M.A. Andrade-Navarro, J.P. von

Kries, T. Stauber, and T.J. Jentsch. 2014. Identification of LRRC8 heteromers as an

essential component of the volume-regulated anion channel VRAC. *Science*. 344:634-638.

Yang, W., Y. Wang, J. Guo, L. He, Y. Zhou, H. Zheng, Z. Liu, P. Zhu, and X.C. Zhang. 2020.

Cryo-electron microscopy structure of CLHM1 ion channel from Caenorhabditis elegans.

Protein Science. 29:1803-1815.

Zimmermann, I., P. Egloff, C.A.J. Hutter, B.T. Kuhn, P. Brauer, S. Newstead, R.J.P. Dawson, E.R.

Geertsma, and M.A. Seeger. 2020. Generation of synthetic nanobodies against delicate

proteins. *Nat Protoc*. 15:1707-1741.

Zivanov, J., T. Nakane, B.O. Forsberg, D. Kimanius, W.J. Hagen, E. Lindahl, and S.H. Scheres.

2018. New tools for automated high-resolution cryo-EM structure determination in

RELION-3. *Elife*. 7.

Zivanov, J., T. Nakane, and S.H.W. Scheres. 2019. A Bayesian approach to beam-induced motion

correction in cryo-EM single-particle analysis. *IUCrJ*. 6:5-17.

Figures and figure legends

Figure 1. Heteromerization and functional properties of placental CALHM paralogs. (**A**) Western

blot (detecting the myc-tag attached to the C-termini of respective constructs) of CALHM subunits

expressed upon co-transfection of HEK293 GnTI⁻ cells with pairs of CALHM subunits. CALHM

channels were isolated by affinity purification of one subunit containing a fusion to Venus and an

SBP-tag (bait, shown at higher molecular weight). The resulting samples contain a mix of

homomers of the purified subunit and heteromers (with the second subunit not containing Venus

and SBP tags, prey at lower molecular weight). The ratio of pray/bait is displayed based on the

integration of the intensity of the displayed Western blots. The experiment has been carried out

once. (**B-E**) Represented patch-clamp electrophysiology recordings (whole cell configuration) of

indicated CALHM subunits expressed in HEK-293 cells measured in buffers containing 2 mM

Ca^{2+} on the extracellular side. (**B**) Comparison of currents from mock transfected cells (left) and

cells expressing CALHM1 (right). Inset shows voltage protocol. (**C**) Current response from cells

co-transfected with DNA coding for CALHM1 and CALHM3 subunits. (**D**) Current response

from cells transfected with CALHM2 (left) or CALHM4 (right) subunits. (**E**) Current response

from cells co-transfected with CALHM2 and CALHM4 subunits. (**F**) Mean current density of

recordings of different CALHM constructs (measured at 100 mV, 190 ms after the voltage step),

errors are s.e.m., values from individual recordings are shown as circles.

Figure supplement 1. Electrophysiology data of CALHM paralogs at low extracellular Ca^{2+}.

Figure 1—figure supplement 1. Electrophysiology data of CALHM paralogs recorded at low

extracellular Ca^{2+}. (**A-D**) Patch clamp electrophysiology recordings (whole cell configuration) of

representative traces of indicated CALHM subunits expressed in HEK-293 cells measured in

buffers not containing Ca^{2+} on the extracellular side. (**A**) Comparison of currents from mock

transfected cells (left) and cells expressing CALHM1 (right). Inset shows voltage protocol. (**B**)

Current response from cells co-transfected with DNA coding for CALHM1 and CALHM3

subunits. (**C**) Current response from cells transfected with CALHM2 (left) or CALHM4 (right)

subunits. (**D**) Current response from cells co-transfected with CALHM2 and CALHM4 subunits.

(**E**) Mean current density of recordings of different CALHM constructs (measured at 100 mV, 400

ms after voltage step), errors are s.e.m., values from individual recordings are shown as circles.

Figure 2. Structural characterization of the CALHM4/SbC4 complex. (**A**) Cryo-EM density at

3.7 Å (after application of D10 symmetry) of a pair of decameric CALHM4/SbC4 complexes

interacting via their cytoplasmic regions. (**B**) Ribbon representation of the same complex viewed

parallel (left) and perpendicular (right) to the membrane plane. Membrane boundaries are indicated

(left). **A**, **B**, CALHM4 subunits are colored in red and green, SbC4 in yellow. (**C**) Cryo-EM density

(left) and ribbon representation (right) of a single subunit of the CALHM4/SbC4 complex.

Transmembrane helices are labeled and shown in unique colors. (**D**) Organization of TM helices

in the decameric channel. The view is from the extracellular side, the coloring is as in C. (**E**) Slab

844 through a low-pass filtered map of the CALHM4/SbC4 complex (viewed from within the

845 membrane) shows bilayer-like density in the lumen of the wide pore.

847 **Figure supplement 1.** Sybody selection and characterization.

848 **Figure supplement 2.** Cryo-EM reconstruction of CALHM4 in complex with sybody SbC4.

Figure 2—figure supplement 1. Sybody selection and characterization. (**A**) Sequence alignment

of the two sybodies SbC4 and SbC2 specifically targeting CALHM4 and CALHM2, respectively.

(**B**) Size exclusion chromatography profile of the CALHM4/SbC4 complex. The elution peak of

the equivalent amount of CALHM4 is displayed as dashed line in comparison. Inset (right) shows

SDS-PAGE gel of the peak fraction. SPR experiments of SbC4 binding to immobilized CALHM4

illustrating specific binding (**C**) and to CALHM2 not showing specific interactions (**D**). Dashed

line in D corresponds to a fit to a 1:1 binding model yielding a K_d of 100 nM. (**E**) Size exclusion chromatography profile of the CALHM2/SbC2 complex. The elution peak of the equivalent amount of CALHM2 is displayed as dashed line in comparison. Inset (right) shows SDS-PAGE gel of the peak fraction. SPR experiments of SbC2 binding to immobilized CALHM2 illustrating specific binding (**F**) and CALHM4 not showing specific interactions (**G**). B-G the experiments werw carried out once.

Figure 2—figure supplement 2. Cryo-EM reconstruction of CALHM4 in complex with sybody

SbC4. (**A**) Representative cryo-EM micrograph acquired with a Titan Krios microscope equipped

with a K3 camera. (**B**) Representative 2D class averages of the CALHM4/Sbc4 complex. (**C**) Data

processing workflow. Particles were picked and subjected to 2D classification. High resolution 2D

class averages were used as input for initial 3D model generation and subsequent 3D classification.

The 3D classification allowed to isolate CALHM4-SbC4 particles, which were further refined with

iterative polishing, and 3D and CTF refinements. The distribution of particles (%) of each class is

indicated. (**D**) FSC plot of the final refined undecameric unmasked (blue), masked (green),

randomized (orange) and corrected for mask convolution effects (purple) cryo-EM density map of

the CALHM4/SbC4 complex. The resolution at which the FSC curve drops below the 0.143

threshold is indicated. (**E**) Final 3D reconstruction of CALHM4/SbC4 complex colored according

to local resolution. The angular distribution of the particles included in the final reconstruction is

indicated. Each cylinder represents one view and the height corresponds to the number of particles

in that view.

Figure 3. Structural characterization of a CALHM2/SbC2 complex. (**A**) Cryo-EM density of the

undecameric CALHM2/SbC2 complex at 3.1 Å (after application of C11 symmetry). (**B**) Ribbon

representation of the same complex viewed parallel to the membrane plane (left) and from the

extracellular side (right). Membrane boundaries are indicated (left). Space filling model shows a

bound lipid. **A**, **B**, CALHM2 subunits are colored in yellow and blue, SbC2 in green, the bound

lipid in magenta. (**C**) Cryo-EM density (left) and ribbon representation (right) of a single subunit

of the CALHM2/SbC2 complex. Transmembrane helices are labeled and shown in unique colors.

(**D**) Subunit (with orientation relative to C indicated) and blow-up of the sybody-interaction region.

The C-terminus of the adjacent subunits contributing to the interaction epitope is colored in yellow

and labeled (CT^{+1}). (**E**) Slab through a low-pass filtered map of the CALHM2/SbC2 complex

(viewed from within the membrane) does not show bilayer-like density in the lumen of the pore.

890   **Figure supplement 1.** Cryo-EM reconstruction of CALHM2 in complex with sybody SbC2.

Figure 3—figure supplement 1. Cryo-EM reconstruction of CALHM2 in complex with sybody SbC2. (**A**) Representative cryo-EM micrograph acquired with a Titan Krios microscope equipped with an energy filter and a K3 camera. (**B**) 2D class averages of CALHM2-SbC2 complex. (**C**) Data processing workflow. Particles were picked and subjected to 2D classification. High resolution 2D class averages were used as an input for ab-initio reconstructions, which allowed to isolate two populations of particles (containing decameric and undecameric assemblies) from a decoy class. The distribution of particles (%) of each class is indicated. Particles were further sorted by heterogenous refinement and undecameric assemblies were improved in several rounds

of iterative non-uniform and CTF refinement. To increase the resolution in the sybody binding site,

a local refinement of the masked interaction region was performed. (**D**) FSC plot of the final

refined cryo-EM density map of undecameric channels (unmasked (blue), loose (green) and tight

(red) mask, and corrected for mask convolution effects (purple)) of the CALHM2/SbC2 complex.

The resolution at which the FSC curve drops below the 0.143 threshold is indicated. (**E**) Final 3D

reconstruction of the CALHM2/sybody complex colored according to local resolution.

Figure 4. Classification of CALHM2/CALHM4 heteromers. Classes correspond to distinct

populations of oligomers in the CALHM2/4/SbC2/SbC4 complex. (**A**) Assemblies containing an

excess of CALHM4 subunits. Shown is the well-resolved halve of interacting channel pairs (with

weak density corresponding to the second channel). (**B**, **C**) Assemblies with an excess of

CALHM2 subunits. (**C**) Dodecameric channels containing three and four CALHM4 subunits. (**D**)

Undecameric channels containing two to four CALHM4 subunits. **A-C**, Shown are low resolution

maps (processed in C1) with bound SbC2 and SbC4 colored in green and yellow , respectively.

Numbering refers to CALHM4 subunits. The relationship between views is indicated. Top panels

show inclined view towards the membrane plane, bottom panels a view from the extracellular side.

Figure supplement 1. Cryo-EM reconstruction of CALHM2/4 in complex with sybody SbC4.

Figure supplement 2. Cryo-EM reconstruction of CALHM2/4 in complex with sybodies SbC2

and SbC4.

Figure 4—figure supplement 1. Cryo-EM reconstruction of CALHM2/4 in complex with sybody

SbC4. (**A**) Representative cryo-EM micrograph acquired with a Titan Krios microscope equipped

with a K3 camera. (**B**) Data processing workflow. Particles were picked and subjected to 2D

classification. 2D class averages allowed for the initial separation of populations of single

channels and their dimers. The distribution of particles (%) of each class is indicated. Further 3D

classification has led to the separation of different assemblies. Ab-initio reconstructions of particles

featuring single channels resulted in low resolution maps of undecameric assemblies. Further steps

of heterogenous refinement have separated undecamers and dodecamers. The smaller channels

were refined to high resolution via several cycles of hetero- and non-uniform refinement. (**C**) FSC

plot of the final refined unmasked (blue), masked (green), randomized (orange) and corrected for

mask convolution effects (purple) cryo-EM density maps of an undecameric CALHM4/SbC4

complex. (**D**) Final 3D reconstruction of a CALHM2/4/SbC4 complex colored according to local

resolution.

Figure 4—figure supplement 2. Cryo-EM reconstruction of CALHM2/4 in complex with sybodies SbC2 and SbC4. (**A**) Representative cryo-EM micrograph acquired with a Titan Krios microscope equipped with a K3 camera. (**B**) Data processing workflow. Particles were picked and subjected to 2D classification. High resolution 2D class averages revealed populations of single channels and dimers of channels. The distribution of particles (%) of each class is indicated. Subsequent 3D classification of all the particles allowed for the separation of different assemblies. Ab-initio reconstructions of particles featuring single channels yielded undecameric assemblies. The subsequent heterogenous refinement separated undecameric and dodecameric channels. The undecameric assemblies were further improved to high resolution by hetero, CTF and non-uniform refinement. (**C**) Fourier shell correlation (FSC) curves of the final refined unmasked (blue), masked (green), randomized (orange) and corrected for mask convolution effects (purple) cryo-EM density maps of an undecameric CALHM2/4/SbC2/SbC4 complex. (**D**) Final 3D reconstruction of the CALHM2/4/SbC2/SbC4 complex colored according to local resolution. Shown is the map of the channel viewed from the side (left) and the slice through the channel (right).

Figure 5. CALHM2/CALHM4 heteromer structure. (**A**) Schematic depiction of the subunit distribution in CALHM2/4 heteromers. 4 refers to CALHM4, 2 to CALHM2 in 'up' conformation, 2d to CALHM2 in 'down' conformation, 2u/d to a mixed population of both conformations of CALHM2 subunits, 4/2 and 2/4 to mixed populations of CALHM2 and 4 with the first number indicating the predominant subunit. (**B**) Cryo-EM density of the CALHM2/4/SbC4 complex at 3.8 Å viewed with an inclination towards the membrane plane. (**C**) Ribbon representation of the CALHM2/4/SbC4 complex viewed from the same perspective as in B (left) and from the extracellular side (right). (**D**) Cryo-EM density of the CALHM2/4/SbC2/SbC4 complex at 3.3 Å

959   viewed with an inclination towards the membrane plane (top) and from the intracellular side

960   (bottom). (**C**) Ribbon representation of the CALHM2/4/SbC2/SbC4 complex viewed from the

961   same perspective as in B (left, top) and from the extracellular side (right). **B-E**, Colors and numbers

962   of channel subunits are as in **A**. SbC2 and SbC4 are colored in yellow and green, respectively.

Figure 6. Conformational properties of CALHM subunits in CALHM2/4 heteromers. (**A**) Cryo-EM density of CALHM2 subunits in 'down' (position 3, left) and 'up' (position 6, top) and of CALHM4 (in 'down' conformation, position 1, bottom). (**B**) Ribbon representation of a superposition of CALHM2 subunits in 'up' and 'down' conformations. Inset (right) shows blowup of the intracellular region around TM1-3. (**C**) Molecular surface of the intracellular region relating TM3 and TM4 viewed from within the membrane plane. A gap between both helices (left, asterisk), which is occupied by a lipid (space-filling representation) in the 'up' conformation (center), is closed in the 'down' conformation (right). **A**-**C** Transmembrane helices and attached sybodies are shown in unique colors. (**D**) Low-pass filtered map of the CALHM2/4/SbC2/SbC4 complex depicting residual bilayer-like density in the pore lumen. Slice through the channel in a view parallel to the membrane (left), and from the extracellular side (right). Subunit conformations are indicated (d, 'down', u, 'up'). CALHM4 subunits are shown in red and magenta, CALHM2 subunits in blue and cyan.

Figure 7. Structural features of subunit interfaces. (**A-C**) Interaction interfaces between pairs of

CALHM2 subunits (**A**), CALHM4 subunits (**B**) and a CALHM2 and a CALHM4 subunit (**C**).

Shown are ribbon representations of neighboring subunits viewed from the inside of the channel

parallel to the membrane and their molecular surfaces that are buried in the interaction interface.

(**D-F**) Subunit interactions in the cytoplasmic domain of CALHM channels. Shown is a central

subunit (x) and its interaction partners in clockwise and counterclockwise position (top). The

region of the intracellular domain that is contacted by neighboring subunits is mapped on the

986   molecular surface of the central subunit. Center, view from the extracellular side (out), bottom

987   from the cytoplasm (in). **A-F** The relative position of subunits within the assembly is indicated.

988   Paralogs are represented in unique colors.

Figure 8. Discussion. (**A**, **B**) Superposition of single subunits of different CALHM paralogs on CALHM2 viewed from within the membrane (top) and of their cytosolic domain viewed from the outside (bottom). (**A**) Superposition with CALHM2-like subunits CALHM4-6. (**B**) Superposition with CALHM1. The difference in the conformation of the intracellular domain is apparent. (**C**) Segregation of subunits in heteromeric channels of the LRRC8 family (left) and of CALHM2/4 channels (right). Proteins are shown as space-filling models. The view is from the extracellular

996   side. (**D**) Schematic depiction of how the conformation of CALHM subunits influences the

997   properties of the bilayer-like density inside the channel. The conformation of helices is indicated.

998　**Table 1.** Cryo-EM Data collection, refinement and validation statistics.

	CALHM4/SbC4	CALHM2/SbC2	CALHM2/CALHM4/SbC4	CALHM2/CALHM4/SbC2/SbC4
	EMD-19365 PDB 8RMN	EMD-19362 PDB 8RMK	EMD-19363 PDB 8RML	EMD-19364 PDB 8RMM
Data collection and processing				
Microscope	FEI Titan Krios G3i	FEI Titan Krios G3i	FEI Titan Krios G3i	FEI Titan Krios G3i
Camera	Gatan K3+GIF	Gatan K3+GIF	Gatan K3+GIF	Gatan K3+GIF
Magnification	130,000	130,000	130,000	130,000
Voltage (kV)	300	300	300	300
Electron exposure (e$^-$/Å^2)	70	72	60	60
Defocus range (μm)	-1.0 to -2.4	-1.0 to -2.4	-1.0 to -2.4	-1.0 to -2.4
Pixel size* (Å)	0.651 (0.3255)	0.651 (0.3255)	0.651 (0.3255)	0.651 (0.3255)
Initial particle images (no.)	424,942	1,068,223	1,345,836	2,800,265
Final particle images (no.)	52,248	37,894	131,652	93,191
Symmetry imposed	D10	C11	C1	C1
Map resolution (Å) FSC threshold 0.143	3.7	3.07	3.84	3.26
Map resolution range (Å)	3.5-5	2.6-5	3.5-6	3-6
Refinement				
Model resolution (Å) FSC threshold 0.5	3.9	3.2	4.1	3.7

Map sharpening b-factor (Å^2)	-152.97	-101.8	-113.8	-67.2
Model composition Non-hydrogen atoms Protein residues Ligand (PLC)	55,460 7199 	36,113 4521 11	24,456 3089 0	32,891 4163 6
B factors (Å^2) Protein Ligand	 74.47	 75.52 59.93	 121.74	 69.29 80.08
R.M.S. deviations Bond lengths (Å) Bond angles (°)	0.004 0.710	0.003 0.611	0.004 0.596	0.003 0.671
Validation MolProbity score Clashscore Poor rotamers (%)	2.00 13.63 1.16	1.86 12.04 0.45	1.87 12.92 0.46	1.85 11.37 0.79
Ramachandran plot Favored (%) Allowed (%) Disallowed (%)	95.60 4.40 0.00	96.05 3.95 0.00	96.28 3.72 0.00	96 3.98 0.02

999 *Values in parentheses indicate the pixel size in super-resolution.