Alteration of long and short-term hematopoietic stem cell ratio causes myeloid-

biased hematopoiesis

Running title; Self-renewal heterogeneity causes myeloid skewing

Katsuyuki Nishi[1,2], Taro Sakamaki[1,2], Akiomi Nagasaka[1,2], Kevin S. Kao[3], Kay

Sadaoka[1,2], Masahide Asano[4], Nobuyuki Yamamoto[5], Akifumi Takaori-Kondo[6],

Masanori Miyanishi[1,2] *

[1]Hematopoietic Stem Cell Biology and Medical Innovation (HSCBMI), Department of

Pediatrics, Kobe University Graduate School of Medicine, Kobe, Hyogo 650-0047,

Japan

[2]RIKEN Center for Biosystems Dynamics Research, Kobe, Hyogo 650-0047, Japan

[3]Weill Cornell, Rockefeller, Sloan-Kettering, Tri-Institutional MD-PhD Program, New

York, NY

[4]Institute of Laboratory Animals, Kyoto University Graduate School of Medicine,

Kyoto, Kyoto, 606-8501, Japan.

[5]Department of Pediatrics, Kobe University Graduate School of Medicine, Kobe, Hyogo

650-0017, Japan

[6]Department of Hematology and Oncology, Kyoto University Graduate School of

Medicine, Kyoto, Kyoto 606-8507, Japan

*Correspondence: Masanori Miyanishi, M.D, Ph.D.

Hematopoietic Stem Cell Biology and Medical Innovation (HSCBMI), Department of

Pediatrics, Kobe University Graduate School of Medicine, Kobe, Hyogo 650-0047,

Japan

E-mail: miya75@med.kobe-u.ac.jp

Phone: 078-304-6037

FAX: 078-304-6100

Abstract

Myeloid-biased hematopoiesis is a well-known age-related alteration. Several possibilities, including myeloid-biased hematopoietic stem cell (HSC) clones, may explain this. However, the precise mechanisms remain controversial.

Utilizing the Hoxb5 reporter system to prospectively isolate long-term HSCs (LT-HSCs) and short-term HSCs (ST-HSCs), we found that young and aged LT-HSCs co-transplanted into the same recipients demonstrated nearly equivalent myeloid lineage output, contrary to the theory of myeloid-biased HSC clones. Transcriptomics indicated no significant myeloid gene enrichment in aged LT-HSCs compared to their young counterparts. Instead, transplanting reconstituted young HSCs with the ratio of LT/ST-HSCs seen in aged mice can significantly skew the lineage output to myeloid cells. In addition, while the niche environment in the bone marrow minimally affects myeloid-biased hematopoiesis, aged thymi and spleens substantially hinder lymphoid hematopoiesis, resulting in further myeloid-domination. Thus, we demonstrate that myeloid-biased hematopoiesis in aged organisms originates due to alteration of the ratio between LT-HSCs and ST-HSCs rather than in heterogeneous HSC clones with various cell fates.

Introduction

Age-associated changes in individuals are deeply correlated with progressive attenuation of cellular functions in tissue stem cells of organs (Jones and Rando 2011). In the hematopoietic system, hematopoietic stem cells (HSCs), which possess self-renewal capacity and multipotency (Spangrude et al., 1988; Weissman and Shizuru 2008; Majeti et al., 2007), are responsible for various hematopoietic alterations with age, such as reduced self-renewal capacity (Morrison et al., 1996) and myeloid-biased hematopoiesis (Muller-Sieburg et al., 2004; Challen et al., 2010) due to their functional decline. For example, myeloid-biased hematopoiesis potentially reduces response to infections (Webster, 2000), reduces vaccination efficacy (Crooke et al., 2019), and increases myeloid malignancy (Rossi et al., 2008) in aged individuals. To comprehend these age-associated physiological changes, myeloid-biased hematopoiesis has been studied in considerable detail.

Based on the experimental observation that transplantation of aged HSCs exhibits more myeloid-biased differentiation in young recipient mice than transplantation of young HSCs, this phenotype has been thought to originate in cell-intrinsic changes in the HSC compartment (Geiger et al., 2013; De Haan and Lazare 2018; Schultz and Sinclair 2016). A myeloid-biased phenotype after aged HSC transplantation leads to the suggestion that myeloid-biased HSC clones selectively expand with age (Cho et al., 2008; Pang et al., 2011). Transcriptome and epigenetic analyses showing that a set of genes related to myeloid differentiation is significantly enriched in aged HSCs compared to young HSCs (Rossi et al., 2005; Grover et al., 2016) further supports this hypothesis.

On the other hand, some reports support a different point of view in light of experimental evidence showing that aged lymphoid-biased HSCs still demonstrate the same level of

lymphopoiesis as their younger counterparts, despite exhibiting a myeloid-biased gene expression pattern (Montecino-Rodriguez et al., 2019). This result highlights the limitations of predicting a pattern of HSC differentiation based upon gene expression patterns. Moreover, myeloid progenitors such as granulocyte-macrophage progenitor (GMP) and common myeloid progenitor (CMP) should also increase with age if the selective expansion of myeloid-biased HSCs leads to an increase of myeloid cells in peripheral blood (PB). However, increment of such progenitors with age was not consistently demonstrated in earlier research (Rossi et al., 2005; Nilsson et al., 2016; Min et al., 2006). Additionally, a mathematical model demonstrated that aging had no effect on the daily production of CMP supplied from multipotent progenitors (MPP) (Busch et al., 2015; Dorshkind et al., 2020).

As such, while selective expansion of myeloid-biased HSC clones is the most widely accepted hypothesis to explain myeloid-biased hematopoiesis in aging (Mejia-Ramirez and Florian, 2020; SanMiguel et al., 2020), other mechanisms may exist in light of the inconsistent cellular behavior of progenitor fractions relative to HSCs. To the best of our knowledge, no reports have analyzed kinetics of age-associated changes in HSCs, progenitors, and PB cells simultaneously. Therefore, we started our investigation by examining the correlation between age-related changes in PB and BM to shed light on the mechanism underlying myeloid-biased hematopoiesis that occurs during aging.

Results

The discrepancy of age-associated alternation in peripheral blood and bone marrow calls into question the existence of a myeloid-biased clone

Mouse HSC research on aging has used mice aged 18 months or older (Challen et al.,

2010; Cho et al., 2008; D. J. Rossi et al., 2005; Grover et al., 2016; Montecino-Rodriguez

et al., 2019; Nilsson et al., 2016). However, given that continuous accumulation of

cellular stress with age causes a gradual decline of cellular functions, a comprehensive

analysis from young to old mice is necessary to unravel mechanisms by which age-

associated, myeloid-biased hematopoiesis progress. Hence, we analyzed changes in PB

(**Fig. S1 A**) at multiple time points from young to old mice. These results showed that the

percentage of myeloid cells began to change as early as 6 months in mice and continued

to increase at a constant rate until the age of ≥ 23 months (**Fig. 1 A**).

It has been reported that myeloid-biased hematopoiesis is caused by a selective increase

of myeloid-biased clones in the immunophenotypically defined (surface-antigen

defined) HSC fraction (**Fig. S1 B**) (Beerman et al., 2010). According to this hypothesis,

age-associated myeloid hematopoiesis progression in PB would be paralleled by an

increase in myeloid-biased HSC clones. Therefore, we examined the frequency of the

HSC and progenitor fractions in the BM at multiple time points (**Fig. S1, B and C**). We

found that the frequency of immunophenotypically defined HSC in BM rapidly increased

up to the age of 12 months. After the age, the rate of increase in their frequency appeared

to slow down. (**Fig. 1 B**). On the other hand, in contrast to what we anticipated, the

frequency of GMP was stable, and the percentage of CMP actually decreased

significantly with age, defying our prediction that the frequency of components of the

myeloid differentiation pathway, such as CMP, GMP, and MEP would increase in aged

mice if myeloid-biased HSC clones increase with age (**Fig. 1 B**).

Finally, we determined the ratio of each fraction in young mice versus aged mice to

compare the age-associated transition pattern of components comprising the myeloid

differentiation pathway. Our analysis of hematopoietic alternations with age revealed that

age-associated transition patterns of immunophenotypically defined HSC and CMP in

BM were not paralleled with myeloid cell in PB (**Fig. 1 C**). These findings called into

question the hypothesis that there is a selective increase of myeloid-biased HSC clones

in the aged BM. We then set out to elucidate the mechanism by which myeloid-biased

phenotypes arise in the PB of aged mice.

The long-term observation of 2-year-old LT-HSC's differentiation does not indicate the

expansion of myeloid-biased clones.

Numerous studies have claimed that the myeloid-skewed phenotype observed in PB of

aged mice is caused by myeloid-biased HSC clones selectively expanded from HSCs with

originally heterogeneous differentiation potentials (Beerman et al., 2010; Dykstra et al.,

2011; Sudo et al., 2000). Other studies have reported that immunophenotypically defined

HSCs possess heterogeneity associated with self-renewal capacity, suggesting the

existence of at least two different populations in the HSC compartment, LT-HSCs and

ST-HSCs (Morrison and Weissman 1994; Spangrude et al., 1995). Since LT-HSCs have

extensive self-renewal capacity, while ST-HSCs lose their self-renewal capacity within a

short period, LT-HSCs are thought to persist in the BM throughout life and to enrich age-

associated changes compared to ST-HSCs. Therefore, we expect that LT-HSC-specific

analyses will help to answer whether myeloid-biased HSC clones exist.

We previously reported that Hoxb5 exclusively marks LT-HSCs in young mice (Chen et

al., 2016). However, we have not tested in aged mice whether Hoxb5 specifically labels

LT-HSCs and helps to distinguish LT-HSCs and ST-HSCs from immunophenotypically

defined HSCs (hereafter, bulk-HSCs). To confirm this, we first analyzed expression of

Hoxb5 in bulk-HSC, MPP, Flk2$^+$, and Lin$^-$Sca1$^-$c-Kit$^+$ populations in aged *Hoxb5*-tri-

mCherry mice. We observed that Hoxb5 was exclusively expressed in bulk-HSCs (**Fig.**

2 A). Then, to verify long-term engraftment, we conducted a transplantation assay

utilizing Hoxb5$^+$ HSCs and Hoxb5$^-$ HSCs, respectively, isolated from 2-year-old mice

(**Fig. 2 B**). We observed that only recipients receiving aged Hoxb5$^+$ HSCs exhibited

continuous hematopoiesis 16 weeks after primary transplantation (**Fig. 2, C and E; Fig.**

S2 A). In secondary transplantation analysis, only recipients receiving Hoxb5$^+$ HSCs

exhibited robust hematopoiesis throughout the period of observation, indicating that

Hoxb5 can be used as a specific marker of LT-HSCs in aged mice, as well as young mice

(**Fig. 2, D and F; Fig. S2 B**).

A serial transplantation assay with a period of observation longer than 8 months should

be long enough to observe further myeloid-biased change. If bulk-HSCs isolated from

aged mice are already enriched by myeloid-biased HSC clones, we should see more

myeloid-biased phenotypes 16 weeks after primary and the secondary transplantation.

However, we found that kinetics of the proportion of myeloid cells in PB were similar

across the primary and the secondary transplantation (**Fig. 2 G**). These results suggest the

following two possibilities: either myeloid-biased HSCs do not expand in the LT-HSC

fraction, or the expansion of myeloid-biased clones in 2-year-old mice has already

peaked.

Direct comparison between young vs. aged LT-HSC differentiation reveals that LT-

HSCs exhibit unchanged lineage output throughout life

We developed a co-transplantation assay to directly compare the differentiation capacity

between young and aged LT-HSCs by co-transplanting ten young GFP$^+$ LT-HSCs and

ten aged GFP- LT-HSCs into the same recipient mice (**Fig. 3 A**). Then, we found that the

myeloid lineage proportions from young and aged LT-HSCs were nearly comparable

during the observation period after transplantation (**Fig. 3, B and C**). Furthermore, by

analyzing the proportion of mature cell types derived from young and aged LT-HSCs in

the same donor, we directly compared the capacity for hematopoietic reconstitution in

each mature cell type between young and aged LT-HSCs. We confirmed again that the

reconstitution ratio was almost the same across all lineages, although bulk hematopoiesis

derived from young LT-HSCs predominated (**Fig. 3 D**). These results indicate that the

differentiation potential of LT-HSCs remains unchanged throughout their lives.

Several reports have demonstrated that transplanting mixed/bulk-HSCs—a combination

of LT-HSCs and ST-HSCs—obtained from old animals results in blatantly myeloid-

biased hematopoiesis (Beerman et al., 2010; Dykstra et al., 2011; Sudo et al., 2000). To

test this, we co-transplanted bulk-HSCs from young and old mice to corroborate this (**Fig.**

S3 A). As previously described, we observed that aged bulk-HSC exhibited a myeloid-

skewed phenotype (**Fig. S3 B and C**). Additionally, we observed that aged bulk-HSC

reconstitution exhibited higher reconstitution of myeloid cells compared to young bulk-

HSCs (**Fig. S3 D**). These findings unmistakably demonstrated that mixed/bulk-HSCs

showed myeloid skewed hematopoiesis in PB with aging, while LT-HSCs exhibited

unchanged lineage output throughout life.

LT-HSCs never show myeloid-related gene set enrichment during aging

Although the results of co-transplantation of young and aged LT-HSCs did not

demonstrate lineage-biased output by LT-HSCs throughout life, enrichment of myeloid

genes in aged bulk-HSCs shown by previous studies supports the idea that myeloid-

biased hematopoiesis is caused by selective expansion of myeloid-biased HSC clones (D.

J. Rossi et al., 2005; Grover et al., 2016). To compare age-associated myeloid gene

enrichment, we isolated bulk-HSC, LT-HSCs, and ST-HSCs from young or aged mice

(**Fig. 4 A**).

To ensure the quality of the sorted fraction for subsequent RNA-seq analyses, we verified

the Hoxb5 read counts (**Fig. S4 A**). Cluster dendrograms using the whole transcriptome

confirmed that cell fractions isolated from young and aged mice clustered into distinct

groups (**Fig. 4 B**). Then, using sets of aging-related genes, such as inflammation

(Liberzon et al., 2015; Pietras 2017), DNA damage (p53 pathway) (Liberzon et al., 2015;

D. J. Rossi et al., 2007), cell cycle progression (E2F target) (Liberzon et al., 2015;

Kowalczyk et al., 2015), and a common aging signature (Flohr Svendsen et al., 2021), we

ran a violin plot analysis on each cell fraction to confirm the occurrence of typical aging-

related gene expression changes. We found age-associated changes in inflammation and

the DNA damage between cells isolated from young and aged mice, but very similar

patterns regardless of cell fraction in young or aged mice. In terms of cell cycle

progression and aging signature genes, however, we discovered that only the young LT-

HSC fraction differed from other young fractions and tended to represent similar gene

expression pattern with aged fractions (**Fig. 4 C**). Then, taking LT-HSC specific gene

expression pattern into account, we looked at expression patterns of specific genes that

had been highly validated by experiments as being associated with myeloid-biased HSC

(Beerman et al., 2010; Flohr Svendsen et al., 2021; Sanjuan-Pla et al., 2013; Mann et al.,

2018). We discovered that expression of these genes is relatively comparable between

young and aged LT-HSCs, but not other fractions (**Fig. S4 B**).

A recent comprehensive analysis of mouse HSC aging using multiple RNAseq data sets

claimed that almost 80% of differentially expressed genes are poorly reproducible across

data sets (Flohr Svendsen et al., 2021). In fact, we found that almost 80% of genes are

not shared when we compare three representative myeloid/lymphoid gene sets (Sanjuan-

Pla et al., 2013; Chambers et al., 2007; Pronk et al., 2007). Additionally, most genes were

only used in a single gene set (**Fig. 4, D and E**). By using only genes that were shared in

these gene sets, we ran a gene set enrichment analysis (GSEA) to see whether

myeloid/lymphoid genes were enriched in aged LT-HSCs or other fractions. Neither aged

LT-HSCs nor aged ST-HSCs exhibited myeloid/lymphoid gene set enrichment, while

shared myeloid-related genes tended to be enriched in aged bulk-HSCs, although this

enrichment was not statistically significant (**Fig. 4, F and G**). On the other hand, GSEA

analysis using original gene sets, respectively, showed inconsistent results (**Fig. S4, C**

and D).

A myeloid-biased phenotype in peripheral blood depends on the relative decrease of

ST-HSC in the HSC compartment with age

While transplantation of aged bulk-HSCs exhibits a myeloid-biased phenotype in PB

shown in previous reports (Beerman et al., 2010; Dykstra et al., 2011; Sudo et al., 2000)

and our results (**Fig. S3**), aged LT-HSCs remain stable in terms of the balance of their

differentiation potential between myeloid and lymphoid production (**Fig. 3**). Comparative

transcriptome analyses demonstrated that neither LT-HSCs nor ST-HSCs, which are

functionally more homogeneous than bulk-HSCs, exhibited myeloid gene set enrichment

with age, whereas aged bulk-HSCs tended to show more myeloid gene set enrichment

than their young counterparts (**Fig. 4**). We then tried to figure out what causes the

myeloid-biased phenotype in PB after transplantation of bulk-HSCs.

Because LT-HSCs in bulk-HSCs exhibit nearly constant level of hematopoiesis

throughout life, we hypothesized that ST-HSCs could be the key to discovering

mechanisms underlying the myeloid-skewing phenomenon. First, we revisited our

previous study (Chen et al., 2016), which demonstrated that transplanted ST-HSCs

maintain lymphocyte production while rapidly losing myeloid lineage production in

recipients. This result may potentially indicate the presence of HSC clones with a

preference for lymphoid differentiation. Given that donor cells obtained from primary

recipients receiving ST-HSCs do not undergo hematopoiesis in secondary transplantation

(Chen et al., 2016), it is likely that all ST-HSCs have already lost their ability to self-

renew and have disappeared from the HSC fraction in primary recipients. To confirm this,

we transplanted ten LT-HSCs or ST-HSCs and then analyzed PB and BM (**Fig. 5 A**). As

we previously reported (Chen et al., 2016), donor-derived hematopoiesis eventually

becomes lymphocyte-dominant in PB of ST-HSC recipients over time (**Fig. 5 B**).

Subsequently, it was determined that the vast majority of T cell subsets are memory cells

(**Fig. 5, C and D**). In addition, unlike recipients receiving LT-HSCs, recipients receiving

ST-HSCs lacked any bulk-HSCs in their BM (**Fig. 5, E and F**). These results strongly

suggest that lymphoid-biased hematopoiesis observed after transplantation of ST-HSCs

is due to persistence of memory-type lymphocytes in PB, rather than to *de novo*

lymphopoiesis from lymphoid-biased HSC clones. Persistence of memory-type

lymphocytes from ST-HSCs may have led us to misinterpret mechanisms underlying

myeloid-biased hematopoiesis after aged bulk-HSC transplantation. To verify this

hypothesis, we examined kinetics of the LT-HSC/ST-HSC ratio in the bulk-HSC

population, revealing the ratio of ST-HSC to LT-HSC decreased with age (**Figure S5**).

Thus, we hypothesized that the relative decrease in the ST-HSC ratio in the aged bulk-

HSC fraction would lead to reduction of memory-type lymphocytes and myeloid-biased

hematopoiesis following transplantation of aged bulk-HSCs. To test this hypothesis, we

isolated LT-HSCs and ST-HSCs from young donor mice and reconstituted them with a

2:8 ratio (as in young mice) or a 5:5 ratio (as in aged mice) prior to transplanting them

(**Fig. 6 A**). Four weeks after transplantation, PB analysis revealed that both groups had

comparable patterns of hematopoiesis. In contrast, recipient mice transplanted with a 5:5

ratio began to exhibit more myeloid-biased hematopoiesis eight weeks after

transplantation, and by week 16, they produced significantly more myeloid cells than the

other group (**Fig. 6, B-D**). Additionally, we conducted further investigations to determine

whether lymphoid hematopoiesis could be accelerated by alteration of LT-HSC/ST-HSC

ratios using cells from aged mice. To verify this, we isolated aged LT-HSCs and ST-

HSCs and reconstituted them with a 5:5 ratio or a 2:8 ratio prior to transplantation (**Fig.**

S6 A). Peripheral blood analysis revealed that lymphoid lineage output in the recipient

mice transplanted with a 2:8 ratio was significantly greater than those with a 5:5 ratio

(**Fig. S6 B-D**). Based on these findings, we concluded that myeloid-biased hematopoiesis

observed following transplantation of aged HSCs was caused by a relative decrease in

ST-HSC in the bulk-HSC compartment in aged mice rather than the selective expansion

of myeloid-biased HSC clones.

Age-associated extramedullary changes accelerate myeloid-biased hematopoiesis

We found that the fluctuating LT-HSC/ST-HSC ratio in the bulk-HSC compartment

corresponded with the myeloid-biased hematopoiesis associated with aging. In contrast,

the degree of myeloid bias observed in mice older than 23 months without transplantation

(**Fig. 1 A**; $49.0 \pm 23.4\%$) was significantly greater than in recipient mice receiving young

mixed HSCs with a 5:5 ratio 16 weeks after transplantation (**Fig. 6 D;** $30.8 \pm 30.8\%$).

This difference indicated that other intrinsic or extrinsic factors might exist in mice older

than 23 months to promote myeloid-biased hematopoiesis. To investigate this further, we

transplanted ten GFP$^+$ young LT-HSCs into young or 2-year-old recipient mice and

examined their PB (**Fig. 7 A**). We limited the observation period to 12 weeks because

many aged recipient mice had died prior to that point (**Fig. 7 B**). PB analysis revealed

that aged recipient mice receiving young LT-HSCs produced significantly more myeloid

cells than young recipient mice (**Fig. 7, C-E**). These results suggested that recipient-

dependent extrinsic factors, rather than intrinsic factors of transplanted HSCs, had a

greater impact on hematopoietic cell differentiation. Then, we examined BM to determine

what extrinsic factors affected differentiation of transplanted HSCs. We performed BM

analysis for the mice denoted by † in Figure 7C because many of the aged mice had died

before the analysis. Percentages of CMP and GMP, downstream components of myeloid

differentiation, were stable or significantly lower in aged recipient mice than in young

recipient mice, just as they were in non-treated aged mice (**Fig. 7 F; Fig. 1 B**). In addition,

the percentage of CLP, a downstream component of lymphoid differentiation, did not

differ between young and aged recipients (**Fig. 7 F**). In contrast to previous studies (Pinho

et al., 2018; Ergen et al., 2012), these findings suggested that intramedullary age-

associated changes may not have a significant impact on LT-HSC differentiation.

Thymi and spleens of recipient mice were then examined, as these are the structures of

lymphoid maturation and production of lymphoid progenitors following migration from

BM and prior to their appearance in PB. The thymus and spleen analyses were also

performed on the mice denoted by † in Figure 7C. Donor cells were not microscopically

detectable in the majority of aged recipient mice (**Fig. 7, G and I**), and quantitative

analysis using a flow cytometer revealed that the frequency of donor cells in spleens and

thymi of aged recipients was significantly lower than in young recipients (**Fig. 7, H and**

J). These results indicate that the process of lymphoid lineage differentiation is impaired

in the spleens and thymi of aged mice compared to young mice, or that differentiating

cells in the bone marrow do not successfully migrate into these secondary lymphoid

organs. These factors contribute to the enhanced myeloid-biased hematopoiesis in

peripheral blood due to a decrease in de novo lymphocyte production.

Discussion

Age-related, myeloid-biased hematopoiesis has been suggested as a potential reason for

the decline in acquired immunity in the elderly (Geiger et al., 2013). This process has

been linked to an increase in myeloid-biased HSC clones in the bulk-HSC fraction

identified by surface antigens through aging. By isolating LT-HSCs and ST-HSCs,

respectively, we discovered that age-related changes in the ratio of ST-HSCs to LT-HSCs

in bulk-HSCs is responsible for myeloid-biased hematopoiesis.

When aged bulk-HSCs were transplanted into young mice, post-transplant hematopoiesis

in recipient mice was significantly biased toward myeloid in PB compared to

transplantation of young bulk-HSCs (Beerman et al., 2010; Dykstra et al., 2011; Sudo et

al., 2000). Historically, this myeloid-biased hematopoiesis derived from aged bulk-HSCs

has been considered as evidence supporting expansion of myeloid-biased HSC clones

with aging (Geiger et al., 2013; de Haan and Lazare 2018). Other studies suggest that

blockage of lymphoid hematopoiesis in aged mice results in myeloid-skewed

hematopoiesis through alternative mechanisms. However, this result should be

interpreted carefully, since Busulfan was used for myeloablative treatment in this study

(Montecino-Rodriguez et al., 2019). To clarify the cause for this disparity, we chose to

examine the LT-HSC fraction, which persists in the BM for long periods of time and is

enriched for aging-related alterations. We discovered that post-transplant hematopoiesis

of aged LT-HSCs did not vary from that of young LT-HSCs in terms of differentiation

capacity. Expression of myeloid-related genes was not significantly altered when LT-

HSCs from young and aged mice were compared. Therefore, we inferred that age-related

myeloid-biased hematopoiesis cannot be attributed to an increase in myeloid-biased

HSCs, at least among LT-HSCs. Given that bulk-HSCs consist of LT-HSCs and ST-

HSCs, and that LT-HSCs exhibit no change in differentiation potential with age, we

hypothesized that ST-HSCs may be responsible for the myeloid-skewing phenotype in

PB. Transcriptomic analysis revealed no enrichment of myeloid-related genes in the ST-

HSC fraction, ruling out the possibility of an increase in myeloid-biased HSCs in the ST-

HSC fraction. Alternatively, we observed a proportionate decrease in ST-HSCs in bulk-

HSCs accompanying an increase in LT-HSCs. We postulated that this relative decrease

in ST-HSC ratio is responsible for myeloid-biased hematopoiesis. When isolated LT-

HSCs and ST-HSCs from young mice were reconstituted to replicate an aged BM type in

terms of the LT-HSC/ST-HSC ratio, we discovered that PB was strongly skewed towards

myeloid cells after transplantation of reconstituted HSCs. In contrast, transplantation of

reconstituted HSCs with a young BM type resulted in a significantly lymphoid-skewed

PB profile.

On the other hand, it has been reported that myeloid-biased HSCs have a more persistent

capacity for self-renewal (Muller-Sieburg et al., 2004). In fact, in our prior investigation

comparing PB 16 weeks post-transplant, myeloid cell production in PB was greater when

LT-HSCs were transplanted, while ST-HSC transplantation resulted in a considerable

increase in lymphoid cell production (Chen et al., 2016). It is therefore possible that the

LT-HSCs and ST-HSCs we isolated utilizing the Hoxb5 reporter system are myeloid- and

lymphoid-biased HSCs, respectively. However, the myeloid-producing capacity of LT-

HSCs and ST-HSCs 4 weeks post-transplantation is equivalent (Chen et al., 2016). In

contrast, 16 weeks after transplantation, ST-HSC transplant recipients demonstrate

negligible myeloid development and a predominance of lymphocytes, unlike LT-HSC

transplant recipients (Chen et al., 2016). These characteristics show that early post-

transplant ST-HSCs lack the phenotypic of lymphoid-biased HSCs. When ST-HSC-

transplanted BM cells are employed as donors for secondary transplantation, no donor-

cell-derived hematopoiesis is detected in recipients (Chen et al., 2016). In addition, HSCs

are not observed in the BM following transplantation of ST-HSCs (**Fig. 5**). These results

indicate that ST-HSCs lose their capacity for self-renewal earlier after transplantation

than LT-HSCs. In addition, phenotypic examination of lymphocytes in PB after ST-HSC

transplantation revealed that nearly all of them were memory cells, and that there was no

de novo supply of T cells. In other words, ST-HSCs may have been erroneously classified

as lymphoid-biased HSCs due to the preservation of memory lymphocytes with a long

half-life in the peripheral circulation, despite the absence of *de novo* hematopoiesis

following transplantation of ST-HSCs. These findings suggest that the bias toward

myeloid and lymphoid lineages in post-transplant PB is not regulated by heterogeneity of

multipotency, but by heterogeneity of self-renewal capacity.

Multiple extrinsic factors have been implicated as HSC-independent causes of myeloid-

biased hematopoiesis associated with aging. Some findings suggest that cell interaction

with niche cells in BM is altered (Pinho et al., 2018), whereas others claim that chronic

inflammation in aged mice lowers production of lymphoid progenitors and induces

myeloid-biased hematopoiesis (Montecino-Rodriguez et al., 2019; Ergen et al., 2012). To

identify HSC-independent causes of myeloid-biased hematopoiesis specifically, we

transplanted young LT-HSCs into young or aged mice. Cellular differentiation observed

in BM revealed that proportions of GMP, MEP, and CLP are unchanged between young

and aged recipients. In contrast, an examination of spleens and thymi, the sites of

lymphocyte maturation, revealed the absence of donor cells in aged recipients. These

findings suggest that myeloid cells increase in aging mice due to a relative decrease in

the ratio of mature lymphocytes in PB, resulting from inhibition of lymphocyte

maturation outside the BM, such as in the thymus and spleen, rather than a significant

change in cell differentiation caused by the intramedullary environment. Attempts have

been made in the past to restore thymus function in aged mice by administering

keratinocyte growth factor (KGF) (S. W. Rossi et al., 2007). Indeed, in aged mice,

lymphocyte hematopoiesis recovers, which is consistent with our results that the

extramedullary environment influences the myeloid-bias phenotype in PB (S. W. Rossi

et al., 2007).

It has been clinically documented that recovery of the lymphocyte fraction in PB is

delayed in elderly donors, compared to young donors (González-Vicent et al., 2017;

Baron et al., 2006). Although it is undeniable that this phenomenon may be due to

decreased HSC hematopoietic potential, it may also be due to a change in the ratio of LT-

HSC/ST-HSCs in bulk-HSCs, as described in our report, which may give the appearance

of a myeloid bias in PB after transplantation. It has also been noted that lymphocyte

recovery is delayed in elderly patients compared to younger patients (Berger et al., 2008;

van der Maas et al., 2021). Age-related loss in thymus and spleen functions has been

recorded in humans, implying that the same mechanism is acting as in our mouse

observations. If it is possible to assess and regulate the LT-HSC/ST-HSC ratio in donors

and thymus/spleen function in recipients in the future, it will help to develop accurate

models for predicting lymphocyte recovery after transplantation as well as new

transplantation strategies.

In contrast, our findings should be considered in light of some limitations. In this report,

we primarily performed ten to twenty cell transplantation assays. Therefore, the current

theory should be revalidated using single-cell technology with lineage tracing system

(Yamamoto et al. 2018; Rodriguez-Fraticelli et al. 2020). This approach will investigate

changes in the self-renewal capacity of individual HSCs and their subsequent

differentiation into progenitor cells and peripheral blood cells. Additionally, in this report

we purified LT-HSCs by Hoxb5 reporter system. In contrast, various LT-HSC markers

have been previously reported. Therefore, it is ideal to validate our findings using other

LT-HSC makers.

In summary, we have demonstrated for the first time that the ratio of LT-HSC/ST-HSC

is important in age-related myeloid skewed hematopoiesis as an HSC-dependent factor

and that aging-related thymus and spleen dysfunction also contributes significantly as an

HSC-independent factor. Furthermore, here we propose a "self-renewal heterogeneity

model" as a new mechanism for hematopoietic heterogeneity, including myeloid-biased

hematopoiesis in aged mice (**Fig. 8**). In addition to this report, we have also shown that

cellular differentiation in the PB after transplantation can vary by changing

transplantation conditions, ultimately altering self-renewal ability (Sakamaki et al., 2021;

Nishi et al., 2022). This highlights the significance of deciphering molecular processes

that regulate the heterogeneity of HSC self-renewal capacity, which will help to provide

a better understanding of the hematopoietic system and the hematopoietic hierarchy in

the future.

Methods

Mice

Mice with an *Hoxb5*-tri-mCherry (C57BL/6J background), derived from our previous

work (Chen et al., 2016), were harvested for donor cells for transplantation, PB, and BM

analysis. CAG-EGFP mice (C57BL/6J background) were bred with *Hoxb5*-tri-mCherry

for transplantation experiments. 8-14-week-old C57BL/6-Ly5.1 mice (purchased from

Sankyo Labo Service) were used as recipients for transplantation assays. For aged

recipients, 2-year-old CD45.2 C57BL/6J mice were purchased from CLEA Japan.

Supporting BM cells were collected from 8-12-week-old C57BL/6-Ly5.1 × C57BL/6J

(F_1 mice CD45.1$^+$/CD45.2$^+$). All mice were housed in specific pathogen-free (SPF)

conditions and were carefully observed by staff members. Mice were bred according to

RIKEN or Kyoto University. All animal protocols were approved by the RIKEN Center

for Biosystems Dynamics Research and Kyoto University.

Flow cytometry, cell sorting, and bone marrow analyses

Flow cytometry and cell sorting were performed on a FACS Aria II cell sorter (BD

Biosciences) and analyzed using FlowJo[TM] software (BD Biosciences). Bone marrow

(BM) cells were collected from bilateral tibias, femurs, humeri, and pelvises in Ca^{2+}- and

Mg^{2+}-free PBS supplemented with 2% heat-inactivated bovine serum (Gibco) and 2 mM

EDTA (Thermo Fisher Scientific). Cells were passed through 100-μm, and 40-μm

strainers (Corning) before analysis and sorting. Prior to staining, samples were blocked

with 50 ug/mL rat IgG (Sigma-Aldrich) for 15 min. To enrich HSCs and progenitor

populations for sorting and bone marrow analyses, cells were stained with APC-

eFluor780-conjugated anti-c-Kit (clone: 2B8) and fractionated using anti-APC magnetic

beads and LS columns (both Miltenyi Biotec). c-Kit[+] cells were then stained with

combinations of antibodies as described in Supplemental Table 1. All staining was

incubated at 4°C for 30 min, except for CD34 staining, which was incubated for 90 min.

Samples were washed twice after staining. Prior to flow cytometry or cell sorting, samples

were stained with SYTOX Red Dead Cell Stain (Life Technologies) or 7-

aminoactinomycin D (BioLegend). Cells were double sorted for purity. For the combined

analysis of the upstream (HSC, MPP, Flk2+) and downstream (CLP, MEP, CMP, GMP)

fractions in Figures 1B and 7F, we normalized by c-Kit+ cells because we performed a

c-Kit enrichment for the bone marrow analysis.

Transplantation and peripheral blood analyses

12-24 hours prior to transplantation, C57BL/6-Ly5.1 mice, or aged C57BL/6J recipient

mice were lethally irradiated with single doses of 8.7 Gy or 9.1Gy. For transplantation

assays, donor cells were first combined with 2×10^5 whole bone marrow supporting cells

(C57BL/6-Ly5.1 × C57BL/6J F_1 mice CD45.1[+]/CD45.2[+]) in 200 μL of PBS with 2%

FBS, and then injected into the retro-orbital venous plexus. For evaluation of post-

transplant kinetics, PB was collected and analyzed. At each time point, 50 µL of blood

were collected from the tail vein and re-suspended in Ca^{2+}- and Mg^{2+}-free PBS

supplemented with 2 mM EDTA. Red blood cells were lysed twice on ice for 3 min with

BD Pharm Lyse Buffer (BD Pharmingen). Identification of leukocyte subsets was

performed by staining with antibodies. Antibody information was described in

Supplemental Table 1. For evaluation of lineage output, the frequency of each lineage

(NK cell, B cell, T cell, neutrophil, and monocyte) was determined in the whole fraction.

The analysis of donor lineage output was restricted to donor cells showing $\geq 0.1\%$ donor

chimerism at the last PB analysis to allow reliable detection. The percentage of donor

chimerism in PB was defined as the percentage of $CD45.1^-CD45.2^+$ cells among total

$CD45.1^-CD45.2^+$ and $CD45.1^+CD45.2^+$ cells. PB data represents mice which survived

until the last PB analysis.

Thymus and Spleen analysis

Thymi and spleens were harvested and disrupted into a single cell suspension. Cells were

passed through 100-µm, and 40-µm strainers (Corning). Isolated thymocytes and

splenocytes were incubated on ice for 30 min with appropriately diluted antibodies in

staining buffer. Identification of B cells in spleens and T cells in thymi was performed

by staining with antibodies described in Supplemental Table 1. Prior to flow cytometry

or cell sorting, samples were stained with SYTOX Red Dead Cell Stain (Life

Technologies).

Tissue imaging

Freshly dissected spleens and thymi were fixed in 4% PFA (Nacalai tesque) in PBS for

24 h at 4°C. After PFA washout with PBS, tissues were cryoprotected with 30% sucrose

in PBS for 24 h at 4°C, embedded in Tissue-Tek O.C.T. compound (Sakura Finetek), and

snap-frozen in liquid nitrogen. Serial 10-μm longitudinal cryostat sections were obtained

using CryoStar NX50 (Thermo Fisher Scientific). Cell nuclei were counterstained with

4',6-diamidino-2-phenylindole (DAPI, 1 μg/mL, Roche). To reduce autofluorescence,

tissue sections were treated with Vector TrueVIEW autofluorescence quenching kit

(Vector Laboratories). Images were obtained from Leica TCS SP8 (Leica).

RNA sequencing

Total RNA was isolated with Trizol (Thermo Fisher Scientific) and cleaned up using

RNeasy MinElute columns (QIAGEN). cDNA libraries were prepared from bulk-HSCs,

ST-HSCs and LT-HSCs using a KAPA RNA HyperPrep kit with RiboErase (HMR)

(Kapa Biosystems) and sequenced using a Hiseq 1500 (Illumina) to obtain 2×127 base-

pair (bp) paired-end reads. Three replicates were sequenced for each population. Raw

transcriptome sequence data were mapped to the genome (mm10) using HISAT2 (ver

2.1.0) (Kim et al., 2015). Alignments were then passed to StringTie (ver 1.3.4d), which

was used to assemble and quantify transcripts in each sample (Pertea et al., 2016). EdgeR

(ver 3.22.5) was then used to compare all transcripts across conditions and to produce

tables and plots of differentially expressed genes and transcripts (Robinson et al., 2009).

Normalization with the TMM method was performed with the edgeR package in

Bioconductor (https://bioconductor.org/). Genes (12,808 genes) were selected for

hierarchical clustering based on the mean of TMM normalized read counts across all

samples' cells (mean $\geq$ 1). Then, clustering was performed by using the hclust function

for R (distance = Spearman's correlation'; method = 'ward.D2'). We depicted Venn

diagrams and performed GSEA analyses in Figures 4D-G and S4C-D using previously

published gene sets after excluding genes that could not be annotated by our

transcriptome dataset. GSEA was performed using GSEA software

(http://www.broadinstitute.org/gsea) with default settings (Subramanian et al., 2005).

Quantification and statistical analyses

Statistical analyses were performed using ggplot2 in R (version 4.1.2) or Microsoft Excel.

Sample size for each experiment and replicate numbers of experiments are included in

figure legends. Statistical significance was determined using Welch's t test. P values <

0.05 were considered significant.

Data and materials availability

Sequencing data has been deposited in the Gene Expression Omnibus under accession

code GSE226803. Correspondence and requests for materials should be addressed to

M.M. (miya75@med.kobe-u.ac.jp).

Acknowledgements

We gratefully acknowledge Hiroshi Kiyonari for animal care and providing recipient

mice at RIKEN BDR. Shigehiro Kuraku, Mitsutaka Kadota, and Chiharu Tanegashima

provided technical support for RNA sequencing and analysis at RIKEN BDR. Momo

Fujii performed imaging experiments at RIKEN BDR. Hitomi Oga, Kayoko Nagasaka,

and Masaki Miyahashi provided laboratory management at Kobe University. The authors

gratefully acknowledge ongoing support for this work: Masanori Miyanishi was

supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant

Number JP17K07407 and JP20H03268, The Mochida Memorial Foundation for Medical

and Pharmaceutical Research, The Life Science Foundation of Japan, The Takeda

Science Foundation, The Astellas Foundation for Research on Metabolic Disorders,

AMED-PRIME, AMED under Grant Number JP18gm6110020. Taro Sakamaki was

supported by the JSPS Core-to-Core Program, JSPS KAKENHI Grant Numbers

JP21K20669, JP22K16334, RIKEN Junior Research Associate Program. Katsuyuki

Nishi was supported by JSPS Grant Numbers KAKENHI JP18J13408 and 23K15326.

Authorship Contributions

K.N. conceived, performed, analyzed the experiments. T.S. and K.S. performed

transplantation experiments. M.M. conceived, performed, analyzed, and oversaw the

experiments. K.N. and M.M. wrote the manuscript. T.S., A.N., K.S.K., K.S., M.A., N.Y.,

and A.T. provided comments on the manuscript.

Disclosure of Conflicts of Interest

The authors declare no conflicts of interest in relation to the present study.

References

Baron, Frédéric, Barry Storer, Michael B. Maris, Jan Storek, Fanny Piette, Monja
Metcalf, Kristen White, et al. 2006. "Unrelated Donor Status and High Donor Age
Independently Affect Immunologic Recovery after Nonmyeloablative
Conditioning." *Biology of Blood and Marrow Transplantation* 12 (11): 1176–87.
https://doi.org/10.1016/j.bbmt.2006.07.004.
Beerman, Isabel, Deepta Bhattacharya, Sasan Zandi, Mikael Sigvardsson, Irving L.
Weissman, David Brydere, and Derrick J. Rossia. 2010. "Functionally Distinct
Hematopoietic Stem Cells Modulate Hematopoietic Lineage Potential during
Aging by a Mechanism of Clonal Expansion." *Proceedings of the National*

Academy of Sciences of the United States of America 107 (12): 5465–70.
https://doi.org/10.1073/pnas.1000834107.
Berger, Massimo, O. Figari, B. Bruno, A. Raiola, A. Dominietto, M. Fiorone, M.
Podesta, et al. 2008. "Lymphocyte Subsets Recovery Following Allogeneic Bone
Marrow Transplantation (BMT): CD4+ Cell Count and Transplant-Related
Mortality." *Bone Marrow Transplantation* 41 (1): 55–62.
https://doi.org/10.1038/sj.bmt.1705870.
Busch, Katrin, Kay Klapproth, Melania Barile, Michael Flossdorf, Tim Holland-Letz,
Susan M. Schlenner, Michael Reth, Thomas Höfer, and Hans Reimer Rodewald.
2015. "Fundamental Properties of Unperturbed Haematopoiesis from Stem Cells in
Vivo." *Nature* 518 (7540): 542–46. https://doi.org/10.1038/nature14242.
Challen, Grant A., Nathan C. Boles, Stuart M. Chambers, and Margaret A. Goodell.
2010. "Distinct Hematopoietic Stem Cell Subtypes Are Differentially Regulated by
TGF-B1." *Cell Stem Cell* 6 (3): 265–78.
https://doi.org/10.1016/j.stem.2010.02.002.
Chambers, Stuart M., Nathan C. Boles, Kuan Yin K. Lin, Megan P. Tierney, Teresa V.
Bowman, Steven B. Bradfute, Alice J. Chen, et al. 2007. "Hematopoietic
Fingerprints: An Expression Database of Stem Cells and Their Progeny." *Cell*
Stem Cell 1 (5): 578–91. https://doi.org/10.1016/j.stem.2007.10.003.
Chen, James Y., Masanori Miyanishi, Sean K. Wang, Satoshi Yamazaki, Rahul Sinha,
Kevin S. Kao, Jun Seita, Debashis Sahoo, Hiromitsu Nakauchi, and Irving L.
Weissman. 2016. "Hoxb5 Marks Long-Term Haematopoietic Stem Cells and
Reveals a Homogenous Perivascular Niche." *Nature* 530 (7589): 223–27.
https://doi.org/10.1038/nature16943.
Cho, Rebecca H., Hans B. Sieburg, and Christa E. Muller-Sieburg. 2008. "A New
Mechanism for the Aging of Hematopoietic Stem Cells: Aging Changes the Clonal
Composition of the Stem Cell Compartment but Not Individual Stem Cells." *Blood*
111 (12): 5553–61. https://doi.org/10.1182/blood-2007-11-123547.
Crooke, Stephen N., Inna G. Ovsyannikova, Gregory A. Poland, and Richard B.
Kennedy. 2019. "Immunosenescence and Human Vaccine Immune Responses."
Immunity & Ageing 16 (1): 1–16. https://doi.org/10.1186/s12979-019-0164-9.
Dorshkind, Kenneth, Thomas Höfer, Encarnacion Montecino-Rodriguez, Peter D. Pioli,
and Hans Reimer Rodewald. 2020. "Do Haematopoietic Stem Cells Age?" *Nature*
Reviews Immunology 20 (3): 196–202. https://doi.org/10.1038/s41577-019-0236-2.
Dykstra, Brad, Sandra Olthof, Jaring Schreuder, Martha Ritsema, and Gerald de Haan.
2011. "Clonal Analysis Reveals Multiple Functional Defects of Aged Murine
Hematopoietic Stem Cells." *The Journal of Experimental Medicine* 208 (13):
2691–2703. https://doi.org/10.1084/jem.20111490.
Dykstra, Brad, Sandra Olthof, Jaring Schreuder, Martha Ritsema, and Gerald De Haan.
2011. "Clonal Analysis Reveals Multiple Functional Defects of Aged Murine
Hematopoietic Stem Cells." *Journal of Experimental Medicine* 208 (13): 2691–
2703. https://doi.org/10.1084/jem.20111490.
Ergen, Aysegul V., Nathan C. Boles, and Margaret A. Goodell. 2012. "Rantes/Ccl5
Influences Hematopoietic Stem Cell Subtypes and Causes Myeloid Skewing."
Blood 119 (11): 2500–2509. https://doi.org/10.1182/blood-2011-11-391730.
Flohr Svendsen, Arthur, Daozheng Yang, Kyung Mok Kim, Seka Lazare, Natalia
Skinder, Erik Zwart, Anna Mura-Meszaros, et al. 2021. "A Comprehensive

Transcriptome Signature of Murine Hematopoietic Stem Cell Aging." *Blood* 138 (6): 439–51. https://doi.org/10.1182/blood.2020009729.

Geiger, Hartmut, Gerald De Haan, and M. Carolina Florian. 2013. "The Ageing Haematopoietic Stem Cell Compartment." *Nature Reviews Immunology* 13 (5): 376–89. https://doi.org/10.1038/nri3433.

González-Vicent, Marta, Blanca Molina, Natalia Deltoro, Julián Sevilla, José Luis Vicario, Ana Castillo, Manuel Ramirez, and Miguel Ángel Díaz. 2017. "Donor Age Matters in T-Cell Depleted Haploidentical Hematopoietic Stem Cell Transplantation in Pediatric Patients: Faster Immune Reconstitution Using Younger Donors." *Leukemia Research* 57: 60–64. https://doi.org/10.1016/j.leukres.2017.03.001.

Grover, Amit, Alejandra Sanjuan-Pla, Supat Thongjuea, Joana Carrelha, Alice Giustacchini, Adriana Gambardella, Iain Macaulay, et al. 2016. "Single-Cell RNA Sequencing Reveals Molecular and Functional Platelet Bias of Aged Haematopoietic Stem Cells." *Nature Communications* 7. https://doi.org/10.1038/ncomms11075.

Haan, Gerald de, and Seka Simone Lazare. 2018. "Aging of Hematopoietic Stem Cells." *Blood* 131 (5): 479–87. https://doi.org/10.1182/blood-2017-06-746412.

Haan, Gerald De, and Seka Simone Lazare. 2018. "Aging of Hematopoietic Stem Cells." www.bloodjournal.org.

Jones, D. Leanne, and Thomas A. Rando. 2011. "Emerging Models and Paradigms for Stem Cell Ageing." *Nature Cell Biology* 13 (5): 506–12. https://doi.org/10.1038/ncb0511-506.

Kim, Daehwan, Ben Langmead, and Steven L. Salzberg. 2015. "HISAT: A Fast Spliced Aligner with Low Memory Requirements." *Nature Methods* 12 (4): 357–60. https://doi.org/10.1038/nmeth.3317.

Kowalczyk, Monika S., Itay Tirosh, Dirk Heckl, Tata Nageswara Rao, Atray Dixit, Brian J. Haas, Rebekka K. Schneider, Amy J. Wagers, Benjamin L. Ebert, and Aviv Regev. 2015. "Single-Cell RNA-Seq Reveals Changes in Cell Cycle and Differentiation Programs upon Aging of Hematopoietic Stem Cells." *Genome Research* 25 (12): 1860–72. https://doi.org/10.1101/gr.192237.115.

Liberzon, Arthur, Chet Birger, Helga Thorvaldsdóttir, Mahmoud Ghandi, Jill P. Mesirov, and Pablo Tamayo. 2015. "The Molecular Signatures Database Hallmark Gene Set Collection." *Cell Systems* 1 (6): 417–25. https://doi.org/10.1016/j.cels.2015.12.004.

Maas, Nicolaas G. van der, Erik G.J. von Asmuth, Dagmar Berghuis, Pauline A. van Schouwenburg, Hein Putter, Mirjam van der Burg, and Arjan C. Lankester. 2021. "Modeling Influencing Factors in B-Cell Reconstitution After Hematopoietic Stem Cell Transplantation in Children." *Frontiers in Immunology* 12 (May): 1–8. https://doi.org/10.3389/fimmu.2021.684147.

Majeti, Ravindra, Christopher Y. Park, and Irving L. Weissman. 2007. "Identification of a Hierarchy of Multipotent Hematopoietic Progenitors in Human Cord Blood." *Cell Stem Cell* 1 (6): 635–45. https://doi.org/10.1016/j.stem.2007.10.001.

Mann, Mati, Arnav Mehta, Carl G de Boer, Monika S Kowalczyk, Kevin Lee, Pearce Haldeman, Noga Rogel, et al. 2018. "Heterogeneous Responses of Hematopoietic Stem Cells to Inflammatory Stimuli Are Altered with Age." *Cell Reports* 25 (11): 2992–3005. https://doi.org/10.1016/j.celrep.2018.11.056.

Mejia-Ramirez, Eva, and Maria Carolina Florian. 2020. "Understanding Intrinsic Hematopoietic Stem Cell Aging." *Haematologica* 105 (1): 22–37. https://doi.org/10.3324/haematol.2018.211342.

Min, Hyeyoung, Encarnacion Montecino-Rodriguez, and Kenneth Dorshkind. 2006. "Effects of Aging on the Common Lymphoid Progenitor to Pro-B Cell Transition." *The Journal of Immunology* 176 (2): 1007–12. https://doi.org/10.4049/jimmunol.176.2.1007.

Montecino-Rodriguez, Encarnacion, Ying Kong, David Casero, Adrien Rouault, Kenneth Dorshkind, and Peter D. Pioli. 2019. "Lymphoid-Biased Hematopoietic Stem Cells Are Maintained with Age and Efficiently Generate Lymphoid Progeny." *Stem Cell Reports* 12 (3): 584–96. https://doi.org/10.1016/j.stemcr.2019.01.016.

Morrison, Sean J., and Irving L. Weissman. 1994. "The Long-Term Repopulating Subset of Hematopoietic Stem Cells Is Deterministic and Isolatable by Phenotype." *Immunity* 1 (8): 661–73. https://doi.org/10.1016/1074-7613(94)90037-X.

Morrison, Sean J, A M Wandycz, K Akashi, A Globerson, and Irving L Weissman. 1996. "The Aging of Hematopoietic Stem Cells." *Nature Medicine* 2 (9): 1011–16. https://doi.org/10.1038/nm0996-1011.

Muller-Sieburg, Christa E., Rebecca H. Cho, Lars Karlsson, Jing F. Huang, and Hans B. Sieburg. 2004. "Myeloid-Biased Hematopoietic Stem Cells Have Extensive Self-Renewal Capacity but Generate Diminished Lymphoid Progeny with Impaired IL-7 Responsiveness." *Blood* 103 (11): 4111–18. https://doi.org/10.1182/blood-2003-10-3448.

Nilsson, Alexandra Rundberg, Shamit Soneji, Sofia Adolfsson, David Bryder, and Cornelis Jan Pronk. 2016. "Human and Murine Hematopoietic Stem Cell Aging Is Associated with Functional Impairments and Intrinsic Megakaryocytic/Erythroid Bias." *PLoS ONE* 11 (7): 1–20. https://doi.org/10.1371/journal.pone.0158369.

Nishi, Katsuyuki, Taro Sakamaki, Kay Sadaoka, Momo Fujii, Akifumi Takaori-Kondo, James Y. Chen, and Masanori Miyanishi. 2022. "Identification of the Minimum Requirements for Successful Haematopoietic Stem Cell Transplantation." *British Journal of Haematology* 196 (3): 711–23. https://doi.org/10.1111/bjh.17867.

Pang, Wendy W., Elizabeth A. Price, Debashis Sahoo, Isabel Beerman, William J. Maloney, Derrick J. Rossi, Stanley L. Schrier, and Irving L. Weissman. 2011. "Human Bone Marrow Hematopoietic Stem Cells Are Increased in Frequency and Myeloid-Biased with Age." *Proceedings of the National Academy of Sciences of the United States of America* 108 (50): 20012–17. https://doi.org/10.1073/pnas.1116110108.

Pertea, Mihaela, Daehwan Kim, Geo M. Pertea, Jeffrey T. Leek, and Steven L. Salzberg. 2016. "Transcript-Level Expression Analysis of RNA-Seq Experiments with HISAT, StringTie and Ballgown." *Nature Protocols* 11 (9): 1650–67. https://doi.org/10.1038/nprot.2016.095.

Pietras, Eric M. 2017. "Inflammation: A Key Regulator of Hematopoietic Stem Cell Fate in Health and Disease." *Blood* 130 (15): 1693–98. https://doi.org/10.1182/blood-2017-06-780882.

Pinho, Sandra, Tony Marchand, Eva Yang, Qiaozhi Wei, Claus Nerlov, and Paul S. Frenette. 2018. "Lineage-Biased Hematopoietic Stem Cells Are Regulated by

Distinct Niches." *Developmental Cell* 44 (5): 634-641.e4.
https://doi.org/10.1016/j.devcel.2018.01.016.

Pronk, Cornelis J.H., Derrick J. Rossi, Robert Månsson, Joanne L. Attema, Gudmundur Logi Norddahl, Charles Kwok Fai Chan, Mikael Sigvardsson, Irving L. Weissman, and David Bryder. 2007. "Elucidation of the Phenotypic, Functional, and Molecular Topography of a Myeloerythroid Progenitor Cell Hierarchy." *Cell Stem Cell* 1 (4): 428–42. https://doi.org/10.1016/j.stem.2007.07.005.

Robinson, Mark D., Davis J. McCarthy, and Gordon K. Smyth. 2009. "EdgeR: A Bioconductor Package for Differential Expression Analysis of Digital Gene Expression Data." *Bioinformatics* 26 (1): 139–40. https://doi.org/10.1093/bioinformatics/btp616.

Rodriguez-Fraticelli, Alejo E., Caleb Weinreb, Shou Wen Wang, Rosa P. Migueles, Maja Jankovic, Marc Usart, Allon M. Klein, Sally Lowell, and Fernando D. Camargo. 2020. "Single-Cell Lineage Tracing Unveils a Role for TCF15 in Haematopoiesis." *Nature* 583 (7817): 585–89. https://doi.org/10.1038/s41586-020-2503-6.

Rossi, Derrick J., David Bryder, Jun Seita, Andre Nussenzweig, Jan Hoeijmakers, and Irving L. Weissman. 2007. "Deficiencies in DNA Damage Repair Limit the Function of Haematopoietic Stem Cells with Age." *Nature* 447 (7145): 725–29. https://doi.org/10.1038/nature05862.

Rossi, Derrick J., Catriona H.M. Jamieson, and Irving L. Weissman. 2008. "Stems Cells and the Pathways to Aging and Cancer." *Cell* 132 (4): 681–96. https://doi.org/10.1016/j.cell.2008.01.036.

Rossi, Derrick J, David Bryder, Jacob M Zahn, Henrik Ahlenius, Rebecca Sonu, Amy J Wagers, and Irving L Weissman. 2005. "Cell Intrinsic Alterations Underlie Hematopoietic Stem Cell Aging." *Proceedings of the National Academy of Sciences of the United States of America* 102 (26): 9194–99. https://doi.org/10.1073/pnas.0503280102.

Rossi, Simona W., Lukas T. Jeker, Tomoo Ueno, Sachiyo Kuse, Marcel P. Keller, Saulius Zuklys, Andrei V. Gudkov, et al. 2007. "Keratinocyte Growth Factor (KGF) Enhances Postnatal T-Cell Development via Enhancements in Proliferation and Function of Thymic Epithelial Cells." *Blood* 109 (9): 3803–11. https://doi.org/10.1182/blood-2006-10-049767.

Sakamaki, Taro, Kevin S. Kao, Katsuyuki Nishi, James Y. Chen, Kay Sadaoka, Momo Fujii, Akifumi Takaori-Kondo, Irving L. Weissman, and Masanori Miyanishi. 2021. "Hoxb5 Defines the Heterogeneity of Self-Renewal Capacity in the Hematopoietic Stem Cell Compartment." *Biochemical and Biophysical Research Communications* 539 (February): 34–41. https://doi.org/10.1016/j.bbrc.2020.12.077.

Sanjuan-Pla, Alejandra, Iain C. Macaulay, Christina T. Jensen, Petter S. Woll, Tiago C. Luis, Adam Mead, Susan Moore, et al. 2013. "Platelet-Biased Stem Cells Reside at the Apex of the Haematopoietic Stem-Cell Hierarchy." *Nature* 502 (7470): 232–36. https://doi.org/10.1038/nature12495.

SanMiguel, Jennifer M., Kira Young, and Jennifer J. Trowbridge. 2020. "Hand in Hand: Intrinsic and Extrinsic Drivers of Aging and Clonal Hematopoiesis." *Experimental Hematology* 91: 1–9. https://doi.org/10.1016/j.exphem.2020.09.197.

Schultz, Michael B., and David A. Sinclair. 2016. "When Stem Cells Grow Old:
Phenotypes and Mechanisms of Stem Cell Aging." *Development (Cambridge)* 143
(1): 3–14. https://doi.org/10.1242/dev.130633.
Spangrude, Gerald J., Diane M. Brooks, and Daniel B. Tumas. 1995. "Long-Term
Repopulation of Irradiated Mice with Limiting Numbers of Purified Hematopoietic
Stem Cells: In Vivo Expansion of Stem Cell Phenotype but Not Function." *Blood*
85 (4): 1006–16. https://doi.org/10.1182/blood.v85.4.1006.bloodjournal8541006.
Spangrude, Gerald J, Shelly Heimfeld, and Irving L Weissman. 1988. "Purification and
Characterization of Mouse Hematopoietic Stem Cells." *Science (New York, N.Y.)*
241 (4861): 58–62. https://doi.org/10.1126/science.2898810.
Subramanian, Aravind, Pablo Tamayo, Vamsi K. Mootha, Sayan Mukherjee, Benjamin
L. Ebert, Michael A. Gillette, Amanda Paulovich, et al. 2005. "Gene Set
Enrichment Analysis: A Knowledge-Based Approach for Interpreting Genome-
Wide Expression Profiles." *Proceedings of the National Academy of Sciences of*
the United States of America 102 (43): 15545–50.
https://doi.org/10.1073/pnas.0506580102.
Sudo, K, H Ema, Y Morita, and H Nakauchi. 2000. "Age-Associated Characteristics of
Murine Hematopoietic Stem Cells [In Process Citation]." *J Exp Med* 192 (9):
1273–80.
Webster, Robert G. 2000. "Immunity to Influenza in the Elderly." *Vaccine* 18 (16):
1686–89. https://doi.org/10.1016/S0264-410X(99)00507-1.
Weissman, Irving L., and Judith A. Shizuru. 2008. "The Origins of the Identification
and Isolation of Hematopoietic Stem Cells, and Their Capability to Induce Donor-
Specific Transplantation Tolerance and Treat Autoimmune Diseases." *Blood* 112
(9): 3543–53. https://doi.org/10.1182/blood-2008-08-078220.
Yamamoto, Ryo, Adam C. Wilkinson, Jun Ooehara, Xun Lan, Chen Yi Lai, Yusuke
Nakauchi, Jonathan K. Pritchard, and Hiromitsu Nakauchi. 2018. "Large-Scale
Clonal Analysis Resolves Aging of the Mouse Hematopoietic Stem Cell
Compartment." *Cell Stem Cell* 22 (4): 600-607.e4.
https://doi.org/10.1016/j.stem.2018.03.013.

815　**Figure 1.**

Figure 1. Comprehensive analysis of hematopoietic alternations with age shows a

discrepancy of age-associated changes between peripheral blood and bone marrow

(A) Average frequency of myeloid cells (neutrophils, monocytes) and lymphoid cells (B

cells, T cells, and NK cells) in PB at the age of 2-3 months ($n = 6$), 6 months ($n = 6$), 12-

13 months ($n = 6$), 18 months ($n = 6$), 20-21 months ($n = 5$), $\geq$ 23 months ($n = 16$).

Abbreviation: PB = Peripheral blood.

(B) Average frequency of immunophenotypically defined HSC and progenitor cells in

BM of 2-3-month mice ($n = 6$), 6-month mice ($n = 6$), 12-13-month mice ($n = 6$), $\geq$ 23-

month mice ($n = 7$).

(C) Age-associated changes of immunophenotypically defined HSC and myeloid

differentiation components (CMP and myeloid cells in the PB). The ratio of aged to young

frequency was calculated as (the fraction frequency at each aged mice (%)) / (the average

fraction frequency at 2-3-month mice (%)). *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$. Data

and error bars represent means ± standard deviation.

Figure 2.

Figure 2. The expansion of myeloid-biased clones was not observed in 2-year-old

LT-HSCs after their transplantation.

(A) Hoxb5 reporter expression in bulk-HSC, MPP, Flk2$^+$, and Lin$^-$Sca1$^-$c-Kit$^+$

populations in the 2-year-old *Hoxb5*-tri-mCherry mice (Upper panel) and 3-month-old

Hoxb5-tri-mCherry mice (Lower panel). Values indicate the percentage of mCherry$^+$ cells

± standard deviation in each fraction (*n* = 3).

(B) Experimental design to assess the long-term reconstitution ability of Hoxb5$^+$ HSCs or

Hoxb5$^-$ HSCs. Hoxb5$^+$ HSCs and Hoxb5$^-$ HSCs were isolated from 2-year-old CD45.2

Hoxb5-tri-mCherry mice and were transplanted into lethally irradiated CD45.1 recipient

mice with 2×10^5 supporting cells (Hoxb5$^+$ HSCs, *n* = 13; Hoxb5$^-$ HSCs, *n* = 10). For

secondary transplants, 1×10^7 whole BM cells were transferred from primary recipient

mice. Abbreviations: PB = Peripheral blood, RT = Radiation therapy.

(C) Percentage chimerism at 16 weeks after receiving ten aged Hoxb5$^-$ HSCs or ten aged

Hoxb5$^+$ HSCs. Each column represents an individual mouse.

(D) Percentage chimerism at 16 weeks after whole BM secondary transplantation. Donor

whole BM cells for secondary transplantation were taken from mice denoted by † in

Figure 2C.

(E) Kinetics of average donor chimerism in each PB fraction after primary

transplantation.

(F) Kinetics of average donor chimerism after secondary transplantation.

(G) Kinetics of average donor myeloid output (myeloid proportion in donor cells) in LT-

HSC recipient mice after primary and secondary transplantation. *P < 0.05. **P < 0.01.

***P < 0.001. Data and error bars represent means ± standard deviation.

Figure 3.

Figure 3. Aged LT-HSCs show balanced hematopoiesis throughout life

(A) Experimental design for competitive co-transplantation assay using young LT-HSCs

sorted from *Hoxb5*-tri-mCherry GFP mice and aged LT-HSCs sorted from *Hoxb5*-tri-

mCherry mice. Ten CD45.2$^+$ young LT-HSCs and ten CD45.2$^+$ aged LT-HSCs were

transplanted with 2×10^5 CD45.1$^+$/CD45.2$^+$ supporting cells into lethally irradiated

CD45.1$^+$ recipient mice ($n = 8$).

(B) Lineage output of young or aged LT-HSCs at 4, 8, 12, 16 weeks after transplantation.

Each bar represents an individual mouse.

(C) Lineage output kinetics of young LT- HSCs or aged LT- HSCs at 4, 8, 12, 16 weeks

post-transplant.

(D) Competitive analysis of young LT-HSCs vs. aged LT-HSCs lineage output at 4, 8,

12, 16 weeks post-transplant. Competitive ratio was calculated as the proportion of young

LT-HSCs derived cells vs. aged LT- HSCs derived cells in each fraction. Abbreviations:

Neut = Neutrophils, Mo = Monocytes, B = B cells, T = T cells, and NK = NK cells. Data

and error bars represent means ± standard deviation. "n.d." stands for "not detected."

Figure 4.

Figure 4. Myeloid-associated genes were not enriched in aged LT-HSCs compared to their young counterparts

(A) Experimental schematic for transcriptome analysis. LT-HSCs (n = 3), ST-HSCs (n =

3), and bulk-HSCs (n = 3) were sorted from young (2-3 months) or aged (23-25 months)

Hoxb5-tri-mCherry mice, after which each RNA was harvested for RNA sequencing.

(B) Hierarchical clustering dendrogram of whole transcriptomes using Spearman distance

and the Ward clustering algorithm.

(C) Violin plots showing normalized gene expression levels for each gene set in young

and aged LT-HSCs, ST-HSCs, and bulk-HSCs. Expression values for each gene were

standardized independently by applying *Z* score transformation.

(D, E) Venn diagram showing the overlap of genes between three myeloid signature gene

sets, and lymphoid signature gene sets (A Sanjuan-Pla et al., 2013, Pronk et al., 2007, SM

Chambers et al., 2007).

(F, G) Signature enrichment plots from GSEA analysis using defined myeloid and

lymphoid signature gene sets that overlapped in the three gene sets. Values indicated on

individual plots are the normalized Enrichment Score (NES) and q-value of enrichment.

***$P < 0.001$.

1000

1001

1002

1003 **Figure 5.**

1004 **A**

Figure 5. The memory-type lymphocytes in the peripheral blood make it look as if

ST-HSCs are lymphoid-biased HSCs

(A) Experimental design for assessing the lineage output of young LT-HSCs or ST-HSCs.

Ten LT-HSCs, or ten ST-HSCs were isolated from 2-month-old CD45.2 *Hoxb5*-tri-

mCherry GFP mice and were transplanted into lethally irradiated CD45.1 recipient mice

with 2×10^5 supporting cells (LT-HSCs, $n = 3$; ST-HSCs, $n = 4$).

(B) Kinetics of average frequency of lymphoid cells (B cells, T cells, and NK cells) in

donor fraction after LT-HSCs or ST-HSCs transplantation.

(C) Gating scheme to identify memory (Central and effector) T cells and naive T cells in

the PB after excluding doublets, dead cells, and non-donor cells.

(D) Percentage of memory (Central and effector) T cells and naive T cells in donor CD4$^+$

fraction 10 months after LT-HSC or ST-HSC transplantation.

(E) Gating scheme to identify donor cells in bulk-HSC fraction in bone marrow analysis.

(F) Donor chimerism in bulk-HSC fraction 12 months after LT-HSCs or ST-HSCs

transplantation. *$P < 0.05$. ***$P < 0.001$. Data and error bars represent means ± standard

deviation.

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

 Figure 6.

Figure 6. **Hematopoiesis after transplantation inclined either toward myeloid or lymphoid cell production by artificially changing the ratio of LT-HSC/ST-HSC**

(A) Experimental design for the transplantation of 2-3-month-old LT-HSCs and ST-HSCs in a 2:8 ratio (the same ratio as in young mice BM) or 5:5 ratio (the same ratio as in aged mice BM). Donor cells were transplanted with 2×10^5 CD45.1$^+$/CD45.2$^+$

supporting cells into lethally irradiated CD45.1$^+$ recipient mice (2:8 ratio, $n = 18$; 5:5

ratio, $n = 23$).

(B) Donor lineage output of young LT-HSCs and ST-HSCs transplanted either in a 2:8

ratio or a 5:5 ratio at 4, 8, 12, 16 weeks post-transplant. Each bar represents an individual

mouse.

(C) Kinetics of average lineage output of young LT-HSCs and ST-HSCs in a 2:8 ratio or

a 5:5 ratio at 4, 8, 12, 16 weeks post-transplant.

(D) Frequency of myeloid cells in donor cell fraction. $*P < 0.05$. $**P < 0.01$. Error bars

represent standard deviation. Data represent two independent experiments.

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

Figure 7.

Figure 7. Age-associated physiological changes drive differentiation of LT-HSCs toward myeloid cells

(A) Experimental design for assessing the impact of age-associated physiological changes on differentiation of LT-HSCs. Ten GFP$^+$ LT-HSCs sorted from young (2-3 months) *Hoxb5*-tri-mCherry GFP mice, were transplanted with 2×10^5 CD45.1$^+$/CD45.2$^+$ supporting cells into lethally irradiated young or aged recipient mice. We defined donor cells as GFP$^+$ cells and supporting cells as CD45.1$^+$/CD45.2$^+$ cells.

(B) Survival rate of recipient mice in each group.

(C) Donor lineage output in young or aged recipient mice 11-12 weeks after transplanting young LT-HSCs (young recipient, $n = 17$; aged recipient, $n = 10$).

(D) Myeloid output (Frequency of donor myeloid cells in donor fraction) in young or aged recipient mice 11-12 weeks after transplantation.

(E) Kinetics of lineage output from donor LT-HSCs in young or aged recipient mice 4, 8, 11-12 weeks after transplantation.

(F) Average frequency of donor bulk-HSC and progenitor cells in donor c-Kit$^+$ cells in BM (young recipient, $n = 8$; aged recipient, $n = 8$). BM samples were taken from mice denoted by † in Figure 7C.

(G) Representative immunofluorescence images of frozen spleen sections derived from young or aged recipient mice. Green: donor cells (GFP fluorescence); blue: DNA (DAPI); Scale bar: 200 μm.

(H) Frequency of donor cells in spleen B cells of young or aged recipient mice (young recipient, $n = 8$; aged recipient, $n = 8$). Spleens are taken from mice denoted by † in Figure 7C.

(I) Representative immunofluorescence images of frozen thymus sections derived from

young or aged recipient mice. Green: donor cells (GFP fluorescence); blue: DNA (DAPI);

Scale bar: 200 μm.

(J) Frequency of donor cells in thymus T cells of young or aged recipient mice (young

recipient, $n = 8$; aged recipient, $n = 8$). Thymi are taken from mice denoted by † in Figure

7C. *$P < 0.05$. ***$P < 0.001$. Error bars represent standard deviation. Data represent two

independent experiments.

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170 **Figure 8.**

1171

Figure 8. Our new model: **Self-renewal heterogeneity model**. It has been thought that

there were myeloid (My-) or lymphoid biased (Ly-) HSCs, and that clonal selection of

My-HSCs caused age-associated myeloid biased hematopoiesis. However, in our model,

LT-HSCs represent unbiased hematopoiesis throughout life. ST-HSCs lose their

hematopoietic ability within a short period and memory-type lymphocytes remains in the

PB after ST-HSC transplantation. These remaining memory-type lymphocytes make it

look as if ST-HSCs are lymphoid-biased (The upper section). As a result, the age-

associated relative decrease of ST-HSCs in bulk-HSC fraction causes myeloid biased

hematopoiesis with age. Additionally, the blockage of lymphoid differentiation at spleen

and thymus accelerates further myeloid biased hematopoiesis in aged mice (The lower

section).

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217