Claudin-4 modulates autophagy via SLC1A5/LAT1 as a tolerance mechanism for genomic instability in ovarian cancer.

Fabian R. Villagomez[1], Julie Lang[2], Patricia Webb[1], Margaret Neville[1], Elizabeth R. Woodruff[1], Benjamin G. Bitler[1, 3, *]

[1]Division of Reproductive Sciences, Department of Obstetrics and Gynecology, School of Medicine, University of Colorado, Anschutz Medical Campus, Aurora, Colorado.
[2]Department of Immunology and Microbiology, University of Colorado Anschutz Medical Campus, Aurora, USA.
[3]Lead contact.
*Correspondence: benjamin.bitler@cuanschutz.edu

Running title. Tolerance to genomic instability

Abstract. Genome instability is key for tumor heterogeneity and derives from defects in cell division and DNA damage repair. Tumors show tolerance for this characteristic, but its accumulation is regulated somehow to avoid catastrophic chromosomal alterations and cell death. Claudin-4 is upregulated and closely associated with genome instability and worse patient outcome in ovarian cancer. This protein is commonly described as a junctional protein participating in processes such as cell proliferation and DNA repair. However, its biological association with genomic instability is still poorly-understood. Here, we used CRISPRi and a claudin mimic peptide (CMP) to modulate the cladudin-4 expression and its function, respectively in *in-vitro* (high-grade serous carcinoma cells) and *in-vivo* (patient-derived xenograft in a humanized-mice model) systems. We found that claudin-4 promotes a protective cellular-mechanism that links cell-cell junctions to genome integrity. Disruption of this axis leads to irregular cellular connections and cell cycle that results in chromosomal alterations, a phenomenon associated with a novel functional link between claudin-4 and SLC1A5/LAT1 in regulating autophagy. Consequently, claudin-4's disruption increased autophagy and associated with engulfment of cytoplasm-localized DNA. Furthermore, the claudin-4/SLC1A5/LAT1 biological axis correlates with decrease ovarian cancer patient survival and targeting claudin-4 *in-vivo* with CMP resulted in increased niraparib (PARPi) efficacy, correlating with increased tumoral infiltration of T CD8+ lymphocytes. Our results show that the upregulation of claudin-4 enables a mechanism that promotes tolerance to genomic instability and immune evasion in ovarian cancer; thus, suggesting the potential of claudin-4 as a translational target for enhancing ovarian cancer treatment.

Keywords. HGSC; genomic instability-tolerance; SLC1A5/LAT1; PARPi; T-CD8+ cells TCR-zeta chain; MHC-ii; PD1-PDL-1.

Abbreviations: HGSC, high-grade serous carcinoma; CMP, claudin mimic peptide; PDX, patient derived xenograft; HIS, human immune system; PARP, Poly (ADP-ribose) polymerase; TIL, tumor infiltrating lymphocytes.

INTRODUCTION.

Ovarian cancer is the overarching term for a collection of unique diseases that represent the leading cause of death among gynecologic cancers. High-grade serous carcinoma (HGSC) is the most common and deadliest of these malignancies. Patients with HGSC initially respond to standard of care (surgery debulking, carboplatin, paclitaxel) and maintenance treatments (e.g., PARP inhibitors, PARPi). However, most of them eventually develop therapy resistance; therefore, overcoming such a resistance is an urgent need. [1,2]

Genome instability is a hallmark of cancer; consequently, there is tolerance for such a characteristic within the physiological state of tumors. This hallmark is one of the factors that contribute to tumor progression, intratumorally heterogeneity, and the development of chemoresistance. [1,3-5] It arises from alterations in cell cycle progression and DNA damage repair (DDR) mechanisms, resulting in an increased rate of mutations and generation of multiple chromosomal modifications, such as amplifications, translocations, and micronuclei formation. Tumors are composed of heterogeneous cell subpopulations (metapopulation) exhibiting varying degrees of alterations in the genome; nevertheless, these cells can restrict the accrual of those alterations to prevent severe chromosomal transformations. Thus, this indicates the existence of a threshold tolerance for genomic instability within the tumor, and exceeding this critical threshold renders the tumor cells susceptible to cell death. However, the cellular mechanisms regulating such a tolerance are not well-known. [3,6-11]

Claudin-4 is not normally expressed in the epithelium of the fallopian tube (which is believed to be the origin of HGSC) but is upregulated in up to 70% of HGSC tumors, [12] and its upregulation is associated with development of therapy resistance and worse patient outcome. This protein is traditionally though to regulate cell-cell junctions, more specifically tight junctions (a type of cell connections that are localized at the apical region of epithelial and endothelial cells); therefore, its primary function has been related to regulation of permeability. [13,14] In spite of that, it is known that claudin-4 also participates in multiple processes other than permeability, including migration, mitosis, apoptosis, and DNA repair. [12,15,16] Recently, we reported that in mitosis claudin-4 is associated in a way with proper dynamics of one of the components of the cytoskeleton, tubulin.[17] Likewise, we showed that claudin-4 participates in a mechanism of DNA repair, the nonhomologous end joining, NHEJ and found that the levels of claudin-4 expression correlate with frequency of genetic modifications as well as therapy resistance in ovarian tumor samples.[12]

However, the molecular mechanims by which claudin-4 regulates genomic instability in ovarian cancer are still poorly understood. Here, we generated *in-vitro* (claudin-4 overexpression [OVCAR8 cells which do not express claudin-4]; and claudin-4 downregulation [OVCA429 and OVCAR3 cells which do express claudin-4] in HGSC cells) and *in-vivo* (PDX-HIS mice; patient-derived xenograft-human immune system mice) systems to modulate the function of claudin-4. We found that claudin-4 drives a dual regulatory mechanism on cell-cell junctions and cell cycle progression, contributing to the threshol-tolerance of genomic instability by ovarian tumor cells. Furthermore, targeting claudin-4 *in vivo* was found to enhance the treatment efficacy of a FDA-approved PARPi, niraparib in inhibiting tumor progression and increasing intratumural infiltration of T CD8+ lymphocytes.

RESULTS.

Claudin-4 controls the heterogeneity of HGSC cells through fine-tuned cell cycle progression and inhibition of chromosomal amplifications.

Tumor heterogeneity is associated with diverse expression programs and biological functions, a phenomenon also observed in tumor-derived cell lines. Among these biological functions, the cell cycle stands out. [18,19] [20] This cellular process is significantly controlled by nutrient availability and [21] is intimately related to chromosomal instability. [22,23] Claudin-4 has been reported to participate in the cell cycle, but its biological role is still poorly-understood. [17] To address in more detail the biological role of claudin-4 in the cell cycle and genomic instability, we modulated its expression in different HGSC cells (*Supplementary table 1*). For this purpose, we used claudin-4 downregulation (knockdown, KD) and overexpression (claudin-4 overexpression, C4) systems, along with a claudin mimic peptide (CMP) designed by our laboratory. The peptide targets a homologous and conserved sequence located in the short extracellular loop in claudin-4 (*Supplementary Figure 1a, b*), thereby interfering with its interaction with other proteins. [16,24-26]

First, we analyzed cycle progression by flow cytometry using wild type (WT) HGSC cells stained with propidium iodide. Under common culture conditions, the phenotype we observed is that HGSC cells cycle among all phases (G0-G1, S, and G2-M). Particularly, over time (24h *vs* 48h) some tumor cells stayed longer in the G0-G1 phase, downregulating their transit to the S and G2-M phases. Thus, we verified that the cell cycle progresses heterogeneously within both similar (same cell line) and different (OVCAR8, OVCA429, and OVCAR3) HGSC cells, fluctuating over time. This resulted in some cells not actively preparing for division (G0-G1) or clearly in mitosis (G2-M) (**Figure 1a**), potentially due to decreased nutrient availability (*Supplementary Figure 1c*). Manipulation of claudin-4 expression modified such heterogeneity in HGSC cells. Overexpression of claudin-4 associated with significant reduction of cells present in the S-phase (**Figure 1b**). Downregulation of claudin-4 correlated with significantly more cells in the G2-M phase and fewer in the G0-G1 phases in OVCA429 cells (**Figure 1c**), with a similar trend observed in OVCAR3 cells (**Figure 1d**). This observation is consistent with the higher number of OVCAR3 cells quantified in the G2-M phase (synchronized using starvation at 6h), as reported previously due to claudin-4 downregulation. [17] Collectively, our results show that claudin-4 significantly influences the heterogeneity of the cell cycle progression in HGSC cells by supporting their transit through various phases. Within this cellular process, it appears that the expression of claudin-4 arrests some

HGSC cells in the G0-G1 phase, as indicated by more cells in the G0-G1 phase at 24h and 48h (**Figure 1b**), resulting in a smaller number of cells transitioning to the S phase (**Figure 1b**). This phenotype is associated with the observation of a finely-tuned exit from the G2-M phase, coupled with decreased entry into the G0-G1 phase observed in OVCA429 cells during claudin-4 downregulation (**Figure 1c**), also mirroring the phenotype observed in OVCAR3 cells (**Figure 2d**, with fewer cells in the G0-G1 and more in the G2-M phases at 24h).

Additionally, it is known that an essential and shared characteristic between cell cycle progression and cancer development is nuclear morphological alteration. Cell cycle progression involves the remodeling of the nucleus as it advances through each phase, and its proper control is key to avoiding genomic instability. [27-33] Cancer development is related to significant changes in the morphology of the nucleus, which are intimately linked with genome instability. Particularly in ovarian cancer, there is a positive correlation between the nuclei size and chromosomic amplifications, a type of genomic instability. [34-39] Claudin-4 is associated with genomic instability [12], and in ovarian tumor samples, there is positive correlation between claudin-4 expression and reduced chromosomic amplifications (**Figure 1e**). Thus, indicating that claudin-4 somehow is limiting genomic instability. To gain further insights into this correlation, we used CMP to interfere with the function of claudin-4 and we examined indicators of genomic instability in HGSC cells through morphometric characterization of nuclei (stained with dapi). Targeting both claudin-4 expression and its function derived in significant morphological modifications in the nuclei of HGSC cells. The nuclear morphological architecture was reduced in claudin-4 overexpressing cells, and this phenotype was reversed by the effect CMP treatment, dependent on claudin-4 expression (**Figure 1f**). Conversely, this nuclear architecture was amplified when claudin-4 was downregulated in different HGSC cells (**Figure 1g, h**). However, the CMP effect on increasing nuclear size, was recapitulated only in OVCA429 cells. In contrast, the effect of CMP in OVCAR3 cells led to reduced nuclei size. This highlights not only the intratumoral diversity and a key biological role of the claudin-4 protein levels but also a significant effect of CMP in modifying genomic instability in HGSC cells, suggesting that the claudin-4-interacting proteins in the plasma membrane of OVCA429 and OVCAR3 could differ as well (*Supplementary Figure 1b*). Given that nuclei size is a signal of genomic instability, these initial results indicate that in our *in vitro* models, claudin-4's expression exhibits a similar phenotype to that in HGSC tumor concerning genomic instability (**Figure 1e**). Additionally, these findings highlight the

157 biological significance of claudin-4 in ovarian tumors, suggesting its possible regulatory role of

158 proper cell cycle progression to prevent one of the indicators of genomic instability, chromosomic

159 amplifications.

Claudin-4 controls a physiological connection between genomic instability and the actin
161
cytoskeleton in HGSC cells.
162

The cytoskeleton is composed of elements such as actin, microtubules, and intermediate filaments,

which are intimately related to cell-cell junctions in maintaining the morphology of epithelial and

endothelial tissues. This structure physically connects nuclei with cell-cell junctions, specifically

through junctional F-actin, and its dynamics are essential during cell cycle. [27-29,31,33,40,41] To further

explore the regulatory role of claudin-4 in cell cycle and genomic instability, we stained HGSC

cells with phalloidin to label the main component of the cytoskeleton, actin [42] and assessed its

intracellular distribution, particularly at the perinuclear and cell junction regions. Our analysis

revealed significant changes in the actin-cytoskeleton, characterized by an increased accumulation

of perinuclear F-actin during claudin-4 overexpression (**Figure 2a**) and the opposite effect during

its downregulation, notably in OVCA429 cells (**Figure 2b, c**). Additionally, all HGSC cells treated

with CMP exhibited a trend of reduced F-actin localization at the perimeter of nuclei. This result

implies that the proper functioning of claudin-4 is important for such localization. Further

differences were observed at the cell junction region, where F-actin accumulated significantly less

when claudin-4 was downregulated in various HGSC cells (**Figure 2d, e**). However, CMP

treatment did not affect this accumulation, indicating that the adequate function of claudin-4 in

maintaining junctional F-actin is less sensitivity to CMP inhibition compared to perinuclear F-

actin, suggesting the relevance of claudin-4 in cell-to-cell connections.

Given that claudin-4 is traditionally described as a junctional protein, we hypothesize that the

reduction of F-actin observed during claudin-4 downregulation in fixed cells could alter cell

connections in living cells, thus changing tissue architecture and the phenotype of tumor cells, as

reported. [43,44] Therefore, we generated HGSC cells expressing lifeact (a marker of F-actin) and

performed time-lapse confocal imaging to analyze junctional F-actin dynamics as previously

reported.[44] As a result of claudin-4 downregulation, cell junctions moved more and irregularly

over time. This observation is consistent with results observed in fixed cells (**Figure 2d, e**) and

strongly suggest that claudin-4 downregulation makes cell-cell junctions more unstable and

dynamic (**Figure 2f, g**; *Supplementary Figure1d, Supplementary video 1 and 2*). We conclude that

the effect of claudin-4 in cell cycle and genomic instability is intimately related to its functional

association with the actin-cytoskeleton, especially actin dynamics at the perinuclear and cell

junction regions.

The disruption of claudin-4 correlates with the formation of micronuclei and the activation

of the cGAS-STING/autophagy molecular axis.

The physical connection of nuclei with the cytoskeleton occurs through the linker of

nucleoskeleton and cytoskeleton (LINC) complex, which binds to nuclear lamina and underlies

the nuclear envelope. [45] Maintaining the integrity of the nuclear envelope (NE) is essential to avoid

the release of nuclear content into the cytoplasm. However, this structure can be ruptured by

pressure exerted by perinuclear F-actin, [30,46,47] which is enhanced when the function of p53 is lost.

[48] These conditions can lead to the formation of micronuclei originating from different sources of

DNA and phases of the cell cycle, such as interphase and mitosis. [49-51] Similar to the nucleus,

micronuclei can also be affected by NE ruptures due to changes in lamin B1 and lamin A/C (the

main component of nuclear lamina) dynamics. Consequently, DNA is more likely to be released

into the cytoplasm, potentially activating the cGAS-STING pathway for an anti-tumor immune

response. [30,51-54] We analyzed the possible association of claudin-4 with micronuclei formation and

the cGAS-STING pathway in HGSC cells by confocal microscopy and immunoblotting.

We identified micronuclei in OVCAR8, OVCA429, and OVCAR3 cells, indicating that this type

of chromosomic alteration is a common characteristic of HGSC cells. However, the expression

and function of claudin-4 was found to be somehow related to micronuclei formation. Both

overexpression and downregulation of claudin-4 correlated with significant changes in the

frequency of micronuclei quantification. Specifically, a positive correlation was observed between

claudin-4 overexpression and an increased number of micronuclei (**Figure 3a**), and a trend towards

the opposite effect was noted during its downregulation in OVCA429 cells (**Figure 3b**). In

OVCAR3 cells, a decrease in claudin-4 expression correlated positively with a higher incidence

of micronuclei (**Figure 3c**). These differences could be attributed to the unique doubling time of

OVCAR8 and OVCAR3 cells (**Figure 1a**) and to the similar trend we observed in the cell cycle

between these cells (OVCAR8 and OVCAR3), both of which were more localized in the G2-M

phase at 24h (**Figure 1b, d**). In addition, we quantified the total levels of lamin B1 and lamin A/C,

finding that these proteins tended to vary before claudin-4 manipulation, particularly during

claudin-4 downregulation (*Supplementary Figure 1e-g*). Furthermore, we characterized the

nuclear envelope surrounding the micronuclei) in all HGSC cells and identified that some of these

structures lacked components of the nuclear lamina (lamin B1, lamin A/C) (**Figure 3d-f**).

Interestingly, during claudin-4 downregulation, the localization of lamin proteins in the nucleus

changed significantly, notably with a higher amount of lamin B1 in HGSC claudin-4 knockdown

cells (**Figure 3g, h**), indicating changes in lamin B1 dynamic. [54] This result correlates with the

reduced perinuclear F-actin observed in KD cells (**Figure 2b, c**). Given that claudin-4 is localized

in the nucleus in OVCAR3 [55] as well as in OVCA429 (not shown), this phenotype suggests that

claudin-4 may modify nuclear lamina dynamics, possibly through a displacing mechanism. This

mechanism is akin to the competition and exclusion observed between the actin bundling proteins,

actinin and fascin, where they compete and exclude each other to bundle actin filaments into

different actin networks. [56] Although the origin of micronuclei formation was not determined, our

results show that claudin-4 impacts its formation or elimination, possibly through autophagy, as

observed in osteosarcoma U2OS cells. [57] Moreover, the lack of nuclear lamina in micronuclei

highlights the possibility that the DNA would be more likely to be released into the cytoplasm and

activate the cGAS-STING pathway for an anti-tumor immune response. [30,51,52]

In ovarian cancer, the cGAS-STING pathway is typically inhibited as a mechanism of tumor

immune evasion. [52,58] However, our data suggest that micronuclei in HGSC cells are likely to

release DNA into the cytoplasm, activating the cGAS-STING pathway. Therefore, we evaluated

the activation status of this pathway using immunoblotting and confocal microscopy. Briefly, upon

cytoplasmic DNA detection by cGAS, cGAMP is produced, which then binds STING, recruiting

TBK1. This process culminates in the phosphorylation TBK1 and STING (pTBK1 and pSTING),

followed by downstream signaling of IRF3 to produce a type I interferon (IFN) response (**Figure**

4a). On the other hand, pSTING can induce autophagy through an independent mechanism of

TBK1.[59,60] Manipulating the function of claudin-4 resulted in changes in the phosphorylation status

of STING and TBK1 (**Figure 4b***); suggesting that the activity of the cGAS-STING pathway was

altered due to modifications in claudin-4 expression. The levels of pSTING were increased in both

claudin-4 overexpression and downregulation, while the levels of pTBK1 (downstream of

pSTING) were reduced under these conditions. This correlation, with more pSTING and less pTBK1, suggests that pSTING might be inactivated, leading to reduced pTBK1 or an increase in the type I IFN response. A recent report described that the levels of pSTING are regulated through direct encapsulation by lysosomes (micro autophagy). [61] Consequently, we stained HGSC cells for pSTING and lamp1 (lysosome marker) and visualized these cells by confocal microscopy. We found that pSTING was notably elevated before the claudin-4 downregulation (**Figure 4c**), consistent with our immunoblotting results. Moreover, in HGSC cells, we observed that pSTING was associated with lamp1, apparently inside lysosomes. This observation suggests that the direct encapsulation of pSTING by lysosomes could be occurring in HGSC cells as well. [61] This finding emphasizes the importance of the lysosome-mediated degradative pathway, as well as the necessity of vesicular intracellular trafficking in such a process, for the proper functioning of the cGAS-STING/Autophagy axis regulation. This notion is supported by our results in OVCA429 WT cells, which were manipulated to express ISRE (interferon-stimulated response element) to measure more directly the activation of a type I interferon IFN response. We treated these cells with cGAMP (an activator of cGAS-STING and type I IFN response) and found that this treatment increased a type I IFN response in OVCA429 cells; however, this induction was ablated when blocking vesicular trafficking using chloroquine (CQ) (*Supplementary Figure 1h*).

On the other hand, we performed immunoblotting for the autophagy marker, LC3 A/B, which has two isoforms: LC3 A/B-i (cytosolic) and LC3 A/B-ii (membrane bound). The lipidated LC3 A/B-ii form binds to autophagosomes, which are subsequently degraded as autophagosomes trafficked intracellularly and fuse with lysosomes. Our immunoblotting indicated that the manipulation of claudin-4 expression also affected autophagy (**Figure 4d**), seemingly by increasing this pathway. This notion is supported by the observation that when OVCA429 cells were treated with chloroquine (an autophagy inhibitor via blocking vesicular trafficking) or rapamycin (an autophagy activator via inhibition of mTORC1), we observed changes in the levels of LC3 A/B-ii, like those observed when autophagy was induced with rapamycin. (**Figure 4e**) Additionally, the phosphorylation status of a downstream target of mTORC1, p70 S6 kinase, changed due to claudin-4 manipulation (overexpression and downregulation) (**Figure 4f-h**). Specifically, these levels of phosphorylation were reduced in claudin-4 knockdown cells (**Figure 4g, h**), suggesting that the function of mTORC1 was inhibited. Together, our results demonstrate that claudin-4

significantly impacts the cGAS-STING/autophagy molecular axis, particularly suggesting its

impact on the fate of micronuclei formation in HGSC cells.

Micronuclei are intimately related to autophagy which is controlled via a functional

connection of claudin-4 with the transporters of amino acids, SLC1A5/LAT1 in HGSC cells.

The axis of cGAS-STING/autophagy has been associated with the clearance of cytosolic DNA, [60]

and removal of micronuclei has been indicated in osteosarcoma U2OS cells. [57] Moreover, claudin-

4 appears to modulate micronuclei formation in some way. To confirm our hypothesis regarding

the function of claudin-4 in micronuclei modulation through autophagy, we generated HGSC cells

expressing mCherry-GFP-LC3 to measure autophagy activity (autophagy flux) using flow

cytometry and confocal microscopy, as previously reported. [62] Briefly, LC3 serves as a marker of

autophagy, localizing in vacuoles named autophagosomes. These vacuoles traffic intracellularly,

fusing with lysosomes, causing a decrease in the intraluminal vacuolar pH and subsequently

quenching of GFP fluorescence, unlike mCherry, that its resistance to such pH changes.

Consequently, autophagy flux results in an increase in mCherry positive cells derived from double

positive GFP-mCherry cells (*Supplementary Figure 2a*). To validate our strategy, HGSC-

mCherry-GFP-LC3 cells were treated with CQ (chloroquine) and rapamycin to block and activate

autophagy, respectively. These cells responded appropriately to these stimuli (*Supplementary*

Figure 2b). Similarly, we treated our cells with CMP and analyzed autophagy using flow

cytometry. We confirmed that claudin-4 participates in autophagy in HGSC cells. Both claudin-4

downregulation and overexpression correlated with increase autophagy activity (**Figure 5a-c**).

This suggests that claudin-4 does not directly favor or limit the autophagy pathway but rather

modulates it through other elements, e.g., mTORC1, the signaling complex regulating this activity,

as suggested by our findings (**Figure 4f-h**). Remarkably, we found that micronuclei are engulfed

through autophagy in HGSC cells (**Figure 5d-f**). Consequently, the possibility of DNA being

released into the cytoplasm due to collapsed micronuclei is reduced, potentially representing a

source to control genetic instability and nutrient supply. [63] Additionally, we found a similar

correlation of autophagy and micronuclei *in vivo*. The marker of this process, LC3 A/B was closely

associated to micronuclei (**Figure 5g**) and targeting claudin-4 with CMP showed signals of

modification for that cellular pathway (**Figure 5h-i**, *Supplementary Figure 3a-d*). Our findings

indicate a strong association of autophagy and the claudin-4's functional association with genome

instability.

Afterwards, we focused on examining the molecular mechanism governing the claudin-4 role in

autophagy. Previously, we identified the claudin-4-interacting proteins in OVCAR3 cells using

BioID. [55] Some of the identified elements (as claudin-4-interacting proteins) are the transporters

of amino acids SLC1A5, LAT1 (also known as SLC7A5), and SLC3A2. [55] The BioID method

strongly suggests that claudin-4 and those proteins are in the same protein complex and possibly

have a functional relationship. [64] It is known that LAT1 forms a heterodimer with SLC3A2 to

enable a bidirectional transport system of amino acids along with SLC1A5. This system regulates

upstream mTOR in cellular processes such as cell growth and autophagy which is mediated by a

key role of L-glutamine influx/efflux.[65]

We employed different methodologies to assess the effect of claudin-4 function on the amino acid

transport system and its association with autophagy regulation. Initially, we stained HGSC cells to

visualize SLC1A5, LAT1, and claudin-4. By confocal microscopy, we determined that claudin-4

co-localized with both transporters of amino acids (**Figure 6a, b**). Additionally, we analyzed the

protein expression of SLC1A5 and LAT by immunoblotting and found changes in SLC1A5

expression, especially in claudin-4 overexpression (*Supplementary Figure 3e*). However, the

changes in LAT1 expression were significantly different derived from claudin-4 downregulation.

There was a trend of less and more LAT1 expression in OVCAR8 claudin-4 cells (claudin-4

overexpression) and in OVCAR3 KD cells (claudin-4 downregulation), respectively. The opposite

effect (more expression) was determined in OVCA429 KD cells (**Figure 6c**, *Supplementary*

Figure 4a). This highlights differences among HGSC tumor cells as well as the relevance of

claudin-4 association with the bidirectional transport system formed by SLC1A5/LAT1 (**Figure**

6d), with a particular association with LAT1 in such tumor cells. Furthermore, for the activity of

the transport system of amino acids constituted by SLC1A5/LAT1, a key regulatory element is L-

glutamine, which is internalized by SLC1A5 to increase its intracellular concentration. The amino

acid then functions as a substrate for LAT1, which internalizes essential amino acids as L-

glutamine is exiting the cell. Consequently, we limit L-glutamine availability in culture media and

measure autophagy flux. All our HGSC cells exhibited increased autophagy flux due to L-

glutamine limitation (**Figure 6e**) which agrees with previous reports. [65] Likewise, we observed by

confocal microscopy changes in SLC1A5 intracellular distribution directly associated to claudin-4 overexpression and downregulation. The most significant difference was identified in OVCA429 KD cells, where SLC1A5 was markedly localized at the level of cell-cell junctions **(Figure 6f)** and interestingly, L-glutamine limitation was associated with changes in the SLC1A5 intracellular distribution, particularly in OVCA429 WT cell. In these cells, SLC1A5 localized clearly at the level of cell junctions because of L-glutamine limitation (**Figure 6g**). Additionally, because previously we observed accumulation of pSTING, which is involved in autophagy, in claudin-4 KD cells we cultured HGSC cells in media without L-glutamine and stained for pSTING and lamp1, as previously indicated (*Supplementary Figure 4b*). L-glutamine limitation correlated with accumulation of pSTING in OVCA429 and OVCAR3 WT cells, similarly to KD cells cultured in media with L-glutamine (**Figure 4c**). Also, autophagy blocking with CQ or autophagy induction with rapamycin was associated with changes in the intracellular distribution of SLC1A5 (*Supplementary Figure 4c, d*).

Finally, to verify specificity, we inhibited the transporter function of SLC1A5/LAT1 as previously described, [65] using L-g-glutamyl-p-nitroanilide (GPNA) and 2-aminobicyclo-(2,2,1) heptane-carboxylic acid (BCH), respectively and analyzed such inhibition on autophagy activity. Inhibition of SLC1A5 and LAT1 (**Figure 6h, i**) resulted in a trend of increase autophagy flux in HGSC cells, particularly in OVCAR3 cells; thus, confirming that specific inhibition of SLC1A5 and LAT1 dysregulates autophagy in HGSC cells. Our data strongly suggests that the bidirectional transport system mentioned before is present in HGSC cells and that claudin-4 is functionally associated with it. For instance, by stabilizing cell-cell junctions when that system is formed at that location, as suggested in **Figure 6d.**

Targeting claudin-4 *in vivo* modifies the tumor immune microenvironment and increases the therapy efficacy of niraparib in PDX-HIS mice.

Previously, we targeted claudin-4 *in vivo* with CMP using immunodeficient tumor-bearing mice. Disruption of claudin-4 resulted in reduction of tumor growth due to cell death.[16] However, the immunological consequences of targeting claudin-4 in tumors, via CMP are not known. To study the outcome of such intervention, we generated a humanized mouse model, the human immune system (HIS)-mice as previously reported.[66] This model is considered a better translational

approach than immunocompromised mice because it incorporates the contribution of a human
immune system in modeling the human tumor progression. [67] We implanted a reported patient-
derived xenograft (PDX, GTFB1016)[68] into the humanized mice to get our *in vivo* (PDX-HIS-
mice) system. After tumor establishment (**Figure 7a**, right; IVIS scan, before treatment), we
applied cycles of treatments to the tumor-bearing mice with CMP (claudin mimic peptide) and
follow-up tumor progression by IVIS scan. This this intervention was compared with a PARPi
therapy, niraparib (FDA-approved) and with a combinatory therapy, CMP/niraparib (see methods
and *Supplementary Figure 6a, b*).

As expected from our previous study, cycles of CMP treatment decreased tumor growth, [16] similar
to cycles of niraparib treatment (*Supplementary Figure 3f*). Remarkably, combinatorial therapy
showed a significant improvement in the effectiveness of niraparib, thus suggesting that claudin-
4 function is associated with resistance to niraparib treatment *in vivo* (**Figure 7a**). Likewise, the
physiological state of tumors (detected by EpCAM expression) was modified due to the
therapeutic intervention, particularly in the CMP/niraparib combination. We quantified lower
percentages of live tumor cells that were recovered from ascites samples (**Figure 7b**). As the same
methodology was applied to process all samples, these results suggest that CMP and niraparib
treatments somehow affected tumor survival, possibly by increasing tumor cell apoptosis. [16,69] In
addition, we identified that the frequency of cells expressing MHC-I (major compatibility complex
class-I) (**Figure 7c**) within the tumors was similar among all conditions, but not MHC-II (major
histocompatibility complex class-II) which was not only expressed by tumors (MHC-II is usually
expressed in professional antigen presenting cells, such as dendritic cells and B-cells which
stimulate the activation of T CD4+ cells), [70] but its localization was significantly decreased in
tumor cells treated with CMP/niraparib (**Figure 7d**). In different types of cancer, including ovarian
cancer, the expression of MHC-II is associated with a better patient outcome, highlighting (as in
other cancer types) the relevance of the T CD4+ lymphocyte activity as well as T CD8+ cells. [70-
75]

Furthermore, PDL-1+ tumor cells were observed more frequently in all groups of treated mice,
especially in niraparib and CMP/niraparib treated tumor-bearing mice, with the particularity that
the expression level of this protein slightly decreased in the CMP group (**Figure 7e**). Additionally,
these tumor cells expressed MHC-II with a similar trend to that observed in total live HGSC cells

(*Supplementary Figure 6a*). The expression of this protein has been reported previously in ovarian

cancer and related to a population of tumor cells with a stem cell phenotype. [76] This protein (PDL-

1) is traditionally associated with suppression of immunity by tumor cells, as a cellular mechanism

of tumor immune evasion, where PDL-1 can interact with its receptor (PD-1), which is upregulated

in T cells when they are activated and functions a mechanism of immune cell regulation. [77] For

example, it has been reported that high expression of PDL-1 in tumor cells and its interaction with

PD-1 in T cells, interfere with signaling cascade (zeta chain of the TCR, T cell receptor and CD28)

necessary to sustain the lymphocyte activation. [78] Strikingly, regardless of the treatment applied,

tumors with PDL-1 expression co-expressed PD-1 as well, approximately 50%, (**Figure 7f**). The

co-expression of PDL-1 and PD-1 has been previously reported in tumors, where such co-

expression is associated with a mechanism of tumor suppression associated to modulation of

mTOR signaling and independent of adaptive immunity. [79-82] Together, our results show that the

combination of CMP and niraparib treatment, modifies the physiological state of tumors by

regulating their cellular composition; particularly limiting the increasing of tumor cells with MHC-

II and increasing the number of tumor cells with PDL-1 and PD-1 co-expression. Thus, affecting

the known systemic immunosuppression induced in cancer [83].

During immune surveillance, immune cells migrate through different lymphoid organs, notably

the spleen [84,85]. It is known that in this organ, T CD4 lymphocytes cooperate with T CD8

lymphocytes to support their activation [86] and enhancement of tumor destruction [87].

However, the spleen also represents a site of tolerance to cancer development which is importantly

associated to the function of some myeloid cells. [88,89] Interestingly, derived from our treatments

we observed reduced proportions of myeloid cells in the spleen, particularly when combining CMP

and niraparib (*Supplementary Figure 7b*). Other immune cells such as B cells and natural killer

cells were quantified as well (*Supplementary Figure 7c and d, respectively*). Specifically in T cells

(**Figure 7g**), we observed increased infiltration of T CD8+ lymphocyte in tumor-bearing mice

treated with CMP and this infiltration was significantly augmented using the CMP/niraparib

combination (**Figure 7h**). This correlates with decreased tumor progression and the beneficial

effect of tumoral infiltration of T CD8+ lymphocyte. [90] Likewise, the expression level of the zeta

chain of the TCR (T cell receptor) was increased in lymphocytes (**Figure 7i**). This chain is usually

decreased in different types of cancer, [91] including ovarian cancer. [92] This decrease is associated

with reduced activity of lymphocytes and therefore, is considered a mechanism of tumor immune
evasion. In the spleen a T CD4+ population of lymphocytes expressing PD-1 was significantly
increased (**Figure 7j**) meanwhile the T CD8+ lymphocytes expressing PD-1 showed a trend to be
reduced (**Figure 7k**). This suggests that these lymphocytes, at least in the spleen, the activation
status of these cells is different, where it seems that T CD4+ cells are more active. Also, to mount
a proper immune response, T CD8+ lymphocytes need the help of T CD4+ cells. [92] Then, is
feasibly that the expression of the zeta chain of the TCR (in lymphocytes of the spleen) is a key
element contributing to the help provided by T CD4+ to T CD8+ lymphocytes.

Collectively, our results suggest that the combined CMP/niraparib treatments select specific
population of tumor cells characterized by co-expression of PLD-1/PD-1 and expression of MHC-
ii. These selected cells appear to be less efficient in maintaining a systemic immunosuppression
state. Consequently, fewer myeloid cells accumulate in the spleen, which hinders their ability to
inhibit the function of lymphocytes. Furthermore, these lymphocytes (localized in the spleen)
express an increased level of zeta chain (of the TCR), making them less susceptible to myeloid
cell inhibition.

In summary, the CMP/niraparib treatment seems to influence the selection of a distinct
composition of tumor cells within the tumor, marked by co-expression of PDL-1 and PD-1, along
with MHC-ii. This selected population demonstrated reduced efficacy in inducing systemic
immune suppression, particularly evident at the level of spleen and myeloid infiltration. Moreover,
these tumor cells show a diminish ability to inhibit the expression of the zeta chain of the TCR in
lymphocytes, rendering them more susceptible to recognition by immune cells.

A notable consideration is that the reduced presence of myeloid cells in the spleen, associated with
decreased tolerance to cancer, coincides with heightened zeta chain expression in lymphocytes
withing this secondary lymphoid organ. In contrast, these lymphocytes, particularly T CD4+
lymphocytes are likely to cooperate more effectively with T CD8+ lymphocytes, facilitating their
infiltration into the tumor (*Supplementary Figure 6e*).

The claudin-4 expression and function are highly linked with LAT1, which is associated with transport of amino acids and decreased survival of patients with HGSC.

Here we show that the claudin-4 association with genomic instability in HGSC is intimately related to its role in cell cycle progression and modulation of genomic instability through autophagy regulation mediated by the transporters of amino acids, SLC1A5/LAT1. Furthermore, targeting claudin-4 modifies the tumor microenvironment resulting in decreased tumor progression and tumor immune evasion. To explore the implications of our *in-vitro* and *in-vivo* findings in HGSC tumor, we looked for data mining. First, we construct the protein-protein network of the claudin-4-interacting proteins we reported previously in OVCAR3 cells. [55] From such a network, we selected the proteins whose interaction is experimentally reported and then constructed another protein-protein network to select proteins with interacting reported experimentally. Together, these selected proteins were analyzed in publicly available data sets of HGSC tumors (cBioPortal). We selected those proteins that show higher mutation rates, the top for highly mutated proteins were BRD4, CCDC130, PTK2, and NDRG1. Afterwards, we included these proteins with those from our BioID results (whose interaction with claudin-4 was reported experimentally) to generate clusters of protein-protein networks (as indicated above) and look for biological processes. The distinctively associated cellular processes with such clusters were regulation of cell-cell junctions and cytoskeleton dynamics, and transport of amino acids through the plasma membrane (**Figure 8a**). Thus, it is highly possible that those cellular mechanisms are key drivers of the correlation of claudin-4 upregulation and a worse patient outcome in HGSC. This notion is supported by our findings regarding the correlation of expression, of the elements forming every functional cluster with claudin-4, with aggressiveness of HGSC tumors and decrease patient survival (**Figure 8b**). Likewise, we used these clusters (all, cluster 4, and cluster 4) to explore if they were correlated with aggressiveness in other types of cancer. We found that all clusters are associated with worse patient outcomes in breast and lung cancer but not in stomach. Notably, cluster 4 correlated with decreased RFS (relapse-free survival) and OS (overall survival) in breast and lung cancer, respectively (**Figure 8c**).

DISCUSSION.

Genome instability is a hallmark of cancer, indicating that tumor cells have cellular mechanisms of tolerance to genomic instability. However, it is not well-know how tumor cells limit the accumulation of such modifications to avoid catastrophic chromosomic alterations and cell death. Here we show that claudin-4, traditionally described as a tight junctional protein, mediates a mechanism of tolerance to genomic instability in HGSC cells. This mechanism is a dual protective cellular mechanism that allows HGSC cells proper transit throughout the cell cycle phases and modulation of the accumulation of genomic instability via induction of autophagy.

Briefly, claudin-4's biological role in cells includes the stabilization of the actin-cytoskeleton, particularly the F actin at the cell-cell junctions and perinuclear regions. All of these impacts the nuclear shape and cell cycle progression, and therefore, the modulation of genomic instability in HGSC cells.

Interestingly, micronuclei generated in HGSC cells are modulated by the function of claudin-4. These forms of genetic alterations are typically associated with the cGAS-STING activation, [52] but we show that they are also intimately related to autophagy. Consequently, in spite that micronuclei generated in HGSC cells can collapsed due to disruption of the nuclear envelope, the possibility of releasing the DNA contained these micronuclei into the cytoplasm is reduced because of autophagy induction, which can prevent micronuclei accumulation. [93]

Furthermore, we identified a novel functional link of claudin-4 with the transporters of amino acids, SLC1A5 and LAT1 which regulate a bidirectional transport system of amino acids that control autophagy activation.[65] Likewise, we found that such a functional link, particularly between claudin-4 and LAT1, strongly suggest that one key cellular process driving aggressiveness in ovarian cancer in the transport of amino acids through the plasma membrane. This is interestingly because formation of ascites in common in advanced stage of ovarian cancer and ascites play a fundamental role in immune regulation via amino acids and their transport [94-96]

Remarkably, *in vivo* targeting of claudin-4 not only affected tumor progression but also such targeting, improved the efficacy of niraparib treatment for HGSC tumors. This effect was related to immune modulation of some tumor clones. For example, tumor cells with double expression of PD-1 and PDL-1 were selected due to treatment. Likewise, some cells from the HGSC tumor expressed MHC-11 but this was modulated differentially before disruption of claudin-4 and niraparib treatment. Thus, suggesting a that modulation of MHC-ii has a biological relevance for ovarian cancer progression. Also, we observed more infiltration of T CD8 lymphocytes in tumor-bearing mice treated with CMP/niraparib. This is consistent with the reported benefit of such infiltration in ovarian cancer. [90] Interestingly, we also identified that the zeta chain of the TCR (T cell receptor) was increased when tumor cells received treatment (CMP, niraparib, and combination), particularly in T CD8+ lymphocytes localized in the spleen. Down regulation of this chain is associated with a mechanism of tumor immune evasion in different types of cancer, [91] including in ovarian cancer. Here, it is reported that downregulation of that chain is due to a

secreted factor of 14 kDa. [92] Therefore, increased zeta chain expression in the TCR suggests

increased activity of lymphocytes in cancer.

Finally, given the translational nature of the *in vivo* system we used (the humanized mouse model),

our data strongly suggests that claudin-4 can be a target of inhibition with CMP to increase the

efficacy of the PARPi, niraparib in HGSC patients. Also, the functional link of claudin-4 with

SLC1A5/LAT1, particularly LAT1 suggests that targeting claudin-4 could be a way to target

indirectly the function of LAT1. Perhaps by modulating the localization of these transporters of

amino acids in plasma membrane by stabilizing cell-cell junctions with forskolin [97-99]

METHODS.

Cell lines and viral vectors.

Human derived cells, OVCA429 (RRID:CVCL_3936), OVCAR3 (RRID:CVCL_DH37),

OVCAR8 (RRID:CVCL_1629) were cultured in RPMI-1640 medium (Gibco, Thermo Fisher

Scientific, Cat. #11875) plus 10% heat-inactivated fetal bovine serum (Phoenix Scientific, Cat. #

PS-100, Lot # 20055-01-01) and 1% penicillin/streptomycin (Corning, Cat. #30-002-CI) at 37°C

and 5% CO2. HEK293-FT (RRID:CVCL_6911) were culture similarly but in DMEM medium

(Gibco, Thermo Fisher Scientific, Cat. #11995040). pCDH-EF1a-mCherry-EGFP-LC3B (RRID:

Addgene_170446); pLenti-Lifeact-tdTomato (RRID: Addgene_64048)

Inhibition of claudin-4 expression by CRISPRi.

Stable transfectant OVCA429 and OVCAR3-dCas9 cells were generated by lipocomplexes (Lipofectamine 2000, ThermoFisher, cat: 11668-019; vector pB-CAGGS-dCas9-KRAB-MeCP2, Add gene: 110824) following the manufacturer's instructions. Selection was carried out by antibiotic resistance (blasticidin). A gRNA (forward: GCTGGCTTGCGCATCAGGAC; reverse: GTCCTGATGCGCAAGCCAGC) specific for human *CLDN4* (claudin-4) was generated in the Broad Institute portal (https://portals.broadinstitute.org/gppx/crispick/public). Subsequently, the pCRISPRia-v2_base (TagBFP) vector was digested (BstXI and BlpI) to insert the gRNA by ligation, followed by cloning using *E. coli* (Stbl3™, ThermoFisher, cat: C737303), and selection by flow cytometry (MoFlo XDP100; CU Cancer Center Flow Cytometry Shared Resource).

Lentivirus production and transduction.

293FT cells were transfected using lipocomplexes (Lipofectamine 2000, ThermoFisher, cat: 11668-019) containing the viral packaging system of second generation as well as the lentiviral construct of interest pCDH-EF1a-mCherry-EGFP-LC3B (RRID:Addgene_170446), respectively. Supernatant of transfected 293 ft cells was collected, filtered (0.45 µm), used or stored (-80°C).

Immunoblot and CMP synthesis.

To analyze levels of claudin-4 protein expression, tumor cells were scraped from culture plates in the presence of lysis buffer (30 mM Tris HCl pH7.4, 150 mM NaCl, 1% TritonX-100, 10% glycerol, 2 mM EDTA, 0.57 mM PMSF, 1X cOmplete™ Protease Inhibitor Cocktail), placed on a shaker for 10 minutes and spun at 13,000 rpm for 10 minutes. Protein was separated by SDS-PAGE and transferred to PVDF membrane using the TransBlot Turbo (BioRad). Membranes were blocked with Intercept Blocking Buffer (LI-COR, #927-60001) for 2 hour at room temperature. Mouse anti-human claudin-4 (Thermo Fisher Scientific Cat# 32-9400, RRID:AB_2533096, 1:500), rabbit anti-human β-actin (Abcam Cat# ab6276, RRID:AB_2223210,1:10,000), primary

antibody incubation was performed overnight at 4 °C. Membranes were washed 3 times for 5 minutes each in TBST (50 mM Tris pH 7.5, 150 mM NaCl, 0.1% Tween-20), followed by secondary antibodies for two hour at room temperature. Membranes were washed again 5 times for 5 minutes each in TBST. For fluorescent detection, bands were visualized using the LI-COR Odyssey Imaging System. CMP was synthesized as previously reported[24]

Cell cycle analysis by flow cytometry. $3x10^5$ cells were seeded onto 6 well-plate, the next day cells were washed (sterile PBS 1x) and changed media. After 24h and 48h incubation cells were detached (trypsin 0.25mM), centrifuged (1800 rpm/5min) and washed with PBS followed by centrifugation. PBS was discarded and cells were fixed using cold ethanol 70% (ethanol, milliQ water, v/v) for 30min (4°C) followed by centrifugation (1200 rpm/5min/4°C). Afterwards, cells were washed twice with PBS and the PBS was discarded after centrifugation (1200 rpm/5min/4°C). Cells were treated with RNAse A (50µL/100µM) for 30min at RT and then stained with propidium iodide 300µL (50µM). Analysis were carried out in the Cancer Center Flow Cytometry Shared Resource, University of Colorado Anschutz Medical Campus

Immunofluorescence. Cells were fixed with paraformaldehyde at 4% (PBS 1X) for 10 minutes, followed by permeabilization (30 minutes, 0.1% Triton X-100, PBS 1X). Blocking was carried out by 2h incubation with BSA at 5% (PBS 1X, room temperature, RT, shaking). Primary antibodies (mouse, anti-claudin-4, ThermoFisher, cat: 32-9400, at 1:200 dilution; rabbit, anti-GM130, Cell signaling, cat: 12480, at 1:1600 dilution) were incubated (BSA at 2%, PBS 1X) overnight at 4 °C/shaking. Secondary antibodies (AlexaFluor546 anti-mouse, ThermoFisher, cat: A-11030, at 2 µg/mL; AlexaFluor647 anti-rabbit, ThermoFisher, cat: A32733, at 2 µg/mL) were incubated 2h/shaking at RT (BSA at 2%, PBS 1X). Nuclei were stained with DAPI at 1 µg/mL (PBS 1X) for 10 minutes. All microscopy acquisition was carried out in the Neurotechnology Center, University of Colorado Anschutz Medical Campus.

Generation of HIS-BRGS mice and chimerism evaluation.

Stem Cell Isolation. HSCs were isolated from PBMCs prepared from clinically-rejected CB units from the University of Colorado Cord Blood bank (Clinimmune Labs, Aurora, CO) using CD34+ magnetic Miltenyi beads, and expanded in short-term cultures with IL6 (10 ng/ml), SCF (40 ng/ml) and FLt3L (20 ng/ml). CD34+ cells, harvested between days 4-6, were frozen in 90% FCS/10%DMSO and stored at -80°C prior to injection into neonate pups. Investigators were

blinded from donor identities and the studies were performed in compliance with University of Colorado Institutional Review boards (COMIRB#16-0541).

HIS-BRGS mice and chimerism evaluation. To generate human immune system mice, neonatal (d1-3) BRGS (BALB/c$Rag2^{null}IL2Rg^{null}Sirpa^{NOD}$) pups, obtained from the laboratory of James Di Santo, were irradiated with 300 rads 2-6 hrs prior to injection with 0.2-0.6×10^6 expanded then thawed CD34+ cells. The number of cells injected is equivalent to 50,000 fresh CD34+ cells per mouse, i.e. cell count prior to *in vitro* expansion. Mice were bred and engrafted in the University of Colorado Denver Anschutz Medical Campus vivarium with prior Institutional Animal Care and Use Committee (IACUC) protocol and in a facility accredited by the American Association for Accreditation of Laboratory Animal Care. BRGS mice, both breeders and engrafted, were maintained on an alternating bi-weekly Septra-enriched (Uniprim, Harlan) diet. Mice were injected in the facial vein, liver, or both. Humanized mice were generated in the Pre-clinical Human Immune System Mouse model (PHISM) Shared Resource, University of Colorado Anschutz Medical Campus.

All mouse work was performed in accordance with the Guide for the Care and Use of Laboratory Animals and were approved of by the University of Colorado's Institutional Animal Care and Use Committee (IACUC protocol #283). Humanized Immune System (HIS) mice were i.p. injected with 5x106 cells of a developed patient-derived xenograft (PDX) model, PDX GTFB 1016, described previously. [68] Briefly, this model is derived from a primary tumor collected from a patient diagnosed with stage IIIC , who was described to have a high volume of ascites, peritoneal carcinomatosis, and was chemonaïve at the time of sample collection. TP53 and BRCA2 mutations were identified, and we subsequently transfected the cells with a GFP-luciferase tag, enabling tumor tracking by In Vivo Imaging (IVIS, Perkin Elmer), as described previously. [100,101] Tumors were allowed to establish for three weeks prior to treatment initiation. After this time, mice were treated with either Niraparib (50 mg/kg, PO) five day per week followed by two consecutive days off or CMP (4 mg/kg, i.p.) every two days. Control mice were treated with respective vehicles on the same treatment schedule (5% DMSO, 30% 300PEG, PO [Niraparib vehicle] and PBS, i.p. [CMP vehicle]). Treatment occurred for 28 days, and mice were IVIS scanned weekly to assess tumor development. One day after the last treatment, mice were euthanized via CO2 inhalation and cervical dislocation, and tissues were collected.

Flow cytometry. Blood, spleen and tumor single cell suspensions were stained, as described

previously,[66] with fluorescently-labeled Abs as listed in Supplemental Table X. Blood samples for

chimerism assessment were collected on a Biorad Yeti 5-laser flow cytometer, and spleen and

tumor harvest samples were collected on the Cytek Aurora 5-laser spectral cytometer at the

University of Colorado Cancer Center Flow Cytometry Shared Resource. All data were analyzed

using FlowJo software.

Statistical Considerations. ImageJ (NIH) and Prism software (v9.0) were used for microscopy

and statistical data analysis, respectively. 3 biologically independent experiments were conducted.

Unpaired t and Mann–Whitney tests; Kruskal–Wallis test and one-way ANOVA with Dunn's or

Tukey's multiple comparisons test, respect. Based on normal data distribution, and number of

variables. The level of significance was $p < 0.05$.

Figure legends.

Figure 1. Claudin-4 accounts for intratumoral diversity through moderation of cell cycle and

genomic instability in HGSC cells. Intratumoral diversity was explored by evaluating the transit

of different HGSC cells through cell cycle. Cells were quantified and seeded without any stimuli

followed by staining with propidium iodide (at 24h and 48h) and flow cytometry analysis. The

association of claudin-4 with genomic instability was analyzed in human tumors (TCGA) and *in*

vitro by treating cells with CMP (400μM) for 48h followed by single cell analysis of fixed (stained

with dapi and phalloidin to mark genomic DNA). **(a)** Left, graphs show WT tumor cells in different

phases of the cell cycle (green symbol at 24h; red symbol at 48h) which also changes as the time

went by; right, drawing highlighting diversity of tumor cells withing the tumor, indicated as

localization in different phases of cell cycle. Also, it highlights differences in doubling time among

HGSC cells. **(b)** Left, representative histograms of cell cycle phases before claudin-4 over-

expression; right, percentages of cells in every cell cycle progression. Similar data is indicated

when claudin-4 expression was downregulated in OVCA429 **(c)** and OVCAR3 **(d)**. (4 independent

experiments; Two-tailed Unpaired t test). **(e),** correlation of genomic instability (indicated as % of

altered chromosomic copy numbers) in human ovarian tumors and claudin-4 expression. **(f),** top,

confocal images showing (maximum projections) genomic instability (indicated as nuclei size,

bottom: corresponding quantification) associated with claudin-4 overexpression or

downregulation in OVCA429 **(g)** and OVCAR3 **(h)**. (n= OVCAR8, 1711 cells; OVCA429, 2630

cells; Two-tailed Mann Whitney test, Kruskal-Wallis test with Dunn's multiple comparisons). (3

independent experiments; significance, $p<0.05$. Graphs show mean and SEM (standard error of

the mean), scale bar 5µm.

Figure 2. Claudin-4 stabilizes perinuclear and junctional F-actin in HGSC cells. The

association of claudin-4 with the actin cytoskeleton was evaluated in fixed and living cells. HGSC

cells were treated with CMP (400µM) for 48h and stained the actin-cytoskeleton with phalloidin.

Also, HGSC cells were transduced to express LifeAtc (to mark actin in living cells). Then, we

carried out a morphometric characterization. **(a)**, **(b)**, and **(c)** show reconstructions of perinuclear

F-actin and genomic DNA (from confocal z-stacks) and corresponding quantification in OVCAR8,

OVCA429, and OVCAR3 cells, respectively. (n= OVCAR8, 1711 cells; OVCA429, 2630 cells;

OVCAR3, 2365 cells; Two-tailed Mann Whitney test, Kruskal-Wallis test with Dunn's multiple

comparisons). **(d)** and **(e)**, left, quantification of junctional F-actin from reconstructions (from

confocal z-stacks); right, confocal images (maximum projection) and zoom, followed by

reconstruction of selected regions of interest, ROIs (at junctional F-actin) from OVCA429 cells

and OVCAR3 cells (bottom), respectively (n= OVCA429, 783 cells; OVCAR3, 825 cells;

Kruskal-Wallis test with Dunn's multiple comparisons). **(f),** kymographs generated from different

ROIs (at junctional F-actin) from confocal live-cell imaging using transduced OVCA429 cells

(n=142) with LifeAct to mark F-actin (without any stimuli and cultured for 24h) and OVCAR3

cells (bottom) (n=116) (Two-tailed Mann Whitney test). 3 independent experiments; significance, p<0.05. Graphs show mean and SEM (standard error of the mean), scale bar 5µm.

Figure 3. Claudin-4 modifies the structure of nuclear lamina. HGSC cells were treated with CMP (400µM; 48h) and fixed and stained with dapi, lamin B1 and lamin A/C. Characterization of nuclear lamina was carried out by confocal microscopy. **(a)** Left, frequency of micronuclei identification before claudin-4 over-expression; right, confocal images (maximum projections) highlighting (dotted squares) nuclei and micronuclei. Similar data is presented for claudin-4 downregulation in OVCA429 **(b)** and OVCAR3 cells **(c)**. (n= 4134 OVCAR8 cells; n=3315 OVCA429 cells; n= 4653 OVCAR3 cells; Two-tailed Mann Whitney test). **(d)** Selected confocal images (OVCAR8 WT cells treated with the vehicle from **(a)**) for the main components of nuclear lamina, lamin B1 and lamin A/C highlighting the lack of nuclear lamina in some micronuclei. Similar data is presented in **(e)** and **(f)** for OVCA429 and OVCAR3 cells, respectively. **(g)** and **(h)** Quantification of lamin B1 and lamin A/C in the nucleus in HGSC cells during claudin-4 downregulation. (two independent experiments; Kruskal-Wallis test with Dunn's multiple comparisons, p<0.05). Graphs show mean and SEM, scale bar 5µm.

Figure 4. Claudin-4 impacts the cGAS-STING/autophagy molecular axis. Evaluation of the effect of claudin-4 on STING activation was performed via immunoblotting and confocal microscopy. Protein lysates were obtained from HGSC cells were treated with CMP (400µM; 48h) or stained for pSTING. **(a)** Drawing indicating the general signaling cascade of STING to activate a type I IFN response. **(b)** Immunoblotting for pSTING and pTBK1 in different HGSC cells. **(c)** Left, confocal microscopy of HGSC cells stained for pSTING and LAMP1; right, line scan from selected regions (dotted squares) showing similar patter of intracellular distribution of lamp1 and pSTING. **(d)** Left, drawing highlighting the association of STING with autophagy (LC3 marker of autophagy), independent of pTBK1; right, immunoblotting for LC3 in different HGSC cells. **(e)** immunoblotting for LC3 using lysates of OVCA429 cells treated with inhibitor (CQ, 40µM) or activator (rapamycin, 8µM) of autophagy. **(f)**, **(g)**, and **(h)** are immunoblotting for a downstream target of mTORC1, p70 S6 kinase (phospho-p70 S6 kinase) in different HGCS cells. Scale bar 10µm.

Figure 5. Claudin-4 modulates autophagy which is associated with micronuclei in HGSC cell lines and HGSC tumors. We analyzed the functional role of claudin-4 to link autophagy and micronuclei in HGSC cells, *in vitro* (HGSC-GFP-mCherry-LC3 expressing cells) and *in vivo* (patient derived xenograft in a humanized mice model) by flow cytometry, confocal microscopy, and multispectral immunofluorescence, IF. **(a)**, **(b)**, and **(c)** show percentages of HGSC cells with autophagy flux (by flow cytometry) before CMP treatment (400µM) and claudin-4 genetic manipulation (4 independent experiments). **(d)**, **(e)**, and **(f)** are confocal images (maximum projections) of HGSC-GFP-mCherry-LC3 expressing cells and stained with dapi (white arrow indicates the association of autophagy flux with micronuclei. **(g)** Ascites was obtained from tumor-bearing mice treated or not with CMP and prepared in histogel, followed paraffin embedding and multispectral staining. Representative multispectral IF images (all markers used are indicated in the blue-light box) are shown, where a region of interest (white square) is amplified to show specific markers (white arrow are highlighting the overlay of nuclei and the marker of autophagy, LC3 A/B). **(h)** and **(i)** indicate selected (+2) H-score quantification corresponding to percentage of positive nuclei with LC3 A/B, and nuclei with claudin-4 and LC3 A/B, respectively. (significance, p<0.05). Graphs show mean and SEM, scale bar 10µm and 50µm.

Figure 6. Claudin-4 forms a regulatory axis with SLC1A5/LAT1 to modulate autophagy in HGSC tumor cells. We studied the reported regulation of autophagy by SLC1A5/LAT1 and its association with claudin-4 in different HGSC cells. Modulation of claudin-4 function was associated with the activity of SLC1A5 and LAT1 using specific inhibitors and by limiting L-glutamine availability (a key regulator of the regulatory activity of SLC1A5/LAT1 on autophagy) all of which was studied by immunoblotting, confocal microscopy, and flow cytometry. **(a)** left, confocal images (maximum projections) showing the intracellular distribution of SLC1A5 and claudin-4 in HGSC cells; right, colocalization of SLC1A5 and claudin-4. **(b)** left, confocal images (maximum projections) showing the intracellular distribution of LAT1 and claudin-4 in HGSC cells; right, colocalization of LAT1 and claudin-4. **(c)** Shows representative immunoblots for LAT1 in HGSC cells before claudin-4 disruption (4 independent experiments). **(d)** Model of a

functional association of claudin-4 with the bidirectional transport system formed by SLC1A5/LAT1 (adapted from [65]) which propose that such association is important for proper transport of amino acids in HGSC cells. **(e)** Percentage of HGSC cells with autophagic flux before L-glutamine withdrawal and measured by flow cytometry. **(f)** and **(g)** show representative confocal images (maximum projections) of the intracellular distribution of SLC1A5 before claudin-4 overexpression (OVCAR8 claudin-4) and downregulation (OVCA429 KD and OVCAR3 KD) in HGSC cells before L-glutamine withdrawal. **(h)** Graphs indicate percentages of cells with autophagy flux when treated with specific inhibitor of SLC1A5 activity (GNPA, 5mM). **(i)** Graphs indicate percentages of cells with autophagy flux when treated with specific inhibitor of LAT1 activity (BCH, 25mM). (Two-tailed Unpaired t test and Mann Whitney test; Kruskal-Wallis test with Dunn's multiple comparisons; One way ANOVA and Tukey's multiple comparison test; 3 independent experiments; significance, $p < 0.05$). Graphs show mean and SEM, scale bar 20µm.

Figure 7. Targeting claudin-4 *in vivo* with CMP increases tumor-infiltrating T CD8+ lymphocytes and improves the therapy efficacy of niraparib in HGSC. We generated humanized mice and implanted a patient derived xenograft. After tumor establishment, mice were treated with CMP (4mg/kg), niraparib (Nira, 50mg/kg), and combination of CMP and niraparib (combo, 4mg/kg; 50mg/kg, respectively). At the end of treatment mice were sacrificed and we analyzed tumors and immune cells by flow cytometry (from *in vivo* experiment #2). **(a)** left, verification of tumor establishment by IVIS scan (before treatment); center, comparison of tumor size before treatment (day 0) *vs* after treatment (day 27); right, follow-up of tumor progression during therapy (indicated as bioluminescence signal over time; red color indicates high tumor growth in the IVIS scan). **(b)** left, live tumor cells recover after treatments; right, drawing highlighting that EpCAM protein was used as a tumor marker and red arrows indicate a gate from this marker. Also, PD-1, PDL-1 and MHC-ii with different color indicate that there are clones of tumors cells expressing these proteins **(c)** and **(d)** show percentage of live tumor cells positive for MHC-I and II, respectively. **(e)** left, percentage of live PDL-1+ tumor cells; right, corresponding protein expression by MFI (median fluorescence intensity). **(f)** left, percentage of live PDL-1+ tumor cells with PDL-1 expression (PDL-1+/PDL-+); right, corresponding PD1 expression by MFI. **(g)** Prevalence of T cells (CD3+) in tumor and spleen. **(h)** top, frequency of T CD4+ cells;

bottom, frequency of T CD8+ cells. **(i)** top, z chain expression (MFI) on T CD4+ cells; bottom, z chain expression (MFI) on T CD8+ cells. **(j)** left, quantification of T CD4+ PD-1+ cells; and right, PD-1 expression (MFI). **(k)** left, numbers of T CD8+ PD-1+ cells; and right, PD-1 expression (MFI). Graphs show mean (every symbol represents data from a different mouse) (n=Control, 4 mice; CMP, 4 mice; niraparib, 4 mice; niraparib/CMP, 5 mice; Brown-Forsythe ANOVA test and Unpaired t with Welch's correction; $p < 0.05$).

Figure 8. A protein cluster formed by claudin-4 associates with transport of amino acids and worse patient outcome in HGSC patient survival and possibly other cancer types. Reported claudin-4-interacting proteins were used to generate protein-protein interacting networks to find key elements and cellular functions associated with claudin-4 upregulation in HGSC tumors and publicly available data from HGSC tumor (cBioPortal and Kaplan-Meier plotter). **(a)** Protein-protein network (based on BioID) of claudin-4 interacting proteins (experimentally determined using STRING) and highly mutated proteins (cBioPortal) in HGSC tumors shows clusters of proteins and distinctive associated cellular functions. **(b)** Median survival of HGSC patients (criteria: p53 mutated) correlated with every cluster (based on clausin-4) identified (continue lines) and without claudin-4 (dotted lines) (p values and false discovery rate, FDR in indicated in *Supplementary table 2 and 3*). **(c)** Correlation analysis of relapse-free survival (RFS) in breast cancer and overall survival (OS) in lung and stomach cancer with identified claudin-4 clusters (all, cluster 4 and cluster 3).

ACKNOWLEDGEMENTS

We acknowledge philanthropic contributions from D. Thomas and Kay L. Dunton Endowed Chair in Ovarian Cancer Research, the McClintock-Addlesperger Family, Karen M. Jennison, Don and Arlene Mohler Johnson Family, Michael Intagliata, Duane and Denise Suess, Mary Normandin, and Donald Engelstad.

DECLARATIONS.

Ethical Approval

Stem Cell Isolation was performed in compliance with the University of Colorado Institutional Review boards (COMIRB#16-0541). All mouse work was performed in accordance with the Guide for the Care and Use of Laboratory Animals and was approved by the University of Colorado's Institutional Animal Care and Use Committee (IACUC protocol #283). The University

831   of Colorado has an Institutional Review Board approved protocol (COMIRB#07-935). Reference:
832   [68]

Competing interests.

The authors declare no competing interests.

Autor's contribution

BB and MN conceived the overall project. BB supervised the research project. FRV performed the *in vitro* experiments and data analysis along with PW. JL generated humanized mice and performed flow cytometry and analysis. EW and FRV carried out *in vivo* treatments, tumor burden and ascites processing. FRV wrote the manuscript with the contribution of all authors.

Funding.

This work was supported by grants from the Ovarian Cancer Research Alliance (BGB: Collaborative Award), the Department of Defense (BGB: OC170228, OC200302, OC200225), the NIH/NCI (BGB, R37CA261987), and the American Cancer Society (BGB: 134106-RSG-19-129-01-DDC), as well as the University of Colorado Chancellor's Discovery Fund, OB-GYN Academic Enrichment Fund, and Gynecologic Oncology Academic Enrichment Fund. This study utilized University of Colorado Cancer Center shared resources, which are supported in part by the National Cancer Institute through Cancer Center Support Grant P30CA046934. FRV was supported by the 2022 Outside-the-Box Grant (HERA award), from HERA Ovarian Cancer Foundation.

Data availability statement.

Data generated in this study are included in this manuscript and in its supplementary information. Datamining and flow cytometry raw data are available upon request to the corresponding author.

Bibliographic references

1. Ippolito, M.R., Martis, V., Martin, S., Tijhuis, A.E., Hong, C., Wardenaar, R., Dumont, M., Zerbib, J.,
Spierings, D.C.J., Fachinetti, D., et al. (2021). Gene copy-number changes and chromosomal
instability induced by aneuploidy confer resistance to chemotherapy. Developmental cell *56*,
2440-2454 e2446. 10.1016/j.devcel.2021.07.006.
2. Ortiz, M., Wabel, E., Mitchell, K., and Horibata, S. (2022). Mechanisms of chemotherapy resistance
in ovarian cancer. Cancer Drug Resist *5*, 304-316. 10.20517/cdr.2021.147.
3. Lukow, D.A., Sausville, E.L., Suri, P., Chunduri, N.K., Wieland, A., Leu, J., Smith, J.C., Girish, V.,
Kumar, A.A., Kendall, J., et al. (2021). Chromosomal instability accelerates the evolution of
resistance to anti-cancer therapies. Dev Cell *56*, 2427-2439 e2424. 10.1016/j.devcel.2021.07.009.
4. Morden, C.R., Farrell, A.C., Sliwowski, M., Lichtensztejn, Z., Altman, A.D., Nachtigal, M.W., and
McManus, K.J. (2021). Chromosome instability is prevalent and dynamic in high-grade serous
ovarian cancer patient samples. Gynecol Oncol *161*, 769-778. 10.1016/j.ygyno.2021.02.038.
5. Vazquez-Garcia, I., Uhlitz, F., Ceglia, N., Lim, J.L.P., Wu, M., Mohibullah, N., Niyazov, J., Ruiz, A.E.B.,
Boehm, K.M., Bojilova, V., et al. (2022). Ovarian cancer mutational processes drive site-specific
immune evasion. Nature *612*, 778-786. 10.1038/s41586-022-05496-1.
6. Andor, N., Graham, T.A., Jansen, M., Xia, L.C., Aktipis, C.A., Petritsch, C., Ji, H.P., and Maley, C.C.
(2016). Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. Nat
Med *22*, 105-113. 10.1038/nm.3984.

7.   Bielski, C.M., and Taylor, B.S. (2021). Homing in on genomic instability as a therapeutic target in
cancer. Nat Commun *12*, 3663. 10.1038/s41467-021-23965-5.
8.   Sonugur, F.G., and Akbulut, H. (2019). The Role of Tumor Microenvironment in Genomic Instability
of Malignant Tumors. Front Genet *10*, 1063. 10.3389/fgene.2019.01063.
9.   Tubbs, A., and Nussenzweig, A. (2017). Endogenous DNA Damage as a Source of Genomic
Instability in Cancer. Cell *168*, 644-656. 10.1016/j.cell.2017.01.002.
10. Comaills, V., Kabeche, L., Morris, R., Buisson, R., Yu, M., Madden, M.W., LiCausi, J.A., Boukhali, M.,
Tajima, K., Pan, S., et al. (2016). Genomic Instability Is Induced by Persistent Proliferation of Cells
Undergoing   Epithelial-to-Mesenchymal   Transition.   Cell   Rep   *17*,   2632-2647.
10.1016/j.celrep.2016.11.022.
11. Andor, N., Maley, C.C., and Ji, H.P. (2017). Genomic Instability in Cancer: Teetering on the Limit of
Tolerance. Cancer Res *77*, 2179-2185. 10.1158/0008-5472.CAN-16-1553.
12. Yamamoto, T.M., Webb, P.G., Davis, D.M., Baumgartner, H.K., Woodruff, E.R., Guntupalli, S.R.,
Neville, M., Behbakht, K., and Bitler, B.G. (2022). Loss of Claudin-4 Reduces DNA Damage Repair
and Increases Sensitivity to PARP Inhibitors. Molecular cancer therapeutics *21*, 647-657.
10.1158/1535-7163.MCT-21-0827.
13. Zihni, C., Mills, C., Matter, K., and Balda, M.S. (2016). Tight junctions: from simple barriers to
multifunctional molecular gates. Nat Rev Mol Cell Biol *17*, 564-580. 10.1038/nrm.2016.80.
14. Watari, A., Kodaka, M., Matsuhisa, K., Sakamoto, Y., Hisaie, K., Kawashita, N., Takagi, T.,
Yamagishi, Y., Suzuki, H., Tsujino, H., et al. (2017). Identification of claudin-4 binder that
attenuates tight junction barrier function by TR-FRET-based screening assay. Sci Rep *7*, 14514.
10.1038/s41598-017-15108-y.
15. Ma, X., Miao, H., Jing, B., Pan, Q., Zhang, H., Chen, Y., Zhang, D., Liang, Z., Wen, Z., and Li, M.
(2015). Claudin-4 controls the proliferation, apoptosis, migration and in vivo growth of MCF-7
breast cancer cells. Oncol Rep *34*, 681-690. 10.3892/or.2015.4037.
16. Hicks, D.A., Galimanis, C.E., Webb, P.G., Spillman, M.A., Behbakht, K., Neville, M.C., and
Baumgartner, H.K. (2016). Claudin-4 activity in ovarian tumor cell apoptosis resistance and
migration. BMC Cancer *16*, 788. 10.1186/s12885-016-2799-7.
17. Breed, C., Hicks, D.A., Webb, P.G., Galimanis, C.E., Bitler, B.G., Behbakht, K., and Baumgartner,
H.K. (2019). Ovarian Tumor Cell Expression of Claudin-4 Reduces Apoptotic Response to Paclitaxel.
Mol Cancer Res *17*, 741-750. 10.1158/1541-7786.MCR-18-0451.
18. Achimas-Cadariu, P., Kubelac, P., Irimie, A., Berindan-Neagoe, I., and Ruhli, F. (2022). Evolutionary
perspectives, heterogeneity and ovarian cancer: a complicated tale from past to present. Journal
of ovarian research *15*, 67. 10.1186/s13048-022-01004-1.
19. Kinker, G.S., Greenwald, A.C., Tal, R., Orlova, Z., Cuoco, M.S., McFarland, J.M., Warren, A.,
Rodman, C., Roth, J.A., Bender, S.A., et al. (2020). Pan-cancer single-cell RNA-seq identifies
recurring programs of cellular heterogeneity. Nature genetics *52*, 1208-1218. 10.1038/s41588-
020-00726-6.
20. Cancer Genome Atlas Research, N. (2011). Integrated genomic analyses of ovarian carcinoma.
Nature *474*, 609-615. 10.1038/nature10166.
21. Zheng, D., Mao, Y., Gao, Y., He, F., and Ma, J. (2023). Daughter cell fate choice instructed
preemptively   by   mother   cells   facing   nutrient   limitation.   iScience   *26*,   107198.
10.1016/j.isci.2023.107198.
22. Kuznetsova, A.Y., Seget, K., Moeller, G.K., de Pagter, M.S., de Roos, J.A., Durrbaum, M., Kuffer, C.,
Muller, S., Zaman, G.J., Kloosterman, W.P., and Storchova, Z. (2015). Chromosomal instability,
tolerance of mitotic errors and multidrug resistance are promoted by tetraploidization in human
cells. Cell Cycle *14*, 2810-2820. 10.1080/15384101.2015.1068482.

23.   Bakhoum, S.F., Silkworth, W.T., Nardi, I.K., Nicholson, J.M., Compton, D.A., and Cimini, D. (2014).
938   The mitotic origin of chromosomal instability. Curr Biol *24*, R148-149. 10.1016/j.cub.2014.01.019.
939   24.   Baumgartner, H.K., Beeman, N., Hodges, R.S., and Neville, M.C. (2011). A D-peptide analog of the
940   second extracellular loop of claudin-3 and -4 leads to mislocalized claudin and cellular apoptosis
941   in mammary epithelial cells. Chem Biol Drug Des *77*, 124-136. 10.1111/j.1747-0285.2010.01061.x.
942   25.   Gehne, N., Lamik, A., Lehmann, M., Haseloff, R.F., Andjelkovic, A.V., and Blasig, I.E. (2017). Cross-
943   over endocytosis of claudins is mediated by interactions via their extracellular loops. PloS one *12*,
944   e0182106. 10.1371/journal.pone.0182106.
945   26.   Piontek, J., Winkler, L., Wolburg, H., Muller, S.L., Zuleger, N., Piehl, C., Wiesner, B., Krause, G., and
946   Blasig, I.E. (2008). Formation of tight junction: determinants of homophilic interaction between
947   classic claudins. FASEB J *22*, 146-158. 10.1096/fj.07-8319com.
948   27.   Lisica, A., Fouchard, J., Kelkar, M., Wyatt, T.P.J., Duque, J., Ndiaye, A.B., Bonfanti, A., Baum, B.,
949   Kabla, A.J., and Charras, G.T. (2022). Tension at intercellular junctions is necessary for accurate
950   orientation of cell division in the epithelium plane. Proc Natl Acad Sci U S A *119*, e2201600119.
951   10.1073/pnas.2201600119.
952   28.   Aguilar-Aragon, M., Bonello, T.T., Bell, G.P., Fletcher, G.C., and Thompson, B.J. (2020). Adherens
953   junction remodelling during mitotic rounding of pseudostratified epithelial cells. EMBO Rep *21*,
954   e49700. 10.15252/embr.201949700.
955   29.   Davidson, P.M., and Cadot, B. (2021). Actin on and around the Nucleus. Trends Cell Biol *31*, 211-
956   223. 10.1016/j.tcb.2020.11.009.
957   30.   Lim, S., Quinton, R.J., and Ganem, N.J. (2016). Nuclear envelope rupture drives genome instability
958   in cancer. Mol Biol Cell *27*, 3210-3213. 10.1091/mbc.E16-02-0098.
959   31.   Garcia, M.A., Nelson, W.J., and Chavez, N. (2018). Cell-Cell Junctions Organize Structural and
960   Signaling Networks. Cold Spring Harbor perspectives in biology *10*. 10.1101/cshperspect.a029181.
961   32.   Maninova, M., and Vomastek, T. (2016). Dorsal stress fibers, transverse actin arcs, and perinuclear
962   actin fibers form an interconnected network that induces nuclear movement in polarizing
963   fibroblasts. FEBS J *283*, 3676-3693. 10.1111/febs.13836.
964   33.   Gundersen, G.G., and Worman, H.J. (2013). Nuclear positioning. Cell *152*, 1376-1389.
965   10.1016/j.cell.2013.02.031.
966   34.   Singh, I., and Lele, T.P. (2022). Nuclear Morphological Abnormalities in Cancer: A Search for
967   Unifying Mechanisms. Results Probl Cell Differ *70*, 443-467. 10.1007/978-3-031-06573-6_16.
968   35.   Zeimet, A.G., Fiegl, H., Goebel, G., Kopp, F., Allasia, C., Reimer, D., Steppan, I., Mueller-Holzner,
969   E., Ehrlich, M., and Marth, C. (2011). DNA ploidy, nuclear size, proliferation index and DNA-
970   hypomethylation in ovarian cancer. Gynecol Oncol *121*, 24-31. 10.1016/j.ygyno.2010.12.332.
971   36.   Jevtic, P., Edens, L.J., Vukovic, L.D., and Levy, D.L. (2014). Sizing and shaping the nucleus:
972   mechanisms and significance. Curr Opin Cell Biol *28*, 16-27. 10.1016/j.ceb.2014.01.003.
973   37.   Efremov, A.K., Hovan, L., and Yan, J. (2022). Nucleus size and its effect on nucleosome stability in
974   living cells. Biophys J *121*, 4189-4204. 10.1016/j.bpj.2022.09.019.
975   38.   Haarer, E.L., Theodore, C.J., Guo, S., Frier, R.B., and Campellone, K.G. (2023). Genomic instability
976   caused by Arp2/3 complex inactivation results in micronucleus biogenesis and cellular
977   senescence. PLoS Genet *19*, e1010045. 10.1371/journal.pgen.1010045.
978   39.   Spichal, M., and Fabre, E. (2017). The Emerging Role of the Cytoskeleton in Chromosome
979   Dynamics. Front Genet *8*, 60. 10.3389/fgene.2017.00060.
980   40.   Hoelzle, M.K., and Svitkina, T. (2012). The cytoskeletal mechanisms of cell-cell junction formation
981   in endothelial cells. Mol Biol Cell *23*, 310-323. 10.1091/mbc.E11-08-0719.
982   41.   Mege, R.M., Gavard, J., and Lambert, M. (2006). Regulation of cell-cell junctions by the
983   cytoskeleton. Curr Opin Cell Biol *18*, 541-548. 10.1016/j.ceb.2006.08.004.

42. Villagomez, F.R., Diaz-Valencia, J.D., Ovalle-Garcia, E., Antillon, A., Ortega-Blake, I., Romero-
Ramirez, H., Cerna-Cortes, J.F., Rosales-Reyes, R., Santos-Argumedo, L., and Patino-Lopez, G.
(2021). TBC1D10C is a cytoskeletal functional linker that modulates cell spreading and
phagocytosis in macrophages. Sci Rep *11*, 20946. 10.1038/s41598-021-00450-z.
43. Almagro, J., Messal, H.A., Elosegui-Artola, A., van Rheenen, J., and Behrens, A. (2022). Tissue
architecture in tumor initiation and progression. Trends Cancer *8*, 494-505.
10.1016/j.trecan.2022.02.007.
44. Esmaeilniakooshkghazi, A., Pham, E., George, S.P., Ahrorov, A., Villagomez, F.R., Byington, M.,
Mukhopadhyay, S., Patnaik, S., Conrad, J.C., Naik, M., et al. (2023). In colon cancer cells fascin1
regulates adherens junction remodeling. FASEB J *37*, e22786. 10.1096/fj.202201454R.
45. Stroud, M.J. (2018). Linker of nucleoskeleton and cytoskeleton complex proteins in
cardiomyopathy. Biophys Rev *10*, 1033-1051. 10.1007/s12551-018-0431-6.
46. Denais, C.M., Gilbert, R.M., Isermann, P., McGregor, A.L., te Lindert, M., Weigelin, B., Davidson,
P.M., Friedl, P., Wolf, K., and Lammerding, J. (2016). Nuclear envelope rupture and repair during
cancer cell migration. Science *352*, 353-358. 10.1126/science.aad7297.
47. Hatch, E.M., and Hetzer, M.W. (2016). Nuclear envelope rupture is induced by actin-based nucleus
confinement. J Cell Biol *215*, 27-36. 10.1083/jcb.201603053.
48. Yang, Z., Maciejowski, J., and de Lange, T. (2017). Nuclear Envelope Rupture Is Enhanced by Loss
of p53 or Rb. Mol Cancer Res *15*, 1579-1586. 10.1158/1541-7786.MCR-17-0084.
49. Utani, K., Okamoto, A., and Shimizu, N. (2011). Generation of micronuclei during interphase by
coupling between cytoplasmic membrane blebbing and nuclear budding. PloS one *6*, e27233.
10.1371/journal.pone.0027233.
50. Shimizu, N., Itoh, N., Utiyama, H., and Wahl, G.M. (1998). Selective entrapment of
extrachromosomally amplified DNA by nuclear budding and micronucleation during S phase. J Cell
Biol *140*, 1307-1320. 10.1083/jcb.140.6.1307.
51. Hatch, E.M., Fischer, A.H., Deerinck, T.J., and Hetzer, M.W. (2013). Catastrophic nuclear envelope
collapse in cancer cell micronuclei. Cell *154*, 47-60. 10.1016/j.cell.2013.06.007.
52. Mackenzie, K.J., Carroll, P., Martin, C.A., Murina, O., Fluteau, A., Simpson, D.J., Olova, N., Sutcliffe,
H., Rainger, J.K., Leitch, A., et al. (2017). cGAS surveillance of micronuclei links genome instability
to innate immunity. Nature *548*, 461-465. 10.1038/nature23449.
53. Shah, P., Wolf, K., and Lammerding, J. (2017). Bursting the Bubble - Nuclear Envelope Rupture as
a Path to Genomic Instability? Trends Cell Biol *27*, 546-555. 10.1016/j.tcb.2017.02.008.
54. Shimi, T., Kittisopikul, M., Tran, J., Goldman, A.E., Adam, S.A., Zheng, Y., Jaqaman, K., and
Goldman, R.D. (2015). Structural organization of nuclear lamins A, C, B1, and B2 revealed by
superresolution microscopy. Mol Biol Cell *26*, 4075-4086. 10.1091/mbc.E15-07-0461.
55. Neville, M.C., Webb, P.G., Baumgartner, H.K., and Bitler, B.G. (2022). Claudin-4 localization in
epithelial ovarian cancer. Heliyon *8*, e10862. 10.1016/j.heliyon.2022.e10862.
56. Winkelman, J.D., Suarez, C., Hocky, G.M., Harker, A.J., Morganthaler, A.N., Christensen, J.R., Voth,
G.A., Bartles, J.R., and Kovar, D.R. (2016). Fascin- and alpha-Actinin-Bundled Networks Contain
Intrinsic Structural Features that Drive Protein Sorting. Curr Biol *26*, 2697-2706.
10.1016/j.cub.2016.07.080.
57. Rello-Varona, S., Lissa, D., Shen, S., Niso-Santano, M., Senovilla, L., Marino, G., Vitale, I., Jemaa,
M., Harper, F., Pierron, G., et al. (2012). Autophagic removal of micronuclei. Cell Cycle *11*, 170-
176. 10.4161/cc.11.1.18564.
58. de Queiroz, N., Xia, T., Konno, H., and Barber, G.N. (2019). Ovarian Cancer Cells Commonly Exhibit
Defective STING Signaling Which Affects Sensitivity to Viral Oncolysis. Molecular cancer research
: MCR *17*, 974-986. 10.1158/1541-7786.MCR-18-0504.

59.   Decout, A., Katz, J.D., Venkatraman, S., and Ablasser, A. (2021). The cGAS-STING pathway as a
1032   therapeutic target in inflammatory diseases. Nat Rev Immunol *21*, 548-569. 10.1038/s41577-021-
1033   00524-z.
1034   60.   Gui, X., Yang, H., Li, T., Tan, X., Shi, P., Li, M., Du, F., and Chen, Z.J. (2019). Autophagy induction via
1035   STING trafficking is a primordial function of the cGAS pathway. Nature *567*, 262-266.
1036   10.1038/s41586-019-1006-9.
1037   61.   Kuchitsu, Y., Mukai, K., Uematsu, R., Takaada, Y., Shinojima, A., Shindo, R., Shoji, T., Hamano, S.,
1038   Ogawa, E., Sato, R., et al. (2023). STING signalling is terminated through ESCRT-dependent
1039   microautophagy of vesicles originating from recycling endosomes. Nat Cell Biol *25*, 453-466.
1040   10.1038/s41556-023-01098-9.
1041   62.   Wheeler, L.J., Watson, Z.L., Qamar, L., Yamamoto, T.M., Sawyer, B.T., Sullivan, K.D., Khanal, S.,
1042   Joshi, M., Ferchaud-Roucher, V., Smith, H., et al. (2019). Multi-Omic Approaches Identify
1043   Metabolic and Autophagy Regulators Important in Ovarian Cancer Dissemination. iScience *19*,
1044   474-491. 10.1016/j.isci.2019.07.049.
1045   63.   He, C. (2022). Balancing nutrient and energy demand and supply via autophagy. Curr Biol *32*,
1046   R684-R696. 10.1016/j.cub.2022.04.071.
1047   64.   Roux, K.J., Kim, D.I., Burke, B., and May, D.G. (2018). BioID: A Screen for Protein-Protein
1048   Interactions. Curr Protoc Protein Sci *91*, 19 23 11-19 23 15. 10.1002/cpps.51.
1049   65.   Nicklin, P., Bergman, P., Zhang, B., Triantafellow, E., Wang, H., Nyfeler, B., Yang, H., Hild, M., Kung,
1050   C., Wilson, C., et al. (2009). Bidirectional transport of amino acids regulates mTOR and autophagy.
1051   Cell *136*, 521-534. 10.1016/j.cell.2008.11.044.
1052   66.   Lang, J., Weiss, N., Freed, B.M., Torres, R.M., and Pelanda, R. (2011). Generation of hematopoietic
1053   humanized mice in the newborn BALB/c-Rag2null Il2rgammanull mouse model: a multivariable
1054   optimization approach. Clin Immunol *140*, 102-116. 10.1016/j.clim.2011.04.002.
1055   67.   Cogels, M.M., Rouas, R., Ghanem, G.E., Martinive, P., Awada, A., Van Gestel, D., and Krayem, M.
1056   (2021). Humanized Mice as a Valuable Pre-Clinical Model for Cancer Immunotherapy Research.
1057   Front Oncol *11*, 784947. 10.3389/fonc.2021.784947.
1058   68.   Sawyer, B.T., Qamar, L., Yamamoto, T.M., McMellen, A., Watson, Z.L., Richer, J.K., Behbakht, K.,
1059   Schlaepfer, I.R., and Bitler, B.G. (2020). Targeting Fatty Acid Oxidation to Promote Anoikis and
1060   Inhibit Ovarian Cancer Progression. Mol Cancer Res *18*, 1088-1098. 10.1158/1541-7786.MCR-19-
1061   1057.
1062   69.   Zhao, Q., Kohut, A., Li, Y.J., Martincuks, A., Austria, T., Zhang, C., Santiago, N.L., Borrero, R.M.,
1063   Phan, X.T., Melstrom, L., et al. (2022). Niraparib-induced STAT3 inhibition increases its antitumor
1064   effects. Front Oncol *12*, 966492. 10.3389/fonc.2022.966492.
1065   70.   Tay, R.E., Richardson, E.K., and Toh, H.C. (2021). Revisiting the role of CD4(+) T cells in cancer
1066   immunotherapy-new insights into old paradigms. Cancer Gene Ther *28*, 5-17. 10.1038/s41417-
1067   020-0183-x.
1068   71.   Axelrod, M.L., Cook, R.S., Johnson, D.B., and Balko, J.M. (2019). Biological Consequences of MHC-
1069   II Expression by Tumor Cells in Cancer. Clin Cancer Res *25*, 2392-2402. 10.1158/1078-0432.CCR-
1070   18-3200.
1071   72.   Schuster, H., Peper, J.K., Bosmuller, H.C., Rohle, K., Backert, L., Bilich, T., Ney, B., Loffler, M.W.,
1072   Kowalewski, D.J., Trautwein, N., et al. (2017). The immunopeptidomic landscape of ovarian
1073   carcinomas. Proc Natl Acad Sci U S A *114*, E9942-E9951. 10.1073/pnas.1707658114.
1074   73.   Tegeler, C.M., Scheid, J., Rammensee, H.G., Salih, H.R., Walz, J.S., Heitmann, J.S., and Nelde, A.
1075   (2022). HLA-DR Presentation of the Tumor Antigen MSLN Associates with Clinical Outcome of
1076   Ovarian Cancer Patients. Cancers (Basel) *14*. 10.3390/cancers14092260.
1077   74.   Johnson, D.B., Estrada, M.V., Salgado, R., Sanchez, V., Doxie, D.B., Opalenik, S.R., Vilgelm, A.E.,
1078   Feld, E., Johnson, A.S., Greenplate, A.R., et al. (2016). Melanoma-specific MHC-II expression

represents a tumour-autonomous phenotype and predicts response to anti-PD-1/PD-L1 therapy.
Nat Commun *7*, 10582. 10.1038/ncomms10582.

75. Forero, A., Li, Y., Chen, D., Grizzle, W.E., Updike, K.L., Merz, N.D., Downs-Kelly, E., Burwell, T.C.,
Vaklavas, C., Buchsbaum, D.J., et al. (2016). Expression of the MHC Class II Pathway in Triple-
Negative Breast Cancer Tumor Cells Is Associated with a Good Prognosis and Infiltrating
Lymphocytes. Cancer Immunol Res *4*, 390-399. 10.1158/2326-6066.CIR-15-0243.

76. Alwosaibai, K., Aalmri, S., Mashhour, M., Ghandorah, S., Alshangiti, A., Azam, F., Selwi, W.,
Gharaibeh, L., Alatawi, Y., Alruwaii, Z., and Alsaab, H.O. (2023). PD-L1 is highly expressed in ovarian
cancer and associated with cancer stem cells populations expressing CD44 and other stem cell
markers. BMC Cancer *23*, 13. 10.1186/s12885-022-10404-x.

77. Sun, C., Mezzadra, R., and Schumacher, T.N. (2018). Regulation and Function of the PD-L1
Checkpoint. Immunity *48*, 434-452. 10.1016/j.immuni.2018.03.014.

78. Zheng, Y., Fang, Y.C., and Li, J. (2019). PD-L1 expression levels on tumor cells affect their
immunosuppressive activity. Oncol Lett *18*, 5399-5407. 10.3892/ol.2019.10903.

79. Wang, X., Yang, X., Zhang, C., Wang, Y., Cheng, T., Duan, L., Tong, Z., Tan, S., Zhang, H., Saw, P.E.,
et al. (2020). Tumor cell-intrinsic PD-1 receptor is a tumor suppressor and mediates resistance to
PD-1 blockade therapy. Proc Natl Acad Sci U S A *117*, 6640-6650. 10.1073/pnas.1921445117.

80. Du, S., McCall, N., Park, K., Guan, Q., Fontina, P., Ertel, A., Zhan, T., Dicker, A.P., and Lu, B. (2018).
Blockade of Tumor-Expressed PD-1 promotes lung cancer growth. Oncoimmunology *7*, e1408747.
10.1080/2162402X.2017.1408747.

81. Kleffel, S., Posch, C., Barthel, S.R., Mueller, H., Schlapbach, C., Guenova, E., Elco, C.P., Lee, N.,
Juneja, V.R., Zhan, Q., et al. (2015). Melanoma Cell-Intrinsic PD-1 Receptor Functions Promote
Tumor Growth. Cell *162*, 1242-1256. 10.1016/j.cell.2015.08.052.

82. Yao, H., Wang, H., Li, C., Fang, J.Y., and Xu, J. (2018). Cancer Cell-Intrinsic PD-1 and Implications in
Combinatorial Immunotherapy. Front Immunol *9*, 1774. 10.3389/fimmu.2018.01774.

83. Hiam-Galvez, K.J., Allen, B.M., and Spitzer, M.H. (2021). Systemic immunity in cancer. Nat Rev
Cancer *21*, 345-359. 10.1038/s41568-021-00347-z.

84. Ribatti, D. (2017). The concept of immune surveillance against tumors. The first theories.
Oncotarget *8*, 7175-7180. 10.18632/oncotarget.12739.

85. Chauveau, A., Pirgova, G., Cheng, H.W., De Martin, A., Zhou, F.Y., Wideman, S., Rittscher, J.,
Ludewig, B., and Arnon, T.I. (2020). Visualization of T Cell Migration in the Spleen Reveals a
Network of Perivascular Pathways that Guide Entry into T Zones. Immunity *52*, 794-807 e797.
10.1016/j.immuni.2020.03.010.

86. Barinov, A., Galgano, A., Krenn, G., Tanchot, C., Vasseur, F., and Rocha, B. (2017).
CD4/CD8/Dendritic cell complexes in the spleen: CD8+ T cells can directly bind CD4+ T cells and
modulate their response. PloS one *12*, e0180644. 10.1371/journal.pone.0180644.

87. Wong, S.B., Bos, R., and Sherman, L.A. (2008). Tumor-specific CD4+ T cells render the tumor
environment permissive for infiltration by low-avidity CD8+ T cells. J Immunol *180*, 3122-3131.
10.4049/jimmunol.180.5.3122.

88. Ugel, S., Peranzoni, E., Desantis, G., Chioda, M., Walter, S., Weinschenk, T., Ochando, J.C.,
Cabrelle, A., Mandruzzato, S., and Bronte, V. (2012). Immune tolerance to tumor antigens occurs
in a specialized environment of the spleen. Cell Rep *2*, 628-639. 10.1016/j.celrep.2012.08.006.

89. Makarenkova, V.P., Bansal, V., Matta, B.M., Perez, L.A., and Ochoa, J.B. (2006). CD11b+/Gr-1+
myeloid suppressor cells cause T cell dysfunction after traumatic stress. J Immunol *176*, 2085-
2094. 10.4049/jimmunol.176.4.2085.

90. Ovarian Tumor Tissue Analysis, C., Goode, E.L., Block, M.S., Kalli, K.R., Vierkant, R.A., Chen, W.,
Fogarty, Z.C., Gentry-Maharaj, A., Toloczko, A., Hein, A., et al. (2017). Dose-Response Association

of CD8+ Tumor-Infiltrating Lymphocytes and Survival Time in High-Grade Serous Ovarian Cancer.
JAMA Oncol *3*, e173290. 10.1001/jamaoncol.2017.3290.

91. Villagomez, F.R., Medina-Contreras, O., Cerna-Cortes, J.F., and Patino-Lopez, G. (2020). The role
of the oncogenic Rab35 in cancer invasion, metastasis, and immune evasion, especially in
leukemia. Small GTPases *11*, 334-345. 10.1080/21541248.2018.1463895.

92. Taylor, D.D., Bender, D.P., Gercel-Taylor, C., Stanson, J., and Whiteside, T.L. (2001). Modulation of
TcR/CD3-zeta chain expression by a circulating factor derived from ovarian cancer patients. Br J
Cancer *84*, 1624-1629. 10.1054/bjoc.2001.1847.

93. Hewitt, G., and Korolchuk, V.I. (2017). Repair, Reuse, Recycle: The Expanding Role of Autophagy
in Genome Maintenance. Trends Cell Biol *27*, 340-351. 10.1016/j.tcb.2016.11.011.

94. Grobben, Y., den Ouden, J.E., Aguado, C., van Altena, A.M., Kraneveld, A.D., and Zaman, G.J.R.
(2023). Amino Acid-Metabolizing Enzymes in Advanced High-Grade Serous Ovarian Cancer
Patients: Value of Ascites as Biomarker Source and Role for IL4I1 and IDO1. Cancers (Basel) *15*.
10.3390/cancers15030893.

95. Gong, Y., Yang, J., Wang, Y., Xue, L., and Wang, J. (2020). Metabolic factors contribute to T-cell
inhibition in the ovarian cancer ascites. Int J Cancer *147*, 1768-1777. 10.1002/ijc.32990.

96. Kaira, K., Nakamura, K., Hirakawa, T., Imai, H., Tominaga, H., Oriuchi, N., Nagamori, S., Kanai, Y.,
Tsukamoto, N., Oyama, T., et al. (2015). Prognostic significance of L-type amino acid transporter
1 (LAT1) expression in patients with ovarian tumors. Am J Transl Res *7*, 1161-1171.

97. Klusmeier, N., Schnittler, H.J., and Seebach, J. (2019). A Novel Microscopic Assay Reveals
Heterogeneous Regulation of Local Endothelial Barrier Function. Biophys J *116*, 1547-1559.
10.1016/j.bpj.2019.02.008.

98. Novak, D., Quiggle, F., and Haafiz, A. (2006). Impact of forskolin and amino acid depletion upon
System A activity and SNAT expression in BeWo cells. Biochimie *88*, 39-44.
10.1016/j.biochi.2005.07.002.

99. Noda, K., Zhang, J., Fukuhara, S., Kunimoto, S., Yoshimura, M., and Mochizuki, N. (2010). Vascular
endothelial-cadherin stabilizes at cell-cell junctions by anchoring to circumferential actin bundles
through alpha- and beta-catenins in cyclic AMP-Epac-Rap1 signal-activated endothelial cells. Mol
Biol Cell *21*, 584-596. 10.1091/mbc.e09-07-0580.

100. Yamamoto, T.M., McMellen, A., Watson, Z.L., Aguilera, J., Ferguson, R., Nurmemmedov, E.,
Thakar, T., Moldovan, G.L., Kim, H., Cittelly, D.M., et al. (2019). Activation of Wnt signaling
promotes olaparib resistant ovarian cancer. Mol Carcinog *58*, 1770-1782. 10.1002/mc.23064.

101. Bitler, B.G., Nicodemus, J.P., Li, H., Cai, Q., Wu, H., Hua, X., Li, T., Birrer, M.J., Godwin, A.K., Cairns,
P., and Zhang, R. (2011). Wnt5a suppresses epithelial ovarian cancer by promoting cellular
senescence. Cancer Res *71*, 6184-6194. 10.1158/0008-5472.CAN-11-1341.

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

Figures.

Figure 1.

1179

1180

1181 **Figure 2.**

1182

1183

1184

1185

1186   **Figure 3.**

1187

1188

1189

1190

1191

Figure 4.

1192

1193

1194

1195

1196   **Figure 5.**

1197   **Figure 6.**

1198

1199

1200   **Figure 7.**

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219   **Figure 8.**

1220

1221

1222   **SUPPLEMENTARY FIGURES.**

1223

1224

1225

Supplementary Figure 1. (a) Immunoblotting to verify the claudin-4 overexpression (top panel) in OVCAR8 cells (which do not express claudin-4) and its downregulation (knockdown using CRISPRi in OVCA429 and OVCAR3 cells) (bottom panel). **(b)** left, drawing highlighting the structure of claudin-4, in red and green colors (where it can be observed two extracellular loops, large and small) as well as *cis* and *trans* interaction at the plasma membrane as a functional regulation. The drawing also highlights a conserved sequence of claudin-4 (DFYNP) located in the second (small) extracellular loop as well as the sequence of CMP (claudin mimic peptide; DFYNP which is in D-conformation and stabilized by Glycine, thus having NH_2 and OH ends flanking the peptide); right, known and predicted phosphosites in claudin-4 (it is highlighted a predicted and uncharacterized phosphosite at Y148, located in the conserved sequence in the small extracellular loop of claudin-4) generated in (http://www.phosphonet.ca/). **(c)** Quantification of cell cycle progression by flow cytometry and propidium iodide-stained OVCAR8 WT cells during normal (RPMI, 10% FBS) culture conditions or starvation (RPMI, 1% FBS). **(d)** Maximum projection of live-cell confocal imaging (xyt) where two different ROIs (region of interest) were used to generate kymographs from junctional F-actin and estimate actin dynamics at multiple time points. **(e)** immunoblotting for lamin B1 and lamin A/C and corresponding quantification relative to loading control (4 independent experiments) in **(f)** and **(g)**, respectively. **(h) Left, t**ransduced OVCA429 WT cells with ISRE vector and treated with cGAPM 1µg/mL (24h) and chloroquine (CQ, 20 and 40µM; 24h) (3 independent experiments; One-way ANOVA and Tukey's multiple comparisons test; $p < 0.5$); pilot test (one independent experiment using OVAC429 WT and OVCA429 KD-ISRE transduced cells; One-way ANOVA and Tukey's multiple comparisons test; $p < 0.5$). Graphs show mean and SEM, scale bar 5µm.

1265

1287

1288

Supplementary Figure 2. (a) Autophagy flux strategy for flow cytometry. **(b)** Confirmation of autophagy flux in HGSC cells (OVCAR8 WT, OVCA429 WT, OVCAR3 WT). CQ, chloroquine (40µM, 24h) was used to show blocking of autophagy; rapamycin, rap (8µM, 24h) was used to show activation of autophagy. Graphs show mean and SEM.

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

Supplementary Figure 3. (a) Phenotypic frequency (from multispectral immunofluorescence; 8 proteins plus dapi) of cells obtained from ascites generated by HGSC primary tumor in humanized mice from ascites samples. (b) Quantification of cells positive for LC3 A/B and pSTING; and LC3 A/B, pSTING plus pTBK1. (c) Mean expression of claudin-4 and LC3 A/B and (d) pSTING/LC3A/B in ascites samples and (from multispectral immunofluorescence). (e) Left, immunoblotting for SLC1A5 in different HGSC cells; right, protein quantification relative to loading control (4 independent experiments). (f) Tumor burden (experiment #1) indicated as bioluminescence signal over time. (Two-way ANOVA and Tukey's multiple comparisons test; p<0.05). Graphs show mean and SEM.

1317

1318

Supplementary Figure 4. (a) Quantification of LAT1 in lysates from HGSC cells relative to loading control (4
independent experiments; One way ANOVA and Tukey's multiple comparisons test; p<0.05). **(b)** left, confocal images
(maximum projections) of HGSC cells cultured without L-glutamine (24h) and stained for pSTING and lamp1; center,
kymographs from selected regions (from z-stacks confocal images); right, line scan of selected regions for lamp1 and
pSTING. Here is highlighted the co-localization of lamp1 with pSTING before L-glutamine withdrawal. Confocal
images (maximum projections) showing SLC1A5 intracellular distribution in HGSC cells before claudin-4
overexpression (OVCAR8-claudin-4) or downregulation (OVCA429 KD and OVCAr3 KD) and autophagy blocking
(c) (using chloroquine, CQ, 40µM, 24h) or autophagy induction **(d)** (using rapamycin, 8µM, 24h). Graphs show mean
and SEM, scale bar 10µm.

1328

1329

Supplementary Figure 5. (a) and **(b)** Experimental design for targeting claudin-4 in primary tumor in a humanized mouse model (two independent experiments; #1, newborn mice were treated to induced development of human immune system; #2, mice without weaning were treated to induced development of human immune system).

1330
1331
1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

Supplementary Figure 6. (a) Percentage of tumor cells with MHC-II that express PDL-1 expression as
well. **(b)** Determination of immune cell populations other than T-cells by flow cytometry. **(c)** and **(d)** Identification
of lymphocytes CD4+ and CD8+ producers of TNF-α and INF-γ by flow cytometry. **(e)** Proposed model of immune
modulation by niraparib and CMP combinatory treatment. Consisting reduced accumulation on myeloid cells in the
spleen (1), increased expression of the TCR-zeta chain, which could favor the lymphocyte activation in the spleen as
well as the cooperation of T CD4+ and T CD8+ lymphocytes to promote lymphocyte infiltration into the tumor, which
show changes in its cellular composition, correlating with reduced tumor growth. **()** quantification of lymphocytes
CD4+ and CD8+ positive for HLA-DR and GrB. **(f)** Identification of lymphocytes CD4+ and CD8+ with expression
of Tim3, FoxP3, and Treg (T regulatory cell) phenotype, respectively (All data derives from gate: live human immune
cells, T cells, CD4+ and CD8+, respectively). Graphs show mean (every symbol represents data from a different

1355   mouse) (n=Control, 4 mice; CMP, 4 mice; niraparib, 4 mice; niraparib/CMP, 5 mice; Brown-Forsythe ANOVA test
1356   and Unpaired t with Welch's correction; $p < 0.05$).