Tmc7 deficiency causes acrosome biogenesis defects and male infertility in mice

Jing Wang[1,2], Yingying Yin[1,3], Lei Yang[1,2], Junchao Qin[1,2], Zixiang Wang[1,2], Chunhong Qiu[1],

Yuan Gao[3], Gang Lu[4], Fei Gao[5], Zi-jiang Chen[3], Xiyu Zhang[1] ✉, Hongbin Liu[3]✉, Zhaojian

Liu[1,2]✉

[1] Key Laboratory of Experimental Teratology, Ministry of Education, Department of Cell

Biology, School of Basic Medical Sciences, Shandong University, Jinan, China; [2] Advanced

Medical Research Institute, Shandong University, Jinan, China; [3] Center for Reproductive

Medicine, Shandong University, Jinan, China. [4] CUHK-SDU Joint Laboratory on

Reproductive Genetics, School of Biomedical Sciences, The Chinese University of Hong

Kong, Hong Kong, China. [5] State Key Laboratory of Stem Cell and Reproductive Biology,

Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.

Correspondence:

Dr. **Zhaojian Liu**, Department of Cell Biology, School of Basic Medical Science, Shandong

University, 44 Wenhua Xi Road, Jinan 250012, China. Email: liujian9782@sdu.edu.cn

Dr. **Hongbin Liu**, Center for Reproductive Medicine, Shandong University, Jinan, China.

Email: hongbin_sduivf@aliyun.com

Dr. **Xiyu Zhang**, Department of Molecular Medicine and Genetics, School of Basic Medical

Sciences, Shandong University School of Medicine, 44 Wenhua Xi Road, Jinan, Shandong

250012, China. Email: xiyuzhang@sdu.edu.cn

Abstract

Transmembrane channel-like (Tmc) proteins are a highly conserved ion channel family consisting of eight members (TMC1–TMC8) in mammals. TMC1/2 are components of the mechanotransduction channel in hair cells, and mutations of TMC1/2 cause deafness in humans and mice. However, the physiological roles of other TMC proteins remain largely unknown. Here, we show that Tmc7 is specifically expressed in the testis and that it is required for acrosome biogenesis during spermatogenesis. $Tmc7^{-/-}$ mice exhibited complete male infertility due to abnormal sperm morphology, similar to human oligo-astheno-teratozoospermia. We further demonstrate that Tmc7 is colocalized with Gm130 at the cis-Golgi region in round spermatids. Tmc7 deficiency leads to aberrant Golgi morphology and impaired fusion of Golgi-derived vesicles to the developing acrosome. Moreover, upon loss of Tmc7 Golgi pH and ion homeostasis is impaired and ROS levels are increased, which in turn causes Golgi and endoplasmic reticulum (ER) stress. Taken together, these results suggest that Tmc7 is required to maintain Golgi pH and ion homeostasis, which is needed for acrosome biogenesis. Our findings unveil a novel role for Tmc7 in acrosome biogenesis during spermiogenesis.

Keywords: Transmembrane channel-like (Tmc) proteins; infertility; spermiogenesis; acrosome biogenesis

Introduction

Spermiogenesis is the post-meiotic phase of male germ cell development during which the haploid spermatids differentiate into mature spermatozoa[1]. Mouse spermiogenesis consists of 16 steps characterized by morphological changes in acrosome formation, nuclear condensation and elongation, flagellum formation, and cytoplasm removal[2]. Acrosome biogenesis is one of the key events of spermiogenesis, and it begins with the initial stage of spermatid development. The acrosome is a specialized organelle with a cap-like structure covering the anterior part of the mature spermatozoon head and is indispensable for the fertilization process[3]. Although acrosome biogenesis is thought to be derived from the Golgi complex in early spermatids, growing evidence suggests that the endoplasmic reticulum (ER) and lysosomes are involved in acrosome biogenesis[4]. Defects in acrosome biogenesis can result in subfertility or infertility.

Acrosome biogenesis is generally classified into four successive phases, namely the Golgi, cap, acrosome, and maturation phases[5]. A number of ER and Golgi-associated proteins have been shown to be involved in acrosome biogenesis. Heat shock protein 90B1 (Hsp90b1) is an ER molecular chaperone that has been shown to be a testis-specific protein, and Hsp90b1 deficiency leads to a globozoospermia-syndrome like phenotype [6]. Gm130 (Golgi matrix protein 130) localize on the cis surface of the Golgi apparatus, and loss of Gm130 causes acrosome malformations and results in male infertility[7]. Gopc (Golgi-associated PDZ and coiled-coil motif-containing) is a trans-Golgi-associated protein that plays a role in vesicle transport from the Golgi apparatus. Gopc deficiency leads to vesicle transport failure, which in turn results in malformation of the acrosome and to infertility[8]. Other vesicle

transport proteins – including Pick1, Ica1l, and Vps13b – are involved in proacrosomal

vesicle trafficking during acrosome formation[9,10]. Recent evidence suggests that autophagy is

involved in proacrosomal vesicle fusion and transport to form the acrosome, and disruption of

Atg7[11,12] or Atg5 causes aberrant acrosome formation and male infertility or subfertility.

Despite these findings, further studies are needed to better understand the complexity of

acrosome biogenesis.

73   Transmembrane channel-like (Tmc) proteins are highly conserved ion channel-like

proteins, with eight family members (Tmc1–Tmc8) identified in humans and mice [13-16]. Tmc

proteins are predicted to have 6–10 transmembrane domains, and all protein members contain

a conserved 120 amino acid Tmc domain[15]. The single-particle cryo-electron microscopy

structure shows that Tmc1 assembles as dimers and that each monomer contains 10

transmembrane domains with four domains (S4–S7) that line the channel pore[17]. Tmc1 and 2

are mechano-electrical transducer (MET) channels in cochlear hair cells, and mutations in

Tmc1/2 cause deafness in humans and mice[18] by disrupting hair cell mechanoelectrical

transduction, and *Tmc* gene therapy can restore auditory function and balance in mice[19,20].

Germline mutations in *Tmc6* and *Tmc8* are associated with epidermodysplasia verruciformis,

which leads to increased susceptibility to HPV infection [14,15]. Among the eight Tmc protein

members, Tmc1 and Tmc2 are well studied due to their role in hearing conduction, and

understanding the physiological and functional roles of other Tmc family members remains a

major challenge.

87   In this study, we found that Tmc7 begins to be expressed in diplotene spermatocytes and

that it is expressed at high levels in round spermatids in mouse testes. By generating *Tmc7*

conventional knockout mice (*Tmc7$^{-/-}$*), we demonstrate that Tmc7 deficiency leads to aberrant

Golgi morphology and impaired fusion of Golgi-derived vesicles to the acrosome, thus

resulting in male infertility with an oligo-astheno-teratozoospermia (OAT) phenotype. Our

findings provide new insights into the physiological role of Tmc family members in

spermiogenesis.

Results

Tmc7 expression is confined to the testis, and deletion of Tmc7 leads to male infertility

To explore the roles of Tmc family members during spermatogenesis, we analyzed the

expression of Tmc1–Tmc8 using the mouse developmental transcriptome (E-MTAB-6798)[21].

Among the eight Tmc family proteins, we found that only Tmc5 and Tmc7 exhibited elevated

expression in the testis during development (Fig. 1a). Tmc7 caught our great interest because

Tmc7 was expressed at higher levels than Tmc5 in the testis during development. We next

analyzed Tmc7 expression in different tissues using single-cell transcriptomic data from the

Mouse Cell Atlas database[22]. We found that Tmc7 was predominantly expressed in the testis

when compared with all other analyzed tissues (Fig. 1b). Further, we analyzed Tmc7 in

different stages of germ cell development using single-cell sequencing data of male-specific

tissues and organs in mice and humans[23]. Interestingly, Tmc7 began to be expressed in

diplotene spermatocytes and was expressed at high levels in round spermatids in mouse testes

(Fig. 1c, d). The Tmc7 expression pattern was confirmed by RT-PCR (Fig. 1e, f), and Tmc7

was again found to be highly expressed in round spermatids of human testis (Fig. S1a, b). The

distinct expression pattern in the testis indicates the potential role of Tmc7 in

spermatogenesis.

We generated *Tmc7*-deficient mice (*Tmc7*$^{-/-}$) by partial deletion of exon 2 (the first 241

bp) using the CRISPR/Cas9 system (Fig. 1g). *Tmc7*$^{-/-}$ mice were confirmed by PCR-based

genotyping and immunoblotting of the separated cytoplasmic and membrane testis proteins

(Fig. S1c, d). We then examined the phenotypes and found that *Tmc7*-deficient mice were

viable and had a normal appearance and exhibited a normal metabolic rate (Fig. S1e-j).

However, *Tmc7*-deficient male mice showed smaller testes (Fig. 1h, i) and a reduced

testis/body weight ratio (Fig. 1j) compared to control mice. Moreover, *Tmc7*$^{-/-}$ male mice

exhibited complete infertility (Table S1). No difference in the number of tubules per testicle

or in the number of Sertoli cells was seen when comparing *Tmc7*$^{-/-}$ and WT mice; Sertoli cell

were identified by immunofluorescence staining of the Sertoli cell marker Sox9 (Fig. S1k, i).

These findings suggest that Tmc7 is essential for spermatogenesis and thus for male fertility.

***Tmc7* deficiency leads to severe defects in spermiogenesis**

To determine the cause of infertility in *Tmc7*$^{-/-}$ mice, we examined the composition of cell

types in seminiferous tubules and cauda epididymis sections using hematoxylin staining in

mouse testes at different developmental stages. In seminiferous tubules, the populations of

germ cells were comparable between wild type (WT) and *Tmc7*$^{-/-}$ mice at postnatal day

(PD)14 and PD21. Intriguingly, the number of elongating spermatids in *Tmc7*$^{-/-}$ mice

decreased significantly compared with WT mice at PD28 (Fig. S2a). We next examined the

testis and cauda epididymis sections of 9-week-old WT and *Tmc7*$^{-/-}$ mice. We observed

spermiogenesis defects in *Tmc7*$^{-/-}$ mice as evidenced by the reduced number of elongating

spermatids, the increased number of sloughed germ cells, and large vacuoles in the

seminiferous tubules and cauda epididymis (Fig. 2a, b). The vacuolation became more serious

with increasing age (Fig. S2a). The total number of sperm in the cauda epididymis was

drastically decreased in $Tmc7^{-/-}$ mice (~0.15×10^6 cells) compared to wild type (WT) mice

(~8.5×10^6 cells) (Fig. 2c). The morphological assessment of spermatozoa from the caudal

epididymis revealed that a large proportion of spermatozoa (~85%) in $Tmc7^{-/-}$ mice exhibited

abnormal head morphology, similar to human OAT (Fig. 2d, e). In parallel with this, periodic

acid Schiff (PAS) staining indicated that a large number of spermatids failed to differentiate

into elongated spermatids with sickle-shaped heads and instead exhibited abnormal head

morphology. The total number of spermatids began to significantly decrease at steps 9–12 of

spermiogenesis (Fig. 2f, g), and we performed Mito-tracker staining and found shortened or

disorganized mitochondrial sheaths in $Tmc7^{-/-}$ spermatozoa collected from the cauda

epididymis (Fig. 2h). Meanwhile, no motile or progressive sperm were found in the $Tmc7^{-/-}$

mice (Fig. 2i, j). Moreover, meiotic progression was not affected in $Tmc7^{-/-}$ mice as evidenced

by immunofluorescence staining of the meiosis markers Sycp3 (red) and γH2ax (green) in the

seminiferous tubules of PD20 WT and $Tmc7^{-/-}$ mice (Fig. S3a). Overall, we concluded that

Tmc7 ablation results in defects in spermiogenesis and in OAT-like phenotypes.

Acrosome biogenesis is impaired starting from the Golgi phase in $Tmc7^{-/-}$ spermatids

Acrosome biogenesis is one of the key events of spermiogenesis. To investigate whether

Tmc7 is required for the proper formation of the acrosome, FITC-labelled peanut agglutinin

(PNA) was used to assess the acrosomal status of spermatids in 9-week-old WT and $Tmc7^{-/-}$

mice. The spermatids of WT mice had normal structures in the Golgi, cap, acrosome, and

maturation phases, while spermatids in $Tmc7^{-/-}$ mice exhibited abnormal acrosome

morphology as early as the Golgi phase. Fragmented and malformed acrosomes were

observed in all four phases of spermatids in the $Tmc7^{-/-}$ mice (Fig. 3a). Absent or fragmented

acrosomes were also found in $Tmc7^{-/-}$ spermatids collected from the cauda epididymis (Fig.

S4a). To better understand the morphologic defects of the acrosome in greater detail,

transmission electron microscopy (TEM) was performed to analyze the acrosome structure in

WT and $Tmc7^{-/-}$ mice. We found that large vesicles accumulated adjacent to developing

acrosomes in all phases of acrosome biogenesis, starting from the Golgi phase (Fig. 3b, left

panel) in the spermatids of $Tmc7^{-/-}$ mice. Additionally, abnormal acrosome structures and

defective nuclear condensation were observed in $Tmc7^{-/-}$ spermatids (Fig. 3b, right panels).

Further investigation revealed that knockout of $Tmc7$ resulted in Golgi apparatus

fragmentation and disorganization, as determined by TEM and immunofluorescence staining

of the Golgi marker Gm130 (Fig. 3b-c). The aspect ratio (height over width) of the Golgi

apparatuses in $Tmc7^{-/-}$ spermatids (R = ~2.9) based on TEM images were also significantly

smaller than in WT spermatids (R = ~4.9) (Fig. 3d). Meanwhile, we found that the Golgi body

failed to correctly orient toward the acrosome (Fig. S4b). We further performed

immunoblotting to measure Golgi stress markers, Hsp47 and Tfe3 in 9-week-old $Tmc7^{-/-}$

testes compared with WT testes. The results showed that the protein levels of Hsp47 and Tfe3

were significantly increased upon loss of Tmc7, indicating the occurrence of Golgi stress (Fig.

3e). Gopc is localized in the trans-Golgi network and is involved in the transport and fusion of

the proacrosomal vesicles with the acrosome[8]. By immunostaining for Gopc and PNA, we

found that Gopc staining (red) was colocalized with PNA (green) in WT spermatids. In

contrast, colocalized Gopc and PNA spermatids were significantly decreased in $Tmc7^{-/-}$ mice

(Fig. 3f). Together these findings suggest that Tmc7 is required for proacrosomal vesicle

trafficking and fusion during acrosome biogenesis.

Tmc7 localizes to the cis-Golgi, and this is required for maintaining Golgi integrity

To further delineate the role of Tmc7 in acrosome formation, we next sought to determine the

subcellular localization of Tmc7 in the testis. We performed immunostaining and found that

Tmc7 was predominantly expressed in the perinuclear regions of pachytene spermatocytes

and round spermatids in WT testes but not in $Tmc7^{-/-}$ testes, suggesting the specificity of the

antibody (Fig. 4a). By immunofluorescence co-staining of Tmc7 with the cis-Golgi marker

Gm130, the trans-Golgi marker Tgn46, and PNA, we found that Tmc7 colocalized closely

with Gm130 but not with Tgn46 (Fig. 4b, c, Fig. S5a), indicating its localization to the

cis-Golgi apparatus. Further evaluation showed that Tmc7 and PNA were closely associated

but not colocalized (Fig. 4d, Fig. S5b). In parallel, we carried out co-immunoprecipitation

with Tmc7 antibody coupled with immunoblotting in mouse testes and found that Tmc7 was

able to pull down Gm130, the Gm130-interacting protein P115, and Grasp65 (Fig. 4e), further

confirming that Tmc7 is localized to the cis-Golgi apparatus.

Gm130 is reported to be involved in maintaining the function and integrity of the Golgi

apparatus[7]. To gain more insight into the role of Tmc7 in the Golgi apparatus, we performed

immunostaining against Gm130, P115, and Grasp65 in the testes of 9-week-old WT and

$Tmc7^{-/-}$ mice. The results showed that the protein levels of Gm130, P115, and Grasp65 were

significantly reduced in $Tmc7^{-/-}$ testes compared to WT testes (Fig. 4f, g, Fig. S5c-e).

Immunoblotting further verified the reduced levels of Gm130, P115, and Grasp65 upon loss

of Tmc7 (Fig. 4h). These results suggest that Tmc7 colocalizes to Gm30 in the cis-Golgi and

that this is required for maintaining Golgi integrity.

Tmc7 depletion impairs cellular homeostasis and leads to spermatid apoptosis

Golgi pH and ion homeostasis is essential for proper function, including membrane

trafficking, protein glycosylation, and protein sorting[24], and the Golgi apparatus has been

shown to be another intracellular Ca^{2+} store in addition to the endoplasmic reticulum (ER)[25].

$Tmc1$ mutation leads to reduced Ca^{2+} permeability, thus triggering hair cell apoptosis and

subsequent deafness[26]. We measured intracellular Ca^{2+} by flow cytometry in WT and $Tmc7^{-/-}$

germ cells isolated from PD30 testes. The results showed that the intracellular Ca^{2+}

concentration was significantly reduced in $Tmc7^{-/-}$ germ cells compared to WT germ cells

(Fig. 5a). In contrast, intracellular pH was increased in $Tmc7^{-/-}$ germ cells compared to WT

germ cells (Fig. 5b). These results suggest that the pH and ion homeostasis was impaired. We

further measured the reactive oxygen species (ROS) levels and observed that ROS levels were

significantly increased upon loss of Tmc7 (Fig. 5c) In addition, we examined ER

stress-related proteins by immunoblotting and found that p-eif2α, Chop, and Grp78 were

significantly increased in 9-week-old $Tmc7^{-/-}$ testes compared with WT testes, indicating the

occurrence of ER stress (Fig. 5d). Moreover, we carried out TUNEL staining in the

seminiferous tubules and cauda epididymis of 9-week-old WT and $Tmc7^{-/-}$ mice. The

TUNEL-positive cells in seminiferous tubules and cauda epididymis were significantly

increased in $Tmc7^{-/-}$ mice compare to WT mice (Fig. 5e). The expression of apoptosis-related

proteins was next measured by immunoblotting, and depletion of Tmc7 increased the levels of

cleaved caspase-3 and Bax and decreased the levels of Bcl2 (Fig. 5f). These results support

the hypothesis that Tmc7 is required to maintain Golgi pH and ion homeostasis and that this is

needed for acrosome biogenesis.

Discussion

Among the eight Tmc protein members, Tmc1 and Tmc2 are localized to the site of MET

channels at the tips of the shorter stereocilia of cochlear hair cells[27,28], and mutations in *Tmc1*

and *Tmc2* cause deafness. However, Tmc1 and Tmc2 expressed in heterologous cell lines

remain in the ER and fail to localize to the plasma membrane[18]. Tmc4 is a voltage-dependent

chloride channel that is predominantly expressed in the taste buds in the posterior region of

the tongue[29]. Tmc6 and Tmc8 are localized to the ER and play crucial roles in zinc transport[30].

However, the physiological function of Tmc7 has not been defined. In this study, we

demonstrated that Tmc7 was predominantly expressed in testes (Fig. 1e), and by generating

knockout mice lacking *Tmc7* we found that Tmc7 deficiency resulted in complete male

infertility with an OAT-like phenotype. Interestingly, we found that Tmc7 co-localized with

Gm130, but not with Tgn46 (Fig. 4b, c), as illustrated by the immunofluorescence co-staining

assay, indicating that Tmc7 located on the cis side of the Golgi apparatus.

Our results demonstrate the critical role of Tmc7 during the formation of the acrosome. The

acrosome is thought to be derived from the Golgi apparatus, so we speculated that Tmc7

deficiency might cause defects in acrosome formation because Tmc7 localized on the Golgi

apparatus. We observed fragmented and malformed acrosomes starting from the Golgi phase

in $Tmc7^{-/-}$ spermatids as analyzed by TEM (Fig. 3b). Additionally, proacrosomal vesicles

were found to accumulate around the acrosomes (Fig. 3b), and spermatids with colocalized

Gopc and PNA were significantly decreased in $Tmc7^{-/-}$ mice (Fig. 3f), suggesting that

Golgi-derived vesicles failed to fuse with developing acrosomes. In accordance with our

findings, other proteins that localize on the Golgi apparatus have also been reported to be

involved in acrosome biogenesis. Gm130 deficiency results in acrosome biogenesis failure

and male infertility because proacrosomal vesicles are unable to fuse to form acrosomes[30].

Our findings showed that Tmc7 was co-localized with Gm130 and that Tmc7 depletion

significantly reduced the protein level of Gm130. Gopc is localized on the trans-Golgi region,

and its deficiency leads to defective fusion of Golgi-derived transport vesicles to the

acrosomal cap and to a complete lack of acrosomes[8]. Colocalized Gopc and PNA spermatids

were significantly decreased in $Tmc7^{-/-}$ spermatids compared to WT spermatids. Further,

Golga3, Pick1, and several other Golgi proteins are also required for acrosome biogenesis[31].

Our work, combined with previous studies, broadens our understanding of role the Golgi

apparatus plays in the formation of acrosomes.

Our findings highlight the role of Tmc7 in maintaining Golgi and cellular homeostasis.

Upon Tmc7 depletion, the Golgi apparatus undergoes significant disorganization and is

unable to correctly orient toward the acrosome. The mechanism behind this is that

intracellular Ca^{2+} concentrations decreased while the pH value was increased in $Tmc7^{-/-}$ germ

cells compared to WT germ cells, suggesting that Tmc7 is essential for maintaining the pH

and ion homeostasis of the Golgi apparatus. Our findings further revealed that Tmc7

deficiency led to increased ROS levels, Golgi and ER stress, ultimately resulting in apoptosis

of spermatids. The Na^+/H^+ exchanger Nhe8 is involved in the maintenance of intracellular pH

and ion homeostasis, and Nhe8 deficiency leads to Golgi-derived vesicles failing to form

large acrosomal granules and the acrosomal cap[32]. Slc9a3 is another Na^+/H^+ exchanger, and it

is expressed in the acrosomal region of spermatids and loss of Slc9a3 causes severe

acrosomal defects[33]. These studies, together with our current findings, support the hypothesis

that pH and ion homeostasis of the Golgi apparatus is needed for the proper progression of

acrosome biogenesis.

In conclusion, our findings show that Tmc7 localizes to the cis-Golgi region in

spermatids and that this is essential for acrosome biogenesis and for male fertility. In the

absence of Tmc7, Golgi pH and ion homeostasis is disrupted, and this leads to Golgi

disorganization and failure to correctly orientate to the acrosome. Subsequently,

Golgi-derived proacrosomal vesicles are unable to fuse with the developing acrosome and

acrosome biogenesis is severely impaired, and thus Tmc7-deficent male mice are completely

infertile. Our study highlights the physiological role of TMC family members in

spermatogenesis.

Materials and methods

Generation of *Tmc7* knockout mice

Tmc7 conventional knockout mice in the C57BL/6J background were obtained from Cyagen

Bioscience (China). Exon 2 was selected as the target site, and Cas9 mRNA and single guide

RNA were co-injected into fertilized eggs for knockout mouse production. The pups were

genotyped by PCR using the following primers: F1:5'-AGA TGG CTT AGT GGG TGC CTT

AGT-3'; R1:5'TGA TGG ACA GAG AGA TGA TGA ATG G-3'; F3: 5'-CAG TCC CAG

GTT CAG TTC AGT GAG-3'. In WT mice, the primer pair amplified a 513 bp PCR product,

while in *Tmc7*$^{-/-}$ mice it amplified an 891 bp product. Mice were housed under a controlled

illumination regime (light for 12 h) at 21°C–22°C and 60% ± 10% relative humidity. Food

and water were freely available, and all animal experiments were conducted in accordance

with the guidelines of the Animal Care and Use Committee at Shandong University

(ECSBMSSDU2022-2-113).

Metabolic assessment in mice

A Columbus Instruments Comprehensive Lab Animal Monitoring System CLAMS6 was used

for the experiment. Nine-week-old WT and *Tmc7*$^{-/-}$ male mice ($n = 4$) were weighed and had

the metabolic rates measured over a period of 48 hours. The rate of oxygen consumption,

RER (breathing entropy), energy expenditure, and food intake were automatically recorded

every eight minutes. These data were analyzed with a 12 h dark:12 h light cycle at 22–23 °C.

RNA isolation and PCR

The tissues were homogenized in TRIzol (Invitrogen, 15596-026) with a grinder. Following

centrifugation ,the RNA in the supernatant was obtained by precipitating with isopropanol and

washing with ethanol. A total of 1 µg RNA was reverse-transcribed into cDNA with HiScript

III RT SuperMix for RT-PCR (+gDNA wiper) (Vazyme, R223-01) according to the

manufacturer's instructions. The cDNA was used to detect the expression of Tmc7 with Taq

Master Mix (Vazyme, P112-03) with the following primers: F – AGC CAA TCC ATC CGC

AAG TAT; R – TGC TGC TG TAC CGT TTC CAC. The products were analyzed by agarose

gel electrophoresis.

Sperm collection

The cauda epididymis was collected and placed in an EP tube containing PBS. Sperm were

released for 30 min in an atmosphere of 5% CO_2 at 37 °C. The suspension was diluted at 1:50

and transferred to a hemocytometer for counting. For the anther experiments, the PBS with

sperm was spread on the slide and fixed with 4% paraformaldehyde (Servicebio, G1101). The

slides were stored at –80°C after drying.

Histological analysis

For the morphology analysis of the sperm from the caudal epididymis, sperm smears were

directly stained with hematoxylin and sealed with neutral balsam. The tissue was fixed in 4%

paraformaldehyde overnight, then dehydrated and embedded in paraffin. Paraffin-embedded

tissues were sectioned at 4 µm. Tissue slides were deparaffinized in xylene and dehydrated in

a series of ethanol. The tissues were then stained by hematoxylin or with periodic acid

staining using a PAS Stain Kit (Abcam, ab150680) according to the manufacturer's

instructions. Finally, tissue slides were rehydrated in a series of ethanol and xylene and sealed

with neutral balsam. An Olympus BX53 microscope system was used for imaging.

Immunofluorescence

We performed immunofluorescence of germ cells in the testes using two methods, including

testis slides and squashed seminiferous tubules. The squashed seminiferous tubules were

prepared according to a previous report[34]. The testis slides were deparaffinized in xylene and

rehydrated in a series of ethanol. Antigen retrieval was performed by heating in EDTA

Antigen Retrieval Solution (pH 9.0) (ZSGB-BIO, ZLI-9096), and nonspecific binding sites

were blocked with EZ-Buffers M 1X Block BSA in PBS (Sangon, C520035-0250). The slides

were incubated with primary antibody overnight at 4℃. Primary antibodies included

anti-Tmc7 (1:200 dilution, Abcam, ab191521), anti-Sox9 (1:500 dilution, Sigma, AB5535),

anti-Gopc (1:200 dilution, Proteintech, 12163-1-AP), anti-Sycp3 (1:500 dilution, Santa Cruz,

sc-33195), anti-H2ax (1:500 dilution, Millipore, 05-636-1), anti-Gm130 (1:100 dilution,

610822, BD Biosciences), anti-P115 (1:200 dilution, Proteintech, 13509-1-AP), and

anti-Grasp65 (1:200 dilution, Proteintech, 10747-2-AP). The next day they were incubated

with the secondary antibody (1:500 dilution, Thermo, A-11070 and A-11012) for 1 h at room

temperature. The Alexa Fluor 488 conjugates of lectin PNA (1:500 dilution, Thermo, L21409),

anti-Gm130 (1:200 dilution, Proteintech, CL488-11308), and anti-Tgn46 (1:200 dilution,

Proteintech, CL488-13573) were used as secondary antibodies. For the staining of Mito-track

(1:5000 dilution, Beyotime, C1048), the sperm released from the caudal epididymis were

directly stained for 30 min at 37 °C. The sperm were then spread on the slide and observed.

A TUNEL assay was performed using a one-step TUNEL Apoptosis Assay Kit (KeyGEN

BioTECH, KGA7072) according to the manufacturer's instructions. Briefly, the slide was

deparaffinized and antigen retrieval was performed. dUTP was used to label the cells, and the

nuclei were stained with DAPI (Abcam, ab285390). Fluorescent images were captured on a

Dragonfly 200 confocal microscopy system (Andor Technology).

Western blotting

The membrane and cytoplasmic proteins of the testis were obtained using a membrane protein

extraction kit (Abmart, A1008) according to the manufacturer's instructions. Total protein in

the testis was obtained by grinding in a tissue grinder in RIPA lysis buffer (Beyotime,

P0013B), and the concentration of protein was measured using a BCA protein assay kit

(Beyotime, P0009). The proteins were separated by SDS-PAGE and then transferred to a

PVDF membrane. The membranes were blocked in 5 % non-fat milk and then incubated with

primary antibodies overnight at 4 °C. Primary antibodies included anti-Tmc7 (Dia-an

Biotech), anti-Atp1a1 (Proteintech, 14418-1-AP), anti-Gapdh (Proteintech, 60004-1-1g),

anti-Cleaved-Caspase3 (Cell Signaling Technology, 9661), anti-Caspase3 (Cell Signaling

Technology, 9662), anti-Bax (Cell Signaling Technology, 41162), anti-Bcl2 (Cell Signaling

Technology, 17447), anti-α-Tubulin (Proteintech,11224-1-AP), anti-P-Eif2α (ABclonal,

AP0745), anti-Eif2α (ABclonal, A0764), anti-Grp78 (Proteintech, 11587-1-AP), anti-Chop

(Proteintech, 15204-1-AP), anti-P115 (Proteintech, 13509-1-AP), and anti-Grasp65

(Proteintech, 10747-2-AP). All antibodies were diluted 1:1000. The next day, the membrane

was incubated with horseradish peroxidase-conjugated secondary antibody for 1 hour, and the

protein was detected using an ECL system (Vazyme, E412-02). The quantitation was

performed in Image J.

Chromosome spreading

Meiotic nuclear spreading was performed as previously reported[35]. Briefly, spermatocytes

released from seminiferous tubules were incubated in hypotonic solution for 1 hour and then

transferred to 100 mM sucrose. The suspension was spread on the glass slides with 1% PFA

and 0.15% Triton X100, and dried slides were stored at –80°C or were used for the

immunofluorescence staining.

Transmission electron microscopy

The testes of 9-week-old WT and $Tmc7^{-/-}$ mice ($n = 3$) were used for the TEM analysis. The

testes were washed in PBS three times and then fixed in 1% O_SO_4 in 0.1 M PBS (pH 7.4)

buffer at 4°C overnight. The testes were then fixed in 1% O_sO_4 at room temperature for 1

hour. The samples were then washed with water three times for 15 min each and dehydrated

in a graded ethanol series (50%, 70%, 95%, and 100%). The samples were finally embedded

with resin, cut into 80 nm sections, and viewed with a JEM-1400 transmission electron

microscope.

Flow cytometry

The testes of PD30 WT and $Tmc7^{-/-}$ mice ($n = 3$) were used to obtain germ cells. The tunica

albuginea was removed and placed into PBS with 120 U/ml Type I Collagenase (Gibco,

17100-017) on a shaker at 37°C and 220 rpm for 10 min. The supernatant was filtered through

a 70 μm cell filter screen into a new tube. The sediment was digested again with 0.25%

trypsin-EDTA (Gibco, 25200–072) on a shaker at 37°C and 220 rpm for 10 min and then

neutralized by serum albumin and filtered. The germ cells were obtained by centrifuging

twice at $1000 \times g$ for 5 min and then removing the supernatant.

The levels of Ca^{2+}, pH, and ROS were measured using fluorescent probes according to the

manufacturer's protocols (Beyotime, S1056, S1006, S0033S). The probes were all diluted

1:2000. Germ cells were stained with the probes at 37°C for 30 min, then washed with PBS

three times and analyzed by flow cytometry (Gallios, US).

Co-immunoprecipitation

Testes of 9-week-old WT mice were ground in a tissue grinder in RIPA Lysis Buffer

(Beyotime, P0013B) to obtain the total protein. Tmc7 (Dia-an Biotech) and IgG (Beyotime,

A7016) antibodies were incubated with the proteins overnight. The next day, they were

incubated with protein A/G magnetic beads (MCE, HY-K0202) for 2 hours. The beads were

wash and boiled for 10 min in 1× SDS loading buffer. Western blotting was used for the

analysis of input and immunoprecipitated samples with anti-P115 and anti-Grasp65

antibodies.

Statistical analysis

GraphPad Prism 9 was used to analyze the data, and the results are shown as the mean $\pm$

standard deviation (S.D.). Statistical significance was determined using two-sided Student's

t-tests or two-way analysis of variance (ANOVA). $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$

indicate the level of statistical significance, and ns stands for not significant.

Acknowledgements

We thank Translational Medicine Core Facility of Shandong University for consultation and

instrument availability that supported this work. We also thank the Laboratory Animal Center

of Shandong University for mouse housing and care. This work was supported by National

Natural Science Foundation of China (82172718, 82301801, 82303055, 81972437).

411
Author Contributions

J.W., Y.Y, performed most experiments, and data analysis, and writing the manuscript. Z.W.,

L.Y., performed public database sequence data analysis. C.Q., Y.G., L.Y., helped with animal

and cell experiment. F.G., G.Lu., Z.C., X.Z., is involved technical, material support and

guided the experiment. H.L., Z.L. Supervised the project and revised the manuscript. Zhaojian

Liu was the lead contact for the study. Final approval: All authors.

Conflict of interest

The authors declare no competing interests.

Reference

1. Kang, J.Y., Wen, Z., Pan, D., Zhang, Y., Li, Q., Zhong, A., Yu, X., Wu, Y.C., Chen, Y.,

Zhang, X., et al. (2022). LLPS of FXR1 drives spermiogenesis by activating translation

of stored mRNAs. Science *377*, eabj6647. 10.1126/science.abj6647.

2. Li, W., Wu, J., Kim, S.Y., Zhao, M., Hearn, S.A., Zhang, M.Q., Meistrich, M.L., and

Mills, A.A. (2014). Chd5 orchestrates chromatin remodelling during sperm

development. Nat Commun *5*, 3812. 10.1038/ncomms4812.

3. Xiong, W., Shen, C., and Wang, Z. (2021). The molecular mechanisms underlying

acrosome biogenesis elucidated by gene-manipulated mice†. Biol Reprod *105*,

789-807. 10.1093/biolre/ioab117.

4. Khawar, M.B., Gao, H., and Li, W. (2019). Mechanism of Acrosome Biogenesis in

Mammals. Front Cell Dev Biol *7*, 195. 10.3389/fcell.2019.00195.

5. Xu, K., Su, X., Fang, K., Lv, Y., Huang, T., Li, M., Wang, Z., Yin, Y., Muhammad, T.,

Liu, S., et al. (2023). The Slingshot phosphatase 2 is required for acrosome

biogenesis during spermatogenesis in mice. Elife *12*. 10.7554/eLife.83129.

6. Audouard, C., and Christians, E. (2011). Hsp90beta1 knockout targeted to male

germline: a mouse model for globozoospermia. Fertil Steril *95*, 1475-1477

e1471-1474. 10.1016/j.fertnstert.2010.12.006.

7. Han, F., Liu, C., Zhang, L., Chen, M., Zhou, Y., Qin, Y., Wang, Y., Chen, M., Duo, S.,

Cui, X., et al. (2017). Globozoospermia and lack of acrosome formation in

GM130-deficient mice. Cell Death Dis *8*, e2532. 10.1038/cddis.2016.414.

8.   Yao, R., Ito, C., Natsume, Y., Sugitani, Y., Yamanaka, H., Kuretake, S., Yanagida, K.,

Sato, A., Toshimori, K., and Noda, T. (2002). Lack of acrosome formation in mice

lacking a Golgi protein, GOPC. Proc Natl Acad Sci U S A *99*, 11211-11216.

10.1073/pnas.162027899.

9.   Xiao, N., Kam, C., Shen, C., Jin, W., Wang, J., Lee, K.M., Jiang, L., and Xia, J. (2009).

PICK1 deficiency causes male infertility in mice by disrupting acrosome formation. J

Clin Invest *119*, 802-812. 10.1172/JCI36230.

10.   He, J., Xia, M., Tsang, W.H., Chow, K.L., and Xia, J. (2015). ICA1L forms

BAR-domain complexes with PICK1 and is crucial for acrosome formation in

spermiogenesis. J Cell Sci *128*, 3822-3836. 10.1242/jcs.173534.

11.   Wang, H., Wan, H., Li, X., Liu, W., Chen, Q., Wang, Y., Yang, L., Tang, H., Zhang, X.,

Duan, E., et al. (2014). Atg7 is required for acrosome biogenesis during

spermatogenesis in mice. Cell Res *24*, 852-869. 10.1038/cr.2014.70.

12.   Huang, Q., Liu, Y., Zhang, S., Yap, Y.T., Li, W., Zhang, D., Gardner, A., Zhang, L.,

Song, S., Hess, R.A., and Zhang, Z. (2021). Autophagy core protein ATG5 is required

for elongating spermatid development, sperm individualization and normal fertility in

male mice. Autophagy *17*, 1753-1767. 10.1080/15548627.2020.1783822.

13.   Keresztes, G., Mutai, H., and Heller, S. (2003). TMC and EVER genes belong to a

larger novel family, the TMC gene family encoding transmembrane proteins. BMC

Genomics *4*, 24. 10.1186/1471-2164-4-24.

14.   Kurima, K., Yang, Y., Sorber, K., and Griffith, A.J. (2003). Characterization of the

transmembrane channel-like (TMC) gene family: functional clues from hearing loss

and epidermodysplasia verruciformis. Genomics *82*, 300-308.

10.1016/s0888-7543(03)00154-x.

15. Yue, X., Sheng, Y., Kang, L., and Xiao, R. (2019). Distinct functions of TMC channels:

a comparative overview. Cell Mol Life Sci *76*, 4221-4232.

10.1007/s00018-019-03214-1.

16. Kurima, K., Peters, L.M., Yang, Y., Riazuddin, S., Ahmed, Z.M., Naz, S., Arnaud, D.,

Drury, S., Mo, J., Makishima, T., et al. (2002). Dominant and recessive deafness

caused by mutations of a novel gene, TMC1, required for cochlear hair-cell function.

Nat Genet *30*, 277-284. 10.1038/ng842.

17. Jeong, H., Clark, S., Goehring, A., Dehghani-Ghahnaviyeh, S., Rasouli, A.,

Tajkhorshid, E., and Gouaux, E. (2022). Structures of the TMC-1 complex illuminate

mechanosensory transduction. Nature *610*, 796-803. 10.1038/s41586-022-05314-8.

18. Nist-Lund, C.A., Pan, B., Patterson, A., Asai, Y., Chen, T., Zhou, W., Zhu, H., Romero,

S., Resnik, J., Polley, D.B., et al. (2019). Improved TMC1 gene therapy restores

hearing and balance in mice with genetic inner ear disorders. Nat Commun *10*, 236.

10.1038/s41467-018-08264-w.

19. Askew, C., Rochat, C., Pan, B., Asai, Y., Ahmed, H., Child, E., Schneider, B.L.,

Aebischer, P., and Holt, J.R. (2015). Tmc gene therapy restores auditory function in

deaf mice. Sci Transl Med *7*, 295ra108. 10.1126/scitranslmed.aab1996.

20. Nist-Lund, C.A., Pan, B., Patterson, A., Asai, Y., Chen, T., Zhou, W., Zhu, H., Romero,

S., Resnik, J., Polley, D.B., et al. (2019). Publisher Correction: Improved TMC1 gene

therapy restores hearing and balance in mice with genetic inner ear disorders. Nat

486   Commun *10*, 734. 10.1038/s41467-019-08716-x.

487   21.   Cardoso-Moreira, M., Halbert, J., Valloton, D., Velten, B., Chen, C., Shao, Y., Liechti,

488   A., Ascencao, K., Rummel, C., Ovchinnikova, S., et al. (2019). Gene expression

489   across mammalian organ development. Nature *571*, 505-509.

490   10.1038/s41586-019-1338-5.

491   22.   Han, X., Wang, R., Zhou, Y., Fei, L., Sun, H., Lai, S., Saadatpour, A., Zhou, Z., Chen,

492   H., Ye, F., et al. (2018). Mapping the Mouse Cell Atlas by Microwell-Seq. Cell *172*,

493   1091-1107 e1017. 10.1016/j.cell.2018.02.001.

494   23.   Zhao, L., Zhao, Y., Yao, C., Dai, Y., Li, Z., and Tang, Y. (2023). MHA, an interactive

495   website for scRNA-seq data of male genitourinary development and disease.

496   Andrology *11*, 1157-1162. 10.1111/andr.13402.

497   24.   Li, J., and Wang, Y. (2022). Golgi Metal Ion Homeostasis in Human Health and

498   Diseases. Cells *11*. 10.3390/cells11020289.

499   25.   Lissandron, V., Podini, P., Pizzo, P., and Pozzan, T. (2010). Unique characteristics of

500   Ca2+ homeostasis of the trans-Golgi compartment. Proc Natl Acad Sci U S A *107*,

501   9198-9203. 10.1073/pnas.1004702107.

502   26.   Fettiplace, R., Furness, D.N., and Beurg, M. (2022). The conductance and

503   organization of the TMC1-containing mechanotransducer channel complex in auditory

504   hair cells. Proc Natl Acad Sci U S A *119*, e2210849119. 10.1073/pnas.2210849119.

505   27.   Beurg, M., Barlow, A., Furness, D.N., and Fettiplace, R. (2019). A Tmc1 mutation

506   reduces calcium permeability and expression of mechanoelectrical transduction

507   channels in cochlear hair cells. Proc Natl Acad Sci U S A *116*, 20743-20749.

10.1073/pnas.1908058116.

28.   Kurima, K., Ebrahim, S., Pan, B., Sedlacek, M., Sengupta, P., Millis, B.A., Cui, R.,

Nakanishi, H., Fujikawa, T., Kawashima, Y., et al. (2015). TMC1 and TMC2 Localize

at the Site of Mechanotransduction in Mammalian Inner Ear Hair Cell Stereocilia. Cell

Rep *12*, 1606-1617. 10.1016/j.celrep.2015.07.058.

29.   Kasahara, Y., Narukawa, M., Ishimaru, Y., Kanda, S., Umatani, C., Takayama, Y.,

Tominaga, M., Oka, Y., Kondo, K., Kondo, T., et al. (2021). TMC4 is a novel chloride

channel involved in high-concentration salt taste sensation. J Physiol Sci *71*, 23.

10.1186/s12576-021-00807-z.

30.   Lazarczyk, M., Pons, C., Mendoza, J.A., Cassonnet, P., Jacob, Y., and Favre, M.

(2008). Regulation of cellular zinc balance as a potential mechanism of

EVER-mediated protection against pathogenesis by cutaneous oncogenic human

papillomaviruses. J Exp Med *205*, 35-42. 10.1084/jem.20071311.

31.   Xiong, W., Shen, C., and Wang, Z. (2021). The molecular mechanisms underlying

acrosome biogenesis elucidated by gene-manipulated micedagger. Biol Reprod *105*,

789-807. 10.1093/biolre/ioab117.

32.   Oberheide, K., Puchkov, D., and Jentsch, T.J. (2017). Loss of the Na(+)/H(+)

exchanger NHE8 causes male infertility in mice by disrupting acrosome formation. J

Biol Chem *292*, 10845-10854. 10.1074/jbc.M117.784108.

33.   Wang, Y.Y., Chiang, H.S., Cheng, C.Y., Wu, Y.N., Lin, Y.C., Liu, H.C., Tsai, W.K.,

Chen, Y.L., and Lin, Y.H. (2017). SLC9A3 Protein Is Critical for Acrosomal Formation

in Postmeiotic Male Germ Cells. Int J Mol Sci *19*. 10.3390/ijms19010103.

34.   Wellard, S.R., Hopkins, J., and Jordan, P.W. (2018). A Seminiferous Tubule Squash

Technique for the Cytological Analysis of Spermatogenesis Using the Mouse Model. J

Vis Exp. 10.3791/56453.

35.   Peters, A.H., Plug, A.W., van Vugt, M.J., and de Boer, P. (1997). A drying-down

technique for the spreading of mammalian meiocytes from the male and female

germline. Chromosome Res *5*, 66-68. 10.1023/a:1018445520117.

Fig 1. Tmc7 expression is confined to the testis, and its deletion leads to male infertility.

(a) Heatmap analysis of the expression of Tmc family proteins at embryonic day (E)10–E18

and different postnatal days (P0, P3, PD14, PD28, and PD63) in mouse testis using the

Evo-devo mammalian organ database (http://apps.Kaessmannlab.org/evodevoapp/). (b) The

analysis of Tmc7 expression in different tissues using the Mouse Cell Atlas database

(https://bis.zju.edu.cn/MCA/). (c, d) T-Distributed Stochastic Neighbor Embedding (t-SNE)

plot and violin plot showing the expression of Tmc7 in different mouse germ cells using the

Male Health Atlas (http://malehealthatlas.cn/). SSC1-3: Spermatogonial stem cells I-III;

Progenitor: spermatogonia progenitor cell; A1–4: A1–4 spermatogonia; In: Intermediate

spermatogonia; B: Type B spermatogonia; PI: Preleptotene; L: Leptotene; Z: Zygotene; P:

pachytene; D: Diplotene; M: Metaphase; RS: Round spermatid. E: Elongating spermatid. (e)

The expression of *Tmc7* in different adult mouse tissues was measured by RT-PCR, including

the heart, liver, spleen, lung, kidney, brain, ovary, and testis. *Gapdh* was used as the control. (f)

The expression of *Tmc7* at different postnatal times (PD1, PD7, PD9, PD12, PD24, PD33,

and 2 months) was measured by RT-PCR. *Gapdh* was used as the control. (g) Schematic

illustrating the generation of *Tmc7*$^{-/-}$ mice using CRISPR/Cas9. (h) Testes from 9-week-old

WT and *Tmc7*$^{-/-}$ mice. (i) Testicular weights of 9-week-old WT and *Tmc7*$^{-/-}$ mice. Four adult

mice of each genotype were used. (j) Testis/body weight ratio of 9-week-old WT and *Tmc7*$^{-/-}$

mice. Four mice of each genotype were used. Images are representative of at least three

independent experiments. The results are presented as the mean ± S.D., and two-sided

unpaired Student's t-tests were used to calculate the P-values (*P < 0.05, **P < 0.01, ***P <

0.001).

Fig 2. Tmc7 deficiency leads to severe defects in spermiogenesis. (a, b) Hematoxylin

staining in testis and caudal epididymis sections of 9-week-old WT and *Tmc7*$^{-/-}$ male mice.

Scale bar, 50 μm. (c) The total number of sperm in the caudal epididymis from 9-week-old

WT and *Tmc7*$^{-/-}$ male mice. Three mice were used. (d) Hematoxylin-eosin staining of the

spermatozoa isolated from the caudal epididymis in 9-week-old WT and *Tmc7*$^{-/-}$ male mice.

H: head, F: flagella. Scale bar, 10 μm, 2 μm. (e) The percentage of spermatozoa with

abnormal heads in the caudal epididymis from WT and *Tmc7*$^{-/-}$ male mice. Three mice were

used. (f) PAS and hematoxylin staining in 9-week-old WT and *Tmc7*$^{-/-}$ mouse seminiferous

tubules. The numbers represent the stages of spermatogenesis. P: pachytene spermatocyte, PL:

preleptotene spermatocyte, L: leptotene spermatocyte, Z: zygotene spermatocyte, D: diplotene

spermatocyte, RS: round spermatid, ES: elongating spermatid. Scale bar, 5 μm. (g)

Quantification of spermatid numbers in seminiferous tubules from spermatid stages 1–16. N =

15 tubules per stage and per genotype; 3 mice per genotype were used and 5 tubules per stage

and per mouse were counted. (h) Mitochondrial sheath staining by Mito-tracker in sperm

collected from the caudal epididymis. Scale bar, 5 μm. (i, j) The percent of progressive and

motile spermatozoa in 9-week-old WT and *Tmc7*$^{-/-}$ mice. Three mice were used. Images are

representative of at least three independent experiments. The results are presented as the mean

± S.D., and two-sided unpaired Student's t-tests were used to calculate the P-values. (*P <

0.05, **P < 0.01, ***P < 0.001).

Figure 3. Acrosome biogenesis is impaired starting from the Golgi phase in *Tmc7*$^{-/-}$

spermatids. (a) Immunofluorescence staining of PNA (green) in the phases of acrosome

biogenesis in 9-week-old WT and *Tmc7*$^{-/-}$ spermatids. Nuclei were stained with DAPI (blue).

The phases of acrosome biogenesis and corresponding spermiogenesis steps are the Golgi

phase (steps 1–3), cap phase (steps 4–7), acrosome phase (steps 8–12), and maturation phase

(steps 13–16). (b) TEM images of testicular sections of 9-week-old WT and *Tmc7*$^{-/-}$ mice.

The red star represents the abnormal plasma membrane space and accumulated vesicles. N:

nucleus, Ac: acrosome; Apx: acroplaxome; pm: plasma membrane. G: Golgi complex; Scale

bar, 2 μm. (c) Immunofluorescence staining with the antibody against Gm130 in spermatids

from the testes of 9-week-old WT and *Tmc7*$^{-/-}$ mice. Scale bar, 2 μm. (d) Calculation of the

aspect ratio (height [X] over width [Y]) of 9-week-old WT and *Tmc7*$^{-/-}$ Golgi apparatuses

based on TEM images. N = 27, and 9 tubules in each testis were counted from 3 mice. (e)

Western blot analysis of protein levels of Golgi stress- associated proteins (Hsp47, Tfe3) in

9-week-old WT and *Tmc7*$^{-/-}$ testes (*n* = 3 per group). Image J was used to quantify the protein

level. (f) Immunofluorescence staining of Gopc (red) and PNA (green) and the quantitative

analysis of Gopc and PNA lectin colocalization in squashed tubules of 9-week-old WT and

Tmc7$^{-/-}$ mice. Nuclei were stained with DAPI (blue). Scale bar, 10 μm. Four mice were used,

and 200 cells were counted per mouse. Images are representative of at least three independent

experiments. The results are presented as the mean ± S.D., and two-sided unpaired Student's

t-tests were used to calculate the P-values. (*P < 0.05, **P < 0.01, ***P < 0.001).

Figure 4. Tmc7 localizes to the cis-Golgi, and this is required for maintaining Golgi

integrity. (a) Immunohistochemical staining with an antibody against Tmc7 in seminiferous

tubules of 9-week-old mouse testes. Scale bar, 50 μm. (b) Immunofluorescence staining of

Tmc7 (red) and Gm130 (green, a marker of the cis-Golgi), with DAPI indicating the nuclei in

seminiferous tubules of 9-week-old WT and $Tmc7^{-/-}$ testis. Scale bar, 10 μm. Co-localization

was quantified by Image J, and 20 cells were analyzed for the Pearson's R-value. (c)

Immunofluorescence staining of Tmc7 (red) and Tgn46 (green, a marker of the trans-Golgi),

with DAPI indicating the nuclei in the seminiferous tubule. Scale bar, 10 μm. (d)

Immunofluorescence staining of Tmc7 (red) and PNA (green, a marker of the acrosome), with

DAPI indicating the nuclei in the seminiferous tubule. Scale bar, 10 μm. (e) Western blot was

used to analyze the results of co-immunoprecipitation and showed the interaction between

Tmc7 and Gm130, P115, and Grasp65 in 9-week-old WT testes. (f, g) Immunofluorescence

staining and quantification of Gm130 (green), P115 (red) and Grasp65 (red) in testicular

sections of 9-week-old WT and $Tmc7^{-/-}$ mice. Scale bar, 50 μm. (h) Western blots of the

protein levels of Gm130 and the Gm130-interacting proteins P115 and Grasp65 in 9-week-old

WT and $Tmc7^{-/-}$ testes. Image J was used to quantify the protein level. Images are

representative of at least three independent experiments. The results are presented as the mean

± S.D., and two-sided unpaired Student's t-tests were used to calculate the P-values. (*P <

0.05, **P < 0.01, ***P < 0.001).

Fig 5. Tmc7 depletion impairs cellular homeostasis and leads to spermatid apoptosis.

(a) The Ca^{2+} levels were measured by flow cytometry in WT and $Tmc7^{-/-}$ germ cells isolated

from PD30 testes. (b) The pH levels were measured by flow cytometry in WT and $Tmc7^{-/-}$

germ cells isolated from PD30 testes. (c) The ROS levels were measured by flow cytometry

in WT and $Tmc7^{-/-}$ germ cells isolated from PD30 testes. (d) Western blot analysis of protein

levels of ER chaperone-associated proteins (P-Eif2α, Eif2α, Chop, and Grp78) in 9-week-old

WT and $Tmc7^{-/-}$ testes. Image J was used to quantify the protein level. Each value was

detected in three mice of each group. (e) TUNEL assay and the number of TUNEL-positive

cells per tubule in seminiferous tubules and cauda epididymis of 9-week-old WT and $Tmc7^{-/-}$

male mice. Scale bar, 50 μm. N = 60, 20 tubules in each testis and cauda epididymis were

counted, and three mice were used. (f) Apoptosis-related proteins (Cleaved-caspase3,

Caspase3, Bax, Bcl2) were detected by western blotting in 9-week-old WT and $Tmc7^{-/-}$ testes.

Image J was used to quantify the protein level. Images are representative of at least three

independent experiments. The results are presented as the mean ± S.D., and two-sided

unpaired Student's t-tests were used to calculate the P-values. (*P < 0.05, **P < 0.01, ***P <

0.001).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

a Ca²⁺ Fluorescence

b pH Fluorescence

c ROS Fluorescence

d $Tmc7^{+/+}$ $Tmc7^{+/+}$ $Tmc7^{-/-}$ $Tmc7^{-/-}$

P-elif2α — 38 kDa
Eif2α — 38 kDa
Chop — 27 kDa
Grp78 — 78 kDa
α-Tubulin — 55 kDa

e $Tmc7^{+/+}$ $Tmc7^{-/-}$

Testis

Cauda Epididymis

f $Tmc7^{+/+}$ $Tmc7^{-/-}$

Cleaved-caspase3 — 19 kDa
Caspase3 — 32 kDa
Bax — 21 kDa
Bcl-2 — 26 kDa
α-Tubulin — 55 kDa