Sac1 links phosphoinositide turnover to cryptococcal virulence

Elizabeth A. Gaylord[1], Hau Lam Choy[1,2], Guohua Chen[1,3], Sydney L. Briner[1], and Tamara L. Doering[1,*]

[1]Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO, USA

[2]Current position: L.E.K. Consulting, Boston, MA, USA

[3]Current position: Pfizer, Inc., Chesterfield, MO, USA

*Corresponding author: doering@wustl.edu

Running head: Sac1 links PI turnover to cryptococcal virulence

KEYWORDS

Cryptococcus neoformans, mycology, phosphoinositide, PI4P, phosphatase, Sac1, virulence

ABSTRACT

Cryptococcus neoformans is an environmentally-acquired fungal pathogen that causes over 140,000 deaths per

year. Cryptococcal infection occurs when infectious particles are deposited into the lung, where they encounter host

phagocytic cells. *C. neoformans* may be engulfed by these phagocytes, an important step of infection that leads to out-

comes ranging from termination of infection to cryptococcal dissemination. To study this critical process, we screened

approximately 4,700 cryptococcal gene deletion mutants for altered uptake, using primary mouse and human phagocytic

cells. Among the hits of these two screens, we identified 93 mutants with perturbed uptake in both systems, as well as

others with differences in uptake by only one cell type. We further screened the hits for changes in thickness of the cap-

sule, a protective polysaccharide layer around the cell which is an important cryptococcal virulence factor. The combina-

tion of our three screens yielded 45 mutants, including one lacking the phosphatidylinositol-4-phosphate phosphatase

Sac1. In this work, we implicate Sac1 in both host cell uptake and capsule production. We found that *sac1* mutants ex-

hibit lipid trafficking defects, reductions in secretory system function, and changes in capsule size and composition. Many

of these changes occur specifically in tissue culture media, highlighting the role of Sac1 phosphatase activity in respond-

ing to the stress of host-like conditions. Overall, these findings show how genome-scale screening can identify cellular

factors that contribute to our understanding of cryptococcal biology and demonstrate the role of Sac1 in determining

fungal virulence. (240/250)

IMPORTANCE

Cryptococcus neoformans is a fungal pathogen with significant impact on global health. Cryptococcal cells inhaled

from the environment are deposited into the lungs, where they first contact the human immune system. The interaction

between *C. neoformans* and host cells is critical because this step of infection can determine whether the fungal cells die

or proliferate within the human host. Despite the importance of this stage of infection, we have limited knowledge of

cryptococcal factors that influence its outcome. In this study, we identify cryptococcal genes that affect uptake by both

human and mouse cells. We also identify mutants with altered capsule, a protective coating that surrounds the cells to

shield them from the host immune system. Finally, we characterize the role of one gene, *SAC1*, in these processes. Overall,

this study contributes to our understanding of how *C. neoformans* interacts with and protects itself from host cells.

(147/150)

Abstract word count: 385

Text word count: 5250

INTRODUCTION

Cryptococcus neoformans is a fungal pathogen that causes significant mortality in immunocompromised individu-

als, particularly those with HIV/AIDS (1). Despite recent advances in healthcare access and treatments, cryptococcal men-

ingitis still leads to 19% of AIDS-related deaths (2). The interaction between *C. neoformans* and host phagocytes begins

when environmentally-acquired fungal cells are deposited in the lung. Subsequent phagocytosis may lead to fungal cell

death and termination of the infection. However, in some cases cryptococcal cells can withstand the pressures exerted by

phagocytes to survive and proliferate within phagolysosomes (3–6). In this scenario, engulfment of the yeast likely en-

hances cryptococcal dissemination from the lung to other sites (7). Due to the pivotal role of this initial phase of crypto-

coccal pathogenesis, extensive efforts have been devoted to studying the interactions of *C. neoformans* with host phago-

cytic cells.

C. neoformans produces multiple factors which modulate its interactions with host cells and thereby provide pro-

tection from phagocytosis. These include the cell wall and surrounding polysaccharide capsule, complex glycan-based

structures which change in size and composition in response to environmental conditions. Strains with perturbations of

these structures often exhibit increased uptake by phagocytic cells (8, 9), demonstrating the importance of capsule for

protecting against phagocytosis.

Various models have been used to interrogate cryptococcal-host interactions. These include invertebrate and ver-

tebrate animals, amoeba, and primary and immortalized cells (3, 10–16). Genetic screens based on libraries of mutant host

or fungal cells have identified gene products that modulate interactions of cryptococci with specific cell lines. In one ex-

ample, Chun and colleagues subjected cryptococcal cells to insertional mutagenesis and identified mutations that affected

their uptake by RAW264.7 macrophages (17). Srikanta et al. used RNA interference and high-throughput imaging to find

host regulatory genes which govern the interactions of *C. neoformans* with THP-1 cells and discovered 25 fungal-specific

regulators of phagocytosis (18). By applying high-throughput imaging to a *C. neoformans* gene deletion collection (19),

Santiago-Tirado et al. identified 56 fungal genes that affect uptake by THP-1 cells (9). A common strength of these studies

is the genetic tractability and accessibility of the immortalized cell lines used. However, such lines lack some features of

primary host cells (20–22), which may limit these approaches. Screens may also be constrained by the availability and size

of *Cryptococcus* mutant collections. For example, the last study cited screened a library of approximately 1,200 single gene

deletion mutants (9), whereas much larger libraries are currently available, increasing our power to identify novel pheno-

types.

This report incorporates several important departures from previous screens designed to elucidate *C. neoformans*-

host interactions. First, we screened a collection of 4,692 single gene deletion mutants, representing over eighty percent

of the non-essential genes in the cryptococcal genome. Second, we applied a flexible medium-throughput imaging method

that avoids pitfalls of earlier studies, such as difficulty distinguishing adherent versus internalized yeast cells (23). Third,

we screened using primary cells from both mouse (murine bone marrow-derived macrophages; BMDM) and human (mon-

ocyte-derived macrophages; HMDM), to take advantage of a model system while also examining the system of greatest

clinical interest. Finally, we reasoned that many of our hits would have altered capsules, so we further tested mutants that

demonstrated altered phagocytosis for changes in capsule thickness. Overall, we identified 45 deletion strains which ex-

hibited altered uptake in both systems as well as altered capsule size.

While it is known that capsule material traffics at least in part through the secretory pathway (24), few proteins

involved in this process have been characterized. To address this gap, we focused on mutants that might have aberrant

capsule trafficking. One critical process in the regulation of secretion is the subcellular distribution of phosphoinositides

(PI). Various cellular PI species are defined by the patterns of phosphorylation on their inositol head groups. These species

are maintained at distinct cellular compartments by kinases and phosphatases, which specifically and reversibly phosphor-

ylate and dephosphorylate inositol (25). In this way, PIs help define membrane identity and regulate membrane-protein

interactions (25), thus playing important roles in secretory trafficking (26, 27). We therefore were particularly interested

to note that one strain, which emerged as a hit in all three of our screens, lacks the gene *SAC1* (*CKF44_06080*). Sac1 is a

phosphatase that is highly conserved in eukaryotes and acts in phosphoinositide metabolism (28). Our observation that

cells lacking Sac1 are hypocapsular thus suggested that modulation of phosphoinositide turnover in the secretory pathway

influences capsule synthesis. Below we use biochemical, imaging, and molecular engineering approaches to establish the

mechanism of this interaction, linking alterations in lipid homeostasis to the production of a critical cryptococcal virulence

factor.

RESULTS

<u>Fungal genes that influence interactions with host cells</u>

To identify fungal gene products that affect interactions with host cells, we screened 4,692 *C. neoformans* single

gene deletion mutants (19). In our assay, fungal cells were stained with Lucifer Yellow, opsonized with serum, and then

incubated with primary phagocytes to allow engulfment. The samples were washed to remove unassociated cells and then

stained with membrane-impermeant Calcofluor White to label externally-associated fungi and propidium iodide to label

phagocytic cell nuclei. We assessed fungal engulfment as the phagocytic index, or number of internalized fungi (stained

with Lucifer Yellow but not Calcofluor White) per host cell (identified by nuclear staining) (23). We define the uptake score

as the ratio of a mutant's phagocytic index to that of wild-type *C. neoformans* (KN99).

We first applied our assay to screen the interactions of *C. neoformans* with murine bone marrow-derived macro-

phages (BMDM), normalizing to fungal cell number to account for differences in mutant growth rates (Figure 1A). Because

these deletion library plates do not include WT control cells, we initially compared strain uptake scores to the median for

each plate and used permissive cutoffs to call roughly 10% of the strains as hits (Table S1A). We then subjected the 458

strains that we identified with increased or decreased uptake (Table S1B) to a second round of BMDM screening (Table

S1C), this time comparing them to in-plate wild-type controls and using more stringent cutoffs to select hits (Figure 1B;

Table S1D). In parallel, we screened them using human monocyte-derived macrophages (HMDM) in place of BMDM (Figure

1C; Tables S1E and S1F).

Our second round of screening narrowed the initial hits to 93 mutants that significantly influenced host interac-

tions in both BMDM and HMDM cells (Figure 1A, overlap of red and blue circles; Table S1G). When we applied Gene

Ontology (GO) analysis to the corresponding genes, we found that genes encoding cellular factors related to cytoskeletal

interactions and intracellular protein localization and transport, among other categories, were particularly enriched rela-

tive to their abundance in the genome (Figure 1D; see Discussion).

Mutants with altered host interactions and capsule size

We were particularly interested in gene products that might act in capsule production, because of the many out-

standing questions that remain about this process (29). As the outermost layer of the cell, the capsule is fundamental in

mediating cryptococcal interactions (7). Therefore, we hypothesized that our uptake screen hits would be enriched for

mutants with altered capsule. To test this idea, we adapted an automated, fluorescence-based capsule quantification assay

(30) to accommodate medium-throughput screening. Of the 458 first round hits from the uptake screen, 131 indeed

showed capsule changes (Figure 1A), with 57 strains identified as hypocapsular and 74 as hypercapsular (Figure 1E; Tables

S2A, 2B). Validating the screen, this group contained multiple previously-identified capsule mutants, including 16 that were

hypocapsular and 3 that were hypercapsular. GO terms for the hits were particularly enriched for roles in capsule organi-

zation, as we expected, and for processes related to vesicular traffic, some of which were also enriched in our uptake

screen hits. Uniquely enriched for the capsule screen hits were ion transport and phosphoinositide binding (Figure 1F).

We identified 45 mutants which were hits in all three of our screens (Table S2C). The combination of perturbed

interactions with phagocytes and altered capsule thickness of these strains suggested that the deleted genes would be

important for virulence. We further speculated that the uptake phenotypes were related to surface changes in the cells,

which might be mechanistically elucidated by analysis of the deleted genes. Of the 45 triple hits, 18 had been previously

characterized. When we used homology to predict the subcellular localization of the remaining gene products, the largest

group (8 of 27) was predicted to reside in the secretory pathway. We suspected these genes might influence secretion of

capsule material.

Among the eight mutants likely impaired in secretion, we particularly noticed one strain that showed increased

phagocytosis by both BMDM and HMDM and a 61% reduction in capsule thickness compared to WT cells (Figure 2A). Its

missing sequence, *CKF44_06080*, encodes a protein with 38% amino acid identity to the phosphatidylinositol phosphatase

Sac1 of *Saccharomyces cerevisiae,* including 85% identity across a 20-amino acid region that encodes a conserved $CX_5R(RT)$

motif necessary for catalytic function (31). Interestingly, despite this similarity, ectopic overexpression of *S. cerevisiae* Sac1

did not restore the capsule defects of *sac1*Δ cells (Figure S1).

The phenotypes of *sac1*Δ cells, together with the unique biology of the capsule and its importance as a virulence

factor, led us to further investigate the role of Sac1 in *C. neoformans*. To pursue these studies, we first engineered a new

Sac1 deletion strain (*sac1*Δ) and complemented the mutant at the endogenous site (*SAC1*). We also identified the con-

served Sac1 catalytic motif by homology to *S. cerevisiae* Sac1 and inactivated it by replacing an aspartic acid residue with

alanine (*SAC1[456A]*, Figure 2B and S2) (32, 33). Using these strains, we confirmed that loss of Sac1 led to thinner capsules, a

phenotype which was maintained after normalization to the smaller diameters of the *sac1*Δ cells (Figure 2C, D). These

traits were phenocopied by the inactivated strain, showing that the phosphatase function of Sac1 is essential for capsule

production, while complementation restored WT phenotypes (Figure 2C, D). While imaging, we also observed striking

round structures in the *sac1* mutant cells (Figure 2C, right panels). These were distinguished from typical organelles, like

the vacuoles visible in WT and *SAC1* cells, by their circular shape, defined edge, and striking refractility and only occurred

in *sac1* mutants grown in host-like conditions (here defined as DMEM, 37°C, 5% CO_2). These structures will be further

discussed below.

<u>Sac1 is required for cryptococcal virulence</u>

Variations in both uptake and capsule size are associated with changes in pathogenicity (9, 34–40), so we suspected

that *sac*1 mutants would have impaired virulence. To test this, we infected mice intranasally and measured organ burden

at 14 days post-infection. Notably, while *sac1*Δ cells were recovered from the lungs, they remained close to the level of the

original inoculum. In sharp contrast, the lung burden of WT and *SAC1* infected animals increased roughly 2000-fold in this

period (Figure 3A). Cells were recovered from the brain in only a minority of mutant infections, although all WT-infected

animals had detectable brain burden by this time point (Figure 3B). We observed similar results in a time course study,

which demonstrated gradual accumulation of cryptococci in the lung up to 18 days post-infection (Figure S3A) and impaired

dissemination to the brain (Figure S3B). These results motivated us to define the mechanism(s) by which Sac1 influences

virulence.

<u>Sac1 controls substrate distribution and sensitivity to stressors</u>

Sac1 has not been characterized in *C. neoformans.* To test its influence on turnover of intracellular PI4P, we engi-

neered strains to express an mNeonGreen-tagged peptide that binds PI4P (FAPP1) and has been used to assess its locali-

zation (41). We then examined the abundance and localization of this fluorescent peptide (Figure 4A). Overall, the FAPP1

signal was brighter in *sac1*Δ cells than in WT, a trend supported by flow cytometry (Figure S4). This is consistent with a role

for Sac1 as a PI4P phosphatase, as loss of this protein would lead to an increase in its substrate. We also noted distinctive

bright perinuclear rings in the *sac1*Δ cells (especially in rich medium), which were more diffuse in WT cells. These corre-

spond to an endoplasmic reticulum distribution (42) and suggest aberrant accumulation of PI4P in the early secretory

pathway. Finally, the mutant strain exhibited intense puncta, absent from WT cells, which are consistent with PI4P accu-

mulation in the yeast Golgi (43).

Secretion is a vital cellular process, so its perturbation could have multiple deleterious effects on the cell, including

effects on the plasma membrane and cell wall. To examine this, we tested strain growth in various stressful conditions.

Although both *sac1* mutants grow slightly slower than WT or complemented cells even on rich medium, they showed much

worse growth on media containing membrane stressors (the detergent SDS and the sterol-binding antifungal amphotericin

B) or the cell wall stressor Congo Red (Figure 4B). These defects were exacerbated when cells were grown at 37°C. In all of

our assays, the complemented strain phenocopied WT and the inactivated construct phenocopied the deletion mutant.

<u>Sac1 regulates cellular lipids during the response to host-like conditions</u>

We sought to explain the capsule defect and stress sensitivities of *sac1* mutants based on the cellular conse-

quences of perturbed PI4P turnover. In other eukaryotes, PI4P is important for establishing the lipid composition of secre-

tory vesicles (44) and its turnover is critical for the progression of post-Golgi secretory traffic (45). We therefore examined

classes of molecules, including lipids, proteins, and capsule polysaccharides, that are directly or indirectly influenced by

this step of secretion. As noted above, we had observed formation of large internal structures in *sac1* mutant cells grown

under host-like conditions (Figure 2C), which were absent from WT and complemented cells. Because of their shape and

refractile appearance in light micrographs, we suspected they were at least in part composed of lipid. Consistent with this

suggestion, they appeared electron-lucent by transmission electron microscopy (Figure 5A, asterisks). We hypothesized

that these structures result from aberrant lipid handling within *sac1* mutant cells that is induced by the stress of host-like

conditions, since they are absent from mutant cells grown in rich medium (Figure 5A). Based on this idea, and our results

below, we termed them Sac1 lipid-accumulating bodies (SLBs).

SLBs resemble previously-characterized "supersized" lipid droplets identified in *S. cerevisiae* (46). Lipid droplets

are neutral lipid-containing structures that represent essential cellular energy reserves; these increase in abundance when

cryptococcal cells are grown in host-like conditions (47). When we stained for neutral lipids using Nile Red, WT cryptococcal

cells showed dispersed punctate signal, while *sac1Δ* cells showed concentrated central staining, frequently colocalized

with SLBs (Figure 5B,C) and thus supporting the presence of neutral lipids within them.

We recently demonstrated that mislocalization of the important fungal lipid ergosterol leads to accumulation of

lipid droplets (47). We had also noted that *sac1* mutants were more sensitive than WT cells to Amphotericin B, which binds

ergosterol (Figure 4B). Consequently, we wondered whether ergosterol localization in *sac1Δ* cells was perturbed. When

we stained WT cells grown in host-like conditions with filipin, which binds non-esterified sterols, we saw the typical ring-

staining expected for a component of the plasma membrane (Figure 5D). By contrast, mutant cells exhibited increased

internal staining, particularly bordering the SLBs (see Discussion). When we tested filipin and Nile Red together, both stains

localized to SLBs in the *sac1* mutants, with a ring of filipin surrounding the Nile Red stain (Figure S5). In contrast, WT cells

showed peripheral filipin staining and Nile Red staining dispersed throughout the cell.

In testing cell growth, we fortuitously discovered that supplementation of host-like conditions (DMEM, 37°C, 5%

CO_2) with various exogenous fatty acids partially restored the growth of *sac1Δ* cells (Figure 6 and Supplemental Figure S6);

this effect was dose-dependent (shown for myristate in Figure 6A). This supplementation also prevented the formation of

SLBs (Figure 6B, bottom right panel). When we visualized capsule in myristate-supplemented cells, we made several ob-

servations. First, the addition of the BSA vehicle alone led to a slight capsule size increase, of similar magnitude, in both

cell types. Second, myristate supplementation yielded significant capsule reduction in both strains, which was more dra-

matic in the mutant (Figure 6B). Lipid supplementation thus restores growth and prevents SLAB formation in *sac1Δ* cells

but does not rescue capsule size (see Discussion).

Sac1 impacts protein secretion and localization

Proteins are critical secretory cargo, so it is likely that accumulation of PI4P in the secretory pathway would affect

their transport and secretion. To assess this, we measured protein secretion directly, using inducible acid phosphatase

production as a proxy for general protein secretion (24). Upon entry into phosphate-depleted conditions, WT, *SAC1*, and

both *sac1* mutants began producing detectable levels of acid phosphatase within one hour. While the general patterns of

secretion were similar, the mutants produced less acid phosphatase on a per cell basis throughout the time course (Figure

7A), suggesting an overall reduction in protein secretion.

Perturbations in PI4P levels may affect not only proteins secreted outside the cell, but also those resident in the

secretory system, including multiple capsule biosynthetic proteins in the Golgi apparatus. We hypothesized that the loss

of PI4P turnover could result in failure to correctly localize these enzymes and thereby explain the altered capsule of *sac1*

mutants. To investigate this question, we localized the best-characterized capsule biosynthetic enzyme, Cxt1 (48), in *sac1Δ*

and WT cells grown in host-like conditions. WT cells displayed clear puncta of Cxt1-mNeonGreen, consistent with the ex-

pected Golgi localization of this protein. In contrast, staining in *sac1Δ* cells was heterogenous, with various cell populations

displaying punctate, diffuse, or little to no fluorescence (Figure 7B, C).

Sac1 affects capsule secretion and composition

Capsule glycans constitute a class of secretory cargo which is unique to *C. neoformans*. The predominant capsule

polysaccharide, glucuronoxylomannan (GXM), traverses the secretory pathway prior to extracellular release (24). We hy-

pothesized that GXM secretion, like protein secretion, is reduced in *sac1Δ* cells, causing their thin capsules. To test this, we

quantified GXM secretion at intervals after transition to host-like conditions. Importantly, WT and *sac1* mutants began to

produce detectable levels of GXM concurrently, showing that the mutants respond normally to the environmental signals

that trigger capsule production (Figure 7D). However, the overall amount of GXM they secreted was markedly reduced.

Reduced GXM secretion could certainly explain the small capsules of *sac1* mutant cells. Another reason for altered

capsule could be mislocalization of glycosyltransferases (such as we observed with Cxt1), since compositional differences

in the polysaccharide also affect capsule size and architecture (8). To investigate this, we analyzed the composition of GXM

purified from WT, *sac1Δ*, and *SAC1* cells. Compared to the WT and *SAC1* controls, *sac1Δ* GXM showed a substantial (~2-

fold) increase in the relative proportion of glucuronic acid, as well as a modest increase in xylose (Figure 7E). These results

were supported by linkage analysis, in which we observed a reduction in 3-linked mannopyrannosyl residues as a fraction

of the total (Supplemental Table S3), indicating fewer unmodified backbone mannose residues.

DISCUSSION

How *C. neoformans* interacts with host cells is a critical determinant of infection. In this study, we screened 4,692

cryptococcal deletion strains and identified 93 that demonstrated altered uptake by both mouse and human primary phag-

ocytes. GO analysis showed that this set was enriched in sequences that encode proteins with roles in cytoskeletal inter-

actions, protein transport, and organization of the cell wall and capsule. This is consistent with our understanding that the

outer layers of the cell and secreted factors modulate interactions with host cells. We also identified strains with defects

in only one host system. Why the species differ in this regard could be a fruitful area for future research.

Capsule is a critical factor in the interaction between *C. neoformans* and host cells, but the mechanisms of its

synthesis and transport remain largely unknown. To approach this question, we performed a secondary screen for altered

capsule thickness, which yielded 131 mutants. For comparison, a previous screen of 1,201 deletion strains, using colony

morphology as the primary screen, yielded 16 capsule-related genes (19). Furthermore, our hits represent approximately

one third of the original set of uptake mutants that we tested. This shows that our strategy of using these mutants as the

starting population to screen for capsule defects was highly successful in enriching for capsule phenotypes. Notably, 42 of

the hits are annotated as encoding hypothetical proteins (49), which may be subjects for future investigation. The others

were enriched for GO terms related to cell wall and capsule organization, which validated our methods, and for terms

involving transporter activity, intracellular transport, and phosphatidylinositol binding. As our results with Sac1

demonstrate, the study of such processes, which are not directly related to capsule production, may still yield insight into

capsule biosynthesis.

Interestingly, over half of our capsule screen hits were hypercapsular strains. This highlights an important strength

of our image-based technique, as prior methods (including colony morphology, antibody blotting, or volume-based meth-

ods) identify strains that lack capsule more readily than hypercapsular cells (50). That said, image-based screens may also

have weaknesses. For example, mutants with general morphological defects may register as false positives, as cells that

fail to divide might appear to be one large cell with a capsule defect. Further, this method cannot identify mutants that are

completely acapsular, as they will not bind antibody.

Of the 45 gene products identified as hits in all three of our screens, the largest subset of the uncharacterized hits

was predicted to be involved in the secretory pathway. We focused on one of these hits, Sac1, a PI4P phosphatase encoded

by *CKF44_06080.* The *CNAG_06080* gene (which encodes Sac1 in the closely-related reference strain H99) was previously

identified as a site where random integration of a drug marker yielded an attenuated *C. neoformans* strain (51). Like our

KN99 mutant, this strain showed reduced virulence, although direct comparison is limited because the mutations, mouse

models, and experimental designs differ. Other investigators had proposed that this gene was essential, because they were

unable to delete it (52). We expect this difference from our findings was due to the challenge of working with cells that

have significant defects, although it could also reflect different strain backgrounds (H99S vs KN99) (53, 54).

By expressing and localizing a fluorescent PI4P-binding peptide (41, 55, 56), we supported a role for Sac1 in PI4P

turnover in *C. neoformans*. Although we did not test this activity directly, based on homology to the well-characterized *S.*

cerevisiae Sac1 protein, conservation of Sac1 function throughout eukaryotes, and our binding data, we believe PI4P is the

main substrate of Sac1 in *C. neoformans*. It is curious that despite its high amino acid similarity to the *S. cerevisiae* Sac1

protein, over-expression of the latter did not rescue the defects of *C. neoformans sac1Δ* cells; this result may warrant

further experimentation.

When s*ac1* mutants are grown in host-like medium, they develop large structures which we termed SLBs. The

interiors of SLBs stain with Nile Red, which binds neutral lipids, and the edges stain with the sterol-binding dye filipin. SLBS

thus share certain features with lipid droplets, which are composed of a neutral lipid core of mainly triacylglycerols and

sterol esters surrounded by a phospholipid monolayer (57). More specifically, SLBs resemble "supersized" lipid droplets,

which form in model yeast strains defective in phospholipid metabolism (46, 58). This phenomenon has also been observed

in *C. neoformans* following treatment with the antidepressant sertraline (59). However, our filipin staining suggests the

presence of ergosterol or sterol intermediates (60) on the perimeter of the SLBs, which has not been reported for regular

or supersized lipid droplets; accordingly we have not adopted this terminology.

Ergosterol is distributed in a steeply increasing gradient from the ER to the plasma membrane (61). One model

proposes that PI4P turnover by Sac1 maintains this gradient (62). If this mechanism occurs in *C. neoformans,* loss of PI4P

turnover would likely result in the inability to shuttle ergosterol up its concentration gradient with consequent accumula-

tion in the ER. Another model suggests that the ergosterol gradient depends on membrane phospholipid composition (61).

In either case, loss of Sac1 could perturb lipid transport and homeostasis and lead to mislocalization of ergosterol.

PI4P is a key signaling lipid, that serves as a pro-secretory marker and modulates the exit of secretory traffic from

the *trans*-Golgi (27, 44, 63). We propose that the lack of PI4P turnover in *sac1* mutants leads to its accumulation in the

early portions of the cryptococcal secretory system. This would be consistent with our observations of reduced protein

secretion and mislocalization of the Golgi-resident protein Cxt1 in *sac1*Δ cells. The Golgi serves as a central hub for traf-

ficking of both proteins and lipids, and in doing so plays a key role in membrane trafficking and lipid metabolism (64). We

hypothesize that SLAB formation in *sac1* mutants represents the failure of lipid trafficking under stressful, nutrient-limited

host-like conditions, leading to the accumulation of lipid cargo and/or lipid metabolic precursors. SLAB formation and the

poor growth of *sac1*Δ cells is prevented by fatty acid supplementation, which somehow bypasses the defect caused by

PI4P accumulation and allows the cells to grow normally. How this occurs will be an interesting question for future research.

Importantly, while fatty acid supplementation rescues SLAB formation in *sac1*Δ cells, capsule production is not restored.

This key finding shows that the adverse effects of Sac1 loss occur in divergent pathways.

Capsule material is exported in secretory vesicles (24), so defects in anterograde traffic would also impair this

process. Beyond reduced GXM secretion, which is consistent with this expectation, we observed alterations in GXM com-

position: the mannose backbone had a large increase in glucuronic acid substitution and a slight increase in xylose. We

hypothesize that altered localization of capsule biosynthetic enzymes changes their access to nascent capsule material and

precursor molecules. It is not known whether capsule polysaccharides traverse the cell as subunits or extended polymers

(65, 66), but in either case increased exposure to biosynthetic enzymes could result in compositional changes such as we

observed. Both the reduction of capsule size and changes in composition in *sac1* mutants likely contribute to their in-

creased uptake by host phagocytes.

In this study, we identified *C. neoformans* mutants impaired in host interactions with primary cells and assessed the hits for capsule defects. This strategy greatly increased the number of sequences known to influence capsule production, including many previously uncharacterized genes. Analysis of these genes and the corresponding mutants will help us understand the processes required for synthesis of this central virulence factor, as exemplified in this work on Sac1. Importantly, two-thirds of our original uptake hits did not have altered capsule thickness, implicating additional mechanisms in this fundamental host-pathogen interaction. Further studies will be required to determine what leads to the altered uptake of these strains, which may include changes to the cell wall or surface mannoproteins (29) or structural alterations in the capsule that do not impact its thickness.

METHODS

Strain construction, validation, and growth

sac1Δ, *SAC1* complement, and *SAC1*[D456A] strains were made with split-marker biolistic transformation (67) in KN99α. Cxt1 was tagged with mNEONGREEN (pBHM2404, Addgene #173441, a gift from Hiten Madhani) using CRISPR with short homology-directed repair (68). FAPP1-mNeonGreen (from pRS406-PHO5-GFP-hFAPP1_PH_domain, Addgene #58723, a gift from Tim Levine) and *S. cerevisiae SAC1* strains were similarly inserted in the Safe Haven 2 region (68, 69).

Transformants were validated by antifungal resistance, PCR and/or whole genome sequencing. Expression of *S. cerevisiae* S288C *SAC1* was confirmed by extracting total RNA with TRIzol Reagent, synthesizing cDNA using SuperScript III (Thermo Fisher, #18080400), and PCR-amplifying a unique region of the gene (Fig. S1B).

For all assays except screening, *C. neoformans* strains were grown on YPD plates (2 days, 30°C) and single colonies were inoculated into YPD and grown overnight (30°C, shaking). For capsule inductions, cells were washed in PBS, inoculated in DMEM at 10^6 cells/mL, and grown for 16-24 hours (static, 37°C, 5% CO_2). For screening, cells were cultured from glycerol stocks in 96-well plates, grown overnight in YPD, and then sub-cultured and grown 24 hours in YPD. Detailed methods, including stress media and lipid supplementation, are in Supplemental Materials.

Screening

For uptake screening, C57BL/6J mouse BMDMs were harvested as in (70) and differentiated for 9 days before isolation. HMDMs were from anonymous donors. PBMCs were isolated by Ficoll-Paque centrifugation and differentiated for 6 days. The screen assay was modified from (23). Briefly, opsonized Lucifer Yellow-stained fungi were added to host

cells at an MOI of 20 (BMDM) or 30 (HMDM) and the cultures were co-incubated for one hour and washed. Adherent cells were stained with Calcofluor White; wells were washed, fixed, and permeabilized; and host cell nuclei were stained with propidium iodide before imaging on a BioTek Cytation 3. The uptake index (internalized cells per macrophage) was corrected for the number of fungal cells added and then normalized to the plate median (first round) or KN99 value (later rounds).

For capsule screening, strains identified as hits in the first round BMDM uptake screen were assessed for capsule size differences using slight modifications of our previous method (30). For this, cells were induced as above, capsules were stained with anti-GXM mAb-302-AF488, and cell walls were stained with Calcofluor White. Images were collected as above, and capsule thickness measured as in (31).

Further details of methods and hit selection for both screens are in Supplemental Material.

Virulence studies

All animal protocols were approved by the Washington University Institutional Animal Care and Use Committee (Protocol #20-0108) and care was taken to minimize animal handling and discomfort. 6-8-week-old female C57BL/6 mice (The Jackson Laboratory) were anesthetized by subcutaneous injection of 1.20 mg ketamine and 0.24 mg xylazine in 110 µL sterile PBS. For 14-day infections, mice were intranasally infected with 5×10^4 cryptococcal cells. For time course infection, mice were infected with 1.25×10^4 cryptococcal cells and sacrificed at 6, 12, and 18-days post infection. For both infections, the lungs and brains were harvested, homogenized, and plated on YPD agar to assess fungal organ burden.

Assays

Acid phosphatase secretion was measured as in (24). 10^7 cells/mL cultures were grown in MM-KCl (0.5% KCl, 15 mM glucose, 10 mM $MgSO_4 \cdot 7H_2O$, 13 mM glycine, 3.0 µM thiamine). Supernatants were sampled, incubated with substrate buffer (2.5 mM para-nitrophenylphosphate in 50 mM acetate buffer, pH 4.0) at 37°C for 5 minutes, and stopped with 800 µL saturated Na_2CO_3. Acid phosphatase secretion was quantified as $A_{420}/OD_{600.}$

GXM ELISAs were performed as in (48). Cultures were induced in DMEM,and the supernatants filtered (0.2 µm). Flat-bottom Immulon 1B plates (Thermo Scientific) were coated with 1 µg/ml anti-GXM antibodies 339 and F12D2γ, washed with PBS with 0.5% TWEEN 20 (PBST), and blocked for 90 min with 1% BSA in PBST. Filtrates were diluted in PBST and incubated on the plate for 90 min. The plate was washed three times, probed with HRP-labeled mAb 339/F12D2γ, incubated with KPL TMB Microwell Peroxidase Substrate (SeraCare), developed, and quantified using A_{450}.

<u>Imaging</u>

For capsule visualization, cells were induced as above, mixed with India ink in PBS (1:2, v:v), and imaged by light

microscopy. Capsules of at least 96 cells from randomly-chosen fields were measured using ImageJ. Nile Red and filipin

staining was performed as in (47), and viewed on a ZEISS Axio Imager M2 fluorescence microscope. FAPP1-mNEONGREEN

and Cxt1-mNEONGREEN stains were imaged with a Zeiss LSM880 confocal microscope. Morphological classification of

Cxt1-mNeonGreen and Nile Red staining was performed on at least 94 cells from randomly-selected fields. Electron mi-

croscopy was as in (47) with one change detailed in Supplement.

<u>GXM analysis</u>

GXM was isolated using precipitation with hexadecyltrimethylammonium bromide as in (71). Isolated material

was hydrolyzed with trifluoroacetic acid (3 hr, 100°C) and composition assessed on a Dionex ICS-6000 instrument. Linkage

analysis was performed by the Complex Carbohydrate Research Center (method modified from (72)).

ACKNOWLEDGEMENTS

We thank Nicole Howard and Suhas Bobba for assistance with host cells, Todd Fehniger and lab for LRS chambers

for HMDM isolation, Yan Xie for statistical analysis, Wandy Beatty for electron microscopy, and Tim Levine for plasmid

sequences. We appreciate the assistance of Jessica Tuck and Alex Caron with phenotypic characterization and microscopy

studies. We thank Doering lab members for discussions, and particularly Liza Loza, Daphne Boodwa-Ko, Thomas Hurtaux,

and Zanetta Chang for comments on the manuscript.

This work was supported by NIH grants AI140979, AI135012, and AI178330 to T.L.D. E.A.G and H.L.C were partially

supported by T32 GM007067, and E.A.G. by a Sondra Schlesinger Graduate Fellowship from the Department of Molecular

Microbiology, Washington University School of Medicine. The cryptococcal deletion library, developed by the Madhani lab

with funding from R01AI100272, was purchased from the Fungal Genetics Stock Center. The CCRC is supported by the U.S.

Department of Energy, Office of Science, Basic Energy Sciences, grant number DE-SC0015662 to Parastoo Azadi.

REFERENCES

1. Denning DW. 2024. Global incidence and mortality of severe fungal disease. Lancet Infect Dis https://doi.org/10.1016/S1473-3099(23)00692-8.

2. Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, Meya DB, Chiller TM, Boulware DR. 2022. The global burden of HIV-associated cryptococcal infection in adults in 2020: a modelling analysis. Lancet Infect Dis 22:1748–1755.

3. Feldmesser M, Kress Y, Novikoff P, Casadevall A. 2000. *Cryptococcus neoformans* is a facultative intracellular pathogen in murine pulmonary infection. Infect Immun 68:4225–4237.

4. Tucker SC, Casadevall A. 2002. Replication of *Cryptococcus neoformans* in macrophages is accompanied by phagosomal permeabilization and accumulation of vesicles containing polysaccharide in the cytoplasm. Proc Natl Acad Sci U S A 99:3165–3170.

5. Del Poeta M. 2004. Role of phagocytosis in the virulence of *Cryptococcus neoformans*. Eukaryot Cell 3:1067–1075.

6. Osterholzer JJ, Surana R, Milam JE, Montano GT, Chen G-H, Sonstein J, Curtis JL, Huffnagle GB, Toews GB, Olszewski MA. 2009. Cryptococcal urease promotes the accumulation of immature dendritic cells and a non-protective T2 immune response within the lung. Am J Pathol 174:932–943.

7. Gaylord EA, Choy HL, Doering TL. 2020. Dangerous liaisons: Interactions of *Cryptococcus neoformans* with host phagocytes. Pathog Basel Switz 9:891.

8. Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. UDP-glucuronic acid transport is required for virulence of *cryptococcus neoformans*. mBio 9:e02319-17.

9. Santiago-Tirado FH, Peng T, Yang M, Hang HC, Doering TL. 2015. A Single Protein S-acyl Transferase Acts through Diverse Substrates to Determine Cryptococcal Morphology, Stress Tolerance, and Pathogenic Outcome. PLoS Pathog 11.

10. Steenbergen JN, Shuman HA, Casadevall A. 2001. *Cryptococcus neoformans* interactions with amoebae suggest an

explanation for its virulence and intracellular pathogenic strategy in macrophages. Proc Natl Acad Sci 98:15245–

15250.

11. Mylonakis E, Moreno R, El Khoury JB, Idnurm A, Heitman J, Calderwood SB, Ausubel FM, Diener A. 2005. *Galleria*

mellonella as a model system to study *Cryptococcus neoformans* pathogenesis. Infect Immun 73:3842–3850.

12. Mukaremera L, McDonald TR, Nielsen JN, Molenaar CJ, Akampurira A, Schutz C, Taseera K, Muzoora C, Meintjes G,

Meya DB, Boulware DR, Nielsen K. 2019. The mouse inhalation model of *Cryptococcus neoformans* infection reca-

pitulates strain virulence in humans and shows that closely related strains can possess differential virulence. Infect

Immun 87:e00046-19.

13. Syme RM, Spurrell JCL, Amankwah EK, Green FHY, Mody CH. 2002. Primary dendritic cells phagocytose *Cryptococ-*

cus neoformans via mannose receptors and Fcγ receptor II for presentation to T lymphocytes. Infect Immun

70:5972–5981.

14. Alvarez M, Casadevall A. 2006. Phagosome extrusion and host-cell survival after *Cryptococcus neoformans* phagocy-

tosis by macrophages. Curr Biol 16:2161–2165.

15. Tenor JL, Oehlers SH, Yang JL, Tobin DM, Perfect JR. 2015. Live imaging of host-parasite interactions in a zebrafish

infection model reveals cryptococcal determinants of virulence and central nervous system invasion. mBio

6:e01425-15.

16. Srikanta D, Yang M, Williams M, Doering TL. 2011. A Sensitive High-Throughput Assay for Evaluating Host-Pathogen

Interactions in *Cryptococcus neoformans* Infection. PLOS ONE 6:e22773.

17. Chun CD, Madhani HD. 2010. Ctr2 links copper homeostasis to polysaccharide capsule formation and phagocytosis

inhibition in the human fungal pathogen *Cryptococcus neoformans*. PLOS ONE 5:e12503.

18. Srikanta D, Hole CR, Williams M, Khader SA, Doering TL. 2017. RNA interference screening reveals host CaMK4 as a

regulator of cryptococcal uptake and pathogenesis. Infect Immun 85:e00195-17.

19. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic genetic analysis of virulence in the

human fungal pathogen *Cryptococcus neoformans*. Cell 135:174–188.

20. Kaur G, Dufour JM. 2012. Cell lines: Valuable tools or useless artifacts. Spermatogenesis 2:1–5.

21. Merly L, Smith SL. 2017. Murine RAW 264.7 cell line as an immune target: are we missing something? Im-

munopharmacol Immunotoxicol 39:55–58.

22. Pan C, Kumar C, Bohl S, Klingmueller U, Mann M. 2009. Comparative proteomic phenotyping of cell lines and pri-

mary cells to assess preservation of cell type-specific functions. Mol Cell Proteomics MCP 8:443–450.

23. Chang AL, Hole CR, Doering TL. 2019. An automated assay to measure phagocytosis of *Cryptococcus neoformans*.

Curr Protoc Microbiol 53:e79.

24. Yoneda A, Doering TL. 2006. A Eukaryotic Capsular Polysaccharide Is Synthesized Intracellularly and Secreted via

Exocytosis. Mol Biol Cell 17:5131–5140.

25. Baba T, Balla T. 2020. Emerging roles of phosphatidylinositol 4-phosphate and phosphatidylinositol 4,5-bisphos-

phate as regulators of multiple steps in autophagy. J Biochem (Tokyo) 168:329–336.

26. Di Paolo G, De Camilli P. 2006. Phosphoinositides in cell regulation and membrane dynamics. Nature. Nature Pub-

lishing Group.

27. Santiago-Tirado FH, Bretscher A. 2011. Membrane-trafficking sorting hubs: cooperation between PI4P and small

GTPases at the trans-Golgi network. Trends Cell Biol 21:515–525.

28. Del Bel LM, Brill JA. 2018. Sac1, a lipid phosphatase at the interface of vesicular and nonvesicular transport. Traffic

19:301–318.

29. Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: The outer layers of *Cryptococcus neoformans*.

Mem Inst Oswaldo Cruz 113:1–8.

30. Reuwsaat JCV, Agustinho DP, Motta H, Chang AL, Brown H, Brent MR, Kmetzsch L, Doering TL. 2021. The transcrip-

tion factor Pdr802 regulates titan cell formation and pathogenicity of *Cryptococcus neoformans*. mBio 12:e03457-

20.

31. Hughes WE, Cooke FT, Parker PJ. 2000. Sac phosphatase domain proteins. Biochem J 350:337–352.

32. Kearns BG, McGee TP, Mayinger P, Gedvilaite A, Phillips SE, Kagiwada S, Bankaitis VA. 1997. Essential role for diacyl-

glycerol in protein transport from the yeast Golgi complex. Nature 387:101–105.

33. Del Bel LM, Griffiths N, Wilk R, Wei H-C, Blagoveshchenskaya A, Burgess J, Polevoy G, Price JV, Mayinger P, Brill JA.

2018. The phosphoinositide phosphatase Sac1 regulates cell shape and microtubule stability in the developing *Dro-*

sophila eye. Dev Camb Engl 145:dev151571.

34. Chang YC, Kwon-Chung KJ. 1994. Complementation of a capsule-deficient mutation of *Cryptococcus neoformans*

restores its virulence. Mol Cell Biol 14:4912–4919.

35. Chang YC, Penoyer LA, Kwon-Chung KJ. 1996. The second capsule gene of *Cryptococcus neoformans*, CAP64, is es-

sential for virulence. Infect Immun 64:1977–1983.

36. Chang YC, Kwon-Chung KJ. 1998. Isolation of the third capsule-associated gene, *CAP60*, required for virulence in

Cryptococcus neoformans. Infect Immun 66:2230–2236.

37. Chang YC, Kwon-Chung KJ. 1999. Isolation, characterization, and localization of a capsule-associated gene, *CAP10*,

of *Cryptococcus neoformans*. J Bacteriol 181:5636–5643.

38. Chun CD, Brown JCS, Madhani HD. 2011. A major role for capsule-independent phagocytosis-inhibitory mecha-

nisms in mammalian infection by *Cryptococcus neoformans*. Cell Host Microbe 9:243–251.

39. Bojarczuk A, Miller KA, Hotham R, Lewis A, Ogryzko NV, Kamuyango AA, Frost H, Gibson RH, Stillman E, May RC,

Renshaw SA, Johnston SA. 2016. *Cryptococcus neoformans* intracellular proliferation and capsule size determines

early macrophage control of infection. Sci Rep 6:1–15.

40. Reuwsaat JCV, Doering TL, Kmetzsch L. 2021. Too much of a good thing: Overproduction of virulence factors impairs

cryptococcal pathogenicity. Microb Cell 8:108–110.

41. Levine TP, Munro S. 2002. Targeting of Golgi-specific pleckstrin homology domains involves both PtdIns 4-kinase-

dependent and -independent components. Curr Biol CB 12:695–704.

42. Horianopoulos LC, Lee CWJ, Hu G, Caza M, Kronstad JW. 2021. Dnj1 promotes virulence in *Cryptococcus neofor-*

mans by maintaining robust endoplasmic reticulum homeostasis under temperature stress. Front Microbiol 12.

43. Wang ZA, Griffith CL, Skowyra ML, Salinas N, Williams M, Maier EJ, Gish SR, Liu H, Brent MR, Doering TL. 2014.

Cryptococcus neoformans dual GDP-mannose transporters and their role in biology and virulence. Eukaryot Cell

13:832–842.

44. Johansen J, Ramanathan V, Beh CT. 2012. Vesicle trafficking from a lipid perspective. Cell Logist 2:151–160.

45. Ling Y, Hayano S, Novick P. 2014. Osh4p is needed to reduce the level of phosphatidylinositol-4-phosphate on secre-

tory vesicles as they mature. Mol Biol Cell 25:3389–3400.

46. Fei W, Shui G, Zhang Y, Krahmer N, Ferguson C, Kapterian TS, Lin RC, Dawes IW, Brown AJ, Li P, Huang X, Parton RG,

Wenk MR, Walther TC, Yang H. 2011. A role for phosphatidic acid in the formation of "supersized" lipid droplets.

PLoS Genet 7.

47. Choy HL, Gaylord EA, Doering TL. 2023. Ergosterol distribution controls surface structure formation and fungal

pathogenicity. mBio 14:e01353-23.

48. Klutts JS, Doering TL. 2008. Cryptococcal xylosyltransferase 1 (Cxt1p) from *Cryptococcus neoformans* plays a direct

role in the synthesis of capsule polysaccharides. J Biol Chem 283:14327–14334.

49. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D, Warrenfeltz S, Aurrecoechea C, Chris-

tian J. Stoeckert J, Kissinger JC, Roos DS, Hertz-Fowler C. 2018. FungiDB: an integrated bioinformatic resource for

fungi and oomycetes. J Fungi 4.

50. Boodwa-Ko D, Doering TL. 2024. A quick reCAP: Discovering *Cryptococcus neoformans* capsule mutants. J Fungi Ba-
sel Switz 10:114.

51. Arras SDM, Chitty JL, Blake KL, Schulz BL, Fraser JA. 2015. A genomic safe haven for mutant complementation in
Cryptococcus neoformans. PLoS ONE 10:e0122916.

52. Jin J-H, Lee K-T, Hong J, Lee D, Jang E-H, Kim J-Y, Lee Y, Lee S-H, So Y-S, Jung K-W, Lee D-G, Jeong E, Lee M, Jang Y-B,
Choi Y, Lee MH, Kim J-S, Yu S-R, Choi J-T, La J-W, Choi H, Kim S-W, Seo KJ, Lee Y, Thak EJ, Choi J, Averette AF, Lee Y-H,
Heitman J, Kang HA, Cheong E, Bahn Y-S. 2020. Genome-wide functional analysis of phosphatases in the pathogenic
fungus *Cryptococcus neoformans*. 1. Nat Commun 11:4212.

53. Nielsen K, Cox GM, Wang P, Toffaletti DL, Perfect JR, Heitman J. 2003. Sexual cycle of *Cryptococcus neoformans* var.
grubii and virulence of congenic a and α isolates. Infect Immun 71:4831–4841.

54. Fernandes KE, Fraser JA, Carter DA. 2022. Lineages derived from *Cryptococcus neoformans* type strain H99 support
a link between the capacity to be pleomorphic and virulence. mBio 13:e00283-22.

55. Godi A, Campli AD, Konstantakopoulos A, Tullio GD, Alessi DR, Kular GS, Daniele T, Marra P, Lucocq JM, Matteis
MAD. 2004. FAPPs control Golgi-to-cell-surface membrane traffic by binding to ARF and PtdIns(4)P. 5. Nat Cell Biol
6:393–404.

56. Dowler S, Currie RA, Campbell DG, Deak M, Kular G, Downes CP, Alessi DR. 2000. Identification of pleckstrin-homol-
ogy-domain-containing proteins with novel phosphoinositide-binding specificities. Biochem J 351:19–31.

57. Walther TC, Chung J, Farese RV. 2017. Lipid droplet biogenesis. Annu Rev Cell Dev Biol 33:491–510.

58. Fei W, Shui G, Gaeta B, Du X, Kuerschner L, Li P, Brown AJ, Wenk MR, Parton RG, Yang H. 2008. Fld1p, a functional
homologue of human seipin, regulates the size of lipid droplets in yeast. J Cell Biol 180:473–482.

59. Breuer MR, Dasgupta A, Vasselli JG, Lin X, Shaw BD, Sachs MS. 2022. The antidepressant sertraline induces the for-
mation of supersized lipid droplets in the human pathogen *Cryptococcus neoformans*. J Fungi 8:642.

60.   Solanko KA, Modzel M, Solanko LM, Wüstner D. 2016. Fluorescent sterols and cholesteryl esters as probes for intra-

520   cellular cholesterol transport. Lipid Insights 8:95–114.

521   61.   Menon AK. 2018. Sterol gradients in cells. Curr Opin Cell Biol 53:37–43.

522   62.   von Filseck JM, Vanni S, Mesmin B, Antonny B, Drin G. 2015. A phosphatidylinositol-4-phosphate powered ex-

523   change mechanism to create a lipid gradient between membranes. 1. Nat Commun 6:6671.

524   63.   Santiago-Tirado FH, Legesse-Miller A, Schott D, Bretscher A. 2011. Phosphatidylinositol 4-phosphate and Rab inputs

525   collaborate in myosin-V-dependent transport of secretory compartments in yeast. Dev Cell 20:47–59.

526   64.   Bankaitis VA, Garcia-Mata R, Mousley CJ. 2012. Golgi membrane dynamics viewed through a lens of lipids. Curr Biol

527   CB 22:R414–R424.

528   65.   McFadden DC, Fries BC, Wang F, Casadevall A. 2007. Capsule structural heterogeneity and antigenic variation in

529   *Cryptococcus neoformans*. Eukaryot Cell 6:1464–1473.

530   66.   Doering TL. 2009. How sweet it is! Cell wall biogenesis and polysaccharide capsule formation in *Cryptococcus

531   neoformans*. Annu Rev Microbiol 63:223–247.

532   67.   Fu J, Hettler E, Wickes BL. 2006. Split marker transformation increases homologous integration frequency in *Crypto-

533   coccus neoformans*. Fungal Genet Biol 43:200–212.

534   68.   Huang MY, Joshi MB, Boucher MJ, Lee S, Loza LC, Gaylord EA, Doering TL, Madhani HD. 2022. Short homology-di-

535   rected repair using optimized Cas9 in the pathogen *Cryptococcus neoformans* enables rapid gene deletion and tag-

536   ging. Genetics 220:iyab180.

537   69.   Upadhya R, Lam WC, Maybruck B, Donlin MJ, Chang AL, Kayode S, Ormerod KL, Fraser JA, Doering TL, Lodge JK.

538   2017. A fluorogenic *C. neoformans* reporter strain with a robust expression of m-cherry expressed from a safe ha-

539   ven site in the genome. Fungal Genet Biol FG B 108:13–25.

70. Chang AL, Kang Y, Doering TL. 2019. Cdk8 and Ssn801 regulate oxidative stress resistance and virulence in *Crypto-*

coccus neoformans. mBio 10:e02818-18.

71. Cherniak R, Morris LC, Anderson BC, Meyer SA. 1991. Facilitated isolation, purification, and analysis of glucuronoxy-

lomannan of *Cryptococcus neoformans*. Infect Immun 59:59–64.

72. Willis LM, Stupak J, Richards MR, Lowary TL, Li J, Whitfield C. 2013. Conserved glycolipid termini in capsular poly-

saccharides synthesized by ATP-binding cassette transporter-dependent pathways in Gram-negative pathogens.

Proc Natl Acad Sci 110:7868–7873.

73. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA,

Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000.

Gene Ontology: tool for the unification of biology. 1. Nat Genet 25:25–29.

74. The Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, Ebert D, Feuermann M,

Gaudet P, Harris NL, Hill DP, Lee R, Mi H, Moxon S, Mungall CJ, Muruganugan A, Mushayahama T, Sternberg PW,

Thomas PD, Van Auken K, Ramsey J, Siegele DA, Chisholm RL, Fey P, Aspromonte MC, Nugnes MV, Quaglia F, Tosatto

S, Giglio M, Nadendla S, Antonazzo G, Attrill H, dos Santos G, Marygold S, Strelets V, Tabone CJ, Thurmond J, Zhou P,

Ahmed SH, Asanitthong P, Luna Buitrago D, Erdol MN, Gage MC, Ali Kadhum M, Li KYC, Long M, Michalak A, Pesala

A, Pritazahra A, Saverimuttu SCC, Su R, Thurlow KE, Lovering RC, Logie C, Oliferenko S, Blake J, Christie K, Corbani L,

Dolan ME, Drabkin HJ, Hill DP, Ni L, Sitnikov D, Smith C, Cuzick A, Seager J, Cooper L, Elser J, Jaiswal P, Gupta P,

Jaiswal P, Naithani S, Lera-Ramirez M, Rutherford K, Wood V, De Pons JL, Dwinell MR, Hayman GT, Kaldunski ML,

Kwitek AE, Laulederkind SJF, Tutaj MA, Vedi M, Wang S-J, D'Eustachio P, Aimo L, Axelsen K, Bridge A, Hyka-Nouspikel

N, Morgat A, Aleksander SA, Cherry JM, Engel SR, Karra K, Miyasato SR, Nash RS, Skrzypek MS, Weng S, Wong ED,

Bakker E, Berardini TZ, Reiser L, Auchincloss A, Axelsen K, Argoud-Puy G, Blatter M-C, Boutet E, Breuza L, Bridge A,

Casals-Casas C, Coudert E, Estreicher A, Livia Famiglietti M, Feuermann M, Gos A, Gruaz-Gumowski N, Hulo C, Hyka-

Nouspikel N, Jungo F, Le Mercier P, Lieberherr D, Masson P, Morgat A, Pedruzzi I, Pourcel L, Poux S, Rivoire C,

Sundaram S, Bateman A, Bowler-Barnett E, Bye-A-Jee H, Denny P, Ignatchenko A, Ishtiaq R, Lock A, Lussi Y, Magrane

M, Martin MJ, Orchard S, Raposo P, Speretta E, Tyagi N, Warner K, Zaru R, Diehl AD, Lee R, Chan J, Diamantakis S,

Raciti D, Zarowiecki M, Fisher M, James-Zorn C, Ponferrada V, Zorn A, Ramachandran S, Ruzicka L, Westerfield M.

2023. The Gene Ontology knowledgebase in 2023. Genetics 224:iyad031.

75.   Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence

alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res

22:4673–4680.

FIGURE LEGENDS

Figure 1. Screen results. (A) Schematic of screening workflow. Asterisk indicates *SAC1* in the diagram at right. (B and C)

Uptake scores (as defined in the text) for the 93 mutants identified as hits when screened against both (B) BMDM and (C)

HMDM. The value under each bar represents the center of the bin and 1.0 is the WT value. (D) Selected GO terms that are

enriched in the 93 dual hits from the uptake screen. X-axis, number of hit gene IDs which fall within each category. (E)

Capsule score (capsule thickness relative to that of WT under inducing conditions) for 131 mutants identified as hits in the

capsule screen, plotted as in (B) and (C). (F) Selected GO terms that are enriched in hits from the capsule screen, plotted

as in (D). For panels D and F, bold text denotes categories uniquely enriched for each screen; note that not all categories

are listed.

Figure 2. Cells lacking active Sac1 show increased uptake by phagocytes and smaller capsules. (A) Uptake and capsule

screen results for cells lacking CKF44_06080. (B) Alignment of *C. neoformans* and *S. cerevisiae* sequences that include

the Sac1 catalytic motif. Blue, identical residues; asterisk, residue mutated to alanine in the $SAC1^{D456A}$ strain. (C) India ink

negative stain of the indicated strains grown for 24 hours in DMEM at 37°C with 5% CO_2. Capsule thickness in μm (mean ±

SD) is noted below each panel. All primary images are at the same magnification; scale bar, 10 μm. For this and other

microscopy figures, any region demarcated by a small square is enlarged in the inset at upper right. (D) Distribution of

normalized capsule thicknesses for the indicated strains. At least 96 cells were measured for each strain as described in

the Methods. *, p<0.0332; ****, p<0.0001 by 2-way ANOVA with Tukey's multiple comparison test; ns, not significant.

Figure 3. Sac1 is important for virulence in a mouse infection model. (A) Lung and (B) burden of C57BL/6 mice (groups of

8-10) sacrificed 14 days after intranasal infection with 5×10^4 fungal cells (upper dotted line). Plots show median (bold black

line) with interquartile range. Lower dotted line, limit of detection. Each symbol represents one mouse. ****, p<0.0001;

comparison with no stars shown are not significant. Significance was calculated using 2-way ANOVA with Tukey's multiple

comparison test.

Figure 4. Sac1 regulates substrate distribution and stress sensitivity. (A) FAPP1-mNeonGreen (mNG) localization in cells

grown in YPD at 30°C. All primary images are at the same magnification; scale bar, 10 μm. (B) Serial 10-fold dilutions of the

indicated strains plated on YPD alone or containing 0.01% SDS (SDS), 0.05% Congo Red (CR), or 2 μg/mL Amphotericin B

(AMB), incubated at 30 and 37°C. *inact = SAC1^{D456A}.*

Figure 5. Sac1 is essential for proper lipid trafficking in host-like conditions. (A) Transmission electron micrographs of cells

grown in the indicated medium. White asterisks, example lipid droplet structures, likely distorted during sample prep; scale

bar, 1 μm. (B) Nile Red stain of the indicated strains with (C) quantification of the observed staining pattern. (D) Filipin

staining of cells grown in DMEM. YPD grown cells in Panel A were grown overnight at 30 °C, all other cells were grown for

24 h in DMEM at 37°C with 5% CO_2. Scale bar, 10 μm. All images within each experiment are shown at the same magnifi-

cation.

Figure 6. Supplementation with exogenous fatty acid partially restores *sac1Δ* growth and morphology but not capsule

biosynthesis. (A) Serial 10-fold dilutions of WT and *sac1Δ* cells grown for 24 hours (at 37°C with 5% CO_2) in DMEM supple-

mented with BSA alone (-), or with 8, 40, or 200 μM myristic acid conjugated to BSA (increasing amounts indicated by the

triangle). (B) India ink negative stain of WT and *sac1Δ* cells grown in DMEM at 37°C with 5% CO_2, supplemented as indicated

with BSA alone or 200 μM myristic acid conjugated to BSA. Capsule thickness in μm (mean ± SD) is noted below each panel.

All primary images are to the same magnification; scale bar, 10 μm.

Figure 7. Sac1 controls the secretory pathway to modulate capsule size and composition. (A) Secreted acid phosphatase

activity, normalized to cell number, over time after a shift to phosphate-depleted medium. (B) Confocal imaging of Cxt1-

mNG in the indicated strain backgrounds with (C) quantification of observed staining patterns. Punctate: 3 or more

puncta present; Intermediate: puncta and diffuse signal; Diffuse/low: diffuse signal or <3 puncta. Images are to the same

scale; scale bar, 10 μm. (D) Shed GXM quantified by ELISA. For both (A) and (D), mean ± SD for one representative experi-

ment of three biological replicate studies is shown. (E) Left, GXM composition; right, subunit structure of GXM.

 SUPPLEMENTAL MATERIAL FOR GAYLORD ET AL

Supplemental Methods

<u>BMDM preparation</u>

BMDMs were isolated as in (70) from 9-week old C57BL/6J mice and frozen in 90% FBS, 10% DMSO at 10^7 cells/mL.

Thawed cells were differentiated for 9 days in RPMI-1640 with L-glutamine (Sigma) with 30% L-cell supernatant and 20%

heat-inactivated fetal bovine serum (Gibco), with regular media changes. BMDM were resuspended in 10 mL ice cold PBS

and isolated using 2.5 µg anti-mouse F4/80 biotin (clone BM8, eBiosciences) plus 0.5 µg mouse BD FC-Block (BD

Pharmingen) with anti-biotin microbeads (Miltenyi Biotec) and a MACS separation column (Miltenyi Biotec).

<u>Human macrophage preparation</u>

Leukoreduction system (LRS) chambers were obtained from anonymous donors and human peripheral blood mon-

onuclear cells (PMBC) were isolated by Ficoll-Paque centrifugation (Lymphoprep; Axis-Shield). $5x10^7$ PMBC were allowed

to adhere to a treated culture dish (Corning) for 2 hours at 37°C with 5% CO_2. Detached cells were removed, and adherent

cells were maintained in RPMI-1640 differentiation media containing L-glutamine (Sigma), 10% fetal bovine serum (Sigma),

1% penicillin streptomycin (Fisher), 1% sodium pyruvate (Corning), 50ng/mL human recombinant M-CSF (Fisher), and 1%

MEM non-essential amino acids (Corning). Medium was replaced every two days and adherent cells were harvested on

day 6 using Cell Stripper (Corning).

<u>Uptake screen</u>

Assays were performed using a modification of our prior method (23). Host cells were seeded in flat clear bottom

black polystyrene TC-treated 96-well plates (Corning) at a density of 10^5 cells/well and incubated for one day at 37°C with

5% CO_2. *C. neoformans* strains were cultured from frozen stocks in glycerol/Yeast Extract-Peptone-Dextrose medium (YPD)

in 96 well format for 18-24 h at 30°C with shaking (800 rpm) (Southwest Science, SBT1500-H). At least 2 µL of cells were

then sub-cultured in 100 µL YPD and grown for 18-24 h as above. After sub-culture, cells were sedimented (3000 g, 2 min,

RT), washed twice with 200 µL of PBS, washed once with 200 µL McIlvaine's Buffer [pH 6], and resuspended in 80 µL of the

same. Cells were then stained with 5 mg/mL Lucifer Yellow (30 min, RT, 650 rpm) and resuspended in 100 µL of PBS. To

account for variation in growth rates of *C. neoformans* mutants, OD_{600} was used to roughly standardize cell number and

maintain results in the linear range of the assay. To do this, individual mutants were categorized as having high, medium,

or low growth based on the OD_{600} after resuspension in PBS and the average value of each group was used to prepare wells

with approximately 10^7 fungal cells/mL in PBS. Fungal cells were then opsonized with 10% by volume C57Bl6/J mouse

serum (for BMDM) or human serum from male AB plasma (for HMDM, Sigma) for 30 minutes at 37°C at 250 rpm. Opso-

nized cells were then added to wells containing RPMI and macrophages at a multiplicity of infection of 20 (BMDM) or 30

(HMDM). Cells were co-incubated for one hour at 37°C with 5% CO_2 to allow uptake and washed. This and the following

washes were performed using a Biotek ELx405 Microplate Washer (2 washes, PBS). Adherent fungal cells were stained

with 150 µL of 5 µg/mL Calcofluor White (Sigma) (15 min, 4°C, static). Samples were then washed, fixed with 4% formal-

dehyde (10 min, RT), washed, permeabilized with 1 mg/mL saponin (20 min, RT), washed, and stained with 50 µg/mL

propidium iodide (15 min, RT). Imaging was performed on a BioTek Cytation 3 imager collecting 12 images per well in a

4x3 grid pattern.

A separate plate of each *C. neoformans* strain alone was counted to allow normalization for the number of cells of

each mutant strain that were used in the assay. The phagocytic index for each mutant was determined by calculating the

number of internalized cells per macrophage and normalizing this to the value for WT KN99 and the number of fungi

added. An uptake score was then calculated for each well as fold-change of the normalized phagocytic index compared

either to the plate median (for the initial screen of Madhani library plates) or to WT (for the follow-up screens with BMDM

and HMDM). For the initial screen of the Madhani deletion collection, we calculated the first quartile (Q1), third quartile

(Q3), and interquartile range (IQR) for each plate. We defined hits as those below Q1-c*IQR or above Q3-c*IQR, where c

was calculated using a targeted error rate, α, of 0.01. For the second round of screening in BMDM and HMDM, we used

an internal WT control and defined hits as those strains whose uptake scores differed from WT in a statistically significant

way, using Dunnett's 1-way ANOVA and a p value cutoff of 0.05.

<u>Capsule screen</u>

Strains identified as hits in the uptake assay and wild type control cells were cultured in YPD from frozen stocks in

glycerol/YPD in 96 well format (ON, 30 °C, 800 rpm); 10 µL aliquots were then sub-cultured into 190 µL fresh YPD and

grown for an additional 24 h. Cells were then sedimented (3000 g, RT) and resuspended in 200 µL DMEM. To achieve the

low cell density necessary for optimum capsule induction, the average OD_{600} of the plate was used to calculate the volume

of cells to resuspend in 150 µL DMEM to achieve an average OD_{600} of 0.01. Capsule induction was performed in a 96-well

glass-bottom plate coated with poly-lysine (Eppendorf) and incubated for 24 h at 37°C with 5% CO_2. KN99α cells grown

overnight in YPD were added to empty wells 10 min before the induction ended to serve as an uninduced control. Follow-

ing induction, cells were stained and imaged as published previously (30). Briefly, the capsules were stained with anti-

capsular monoclonal antibody 302 conjugated to Alexa Fluor 488 (Molecular Probes) and the cell walls stained with Calco-

fluor White (MP Biomedicals). Imaging was performed on a BioTek Cytation 3 imager collecting 64 images per well in an

8x8 grid pattern. Images were prepared, annotated, and cell wall and capsule diameters for annotated cells calculated as

described previously (30). Capsule thickness was defined as the difference between cell wall and capsule diameters. A

negative control value (mean capsule thickness of WT cells grown in YPD in the same plate) was subtracted and the result

normalized to the value for similarly-corrected positive control samples (WT cells induced in the same plate) to yield a

capsule score. Hypocapsular and hypercapsular hits were categorized as those with capsule scores of less than 0.5 or more

than 2.0, respectively.

Gene ontology enrichment analysis

Gene ontology (GO) analysis was performed using FungiDB, using both manually curated and computationally

predicted GO terms (49, 73, 74). GO categories that were enriched with a P-value cutoff of 0.01 were manually selected

for each screen to minimize over-representation of redundant phenotypes.

Strain construction and validation

sac1Δ, *SAC1* complement, and *SAC1^{D456A}* strains were engineered using biolistic transformation with a split-marker

strategy (67) in a KN99α background. Design of the catalytically-inactivated construct was based on alignment (75) to the

S. cerevisiae S288C Sac1 amino acid sequence. mNeonGreen-tagged strains were generated using pBHM2404, a gift from

Hiten Madhani (Addgene 173441). PI4P-binding constructs were generated using pRS406-PHO5-GFP-hFAPP1 PH domain,

a gift from Tim Levine (Addgene 58723) (41). This sequence was placed between a *TEF1* promoter and terminator and

inserted into the Safe Haven 2 (SH2) region of the genome (69). All fluorescent constructs were generated using CRISPR

with short homology-directed repair (68). Transformants were confirmed by appropriate antifungal resistance, PCR, and

whole genome sequencing.

To test complementation of *sac1Δ* by the corresponding *S. cerevisiae* sequence, the *SAC1* gene was amplified from

S. cerevisiae strain S288C and inserted into plasmid pMSC042-neo between an *ACT1* promoter and *TRP1* terminator. This

construct (containing the promoter, *SAC1* sequence, terminator and antifungal resistance marker) was inserted into the

SH2 region of WT and *sac1Δ* cells using CRISPR (68, 69). Total RNA was extracted from cells grown in YPD and DMEM using

TRIzol Reagent (Thermo Fisher) and cDNA was synthesized using the SuperScript III First-Strand Synthesis SuperMix kit

(Thermo Fisher). PCR amplification of an S288C Sac1-specific sequence from the cDNA confirmed expression of the S288C

SAC1 gene in both backgrounds (Fig. S1B).

Fungal cell growth and capsule induction

C. neoformans strains were grown from glycerol stocks on YPD plates for 2 days at 30°C. Overnight cultures were

inoculated from single colonies into 4 mL of YPD medium and grown overnight at 30°C with shaking (unless otherwise

noted). For all capsule induction experiments, overnight cultures were washed in sterile PBS, inoculated into DMEM at

10^6 cells/mL, and grown statically at 37°C with 5% CO_2 for 16-24 hours (as indicated in the text).

Virulence studies

All animal protocols were approved by the Washington University Institutional Animal Care and Use Committee

(Protocol #20-0108) and care was taken to minimize animal handling and discomfort. 6-8-week old female C57BL/6 mice

(The Jackson Laboratory) were anesthetized by subcutaneous injection of 1.20 mg ketamine and 0.24 mg xylazine in 110

μL sterile PBS. For 14-day infections, mice were intranasally infected with 5×10^4 cryptococcal cells. For time course infec-

tion, mice were infected with 1.25×10^4 cryptococcal cells and sacrificed at 6-, 12-, and 18-days post infection. For both

infections, the lungs and brains were harvested, homogenized, and plated on YPD agar for calculation of fungal organ

burden.

Stress plate dot spotting

Overnight cultures were pelleted and washed twice in PBS, then diluted to 10^7 cells/mL. 4 μL of this stock and four

serial 1:10 dilutions were spotted onto YPD agar and incubated at both 30 and 37°C for 48 hours. To impose plasma

membrane stress, the YPD was supplemented with 0.01% sodium dodecyl sulfate or 1 μg/ml amphotericin B; to impose

cell wall stress, the YPD was supplemented with 0.05% Congo red.

Acid phosphatase secretion assay

Single colonies were inoculated into 5 mL YPD, grown shaking at 30°C for 6 h, sub-cultured into 25 mL MM-KH$_2$PO$_4$

(0.5% KH_2PO_4, 15 mM glucose, 10 mM $MgSO_4 \cdot 7H_2O$, 13 mM glycine, 3.0 μM thiamine), and grown overnight at 30°C. Cells

were then pelleted, washed twice with MM-KCl (0.5% KCl, 15 mM glucose, 10 mM $MgSO_4 \cdot 7H_2O$, 13 mM glycine, 3.0 μM

thiamine), and resuspended in 50 mL MM-KCl at 10^7 cells/mL. Cultures were grown at 30°C with shaking, and 2 mL aliquots

were collected at each timepoint. For each, pelleted cells were resuspended in 400 μL substrate buffer (2.5 mM para-

nitrophenylphosphate in 50 mM acetate buffer, pH 4.0, Sigma 71768-5G), incubated at 37°C for 5 minutes, and the reac-

tion stopped with 800 μL saturated Na_2CO_3. Acid phosphatase secretion was quantified using absorbance at 420 nm, cor-

rected by subtraction of a cell-free blank value, and normalized to cell density using OD_{600}.

<u>GXM ELISA</u>

Cultures were induced in DMEM for the indicated times, at which point cells were removed by centrifugation

(3000 x g, 5 min, RT), the supernatant fraction filtered through a 0.2μm filter, and the filtrate diluted 1:500 in PBS. Flat-

bottom Immulon 1B microtiter plates (Thermo Scientific) were coated overnight with a 1 μg/ml mix of anti-GXM antibodies

339 and F12D2γ in PBS. This and the following incubations were performed at room temperature. The plates were washed

twice with 200 μL PBS with 0.5% TWEEN 20 (PBST), blocked for 90 min with 200 μL 1% BSA in PBST. The blocking buffer

was replaced with 100 μL PBST and 100 μL of filtered sample was added and incubated for 90 min. The plate was then

washed three times as above and the bound GXM was probed with HRP-labeled mAb 339/F12D2γ, incubated for 1 minute

with KPL TMB Microwell Peroxidase Substrate (SeraCare), developed, and quantified using absorbance at 450 nm.

<u>Imaging</u>

For fluorescence and confocal microscopy, cells were grown as indicated above and washed twice in PBS. For

capsule visualization, cells were resuspended in India ink in PBS (1:2, v:v) and imaged on a ZEISS Axio Imager M2 fluores-

cence microscope. Capsule thickness was measured manually in ImageJ, using a total of at least 96 cells from at least six

randomly-chosen fields of view. To stain neutral lipids, a 0.01% Nile Red stock in acetone was diluted to a working con-

centration of 0.0005% in PBS. Cells were incubated in 1 mL working Nile Red solution for 15 min in the dark, rotating. Cells

were washed twice in PBS, resuspended in PBS, and imaged as above. For quantification of morphology, a minimum of 60

cells from randomly-chosen slide coordinates were examined. For filipin staining, cells were incubated in a freshly pre-

pared solution of 5 μg/mL filipin (Millipore Sigma, F9765-25MG) in DMSO for 15 minutes at room temperature, rotating,

then washed twice with PBS before imaging as above. To visualize FAPP1-mNeonGreen and Cxt1- mNeonGreen, cells were

imaged using a Zeiss LSM880 confocal microscope. For quantification of morphology, a minimum of 94 cells from ran-

domly-chosen slide coordinates were examined. For electron microscopy, cells were grown overnight in YPD or induced

for 24 hours in DMEM before processing as in (47), except that the first incubation in 0.1 M sodium cacodylate buffer was

performed for 2 h at RT.

<u>Lipid supplementation</u>

For lipid supplementation, a fatty acid stock solution (10 mM in ethanol) was diluted 1:10 into 4 mg/ml BSA in

DMEM and then sonicated (Bransonic 2 Ultrasonic Bath Sonicator) before addition to cultures as indicated. After 24-hour

of incubation (37°C, 5% CO_2), cells were serially diluted 1:10 and plated onto YPD agar, with the top spot representing the

undiluted culture.

<u>GXM purification and analysis</u>

GXM was isolated using selective precipitation of culture supernatants with hexadecyltrimethylammonium bro-

mide (CTAB) as described previously (48). Isolated material was hydrolyzed in trifluoroacetic acid for 3 hours at 100°C and

compositional analysis performed using high-performance anion-exchange chromatography with pulsed amperometric

detection on a Dionex ICS-6000 instrument (Thermo Scientific). Glycosyl linkage analysis was performed by the Complex

Carbohydrate Research Center at the University of Georgia using combined gas chromatography/mass spectrometry of

partially methylated alditol acetate derivatives following carboxyl reduction to detect uronic acids (modified from (72)).

Supplemental Table Legends

Supplemental Table S1. Uptake screen results. (S1A) All results from the initial BMDM uptake screen. (S1B) Hits from the

initial BMDM uptake screen. (S1C) All results from the second BMDM uptake screen. (S1D) Hits identified from the second

BMDM uptake screen. (S1E) All results from the HMDM uptake screen. (S1F) Hits from the HMDM uptake screen.

Supplemental Table S2. Capsule screen results. (S2A) All results from the capsule screen. (S2B) Hits identified from the

capsule screen. (S2C) Overlap hits identified in all three screens: HMDM, BMDM, and capsule.

Supplemental Table S3. Linkage analysis of WT, *sac1Δ*, and *SAC1* strains. Values are the percent area of relevant residue

peaks. Numbers in the header row refer to independently isolated samples.

Supplemental Figure Legends

Supplemental Figure S1. Overexpression of the *S. cerevisiae* Sac1 protein in *sac1Δ* does not restore *C. neoformans* capsule

synthesis. (A) Negative stain of the indicated strains grown for 24 hours in DMEM at 37°C with 5% CO_2, with the capsule

thickness in µm (mean ± SD). Scale bar, 10 µm. (B) Amplification of a unique 651 bp region of the S288C Sac1 coding

sequence from cDNA generated from fungal cells of the indicated strain grown in YPD shows that the *S. cerevisiae SAC1*

gene is expressed in both *C. neoformans* strains.

Supplemental Figure S2. Construction and validation of the $SAC1D^{456A}$ strain. (A) Strain construction strategy for *SAC1*

complement and SAC^{D456A} inactivated complement strains. (B) Normalized expression of *SAC1* in WT and $SAC1^{D456A}$ cells

grown for 24 hours in DMEM at 37°C with 5% CO_2. (C) Nucleotide sequence of $SAC1^{D456}$. Red, nucleotide differences from

WT *SAC1*.

Supplemental Figure S3. Cells lacking functional Sac1 have reduced virulence. (A) Lung and (B) brain burden of C57BL/6

mice sacrificed at the indicated days post infection (dpi) with 1.25×10^4 fungal cells of *ckf44_06080Δ* from the Madhani *C.*

neoformans deletion collection. Upper dotted line, inoculum. Lower dotted line, limit of detection. Each symbol represents

one mouse.

Supplemental Figure S4. PI4P content of WT and *sac1Δ* strains. Difference in median fluorescence intensity between

FAPP1-mNG expressing and control cells (ΔMFI) in WT and mutant backgrounds, grown in the indicated medium. ns, not

significant by unpaired t test.

Supplemental Figure S5. Dual staining with filipin and Nile Red. All cells were grown in DMEM for 24 hours at 37°C with

5% CO_2. DIC, Differential interference contrast. Scale bar, 10 μm.

Supplemental Figure S6. Supplementation with exogenous fatty acids partially restores growth of *sac1Δ* cells. Serial dilu-

tions of (left) WT or (right) *sac1Δ* cells grown for 24 hours in DMEM alone, DMEM supplemented with BSA, or DMEM with

100 μM of the indicated fatty acid sodium salt conjugated to 0.4 mg/mL BSA.

Figure 1. Screen results. (A) Left, schematic of screening workflow, and right, overlap of screen hits. Asterisk indicates *SAC1* in the diagram at right. **(B and C)** Uptake scores (as defined in the text) for the 93 mutants identified as hits when screened against both **(B)** BMDM and **(C)** HMDM. The value under each bar represents the center of the bin and 1.0 is the WT value. **(D)** Selected GO terms that are enriched in the 93 dual hits from the uptake screen. X-axis, number of hit gene IDs which fall within each category. **(E)** Capsule score (capsule thickness relative to that of WT under inducing conditions) for 131 mutants identified as hits in the capsule screen, plotted as in **(B)** and **(C)**. **(F)** Selected GO terms that are enriched in hits from the capsule screen, plotted as in **(D)**. For panels D and F, bold text denotes categories uniquely enriched for each screen; note that not all categories are listed.

A

Deleted gene	Annotation	Uptake score		Normalized capsule size
		BMDM	HMDM	
CKF44_06080	inositol/phosphatidylinositol phosphatase	2.25	1.83	0.39

B

Catalytic motif

*

Sc Sac1 V E I V N E Q H S V V R T N C M D C L D R T N V V Q S **(378-404)**
Cn Sac1 T Q V L T K Q T G V V R T N C M D C L D R T N V A Q A **(440-466)**

C

WT — 3.50 ± 0.73 µm
sac1Δ — 1.94 ± 0.57 µm
SAC1 — 3.59 ± 0.74 µm
SAC1^{D456A} — 2.22 ± 0.61 µm

D

Capsule thickness relative to cell diameter

Figure 2. Cells lacking active Sac1 show increased uptake by phagocytes and smaller capsules. (A) Uptake and capsule screen results for cells lacking CKF44_06080. **(B)** Alignment of *C. neoformans* and *S. cerevisiae* sequences that include the Sac1 catalytic motif. Blue, identical residues; asterisk, residue mutated to alanine in the *SAC1^{D456A}* strain. **(C)** India ink negative stain of the indicated strains grown for 24 hours in DMEM at 37°C with 5% CO_2. Capsule thickness in µm (mean ± SD) is noted below each panel. All primary images are at the same magnification; scale bar, 10 µm. For this and other microscopy figures, any region demarcated by a small square is enlarged in the inset at upper right. **(D)** Distribution of normalized capsule thicknesses for the indicated strains. At least 96 cells were measured for each strain as described in the Methods. *, $p<0.0332$; ****, $p<0.0001$ by 2-way ANOVA with Tukey's multiple comparison test; ns, not significant.

Figure 3. **Sac1 is important for virulence in a mouse infection model. (A)** Lung and **(B)** burden of C57BL/6 mice (groups of 8-10) sacrificed 14 days after intranasal infection with 5×10^4 fungal cells (upper dotted line). Plots show median (bold black line) with interquartile range. Lower dotted line, limit of detection. Each symbol represents one mouse. ****, $p < 0.0001$; comparison with no stars shown are not significant. Significance was calculated using 2-way ANOVA with Tukey's multiple comparison test.

Figure 4. Sac1 regulates substrate distribution and stress sensitivity. (A) FAPP1-mNeonGreen (mNG) localization in cells grown in YPD at 30°C. All primary images are at the same magnification; scale bar, 10 μm. **(B)** Serial 10-fold dilutions of the indicated strains plated on YPD alone or containing 0.01% SDS (SDS), 0.05% Congo Red (CR), or 2 μg/mL Amphotericin B (AMB), incubated at 30 and 37°C. *inact = SAC1^{D456A}*.

Figure 5. Sac1 is essential for proper lipid trafficking in host-like conditions. (A) Transmission electron micrographs of cells grown in the indicated medium. White asterisks, example lipid droplet structures, likely distorted during sample prep; scale bar, 1 μm. **(B)** Nile Red stain of the indicated strains with **(C)** quantification of the observed staining pattern. **(D)** Filipin staining of cells grown in DMEM. YPD grown cells in Panel A were grown overnight at 30 °C, all other cells were grown for 24 h in DMEM at 37°C with 5% CO_2. Scale bar, 10 μm. All images within each experiment are shown at the same magnification.

A

WT   sac1Δ

sodium myristate   sodium myristate

B

	DMEM	DMEM + BSA vehicle	DMEM + sodium myristate

WT — 3.13 ± 0.70 μm   3.97 ± 0.81 μm   2.20 ± 0.63 μm

sac1Δ — 2.03 ± 0.52 μm   2.41 ± 0.71 μm   0.67 ± 0.47 μm

Figure 6. Supplementation with exogenous fatty acid partially restores *sac1*Δ growth and morphology but not capsule biosynthesis. (A) Serial 10-fold dilutions of WT and *sac1*Δ cells grown for 24 hours (at 37°C with 5% CO_2) in DMEM supplemented with BSA alone (-), or with 8, 40, or 200 μM myristic acid conjugated to BSA (increasing amounts indicated by the triangle). **(B)** India ink negative stain of WT and *sac1*Δ cells grown in DMEM at 37°C with 5% CO_2, supplemented as indicated with BSA alone or 200 μM myristic acid conjugated to BSA. Capsule thickness in μm (mean ± SD) is noted below each panel. All primary images are to the same magnification; scale bar, 10 μm.

Figure 7. Sac1 controls the secretory pathway to modulate capsule size and composition. (A) Secreted acid phosphatase activity, normalized to cell number, over time after a shift to phosphate-depleted medium. **(B)** Confocal imaging of Cxt1-mNG in the indicated strain backgrounds with **(C)** quantification of observed staining patterns. Punctate: 3 or more puncta present; Intermediate: puncta and diffuse signal; Diffuse/low: diffuse signal or <3 puncta. Images are to the same scale; scale bar, 10 μm. **(D)** Shed GXM quantified by ELISA. For both **(A)** and **(D)**, mean ± SD for one representative experiment of three biological replicate studies is shown. **(E)** Left, GXM composition; right, subunit structure of GXM.

A

B

1 – ladder
2 – ScSac1 *sac1*Δ
3 – ScSac1 KN99

Supplemental Figure S1. Overexpression of the *S. cerevisiae* Sac1 protein in *sac1*Δ does not restore *C. neoformans* capsule synthesis. (A) Negative stain of the indicated strains grown for 24 hours in DMEM at 37°C with 5% CO_2, with the capsule thickness in μm (mean ± SD). Scale bar, 10 μm. **(B)** Amplification of a unique 651 bp region of the S288C Sac1 coding sequence from cDNA generated from fungal cells of the indicated strain grown in YPD shows that the *S. cerevisiae SAC1* gene is expressed in both *C. neoformans* strains.

B

SAC1

ns

WT $SAC1^{D456A}$

Normalized $2^{-\Delta\Delta Ct}$ relative to WT

C

```
ATGGCGTTTAATGCTCTACCTGTATGTGACCTTCGCGAAGTGAACTGGCTGCCGCTAACAAGACTTAGCTTCACGAAACC
TTAAACCTCTACGTCTCTCCTACGGCTTACATTTTCGAGCCTGCTTCCTCCAGTGCTGGCCATGTGGGAATCGATGGAAC
TATATTCGTTGATGAAAGAATGTAAGGGAGAGTATGGTTGTGGACCGTCAGACTGGCCAGATTAGCTTGAGTAGTGAGT
GACAACCCTTGCAATGCGTGGAGCAGAATATCACTGATGCGACAGAACATAACAGCGGCTTCTCATATACCATTTGGAAA
GGAAAAGGTTATTACTTGCTACGGTATTATTGGCATTCTCAGCCTTGCTACTAGTAAGTTTTGTTGCTCCTCAGACACAA
TTGCCCCCTGCTGATTTTGTGCTTTCTAATAGCCGATTTTCTTCTTATTGTCACTTCCCGCTCTCCTTCATGTCGCCTTC
TTTCTCACCCTATCTATCTTGCGAATGATTATCGACTTCTTCCTATTTCCCCTCTGTCTACTTCCTCCGCCATTCTTGAA
CATCCTGTGGAGAAGGAACTCATCTCTCTTGTTGAGCAGGGTCTCAAATCCAGTAAACTCTGGTTCTCGTATGGTTGGGA
CCTTACCAATTCTTTGCAGAGGCAACAAGAAATCGATTTGAAGCTCTCACAGTCCGGAGAACAATGGCCCGCCTGGAGAC
GGGCAGATGAACGTTTCTTTTGGAACAGGTTTCTCATGGACAAGATGATTGACGTGACCGAGAGCGGTGAGGCAGATCTG
AGCCGATTTATCCTTCCTATAATGTACGGCTCTATTGAGCTTCGGTCATCGACTCTTAACTCTCGGGATCTCCTTTTCCT
ACTCATCTCTCGGCGATCTCGTTACCGAGCTGGTACCAGATACTTCACGCGAGGCATCAACCCCTCTGGCCACGTCGCCA
ATTTCAACGAGACTGAGCAAATTGTTATGTATGACCCTATCCCGGAGAACGGAGAAGCCTACGGAAGGGGAAGGGTAGAC
GGTAGGGAAAGGCTGAGCTTTGTGCAGACAAGAGGGAGTGTTCCGCTATTCTGGGCCGAGGTCAACAACTTGAGGTACAA
GCCCGATTTGCAAATTATGGATTACACTGAGACGGTGAGTGAATTGTATTGAAAGGCATGACATGTCCTGACAATGAATT
AGCCTCAAGCTCTCAAAGCACATCTCAACAGTATGATTAAGACCTATGGCCATACATATCTTGTGAATCTTGTAAATCAA
AAGGGACACGAACAACCTGTCAAGGAAGCTTTTGAGCGCTACATGTCCCTTGTCGCCTCATCCGACCCGTCCATTCAGGA
GAAGGCTCACTACCTCTACTTTGACTTTCACCACGAATGCAAAGGTCTAAGGTTTGACCGAATTTCCCTTTTGGTCGAGA
AGCTTGCGACAGCACTTGAGGATATGAGCTGGTACCATTCTGTCAACCCCGACTCGTCAACCTATGCTCTCTTGCAACCA
AACTCTGCAGACACCCAGGTACTTACTAAGCAAACGGGCGTCGTCAGAACCAACTGTATGGCGCGTGTCTAGACAGGACCAA
TGTCGCACAAGCGGCTCTGGCTAGATGGGTTTTGAACCGGCAATTAAGGAAGGTCGGGATCCTGAGCGTGAAGGAAGGTG
TGGAGGATCACGAAGAGTTTATGATTATGTTCCGGAATGGTACGTGCTTTTCATTTTGACTAGAAATTAGACTAACTTGA
GAAAATAGTGTGGGCCGATCATGGTGACACCGTCTCCCGAGCTTATGCAGGAACTGGTGCTCTCAAATCCGACTTTACTC
GTACCGGTAAACGATCCAAGGAGGGTCTCTTGGAAGATGGTTACAAAAGCGTGATGAGATACTTTAGGAATAACTTCTTC
GATGGCGATCGTCAGGTTAGTCCCTACTGTTGTGAGCTGAGAGAGACATCTGACCGGAGAACAGGATGGTTTTGATATTC
TAACTGGAGCATGGGTAGCCCAAAGGGGTGGCATTCCTCCATTGACTGATACAAGACCTCTTATCATGCGCTCGGTAAGT
CCATGGTTTGTCCGACTAATGTTGACCCGAGAAGATGCCGTACATACTGGCGTTTGCTCTCACAATGATCACCGCAGCTC
TCACATTGCCACGAACCTCGGAGATGTCCATATATTCCTTCCTCGTCCTTTGGTTCTTCCTTGCATTCTTCTCAGGAAGT
TATATTTGGGGCAATGTAAGTTATTTTCTTCTTTCATATATTGACTTTGGCCGTTGACCGCCATTTGTCACTCTTAGGGT
ACATCATACGTCTCCTGGCCGCGTTTGAACCCTCCTCTAGAAATTCTCTCTTACTCTGGACCGGGTCACCGCTCTCCGGT
CCGCGGTCGAGGGATGTCGTTCGCAGCCATTGTGCCCCTGTTCGATCGTCGACGCCAAAATGGTCCTGGAGCTGGCGTAC
TTGGTGCCACCGGCGGCTTCGGAGGAAGGAACGATGGAAGACCTGCAATGTATAAGATGGAGGAGTCGGAGTTGGGGAGG
AGGAAGGGTGCGTTGATTGATTAA
```

Supplemental Figure S2. Construction and validation of the $SAC1D^{456A}$ strain. **(A)** Strain construction strategy for *SAC1* complement and SAC^{D456A} inactivated complement strains. **(B)** Normalized expression of *SAC1* in WT and $SAC1^{D456A}$ cells grown for 24 hours in DMEM at 37°C with 5% CO_2. **(C)** Nucleotide sequence of $SAC1^{D456A}$. Red, nucleotide differences from WT *SAC1*.

Supplemental Figure S3. Cells lacking functional Sac1 have reduced virulence. (A) Lung and **(B)** brain burden of C57BL/6 mice sacrificed at the indicated days post infection (dpi) with 1.25x10⁴ fungal cells of *ckf44_06080Δ* from the Madhani *C. neoformans* deletion collection. Upper dotted line, inoculum. Lower dotted line, limit of detection. Each symbol represents one mouse.

Supplemental Figure S4. PI4P content of WT and *sac1Δ* strains. Difference in median fluorescence intensity between FAPP1-mNG expressing and control cells (ΔMFI) in WT and mutant backgrounds, grown in the indicated medium. ns, not significant by unpaired t test.

	DIC	filipin	Nile Red	filipin + Nile Red
WT				
sac1Δ				
SAC1				
SAC1^{D456A}				

Supplemental Figure S5. Dual staining with filipin and Nile Red. All cells were grown in DMEM for 24 hours at 37°C with 5% CO_2. DIC, Differential interference contrast. Scale bar, 10 μm.

Supplemental Figure S6. Supplementation with exogenous fatty acids partially restores growth of *sac1*Δ cells. Serial dilutions of (left) WT or (right) *sac1*Δ cells grown for 24 hours in DMEM alone, DMEM supplemented with BSA, or DMEM with 100 µM of the indicated fatty acid sodium salt conjugated to 0.4 mg/mL BSA.

Residue	WT #1	*sac1Δ* #1	WT #2	*sac1Δ* #2	*SAC1* #2
Terminal Xylopyranosyl residue (t-Xyl)	12.0	19.7	10.4	16.2	11.1
Terminal Mannopyranosyl residue (t-Man)	1.3	0.3	0.3	0.3	0.4
Terminal Glucopyranosyl uronic acid residue (t-GlcA)	17.9	15.4	19.1	16.2	18.2
3-linked Mannopyranosyl residue (3-Man)	29.9	23.3	27.8	21.1	30.6
2,3-linked Mannopyranosyl residue (2,3-Man)	36.0	38.5	37.5	42.8	34.9
2,3,4-linked Mannopyranosyl residue (2,3,4-Man)	2.9	2.8	4.9	3.5	4.8

Supplemental Table S3. Linkage analysis of independently isolated WT, sac1Δ, and *SAC1* strains. Values are percent area of relevant residue peaks. Numbers in header row refer to independently isolated samples.