Orthogonalized human protease control of secreted signals

Carlos A. Aldrete[†,1], Connor C. Call[†,1], Lucas E. Sant'Anna[2],
Alexander E. Vlahos[1], Jimin Pei[3], Qian Cong[3,4,5], and Xiaojing J.
Gao[1]

† These authors contributed equally to this work.
1 Department of Chemical Engineering, Stanford University, Stanford CA 94305, USA
2 Department of Bioengineering, Stanford University, Stanford CA 94305, USA
3 Eugene McDermott Center for Human Growth and Development, University of Texas
Southwestern Medical Center, Dallas, TX 75390, USA
4 Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX
75390, USA
5 Harold C. Simmons Comprehensive Cancer Center, University of Texas Southwestern Medical
Center, Dallas, TX 75390, USA

Abstract

Synthetic circuits that regulate protein secretion in human cells could support cell-based
therapies by enabling control over local environments. While protein-level circuits enable such
potential clinical applications, featuring orthogonality and compactness, their non-human origin
poses a potential immunogenic risk. Here, we developed Humanized Drug Induced Regulation of
Engineered CyTokines (hDIRECT) as a platform to control cytokine activity exclusively using
human-derived proteins. We sourced a specific human protease and its FDA-approved inhibitor.
We engineered cytokines (IL-2, IL-6, and IL-10) whose activities can be activated and abrogated
by proteolytic cleavage. We utilized species specificity and re-localization strategies to
orthogonalize the cytokines and protease from the human context that they would be deployed
in. hDIRECT should enable local cytokine activation to support a variety of cell-based therapies
such as muscle regeneration and cancer immunotherapy. Our work offers a proof of concept for
the emerging appreciation of humanization in synthetic biology for human health.

Intercellular communication, mediated by the secretion and display of soluble factors
(cytokines and growth factors) and receptors, plays a crucial role in health and diseases. Cell
therapies aim to treat complex diseases using human cells through the modulation of intercellular
communication[1], with the chimeric antigen receptor (CAR) T cell therapy being the most notable
example[2]. Aside from treating cancer, other promising examples are the treatment of
autoimmune diseases and supporting muscle regeneration using mesenchymal stem cells
(MSCs)[3–6]. Despite their promise, many cellular therapeutics are susceptible to changes in the
local microenvironment[7–9]. To address this hurdle, researchers have "armed" cell therapies to
constitutively secrete cytokines to tune their local environments to promote proliferation and
efficacy, such as driving CAR-T activation in solid tumors using IL-12[10], and improving the
survival of transplanted MSCs using the anti-inflammatory IL-10[11]. While constant expression of
these powerful signals improve efficacy, increased toxicity was also observed, indicating a need
for further control[10].
The field of synthetic biology offers a plethora of tools and methods for localized,
conditional control of cytokine activity. In particular, compared to the more conventional
transcriptional control, protein-level controls hold advantages for potential clinical use such as
fast operation, compact single-transcript delivery, and context-independent performance[12].
Notably, orthogonal viral proteases have emerged as useful post-translational tools due to their
ability to control the activity, degradation, and localization of target proteins with high substrate
specificity[13,14]. These orthogonal proteases are highly versatile and can be combined to
implement robust sense-and-response behaviors in mammalian cells. They are uniquely
positioned to serve as cargos on mRNA vectors that are safer and more scalable than
conventional DNA vectors[15]. Moreover, they have been successfully applied to the control of
protein secretion and display[16,17].
However, despite the promise of proteases as control knobs, there remains a critical
constraint: the non-human origin and potential immunogenicity of viral proteases may preclude
their translational potential. In analogous scenarios, immune responses against other synthetic
biology tools such as *Streptococcus pyogenes* Cas9 protein (SpCas9) and synthetic CAR-T
receptors have been documented with compromised efficacy and duration, highlighting the need
to reduce immunogenicity of synthetic tools[18,19]. Even more notably, we are inspired by the
decades of efforts to "humanize" monoclonal antibodies[20,21], and we reason that, if synthetic
biology is to deliver its promise in human health, it is imperative that we preempt similar
challenges as we engineer the fundamental tools. In an emerging consensus to "humanize"
synthetic biology, groups have developed tools derived largely from human-derived components
such as zinc finger transcriptional regulators (synZiFTRs)[22], CRISPR-CAS-inspired RNA
targeting systems (CIRTS)[23], and synthetic intramembrane proteolysis receptors (SNIPRs)[24].
While these papers have demonstrated humanized transcriptional and translational regulation, the
humanization efforts have yet to reach the post-translational level.
Here, we sought to develop protease-based control of intercellular communication for
potential application in cellular therapeutics. We developed our system with five features for
potential clinical suitability, combining general humanization considerations[22] with the unique
benefits of protease control: (i) use of human-derived proteins to minimize the risk of
immunogenicity; (ii) use of orthogonalized components with high specificity and minimal
crosstalk with native signaling pathways; (iii) ability to control exogenously using FDA-
approved small molecules; (iv) compact, single-transcript design to facilitate delivery into

therapeutic cells; (v) compatibility with mRNA delivery for potential transient *in vivo* delivery
applications.
We identified a human protease involved in blood pressure regulation, renin, as the
control knob. Renin, in contrast to most human proteases, exhibits high substrate specificity[25].
Additionally, renin comes with a clinically-viable mode of exogenous control in the form of its
FDA-approved small-molecule inhibitor, aliskiren[26]. In contrast to previously used cytosolic
viral proteases, renin functions extracellularly and in the secretory pathway, offering an external
interface to directly control intercellular communication.
In this paper, we introduce Humanized Drug Induced Regulation of Engineered
CyTokines (hDIRECT), using renin to control intercellular communication. We engineered a
variety of clinically relevant cytokines (IL-2, IL-6, and IL-10), whose activities are increased or
decreased by renin-mediated proteolysis and can be tuned exogenously via aliskiren. We used
structure-guided mutagenesis and re-localization strategies to create renin mutants with new and
orthogonal substrate specificity from wildtype proteins. Finally, we demonstrate chemically
dependent and bidirectional control of simultaneous cytokine activity, compact delivery using
various methods and in multiple cell types, as well as small-molecule control *in vivo*. We
envision that conditional control of cytokine activity has immediate use for supporting and
controlling cell therapies as well as dissecting biological functions of dynamic cytokine profiles
– because most of the specific protease-regulated cytokine designs are reported for the first time
in this study. This study demonstrates a proof-of-principle for the orthogonalization of human
proteases and serves as an example of making biomolecular controls more clinically compatible.
97
Results
Engineering caged cytokines.
hDIRECT consists of three components: a membrane-bound renin, renin's FDA-
approved small-molecule inhibitor, aliskiren, and engineered cytokines modulated by renin
cleavage (Fig. 1a, b). Renin is known to function extracellularly as it is natively secreted by
juxtaglomerular cells in the kidney[25]. To localize renin to the plasma membrane to enable the
autonomous control of engineered cells, the active form of renin was fused to the transmembrane
domain of human CD28 (Fig. 1b). Here, constitutive expression of membrane-bound renin can
control the activity of co-expressed secreted cytokines. We first evaluated a variety of strategies
to place cytokines under the control of renin.
To engineer cytokines activated by renin cleavage, we drew inspiration from previous
work on attenuated cytokines, specifically, receptor-masked interleukin-2 (IL-2).[27,28] Here, IL-2
activity was tied to proteolytic cleavage by fusing IL-2 with a receptor subunit via a matrix
metalloproteinase (MMP)-cleavable linker. Cytokine activity is inhibited ("caged") in the
absence of protease, and conditionally activated ("uncaged") by proteolytic cleavage. Using this
general framework, we demonstrated the possibility to cage structurally and functionally distinct
cytokines. Inspired by MSC and CAR T applications, we chose interleukin-10 (IL-10, an
immuno-suppressive cytokine that plays a role in limiting host immune responses as well as
promoting wound healing and muscle regeneration[29]), interleukin-6 (IL-6, generally associated
with inflammation, but also implicated with the activation of satellite cells and promotion of
myotube regeneration for muscle regeneration[30–32]), and IL-2 (a potent factor for the proliferation
and activation of lymphocytes, and widely studied for activating effector T cells against
tumors[33–35]).

We started by designing a caged IL-10. IL-10 is composed of a homodimer that binds to
a tetrameric IL-10 receptor complex formed by two IL-10Ra and two IL-10Rb subunits.[36] Since
effectively caging a dimeric protein may be difficult due to having multiple epitopes to interact
with receptors, we also tested a previously engineered monomeric variant of IL-10 (R5A11M)
with boosted affinity for IL-10Rb and potent immunomodulatory effects.[37] We created two
caged IL-10 constructs. Both designs contain a renin cleavable linker (cut site and glycine-serine,
GS, linker) and the extracellular portion of specific IL-10R subunits for caging. The first
construct was created by fusing two WT IL-10 monomers together with IL-10Ra (CageIL-10-
dimer – Fig. 1a). The second construct used the monomeric R5A11M IL-10 and IL-10Rb
(CageIL-10-mono – Fig. 1a). We hypothesized that aliskiren could be used to control renin
activity and removal of the caged portion of hDIRECT to control cytokine activity (Fig. 1b).
Taking advantage of our fast design-build-test cycle, we transiently transfected human
embryonic kidney (HEK) 293 cells using plasmids encoding the caged cytokines and renin.
Supernatant was transferred to HEK-Blue™ IL-10 reporter cells to assess functional activity.
Here, signal transduction from IL-10 receptor activation results in expression of secreted
embryonic alkaline phosphatase (SEAP), which can be assayed via absorbance following
incubation with a colorimetric substrate. A schematic for this cytokine reporter assay can be
found in Supplementary Fig. 1 and links to plasmid maps as well as amounts of plasmid for each
transient transfection can be found in Supplementary Tables 1 and 2. Unless specified otherwise,
the same workflow for the cytokine reporter assay was used throughout this work. In the absence
of renin, cageIL-10-mono was caged more efficiently than the cageIL-10-dimer (Fig. 1c).
CageIL-10-mono was also able to fully reconstitute activity with protease present (Fig. 1c). From
here on, we refer to this optimal variant as "cageIL-10". Next, to explore the extent of IL-10
caging and linker cleavability, we titrated the amount of transfected cageIL-10 plasmid DNA.
We observed efficient caging even at high doses of cageIL-10 (Fig. 1d), and renin-activated IL-
10 activity at low doses of cageIL-10 (Fig. 1d). We then validated that aliskiren enables dose-
dependent control of hDIRECT (Fig. 1e) at therapeutically relevant concentrations <500 nM[26].
Decreasing the amount of renin plasmid increased the sensitivity of the system to aliskiren
(Supplementary Fig. 2). Western blot analysis confirms cageIL-10 cleavage by renin and
inhibition by aliskiren (Supplementary Fig. 3a).
To expand the cytokine output of hDIRECT, we next developed caged designs for IL-6.
Structurally, IL-6 is a monomeric four-α helical bundle typical of most cytokines (distinct from
the dimeric IL-10). Canonically, IL-6 binds to its high affinity receptor, IL-6Ra, on cell surfaces,
followed by complexing on the membrane with IL-6Rb (gp130) which transduces signal
activation across the membrane[38]. However, IL-6 activity has also been implicated in an
alternative pro-inflammatory pathway wherein IL-6 binds to soluble IL-6Ra (sIL-6Ra) and
mediates transactivation on cells expressing IL-6Rb[39]. To avoid transactivation of our caged IL-6
complex and bias IL-6 signaling to a regenerative phenotype, we sought to cage IL-6 with an IL-
6Ra mutant (A228D/N230D/H280S/D281V) that has abrogated activation of IL-6Rb. We denote
this IL-6Ra mutant as IL6Ra_M4.
Inspection of the IL-6-receptor complex structures suggested that placing IL-6Ra at the
N-terminus of the fusion protein would allow IL-6 to bind IL-6Ra in its canonical conformation.
Consistent with this observation, our data shows improved caging when the receptor is fused to
the N-terminus, although fusion at either terminus exhibits substantial renin-independent
activation (Supplementary Fig. 4). We hypothesized that caging could be improved if we fused
an additional receptor subunit. Therefore, we created a multi-cageIL-6 by fusing IL-6Ra_M4 to

the N-terminus and IL-6Rb to the C-terminus of IL-6 with renin-cleavable linkers (Fig. 2a).
Indeed, multi-cageIL-6 demonstrated vastly improved caging over the initial design (Fig. 2b),
albeit with reduced activity in the presence of renin (Fig. 2b). However, we found that IL-6
activity in response to renin can be increased at higher doses of multi-cageIL-6 plasmid (Fig. 2c).
Finally, to show broad generalizability in cytokines and therapeutic applications, we
ported over previous caged IL-2 constructs to hDIRECT[27]. A mutant IL-2 (sumIL-2) was
recently developed with enhanced affinity for IL-2Rb (i.e., effector T cells) over IL-2Ra (i.e.,
regulatory T cells)[28]. We leveraged this insight to design a caged sumIL-2 with a renin-cleavable
linker and C-terminal IL-2Rb (Fig. 2d). We also investigated whether varying the linker length
on either side of the renin cut site would improve cleavability but did not find any significant
effect (Supplementary Fig. 5). CageIL-2 is efficiently caged at high doses of cytokine (Fig. 2e)
and exhibits renin-activation at low doses (Fig. 2e). IL-2 activity was tuned sensitively by
aliskiren upon drug titration (Fig. 2f), indicating that aliskiren control is also generalizable
between different cytokines. Altogether, we established three different caged cytokines
compatible with renin-aliskiren control.
182
Engineering cleavable cytokines.
To expand the potential of protease regulation, we sought to design cytokines whose
activities would be abrogated (rather than activated) by cleavage. Compared to the caged design,
this modality would allow cytokine activity to be "turned on" or restored by a small-molecule
input. It could be particularly useful in applications where temporary, rather than constitutive,
cytokine activity is desired, such as transient IL-2 activation to support CAR-T cell proliferation
and cytotoxicity while avoiding exhaustion and overstimulation[40].
We hypothesized that cytokines could be directly inactivated if they were engineered to
include protease cut sites. We inserted protease cut sites within flexible loops of cytokines so as
not to perturb receptor binding regions. Like the caged cytokines, we envision engineered cells
constitutively expressing cleavable cytokines and membrane-bound renin. In this fashion,
cytokine activity can be tuned up on demand upon drug input and protease inhibition (Fig. 3a).
We started designing cleavable cytokines using the previously mentioned mono-IL-10.
Researchers engineered this functional mono-IL-10 by inserting a flexible, 6 amino acids,
GGGSGG linker between the D and E helices of human IL-10[37]. We leveraged this insight to
create a cleavable mono-IL-10 by replacing this glycine linker with the 13 amino acid renin
cleavage site. CleaveIL-10 retained activity in the absence of protease and was abrogated upon
co-transfection with renin (Fig. 3b). To explore the range at which cleaveIL-10 is sensitive to
renin cleavage, we co-transfected cells with or without renin while titrating the amount of
cleaveIL-10 DNA and observed a substantial shift of the titration curves by renin (Fig. 3c).
Next, we sought to demonstrate the generalizability of this method to native cytokines
that lack synthetic linkers to substitute. Our general method was to find sites in human IL-6 and
IL-2 tolerable (retaining activity) to insertion of a protease cut site. We scanned various solvent
exposed (protease accessible) flexible loops between helices to find tolerable insertions
(Supplementary Fig. 6) and then optimized linkers to improve their cleavability. Cleave IL-6 and
IL-2 were designed through direct insertion of the renin cut site with or without a five amino acid
GS linker into flexible loops at residues 127-140 in IL-6 and 74-81 in IL-2. Both cleaveIL-6
(Fig. 3d) and cleaveIL-2 (Fig. 3e) retained their functional activity in the absence of protease.
Additionally, we observed substantial shifts in response to renin in both cleaveIL-6 (Fig. 3d) and
cleaveIL-2 (Fig. 3e).

To demonstrate that this cleavable method can be used to turn on cytokine activity by a
small-molecule input, we determined that cleaveIL-10 activity can be tuned via aliskiren (Fig.
3f). Western blot analysis also confirms the cleavage of cleaveIL-10 by renin and inhibition by
aliskiren (Supplementary Fig. 3b). Taken together, these results demonstrate successful
engineering of cleavable cytokine constructs for three structurally distinct cytokines also
compatible with renin-aliskiren control.
219
220 Orthogonalization of hDIRECT from its human counterparts.
After establishing renin-regulated secreted cytokines, our next goal was to orthogonalize
the system from human hosts. In contrast to parts sourced from other organisms (e.g., viral
proteases), utilizing human parts bears the distinct and critical risk of off-target effects due to
crosstalk with other human proteins. In particular, renin is endogenously secreted into the blood
to regulate blood pressure via proteolytic cleavage of angiotensinogen[25]. Therefore, to avoid
crosstalk with the host, there are two issues to address: the activation of angiotensinogen by cell-
surface, exogenous renin and the cleavage of engineered cytokines by endogenous renin.
To address the first problem, we took inspiration from our previous work and
hypothesized that we could spatially separate renin from circulation by localizing it to the
endoplasmic reticulum (ER) of the engineered cell[16,41]. We fused renin to the CD4
transmembrane domain and a cytosolic ER-retention motif (RXR). Here, renin regulates
engineered cytokines in cis as the cytokines move through the secretory pathway, while
remaining sequestered from renin's endogenous substrate in the blood (Fig. 4a). Using cageIL-10
as a test case, we validated that ER-retained renin maintained the ability to activate secreted
cageIL-10 (Fig. 4b). Next, to ensure that aliskiren could permeate engineered cells and inhibit
ER-renin with a similar sensitivity (as it typically acts extracellularly), we performed aliskiren
titration and validated that ER-renin retains aliskiren sensitivity (Fig. 4c).
To address the second problem of endogenous renin cleaving the engineered cytokines,
we took inspiration from renin's diversity of specificities across species. In particular, human and
mouse angiotensinogen are not recognized by the opposing renins[42]. We hypothesized that we
could use the mouse cut site in engineered cytokines to avoid their cleavage by circulating
human renin, and in parallel minimally engineer human renin to recognize the mouse site such
that the protease is still unlikely to be recognized as non-human. To introduce mouse substrate
recognition, we compared substrate bound renin structures across species. Compared to human
renin, mouse renin has a larger binding pocket to accommodate bulkier substrate residues (L35
and Y36). We hypothesized that by mutating four substrate-binding residues in human renin to
corresponding residues in mouse renin with smaller side chains (L147I, R148H, I203V, L290V)
we could expand the binding pocket to accommodate the mouse substrate. We denote this mutant
renin as human-to-mouse (H2M) renin. A more detailed structural analysis can be found in
Supplementary Fig. 7. We then tested the orthogonality of the renin protease and found that the
human and mouse renins only recognize their corresponding cut sites while the H2M renin's
specificity is expanded to recognize both (Fig. 4d). We confirmed that H2M, ER-localized renin
remains sensitive to aliskerin (Fig. 4e) and that ER-localization does not affect orthogonality
(Fig. 4f).
In summary, human renin was successfully orthogonalized using structure-guided
mutation and localization motifs. Importantly, these orthogonalization strategies did not
compromise renin function or sensitivity to aliskiren. We expect that ER-retained H2M renin and

caged and cleavable cytokines harboring the mouse cut site will have minimal crosstalk with
endogenous human renin and angiotensinogen.
260
Therapeutically relevant applications of hDIRECT in mammalian cells.
Clinically, *ex vivo* gene delivery to engineer therapeutic cells is bottlenecked by viral
packaging limits of ~5 kb (adeno-associated virus) to ~10 kb (lentivirus)[43,44]. Compact genetic
constructs are therefore a critical requirement for translational applications. To demonstrate that
hDIRECT enables compact delivery, we created single-transcript constructs encoding engineered
cytokines and renin separated by 2A "self-cleaving" peptides[45].
We created a polycistronic construct containing cageIL-10 (mRens), ER-retained H2M
renin, and a fluorescent co-transfection marker, mCherry (Fig. 5a). Transient transfection of this
construct in HEK293 cells enables inhibition of IL-10 activity upon dosing with aliskiren, albeit
with reduced aliskiren sensitivity (Fig. 5a). We hypothesize this reduced sensitivity could be due
to overexpression of the renin protease, as had been seen in Supplementary Fig. 2. While it is
difficult to fine tune protein ratios in a polycistronic construct, we hypothesize methods to reduce
overall protein levels will improve aliskiren sensitivity, such as controlling the number of
genomic integrations or number of cells delivered. Next, we sought to demonstrate simultaneous
control over the activities of multiple cytokines, as immune and regenerative responses are often
mediated by a variety of cytokine signals with characteristic dosing and temporal profiles.[46,47]
For example, muscle regeneration is mediated by an initial pro-inflammatory regime (i.e. IL-6),
stimulating and recruiting the innate immune system, followed by an anti-inflammatory regime
(i.e. IL-10) to indirectly promote the differentiation of muscle stem cells.[48] To demonstrate that
hDIRECT can simultaneously control the activity of multiple cytokines, we created single-
transcript constructs containing both cageIL-10 and cleaveIL-6 (Fig. 5b) to mimic a cytokine
response for promoting muscle regeneration. In this manner, delivery with drug should allow an
initial regime of IL-6 activity, followed by a switch to IL-10 activity when drug is removed.
HEK293 cells transiently transfected with this single-transcript construct demonstrated
bidirectional control of IL-10 and IL-6 activity with aliskiren (Fig. 5b).
Next, we tested whether hDIRECT's performance is robust against different delivery
methods or cellular contexts. First, we created lentiviral constructs using a weaker constitutive
promoter, EF1-α, to potentially improve the aliskiren sensitivity of the polycistronic cageIL-10
design (from Fig. 5a). Indeed, stable transduction of HEK293 cells enabled regulation of IL-10
activity with improved aliskiren sensitivity (Fig. 5c). Additionally, we observed greater aliskiren
sensitivity at lower seeding densities of stably transduced cells (Supplementary Fig. 9). We also
conducted kinetic experiments of stable cageIL-10 cells to characterize the induction and
reversibility with aliskiren (Supplementary Fig. 10). Next, we sought to test hDIRECT in another
cell line to validate its generalizability to different cell types. We chose Jurkat cells as an
accessible proof of concept model for regulating IL-2 activity in the context of a CAR-T cell
therapy. We designed lentiviral constructs using the EF1-α promoter and a polycistronic cageIL-
2 design (Fig. 5d). Stably transduced Jurkats exhibited lower levels of secretion compared to
HEK293s but retained high sensitivity to aliskiren (Fig. 5d). To assess whether the level of
cytokine activities from our cells were therapeutically relevant, we conducted a series of
recombinant cytokine titrations on the HEKBlue reporter cells (Supplementary Figure 11). IL-6
and IL-10 have been reported to promote muscle regeneration at 10 ng/mL, within the detection
range of their respective reporter cells[49,50]. Notably, the apparent uncaged IL-2 concentration
from the stably transduced Jurkat cells corresponds with a cytokine concentration of 20 ng/mL

which is sufficient to stimulate primary T cell expansion (2-15 ng/mL)[51]. We note that while this
reporter assay may not be suited for granular cytokine quantification, it does give a general
indication whether cytokine activities are at or below these relevant concentrations.
Finally, we sought to demonstrate whether hDIRECT could function in therapeutically
relevant contexts, as well as more precisely quantify cytokine levels using enzyme-linked
immunosorbent assays (ELISAs). One potential application for hDIRECT could local, aliskiren-
switchable stimulation of immune cells for immunotherapy. To demonstrate this, we stably
engineered K-562 cells using the cageIL-2 lentiviral construct from Fig. 5d, so that the K-562s
will switch off IL-2 activity in response to aliskiren. The hDIRECT K-562s were co-cultured
with primary T cells in IL-2 free media (Fig. 6a), and, indeed, we observed aliskiren-regulated T
cell proliferation in response to the hDIRECT K-562s (Fig. 6b). Similarly, quantification of
supernatant IL-2 levels demonstrated aliskiren-dependent IL-2 activity (Fig. 6c). We note that
uncaged IL-2 levels were higher than the 200 U/mL recombinant IL-2 control typically used in T
cell cultures, and the residual caged IL-2 levels probably contributed to the proliferation in the
aliskiren– condition. The latter could be further tuned in the future by decreasing the number of
engineered cells delivered.
To further demonstrate hDIRECT's compact design and unqiue compatibility with
mRNA delivery for potential *in vivo* gene delivery applications, we synthesized an mRNA
transcript encoding the same cageIL-2 construct as above via *in vitro* transcription (Fig. 6d).
mRNA transfection of HEK293 cells with this construct retained therapeutically relevant
expression levels of IL-2 and aliskiren-dependent activity measured via ELISA (Fig. 6e).
To explore the potential for the core mechanism of hDIRECT to be tuned by aliskiren *in*
vivo, we conducted a pilot mouse study. We opted to use luciferase activity as a sensitive and
orthogonal output that can be easily measured from mouse sera. We developed a membrane-
tethered nanoluciferase that can be released by human renin, but not mouse renin, allowing for
aliskiren-regulated luciferase activity from engineered cells (Fig. 6f). To demonstrate aliskiren-
regulation *in vivo*, we used hydrodynamic gene delivery to introduce into the mouse liver
plasmids encoding membrane-bound renin-responsive luciferase, membrane-bound human renin,
and a constitutive co-secretion marker, SEAP, for signal normalization (Fig. 6g). Mice were
given 100 mg/kg doses of aliskiren via oral gavage 2 hours prior and after tail vein injection of
plasmids (Fig. 6h). Mice receiving aliskiren had reduced luciferase activity similar to that of the
negative control without renin expression (Fig. 6i).
These results demonstrate that hDIRECT can be delivered compactly using various
delivery methods and to multiple cell types, regulate cytokines at therapeutically relevant
concentrations, and be tuned exogenously *in vivo* through oral dosing of aliskiren.
339

Discussion

Here, we have presented a post-translational, humanized platform, hDIRECT, for drug-
mediated regulation of secreted cytokine activity in mammalian cells. Exclusively using human
proteins and domains, hDIRECT confers protease control with opposing signs ("caged" and
"cleavable") over cytokine activity. Co-expression of engineered cytokines renin allows the
control of cytokine activity using renin's FDA-approved inhibitor, aliskiren. We used species
specificity and re-localization strategies to orthogonalize both renin and engineered cytokines to
avoid crosstalk with their human counterparts. We showed that hDIRECT is compact and can
regulate the activities of multiple cytokines encoded on a single transcript in different
mammalian cells. Lastly, we demonstrated hDIRECT can control primary T cell proliferation, be

delivered compactly on an mRNA, and can be tuned via oral administration of aliskiren in mice.
hDIRECT therefore represents a proof-of-concept towards humanized engineering for post-
translational tools to control next-generation cell therapies.
353   While we have shown that hDIRECT can regulate cytokines, we expect that the
proteolytic control module can be extended to a variety of secreted and membrane proteins of
interest. Growth factors and chemokines, which regulate the proliferation, differentiation, and
migration of cells, could be similarly engineered for proteolytic regulation. In addition, we
envision that receptors with cleavable pro-domains or competitive caging domains might offer
small-molecule control of receptor availability which, for example, could tune the activity of
CAR-T cells to limit exhaustion[40].
360   We envision that hDIRECT will find immediate applications in cellular therapeutics.
hDIRECT could be used to temporarily supply CAR-T cells with IL-2 to improve proliferation
and cytotoxicity. In applications where the temporal profile of cytokine activity is critical, such
as MSC therapy for muscle regeneration, hDIRECT can be used to activate pro-inflammatory IL-
6 followed by regenerative, anti-inflammatory IL-10. While this work demonstrates the control
of two cytokines simultaneously, lentiviral packaging could allow for the delivery of a single
transcript encoding up to five engineered cytokines. hDIRECT could also be used for research in
basic science investigating the timing and dosage of various cytokine regimes by imparting
temporal control over the activity of multiple cytokines.
369   While our system is built from native human proteins, the presence of junctions from
fusion of domains and a small number of mutations introduced for improving function and
orthogonalization may still be recognized as foreign. Computational analysis of the ER-H2M
renin protein using peptide-MHC I binding and immunogenicity prediction algorithms[52,53]
confirmed that the protein contained less potentially immunogenic regions when compared with
commonly used viral proteases, TEVp and HCVp (Supplementary Fig. 12). While we note that
immunogenicity can only be definitively assessed in patients, *in vitro* assays, such as IFN-γ
ELISpots, can serve as a proxy to monitor donor T cell responses to antigenic peptides. Various
doses of a peptide pool containing all 9-mer peptides of the orthogonalizing H2M-renin
mutations did not elicit an immune response from T cells from any of the four healthy (HLA-
A*02:01) donors tested (Supplementary Fig. 13). Although this currently only assesses pre-
existing immunity, it could also be used to track acquired immunity in future patients. We
believe that this workflow can be used to screen, test, and iterate functional and immune-
tolerized portions over the rest of the junctions from the fusion of domains. We anticipate that
our humanization effort is a step towards immune-tolerized synthetic circuits.
384   While clinically suitable, hDIRECT still requires further work for potential use in
humans. Notably, most experiments here were conducted in the model mammalian cell line,
HEK293. Future work in primary cells more aligned with an intended clinical application, such
as CAR-T cells, could further demonstrate the functional efficacy of this platform. Subsequently,
in vivo experiments using mouse cytokines will help to determine the pharmacokinetics of small-
molecule dosing and localized cytokine activity in a more complex environment. It is important
to note that aliskiren has been shown to be tolerable and safe at oral doses of up to 600 mg in
healthy patients, including elderly[54,55]. This dose corresponds to a C_{max} of 300 ng/mL, which
relates to a drug concentration of 543 nM that we use to benchmark the sensitivity of
hDIRECT[26]. The most common adverse events reported at this dosing are headaches, dizziness,
and nausea but these occur at low incidence rates (1-5%) similar to placebo. To minimize side

effects, renin could be further engineered to be more sensitive to aliskiren so lower doses can be
used or the dose of cells delivered can be lowered.
In summary, hDIRECT is the first post-translational control platform using
orthogonalized humanized components in mammalian cells. Using hDIRECT, we demonstrate
small-molecule control of multiple secreted cytokines simultaneously, compact delivery using
various methods and in multiple cell types, as well as small-molecule control *in vivo*. We
envision that hDIRECT is generalizable towards a wide variety of extracellular factors and
receptors to tune them. Ultimately, we believe our novel engineered cytokines will facilitate
basic biology research by dynamically controlling cytokine activities, and the hDIRECT
platform holds the promise to enhance cell and gene therapies.
405

References

1.   Fischbach, M. A., Bluestone, J. A. & Lim, W. A. Cell-based therapeutics: the next pillar
of medicine. *Sci. Transl. Med.* **5**, 179ps7 (2013).
2.   Labanieh, L., Majzner, R. G. & Mackall, C. L. Programming CAR-T cells to kill cancer.
Nat. Biomed. Eng. **2**, 377–391 (2018).
3.   Ninagawa, N. T. *et al.* Transplantated mesenchymal stem cells derived from embryonic
stem cells promote muscle regeneration and accelerate functional recovery of injured
skeletal muscle. *Biores. Open Access* **2**, 295–306 (2013).
4.   Winkler, T. *et al.* Dose-response relationship of mesenchymal stem cell transplantation
and functional regeneration after severe skeletal muscle injury in rats. *Tissue Eng. Part A*
15, 487–492 (2009).
5.   Munir, H. & McGettrick, H. M. Mesenchymal stem cell therapy for autoimmune disease:
risks and rewards. *Stem Cells Dev.* **24**, 2091–2100 (2015).
6.   Squillaro, T., Peluso, G. & Galderisi, U. Clinical trials with mesenchymal stem cells: an
update. *Cell Transplant.* **25**, 829–848 (2016).
7.   Marofi, F. *et al.* CAR T cells in solid tumors: challenges and opportunities. *Stem Cell
Res. Ther.* **12**, 81 (2021).
8.   Preda, M. B. *et al.* Short lifespan of syngeneic transplanted MSC is a consequence of in
vivo apoptosis and immune cell recruitment in mice. *Cell Death Dis.* **12**, 566 (2021).
9.   Kim, N. & Cho, S.-G. New strategies for overcoming limitations of mesenchymal stem
cell-based immune modulation. *Int. J. Stem Cells* **8**, 54–68 (2015).
10.   Kerkar, S. P. *et al.* Tumor-specific CD8+ T cells expressing interleukin-12 eradicate
established cancers in lymphodepleted hosts. *Cancer Res.* **70**, 6725–6734 (2010).
11.   Nitahara-Kasahara, Y. *et al.* Enhanced cell survival and therapeutic benefits of IL-10-
expressing multipotent mesenchymal stromal cells for muscular dystrophy. *Stem Cell
Res. Ther.* **12**, 105 (2021).
12.   Chen, Z. & Elowitz, M. B. Programmable protein circuit design. *Cell* **184**, 2284–2301
(2021).
13.   Gao, X. J., Chong, L. S., Kim, M. S. & Elowitz, M. B. Programmable protein circuits in
living cells. *Science* **361**, 1252–1258 (2018).
14.   Fink, T. *et al.* Design of fast proteolysis-based signaling and logic circuits in mammalian
cells. *Nat. Chem. Biol.* **15**, 115–122 (2019).
15.   Sahin, U., Karikó, K. & Türeci, Ö. mRNA-based therapeutics--developing a new class of
drugs. *Nat. Rev. Drug Discov.* **13**, 759–780 (2014).

16.   Vlahos, A. E. *et al.* Protease-controlled secretion and display of intercellular signals. *Nat.*
Commun. **13**, 912 (2022).

17.   Praznik, A. *et al.* Regulation of protein secretion through chemical regulation of
endoplasmic reticulum retention signal cleavage. *Nat. Commun.* **13**, 1323 (2022).

18.   Ferdosi, S. R. *et al.* Multifunctional CRISPR-Cas9 with engineered immunosilenced
human T cell epitopes. *Nat. Commun.* **10**, 1842 (2019).

19.   Lamers, C. H. J. *et al.* Immune responses to transgene and retroviral vector in patients
treated with ex vivo-engineered T cells. *Blood* **117**, 72–82 (2011).

20.   Riechmann, L., Clark, M., Waldmann, H. & Winter, G. Reshaping human antibodies for
therapy. *Nature* **332**, 323–327 (1988).

21.   Jones, P. T., Dear, P. H., Foote, J., Neuberger, M. S. & Winter, G. Replacing the
complementarity-determining regions in a human antibody with those from a mouse.
Nature **321**, 522–525 (1986).

22.   Li, H.-S. *et al.* Multidimensional control of therapeutic human cell function with
synthetic gene circuits. *Science* **378**, 1227–1234 (2022).

23.   Rauch, S. *et al.* Programmable RNA-Guided RNA Effector Proteins Built from Human
Parts. *Cell* **178**, 122-134.e12 (2019).

24.   Zhu, I. *et al.* Design and modular assembly of synthetic intramembrane proteolysis
receptors for custom gene regulation in therapeutic cells. *BioRxiv* (2021)
doi:10.1101/2021.05.21.445218.

25.   Danser, A. H. J. & Deinum, J. Renin, prorenin and the putative (pro)renin receptor.
Hypertension **46**, 1069–1076 (2005).

26.   Vaidyanathan, S., Jarugula, V., Dieterich, H. A., Howard, D. & Dole, W. P. Clinical
pharmacokinetics and pharmacodynamics of aliskiren. *Clin. Pharmacokinet.* **47**, 515–531
(2008).

27.   Puskas, J. *et al.* Development of an attenuated interleukin-2 fusion protein that can be
activated by tumour-expressed proteases. *Immunology* **133**, 206–220 (2011).

28.   Hsu, E. J. *et al.* A cytokine receptor-masked IL2 prodrug selectively activates tumor-
infiltrating lymphocytes for potent antitumor therapy. *Nat. Commun.* **12**, 2768 (2021).

29.   Wang, X., Wong, K., Ouyang, W. & Rutz, S. Targeting IL-10 Family Cytokines for the
Treatment of Human Diseases. *Cold Spring Harb. Perspect. Biol.* **11**, (2019).

30.   Sarvas, J. L., Khaper, N. & Lees, S. J. The IL-6 Paradox: Context Dependent Interplay of
SOCS3 and AMPK. *J. Diabetes Metab.* **Suppl 13**, (2013).

31.   Serrano, A. L., Baeza-Raja, B., Perdiguero, E., Jardí, M. & Muñoz-Cánoves, P.
Interleukin-6 is an essential regulator of satellite cell-mediated skeletal muscle
hypertrophy. *Cell Metab.* **7**, 33–44 (2008).

32.   Zhang, C. *et al.* Interleukin-6/signal transducer and activator of transcription 3 (STAT3)
pathway is essential for macrophage infiltration and myoblast proliferation during muscle
regeneration. *J. Biol. Chem.* **288**, 1489–1499 (2013).

33.   Liao, W., Lin, J.-X. & Leonard, W. J. Interleukin-2 at the crossroads of effector
responses, tolerance, and immunotherapy. *Immunity* **38**, 13–25 (2013).

34.   Rosenberg, S. A. Interleukin 2 for patients with renal cancer. *Nat. Clin. Pract. Oncol.* **4**,
497 (2007).

35.   Hawkins, E. R., D'Souza, R. R. & Klampatsa, A. Armored CAR T-Cells: The Next
Chapter in T-Cell Cancer Immunotherapy. *Biologics* **15**, 95–105 (2021).

36.   Logsdon, N. J., Jones, B. C., Josephson, K., Cook, J. & Walter, M. R. Comparison of
interleukin-22 and interleukin-10 soluble receptor complexes. *J. Interferon Cytokine Res.*
22, 1099–1112 (2002).
37.   Gorby, C. *et al.* Engineered IL-10 variants elicit potent immunomodulatory effects at low
ligand doses. *Sci. Signal.* **13**, (2020).
38.   Garbers, C. *et al.* Plasticity and cross-talk of interleukin 6-type cytokines. *Cytokine*
Growth Factor Rev. **23**, 85–97 (2012).
39.   Wolf, J., Rose-John, S. & Garbers, C. Interleukin-6 and its receptors: a highly regulated
and dynamic system. *Cytokine* **70**, 11–20 (2014).
40.   Kwon, B. The two faces of IL-2: a key driver of CD8+ T-cell exhaustion. *Cell. Mol.*
Immunol. **18**, 1641–1643 (2021).
41.   Shikano, S. & Li, M. Membrane receptor trafficking: evidence of proximal and distal
zones conferred by two independent endoplasmic reticulum localization signals. *Proc*
Natl Acad Sci USA **100**, 5783–5788 (2003).
42.   Fukamizu, A. *et al.* Dependence of angiotensin production in transgenic mice carrying
either the human renin or human angiotensinogen genes on species-specific kinetics of
the renin-angiotensin system. *Arzneimittelforschung* **43**, 222–225 (1993).
43.   Wu, Z., Yang, H. & Colosi, P. Effect of genome size on AAV vector packaging. *Mol.*
Ther. **18**, 80–86 (2010).
44.   Kalidasan, V. *et al.* A guide in lentiviral vector production for hard-to-transfect cells,
using cardiac-derived c-kit expressing cells as a model system. *Sci. Rep.* **11**, 19265
(2021).
45.   Tang, W. *et al.* Faithful expression of multiple proteins via 2A-peptide self-processing: a
versatile and reliable method for manipulating brain circuits. *J. Neurosci.* **29**, 8621–8629
(2009).
46.   Subramanian, N., Torabi-Parizi, P., Gottschalk, R. A., Germain, R. N. & Dutta, B.
Network representations of immune system complexity. *Wiley Interdiscip. Rev. Syst. Biol.*
Med. **7**, 13–38 (2015).
47.   Orzechowski, A. Cytokines in skeletal muscle growth and decay. in *The plasticity of*
skeletal muscle (ed. Sakuma, K.) 113–139 (Springer Singapore, 2017). doi:10.1007/978-
981-10-3292-9_5.
48.   Yang, W. & Hu, P. Skeletal muscle regeneration is modulated by inflammation. *J.*
Orthop. Translat. **13**, 25–32 (2018).
49.   Steyn, P. J., Dzobo, K., Smith, R. I. & Myburgh, K. H. Interleukin-6 Induces Myogenic
Differentiation via JAK2-STAT3 Signaling in Mouse C2C12 Myoblast Cell Line and
Primary Human Myoblasts. *Int. J. Mol. Sci.* **20**, (2019).
50.   Deng, B., Wehling-Henricks, M., Villalta, S. A., Wang, Y. & Tidball, J. G. IL-10 triggers
changes in macrophage phenotype that promote muscle growth and regeneration. *J.*
Immunol. **189**, 3669–3680 (2012).
51.   Coppola, C. *et al.* Investigation of the Impact from IL-2, IL-7, and IL-15 on the Growth
and Signaling of Activated CD4+ T Cells. *Int. J. Mol. Sci.* **21**, (2020).
52.   Nielsen, M. & Andreatta, M. NetMHCpan-3.0; improved prediction of binding to MHC
class I molecules integrating information from multiple receptor and peptide length
datasets. *Genome Med.* **8**, 33 (2016).
53.   Calis, J. J. A. *et al.* Properties of MHC class I presented peptides that enhance
immunogenicity. *PLoS Comput. Biol.* **9**, e1003266 (2013).

54. Vaidyanathan, S. *et al.* Pharmacokinetics, safety, and tolerability of the novel oral direct renin inhibitor aliskiren in elderly healthy subjects. *J. Clin. Pharmacol.* **47**, 453–460 (2007).

55. Gradman, A. H. *et al.* Aliskiren, a novel orally effective renin inhibitor, provides dose-dependent antihypertensive efficacy and placebo-like tolerability in hypertensive patients. *Circulation* **111**, 1012–1018 (2005).

56. Kishina, M. *et al.* Therapeutic effects of the direct renin inhibitor, aliskiren, on non-alcoholic steatohepatitis in fatty liver Shionogi ob/ob male mice. *Hepatol. Res.* **44**, 888–896 (2014).

Methods

Plasmid generation. All plasmids were constructed using general practices. Backbones were linearized via restriction digestion, and inserts were generated using PCR, phosphorylated annealing, or purchased from Twist Biosciences. A complete list of plasmids used for each experiment (Supplementary Data 1) and the respective amounts used for transfection can be found in Supplementary Data 2. In addition, the DNA sequences of all the plasmids used in this study can be found in the source data, and all new plasmids with annotations will be available on Addgene.

mRNA synthesis. The DNA templates for in vitro transcription were prepared by PCR on a plasmid template containing hDIRECT coding sequences flanked by optimal 5' and 3' UTRs as well as a T7 promoter. The plasmid template with optimal UTRs was a gift from Prof. Michael Elowitz. Poly-A tail was added to the DNA template by PCR using Q5 PCR master mix (New England Biosciences, M0494L) with reverse primer containing a 200 bp polyA sequence. PCR products were cleaned and concentrated using DNA Clean & Concentrator kit (Zymo Research, D4033).

In vitro transcript mRNA was synthesized from DNA template using HiScribe IVT kit (New England Biosciences, E2040S) and using 100% N1-methyl-psuedouridine-5'-triphosphate modified (Trilink, N-1081-10) bases to improve RNA expression and with murine RNase inhibitor to improve RNA yield (New England Biosciences, M0314). mRNA was co-transcriptionally capped with a Cap 1 structure using CleanCapAG (Trilink, N-7113). Synthesized RNA was purified using Monarch RNA Cleanup kit (New England Biosciences, T2040).

Tissue culture. WT Human Embryonic Kidney (HEK) 293 cells (ATCC, catalog no. CRL-1573) and HEK293T-LentiX (Takara Biosciences, catalog no. 632180), were cultured in a humidity-controlled incubator under standard culture conditions (37 °C with 5% CO_2) in Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% fetal bovine serum (Fisher Scientific, catalog no. FB12999102), 1 mM sodium pyruvate (EMD Millipore, catalog no. TMS-005-C), 1× penicillin–streptomycin (Genesee, catalog no. 25-512), 2 mM l-glutamine (Genesee, catalog no. 25-509) and 1× MEM non-essential amino acids (Genesee, catalog no. 25-536). HEKBlue™ IL-2, IL-6, and IL-10 cells were purchased from Invivogen (catalog no. hkb-il2, hkb-hil6, hkb-il10, respectively). Cells were cultured as per manufacturer's protocol. T-REx™ Jurkat cells (Invitrogen, catalog no. R72207) and WT K-562 cells (gift from the Prof. Lacramioara Bintu, ATCC, catalog no. CCL-243) were cultured in a humidity-controlled

577   incubator under standard culture conditions (37 °C with 5% CO_2) in RPMI 1640 (Gibco, catalog
578   no.11875093, supplemented with 10% heat-inactivated fetal bovine serum (Fisher Scientific,
579   catalog no. FB12999102) and 1× penicillin–streptomycin (Genesee, catalog no. 25-512).
580
Transient transfection. Plasmid DNA. HEK293 cells were cultured in 96-well tissue culture-
treated plates under standard culture conditions. When cells were 70-90% confluent, the cells
were transiently transfected with plasmid constructs using the jetOPTIMUS DNA transfection
reagent (Polyplus, catalog no. 117-15), as per manufacturer's instructions using 0.375 μL of
reagent per 50 μL of jetOPTIMUS buffer for 500 ng total DNA transfections. Experiments
requiring incubation with aliskiren (Sigma Aldrich, catalog no. SML2077) were treated with
drug at time of transfection through media exchange. Drug was titrated from 0.5-500,000 nM of
aliskiren in 10x increments.
589
mRNA. HEK293 cells were cultured in 24-well tissue culture-treated plates under standard
culture conditions. When cells were 70-90% confluent, the cells were transiently transfected with
mRNA using the Lipofectamine™ MessengerMAX mRNA Transfection Reagent (Invitrogen,
LMRNA001), as per manufacturer's instructions using 1 μL reagent per 50 μL reaction/well for
a 500 ng total mRNA transfection. Experiments requiring incubation with aliskiren were treated
with drug at time of transfection through media exchange. Drug was titrated from 0.5-500,000
nM of Aliskiren in 10x increments.
597
Lentiviral transduction. LentiX cells were cultured in 6-well tissue culture-treated plates under
standard culture conditions. When cells were 70-90% confluent, the cells were transiently
transfected with plasmid constructs (600 ng PAX2, pMD2g, and 1,100 ng transfer target) using
the jetOPTIMUS DNA transfection reagent (Polyplus catalog no. 117-15), as per manufacturer's
instructions using 1.5 μL of reagent per 200 μL of jetOPTIMUS buffer for 2,000 ng total DNA
transfections. Cells were incubated for 24 h under standard culture conditions. Afterwards an
additional 3 mL of DMEM media was added carefully to each well. Lentivirus was concentrated
24 hrs afterward using viral precipitation. For each lentiviral prep, media was filtered using a
syringe and 0.45 μm filter into 15 mL conical tubes. 5x Lentivirus Precipitation Solution (Alstem
catalog no. VC100) was mixed with each prep and incubated at 4 °C overnight. Then, virus was
spun down at 1500xg for 30 min at 4 °C. Supernatant was aspirated, and virus was resuspended
using 200 μL Dulbecco's PBS (Genesee catalog no. 25-508). Virus was added dropwise onto 24-
well tissue culture plates containing HEK293, K-562, or Jurkat cells seeded at 250k cells/well.
72 h after incubation, the percentage of mCherry-positive cells were quantified using flow
cytometry. The cells were selected using 1000 ng/mL of puromycin (ThermoFisher Scientific;
catalog# J61278-MB) for one week until >90% of cells were mCherry-positive before being used
for experiments.
615
Cytokine reporter cell assay. Supernatant (10 μL) from HEK293 cells transiently transfected
24 h prior were incubated with 53,200 HEKBlue™ cells (190 μL) in 96-well tissue culture-
treated plates for 24 h under standard culture conditions. Afterwards, 20 μL HEKBlue™ cell
supernatant was collected and incubated with 180 μL Quanti-Blue reagent (Invivogen) for 2 h at
37 C as per manufacturer protocol. The mixture was read at 630 nm using the Spark® Multimode
microplate reader (TECAN). Cartoon schematic can be found in Supplementary Fig. 1.
622

For stable transductions, stable HEK293 cells were seeded at 7,500 cells (unless otherwise specified) in 96-well tissue culture-treated plates for 48 h under standard culture conditions. Media was changed either with or without aliskiren 24 h into the incubation. Supernatant transfer to HEKBlue^{TM} cells and assay proceeded the same as prior. Jurkat cells were seeded at 50,000 cells in U-bottom 96-well plates for 48 h under standard culture conditions. Cells were spun down in a centrifuge at 300 xg for 7 min and media was replaced with or without aliskiren 24 h into the incubation. 48 h after seeding, cells were spun down in a centrifuge at 300 xg for 7 min again and supernatant was transferred to HEKBlue^{TM} cells and assay proceeded the same as prior.

Western blot. HEK293 cells in 24-well tissue culture-treated plates were transiently transfected using transfection protocols mentioned prior. 24 hours after transient transfection, media was exchanged with serum-free, supplemented DMEM at half the volume (250 uL). Conditions requiring incubation with Aliskiren were treated with drug during media exchange. 48 hours after transient transfection, 16 μL of supernatant was incubated with 20 μL Tris-Glycine SDS sample buffer (2X) (ThermoFisher LC2676) and 4 μL NuPage Sample Reducing Agent (10X) (ThermoFisher NP0009) at 85 °C for 3 min. Samples and 5 μL PageRuler Prestained Protein Ladder (ThermoFisher 26616) were loaded onto 10-well Novex Tris-Glycine mini gels (16%) (ThermoFisher XP00160BOX).

Gels were transferred using a Mini Blot Module (ThermoFisher B1000) and a PVDF/Filter paper membrane sandwich (ThermoFisher LC2002). Sponges were soaked in 1x Tris-Glycine Transfer Buffer (ThermoFisher LC3675) and the PVDF membrane was activated by submerging in 100% ethanol prior to stacking. After blotting, the membrane was blocked using 5% nonfat dry milk (Apex catalog no. 20-241) in 1x DPBS (Genesee catalog no. 25-508) for 1 hour at room temperature with shaking. After blocking, the membrane was incubated with a primary Rb anti-myc antibody (Abcam, catalog AB9106) at a 1/2500 dilution in a 1% bovine serum albumin (BSA) (UniRegion, UR-BSA001-100G) solution with DPBS overnight at 4 °C with shaking. The following day, the membrane was washed in 1x DPBS with 0.1% Polysorbate 20 (ThermoScientific, catalog J66278.AE) for 15 min at room temperature with shaking. The wash was repeated three more times, 5 minutes each, with shaking at room temperature. After the final wash, the membrane was incubated with a secondary anti-Rb IgG antibody (Abcam, AB205718) (1/1000 dilution) in a 1% BSA solution with DPBS for 1 hour at room temperature with shaking. After incubation, the membrane was washed again for 3 additional times, 5 minutes each, at room temperature with shaking. The membrane was incubated with an ECL substrate kit (Abcam, AB133406) for 2 minutes prior to imaging on an iBright CL1500 imager (Invitrogen) using the ChemiBlot protocol.

Kinetic experiments. Induction. Stable HEK293 cells expressing the single transcript construct were split at 15,000 cells/well in a tissue-culture treated 96-well plate. 24 hours later, media was exchanged with fresh media or media containing 5,000 nM aliskiren. 10 μL supernatant was transferred from individual biological replicates at varying timepoints (30 min to 24 hrs) to another 96-well plate which was stored at 4 °C to preserve the samples. After the final supernatant collection, 50,000 HEKBlue^{TM} IL-10 cells/well (190 μL) was transferred to the 96-well plates containing the supernatant samples and incubated for 24 h under standard culture conditions. Afterwards, 20 μL HEKBlue^{TM} IL-10 cell supernatant was collected and incubated

with 180 µL Quanti-Blue reagent (Invivogen) for 2 h at 37 C as per manufacturer protocol. The
mixture was read at 630 nm using the Spark® Multimode microplate reader (TECAN).

Reversibility. Cells were prepared in the same manner as the "Induction" section. 2 hours after
seeding, media was exchanged with fresh media or media containing 5,000 nM aliskiren. 24
hours later, drug was removed from all conditions with a fresh media change. Supernatant was
transferred, stored, and assayed in the same manner as the "Induction" section.

HEKBlue reporter cell recombinant cytokine titration. Fresh serial dilutions of recombinant
human IL-2 (PeproTech, AF-200-02), IL-6 (PeproTech, 200-06), and IL-10 (PeproTech, 200-10)
were prepared with PBS. 10 µL samples at each concentration were incubated with 50,000
HEKBlue™ cells (190 µL) in 96-well tissue culture-treated plates for 24 h under standard culture
conditions. HEKBlue™ cell supernatant was assayed for reporter cell activity as previously
mentioned in "cytokine reporter cell assay" above.
HEKBlue™ IL-2 activity standard curve was obtained using a three-parameter nonlinear fit with
GraphPad Prism 10 to relate OD (630 nM) measurements to cytokine concentration (ng/mL).

$$Y = Bottom + X * \frac{Top - Bottom}{EC50 + X}$$

686 Or

$$X = EC50 * \frac{Y - Bottom}{Top - Y}$$

Where Y = OD (630 nM) and X = [Cytokine concentration] (ng/mL).

Flow cytometry. Flow Cytometry was performed on ZE5 Cell analyzer (BioRad). Samples were
prepared in 50 µL culture media and diluted 1:1 with Hank's Balanced Salt Solution (Cytiva,
SH30588.01) + 0.25% BSA and measured in biological triplicate. Live cells were gated via
forward (FSC) and side (SSC) scatter. For mCherry-expressing cells, fluorescence was measured
using 561nm excitation laser with a 615/24 nm emission filter. Flow cytometry data was
analyzed using FlowJo (BD). T cells were gated from engineered K-562 cells by gating on the
presence of mCherry and total cell counts for mCherry⁻ cells were determined. Flow cytometry
data was further processed using Prism 9 software (GraphPad).

Fluorescence activated cell sorting (FACS). To sort K-562 cells stably integrated with
hDIRECT-IL2 cassettes, samples were concentrated to $5x10^6$ cells/mL in 250 uL. Lives cells
were gated by forward scatter (FSC) and side scatters (SSC). Fluorescence data was collected for
mCherry (excitation laser: 561nm) Gated populations of cells expressing mCherry were sorted
on a FACSAria II (BD Biosciences) into fresh RPMI 1640 + 10% heat inactivated FBS + 1 %
penicillin–streptomycin. K-562 cells were cultured for 3 days and allowed to reach confluency of
$1x10^6$ cells/mL before staring co-culture experiments.

Human primary T cell culture. Primary T cells were isolated from a peripheral blood
mononuclear cells (PBMCs) Leuko Pak from a healthy donor (StemCell, catalog no. 70500.1) by
using the EasySep Human T Cell Isolation kit (STEMCELL, catalog no. 17951). Cells were
frozen down using Cellbanker 1 (AimsBio, 11910). Cells were cultured in X-Vivo15 medium,
(Lonza, catalog no. 04418Q) supplemented with 1 mM N-Acetyl-L-cysteine (Thermo scientific,
A15409.22), 55 uM 2-Mercaptoethanol (Gibco, 21985-023), 5% heat inactivated human AB

serum (BioIVT, HP1022HI), pH balanced with 1N NaOH (Fisher Scientific, SS255-1), and 200
U/mL IL-2 (Fredrick National Lab BRB Preclinical Repository).
715

In vitro T cell proliferation co-culture. T cells were thawed and plated in 24-well plates at a
density of 1×10^6 cells/mL. T cells were activating using CD3/CD28 Human T-Activator
Dynabeads (Gibco, 11131D) at a 3:1 bead:cell ratio for 48 hours. Beads were removed via
magnetic separation, and cells were cultured for 48 hours prior to proliferation assay.
T cells were counted using Denovix CellDrop automated cell counter and plated in a 96-well
untreated round-bottom plate at 25,000 cells/well in 50 µL media. T cells were co-cultured with
25,000 cells of either sorted, hDIRECT engineered K-562 cells or wild-type K-562 cells, both
engineered to express mCherry. Cell culture was brought up to 100 uL X-Vivo 15 media per
well. Populations were treated with either 5000 nM Aliksiren, vehicle control, or 200 U/mL IL-2.
725

726   T cell counts were analyzed using flow cytometry to count live cells by gating on FSC/SSC and
727   mCherry. Cells were prepared for flow cytometry on days 0, 2, 4, and 6 by removing 50 uL of
728   well-dispersed cell culture and diluting 1:1 in HBSS + 0.25% BSA. On day 3, cells were
729   centrifuged at 100xg for 10 minutes and media was replaced. Spent media was analyzed for IL-2
730   concentration via ELISA.
731

Human IL-2 ELISA. ELISA was performed with 50 µL of supernatant from cultured T cells
using Human IL-2 ELISA (Abcam, ab270883) according to manufactures instructions.
Absorbance was measured at 450 nm using Tecan Infinite M Plex microplate reader. All ELISA
measurements were performed in biological triplicates. Samples with absorbance higher than
standard ladders provided were diluted until within range of the provided standards. IL-2
concentrations were calculated via interpolation of a four-parameter curve fit, and diluted
samples were adjusted via linear scaling in accordance with the manufacturer recommendation.
739

In vivo control of human renin via aliskiren.
Animal studies were conducted at Stanford University with the approval of the Stanford
Administrative Panel on Laboratory Animal Care (#34743). Female BALB/c mice 12-14 weeks
of age were randomly assigned to each experimental group. Aliskiren hemifumarate
(MedChemExpress, HY-12177) was prepared at 20 mg/mL in PBS pH 7.4 (Gibco, 10010023)
and given at a dose of 100 mg/kg p.o. 2 hours prior to plasmid delivery via 1 mL syringe with a
20G 1.5" feeding tube (BrainTree, DT9920). Following initial drug dosing, plasmids were
delivered via hydrodynamic gene delivery. All mice received plasmid delivery of 8 µg of
membrane-bound, renin cleavable NanoLuciferase and 4 µg of CMV-driven secreted embryonic
alkaline phosphatase (SEAP) as a co-secretion marker. Treatment groups received an additional
4 ug of CMV-driven human renin plasmid. Plasmids were diluted in 2 mL Trans-IT EE delivery
solution (Mirus, MIR 5340) were delivery intravenously via tail vein using 5 mL syringes with a
27G needle (Air-Tite, N2712) in less than 10 seconds. Aliskiren dosing was repeated at 100
mg/kg 2 hours following plasmid delivery. 4 hours following final aliskiren administration,
blood samples were collected via saphenous vein puncture into a clotting capillary tube for
serum collection. High doses of aliskiren were given to mice due to the low bioavailability of
aliskiren in mice relative to humans[56]. Samples were centrifuged at 2000xg for 10 minutes to
isolate serum. Serum levels of cleaved NanoLuciferase were measured by diluting 5 µg of serum
in 45 µL of DI water and NanoLuciferase was measured using NanoGlo Luciferase Assay

system (Promega, N1120). Luminescence was measured using Infinite M Plex microplate reader
(TECAN) with a 1000 ms integration time. Serum luciferase levels were normalized to SEAP
serum values measured using Phospha-Light SEAP Reporter Gene Assay system (Applied
Biosciences, T1015) by diluting 5 µL serum in 45 µL of the provided diluent.
Chemiluminescence was measured using Infinite M Plex microplate reader (TECAN) with a 100
ms integration time.
765
Bioinformatic Analysis of Predicted MHC-I Binding and Immunogenicity. Immunogenicity
of viral proteases was compared to engineered human proteases using the Immune Epitope
Database & Tools (IEDB) platform (https://www.iedb.org/). Peptide sequences were obtained
from literature for TEVp and HCVp and directly compared with the sequence from ER-H2M-
Renin, the most engineered human protease reported here. The sequence for each peptide was
sequentially tiled into peptides of length 9, and a Protein BLAST using NCBI blastp
(https://blast.ncbi.nlm.nih.gov/Blast.cgi) was performed on each 9-mer peptide to search for
exact sequence matches to human proteins in the Protein Data Bank. Any exact matches to
human protein sequences were excluded from further analysis, as these peptides were assumed to
be non-immunogenic and therefore a source of potential false positives. Nonhuman 9-mer
peptide sequences were then analyzed using IEDB's MHC I Binding Affinity predictor
(NetMHCpan4.1 BA), where peptides with binding affinities less than or equal to 500 nM are
considered likely to bind to and be presented by MHC I complexes[52]. HLA-A*02:01 was
selected for our analysis since it is the most commonly analyzed HLA allele. The MHC I
Immunogenicity predictor was also used with default settings to get predictions of how likely
each pMHC complex would be to elicit an immune response.
782
IFN-γ ELISPOT. H2M-renin peptide immunogenicity was assayed using a human IFN-γ single-
color enzymatic ELISPOT assay kit (CTL, hIFNg-1M/2) according to the manufacturer's
recommendation. A custom synthesized H2M-renin peptide pool was purchased from Elim
Biopharm (28 9-mer peptides, 1mg/peptide). Peptide pool was initially reconstituted to 1 mg/mL
in sterile PBS. Frozen vials of PBMCs from four healthy donors (HLA-A*02:01) were purchased
from (StemCell, catalog no. 70107). 96-well plates were coated with a human IFN-γ capture
solution and sealed overnight with parafilm at 4 C.
790
The next day, donor PBMCs were rapidly thawed in a 37C water bath and washed with RPMI
1640 + 1% L-glutamine (Genesee, catalog no. 25-509) prior to 10 minutes of centrifugation at
330 xg. Cells were counted using Denovix CellDrop automated cell counter and resuspended in
CTL-Test medium (CTL, CTL-010) to $3x10^6$ cells/mL. 2x antigen solutions of the peptide pool
were prepared at 0.2, 2, 20 µg/mL using CTL-Test media. CERI-MHC Class I Control Peptide
Pool (CTL, CTL-CERI-300) was diluted 1:20 using CTL-Test media. The coated plates were
decanted and washed with sterile PBS. Antigen (H2M-renin peptide pool dilutions), positive
control (CERI dilution), and negative control (CTL-Test medium) were plated at 100 µL/well in
triplicates and incubated for 10 minutes in a 37 C incubator. PBMCs were plated at 100 µL/well
(300k cells/well) onto sample wells and incubated for 24 hours at 37 C.
The next day, plates were washed a total of three times with sterile PBS and 0.05% Tween-PBS
prior to incubating with 80 µL/well of anti-human IFN-γ detection solution for 2 hours at room
temperature. Plates were washed three times again with 0.05% Tween-PBS prior to incubating
with 80 µL/well of a Tertiary Solution for 30 minutes at room temperature. Plates were washed

with 0.05% Tween-PBS twice, followed by two washes with distilled water. Plates were incubated with 80 µL/well of Blue Developer Solution for 15 minutes at room temperature and the reaction was stopping by rinsing the membrane with tap water three times. Plates were air-dried overnight at room temperature, avoiding exposure to light. The plates were sent to Cellular Technology Limited (CTL) for scanning and analysis of spot forming units (SFU). Wells unable to be precisely quantified due to overly numerous spot formations are labeled too numerous to count (TNTC).

Statistical analysis. For experiments comparing two groups, an unpaired two-tailed Student's *t*-test was used to assess significance. Single, double, triple, and quadruple asterisks indicate $P < 0.05$, $P < 0.01$, $P < 0.001$ and $P < 0.0001$ unless corrected for multiple comparisons using a Bonferroni correction. EC50 and IC50 measurements for curves were calculated using GraphPad Prism and reported as mean +- standard error of the mean (s.e.m.). A one-way ANOVA was used to compare the means among different experiment groups. All statistical analyses were performed using GraphPad Prism 10.0.0.

Data availability.

All data reported in this paper are available from the corresponding author on request. Plasmid maps for all plasmids used in this study will be available as GenBank files. The plasmids used in this study will be deposited on Addgene, with complete and annotated GenBank files on their website. All figures and supplemental figures have associated raw data. There are no restrictions on data availability.

Acknowledgements.

This research was supported by NIH (R00EB027723, DP2OD034951; X.J.G), Longevity Impetus Grants (X.J.G.), Wu Tsai Human Performance Alliance Agility Project Grant (X.J.G.), Stanford Bio-X Interdisciplinary Initiatives Seed Grant Program (IIP) [R11-7] (X.J.G.), the Stanford Graduate Fellowship (C.A.), the Sarafan ChEM-H CBI training program (C.A.), the Bio-X Stanford Interdisciplinary Graduate Fellowship (C.A.), the CMB training grant NIH (T32 GM007276; C.C.C.), the National Science Foundation Graduate Research Fellowship (DGE-2146755; C.C.C.), and the International Human Frontier Science Program Organization (LT000221/2021-L; A.E.V.).

Author contributions.

C.A.A. and X.J.G. conceived and directed the study. C.A.A. designed and performed all experiments for the protein engineering and characterization. C.C.C. designed and performed the T cell proliferation and in vivo study. A.E.V. created the transmembrane domain and ER-retention motifs for renin constructs, as well as performed tail vein injections for the in vivo study. J.P. and Q.C. provided recommended mutations for the H2M renin. L.E.S. conducted the computational immunogenicity analysis. C.A.A. and C.C.C. analyzed the data for the manuscript. C.A.A., X.J.G., and C.C.C. wrote the manuscript. All authors provided feedback on the manuscript.

Competing interests.

The board of trustees of the Leland Stanford Junior University have filed a patent on behalf of the inventors (C.A.A. and X.J.G.) of the small-molecule control of membrane and secreted

851   proteins using human proteases platform described (US provisional Application No. 63/458833).
852   An additional provisional patent application has also been filed on behalf of the inventors
853   (C.A.A., X.J.G., J.P, and Q.C.) for the orthogonalization of human renin. X.J.G. is a co-founder
854   and serves on the scientific advisory board of Radar Tx.
855

Figures:

Fig. 1: Engineering caged IL-10. a, Caged cytokine design for dimeric and monomeric IL-10. Cytokines are fused with renin-cleavable linkers to cognate receptor subunits, inhibiting binding to cellular receptors. **b,** hDIRECT circuit using caged cytokines to control cytokine activity. Cells constitutively secrete attenuated caged cytokines and cytokine activity can be tuned down via small-molecule inhibitor for co-transfected protease, renin. **c,** Functional activity of caged IL-10 constructs with or without renin co-expression. **d,** Functional activity of cageIL-10 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **e,** Functional activity of cageIL-10 co-expressed with renin at increasing aliskiren concentrations. (For **c-e**: Cytokine activity measured from supernatant using HEKBlue IL-10^{TM} reporter cells. For **c**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **d-e**: n = 2 biological replicates; data represent mean ± s.e.m. *** p ≤ 0.001, **** p < 0.0001.)

857
858

Fig. 2: Engineering caged IL-6 and IL-2. a, Caged cytokine design for single and multi-caged IL-6 constructs. **b,** Functional activity of caged IL-6 constructs with or without renin co-expression. **c,** Functional activity of multi-cageIL-6 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **d,** Caged cytokine design for caged IL-2. **e,** Functional activity of cageIL-2 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **f,** Functional activity of cageIL-2 co-expressed with renin at increasing aliskiren concentrations. (Cytokine activity measured from supernatant using HEKBlue IL-6^{TM} (**b-c**) and IL-2^{TM} (**e-f**) reporter cells. For **b**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **c,e-f**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

859
860

Fig. 3: Engineering of cleavable cytokines. a, hDIRECT circuit using cleavable cytokines to control cytokine activity. Cells constitutively secrete cleavable cytokines and cytokine activity can be tuned via small-molecule inhibitor for co-transfected protease, renin. **b,** Functional activity of cleaveIL-10 with or without renin co-expression. **c,** Functional activity of cleaveIL-10 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **d,** Functional activity of cleaveIL-6 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **e,** Functional activity of cleaveIL-2 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **f,** Functional activity of cleaveIL-10 co-expressed with renin at increasing aliskiren concentrations. (Cytokine activity measured from supernatant using HEKBlue IL-10[TM] (**b-c,f**), IL-6[TM] (**d**), and IL-2[TM] (**e**) reporter cells. For **b**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **c-f**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

861
862
863
864

Fig. 4: Orthogonalization of renin. a, hDIRECT circuit using caged cytokines and ER-retained renin to control cytokine activity. Cells constitutively secrete caged cytokines and cytokine activity can be tuned via small-molecule inhibition of a co-expressed protease, renin. ER-renin is retained in ER via cytosolic motif and can cleave secreted cytokines. **b,** Activity of cageIL-10 transiently transfected with membrane (memRenin) or ER-retained (ER-renin) renin. **c,** Activity of cageIL-10 co-expressed with ER-renin at increasing aliskiren concentrations. **d,** Orthogonal renin matrix. Functional activity of cageIL-10 constructs containing human (h) or mouse (m) cut sites transiently transfected with membrane-bound human, mouse, or engineered H2M renin. **e,** Functional activity of cageIL-10 with an mRenin cut-site co-expressed with H2M ER-renin at increasing aliskiren concentrations. **f,** Orthogonal renin matrix. Functional activity of cageIL-10 constructs containing human (h) or mouse (m) cut sites transiently transfected with ER-retained human, mouse, or engineered H2M renin. Replicate data for **d,f** can be found in Supplementary Fig. 8. (Cytokine activity measured from supernatant using HEKBlue IL-10™ (**b-f**) reporter cells. For **b,d,f**: mean of n = 3 biological replicates; Individual replicate data for **d,f** can be found in Supplementary Fig. 5. Unpaired two-tailed Student's t-test. For **c,e**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

Fig. 5: Compact delivery of hDIRECT in mammalian cells. a, Polycistronic design of co-expressed cageIL-10 (mRens) and ER-H2M renin (top cartoon). Aliskerin-regulated activity of cageIL-10 in HEK293 cells transiently transfected with cageIL-10 single-transcript plasmid. **b,** Polycistronic design of cageIL-10 (mRens), cleaveIL-6 (mRens), and ER-H2M renin (top). Aliskiren-regulated activity of cageIL-10 (left-axis) and cleaveIL-6 (right-axis) in HEK293 cells transiently transfected with cageIL-10/cleaveIL-6 single-transcript plasmid. **c,** Aliskiren-regulated activity of cageIL-10 in HEK293 cells stably transduced with cageIL-10 single-transcript design (from **a**). **d,** Polycistronic design of co-expressed cageIL-2 (mRens) and ER-H2M renin (top cartoon). Aliskiren-regulated activity of cageIL-2 at increasing aliskiren concentrations from Jurkat cells stably transduced with cageIL-2 single-transcript design. (Cytokine activity measured from supernatant using HEKBlue IL-10[TM] (**a-c**), IL-6[TM] (**b**), and IL-2[TM] (**d**) reporter cells. For **a-d**: n = 2 biological replicates; data represent mean ± s.e.m.)

Fig. 6: hDIRECT primary T cell proliferation co-culture. a, CageIL-2 hDIRECT circuit to control primary T cell proliferation using stably engineered K-562 cells via *in vitro* co-culture. **b,** Primary T cells were co-cultured with engineered K-562 cells (single-transcript lentivirus from Fig. 5d) in IL-2-free media at a 1:1 ratio. Cells were plated and induced with aliskiren (0 or 5000 nM) on Day 0 and live-cell counts were quantified via flow cytometry at the indicated days. Untransduced (wild-type) and cells supplemented with recombinant IL-2 (200 U/mL) were cultured and included as positive and negative controls. Asterisks represent p-values for t-tests between -Aliskiren and +Aliskiren conditions. **c,** Aliskiren-regulated IL-2 activity from stably engineered K-562 cells. Supernatant on Day 3 from T cell proliferation co-culture wells were assayed for IL-2 binding via ELISA. **d,** Single-transcript design of cageIL-2 hDIRECT circuit for mRNA delivery. **e,** Aliskerin-regulated level of cageIL-2 secreted from HEK293 cells transiently transfected with mRNA from (**d**) measured via IL-2 ELISA. **f,** hDIRECT circuit using membrane-bound nanoluciferase (NLuc) to control luciferase activity. Cells constitutively display NLuc on their surface which is cleaved free by renin. Luciferase activity can be tuned down via aliskiren. **g,** Testing exogenous renin's response to aliskiren *in vivo*. Cartoon of hydrodynamic gene delivery (HGD) of plasmids mice were given depending on condition. SEAP plasmid was used as a co-secretion factor to normalize total secretion. **h,** Timeline of *in vivo* mouse experiment, in which BALB/c mice were injected i.v. with plasmids via HGD. Mice were treated with 100 mg/kg of aliskiren or vehicle in two doses with oral gavage prior to and after plasmid delivery. Serum was collected to assay for target protein activity. **i,** Aliskiren-regulated activity of membrane-bound luciferase in mice. Serum was assayed for luminescence and normalized to SEAP activity. (For **b:** Lines represent mean values for n = 3 biological replicates ± SD. Significance was tested by unpaired two-tailed Student's t-test among multiple conditions with a Bonferroni correction for m = 3 conditions. * p<0.0167, ** p<0.0033, *** p<0.0003. For **c:** mean of n = 3 biological replicates. Unpaired two-tailed Student's t-test. **** p<0.0001. For **e:** n = 2 biological replicates; data represent mean ± s.e.m.. For **g:** error bars represent mean and standard deviation of n = 3 or 4 mice. One-way ANOVA with Tukey's multiple comparisons test. * p<0.05, *** p<0.001.)

871