Orthogonalized human protease control of secreted signals

Carlos A. Aldrete[1], Alexander E. Vlahos[1], Jimin Pei[2], Qian Cong[2,3,4], and Xiaojing J. Gao[1]

1 Department of Chemical Engineering, Stanford University, Stanford CA 94305, USA
2 Eugene McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA
3 Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA
4 Harold C. Simmons Comprehensive Cancer Center, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

Abstract

Synthetic circuits that regulate protein secretion in human cells could support cell-based therapies by enabling control over local environments. While protein-level circuits enable such potential clinical applications, featuring orthogonality and compactness, their non-human origin poses a potential immunogenic risk. Here, we developed Humanized Drug Induced Regulation of Engineered CyTokines (hDIRECT) as a platform to control cytokine activity exclusively using human-derived proteins. We sourced a specific human protease and its FDA-approved inhibitor. We engineered cytokines whose activities can be activated and abrogated by proteolytic cleavage. We utilized species specificity and re-localization strategies to orthogonalize the cytokines and protease from the human context that they would be deployed in. hDIRECT should enable local cytokine activation to support a variety of cell-based therapies such as muscle regeneration and cancer immunotherapy. Our work offers a proof of concept for the emerging appreciation of humanization in synthetic biology for human health.

Intercellular communication, mediated by the secretion and display of soluble factors
(cytokines and growth factors) and receptors, plays a crucial role in health and diseases. Cell
therapies aim to treat complex diseases using human cells through the modulation of intercellular
communication[1], the chimeric antigen receptor (CAR) T cell therapy being the most notable
example[2]. Aside from treating cancer, another promising example is treating autoimmune
diseases and support muscle regeneration using mesenchymal stem cells (MSCs)[3–6]. Despite their
promise, many cellular therapeutics are susceptible to changes in the local microenvironment[7–9].
To address this hurdle, researchers have "armed" cell therapies to constitutively secrete
cytokines to tune their local environments to promote proliferation and efficacy, such as driving
CAR-T activation in solid tumors using IL-12[10], and improving the survival of transplanted
MSCs using the anti-inflammatory IL-10[11]. While constant expression of these powerful signals
improve efficacy, increased toxicity was also observed, indicating a need for further control[10].
The field of synthetic biology offers a plethora of tools and methods for localized,
conditional control of cytokine activity. In particular, compared to the more conventional
transcriptional control, protein-level controls hold advantages for potential clinical use such as
fast operation, compact single-transcript delivery, and context-independent performance[12].
Notably, orthogonal viral proteases have emerged as useful post-translational tools due to their
ability to control the activity, degradation, and localization of target proteins[13,14]. These
orthogonal proteases are highly versatile and can be combined to implement robust sense-and-
response behaviors in mammalian cells. They are uniquely positioned to serve as cargos on
mRNA vectors that are safer and more scalable than conventional DNA vectors[15]. They have
been successfully applied to the control of protein secretion and display[16,17].
However, despite the promise of proteases as control knobs, there remains a critical
constraint: the non-human origin and potential immunogenicity of viral proteases may preclude
their translational potential. In analogous scenarios, immune responses against other synthetic
biology tools such as *Streptococcus pyogenes* Cas9 protein (SpCas9) and synthetic CAR-T
receptors have been documented with compromised efficacy and duration, highlighting the need
to reduce immunogenicity of synthetic tools[18,19]. Even more notably, we are inspired by the
decades of efforts to "humanize" monoclonal antibodies[20,21], and we reason that, if synthetic
biology is to deliver its promise in human health, it is imperative that we preempt similar
challenges as we engineer the fundamental tools. In an emerging concensus to "humanize"
synthetic biology, groups have developed tools derived largely from human-derived components
such as zinc finger transcriptional regulators (synZiFTRs)[22], CRISPR-CAS-inspired RNA
targeting systems (CIRTS)[23], and synthetic intramembrane proteolysis receptors (SNIPRs)[24].
While these papers have demonstrated humanized transcriptional and translational regulation, the
humanization efforts have yet to reach the post-translational level.
Here, we sought to develop protease-based control of intercellular communication for
potential application in cellular therapeutics. We developed our system with five features for
potential clinical suitability, combining general humanization considerations[22] with the unique
benefits of protease control: (i) use of human-derived proteins to minimize the risk of
immunogenicity; (ii) use of orthogonalized components with high specificity and minimal
crosstalk with native signaling pathways; (iii) ability to control exogenously using FDA-
approved small molecules; (iv) compact, single-transcript design to facilitate delivery into
therapeutic cells; (v) compatibility with mRNA delivery for potential *in vivo* gene delivery
applications.

We identified a human protease, renin, as the control knob. Renin, in contrast to most
human proteases, exhibits high substrate specificity[25]. Additionally, renin comes with a
clinically-viable mode of exogenous control in the form of its FDA-approved small-molecule
inhibitor, aliskiren[26]. In contrast to previously used cytosolic proteases, renin functions
extracellularly and in the secretory pathway, offering an external interface to directly control
intercellular communication.
In this paper, we introduce Humanized Drug Induced Regulation of Engineered
CyTokines (hDIRECT), using renin to control intercellular communication. We engineered a
variety of cytokines, whose activities are increased or decreased by renin-mediated proteolysis
and can be tuned exogenously via aliskiren. We used structure-guided mutagenesis and re-
localization strategies to create renin mutants with new and orthogonal substrate specificity from
wildtype proteins. Finally, we showed that cytokine activity can be tuned with aliskiren in stably
transduced mammalian cells. We envision that conditional control of cytokine activity has
immediate use for supporting and controlling cell therapies. This study demonstrates a proof-of-
principle for the orthogonalization of human proteases and serves as an example of making
biomolecular controls more clinically compatible.
89

Results

Engineering caged cytokines.
hDIRECT consists of three components: a membrane-bound renin, renin's FDA-
approved small-molecule inhibitor, aliskiren, and engineered cytokines modulated by renin
cleavage (Fig. 1a, b). Renin is known to function extracellularly as it is natively secreted by
juxtaglomerular cells in the kidney[25]. To localize renin to the plasma membrane to enable the
autonomous control of engineered cells, the active form of renin was fused to the transmembrane
domain of human CD28 (Fig. 1b). Here, constitutive expression of membrane-bound renin can
control the activity of co-expressed secreted cytokines. We first evaluated a variety of strategies
to place cytokines under the control of renin.
To engineer cytokines activated by renin cleavage, we drew inspiration from previous
work on attenuated cytokines, specifically, receptor-masked interleukin-2 (IL-2).[27,28] Here, IL-2
activity was tied to proteolytic cleavage by fusing IL-2 with a receptor subunit via a matrix
metalloproteinase (MMP)-cleavable linker. Cytokine activity is inhibited ("caged") in the
absence of protease, and conditionally activated ("uncaged") by proteolytic cleavage. Using this
general framework, we demonstrated the possibility to cage structurally and functionally distinct
cytokines. Inspired by MSC and CAR T applications, we chose interleukin-10 (IL-10, an
immuno-suppressive cytokine that plays a role in limiting host immune responses as well as
promoting wound healing and muscle regeneration[29]), interleukin-6 (IL-6, generally associated
with inflammation, but also implicated with the activation of satellite cells and promotion of
myotube regeneration for muscle regeneration[30–32]), and IL-2 (a potent factor for the proliferation
and activation of lymphocytes, and widely studied for activating effector T cells against
tumors[33–35]).
We started by designing a caged IL-10. IL-10 is composed of a homodimer that binds to
a tetrameric IL-10 receptor complex formed by two IL-10Ra and two IL-10Rb subunits.[36] Since
effectively caging a dimeric protein may be difficult due to having multiple epitopes to interact
with receptors, we also tested a previously engineered monomeric variant of IL-10 (R5A11M)
with boosted affinity for IL-10Rb and potent immunomodulatory effects.[37] We created two
caged IL-10 constructs. Both designs contain a renin cleavable linker (cut site and glycine-serine,

Fig. 1: Engineering caged IL-10. a, Caged cytokine design for dimeric and monomeric IL-10. Cytokines are fused with renin-cleavable linkers to cognate receptor subunits, inhibiting binding to cellular receptors. **b,** hDIRECT circuit using caged cytokines to control cytokine activity. Cells constitutively secrete attenuated caged cytokines and cytokine activity can be tuned down via small-molecule inhibitor for co-transfected protease, renin. **c,** Functional activity of caged IL-10 constructs with or without renin co-expression. **d,** Functional activity of cageIL-10 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **e,** Functional activity of cageIL-10 co-expressed with renin at increasing aliskiren concentrations. (For **c-e**: Cytokine activity measured from supernatant using HEKBlue IL-10^{TM} reporter cells. For **c**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **d-e**: n = 2 biological replicates; data represent mean ± s.e.m. *** p ≤ 0.001, **** p < 0.0001.)

GS, linker) and the extracellular portion of specific IL-10R subunits for caging. The first
construct was created by fusing two WT IL-10 monomers together with IL-10Ra (CageIL-10-
dimer – Fig. 1a). The second construct used the monomeric R5A11M IL-10 and IL-10Rb
(CageIL-10-mono – Fig. 1a). We hypothesized that aliskiren could be used to control renin
activity and removal of the caged portion of hDIRECT to control cytokine activity (Fig. 1b).
Taking advantage of our fast design-build-test cycle, we transiently transfected human
embryonic kidney (HEK) 293 cells using plasmids encoding the caged cytokines and renin.
Supernatant was transferred to HEK-Blue^{TM} IL-10 reporter cells to assess functional activity.
Here, signal transduction from IL-10 receptor activation results in expression of secreted
embryonic alkaline phosphatase (SEAP), which can be assayed via absorbance following

incubation with a colorimetric substrate. A schematic for this cytokine reporter assay can be
found in Supplementary Fig. 1. Unless specified otherwise, the same workflow was used
throughout this work. In the absence of renin, cageIL-10-mono was caged more efficiently than
the cageIL-10-dimer (Fig. 1c). CageIL-10-mono was also able to fully reconstitute activity with
protease present (Fig. 1c). From here on, we refer to this optimal variant as "cageIL-10". Next, to
explore the extent of IL-10 caging and linker cleavability, we titrated the amount of transfected
cageIL-10 plasmid DNA. We observed efficient caging even at high doses of cageIL-10 (Fig.
1d), and renin-activated IL-10 activity at low doses of cageIL-10 (Fig. 1d). We then validated
that aliskiren enables dose-dependent control of hDIRECT (Fig. 1e).
To expand the cytokine output of hDIRECT, we next developed caged designs for IL-6.
Structurally, IL-6 is a monomeric four-α helical bundle typical of most cytokines (distinct from
the dimeric IL-10). Canonically, IL-6 binds to its high affinity receptor, IL-6Ra, on cell surfaces,
followed by complexing on the membrane with IL-6Rb (gp130) which transduces signal
activation across the membrane[38]. However, IL-6 activity has also been implicated in an
alternative pro-inflammatory pathway wherein IL-6 binds to soluble IL-6Ra (sIL-6Ra) and
mediates transactivation on cells expressing IL-6Rb[39]. To avoid transactivation of our caged IL-6
complex and bias IL-6 signaling to a regenerative phenotype, we sought to cage IL-6 with an IL-
6Ra mutant (A228D/N230D/H280S/D281V) that has abrogated activation of IL-6Rb. We denote
this IL-6Ra mutant as IL6Ra_M4.
Inspection of the IL-6-receptor complex structures suggested that placing IL-6Ra at the
N-terminus of the fusion protein would allow IL-6 to bind IL-6Ra in its canonical conformation.
Consistent with this observation, our data shows improved caging when the receptor is fused to
the N-terminus, although fusion at either terminus exhibits substantial renin-independent
activation (Supplementary Fig. 2). We hypothesized that caging could be improved if we fused
an additional receptor subunit. Therefore, we created a multi-cageIL-6 by fusing IL-6Ra_M4 to
the N-terminus and IL-6Rb to the C-terminus of IL-6 with renin-cleavable linkers (Fig. 2a).
Indeed, multi-cageIL-6 demonstrated vastly improved caging over the initial design (Fig. 2b),
albeit with reduced activity in the presence of renin (Fig. 2b). However, we found that IL-6
activity in response to renin can be increased at higher doses of multi-cageIL-6 plasmid (Fig. 2c).
Finally, to show broad generalizability in cytokines and therapeutic applications, we
ported over previous caged IL-2 constructs to hDIRECT[27]. A mutant IL-2 (sumIL-2) was
recently developed with enhanced affinity for IL-2Rb (i.e., effector T cells) over IL-2Ra (i.e.,
regulatory T cells)[28]. We leveraged this insight to design a caged sumIL-2 with a renin-cleavable
linker and C-terminal IL-2Rb (Fig. 2d). We also investigated whether varying the linker length
on either side of the renin cut site would improve cleavability but did not find any significant
effect (Supplementary Fig. 3). CageIL-2 is efficiently caged at high doses of cytokine (Fig. 2e)
and exhibits renin-activation at low doses (Fig. 2e). IL-2 activity was tuned sensitively by
aliskiren upon drug titration (Fig. 2f), indicating that aliskiren control is also generalizable
between different cytokines. Altogether, we established three different caged cytokines
compatible with renin-aliskiren control.

Fig. 2: Engineering caged IL-6 and IL-2. a, Caged cytokine design for single and multi-caged IL-6 constructs. **b,** Functional activity of caged IL-6 constructs with or without renin co-expression. **c,** Functional activity of cageIL-6 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **d,** Caged cytokine design for caged IL-2. **e,** Functional activity of cageIL-2 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **f,** Functional activity of cageIL-2 co-expressed with renin at increasing aliskiren concentrations. (Cytokine activity measured from supernatant using HEKBlue IL-6™ (**b-c**) and IL-2™ (**e-f**) reporter cells. For **b**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **c,e-f**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

Engineering cleavable cytokines.

To expand the potential of protease regulation, we sought to design cytokines whose
activities would be abrogated (rather than activated) by cleavage. Compared to the caged design,
this modality would allow cytokine activity to be "turned on" or restored by a small-molecule
input. It could be particularly useful in applications where temporary, rather than constitutive,
cytokine activity is desired, such as transient IL-2 activation to support CAR-T cell proliferation
and cytotoxicity while avoiding exhaustion and overstimulation[40].

We hypothesized that cytokines could be directly inactivated if they were engineered to
include protease cut sites. We inserted protease cut sites within flexible loops of cytokines so as
not to perturb receptor binding regions. Like the caged cytokines, we envision engineered cells
constitutively expressing cleavable cytokines and membrane-bound renin. In this fashion,
cytokine activity can be tuned up on demand upon drug input and protease inhibition (Fig. 3a).

We started designing cleavable cytokines using the previously mentioned mono-IL-10.
Researchers engineered this functional mono-IL-10 by inserting a 6 amino acid GGGSGG linker
between the D and E helices of human IL-10. We leveraged this insight to create a cleavable
mono-IL-10 by replacing this glycine linker with the 13 amino acid renin cleavage site.
CleaveIL-10 retained activity in the absence of protease and was abrogated upon co-transfection
with renin (Fig. 3b). To explore the range at which cleaveIL-10 is sensitive to renin cleavage, we
co-transfected cells with or without renin while titrating the amount of cleaveIL-10 DNA and
observed a substantial shift of the titration curves by renin (Fig. 3c).
Next, we sought to demonstrate the generalizability of this method to native cytokines
that lack synthetic linkers to substitute. Our general method was to find sites in human IL-6 and
IL-2 tolerable (retaining activity) to insertion of a protease cut site. We scanned various solvent
exposed (protease accessible) flexible loops between helices (Supplementary Fig. 4) and then
optimized linkers to improve their cleavability. Cleave IL-6 and IL-2 were designed through
direct insertion of the renin cut site with or without a five amino acid GS linker into flexible
loops at residues 127-140 in IL-6 and 74-81 in IL-2. Both cleaveIL-6 (Fig. 3d) and cleaveIL-2
(Fig. 3e) retained their functional activity in the absence of protease. Additionally, we observed
substantial shifts in response to renin in both cleaveIL-6 (Fig. 3d) and cleaveIL-2 (Fig. 3e).
To demonstrate that this cleavable method can be used to turn on cytokine activity by a
small-molecule input, we determined that cleaveIL-10 activity can be tuned via aliskiren (Fig.
3f). Taken together, these results demonstrate successful engineering of cleavable cytokine
constructs for three structurally distinct cytokines also compatible with renin-aliskiren control.

Fig. 3: Engineering of cleavable cytokines. a, hDIRECT circuit using cleavable cytokines to control cytokine activity. Cells constitutively secrete cleavable cytokines and cytokine activity can be tuned via small-molecule inhibitor for co-transfected protease, renin. **b,** Functional activity of cleaveIL-10 with or without renin co-expression. **c,** Functional activity of cleaveIL-10 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **d,** Functional activity of cleaveIL-6 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **e,** Functional activity of cleaveIL-2 plasmid transiently transfected with increasing amounts, with or without renin co-expression. **f,** Functional activity of cleaveIL-10 co-expressed with renin at increasing aliskiren concentrations. (Cytokine activity measured from supernatant using HEKBlue IL-10[TM] (**b-c,f**), IL-6[TM] (**d**), and IL-2[TM] (**e**) reporter cells. For **b**: mean of n = 3 biological replicates; Unpaired two-tailed Student's t-test. For **c-f**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

Orthogonalization of hDIRECT from its human counterparts.

204
After establishing renin-regulated secreted cytokines, our next goal was to orthogonalize
the system from human hosts. In contrast to parts sourced from other organisms (e.g., viral
proteases), utilizing human parts bears the distinct and critical risk of off-target effects due to
crosstalk with other human proteins. In particular, renin is endogenously secreted into the blood
to regulate blood pressure via proteolytic cleavage of angiotensinogen[25]. Therefore, to avoid
crosstalk with the host, there are two issues to address: the activation of angiotensinogen by cell-
surface, exogenous renin and the cleavage of engineered cytokines by endogenous renin.
To address the first problem, we took inspiration from our previous work and
hypothesized that we could spatially separate renin from circulation by localizing it to the
endoplasmic reticulum (ER) of the engineered cell[16,41]. We fused renin to the CD4

transmembrane domain and a cytosolic ER-retention motif (RXR). Here, renin regulates
engineered cytokines in cis as the cytokines move through the secretory pathway, while
remaining sequestered from renin's endogenous substrate in the blood (Fig. 4a). Using cageIL-10
as a test case, we validated that ER-retained renin maintained the ability to activate secreted
cageIL-10 (Fig. 4b). Next, to ensure that aliskiren could permeate engineered cells and inhibit
ER-renin with a similar sensitivity (as it typically acts extracellularly), we performed aliskiren
titration and validated that ER-renin retains aliskiren sensitivity (Fig. 4c).
To address the second problem of endogenous renin cleaving the engineered cytokines,
we took inspiration from renin's diversity of specificities across species. In particular, human and
mouse angiotensinogen are not recognized by the opposing renins[42]. We hypothesized that we
could use the mouse cut site in engineered cytokines to avoid their cleavage by circulating
human renin, and in parallel minimally engineer human renin to recognize the mouse site such
that the protease is still unlikely to be recognized as non-human. To introduce mouse substrate
recognition, we compared substrate bound renin structures across species. Compared to human
renin, mouse renin has a larger binding pocket to accommodate bulkier substrate residues (L35
and Y36). We hypothesized that by mutating four substrate-binding residues in human renin to
corresponding residues in mouse renin with smaller side chains (L147I, R148H, I203V, L290V)
we could expand the binding pocket to accommodate the mouse substrate. We denote this mutant
renin as human-to-mouse (H2M) renin. We then tested the orthogonality of the renin protease
and found that the human and mouse renins only recognize their corresponding cut sites while
the H2M renin's specificity is expanded to recognize both (Fig. 4d). We confirmed that H2M,
ER-localized renin remains sensitive to aliskerin (Fig. 4e) and that ER-localization does not
affect orthogonality (Fig. 4f).
In summary, human renin was successfully orthogonalized using structure-guided
mutation and localization motifs. Importantly, these orthogonalization strategies did not
compromise renin function or sensitivity to aliskiren. We expect to use ER-retained H2M renin
and caged and cleavable cytokines harboring the mouse cut site, with minimal crosstalk with
endogenous human renin and angiotensinogen.

Fig. 4: Orthogonalization of renin. a, hDIRECT circuit using caged cytokines and ER-retained renin to control cytokine activity. Cells constitutively secrete caged cytokines and cytokine activity can be tuned via small-molecule inhibition of a co-expressed protease, renin. ER-renin is retained in ER via cytosolic motif and can cleave secreted cytokines. **b,** Activity of cageIL-10 transiently transfected with membrane or ER-retained renin. **c,** Activity of cageIL-10 co-expressed with ER-renin at increasing aliskiren concentrations. **d,** Orthogonal renin matrix. Functional activity of cageIL-10 constructs containing human (h) or mouse (m) cut sites transiently transfected with membrane-bound human, mouse, or engineered H2M renin. **e,** Functional activity of cageIL-10 with an mRenin cut-site co-expressed with H2M ER-renin at increasing aliskiren concentrations. **f,** Orthogonal renin matrix. Functional activity of cageIL-10 constructs containing human (h) or mouse (m) cut sites transiently transfected with ER-retained human, mouse, or engineered H2M renin. (Cytokine activity measured from supernatant using HEKBlue IL-10^{TM} (**b-f**) reporter cells. For **b,d,f**: mean of n = 3 biological replicates; Individual replicate data for **d,f** can be found in Supplementary Fig. 5. Unpaired two-tailed Student's t-test. For **c,e**: n = 2 biological replicates; data represent mean ± s.e.m. **** p < 0.0001.)

Compact and stable transduction of hDIRECT in mammalian cells.

Clinically, *ex vivo* gene delivery to engineer therapeutic cells is bottlenecked by viral packaging limits of ~5 kb (adeno-associated virus) to ~10 kb (lentivirus)[43,44]. Compact genetic constructs are therefore a critical requirement for translational applications. To demonstrate that hDIRECT enables compact delivery, we created single-transcript constructs encoding engineered cytokines and renin separated by 2A "self-cleaving" peptides[45].

We created a polycistronic construct containing cageIL-10, membrane-bound renin, and a fluorescent co-transfection marker, mCherry (Fig. 5a). Transient transfection of this construct in HEK293 cells enables inhibition of IL-10 activity upon dosing with aliskiren (Fig. 5a). However, immune and regenerative responses are often mediated by the action of multiple cytokines with characteristic dosing and temporal profiles, rather than the activity of a single cytokine.[46,47] For example, muscle regeneration is mediated by an initial pro-inflammatory regime (i.e. IL-6), stimulating and recruiting the innate immune system, followed by an anti-inflammatory regime (i.e. IL-10) to indirectly promote the differentiation of muscle stem cells.[48] To demonstrate that hDIRECT can simultaneously control the activity of multiple cytokines, we created single-transcript constructs containing both cageIL-10 and cleaveIL-6 (Fig. 5b) to mimic a cytokine response for promoting muscle regeneration. In this manner, delivery with drug should allow an initial regime of IL-6 activity, followed by a switch to IL-10 activity when drug is removed. HEK293 cells transiently transfected with this single-transcript construct demonstrated bidirectional control of IL-10 and IL-6 activity with aliskiren (Fig. 5b).

Finally, we tested whether hDIRECT's performance is robust against different delivery methods or cellular contexts. First, we created lentiviral constructs using a weaker constitutive promoter, EF1-α, and the polycistronic cageIL-10 design (from Fig. 5a). We chose a weak promoter since stronger promoters during our pilot study exhibited leaky cytokine activity (data not shown). Stable transduction of HEK293 cells enabled regulation of IL-10 activity with aliskiren (Fig. 5c). Next, we sought to test hDIRECT in another cell line to validate its generalizability to different cell types. We chose Jurkat cells as an accessible proof of concept model for regulating IL-2 activity in the context of a CAR-T cell therapy. We designed lentiviral constructs using the EF1-α promoter and a polycistronic cageIL-2 design (Fig. 5d). Stably transduced Jurkats exhibited lower levels of secretion compared to HEK293s, but retained high sensitivity to aliskiren regulation of IL-2 activity (Fig. 5d). These results demonstrate that hDIRECT can be delivered as a stably expressed, single transcript in a commonly used viral vector, one step closer to the engineering of therapeutic cells.

Fig. 5: Compact delivery of hDIRECT in mammalian cells. a, Polycistronic design of co-expressed cageIL-10 and renin (top cartoon). Aliskerin-regulated activity of cageIL-10 in HEK293 cells transiently transfected with cageIL-10 single-transcript design. **b,** Polycistronic design of cageIL-10, cleaveIL-6, and renin (top). Aliskiren-regulated activity of cageIL-10 (left-axis) and cleaveIL-6 (right-axis) in HEK293 cells transiently transfected with cageIL-10/cleaveIL-6 single-transcript design. **c,** Aliskiren-regulated activity of cageIL-10 in HEK293 cells stably transduced with cageIL-10 single-transcript design (from **a**). **d,** Polycistronic design of co-expressed cageIL-2 and renin (top cartoon). Aliskiren-regulated activity of cageIL-2 at increasing aliskiren concentrations from Jurkat cells stably transduced with cageIL-2 single-transcript design. (Cytokine activity measured from supernatant using HEKBlue IL-10[TM] (**a-c**), IL-6[TM] (**b**), and IL-2[TM] (**d**) reporter cells. For **a-d**: n = 2 biological replicates; data represent mean ± s.e.m.)

Discussion

Here, we have presented a post-translational, humanized platform, hDIRECT, for drug-
mediated regulation of secreted cytokine activity in mammalian cells. Exclusively using human
proteins and domains, hDIRECT confers protease control with opposing signs ("caged" and
"cleavable") over cytokine activity. Co-expression of engineered cytokines renin allows the
control of cytokine activity using renin's FDA-approved inhibitor, aliskiren. We used species
specificity and re-localization strategies to orthogonalize both renin and engineered cytokines to
avoid crosstalk with their human counterparts. Lastly, we showed that hDIRECT is compact and
can regulate the activities of multiple cytokines encoded on a single transcript in different

mammalian cells. hDIRECT therefore represents a proof-of-concept towards humanized
engineering for post-translational tools to control next generation cell therapies.
While we have shown that hDIRECT can regulate cytokines, we expect that the
proteolytic control module can be extended to a variety of secreted and membrane proteins of
interest. Growth factors and chemokines, which regulate the proliferation, differentiation, and
migration of cells, could be similarly engineered for proteolytic regulation. In addition, we
envision that receptors with cleavable pro-domains or competitive caging domains might offer
small-molecule control of receptor availability which, for example, could tune the activity of
CAR-T cells to limit exhaustion[40].
We envision that hDIRECT will find immediate applications in cellular therapeutics.
hDIRECT could be used to temporarily supply CAR-T cells with IL-2 to improve proliferation
and cytotoxicity. In applications where the temporal profile of cytokine activity is critical, such
as MSC therapy for muscle regeneration, hDIRECT can be used to activate pro-inflammatory IL-
6 followed by regenerative, anti-inflammatory IL-10. While this work demonstrates the control
of two cytokines simultaneously, lentiviral packaging could allow for the delivery of a single
transcript encoding up to five engineered cytokines. hDIRECT could also be used for research in
basic science investigating the timing and dosage of various cytokine regimes by imparting
temporal control over the activity of multiple cytokines.
While our system is built from native human proteins, the presence of junctions from
fusion of domains and small number of mutations introduced for improving function and
orthogonalization may still be recognized as foreign. Although immunogenicity can be estimated
in silico using methods such as MHC binding algorithms, empirically validating these
predictions will require clinical studies on human patients[49]. Nonetheless, we anticipate that
human-derived engineering is a step towards immune-tolerized synthetic circuits.
While clinically suitable, hDIRECT still requires further work for potential use in
humans. Notably, most experiments here were conducted in the model mammalian cell line
HEK293. Future work in primary cell lines more aligned with an intended clinical application,
such as CAR-T cells, could demonstrate the functional efficacy of this platform more.
Subsequently, *in vivo* experiments in mice will help to determine the pharmacokinetics of small-
molecule dosing and localized cytokine activity in a more complex environment. Furthermore,
the use of aliskiren (approved as an antihypertensive) in a patient requiring a cellular therapeutic
might have possible side effects. To minimize side effects, renin could be further engineered to
be more sensitive to aliskiren so lower doses can be used.
In summary, hDIRECT is the first post-translational control platform using
orthogonalized humanized components in mammalian cells. Using hDIRECT, we demonstrate
small-molecule control of multiple secreted cytokines simultaneously. We envision that
hDIRECT is generalizable towards a wide variety of extracellular factors and receptors to tune
the. We expect our future engineering efforts to establish hDIRECT as a powerful tool routinely
used in cell therapies.
329

References

1. Fischbach, M. A., Bluestone, J. A. & Lim, W. A. Cell-based therapeutics: the next pillar of
medicine. *Sci. Transl. Med.* **5,** 179ps7 (2013).
2. Labanieh, L., Majzner, R. G. & Mackall, C. L. Programming CAR-T cells to kill cancer.
Nat. Biomed. Eng. **2,** 377–391 (2018).
3. Ninagawa, N. T. *et al.* Transplantated mesenchymal stem cells derived from embryonic stem

cells promote muscle regeneration and accelerate functional recovery of injured skeletal
muscle. *Biores. Open Access* **2,** 295–306 (2013).

4. Winkler, T. *et al.* Dose-response relationship of mesenchymal stem cell transplantation and
functional regeneration after severe skeletal muscle injury in rats. *Tissue Eng. Part A* **15,**
487–492 (2009).

5. Munir, H. & McGettrick, H. M. Mesenchymal stem cell therapy for autoimmune disease:
risks and rewards. *Stem Cells Dev.* **24,** 2091–2100 (2015).

6. Squillaro, T., Peluso, G. & Galderisi, U. Clinical trials with mesenchymal stem cells: an
update. *Cell Transplant.* **25,** 829–848 (2016).

7. Marofi, F. *et al.* CAR T cells in solid tumors: challenges and opportunities. *Stem Cell Res*
Ther **12,** 81 (2021).

8. Preda, M. B. *et al.* Short lifespan of syngeneic transplanted MSC is a consequence of in vivo
apoptosis and immune cell recruitment in mice. *Cell Death Dis.* **12,** 566 (2021).

9. Kim, N. & Cho, S.-G. New strategies for overcoming limitations of mesenchymal stem cell-
based immune modulation. *Int J Stem Cells* **8,** 54–68 (2015).

10. Kerkar, S. P. *et al.* Tumor-specific CD8+ T cells expressing interleukin-12 eradicate
established cancers in lymphodepleted hosts. *Cancer Res.* **70,** 6725–6734 (2010).

11. Nitahara-Kasahara, Y. *et al.* Enhanced cell survival and therapeutic benefits of IL-10-
expressing multipotent mesenchymal stromal cells for muscular dystrophy. *Stem Cell Res*
Ther **12,** 105 (2021).

12. Chen, Z. & Elowitz, M. B. Programmable protein circuit design. *Cell* **184,** 2284–2301
(2021).

13. Gao, X. J., Chong, L. S., Kim, M. S. & Elowitz, M. B. Programmable protein circuits in
living cells. *Science* **361,** 1252–1258 (2018).

14. Fink, T. *et al.* Design of fast proteolysis-based signaling and logic circuits in mammalian
cells. *Nat. Chem. Biol.* **15,** 115–122 (2019).

15. Sahin, U., Karikó, K. & Türeci, Ö. mRNA-based therapeutics--developing a new class of
drugs. *Nat. Rev. Drug Discov.* **13,** 759–780 (2014).

16. Vlahos, A. E. *et al.* Protease-controlled secretion and display of intercellular signals. *Nat.*
Commun. **13,** 912 (2022).

17. Praznik, A. *et al.* Regulation of protein secretion through chemical regulation of
endoplasmic reticulum retention signal cleavage. *Nat. Commun.* **13,** 1323 (2022).

18. Ferdosi, S. R. *et al.* Multifunctional CRISPR-Cas9 with engineered immunosilenced human
T cell epitopes. *Nat. Commun.* **10,** 1842 (2019).

19. Lamers, C. H. J. *et al.* Immune responses to transgene and retroviral vector in patients
treated with ex vivo-engineered T cells. *Blood* **117,** 72–82 (2011).

20. Riechmann, L., Clark, M., Waldmann, H. & Winter, G. Reshaping human antibodies for
therapy. *Nature* **332,** 323–327 (1988).

21. Jones, P. T., Dear, P. H., Foote, J., Neuberger, M. S. & Winter, G. Replacing the
complementarity-determining regions in a human antibody with those from a mouse. *Nature*
321, 522–525 (1986).

22. Li, H.-S. *et al.* Multidimensional control of therapeutic human cell function with synthetic
gene circuits. *Science* **378,** 1227–1234 (2022).

23. Rauch, S. *et al.* Programmable RNA-Guided RNA Effector Proteins Built from Human
Parts. *Cell* **178,** 122–134.e12 (2019).

24. Zhu, I. *et al.* Design and modular assembly of synthetic intramembrane proteolysis receptors

for custom gene regulation in therapeutic cells. *BioRxiv* (2021).
doi:10.1101/2021.05.21.445218

25. Danser, A. H. J. & Deinum, J. Renin, prorenin and the putative (pro)renin receptor.
Hypertension **46,** 1069–1076 (2005).

26. Vaidyanathan, S., Jarugula, V., Dieterich, H. A., Howard, D. & Dole, W. P. Clinical
pharmacokinetics and pharmacodynamics of aliskiren. *Clin Pharmacokinet* **47,** 515–531
(2008).

27. Puskas, J. *et al.* Development of an attenuated interleukin-2 fusion protein that can be
activated by tumour-expressed proteases. *Immunology* **133,** 206–220 (2011).

28. Hsu, E. J. *et al.* A cytokine receptor-masked IL2 prodrug selectively activates tumor-
infiltrating lymphocytes for potent antitumor therapy. *Nat. Commun.* **12,** 2768 (2021).

29. Wang, X., Wong, K., Ouyang, W. & Rutz, S. Targeting IL-10 Family Cytokines for the
Treatment of Human Diseases. *Cold Spring Harb. Perspect. Biol.* **11,** (2019).

30. Sarvas, J. L., Khaper, N. & Lees, S. J. The IL-6 Paradox: Context Dependent Interplay of
SOCS3 and AMPK. *J. Diabetes Metab.* **Suppl 13,** (2013).

31. Serrano, A. L., Baeza-Raja, B., Perdiguero, E., Jardí, M. & Muñoz-Cánoves, P. Interleukin-
6 is an essential regulator of satellite cell-mediated skeletal muscle hypertrophy. *Cell Metab.*
7, 33–44 (2008).

32. Zhang, C. *et al.* Interleukin-6/signal transducer and activator of transcription 3 (STAT3)
pathway is essential for macrophage infiltration and myoblast proliferation during muscle
regeneration. *J. Biol. Chem.* **288,** 1489–1499 (2013).

33. Liao, W., Lin, J.-X. & Leonard, W. J. Interleukin-2 at the crossroads of effector responses,
tolerance, and immunotherapy. *Immunity* **38,** 13–25 (2013).

34. Rosenberg, S. A. Interleukin 2 for patients with renal cancer. *Nat. Clin. Pract. Oncol.* **4,** 497
(2007).

35. Hawkins, E. R., D'Souza, R. R. & Klampatsa, A. Armored CAR T-Cells: The Next Chapter
in T-Cell Cancer Immunotherapy. *Biologics* **15,** 95–105 (2021).

36. Logsdon, N. J., Jones, B. C., Josephson, K., Cook, J. & Walter, M. R. Comparison of
interleukin-22 and interleukin-10 soluble receptor complexes. *J. Interferon Cytokine Res.*
22, 1099–1112 (2002).

37. Gorby, C. *et al.* Engineered IL-10 variants elicit potent immunomodulatory effects at low
ligand doses. *Sci. Signal.* **13,** (2020).

38. Garbers, C. *et al.* Plasticity and cross-talk of interleukin 6-type cytokines. *Cytokine Growth*
Factor Rev. **23,** 85–97 (2012).

39. Wolf, J., Rose-John, S. & Garbers, C. Interleukin-6 and its receptors: a highly regulated and
dynamic system. *Cytokine* **70,** 11–20 (2014).

40. Kwon, B. The two faces of IL-2: a key driver of CD8+ T-cell exhaustion. *Cell Mol Immunol*
18, 1641–1643 (2021).

41. Shikano, S. & Li, M. Membrane receptor trafficking: evidence of proximal and distal zones
conferred by two independent endoplasmic reticulum localization signals. *Proc. Natl. Acad.*
Sci. USA **100,** 5783–5788 (2003).

42. Fukamizu, A. *et al.* Dependence of angiotensin production in transgenic mice carrying either
the human renin or human angiotensinogen genes on species-specific kinetics of the renin-
angiotensin system. *Arzneimittelforschung.* **43,** 222–225 (1993).

43. Wu, Z., Yang, H. & Colosi, P. Effect of genome size on AAV vector packaging. *Mol. Ther.*
18, 80–86 (2010).

44. Kalidasan, V. *et al.* A guide in lentiviral vector production for hard-to-transfect cells, using
cardiac-derived c-kit expressing cells as a model system. *Sci. Rep.* **11,** 19265 (2021).
45. Tang, W. *et al.* Faithful expression of multiple proteins via 2A-peptide self-processing: a
versatile and reliable method for manipulating brain circuits. *J. Neurosci.* **29,** 8621–8629
(2009).
46. Subramanian, N., Torabi-Parizi, P., Gottschalk, R. A., Germain, R. N. & Dutta, B. Network
representations of immune system complexity. *Wiley Interdiscip Rev Syst Biol Med* **7,** 13–38
(2015).
47. Orzechowski, A. in *The plasticity of skeletal muscle* (ed. Sakuma, K.) 113–139 (Springer
Singapore, 2017). doi:10.1007/978-981-10-3292-9_5
48. Yang, W. & Hu, P. Skeletal muscle regeneration is modulated by inflammation. *J. Orthop.*
Translat. **13,** 25–32 (2018).
49. Calis, J. J. A. *et al.* Properties of MHC class I presented peptides that enhance
immunogenicity. *PLoS Comput. Biol.* **9,** e1003266 (2013).

Methods

Methods

Plasmid generation. All plasmids were constructed using general practices. Backbones were
linearized via restriction digestion, and inserts were generated using PCR, phosphorylated
annealing, or purchased from Twist Biosciences. A complete list of plasmids used for each
experiment (Supplementary Data 1) and the respective amounts used for transfection can be
found in Supplementary Data 2. In addition, the DNA sequences of all the plasmids used in this
study can be found in the source data, and all new plasmids with annotations will be available on
Addgene.

Tissue culture. WT Human Embryonic Kidney (HEK) 293 cells (ATCC, catalog no. CRL-1573)
and HEK293T-LentiX (Takara Biosciences, catalog no. 632180) were cultured in a humidity-
controlled incubator under standard culture conditions (37 °C with 5% CO_2) in Dulbecco's
modified Eagle's medium, supplemented with 10% fetal bovine serum (Fisher Scientific, catalog
no. FB12999102), 1 mM sodium pyruvate (EMD Millipore, catalog no. TMS-005-C), 1×
penicillin–streptomycin (Genesee, catalog no. 25-512), 2 mM l-glutamine (Genesee, catalog no.
25-509) and 1× MEM non-essential amino acids (Genesee, catalog no. 25-536). HEKBlue™ IL-
2, IL-6, and IL-10 cells were purchased from Invivogen (catalog no. hkb-il2, hkb-hil6, hkb-il10,
respectively). Cells were cultured as per manufacturer's protocol. T-REx™ Jurkat cells
(Invitrogen, catalog no. R72207) were cultured as per manufacturer's protocol.

Transient transfection. HEK293 cells were cultured in 96-well tissue culture-treated plates
under standard culture conditions. When cells were 70-90% confluent, the cells were transiently
transfected with plasmid constructs using the jetOPTIMUS DNA transfection reagent (Polyplus,
catalog no. 117-15), as per manufacturer's instructions using 0.375 μL of reagent per 50 μL of
jetOPTIMUS buffer for 500 ng total DNA transfections. Experiments requiring incubation with
aliskiren (Sigma Aldrich, catalog no. SML2077) were treated with drug at time of transfection
through media exchange.

Lentiviral transduction. LentiX cells were cultured in 6-well tissue culture-treated plates under
standard culture conditions. When cells were 70-90% confluent, the cells were transiently
transfected with plasmid constructs (600 ng PAX2, pMD2g, and 1,100 ng transfer target) using

the jetOPTIMUS DNA transfection reagent (Polyplus catalog no. 117-15), as per manufacturer's
instructions using 1.5 μL of reagent per 200 μL of jetOPTIMUS buffer for 2,000 ng total DNA
transfections. Cells were incubated for 24 h under standard culture conditions. Afterwards an
additional 3 mL of DMEM media was added carefully to each well. Lentivirus was concentrated
24 hrs afterward using viral precipitation. For each lentiviral prep, media was filtered using a
syringe and 0.45 μm filter into 15 mL conical tubes. 5x Lentivirus Precipitation Solution (Alstem
catalog no. VC100) was mixed with each prep and incubated at 4 °C overnight. Then, virus was
spun down at 1500xg for 30 min at 4 °C. Supernatant was aspirated, and virus was resuspended
using 200 μL Dulbecco's PBS (Genesee catalog no. 25-508). Virus was added dropwise onto 24-
well tissue culture plates containing HEK293 or Jurkat cells seeded at 250k cells/well. 72 h after
incubation, the percentage of mCherry-positive cells were quantified using flow cytometry. The
cells were selected using 1000 ng/mL of puromycin (ThermoFisher Scientific; catalog# J61278-
MB) for one week until >90% of cells were mCherry-positive before being used for experiments.
487
Cytokine reporter cell assay. Supernatant (10 μL) from HEK293 cells transiently transfected
24 h prior were incubated with 5 x 10^4 HEKBlue^{TM} cells (190 uL) in 96-well tissue culture-
treated plates for 24 h under standard culture conditions. Afterwards, 20 μL HEKBlue^{TM} cell
supernatant was collected and incubated with 180 μL Quanti-Blue reagent (Invivogen) for 2 h at
37 C as per manufacturer protocol. The mixture was read at 630 nm using the Spark® Multimode
microplate reader (TECAN). Cartoon schematic can be found in Supplementary Fig. 1.
494
For stable transductions, stable HEK293 cells were seeded at 1.5 x 10^4 cells in 96-well tissue
culture-treated plates for 48 h under standard culture conditions. Media was changed either with
or without aliskiren 24 h into the incubation. Supernatant transfer to HEKBlue^{TM} cells and assay
proceeded the same as prior. Jurkat cells were seeded at 5 x 10^4 cells in U-bottom 96-well plates
for 48 h under standard culture conditions. Cells were spun down in a centrifuge at 300 xg for 7
min and media was replaced with or without aliskiren 24 h into the incubation. 48 h after
seeding, cells were spun down in a centrifuge at 300 xg for 7 min again and supernatant was
transferred to HEKBlue^{TM} cells and assay proceeded the same as prior.
503
504 **Statistical analysis.** An unpaired *t*-test was used to determine the statistically significant
505 difference between two groups. Single, double, triple, and quadruple asterisks
506 indicate $P < 0.05$, $P < 0.005$, $P < 0.001$ and $P < 0.0001$. All statistical analyses were performed
507 using GraphPad Prism 10.0.0.
508

Data availability.
All data reported in this paper are available from the corresponding author on request. Plasmid
maps for all plasmids used in this study will be available as GenBank files. The plasmids used in
this study will be deposited on Addgene, with complete and annotated GenBank files on their
website. All figures and supplemental figures have associated raw data. There are no restrictions
on data availability.
515

Acknowledgements.
This research was supported by NIH (R00EB027723, DP2OD034951; X.J.G), Longevity
Impetus Grants (X.J.G.), Wu Tsai Human Performance Alliance Agility Project Grant (X.J.G.),
Stanford Bio-X Interdisciplinary Initiatives Seed Grant Program (IIP) [R11-7] (X.J.G.), the

Stanford Graduate Fellowship (C.A.), the Sarafan ChEM-H CBI training program (C.A.), and the
Bio-X Stanford Interdisciplinary Graduate Fellowship (C.A.).
522

Author contributions

524   C.A. and X.J.G. conceived and directed the study. C.A. performed all the experiments. A.E.V.
525   created the transmembrane domain and ER-retention motifs for renin constructs. J.P. and Q.C.
526   provided recommended mutations for the H2M renin. C.A. analyzed the data for the manuscript.
527   C.A. and X.J.G. wrote the manuscript. All authors provided feedback on the manuscript.
528

Competing interests

The board of trustees of the Leland Stanford Junior University have filed a patent on behalf of
the inventors (C.A.A. and X.J.G.) of the small-molecule control of membrane and secreted
proteins using human proteases platform described (US provisional Application No. 63/458833).
An additional provisional patent application has also been filed on behalf of the inventors
(C.A.A., X.J.G., J.P, and Q.C.) for the orthogonalization of human renin. X.J.G. is a co-founder
and serves on the scientific advisory board of Radar Tx.