Navigating contradictions: Salmonella Typhimurium chemotactic responses to conflicting effector

stimuli

Kailie Franco [1], Zealon Gentry-Lear [2], Michael Shavlik [3], Michael J. Harms [2,3], and Arden Baylink

[1*]

[1] Washington State University, Department of Veterinary Microbiology and Pathology, Pullman,

WA 99164

[2] University of Oregon, Institute of Molecular Biology, Eugene, OR, 97403

[3] University of Oregon, Department of Chemistry & Biochemistry, Eugene, OR, 97403

[*] Corresponding author: arden.baylink@wsu.edu (A. Baylink)

Summary

Chemotaxis controls swimming motility and colonization of many intestinal bacteria, but how

enteric pathogens navigate the complex chemical landscape of the gut, which contains

contradictory chemoattractant and chemorepellent stimuli, remains poorly understood. We find

Salmonella Typhimurium requires chemotactic sensing of two opposing signals present in the

intestinal lumen—the microbiota metabolite and bacteriostatic chemorepellent indole, and the

nutrient chemoattractant L-Ser—for efficient invasion of colonic tissue. Despite feces being the

major biological source of indole, accumulating to millimolar levels, non-typhoidal *Salmonella*

are strongly attracted to fecal material because chemoattraction to L-Ser and other attractants

override indole chemorepulsion. This behavior is orchestrated through the chemoreceptor Tsr,

which coordinates a spectrum of distinct rearrangements in the bacterial population structure based

on the ratio of L-Ser to indole. Through seeking niches with the highest L-Ser to indole ratio, *S.*

Typhimurium presumably optimizes nutrient access and avoids regions of high-competitor

density.

Keywords:

29 Chemotaxis, *Salmonella*, chemoeffector, chemohalation, serine, indole, chemoreceptor, Tsr,

30 microbiome, gastrointestinal pathogen

Introduction

Many bacteria that colonize the gastrointestinal tracts of humans and other animals employ chemotaxis to sense chemical effectors in the gut lumen and swim to environments conducive for growth and colonization [1–5]. Chemotaxis enables motile bacteria to seek nutrients, avoid toxins, and respond to molecular cues [1,2,4,5]. This process is controlled by chemoreceptor proteins, which recognize chemical effectors and transduce signals through a phosphorylation cascade to regulate flagellar rotation and swimming direction, ultimately determining the spatial and temporal patterns of bacterial colonization (Fig. 1A) [1,2,6,7]. While many effectors have been studied and characterized in isolation as attractants or repellents [4,7], natural environments contain mixtures of opposing signals. In the enteric lumen, bacteria encounter a complex milieu of conflicting effector signals, and little is known about how they prioritize these signals to direct their movement and colonization (Fig. 1B). Ultimately, the colonization topography of bacteria within the gut influences the health of the host through nutrient absorption, developmental regulation, and resistance to pathogens [8–10].

A chemical effector of major importance for enteric bacterial communities is indole, an interbacterial signaling molecule that regulates diverse aspects of bacterial physiology and lifestyle [11–13]. Indole is excreted by gut microbiota as a byproduct of tryptophan metabolism and accumulates to millimolar levels in human feces (Fig. 1A-B) [11,14,15]. Indole is amphipathic and can transit bacterial membranes to regulate biofilm formation and motility, suppress virulence programs, and exert bacteriostatic and bactericidal effects at high concentrations [11–13,15–17]. Indole was one of the earliest identified chemorepellents, and subsequent work has extensively explored its role in *Escherichia coli* chemotaxis (Table S1) [13,18–22]. The molecular mechanism by which *E. coli* senses indole remains unclear, but is known to involve the chemoreceptor taxis to serine and repellents (Tsr) (Fig. 1A-B) [18,20,21]. From this body of work, the hypothesis emerged that indole repels pathogens and restricts their growth as a mechanism of colonization resistance conferred by the microbiota [11–13,18,23]. However, no prior work has actually tested whether human fecal material, the major biological source of indole in the gut, which contains a complex mixture of effectors, induces chemorepulsion or inhibits pathogen growth at physiologically relevant levels.

We were interested in further studying the role of Tsr in navigating contradicting effector stimuli because this chemoreceptor has an interesting dual function: it both senses chemorepellents and also recognizes the amino acid and nutrient, L-Ser as a chemoattractant (Fig. 1A-B) [1,24–26].

Prior *in vitro* work with purified effectors showed that when both attractants and repellents are
present, Tsr facilitates "intermediate" responses between chemoattraction and chemorepulsion,
suggesting that, in some cases, navigating conflicting chemotactic stimuli is regulated at the single
chemoreceptor level [20,21]. The mechanisms and temporal dynamics of these intermediate
chemotactic behaviors are mostly uncharacterized, but could be relevant in natural settings that
contain conflicting effectors. We recently reported that many enteric pathogens and pathobionts
possess Tsr orthologues, including the genera *Salmonella*, *Citrobacter*, and *Enterobacter* [24].
Whether these and other bacteria respond to indole as a chemorepellent has remained unclear
because all prior studies of indole taxis focused on *E. coli* (Table S1).

Salmonella Typhimurium is a frank pathogen that relies on chemotaxis for enteric invasion
[27–30], and differs fundamentally from *E. coli* because it lacks tryptophanase genes and cannot itself
produce indole, which could provide a novel perspective on indole taxis [31,32]. In this study, we
used the pathogen *S.* Typhimurium as a model to: (1) test the hypothesis that the microbiota
secretion product indole is protective against enteric invasion, (2) investigate if pathogens are
repelled by indole-containing fecal material, and (3) examine how the chemoreceptor Tsr regulates
pathogen spatial localization in response to conflicting chemotactic stimuli of the intestinal
environ. Our study is the first to employ live imaging to directly visualize how enteric pathogen
populations dynamically restructure in response to physiological mixtures of attractants and
repellents. We demonstrate that the ability to navigate conflicting effector signals in fact mediates
efficient pathogen invasion, and that chemotaxis responses to natural biological combinations of
effectors and their impacts on infection outcomes, are not easily predicted based on responses to
individual effectors.

Results & Discussion

Opposing chemoeffector stimuli mediate efficient pathogen invasion in swine colonic explants

We sought to determine whether indole in human fecal matter protects against *S.*
Typhimurium infection and whether this involves chemotaxis mediated by the chemoreceptor Tsr.
To address this, we developed a swine colonic explant model that mimics the architecture and size
of adult human colonic tissue [33–37]. The explant tissue was gently cleaned and treated with various
effector solutions: solubilized human feces, purified indole and/or L-Ser at fecal-relevant

concentrations, or buffer as a control (see Method Details). The tissue was then exposed to co-
infection with wild-type (WT) *S.* Typhimurium strain IR715 and either a *cheY* mutant (motile but
non-responsive to chemoeffector stimuli) or a *tsr* deletion mutant (Fig. 1C-D, Key Resources
Table) [28]. To assess the role of chemotaxis in infection, we quantified total bacteria harvested from
tissue homogenates and enumerated intracellular bacteria using a gentamicin wash at 1-, 3-, and
6-h post-infection (Fig. 1C, Method Details) [38,39]. For buffer-treated explants, WT *S.*
Typhimurium showed a modest time-dependent advantage in colonization and invasion compared
to chemotactic mutants, indicating that chemotaxis, and specifically Tsr, promotes tissue
colonization under baseline conditions (Fig. 1E, Fig. S1A-B).

Contrary to the hypothesis that indole-rich fecal matter would inhibit pathogen
colonization, we found that fecal treatment significantly enhances intracellular invasion of WT *S.*
Typhimurium, providing a >100-fold competitive advantage mediated by Tsr (Fig. 1F, Fig. S1C-
D). Analysis of the liquified human fecal matter used in this study revealed an indole concentration
of 862 μM, consistent with previously reported ranges (0.5–5 mM) (Fig. 1D, Method Details) [11,14–
17]. However, when colonic tissue was treated with purified indole at the same concentration, the
competitive advantage of WT over the chemotactic mutants was abolished (Fig. 1G, Fig. S1E-F).
Given that Tsr mediates attraction to L-Ser in both *E. coli* and *S.* Typhimurium, we hypothesized
that L-Ser present in feces might be responsible for increased colonization of fecal-treated tissue
[1,24,40,41]. However, treatment with 338 μM L-Ser, the concentration present in our fecal sample
(Fig. 1D, Method Details), actually reduces the competitive advantage of WT bacteria, similar to
the effect of indole alone (Fig. 1H). While treatments with either indole or L-Ser alone reduce the
WT advantage, total bacterial loads are similar, suggesting that neither effector is "protective" in
terms of total infection burden (Fig. S1). We then wondered whether Tsr might require sensing of
both indole and L-Ser in combination to coordinate infection. Interestingly, we found that treatment
containing a mixture of indole and L-Ser restores the competitive advantage of WT (Fig. 1I, Fig.
S1I-J). Thus, we discovered that fecal material, the major biological source of indole in the gut,
does not protect against *S.* Typhimurium invasion, and that the fecal effectors indole and L-Ser, in
combination, direct efficient colonic invasion through chemotaxis and the chemoreceptor Tsr (Fig.
1, Fig. S1).

Fig. 1. Tsr and chemotaxis mediate efficient pathogen invasion of colonic tissue in the presence of conflicting chemotactic stimuli. A-B. Overview of the role of Tsr in chemotactic responses and premise of this study. C. Experimental design of colonic explant infections. D. Serine (presumed to be nearly 100% L-Ser, see Materials & Methods) and indole content of liquid human fecal treatments. E-I. Competitive indices (CI) of colony-forming units (CFU) recovered from co-infected swine explant tissue, either from the total homogenate (open box and whiskers plots), or the invaded intracellular population (checkered box and whisker plots), as indicated (n=5). Boxes show median values (line) and upper and lower quartiles, and whiskers show max and min values. Effect size (Cohen's *d*) and statistical significance are noted (not significant, ns; * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001, **** *p* < 0.0001). See also Figure S1.

Non-typhoidal S. enterica are attracted to human feces despite high indole content

Having found that Tsr mediates colonic invasion for *S.* Typhimurium, we next sought to learn what

chemotactic behaviors this chemoreceptor orchestrates in response to human fecal material; given

the high concentration of indole, we expected to observe chemorepulsion based on earlier studies

of Tsr function in *E. coli* (Fig. 1D, Table S1) [18–21]. We employed the chemosensory injection rig

assay (CIRA) for live-imaging of bacterial chemotaxis responses to a source of effectors injected

through a glass microcapillary [24]. In this assay, chemoattraction is observed as an influx of cells

toward the effector source and chemorepulsion as decreasing cells (Figure S2). As described

previously, effector injection introduces a steep microgradient, and using mathematical modeling

of the diffusion of the fecal sources of indole and L-Ser, we can approximate the local

concentrations experienced by bacteria at a given distance from the injection source, which for

most of the field of view is in the picomolar to low nanomolar range (Fig. 2A, Method Details) [24].

Over five minutes, we found both WT and *tsr* exhibit strong chemoattraction to fecal

material, whereas *cheY* remains randomly distributed (Fig. 2B, Movie S1). By examining the radial

distribution of the bacterial populations, we found WT more tightly centers around the treatment

source than *tsr* (Fig. 2C-E, Movie S1). In terms of the rate of accumulation of bacteria at the

treatment source, the chemoattraction of *tsr* lags behind the WT for the first 120 s (Fig. 2F-G,

Movie S1). We wondered how these deficiencies in fecal attraction might translate to direct

competition, where different strains are experiencing the same treatment source simultaneously.

To address this, we performed CIRA with solubilized human feces and two strains present in the

same pond, which we tracked independently through fluorescent markers (Fig. 3). As expected,

WT shows a strong chemoattraction response versus *cheY* (Fig. 3A, Movie S2). Interestingly, we

found that when competed directly, WT vastly outperforms *tsr*, with the maximal bacterial

distribution in proximity to the treatment source higher by about 4-fold (Fig. 3B, Movie S2).

Fig. 2. *Salmonella* Typhimurium exhibits attraction toward liquid human fecal material. A. Diffusion modeling showing calculated local concentrations in CIRA experiments with liquid human fecal material. B. Max projections of representative *S.* Typhimurium IR715 responses to a central source of liquid human fecal material. C-E. Mean bacterial distribution at 10 s intervals. F-G. Temporal analyses of area under the curve (AUC) or relative number of bacteria within 150 μm of the source. Effect size (Cohen's *d*) comparing responses of WT and *tsr* attraction at 120 s post-treatment is indicated. Data are means and error bars are standard error of the mean (SEM, n=3-5). See also Movie S1, Table S1 and Figure S2.

These data confirm that despite its high indole content, *S.* Typhimurium is attracted to human fecal

material through chemotaxis, and this response involves Tsr, although not as the sole mediator.

We expect the attraction of the *tsr* mutant is explained by the fact that *S.* Typhimurium possesses

other chemoreceptors that detect glucose, galactose, and L-Asp as chemoattractants, which are

present in human feces [1,7,42–45].

To understand the broader relevance of these behaviors to enteric infections, we examined

chemotaxis responses to fecal material among diverse *Salmonella* serovars and strains responsible

for human infections. Using dual-channel imaging, we compared *S.* Typhimurium IR715 with a

clinical isolate of *S.* Typhimurium, SARA1, and found both strains exhibit fecal attraction,

although SARA1 shows a slightly weaker response (Fig. 3C, Movie S3). We then tested a clinical

isolate of *S.* Newport, an emerging cause of salmonellosis in the United States and Europe [46,47].

This strain is strongly attraction to fecal material, with a tighter accumulation of cells at the

treatment source than *S*. Typhimurium IR715 (Fig. 3D, Movie S3). Lastly, we examined a clinical

isolate of *S*. Enteritidis, a zoonotic pathogen commonly transmitted from poultry, which displays

weak attraction to fecal material (Fig. 3E, Movie S3) [47]. Overall, we found that chemoattraction to

fecal material is conserved among non-typhoidal *Salmonella* serovars responsible for human

infections, although the degree of attraction varies. Notably, despite the high indole content in

feces, none of the strains tested exhibit chemorepulsion.

Mediation of opposing chemotactic responses by Tsr

We wondered if our inability to observe repulsion from fecal material might be due to *S*.

Typhimurium not sensing indole as a chemorepellent since this chemotactic response has only

been previously described for *E. coli* (Table S1). We again employed CIRA to address this

question, comparing chemotaxis responses to either 5 mM L-Ser or 5 mM indole, and found that

S. Typhimurium responded rapidly to these two effectors as chemoattractants and chemorepellents,

respectively (Fig. S2H-I). Treatment with 5 mM indole, a concentration at the upper end of what

occurs in the human gut [17], induces rapid chemorepulsion with the bacteria vacating the region

proximal to the source (Fig. S2B). Interestingly, the chemorepulsion response occurs faster than

chemoattraction, with a zone of avoidance clearly visible within the first 10 s of indole exposure

(Fig. S2I, Movie S4).

We next wondered if perhaps our fecal treatments contained insufficient indole to elicit

chemorepulsion from *S*. Typhimurium. To identify the effective source concentrations that drive

indole chemorepulsion and understand the temporal dynamics of this response, we performed a

titration of indole across 0.05-10 mM (Fig. 3F). At all concentrations tested, indole induces

chemorepulsion, and the bacteria avoid the treatment source for the duration of the 5-minute

experiment (Fig. 3F-G). At source concentrations exceeding 3 mM, essentially all motile cells

vacate the field of view within 60 s (Fig. 3F-G). Integrating these chemorepulsion responses and

fitting them to a Monad curve suggests an indole source concentration of approximately 67 µM is

sufficient for half-maximal ($K_{1/2}$) chemorepulsion (Fig. 3H). These data show that even though we

observed a strong chemoattraction response to fecal material, indole at the concentration present

in fecal material, and far lower, is indeed a strong chemorepellent for *S*. Typhimurium.

Based on its function in *E. coli*, we hypothesized that both indole chemorepulsion and L-

Ser chemoattraction for *S*. Typhimurium could be partly or fully mediated through Tsr [7,18,26].

Fig. 3. Fecal indole is insufficient for chemorepulsion but indole in isolation is a strong chemorepellent. A-E. Dual-channel imaging of chemotactic responses to solubilized human feces by WT *S.* Typhimurium IR715 (pink) and isogenic mutants or clinical isolate strains, as indicated. Shown are max projections at time 295-300 s post-treatment. Data are means and error bars are standard error of the mean (SEM, n=3-5). See also Movies S2-S3. F. Representative max projections of responses at 295-300 s of indole treatment. G-H. Quantification of chemorepulsion as a function of indole concentration (n=3-5). I-K. Comparison of WT and *tsr* mutant responses to L-Ser or indole. See also Fig. S2. L-M. Isothermal titration calorimetry (ITC) experiments with 50 μM *S.* Typhimurium Tsr ligand-binding domain (LBD) and indole, or with L-Ser in the presence of 500 μM indole. Data are means and error bars are standard error of the mean (SEM, n=3-5). AUC indicates area under the curve. Scale bars are 100 μm. See also Movies S2-S3.

We compared the chemotactic responses of the WT and *tsr* strains when exposed to sources of
these effectors, and found Tsr to be required for both chemorepulsion from indole and
chemoattraction to L-Ser (Fig. 3I-K). The canonical mode of chemoreceptor effector recognition
involves binding of the effector to the ligand-binding domain (LBD) [7,48], but the mechanism by
which indole is sensed through Tsr has not been elucidated. We recently reported the first crystal
structure of *S.* Typhimurium Tsr LBD, which clearly defines how the binding site recognizes the
L-Ser ligand (PDB code: 8fyv), and we thought it unlikely indole can be accommodated at the
same site [24]. Nevertheless, to test whether the Tsr LBD binds indole directly, we expressed and
purified the LBD, corresponding to the soluble periplasmic portion, and performed isothermal
titration calorimetry (ITC). These data show that no binding occurs between the Tsr LBD and
indole (Fig. 3L). We next wondered if indole acts as an allosteric regulator, possibly through
interacting with the L-Ser bound form or interfering with L-Ser recognition. To address these
possibilities, we performed ITC of 50 μM Tsr LBD with L-Ser in the presence of 500 μM indole
and observed a robust exothermic binding curve and K_D of 5 μM, identical to the binding of L-Ser
alone (Fig. 3M) [24]. These data indicate that indole does not alter the Tsr LBD affinity for L-Ser.

We conclude that Tsr senses indole through an atypical mechanism, which might either
involve regulation through a solute-binding protein [18,49], responsiveness to perturbation in the
proton motor force [13], or binding to a different region other than the periplasmic LBD. Our findings
reveal that while indole acts as a chemorepellent for *S.* Typhimurium in isolation, sensed through
Tsr, its presence within fecal material mixed with other effectors is insufficient to elicit
chemorepulsion.

Compromising between conflicting effector signals through chemohalation
Earlier work with *E. coli* revealed that exposure to mixtures of L-Ser and indole can generate

intermediate chemotactic responses between chemoattraction and chemorepulsion after prolonged exposure (1-6 h) (Table S1) [20]. Having confirmed that Tsr in *S.* Typhimurium mediates opposing chemotactic responses to the chemoattractant L-Ser and the chemorepellent indole in isolation, we next sought to learn how the bacterial population behaves when confronted with physiological combinations of these effectors. To address this, we performed a series of CIRA experiments with 500 µM L-Ser and increasing concentrations of indole at L-Ser:indole molar ratios of 10:1, 1:1, or 1:10 (Fig. 4A-D, Movie S4).

These experiments reveal a fascinating transition in the distribution of the pathogen population as a function of increasing chemorepellent, which occurs within minutes of exposure (Fig. 4A-D, Movie S4). With only chemoattractant present, the bacterial population organizes tightly around the effector source (Fig. 4A, Movie S4). When indole is introduced at a concentration 10-fold lower than L-Ser, the bacterial distribution still exhibits chemoattraction but becomes more diffuse (Fig. 4B, Movie S4). At a 1:1 ratio of chemoattractant and chemorepellent, a novel population structure emerges in which the swimming bacteria are attracted toward the source but form a halo around the treatment with an interior region of avoidance (Fig. 4C, Fig. 4E, Movie S4). When the concentration of indole is 10-fold higher than L-Ser, the bacteria exhibit a wider zone of avoidance (Fig. 4D-E, Movie S4). Interestingly, whereas 5 mM indole on its own induces strong chemorepulsion (Fig. S2I, Movie S4), the addition of 10-fold lower L-Ser effectively converts the behavior to a null response (Fig. 4D-E, Movie S4). This demonstrates that even at the highest concentrations of indole *S.* Typhimurium might encounter in the gut, the presence of chemoattractant can override indole chemorepulsion.

The intermediate responses to opposing effector mixtures bear similarities to CIRA experiments with fecal material, some of which also exhibited a halo-like structure around the treatment source (Fig. 2, Movies S2-S3). To our knowledge, there exists no consensus term for intermediate chemotaxis responses of this nature, so here we introduce "chemohalation," in reference to the halo formed by the cell population, and which is congruent with the established nomenclature of chemoattraction and chemorepulsion. We expect chemohalation is a compromise in positional location at the population level between the chemoattraction driven by L-Ser and the chemorepulsion driven by indole. Across these experiments, the interior zone of avoidance roughly corresponds to where the local concentration of indole exceeds 10 nM (Fig. 4E-F). In a biological setting, we presume that the distribution bias orchestrated by chemohalation regulates the

probability that cells will colonize, adhere, and transition to sessility at a given site; the greater the

local indole content, the wider the zone of avoidance and the less likely tissue invasion occurs.

We questioned why non-typhoidal *Salmonella* are attracted to a biological solution with

high concentrations of indole, a chemical reported to inhibit bacterial growth [12,50,51]. We examined

how bacterial growth is affected by 0-25 mM indole or L-Ser in a background of minimal media

(MM). As expected, increasing amounts of the nutrient L-Ser provide a growth advantage for all

Salmonella strains analyzed, with maximal benefit achieved by approximately 500 µM (Fig. 4G).

Equivalent treatments with indole show tolerance up to approximately 1 mM, with growth

inhibition occurring in the 1-5 mM range and lethality occurring at indole concentrations greater

than 5 mM (Fig. 4H). However, adding L-Ser in a background of 500 µM indole provides only a

small growth enhancement (Fig. 4I), and addition of 500 µM L-Ser increases tolerance for indole

up to about 1 mM, above which indole toxicity is unavoidable (Fig. 4J). So, we conclude that

mixtures of these effectors also impact growth differently than the effectors in isolation, and the

relative attraction to combinations of these effectors relates to their propensity to enhance or inhibit

growth.

Conclusions

Bacteria in the human gastrointestinal tract encounter complex chemical landscapes that contain

both chemoattractants and chemorepellents. However, chemotaxis responses are often studied in

isolation, outside of their biological and ecological contexts, which can lead to an over- or

underestimation of the roles specific interactions play in natural settings. In the present work, we

contribute to an emerging understanding that bacteria exhibit rapid and well-orchestrated

responses to conflicting stimuli distinct from chemoattraction or chemorepulsion and relate these

chemotactic compromises to enteric infection and pathogen growth (Fig. S3) [20,52]. Previously, no

study had addressed whether bacteria other than *E. coli* sense indole as a chemorepellent. In the

model system we investigated, we confirm that *S.* Typhimurium utilizes the chemoreceptor Tsr to

respond to indole as a chemorepellent and L-Ser as a chemoattractant (Fig. 3). Further, we show

that physiological mixtures of these effectors induce the behavior we define here as chemohalation,

where the bacteria accumulate at a distance from the treatment source and form a halo with an

interior zone of avoidance (Fig. 3, Fig. 4, Fig. S2, Movie S4). Our study is the first to capture real-

time videos of this phenomenon for an enteric pathogen and visualize how the population structure

Fig. 4. *S.* Typhimurium mediates distinct chemotactic responses based on the ratio of L-Ser to indole. A-D. Representative max projections of responses to treatments of L-Ser and indole at 295-300 s, as indicated. E. Relative bacterial distribution in response to treatments of 500 μM L-Ser and varying amounts of indole, from panels A-D, with the mean value normalized to 100%. F. Diffusion modeling of local effector concentrations based on sources of 5 mM indole (dark brown), 500 μM L-Ser (blue), 500 μM indole (light brown), and 50 μM indole (yellow) are shown as dashed lines. The approximate local concentration of indole that elicits a transition in chemotactic behavior is highlighted in light blue. Data are means and error bars are standard error of the mean (SEM, n=3-5). Scale bars are 100 μm. G-H. Bacterial growth as a function of L-Ser or indole, at the time point where the untreated culture reaches A_{600} of 0.5. I-J. Bacterial growth +/- pretreatment with 500 μM indole or L-Ser, and increasing concentrations of indole or L-Ser, as indicated at the time point where the untreated culture reaches A_{600} of 0.5. Data are means and error bars are standard error of the mean (SEM, n=8-24). See also Figure S3.

changes based on the ratio of attractant to repellent, ranging from chemoattraction, diffuse
chemoattraction, chemohalation, diffuse chemohalation, and chemorepulsion (Fig. 4, Fig. S2, Fig.
S3, Movie S4). These dynamic micron-scale population structures would be difficult or impossible
to detect and quantify without directly viewing them through live imaging.

We predicted that human fecal material, rich in indole, would elicit chemorepulsion, inhibit
pathogen growth, and protect against infection. Instead, we found that chemotactic sensing of
human fecal material promotes colonic invasion, predominantly eliciting chemoattraction or
chemohalation (Fig. 1, Fig. S1, Fig. 2, Movies S1-S3). We also found that chemotactic sensing of
opposing effectors mediates efficient colonic invasion but provides no advantage when only a
single effector is present (Fig. 1, Fig. S1). As evidenced by the phenomenon of chemohalation, the
bacteria bias their spatial location based on the ratio of chemoattractant to chemorepellent, and we
expect this behavior functions to rank colonization niches and regulate the probability of invading
specific sites. In the context of *S.* Typhimurium infection, we propose that Tsr orchestrates a
compromise between seeking niches rich in nutrients, signaled by local L-Ser concentrations, and
avoiding niches with high microbial competition, indicated by local indole concentrations (Fig.
S3). Chemorepulsion from indole can be overridden by the presence of chemoattractants, and *S.*
Typhimurium growth is quite tolerant of indole within physiological ranges, suggesting the
bacteria generally prioritize nutrient acquisition over the inhibitory effects of indole (Fig. 4). Since
the enteric lumen may never be devoid of attractant stimuli, it is possible that outright
chemorepulsion from a source of indole may not actually occur *in vivo*.

Having characterized and confirmed the dual sensing role of Tsr for *S.* Typhimurium, we
speculate that the diverse bacterial species that possess Tsr orthologues, particularly common
among *Enterobacteriaceae* [24], are capable of similar chemohalation behaviors and regulating taxis
based on local indole content, further supporting indole as a key regulator of polymicrobial
communities of the gut [12,16]. Recently, we reported on *Enterobacteriaceae* chemotactic sensing of
blood serum, another complex biological effector source at the host-pathogen interface, and those
responses appear to involve chemohalation [24]. Evidence of chemohalation is also seen in the case
of the gastric pathogen *Helicobacter pylori* responding to mixtures of urea, a chemoattractant, and
acid, a chemorepellent [53,54]. Continuing to investigate chemohalation behaviors and understanding
how they coordinate bacterial colonization may provide important insights into how chemotaxis
confers fitness advantages in natural environments.

STAR Methods

RESOURCE AVAILABILITY

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial Strains		
S. enterica Typhimurium IR715 nalidixic acid derivative of ATCC 14028	Rivera-Chávez, F. et al.[28]	N/A
S. enterica Typhimurium IR715 Δ*cheY::Tn10* (Tet^R)	Rivera-Chávez, F. et al.[28]	N/A
S. enterica Typhimurium IR715 Δ*tsr::pFR3* (Cm^R)	Rivera-Chávez, F. et al.[28]	N/A
S. enterica SARA1	Beltran, P. et al.[55]	N/A
S. enterica Newport	Shariat, N. et al.[46]	M11018046001A
S. enterica Enteriditis	Shariat, N. et al.[46]	05E01375
E. coli BL21-DE3	Millipore-Sigma	Cat# 70954-3
Biological Samples		
Single Human Donor Fecal Sample	Lee BioSolutions	Cat# 991-18
Chemicals, Peptides, and Recombinant Proteins		
L-Serine	Fisher	Cat# 56-45-1
Indole	Sigma	Cat# I3408-100G
Hydroxylamine hydrochloride	Sigma	Cat# 159417-100G
Recombinant DNA		
pXS-sfGFP	Glenn et al.[24]	N/A
pXS-mPlum	Glenn et al.[24]	N/A
pET-30a(+)-SeTsrLBD	Glenn et al.[24]	N/A
Software and Algorithms		
Matlab R2022a	The MathWorks Inc., Natick, Massachusetts, USA	Mathworks.com
Fiji	ImageJ, Bethesda, Maryland, USA[56]	Fiji Home
TrackingGUI_rp	R. Parthasarathy [57]	TrackingGUI Public

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Arden Baylink (arden.baylink@wsu.edu).

Materials availability

Strains and plasmids generated in this study will be made available upon request by the Lead

Contact with a completed Materials Transfer Agreement.

 Data availability

Source data from this work are archived and available upon request by the Lead Contact. This

paper does not report original code. Any additional information required to reanalyze the data

reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY DETAILS

All methods were carried out in accordance with relevant guidelines, regulations, and state and

federal law. Experimental protocols were approved by the Institutional Biosafety Committee (IBC)

of Washington State University (#1372).

Bacterial strains and growth conditions

Bacterial strains and plasmids used in this study are listed in Table 1. As previously described [24],

bacteria intended for chemotaxis assays were grown overnight in tryptone broth (TB) with

antibiotic selection, as appropriate. Motile bacteria were prepared with a 1:1000 back-dilution and

grown shaking for approximately 4 hours at 37° C to reach A_{600} of 0.5. Cells were centrifuged,

washed, and resuspended in a chemotaxis buffer (CB) containing 10 mM potassium phosphate

(pH 7), 10 mM sodium lactate, and 100 µM EDTA to A_{600} of 0.2 and rocked gently at room

temperature until fully motile. For *in vitro* growth analyses, cultures were grown overnight in

Lysogeny Broth (LB) at 37° C. Subsequently, 5 µl of A_{600} 2.0 cells were used to inoculate 200 µl

of minimal media (MM), containing 47 mM Na_2HPO_4, 22 mM KH_2PO_4, 8 mM NaCl, 2mM

$MgSO_4$, 0.4% glucose (w/v) 11.35 mM $(NH_4)_2SO_4$, 100 µM $CaCl_2$ and L-Ser and/or indole at the

described concentrations, and cultured in a 96-well microtiter plate. Cultures were grown at 37° C

and monitored by A_{600} readings at 5-minute intervals.

METHOD DETAILS

Chemosensory injection rig assay (CIRA)

CIRA was performed as described previously [24]. Briefly, an Eppendorf Femtotip 2 microcapillary

containing the treatment of interest was lowered into a pond of 50 µl of motile cells using a Sutter

micromanipulator. An injection flow of effector into the pond at approximately 300 fl per minute

was achieved using a Femtojet 4i set to P_c 35. Solubilized fecal treatments were prepared by

dissolving 1 g of commercially obtained human feces (Innovative Research) in 10 ml of CB. The

solution was clarified by centrifugation at 10,000 g for 20 minutes, followed by sterile filtration

through a 0.2 μm filter. Treatment solutions of indole and L-Ser were also diluted into CB and

sterile-filtered before application. Videos were recorded using an inverted Nikon Ti2 microscope

with heated sample chamber at 37 ˚C.

CIRA microgradient modeling

Modeling the microgradient generated through CIRA was performed as described earlier [24], based

on the continual injection and diffusion of an effector from a fixed-point source. Briefly, diffusion

is modeled as a 3D process where the diffusing substance is gradually and continuously introduced

at a fixed point within a large surrounding fluid volume. The substance is prepared at a

concentration of M_s (typically between 0.5 μM and 5 mM) and injected at a volume rate of $Q =$

305.5 fl/min. The species then diffuses into the ambient fluid with a diffusion constant D.

368
$$C(r,t) = \frac{q}{4\pi Dr} erfc \frac{r}{2\sqrt{Dt}}$$

Here, r is the distance from the point source, t is the time from initial injections, q is the injection

rate of the species (equal to M_sQ), and C is the species concentration.

Purification of recombinant S. Typhimurium Tsr LBD

Purification of *S.* Typhimurium Tsr LBD was performed as described previously [24]. Rosetta

BL21(DE3) *E. coli* cells with a Tsr-LBD-pet-30a(+) vector were grown with LB and 20 μg

kanamycin and induced at A_{600} of 0.8 with 0.4 mM isopropyl β-D-1-thiogalactopyranoside (IPTG).

Cells were harvested after 3 h of growth at 37˚ C. Cells were lysed, the lysate was clarified through

centrifugation, and the soluble fraction was subjected to an ammonium sulfate precipitation, with

Tsr LBD precipitating in the 20-40% fraction. The fractions were pooled, treated with TEV

protease to remove the N-terminal expression sequence, and purified using an anion exchange

column and Akta FPLC. Lastly, the protein was purified by gel filtration using an S200 column

with a final buffer of 50 mM Tris pH 7.5, 1 mM EDTA, and 150 mM NaCl and stored at 7 mg/ml

at -80 C.

Isothermal titration calorimetry ligand binding studies (ITC)

ITC experiments were performed using a Microcal ITC200 instrument (GE Healthcare). Either

500 μM indole or L-Ser was titrated in 2.5 μL injections into a 200 μL sample cell containing 50

μM Tsr LBD. For the indole/L-Ser competition experiment, 500 μM indole was added to both the

titrant and sample cell, thus providing a constant excess background concentration of indole. For

all experimental conditions, blank titrations were also collected in which indole or L-Ser was

titrated into a cell containing buffer alone. All experiments were performed using thoroughly

degassed samples at 25 °C in 50 mM Tris, 150 mM NaCl, 1 mM EDTA, pH 7.5. The reference

power was set to 5 μcal/sec. The resulting power curves were integrated using the Origin analysis

software included with the instrument. The heat of dilution was subtracted from each point using

the blank. A single-site binding model was then fit to the data, floating parameters describing the

binding enthalpy (ΔH), equilibrium constant (K_D), and apparent binding stoichiometry (n). The

instrument software was used for this purpose.

Quantification of indole and serine in human fecal samples

Solubilized human feces was prepared as described above for CIRA and analyzed by mass

spectrometry to determine the molar serine content as a service through the University of

Washington Mass Spectrometry Center. This measurement reflects total serine, of which close to

100% is expected to be L-Ser [24]. As described in earlier work, the indole content of solubilized

human fecal samples was determined using a hydroxylamine-based calorimetric assay with

purified indole as a reference and standard [58].

Explant infection assays

Swine intestinal tissue was acquired from the descending colon of an 8-week-old animal, pursuant

to animal protocol ASAF #7128, approved through the Washington State University IACUC.

Before infection, the luminal side of an approximately 20 by 20 mm piece of swine intestinal

explant tissue was gently washed with PBS to remove fecal matter. Next, the tissue section was

bathed in 2 ml of chemoeffector solution (solubilized human fecal matter, a mixture of 338 μM L-

Ser and 862 μM indole, 338 μM L-Ser alone, or 862 μM indole alone) in a 6-well tissue culture

plate (Celltreat) and incubated at 4° C for 1 h. Then, tissue was transferred to a 35 mm Mattek dish

where the luminal side of the tissue was exposed to a bacterial solution containing a 1:1 mixture

(10^9 CFU each) of WT *S.* Typhimurium IR715 and either the isogenic *tsr* or *cheY* mutant,

suspended in CB at a volume of 300 µl. The tissue was then incubated in the dish with the

competing bacteria at 37 °C and 5% CO_2 for 1-, 3-, or 6-h. After, half of the tissue was transferred

into screwcap tubes containing 500 µl LB media and 5-10 2.3 mm zirconia beads (BioSpec

Products) on ice and homogenized using a Bead Mill 24 (Fisher Scientific, 6.5 m/s for 60 s, four

times). To enumerate the intracellular bacteria, the other half of the tissue was washed in PBS and

incubated in PBS containing 100 g/ml gentamicin for 1 h at 37 °C and 5% CO_2, then washed twice

in PBS, as done previously [39,59,60]. The homogenization process was then repeated for the

gentamicin-treated tissue. CFUs were enumerated by 100 µl spot plating of 10-fold dilutions on

LB agar plates containing the appropriate antibiotic [39,61]. Competitive index values were calculated

by dividing the number of mutant CFUs by the number of WT CFUs for each treatment and time

point [62,63].

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantification of CIRA data

Videos of chemotactic responses were quantified as described previously [24]. The number of cells

in each frame was calculated by determining a fluorescence intensity ratio per cell for frames pre-

treatment and extrapolated using the 'plot profile' function of ImageJ. The distribution of the

bacteria was calculated using the Radial Profile ImageJ plugin. Local background subtraction was

performed based on experiments with the non-chemotactic *cheY* strain to control for

autofluorescence in solubilized fecal samples.

Statistical Analyses

Competitive indices (CIs) for explant experiments were calculated for each treatment group at

each time point. Log-transformed CI values were obtained by taking the logarithm ($\log_{10}$) of the

original CI measurements. These log-transformed values were then subjected to statistical analysis.

First, a one-sample t-test was performed to determine whether the mean of the log-transformed

CIs significantly differed from zero. In cases where the assumption of normality was violated, the

non-parametric Wilcoxon rank sum test was applied as an alternative. Effect size was assessed

using Cohen's *d and* calculated using the same log-transformed CIs.

The formula for Cohen's *d* value is as follows:

450
$$d = \frac{M_1 - M_2}{\sigma_{\text{pooled}}}$$

451   Where M_1 serves as the mean of the treatment group, M_2 serves as the mean of the control group,

452   and σ_{pooled} is the pooled standard deviation:

454
$$\sigma_{pooled} = \sqrt{\frac{\sigma_1{}^2 + \sigma_2{}^2}{2}}$$

Here, σ_1 is the standard deviation of the treatment group, and σ_2 is the standard deviation of the

control group.

Acknowledgments

Funding for this research was provided by NIAID through awards 1K99AI148587 and

4R00AI148587-03, and funding from Washington State University to AB. Bacterial strains were

provided by Nikki Shariat (University of Georgia, Athens), Nkuchia Mikanatha and Pennsylvania

NARMS and GenomeTrakr Programs, and Andreas Bäumler (University of California, Davis).

All research on human samples was performed in accordance with, and approval of, the

Institutional Biosafety Committee and Institutional Animal Care and Use Committee at

Washington State University.

Author Contributions

K.F. performed the microgradient modeling and explant experiments. A.B. conducted the CIRA

experiments and purification of Tsr-LBD. Z.G. performed bacterial growth experiments. M.S. and

M.J.H. performed the ITC experiments. All authors contributed to data analyses and writing of the

manuscript.

Declaration of Interests

A.B. owns Amethyst Antimicrobials, LLC.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT in order to proof-read and enhance the clarity and organization of the text. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Supplemental Information

Document S1. Table S1 and Figures S1–S3.

Movie S1. Chemotactic response of *S.* Typhimurium IR715 to solubilized human feces, related to Figure 2. Representative CIRA experiments showing *S.* Typhimurium IR715 WT and mutant strains responding to a source over 300 s (shown at 10x speed) Viewable at: https://www.youtube.com/watch?v=BqUcRN3YwjU

Movie S2. Chemotactic response of *S.* Typhimurium IR715 WT and chemotactic mutant strains to solubilized human feces, related to Figure 3. Representative CIRA experiments showing competition between *S.* Typhimurium IR715 (mPlum) and *cheY*, or *tsr*, as indicated (GFP), over 300 s. Viewable at: https://www.youtube.com/watch?v=D5JL46b4lsI

Movie S3. Chemotactic response of *S. enterica* clinical isolates to solubilized human feces, related to Figure 3. Representative CIRA experiments showing competition between *S.* Typhimurium IR715 (mPlum) and clinical isolates, as indicated (GFP), responding to a source of solubilized human feces over 300 s. Viewable at: https://www.youtube.com/watch?v=dLsFDV0XgpY

Movie S4. Chemotactic response of *S.* Typhimurium IR715 to L-Ser and indole treatments, related to Figure 4 and Figure S2. Representative CIRA experiments with treatment sources as indicated, over 300 s. Viewable at: https://www.youtube.com/watch?v=bNQMqF2QMek

References

1. Zhou, B., Szymanski, C.M., and Baylink, A. (2023). Bacterial chemotaxis in human diseases. Trends in Microbiology *31*, 453–467. https://doi.org/10.1016/j.tim.2022.10.007.

2. Keegstra, J.M., Carrara, F., and Stocker, R. (2022). The ecological roles of bacterial chemotaxis. Nature Reviews Microbiology. https://doi.org/10.1038/s41579-022-00709-w.

3. Bi, S., and Sourjik, V. (2018). Stimulus sensing and signal processing in bacterial
chemotaxis. Current Opinion in Microbiology *45*, 22–29.
https://doi.org/10.1016/j.mib.2018.02.002.

4. Matilla, M.A., and Krell, T. (2018). The effect of bacterial chemotaxis on host infection and
pathogenicity. FEMS Microbiology Reviews *42*. https://doi.org/10.1093/femsre/fux052.

5. Matilla, M.A., Gavira, J.A., and Krell, T. (2023). Accessing nutrients as the primary benefit
arising from chemotaxis. Current Opinion in Microbiology *75*, 102358.
https://doi.org/10.1016/j.mib.2023.102358.

6. Matilla, M.A., and Krell, T. (2017). Chemoreceptor-based signal sensing. Current Opinion in
Biotechnology *45*, 8–14. https://doi.org/10.1016/j.copbio.2016.11.021.

7. Ortega, Á., Zhulin, I.B., and Krell, T. (2017). Sensory Repertoire of Bacterial
Chemoreceptors. Microbiol Mol Biol Rev *81*, e00033-17.
https://doi.org/10.1128/MMBR.00033-17.

8. Rogers, A.W.L., Tsolis, R.M., and Bäumler, A.J. (2020). Salmonella versus the Microbiome.
Microbiology and Molecular Biology Reviews *85*, e00027-19.
https://doi.org/10.1128/MMBR.00027-19.

9. Vos, W.M. de, Tilg, H., Hul, M.V., and Cani, P.D. (2022). Gut microbiome and health:
mechanistic insights. Gut *71*, 1020–1032. https://doi.org/10.1136/gutjnl-2021-326789.

10. Hill, J.H., Massaquoi, M.S., Sweeney, E.G., Wall, E.S., Jahl, P., Bell, R., Kallio, K., Derrick,
D., Murtaugh, L.C., Parthasarathy, R., et al. (2022). BefA, a microbiota-secreted membrane
disrupter, disseminates to the pancreas and increases β cell mass. Cell Metabolism *34*, 1779-
1791.e9. https://doi.org/10.1016/j.cmet.2022.09.001.

11. Li, X., Zhang, B., Hu, Y., and Zhao, Y. (2021). New Insights Into Gut-Bacteria-Derived
Indole and Its Derivatives in Intestinal and Liver Diseases. Front. Pharmacol. *12*, 769501.
https://doi.org/10.3389/fphar.2021.769501.

12. Lee, J.-H., and Lee, J. (2010). Indole as an intercellular signal in microbial communities.
FEMS Microbiol Rev *34*, 426–444. https://doi.org/10.1111/j.1574-6976.2009.00204.x.

13. Gupta, R., Rhee, K.Y., Beagle, S.D., Chawla, R., Perdomo, N., Lockless, S.W., and Lele,
P.P. (2022). Indole modulates cooperative protein–protein interactions in the flagellar motor.
PNAS Nexus *1*, pgac035. https://doi.org/10.1093/pnasnexus/pgac035.

14. Darkoh, C., Chappell, C., Gonzales, C., and Okhuysen, P. (2015). A Rapid and Specific
Method for the Detection of Indole in Complex Biological Samples. Appl Environ Microbiol
81, 8093–8097. https://doi.org/10.1128/AEM.02787-15.

15. Chappell, C.L., Darkoh, C., Shimmin, L., Farhana, N., Kim, D.-K., Okhuysen, P.C., and
Hixson, J. (2016). Fecal Indole as a Biomarker of Susceptibility to Cryptosporidium
Infection. Infect Immun *84*, 2299–2306. https://doi.org/10.1128/IAI.00336-16.

16. Ye, X., Li, H., Anjum, K., Zhong, X., Miao, S., Zheng, G., Liu, W., and Li, L. (2022). Dual
Role of Indoles Derived From Intestinal Microbiota on Human Health. Front. Immunol. *13*,
903526. https://doi.org/10.3389/fimmu.2022.903526.

17. Kumar, A., and Sperandio, V. (2019). Indole Signaling at the Host-Microbiota-Pathogen
Interface. mBio *10*, e01031-19. https://doi.org/10.1128/mBio.01031-19.

18. Yang, J., Chawla, R., Rhee, K.Y., Gupta, R., Manson, M.D., Jayaraman, A., and Lele, P.P.
(2020). Biphasic chemotaxis of *Escherichia coli* to the microbiota metabolite indole. Proc.
Natl. Acad. Sci. U.S.A. *117*, 6114–6120. https://doi.org/10.1073/pnas.1916974117.

19. Bansal, T., Englert, D., Lee, J., Hegde, M., Wood, T.K., and Jayaraman, A. (2007).
Differential Effects of Epinephrine, Norepinephrine, and Indole on *Escherichia coli*
O157:H7 Chemotaxis, Colonization, and Gene Expression. Infect Immun *75*, 4597–4607.
https://doi.org/10.1128/IAI.00630-07.

20. Livne, N., and Vaknin, A. (2022). Collective responses of bacteria to a local source of
conflicting effectors. Sci Rep *12*, 4928. https://doi.org/10.1038/s41598-022-08762-4.

21. Englert, D.L., Manson, M.D., and Jayaraman, A. (2009). Flow-Based Microfluidic Device
for Quantifying Bacterial Chemotaxis in Stable, Competing Gradients. Appl Environ
Microbiol *75*, 4557–4564. https://doi.org/10.1128/AEM.02952-08.

22. Adler, J., and Tso, W.-W. (1974). "Decision"-Making in Bacteria: Chemotactic Response of
Escherichia coli to Conflicting Stimuli. Science *184*, 1292–1294.
https://doi.org/10.1126/science.184.4143.1292.

23. Kohli, N., Crisp, Z., Riordan, R., Li, M., Alaniz, R.C., and Jayaraman, A. (2018). The
microbiota metabolite indole inhibits Salmonella virulence: Involvement of the PhoPQ two-
component system. PLOS ONE *13*, e0190613. https://doi.org/10.1371/journal.pone.0190613.

24. Glenn, S.J., Gentry-Lear, Z., Shavlik, M., Harms, M.J., Asaki, T.J., and Baylink, A. (2024).
Bacterial vampirism mediated through taxis to serum. eLife *12*, RP93178.
https://doi.org/10.7554/eLife.93178.3.

25. Piñas, G.E., DeSantis, M.D., Cassidy, C.K., and Parkinson, J.S. (2022). Hexameric rings of
the scaffolding protein CheW enhance response sensitivity and cooperativity in Escherichia
coli chemoreceptor arrays. Science Signaling *15*, eabj1737.
https://doi.org/10.1126/scisignal.abj1737.

26. Burt, A., Cassidy, C.K., Ames, P., Bacia-Verloop, M., Baulard, M., Huard, K., Luthey-
Schulten, Z., Desfosses, A., Stansfeld, P.J., Margolin, W., et al. (2020). Complete structure
of the chemosensory array core signalling unit in an E. coli minicell strain. Nat Commun *11*,
743. https://doi.org/10.1038/s41467-020-14350-9.

27. Winter, S.E., Lopez, C.A., and Bäumler, A.J. (2013). The dynamics of gut-associated
microbial communities during inflammation. EMBO Reports *14*, 319–327.
https://doi.org/10.1038/embor.2013.27.

28. Rivera-Chávez, F., Winter, S.E., Lopez, C.A., Xavier, M.N., Winter, M.G., Nuccio, S.-P.,
Russell, J.M., Laughlin, R.C., Lawhon, S.D., Sterzenbach, T., et al. (2013). Salmonella Uses
Energy Taxis to Benefit from Intestinal Inflammation. PLoS Pathog *9*, e1003267.
https://doi.org/10.1371/journal.ppat.1003267.

29. Winter, S.E., Thiennimitr, P., Winter, M.G., Butler, B.P., Huseby, D.L., Crawford, R.W.,
Russell, J.M., Bevins, C.L., Adams, L.G., Tsolis, R.M., et al. (2010). Gut inflammation
provides a respiratory electron acceptor for Salmonella. Nature *467*, 426–429.
https://doi.org/10.1038/nature09415.

30. Cooper, K.G., Chong, A., Kari, L., Jeffrey, B., Starr, T., Martens, C., McClurg, M., Posada,
V.R., Laughlin, R.C., Whitfield-Cargile, C., et al. (2021). Regulatory protein HilD stimulates
Salmonella Typhimurium invasiveness by promoting smooth swimming via the methyl-
accepting chemotaxis protein McpC. Nat Commun *12*, 348. https://doi.org/10.1038/s41467-
020-20558-6.

31. Vega, N.M., Allison, K.R., Samuels, A.N., Klempner, M.S., and Collins, J.J. (2013).
Salmonella typhimurium intercepts *Escherichia coli* signaling to enhance antibiotic
tolerance. Proc. Natl. Acad. Sci. U.S.A. *110*, 14420–14425.
https://doi.org/10.1073/pnas.1308085110.

32. Roager, H.M., and Licht, T.R. (2018). Microbial tryptophan catabolites in health and disease.
Nat Commun *9*, 3294. https://doi.org/10.1038/s41467-018-05470-4.

33. Kronsteiner, B., Bassaganya-Riera, J., Philipson, C., Viladomiu, M., Carbo, A., Pedragosa,
M., Vento, S., and Hontecillas, R. (2013). Helicobacter pylori Infection in a Pig Model Is
Dominated by Th1 and Cytotoxic CD8[+] T Cell Responses. Infect Immun *81*, 3803–3813.
https://doi.org/10.1128/IAI.00660-13.

34. Smith, A.C., and Swindle, M.M. (2006). Preparation of Swine for the Laboratory. ILAR
Journal *47*, 358–363. https://doi.org/10.1093/ilar.47.4.358.

35. Poutahidis, T., Tsangaris, T., Kanakoudis, G., Vlemmas, I., Iliadis, N., and Sofianou, D.
(2001). *Helicobacter pylori* -induced Gastritis in Experimentally Infected Conventional
Piglets. Vet Pathol *38*, 667–678. https://doi.org/10.1354/vp.38-6-667.

36. Boyen, F., Pasmans, F., Van Immerseel, F., Donné, E., Morgan, E., Ducatelle, R., and
Haesebrouck, F. (2009). Porcine *in vitro* and *in vivo* models to assess the virulence of
Salmonella enterica serovar Typhimurium for pigs. Lab Anim *43*, 46–52.
https://doi.org/10.1258/la.2007.007084.

37. Boyen, F., Haesebrouck, F., Vanparys, A., Volf, J., Mahu, M., Van Immerseel, F., Rychlik,
I., Dewulf, J., Ducatelle, R., and Pasmans, F. (2008). Coated fatty acids alter virulence
properties of Salmonella Typhimurium and decrease intestinal colonization of pigs.
Veterinary Microbiology *132*, 319–327. https://doi.org/10.1016/j.vetmic.2008.05.008.

38. MacBeth, K.J., and Lee, C.A. (1993). Prolonged inhibition of bacterial protein synthesis
abolishes Salmonella invasion. Infect Immun *61*, 1544–1546.
https://doi.org/10.1128/iai.61.4.1544-1546.1993.

39. Sharma, A., and Puhar, A. (2019). Gentamicin Protection Assay to Determine the Number of
Intracellular Bacteria during Infection of Human TC7 Intestinal Epithelial Cells by Shigella
flexneri. BIO-PROTOCOL *9*. https://doi.org/10.21769/BioProtoc.3292.

40. Kitamoto, S., Alteri, C.J., Rodrigues, M., Nagao-Kitamoto, H., Sugihara, K., Himpsl, S.D.,
Bazzi, M., Miyoshi, M., Nishioka, T., Hayashi, A., et al. (2020). Dietary l-serine confers a
competitive fitness advantage to Enterobacteriaceae in the inflamed gut. Nature
Microbiology *5*, 116–125. https://doi.org/10.1038/s41564-019-0591-6.

41. Sugihara, K., and Kamada, N. (2020). THE ROLE OF DIETARY L-SERINE IN THE
REGULATION OF INTESTINAL MUCUS BARRIER DURING INFLAMMATION.
Inflammatory Bowel Diseases *26*, S42–S42. https://doi.org/10.1093/ibd/zaa010.110.

42. Ranque, C.L., Stroble, C., Amicucci, M.J., Tu, D., Diana, A., Rahmannia, S., Suryanto, A.H.,
Gibson, R.S., Sheng, Y., Tena, J., et al. (2020). Examination of Carbohydrate Products in
Feces Reveals Potential Biomarkers Distinguishing Exclusive and Nonexclusive
Breastfeeding Practices in Infants. J Nutr *150*, 1051–1057.
https://doi.org/10.1093/jn/nxaa028.

43. Ke, Z., Lu, Z., Li, Q., and Tong, W. (2024). Intestinal glucose excretion: A potential
mechanism for glycemic control. Metabolism *152*, 155743.
https://doi.org/10.1016/j.metabol.2023.155743.

44. Larke, J.A., Bacalzo, N., Castillo, J.J., Couture, G., Chen, Y., Xue, Z., Alkan, Z., Kable,
M.E., Lebrilla, C.B., Stephensen, C.B., et al. (2023). Dietary Intake of Monosaccharides
from Foods is Associated with Characteristics of the Gut Microbiota and Gastrointestinal
Inflammation in Healthy US Adults. The Journal of Nutrition *153*, 106–119.
https://doi.org/10.1016/j.tjnut.2022.12.008.

45. Shen, X., Jain, A., Aladelokun, O., Yan, H., Gilbride, A., Ferrucci, L.M., Lu, L., Khan, S.A.,
and Johnson, C.H. (2022). Asparagine, colorectal cancer, and the role of sex, genes,
microbes, and diet: A narrative review. Front Mol Biosci *9*, 958666.
https://doi.org/10.3389/fmolb.2022.958666.

46. Shariat, N., Kirchner, M.K., Sandt, C.H., Trees, E., Barrangou, R., and Dudley, E.G. (2013).
Subtyping of Salmonella enterica Serovar Newport Outbreak Isolates by CRISPR-MVLST
and Determination of the Relationship between CRISPR-MVLST and PFGE Results. J Clin
Microbiol *51*, 2328–2336. https://doi.org/10.1128/JCM.00608-13.

47. Ferrari, R.G., Rosario, D.K.A., Cunha-Neto, A., Mano, S.B., Figueiredo, E.E.S., and Conte-
Junior, C.A. (2019). Worldwide Epidemiology of *Salmonella* Serovars in Animal-Based
Foods: a Meta-analysis. Appl Environ Microbiol *85*, e00591-19.
https://doi.org/10.1128/AEM.00591-19.

48. Gavira, J.A., Gumerov, V.M., Rico-Jiménez, M., Petukh, M., Upadhyay, A.A., Ortega, A.,
Matilla, M.A., Zhulin, I.B., and Krell, T. (2020). How Bacterial Chemoreceptors Evolve
Novel Ligand Specificities. mBio *11*, e03066-19. https://doi.org/10.1128/mBio.03066-19.

49. Matilla, M.A., Ortega, Á., and Krell, T. (2021). The role of solute binding proteins in signal
transduction. Computational and Structural Biotechnology Journal *19*, 1786–1805.
https://doi.org/10.1016/j.csbj.2021.03.029.

50. Chimerel, C., Field, C.M., Piñero-Fernandez, S., Keyser, U.F., and Summers, D.K. (2012).
Indole prevents Escherichia coli cell division by modulating membrane potential. Biochim
Biophys Acta *1818*, 1590–1594. https://doi.org/10.1016/j.bbamem.2012.02.022.

51. Melander, R.J., Minvielle, M.J., and Melander, C. (2014). Controlling bacterial behavior
with indole-containing natural products and derivatives. Tetrahedron *70*, 6363–6372.
https://doi.org/10.1016/j.tet.2014.05.089.

52. Moore, J.P., Kamino, K., Kottou, R., Shimizu, T.S., and Emonet, T. (2024). Signal
integration and adaptive sensory diversity tuning in *Escherichia coli* chemotaxis. Cell
Systems *15*, 628-638.e8. https://doi.org/10.1016/j.cels.2024.06.003.

53. Huang, J.Y., Sweeney, E.G., Sigal, M., Zhang, H.C., Remington, S.J., Cantrell, M.A., Kuo,
C.J., Guillemin, K., and Amieva, M.R. (2015). Chemodetection and Destruction of Host
Urea Allows Helicobacter pylori to Locate the Epithelium. Cell Host & Microbe *18*, 147–
156. https://doi.org/10.1016/j.chom.2015.07.002.

54. Huang, J.Y., Sweeney, E.G., Guillemin, K., and Amieva, M.R. (2017). Multiple Acid
Sensors Control Helicobacter pylori Colonization of the Stomach. PLOS Pathogens *13*,
e1006118. https://doi.org/10.1371/journal.ppat.1006118.

55. Beltran, P., Plock, S.A., Smith, N.H., Whittam, T.S., Old, D.C., and Selander, R.K. (1991).
Reference collection of strains of the Salmonella typhimurium complex from natural
populations. Journal of General Microbiology *137*, 601–606.
https://doi.org/10.1099/00221287-137-3-601.

56. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

57. Parthasarathy, R. (2012). Rapid, accurate particle tracking by calculation of radial symmetry
centers. Nat Methods *9*, 724–726. https://doi.org/10.1038/nmeth.2071.

58. Sugihara, K., Kitamoto, S., Saraithong, P., Nagao-Kitamoto, H., Hoostal, M., McCarthy, C.,
Rosevelt, A., Muraleedharan, C.K., Gillilland, M.G., Imai, J., et al. (2022). Mucolytic
bacteria license pathobionts to acquire host-derived nutrients during dietary nutrient
restriction. Cell Reports *40*, 111093. https://doi.org/10.1016/j.celrep.2022.111093.

59. Chandra, K. (2022). Gentamicin Protection assay (Intracellular Survival Assay) for
Salmonella Typhimurium/Typhi v1. Preprint,
https://doi.org/10.17504/protocols.io.b4zmqx46
https://doi.org/10.17504/protocols.io.b4zmqx46.

60. Rivera Calo, J., Rubinelli, P.M., and Ricke, S.C. (2024). Gentamicin Susceptibility and
Comparison of Adhesion and Invasion of Caco-2 and HD11 Cell Lines by Salmonella
enterica Serotypes. Applied Sciences *14*, 3305. https://doi.org/10.3390/app14083305.

61. Ben-David, A., and Davidson, C.E. (2014). Estimation method for serial dilution
experiments. Journal of Microbiological Methods *107*, 214–221.
https://doi.org/10.1016/j.mimet.2014.08.023.

62. Macho, A.P., Zumaquero, A., Ortiz-Martín, I., and Beuzón, C.R. (2007). Competitive index
in mixed infections: a sensitive and accurate assay for the genetic analysis of *Pseudomonas*
syringae –plant interactions. Molecular Plant Pathology *8*, 437–450.
https://doi.org/10.1111/j.1364-3703.2007.00404.x.

63. Auerbuch, V., Lenz, L.L., and Portnoy, D.A. (2001). Development of a competitive index
assay to evaluate the virulence of Listeria monocytogenes actA mutants during primary and
secondary infection of mice. Infect Immun *69*, 5953–5957.
https://doi.org/10.1128/IAI.69.9.5953-5957.2001.