Navigating contradictions: *Salmonella* Typhimurium chemotaxis amidst conflicting stimuli of

the intestinal environment

Kailie Franco,[a] Zealon Gentry-Lear,[b*] Michael Shavlik,[c] Michael J. Harms,[b,c] Arden Baylink[a,d#]

[a] Department of Veterinary Microbiology and Pathology, Washington State University, Pullman,

Washington, USA

[b] Institute of Molecular Biology, University of Oregon, Eugene, Oregon, USA

[c] Department of Chemistry & Biochemistry, University of Oregon, Eugene, Oregon, USA

[d] Amethyst Antimicrobials, LLC, Pullman, USA

[#] Address correspondence to Arden Baylink, PhD, arden.baylink@wsu.edu

[*] Present address: Zealon Gentry-Lear, Department of Microbiology, University of Washington,

Seattle, Washington, USA

Abstract

Motile bacteria sense and avoid deleterious stimuli in their environment through chemorepulsion, a behavior that helps them locate permissive ecological niches. In the gut, indole is a bacteriostatic compound produced by the microbiota and is thought to act as a chemorepellent for invading pathogens, thereby protecting the host against infection. The principal reservoir of intestinal indole is fecal matter, a complex biological material that contains both attractant and repellent stimuli. Whether indole in its natural context is sufficient for pathogen chemorepulsion or host protection has remained unknown. Using an intestinal explant system, we show that while pure indole indeed suppresses an infection advantage mediated through chemotaxis for the enteric pathogen *Salmonella enterica* serovar Typhimurium, this effect is abolished in the presence of other chemoeffectors present in feces, including the chemoattractant L-Serine (L-Ser), in a manner dependent on the chemoreceptor Tsr. Live imaging reveals that although *S.* Typhimurium is repelled by pure indole, the pathogen is actually strongly attracted to human fecal matter despite its high indole content, and that this response is mediated by Tsr, which simultaneously senses both indole and L-Ser. Fecal attraction is conserved across diverse *Enterobacteriaceae* species that harbor Tsr orthologues, including *Escherichia coli*, *Citrobacter koseri*, *Enterobacter cloacae*, and clinical isolates of non-typhoidal *Salmonella*. In a defined system of fecal chemoeffectors, we find that L-Ser and other fecal chemoattractants override indole chemorepulsion, but the magnitude of bacterial chemoattraction is controlled by indole levels. Together, these findings suggest that indole in its native context is not protective against enteric infection and that indole taxis actually benefits pathogens during infection by locating niches with low competitor density. Our study highlights the limitations of applying single-effector studies in predicting bacterial behavior in natural environments, where chemotaxis is shaped by the integration of multiple, often opposing, chemical signals.

Introduction

Motile bacteria that colonize the gastrointestinal tracts of humans and other animals employ chemotaxis to sense chemical effectors in the gut lumen and navigate to environments conducive to growth and colonization [1–5]. This process is controlled by chemoreceptor proteins, which recognize chemical effectors and transduce signals through a phosphorylation cascade to regulate flagellar rotation and swimming direction, ultimately determining the spatial and temporal patterns of bacterial colonization (Fig. 1A) [1,2,6,7]. While many effectors have been studied and characterized in isolation as chemoattractants or chemorepellents [4,7], natural environments like the gut contain complex mixtures of opposing signals. Only a handful of studies have investigated how bacteria navigate conflicting chemical gradients, and much remains to be learned about how bacteria prioritize these signals to direct their movement and colonization (Fig. 1A) [1,4,8–15].

A chemical effector of major importance for enteric bacterial communities is indole, an interbacterial signaling molecule that regulates diverse aspects of physiology and lifestyle [16–18]. Indole is excreted by gut microbiota as a byproduct of tryptophan metabolism and accumulates to millimolar levels in human feces (Fig. 1A) [16,19,20]. Indole is amphipathic and can transit bacterial membranes to regulate biofilm formation and motility, suppress virulence programs, and exert bacteriostatic and bactericidal effects at high concentrations [16–18,20–24]. Indole was one of the earliest identified chemorepellents, and subsequent work has extensively explored its role in *Escherichia coli* chemotaxis, mostly examining responses to indole as a singular effector (Table S1) [8,9,11,14,18,25]. Recent studies have advanced understanding of the molecular mechanisms underlying *E. coli* indole taxis and the involvement of the chemoreceptor taxis to serine and repellents (Tsr) (Fig. 1A) [8,9,14,18].

From this body of research, the hypothesis emerged that indole from the gut microbiota functions to repel pathogens and restrict their growth as a mechanism of colonization resistance [8,16–18,23,24]. If true, this could represent an intriguing avenue for cultivating microbiomes that are more robust against pathogen infiltration—a major area of interest for improving gut health [26,27]. However, this hypothesis is based largely on observations of bacterial chemorepulsion in response to indole as a single effector, and it remains unclear whether chemotactic behavior is similar or altered in the presence of other intestinal effectors. For instance, fecal material, while rich in indole, also contains high concentrations of sugars and amino acids that serve as nutrients

and chemoattractants—factors that could diminish, nullify, or have no influence on indole

chemorepulsion [1,21,22,28]. Although indole may suppress bacterial growth, nutrients derived from

the host diet could offset this effect, allowing bacteria to tolerate indole and still benefit from

colonizing indole-rich niches. Indeed, pathogens frequently succeed in establishing enteric

infections, suggesting that they can tolerate indole or circumvent its effects under certain

conditions. Thus, bacteria in the intestinal environment must navigate contradictory chemotaxis

signals, and how they resolve these conflicts influences infection, pathogenesis, and host health

in ways that remain to be elucidated. Furthermore, because indole taxis has only been studied in

E. coli, it remains unconfirmed whether other enteric species even chemotactically sense or

respond to indole (Table S1).

In this study, we aimed to (1) test the hypothesis that indole protects against intestinal

infection and (2) determine how enteric pathogens use chemotaxis to navigate the complex

mixture of opposing chemical cues present in fecal material, a major source of both indole and

nutrients within the gut. We used *Salmonella enterica* serovar Typhimurium as a model

pathogen, as it requires chemotaxis and the chemoreceptor Tsr for efficient infection and cellular

invasion of intestinal tissue [29–35]. Tsr is of particular interest because it mediates responses to

both chemorepellents and the chemoattractant L-Serine (L-Ser), suggesting an important role in

integrating contradictory chemotactic signals for *S.* Typhimurium and other *Enterobacteriaceae*

that possess Tsr orthologues (Fig. 1A-B) [1,8,9,14,18,25,36–39]. *S.* Typhimurium also differs from *E.*

coli in that it lacks tryptophanase and cannot itself produce indole, thereby offering a novel

perspective on indole taxis [40,41]. Our findings reveal that bacterial chemotaxis within biologically

relevant mixtures of effectors cannot be reliably inferred from studies of individual compounds

alone, with important implications for understanding how chemotaxis influences pathogen

behavior within the gut.

Results

Fecal indole is insufficient to protect against pathogen invasion in a colonic explant model

We sought to test whether indole in human fecal matter protects against *Salmonella enterica*

serovar Typhimurium infection and whether this involves indole chemorepulsion mediated by

the chemoreceptor Tsr. *Salmonella* Typhimurium preferentially infects tissue of the distal ileum

but also infects the cecum and colon in humans and animal models [42–46]. We presume that the

amount of indole is greatest in the lower gastrointestinal tract, where tryptophan has been

Fig. 1. Chemotaxis-mediated infection advantages in the presence of fecal effectors. A. Overview of the role of Tsr in coordinating responses to conflicting stimuli. B. Experimental design of colonic explant infections. See Materials & Methods for experimental details such as tissue dimensions. C. Conceptual model of the explant infection system. The effectors from the treated tissue (gray) diffuse into the surrounding buffer solution providing a gradient. Note that the bacteria are not immersed in the effector solution, and experience a local concentration during infection much lower than the effector pretreatment. Quantifications of tissue-associated bacteria reflect the ability of chemotaxis to provide an advantage (black arrow) in accessing the intestinal mucosa (reddish brown). D. Serine (presumed to be nearly 100% L-Ser, see Materials & Methods) and indole content of liquid human fecal treatments, as measured by mass spectrometry. E-I. Competitive indices (CI) of colony-forming units (CFUs) recovered from co-infected swine explant tissue, either from the total homogenate (open box and whiskers plots), or from tissue washed with gentamicin to kill extracellular and attached cells, which we refer to as the "invaded" intracellular population (checkered box and whisker plots), as indicated. Each data point represents a single experiment of a section of tissue infected with bacteria, normalized by tissue weight, and the CI of CFUs recovered from that tissue (n=7-10). Boxes show median values (line) and upper and lower quartiles, and whiskers show max and min values. Effect size (Cohen's *d*) and statistical significance are noted for each experiment in relation to competitive advantage, i.e. deviation from a CI of 1 (not significant, ns; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). See also Figure S1 for competition between WT and an invasion-inhibited mutant *invA*, and Figure S2 for disaggregated CFU enumerations for each experimental group prior to CI calculation. Data S1 contains all numerical CFU measurements.

maximally digested by microbial tryptophanases. To mimic this environment, we developed a

swine colonic explant model that simulates the architecture and size of adult human colonic
tissue (Fig. 1B) [41,47–51]. This model was based on a prior study using explant tissue to
characterize cellular invasion via gentamicin washes, which kill extracellular and surface-
attached bacteria (Fig. 1B) [52]. Gentamicin washing is also a commonly used method to quantify
intracellular *Salmonella* Typhimurium populations in cell culture experiments [53,54].

A section of colonic tissue was prepared for each experiment by gentle cleaning and then
soaked with different effector solutions for 1 h: solubilized human feces, purified indole and/or
L-Ser at fecal-relevant concentrations, or buffer as a control (see Materials & Methods for tissue
dimensions and additional experimental details). Subsequently, the tissue was removed from the
effector treatment and oriented with the luminal side downward onto a Mattek dish containing
300 µl of buffer with motile *S*. Typhimurium (Fig. 1B). To be clear, in this system the bacteria
are not immersed in the effector treatment and experience an effector concentration far lower
than used for the soak of the tissue prior to infection where the residual effector diffuses outward
from the tissue into the much larger volume of buffer in which the cells are swimming (Fig. 1C).
This establishes a chemical gradient which we can use to quantify the degree to which different
effector treatments are permissive of pathogen association with, and cellular invasion of, the
intestinal mucosa (Fig. 1C). Using this approach we sought to test infection based on fecal
treatments and fecal-relevant concentrations of L-Ser and indole (Fig. 1D).

We employed a strategy of co-infections in order to compete and compare the advantages
conferred by chemotaxis using *S*. Typhimurium strain IR715 wildtype (WT) and either a *cheY*
mutant (motile but non-responsive to chemoeffector stimuli) or *tsr* deletion mutant (Fig. 1B,
Table 1, see Materials & Methods) [30]. The functionality of these mutants has been previously
confirmed through *in vivo* infection studies using genetically complemented strains [30,31]. To
assess the role of chemotaxis in infection, we quantified bacteria harvested from tissue
homogenates at 1, 3, and 6 h post-infection (Fig. 1C, Materials & Methods) [55,56]. Using this
experimental setup, we found for tissue pretreated with fecal material that WT had a competitive
advantage over an invasion-inhibited mutant (*invA*) in homogenates from gentamicin-washed
tissue, but no advantage in unwashed homogenates, supporting that gentamicin washing selects
for intracellular bacteria [57]. For simplicity in discussing the explant infection data we refer to
these two types of quantifications as "invaded" (i.e. *Salmonella* that have entered non-phagocytic
host cells) and "total" bacteria in the figures, respectively (Fig. S1) [58].

In buffer-treated explant experiments, WT *S.* Typhimurium exhibits a 5- to 10-fold time-
dependent advantage in colonization and cellular invasion compared to chemotactic mutants,
indicating that chemotaxis, and specifically Tsr, promotes tissue colonization in this system (Fig.
1E, Fig. S2A–B, Data S1). The mechanism mediating this advantage isn't clear, but could arise
from a combination of factors, including sensing of effectors emitted from the tissue, redox or
energy taxis, and/or swimming behaviors that enhance infection [5,30,31,35]. This experiment
indicates that under baseline conditions, the intestinal mucosa is accessible to the pathogen. The
hypothesis that indole protects against pathogen colonization predicts that feces, the major
biological reservoir of gut indole, should confer protection against infection. Contrary to this
prediction, we found that fecal treatment provided a similar infection advantage as buffer
treatment, and this effect was mediated by chemotaxis and Tsr (Fig. 1F, Fig. S2C–D). One
notable difference, however, was that fecal treatments yielded a higher competitive advantage for
the WT invaded population over the total population at 3 hours compared to buffer treatment
(comparing buffer and fecal treatments: WT vs *cheY*, $p = 0.18$ and $p = 0.02$; WT vs *tsr*, $p = 0.35$
and $p = 0.0004$, respectively; Fig. 1E–F, Data S1). That the phenotype fades at longer time points
could relate to the effector gradient being eliminated by diffusion.

Analysis of the liquefied human fecal matter used in this study revealed an indole
concentration of 862 µM, consistent with previously reported ranges (0.5–5 mM) (Fig. 1D;
Materials & Methods) [16,19–22]. When colonic tissue was treated with purified indole at this
concentration, WT loses its competitive advantage over the chemotactic mutants (Fig. 1G, Fig.
S2E–F, Data S1). Given that Tsr mediates attraction to L-Ser in both *Escherichia coli* and
Salmonella Typhimurium, we hypothesized that L-Ser present in feces might negate the
protective effect of indole [1,36,59,60]. Treatment with 338 µM L-Ser alone, the concentration present
in our fecal sample (Fig. 1D; Materials & Methods), conferred a WT advantage similar to buffer
and fecal treatments (Fig. 1H) and WT also exhibits a colonization advantage when L-Ser is co-
administered with indole (Fig. 1I, Data S1).

Our key takeaway from these experiments is that pretreatment of intestinal tissue with
indole alone is unique in that the WT strain gains no infection advantage in this context (Fig.
1G). In contrast, under all other treatment conditions, the WT infects the tissue to a greater extent
than the chemotactic mutants (Fig. 1E-I). Put another way, chemotaxis and Tsr enhance pathogen
transit of the chemical gradient and increase access to the intestinal tissue in all conditions except

when indole is the sole effector. This was surprising, given that other treatments contain the

same concentration of indole and still permit a chemotaxis- and Tsr-mediated advantage (Fig. 1).

To us, this suggests that bacterial perception of indole via chemotaxis differs fundamentally

depending on whether indole is the only effector present or amidst other fecal effectors. Notably,

only fecal treatment resulted in a greater competitive advantage for the invaded population than

for the total population (Fig. 1F).

Enterobacteriaceae species are attracted to human feces despite high indole content

Having found that chemotaxis and Tsr mediate efficient infection, but do not confer an

advantage when indole is the only effector, we next sought to understand the chemotactic

behaviors orchestrated by Tsr in response to indole-rich feces. Given that feces represents the

highest concentration of indole that *S*. Typhimurium is likely to encounter natively, we expected

to observe chemorepulsion (Fig. 1D, Table S1) [8,9,14,25,39]. We employed the chemosensory

injection rig assay (CIRA) for live-imaging of bacterial chemotaxis responses to a source of

effectors injected through a glass microcapillary [36]. The flow and dynamic nature of the gut

lumen make this a suitable *in vitro* approach for modeling and studying enteric chemotaxis

responses [36,61].

In this assay, chemoattraction is observed as an influx of cells toward the effector source,

whereas chemorepulsion is indicated by a decrease in local cell density (Fig. S3). As described

previously, effector injection creates a steep chemical microgradient [36]. Using mathematical

modeling of the diffusion of fecal-relevant concentrations of indole and L-Ser, we approximated

the local concentrations experienced by bacteria at varying distances from the injection site. For

most of the field of view, these concentrations fall within the picomolar to low nanomolar range

(Fig. 2A; Materials & Methods) [36].

Over five minutes, we found both WT and *tsr* exhibit strong chemoattraction to fecal

material, whereas *cheY* remains randomly distributed (Fig. 2B, Movie 1). By examining the

radial distribution of the bacterial populations, we found WT more tightly centers around the

treatment source than *tsr* (Fig. 2C-E, Movie 1). In terms of the rate of bacterial accumulation, the

chemoattraction of *tsr* lags behind the WT for the first 120 s (Fig. 2F-G, Movie 1). We wondered

how these deficiencies in fecal attraction might translate to direct competition, where different

strains are experiencing the same treatment source simultaneously. To address this, we

Fig. 2. *Salmonella* Typhimurium exhibits attraction toward indole-rich liquid human fecal material. A. Diffusion modeling showing calculated local concentrations in CIRA experiments with liquid human fecal material based on distance from the central injection source. B. Max projections of representative *S.* Typhimurium IR715 responses to a central source of injected liquid human fecal material. C-E. Bacterial population density over time in response to fecal treatment. The initial uniform population density in these plots is indicated with the blue line (time 0), and the final mean distributions with the red line (time 280 s), with the mean distributions between these displayed as a blue-to-red spectrum at 10 s intervals. F-G. Temporal analyses of area under the curve (AUC) or relative number of bacteria within 150 µm of the source. Effect size (Cohen's *d*) comparing responses of WT and *tsr* attraction at 120 s post-treatment is indicated. Data were collected at 30 °C. Data are means and error bars are standard error of the mean (SEM, n=3-5). See also Movie 1, Table S1, and Figure S3.

performed CIRA with solubilized human feces and two strains present in the same pond, which we tracked independently through fluorescent markers (Fig. 3) [36]. As expected, WT shows a strong chemoattraction response versus *cheY* (Fig. 3A, Movie 2). Interestingly, we found that when competed directly, WT vastly outperforms *tsr*, with the maximal bacterial distribution in proximity to the treatment source higher by about 4-fold (Fig. 3B, Movie 2). These data confirm that despite its high indole content, *S.* Typhimurium is attracted to human fecal material through chemotaxis, and this response involves Tsr, although not as the sole mediator. We expect the attraction of the *tsr* mutant is explained by the fact that *S.* Typhimurium possesses other chemoreceptors that detect glucose, galactose, ribose, and L-Asp as chemoattractants, which are also present in human feces [1,7,62–66].

Fig. 3. Non-typhoidal *Salmonella* exhibit fecal attraction. A-E. Dual-channel imaging of chemotactic responses to solubilized human feces by WT *S.* Typhimurium IR715 (pink) and isogenic mutants or clinical isolate strains (green), as indicated. Shown are max projections at time 295-300 s post-treatment. Data were collected at 37 °C. Data are means and error bars are standard error of the mean (SEM, n=3-5). See also Movie 2, Movie 3.

Recent work highlights how genetic diversity among *Salmonella* strains and differences

in Tsr expression, even within isogenic populations, modulate chemotaxis function [9,32]. To gain a

broader perspective on fecal taxis, we examined the responses among diverse non-typhoidal

Salmonella serovars and strains responsible for human infections. Using dual-channel imaging,

we compared *S.* Typhimurium IR715 with a clinical isolate of *S.* Typhimurium, SARA1, and

found both strains exhibit attraction to feces, although SARA1 shows a slightly weaker response

(Fig. 3C, Movie 3) [67]. We then tested a clinical isolate of *S.* Newport, an emerging cause of

salmonellosis in the United States and Europe [68,69]. This strain is strongly attracted to fecal

material, with a tighter accumulation of cells at the treatment source than *S.* Typhimurium IR715

(Fig. 3D, Movie 3). We also examined a clinical isolate of *S.* Enteritidis, a zoonotic pathogen

commonly transmitted from poultry, which displays weak attraction to fecal material (Fig. 3E,

Movie 3) [69].

Next, we extended this analysis to other disease-causing *Enterobacteriaceae* that possess

Tsr orthologues [36]. *E. coli* strain MG1655, commonly used for *in vitro* experiments, and *E. coli*

NCTC 9001, a strain isolated from human urine and associated with urinary tract infections, both

exhibited fecal attraction, although the response was more diffuse than that observed for

Salmonella (Fig. 4A–D; Movies 4–5) [70]. The clinical isolate *Citrobacter koseri* strain 4225-83

also showed fecal attraction, with a tight association near the effector source (Fig. 4E–F; Movie

6). Lastly, *Enterobacter cloacae* CDC 442-68, a clinical isolate with uncharacterized chemotaxis

behavior, appeared to exhibit fecal attraction as well, although this strain was not extensively

tested due to limited motility under laboratory conditions (Fig. S4).

Overall, we find that Tsr mediates fecal attraction in *Salmonella*, and that this behavior is

conserved among diverse *Enterobacteriaceae* that possess Tsr and are associated with human

infections. Although the degree of attraction varies, none of the enteric pathogens or pathobionts

Fig. 4. Representative *Enterobacteriaceae* exhibit fecal attraction. Shown are max projections from CIRA experiments over 5 s before fecal treatment and after 5 minutes of treatment, as well as quantifications of bacteria within 500 µm of the treatment source at these same time points for *E. coli* MG1655 (A-B, GFP-reporter), *E. coli* NCTC 9001 (C-D, phase), and *C. koseri* CDC 4225-83 (E-F, phase). Data were collected at 37 °C. Data are means and error bars are standard error of the mean (SEM, n=3-5). See also Movie 4, Movie 5, Movie 6.

tested exhibited chemorepulsion from feces, despite its high indole content.

Fecal chemoattractants override indole chemorepulsion

To better understand the relationship between indole and other fecal effectors in directing *S.*

Typhimurium chemotaxis, we next employed a reductionist approach and developed a mixture of

fecal effectors at physiological concentrations based on our measurements and the Human

Metabolome Database (Fig. 5) [28]. Along with the chemorepellent indole (862 µM), we tested

Fig. 5. Chemotactic responses to defined fecal effector mixtures. CIRA experiments with *S.* Typhimurium IR715 were performed with different combinations of fecal effectors (n=3-5). Shown are max projections from experiments over 5 s before fecal treatment and after 5 minutes of treatment as well as quantifications of bacteria within 500 µm of the treatment source at these same time points. Data are means and error bars are standard error of the mean (SEM, n=3-5). To achieve the greatest degree of sensitivity to differences in responses, experiments were performed using the same culture on the same day. The complete fecal effector mixture consists of indole (862 µM), L-Ser (338 µM), D-Glucose (970 µM), D-Galactose (78 µM), ribose (28.6 µM), and L-Asp (13 µM), modified to include or exclude certain effectors as indicated. See also Movie 7, Movie 8, Movie 9, and Movie 10 Data were collected at 30 °C.

combinations of fecal chemoattractants including L-Ser (338 µM), sensed through Tsr; D-glucose

(970 µM), D-galactose (78 µM), and ribose (28.6 µM), sensed through the chemoreceptor Trg;

and L-aspartate (L-Asp, 13 µM), sensed through the chemoreceptor Tar [1,7,28,36,71,72]. A low

density of motile cells (A□□□ ~ 0.05) was used in the CIRA experiments to increase sensitivity

for detecting attraction in response to different combinations of these fecal effectors (Fig. 5).

We observed that L-Ser was sufficient to negate indole chemorepulsion, and may even

elicit attraction, although this was not statistically significant (Fig. 5A–B; Movie 7). When all

effectors were present, bacteria were clearly attracted to the treatment (Fig. 5C–D; Movie 8),

with a slightly reduced attraction in the absence of L-Ser (Fig. 5E–F; Movie 9). Interestingly,

when all effectors were present but the concentration of indole was halved (431 µM), cells

exhibited the greatest degree of attraction (Fig. 5G–H; Movie 10).

From these data, we conclude that the Tsr ligand L-Ser can override chemorepulsion from

indole. However, this effect can also be mediated by other fecal effectors sensed through

different chemoreceptors, providing an explanation for the reduced, but still appreciable, fecal

attraction observed for the *tsr* mutant (Fig. 3B). While the overall responses to this mixture of

fecal effectors can be characterized as attraction, the bacteria remain sensitive to indole levels, as

reflected in the enhanced attraction observed in treatments with lower indole concentrations (Fig.

5G–H).

Mediation of opposing chemotactic responses by Tsr

We considered whether our inability to observe repulsion from fecal material and mixtures of

fecal effectors might be due to *S*. Typhimurium not sensing indole as a chemorepellent, since this

chemotactic response has only been previously described for *E. coli* (Table S1). We compared

chemotaxis responses to either 5 mM L-Ser or 5 mM indole and found that *S*. Typhimurium

responds rapidly to these two effectors as chemoattractants and chemorepellents, respectively

(Fig. S3H-I). Treatment with 5 mM indole, a concentration at the upper end of what occurs in the

human gut [22], induces rapid chemorepulsion with the bacteria vacating the region proximal to the

source (Fig. S3I) . Interestingly, the chemorepulsion response occurs faster than chemoattraction,

with a zone of avoidance clearly visible within the first 10 s of indole exposure (Fig. S3H-I,

Movie 11).

We next wondered if perhaps our fecal treatments contained insufficient indole to elicit

chemorepulsion from *S*. Typhimurium. To identify the effective source concentrations that drive

indole chemorepulsion and understand the temporal dynamics of this response, we performed a

titration of indole across 0.05-10 mM (Fig. 6A-B). At all concentrations tested, indole induces

chemorepulsion, and the bacteria avoid the treatment source for the duration of the 5-minute

experiment (Fig. 6A-B). At source concentrations exceeding 3 mM most motile cells vacate the
field of view within 60 s (Fig. 6A). Integrating these chemorepulsion responses and fitting them
to a Monod curve suggests that an indole source concentration of approximately 67 µM is
sufficient for half-maximal ($K_{1/2}$) chemorepulsion (Fig. 6C). These data show that even though
we observe strong attraction to fecal material, pure indole at the concentration present in fecal
material, and far lower, is indeed a strong chemorepellent for *S*. Typhimurium.

Based on its function in *E. coli*, we hypothesized that both indole chemorepulsion and L-
Ser chemoattraction for *S*. Typhimurium could be partly or fully mediated through Tsr [7,8,38]. We
compared the chemotactic responses of the WT and *tsr* strains when exposed to sources of these
effectors and found Tsr to be required for both chemorepulsion from indole and chemoattraction
to L-Ser (Fig. 6E-F). The canonical mode of chemoreceptor effector recognition involves binding
of the effector to the periplasmic ligand-binding domain (LBD) [7,73], but the mechanism by which

Fig. 6. Tsr mediates indole chemorepulsion in *S*. Typhimurium. A. Representative max projections of responses at 295-300 s of indole treatment. B-C. Quantification of chemorepulsion as a function of indole concentration (n=3-5). D-F. Comparison of WT and *tsr* mutant responses to L-Ser or indole. E. Shows a quantification of the relative number of cells in the field of view over time following treatment with 5 mM indole for a competition experiment with WT and *tsr* (representative image shown in F). Data were collected at 30 °C. G-H. Isothermal titration calorimetry (ITC) experiments with 50 µM *S*. Typhimurium Tsr ligand-binding domain (LBD) and indole, or with L-Ser in the presence of 500 µM indole. Data are means and error bars are standard error of the mean (SEM, n=3-5). AUC indicates area under the curve.

indole is sensed through Tsr in *Salmonella* has not been elucidated. We recently reported the first

crystal structure of *S*. Typhimurium Tsr LBD, which clearly defines how the binding site

recognizes the L-Ser ligand (PDB code: 8fyv), and we thought it unlikely indole can be

accommodated at the same site [36]. To our knowledge, no prior study has tested whether the Tsr

LBD binds indole directly, so we expressed and purified the LBD, corresponding to the soluble

periplasmic portion, and performed isothermal titration calorimetry (ITC). These data show that

no binding occurs between the Tsr LBD and indole (Fig. 6G).

We next wondered if indole acts as an allosteric regulator of the LBD, possibly through

interacting with the L-Ser-bound form or interfering with L-Ser recognition. To address these

possibilities, we performed ITC of 50 µM Tsr LBD with L-Ser in the presence of 500 µM indole

and observed a robust exothermic binding curve and K_D of 5 µM, identical to the binding of L-

Ser alone, which we reported previously (Fig. 6H) [36]. These data indicate that indole does not

alter the Tsr LBD affinity for L-Ser. We conclude that Tsr senses indole through an atypical

mechanism, which might either involve regulation through a solute-binding protein [8,74],

responsiveness to perturbation in the proton motor force [18], or binding to a different region other

than the periplasmic LBD. Our findings reveal that while indole acts as a chemorepellent for *S*.

Typhimurium in isolation, and is sensed through Tsr, its presence within fecal material mixed

with other effectors is insufficient to elicit chemorepulsion.

Compromising between conflicting effector signals through chemohalation

Since Tsr mediates both chemoattraction to L-Ser and chemorepulsion from indole, we wondered

at what threshold each response dominates, and how this behavior is regulated at the point of

transition. To address these questions, we assessed responses to physiological mixtures of these

effectors using 500 µM L-Ser and increasing concentrations of indole at L-Ser:indole molar ratios

of 10:1, 1:1, or 1:10 (Fig. 7A-D, Movie 11). These experiments reveal a fascinating transition in

the distribution of the pathogen population as a function of increasing chemorepellent, which

occurs within minutes of exposure (Fig. 7A-D, Movie 11).

With only chemoattractant present, the bacterial population organizes tightly around the

effector source (Fig. 7A, Movie 11). When indole is introduced at a concentration 10-fold lower

than L-Ser, the bacterial distribution still exhibits chemoattraction but becomes more diffuse

(Fig. 7B, Movie 11). At a 1:1 ratio of chemoattractant and chemorepellent, a different population

structure emerges, in which the swimming bacteria are attracted toward the source but form a
halo around the treatment with an interior region of avoidance (Fig. 7C, Fig. 7E, Movie 11).
When the concentration of indole is 10-fold higher than L-Ser, the bacteria exhibit a wider zone
of avoidance (Fig. 7D-E, Movie 11). Interestingly, whereas 5 mM indole on its own induces
strong chemorepulsion (Fig. S3I, Movie 11), the addition of 10-fold lower L-Ser effectively
converts the behavior to a null response (Fig. 7D-E, Movie 11). This demonstrates that even at
the highest concentrations of indole *S.* Typhimurium might encounter in the gut, the presence of
chemoattractant can override indole chemorepulsion.

The intermediate responses to opposing effector mixtures bear similarities to CIRA
experiments with fecal material, some of which also exhibited a halo-like structure around the
treatment source (Fig. 3, Movie 2, Movie 3). Previous studies have also observed responses that
are an intermediate behavior between chemattraction and chemorepulsion, and have been
referred to by a variety of names [9,14,15]. There exists no consensus descriptor for taxis of this
nature, and so we suggest expanding the lexicon with the term "chemohalation," in reference to
the halo formed by the cell population, and which is congruent with the commonly used terms
chemoattraction and chemorepulsion. We expect chemohalation is a compromise between the
chemoattraction driven by L-Ser and the chemorepulsion driven by indole. Across these
experiments, the interior zone of avoidance roughly corresponds to where the local concentration
of indole is calculated to exceed 10 nM (Fig. 4E-F).

L-Ser enables resilience to indole-mediated growth inhibition

Fig. 7. *S.* Typhimurium mediates distinct chemotactic responses based on the ratio of L-Ser to indole. A-D. Representative max projections of responses to treatments of L-Ser and indole at 295-300 s, as indicated. E. Relative bacterial distribution in response to treatments of 500 µM L-Ser and varying amounts of indole, from panels A-D, with the mean value normalized to 100%. Data were collected at 30 °C. Data are means and error bars are standard error of the mean (SEM, n=3-5). F. Diffusion modeling of local effector concentrations based on sources of 5 mM indole (dark brown), 500 µM L-Ser (blue), 500 µM indole (light brown), and 50 µM indole (yellow) are shown as dashed lines. The approximate local concentration of indole that elicits a transition in chemotactic behavior is highlighted in light blue. G-H. Bacterial growth as a function of L-Ser or indole, at the time point where the untreated culture reaches A_{600} of 0.5. I-J. Bacterial growth +/- pretreatment with 500 µM indole or L-Ser, and increasing concentrations of indole or L-Ser, as indicated at the time point where the untreated culture reaches A_{600} of 0.5. Data are means and error bars are standard error of the mean (SEM, n=8-24). See also Movie 11.

We questioned why non-typhoidal *Salmonella* are attracted to a biological solution with high
amounts of indole, a chemical reported to inhibit bacterial growth [17,75,76]. We examined how
growth is affected by 0-25 mM indole or L-Ser in a background of minimal media (MM,
Materials & Methods). As expected, increasing amounts of the nutrient L-Ser provide a growth
advantage for all *Salmonella* strains analyzed, with maximal benefit achieved by approximately
500 µM (Fig. 7G). Equivalent treatments with indole show tolerance up to approximately 1 mM,
with growth inhibition occurring in the 1-5 mM range and lethality occurring at indole
concentrations greater than 5 mM (Fig. 7H). However, adding L-Ser in a background of 500 µM
indole provides only a small growth enhancement (Fig. 7I), and addition of 500 µM L-Ser
increases tolerance for indole up to about 1 mM, above which indole toxicity is unavoidable
(Fig. 7J). It appears that the relative attraction to combinations of these effectors relates to their
propensity to enhance or inhibit growth, with more permissive conditions eliciting a greater
degree of chemoattraction. Overall, the bacteria still obtain growth benefits from L-Ser so long as
the concentration of indole is under 1 mM.

Discussion

Bacteria in the human gastrointestinal tract encounter complex chemical landscapes that contain

both chemoattractants and chemorepellents [4,28,36]. However, chemotaxis responses are often

studied in isolation, outside of their biological and ecological contexts, which can lead to an

over- or underestimation of the roles specific interactions play in natural settings. In the present

work, we contribute to an emerging understanding of how bacteria navigate conflicting

chemotaxis stimuli and relate these chemotactic compromises to enteric infection and pathogen

growth [9,15,77].

In this study, we show that despite the microbial metabolite indole being a strong

chemorepellent in isolation (Fig. 6), fecal indole is insufficient to elicit pathogen

chemorepulsion, i.e. pathogens do not swim away from fecal material, or protect against cellular

invasion in an explant model (Fig. 1, Fig. 2, Fig. 3, Fig. 4). Instead, it appears pathogens employ

indole taxis as a means to regulate the magnitude of their attraction toward sources of intestinal

nutrients (Fig. 5). *In vivo*, we expect that the bacteria are attracted to indole-rich fecal material,

and it is simply a matter of the degree of attraction and which sites are prioritized among those

accessible to the invading pathogen. This finding revises our understanding of indole taxis during

enteric infection, suggesting that, rather than impairing pathogen infection as others have

proposed [8,23], indole chemorepulsion serves a useful function for pathogens and enables them to

integrate information about local microbial competitors into their chemotaxis responses. This, in

turn, allows pathogens to prioritize niches with abundant nutrients and reduced microbial

competition.

Interpretations of explant infections and the functions of chemotaxis and Tsr

Our explant experiments can be thought of as testing whether a layer of effector solution is

permissive to pathogen entry to the intestinal mucosa, and whether chemotaxis provides an

advantage in transiting this chemical gradient to associate with, and invade, the tissue (Fig. 1C,

Fig. S5). This behavior is probabilistic, and given sufficient time even chemotactic-deficient

cells will contact the tissue. This is reflected in that all treatments showed substantial infection

by all strains in terms of absolute CFUs isolated from homogenates (Fig. S2). If we compare the

probability of chemotaxis-mediated transit of the effector gradients we tested, greatest is for fecal

treatment, which among all treatments showed the highest degree of intracellular invasion (at 3 h

post-infection, Fig. 1). Then, buffer, L-Ser, and L-Ser + indole treatments are similarly

permissive and chemotaxis enhances infection in these backgrounds as well (Fig. 1, Fig. S5).

That chemotaxis provides an advantage in the buffer treated background, without any added

chemoeffector, is interesting and could simply be from effectors emitted from the host tissue

(Fig. S5). For instance, there is evidence that intestinal tissue, and host cells more broadly, can

release L-Ser and other amino acids, particularly in the context of tissue injury which could be

caused by *Salmonella* epithelial invasion in these experiments [1,36,59,78]. Altogether, chemotaxis

enhances the transit of the effector gradients mentioned above to access the host tissue (Fig. S5).

The explant system offers new insights into whether indole is protective against pathogen

infection. First, indole treatment does negate the infection advantage conferred by chemotaxis,

which was a unique effect among the treatments we tested (Fig. 1, Fig. S5). This is an interesting

result, somewhat mirroring what others have seen in cell culture [23], and indicates that the indole

gradient does not increase the likelihood of transit for chemotactic cells. In assessing the total

bacteria isolated from homogenates we see no evidence that indole protects against infection,

since the bacteria counts are prevalent and similar to other treatments, though this could be

different at lower multiplicities of infection (Fig. S2). Second, this effect is only observed with

indole as the sole effector, but not when the same concentration of indole is present within fecal

material, as it exists in the intestinal environment, or co-treatment with the fecal effector L-Ser

(Fig. 1, Fig. S5). Thus, the loss of the chemotactic advantage observed with indole treatment is

reversed in the presence of chemoattractant stimuli, suggesting that this effect is unlikely to

occur *in vivo*. It is also worth noting that the residual effector concentrations experienced by the

bacteria in the explant experiments were very low (Fig. 1C), and so we do not think the effects

we see are due to impacts on bacterial growth. It is unclear whether there would ever be a

situation *in vivo* where indole is the dominant effector, and so the behavior of bacteria swimming

away from a source of pure indole may be somewhat artificial (Fig. 6). These data, overall, do

not support indole chemorepulsion as a mechanism of colonization resistance against pathogens,

although indole is known to reduce virulence through other mechanisms [8,22,23,75].

New insights into indole taxis from non-E. coli systems

Indole is a key regulator of enteric microbial communities, known to modulate motility and
virulence, and is highly abundant in fecal matter due to the metabolic activity of the microbiota
[8,16–18,21–24]. While *E. coli* has served as an important model system for elucidating the
mechanisms of indole chemorepulsion (Table S1) [8,18,23,25], no prior work has examined how
indole sensing is integrated alongside multiple other intestinal effectors, nor whether these
behaviors are conserved across clinical isolates of disease-causing species. Here, we address
these gaps by providing confirmatory evidence for some earlier predictions and evidence that
challenges others, refining our understanding of how indole influences pathogen behavior
through chemotaxis within the intestinal environment.

Perhaps our most striking finding is that fecal material, the native reservoir of the strong
chemorepellent indole, does not elicit chemorepulsion in the form of bacteria swimming away
(Fig. 2, Fig. 3, Fig. 4). Instead, a representative panel of diverse *Enterobacteriaceae* pathogens
and pathobionts exhibited fecal attraction (Fig. 3, Fig. 4), demonstrating that enteric species
associated with disease are undeterred by indole in its natural context, i.e., when mixed with
other fecal chemoattractants. Through analyses showing that indole chemorepulsion is easily
overridden by the presence of intestinal nutrients we surmise that the benefits associated with
fecal material typically outweigh the deleterious effects of indole (Fig. 5, Fig. 7). This conclusion
is further supported by growth analyses indicating that *Salmonella* tolerates indole when
sufficient nutrients are available (Fig. 7).

We used *Salmonella* Typhimurium as a model to dissect the mechanisms underlying fecal
attraction and indole sensing. Regarding the latter, our findings largely confirm previous studies
in *E. coli*, showing that the chemoreceptor Tsr is required for indole taxis (Fig. 6, Table S1)
[8,18,25]. However, we do add some new dimensions to understanding indole taxis. First, for *E. coli*
the involvement of Tar in indole sensing has been reported [8], but we see no equivalent function
for *S.* Typhimurium. Yet, we previously showed *S.* Typhimurium WT and *tsr* are both readily
attracted to L-Asp, supporting the presence of a functional Tar under the same experimental
conditions as we test here (Fig. 6) [36]. We do not know the reason for this outcome, but note
different assays were used, and this could also reflect variation between the chemotaxis systems
of these two bacteria. Second, while Tsr serves as the sensor for indole in *Salmonella*, it is also a
key mediator of fecal attraction through its role in sensing L-Ser, which is also abundant in fecal
material (Fig. 1, Fig. 2, Fig. 3, Fig. 6). Third, we are the first to visualize and quantify the rapid

temporal dynamics of indole chemorepulsion (Fig. 5, Fig. 6, Fig. S3I; Movie 11). For responses
to pure indole a clear zone of avoidance around the treatment appears within 10 seconds of
exposure, much faster than chemoattraction to L-Ser, suggesting the cells have the ability to
rapidly flee deleterious conditions (Fig. 6, Fig. S3I; Movie 11). Lastly, we also investigated
whether indole sensing occurs through the canonical chemoreceptor mechanism of direct binding
to the Tsr ligand-binding domain (LBD). Our data shows it does not, nor does indole antagonize
or inhibit L-Ser binding to the LBD (Fig. 6). While these findings do not resolve the molecular
mechanism of indole sensing, they eliminate two plausible models that, to our knowledge, have
not been previously tested. Overall, our data support the hypothesis that Tsr employs a non-
canonical mechanism to sense indole [8,18,74].

Having confirmed the role of Tsr in mediating indole chemorepulsion in *S.* Typhimurium,
and shown it to function similarly as in *E. coli* [8,9,14], and having demonstrated that diverse
Enterobacteriaceae are attracted to indole-rich fecal material, we expect that the behaviors
described here are representative of the many enteric species that possess Tsr orthologues, which
we mapped in a previous study [36]. As we report here, there does seem to be a large variety in the
magnitude of fecal attraction by different 'wild' enteric pathogens, which could reflect
adaptations to different host intestinal environments and microbiota communities and may
influence pathogenesis (Fig. 3, Fig. 4). While foundational insights into indole taxis have come
from model bacterial systems, continued progress in understanding the role of chemotaxis in
human disease will benefit from extending such analyses to a broader range of clinically relevant
bacterial species and strains.

Function of indole taxis in enteric invasion
In the context of non-typhoidal *Salmonella* infections, it is clear there exists complex
relationships between chemotactic sensing of effectors, bacterial growth, and invasion (Fig. S5).
In addition to the factors we have investigated, it is already well-established in the literature that
the vast metabolome in the gut contains many chemicals that modulate *Salmonella* cellular
invasion, virulence, growth, and pathogenicity [79–81]. As it pertains specifically to sensing the
opposing effectors L-Ser and indole, we propose that Tsr directs bacteria toward the highest ratio
of attractant to repellent accessible in the local environment, with fine-tuning of navigation
occurring through regulation of the magnitude of attraction and chemohalation. In addition to

sensing these two effectors, our data indicate that fecal attraction involves other stimuli,

including L-Asp sensing through Tar and sugar sensing through Trg (Fig. 5). Ultimately, the dual

sensing of opposing effectors by Tsr serves to improve pathogen fitness through colonizing

niches rich in nutrients, signaled by local L-Ser concentrations, and seeking niches with low

microbial competition, indicated by local indole concentrations.

Navigating contradictory stimuli in nature

The scenario we investigated here of *S.* Typhimurium encountering high concentrations of

opposing chemotactic stimuli in the intestinal environment is just one example of the complex

chemical landscapes that bacteria navigate in nature. To better understand the "decision-making"

process underlying chemotaxis in the presence of conflicting effectors, we examined

physiological mixtures of the fecal metabolites indole and L-Ser and recorded a series of real-

time videos capturing behavioral transitions as a function of effector concentration (Fig. 7;

Movie 11). These videos reveal that, upon sensing conflicting stimuli, the bacterial population

structure rapidly evolves based on the attractant-to-repellent ratio, displaying a spectrum of

behaviors: chemoattraction, diffuse chemoattraction, chemohalation, diffuse chemohalation, and

chemorepulsion (Fig. 7; Movie 11).

These dynamic, micron-scale chemohalation patterns reflect a behavioral compromise

between attraction and repulsion and would be difficult or impossible to detect without live

imaging, which may explain why they were previously unappreciated in binary models of

chemotaxis [11]. In fact, many chemotaxis assays that use indirect methods of quantification, such

as growth, would not be able to distinguish between chemoattraction and chemohalation since

they both involve in increase in bacteria overt time. To be clear, we suggest chemohalation as a

new term to generally describe intermediate chemotaxis responses to conflicting stimuli that are

neither chemoattraction nor chemorepulsion, but others have contributed to studying how

chemotaxis functions in confounding chemical landscapes [9,14,15]. The chemohalation responses

reported here most closely resemble the "trade-off" response previously described in *E. coli*

exposed to attractant–repellent mixtures [9]. Interestingly, that study also described a "bet-

hedging" response, in which a subpopulation remained attracted despite the presence of a

chemorepellent, but we did not observe this behavior in our system [9].

In addition to our work here, there are other examples of chemohalation responses to
complex biological stimuli of the gastrointestinal environment. Recently, we reported on
Enterobacteriaceae chemotactic sensing of blood serum, which bacteria encounter during
intestinal bleeding events, and those responses appear to involve chemohalation [36].
Chemohalation is also seen in the case of the gastric pathogen *Helicobacter pylori* responding to
mixtures of urea, a chemoattractant, and acid, a chemorepellent, conflicting effectors it
encounters near the stomach mucosa [15,82]. The functional significance of chemohalation remains
to be understood, but could be a method of fine-tuning colonization bias such that nutrients can
be acquired while not approaching too closely to a deleterious stimulus. Continuing to
investigate chemohalation behaviors and understanding how they coordinate bacterial
colonization may provide important insights into how chemotaxis confers fitness advantages in
natural environments.
520
Limitations of this study
This study provides insights into the roles of chemotaxis in directing the behaviors of
Enterobacteriaceae species in response to fecal material and indole, however, several limitations
should be considered. First, our analyses of pathogen enteric infection were performed using
swine colonic explants, which do not fully recapitulate the complexity of *in vivo* infection
dynamics in the human gut. While explant assays offered insights into the relationship between
chemotaxis and tissue colonization, these experiments exhibited variability. To mitigate this, we
used multiple tissue sections from a single animal to improve experimental consistency.
However, this approach limits our ability to assess how inter-host variability might influence
bacterial responses. Future studies using distal ileum tissue, a major site of *S.* Typhimurium
cellular invasion known to contain distinct chemical features, may provide further insight into
the functions of indole taxis during infection [42]. Another experimental limitation is the difference
in timescales between our assays. Chemotaxis experiments were conducted over approximately
5-minutes, whereas tissue explant experiments required several hours for significant differences
in colonization and cellular invasion to be observed. Thus, there are effects from chemotactic
adaptation and replication that we do not elucidate here. Lastly, while we confirmed that non-
typhoidal *Salmonella* are attracted to human fecal material, we only determined the dependency
on Tsr, and sensing of L-Ser, in our model strain (IR715). Although we predict that

Enterobacteriaceae also use Tsr for fecal attraction, this remains uncertain without targeted

genetic analyses in each strain background.

Materials & Methods

All methods were carried out in accordance with relevant guidelines, regulations, and state and federal law. Experimental protocols were approved by the Institutional Biosafety Committee (IBC) of Washington State University (#1372).

Bacterial strains and growth conditions

Bacterial strains and plasmids used in this study are listed in Table 1. As previously described [36], bacteria intended for chemotaxis assays were grown overnight in tryptone broth (TB) with antibiotic selection, as appropriate. Motile bacteria were prepared with a 1:1000 back-dilution and grown shaking for approximately 4 hours at 37° C to reach A_{600} of 0.5. Cells were centrifuged, washed, and resuspended in a chemotaxis buffer (CB) containing 10 mM potassium phosphate (pH 7), 10 mM sodium lactate, and 100 μM EDTA to A_{600} of 0.2 and rocked gently at the temperatures indicated in figure legends until fully motile, typically 1-2 h. For *in vitro* growth analyses, cultures were grown overnight in Lysogeny Broth (LB) at 37° C. Subsequently, 5 μl of A_{600} 2.0 cells were used to inoculate 200 μl of minimal media (MM), containing 47 mM Na_2HPO_4, 22 mM KH_2PO_4, 8 mM NaCl, 2 mM $MgSO_4$, 0.4% glucose (w/v) 11.35 mM $(NH_4)_2SO_4$, 100 μM $CaCl_2$ and L-Ser and/or indole at the described concentrations, and cultured in a 96-well microtiter plate. Cultures were grown at 37° C and monitored by A_{600} readings at 5-minute intervals.

Table 1: Bacterial Strains

Strain	Reference/Source
S. enterica Typhimurium IR715 nalidixic acid-resistant derivative of ATCC 14028	Rivera-Chávez, F. et al.[30]
S. enterica Typhimurium IR715 Δ*cheY::Tn10* (Tet^R)	Rivera-Chávez, F. et al.[30]
S. enterica Typhimurium IR715 Δ*tsr::pFR3* (Cm^R)	Rivera-Chávez, F. et al.[30]
S. enterica Typhimurium IR715 invA::pSW127 (Carb^R)	Thiennimitr, P. et al.[57]
S. enterica SARA1	Beltran, P. et al.[67]
S. enterica Newport M11018046001A	Shariat, N. et al.[68]
S. enterica Enteriditis 05E01375	Shariat, N. et al.[68]
E. coli BL21-DE3 Cat# 70954-3	Millipore-Sigma
E. coli MG1655 pXS-GFP (Amp^R)	This study
Enterobacter cloacae subsp. *cloacae* strain CDC 442-68 Cat# 13047	ATCC
Citrobacter koseri Frederiksen 4225-83 Cat# BAA-895	ATCC

 Escherichia coli strain NCTC 9001 Cat# 11775   ATCC

Chemosensory injection rig assay (CIRA)

CIRA was performed as described previously [36]. Briefly, an Eppendorf Femtotip 2

microcapillary containing the treatment of interest was lowered into a pond of 50 µl of motile

cells using a Sutter micromanipulator. An injection flow of effector into the pond at

approximately 300 fl per minute was achieved using a Femtojet 4i set to P_c 35. Solubilized fecal

treatments were prepared by dissolving 1 g of commercially obtained human feces (Lee

Biosolutions) in 10 ml of CB. The solution was clarified by centrifugation at 10,000 g for 20

minutes, followed by sterile filtration through a 0.2 µm filter. Treatment solutions of indole and

L-Ser were also diluted into CB and sterile-filtered before application. Videos were recorded

using an inverted Nikon Ti2 microscope with heated sample chamber at 37 ˚C.

CIRA microgradient modeling

Modeling the microgradient generated through CIRA was performed as described earlier [36],

based on the continual injection and diffusion of an effector from a fixed-point source. Briefly,

diffusion is modeled as a 3D process where the diffusing substance is gradually and continuously

introduced at a fixed point within a large surrounding fluid volume. The substance is prepared at

a concentration of M_s (typically between 0.5 µM and 5 mM) and injected at a volume rate of $Q =$

305.5 fl/min. The species then diffuses into the ambient fluid with a diffusion constant D:

$$C(r,t) = \frac{q}{4\pi Dr} erfc \frac{r}{2\sqrt{Dt}}$$

Here, r is the distance from the point source, t is the time from initial injections, q is the injection

rate of the species (equal to M_sQ), and C is the species concentration. In earlier work [36], we

reported using fluorescent dye that the concentrations predicted by this model appear to be

accurate within 5% in the range of 70-270 µm from the source, whereas at distances less than 70

µm the measured concentrations are about 10% lower than predicted. At the point where the

effector treatment is injected into the larger volume the local concentration drops precipitously,

hence why the concentration reported at distance 0 is not that of the concentration within the

microcapillary.

Isothermal titration calorimetry ligand binding studies (ITC)

592 Purification of *S*. Typhimurium Tsr LBD was performed as described previously [36]. ITC

593 experiments were performed using a Microcal ITC200 instrument (GE Healthcare). Either 500

594 μM indole or L-Ser was titrated in 2.5 μL injections into a 200 μL sample cell containing 50 μM

595 Tsr LBD. For the indole/L-Ser competition experiment, 500 μM indole was added to both the

596 titrant and sample cell, thus providing a constant excess background concentration of indole. For

597 all experimental conditions, blank titrations were also collected in which indole or L-Ser was

598 titrated into a cell containing buffer alone. All experiments were performed using thoroughly

599 degassed samples at 25 °C in 50 mM Tris, 150 mM NaCl, 1 mM EDTA, pH 7.5. The reference

600 power was set to 5 μcal/sec. The resulting power curves were integrated using the Origin

601 analysis software included with the instrument. The heat of dilution was subtracted from each

602 point using the blank. A single-site binding model was then fit to the data, floating parameters

603 describing the binding enthalpy (ΔH), equilibrium constant (K_D), and apparent binding

604 stoichiometry (n). The instrument software was used for this purpose.

606 *Quantification of indole and serine in human fecal samples*

607 Solubilized human feces was prepared as described above for CIRA and analyzed by mass

608 spectrometry to determine the molar serine content as a service through the University of

609 Washington Mass Spectrometry Center. This measurement reflects total serine, of which close to

610 100% is expected to be L-Ser [36]. As described in earlier work, the indole content of solubilized

611 human fecal samples was determined using a hydroxylamine-based calorimetric assay with

612 purified indole as a reference and standard [83].

614 *Explant infection assays*

615 Swine intestinal tissue was acquired from the descending colon of an 8-week-old animal,

616 pursuant to animal protocol ASAF #7128, approved through the Washington State University

617 IACUC. Before infection, an approximately 20 by 20 mm piece of swine intestinal explant tissue

618 was gently washed with PBS to remove fecal matter. Next, the tissue section was bathed in

619 chemoeffector solution (solubilized human fecal matter (Lee Biosolutions), a mixture of 338 μM

620 L-Ser and 862 μM indole, 338 μM L-Ser alone, 862 μM indole alone, or chemotaxis buffer) in a

621 6-well tissue culture plate (Celltreat) and incubated at 4° C for 1 h. Then, tissue was transferred

622 to a 35 mm Mattek dish where the luminal side of the tissue was exposed to a bacterial solution

containing a 1:1 mixture (~10^9 CFU each) of WT *S*. Typhimurium IR715 and either the isogenic

tsr or *cheY* mutant, suspended in CB at a volume of 300 µl. The tissue was then incubated in the

dish with the competing bacteria at 37 °C and 5% CO_2 for 1, 3, or 6 h. After, half of the tissue

was transferred into screwcap tubes containing 500 µl LB media and 5-10 2.3 mm zirconia beads

(BioSpec Products) on ice and homogenized using a Bead Mill 24 (Fisher Scientific) at 6.5 m/s

for 60 s, repeated four times. To enumerate the "invaded" bacteria, the other half of the tissue

was washed in PBS and incubated in PBS containing 100 µg/ml gentamicin for 1 h at 37 °C and

5% CO_2, then washed twice in PBS, as done previously [56,84,85]. The homogenization process was

then repeated for the gentamicin-treated tissue. CFUs were enumerated by plating 10-fold

dilutions on LB agar plates containing the appropriate antibiotic [56,86]. Competitive index values

were calculated by dividing the number of mutant CFUs by the number of WT CFUs for each

treatment and time point [87,88].

Quantification of CIRA data

Videos of chemotactic responses were quantified as described previously [36]. The number of cells

in each frame were calculated by determining a fluorescence intensity ratio per cell for frames

pre-treatment and extrapolated using the 'plot profile' function of ImageJ. The distribution of the

bacteria was calculated using the Radial Profile ImageJ plugin. Local background subtraction

was performed based on experiments with the non-chemotactic *cheY* strain to control for

autofluorescence in solubilized fecal samples.

Statistical Analyses

Competitive indices (CIs) for explant experiments were calculated for each treatment group at

each time point. Log-transformed CI values were obtained by taking the logarithm (log_{10}) of the

original CI measurements. These log-transformed values were then subjected to statistical

analysis. First, a one-sample t-test was performed to determine whether the mean of the log-

transformed CIs significantly differed from zero. In cases where the assumption of normality was

violated, the non-parametric Wilcoxon rank sum test was applied as an alternative. Effect size

was assessed using Cohen's *d* and calculated using the same log-transformed CIs. To determine

p-values between total and invaded populations at 3 and 6 h and for comparing relative bacteria

% within 500 µm of the treatment source in Figs. 5-6, unpaired t-tests were employed.

Acknowledgments

Funding for this research was provided by NIAID through awards 1K99AI148587 and 4R00AI148587-03, and funding from the College of Veterinary Medicine at Washington State University to AB. Bacterial strains were provided by Nikki Shariat (University of Georgia, Athens), Nkuchia Mikanatha and Pennsylvania NARMS and GenomeTrakr Programs, and Andreas Bäumler (University of California, Davis). All research on human and animal samples was performed in accordance with, and approval of, the Institutional Biosafety Committee and Institutional Animal Care and Use Committee at Washington State University.

Author Contributions

K.F. performed the microgradient modeling and explant experiments. A.B. conducted the CIRA experiments and purification of Tsr-LBD. Z.G. performed bacterial growth experiments. M.S. and M.J.H. performed the ITC experiments. All authors contributed to data analyses and writing of the manuscript.

Declaration of Interests

A.B. owns Amethyst Antimicrobials, LLC.

Movie figure legends

Movie 1. Chemotactic response of *S.* Typhimurium IR715 to solubilized human feces. Representative CIRA experiments showing *S.* Typhimurium IR715 WT and mutant strains responding to a source over 300 s (shown at 10x speed) Viewable at: https://www.youtube.com/watch?v=BqUcRN3YwjU

Movie 2. Chemotactic response of *S.* Typhimurium IR715 WT and chemotactic mutant strains to solubilized human feces. Representative CIRA experiments showing competition between *S.* Typhimurium IR715 (mPlum) and *cheY*, or *tsr*, as indicated (GFP), over 300 s. Viewable at: https://www.youtube.com/watch?v=D5JL46b4lsI

684 **Movie 3.** Chemotactic response of *S. enterica* clinical isolates to solubilized human feces.
685 Representative CIRA experiments showing competition between *S.* Typhimurium IR715
686 (mPlum) and clinical isolates, as indicated (GFP), responding to a source of solubilized human
687 feces over 300 s. Viewable at: https://www.youtube.com/watch?v=dLsFDV0XgpY
688
689 **Movie 4.** Chemotactic response of *E. coli* MG1655 to solubilized human feces. Representative
690 CIRA experiment over 300 s. Viewable at:
691 https://youtube.com/shorts/WH6tabDbrw4?feature=share
692
693 **Movie 5.** Chemotactic response of *E. coli* NCTC 9001 to solubilized human feces.
694 Representative CIRA experiment over 300 s. Viewable at:
695 https://youtube.com/shorts/yzU2M4Z_Yf4?feature=share
696
697 **Movie 6.** Chemotactic response of *C. koseri* 4225-83 to solubilized human feces. Representative
698 CIRA experiments with treatment sources as indicated, over 300 s. Viewable at:
699 https://youtube.com/shorts/s_ybO0xcIDw?feature=share
700
701 **Movie 7.** Chemotactic response of *S.* Typhimurium IR715 to L-Ser and indole treatment at fecal-
702 relevant concentrations. Representative CIRA experiment over 300 s. Viewable at:
703 https://youtube.com/shorts/4UEYoBS6jIQ?feature=share
704
705 **Movie 8.** Chemotactic response of *S.* Typhimurium IR715 to complete mixture of fecal effectors.
706 Representative CIRA experiment over 300 s. Viewable at:
707 https://youtube.com/shorts/Yd14m3sI6Pw?feature=share
708
709 **Movie 9.** Chemotactic response of *S.* Typhimurium IR715 to mixture of fecal effectors lacking
710 L-Ser. Representative CIRA experiment over 300 s. Viewable at:
711 https://youtu.be/5QM116BrHhQ
712
Movie 10. Chemotactic response of *S.* Typhimurium IR715 to mixture of fecal effectors with
0.5x indole. Representative CIRA experiment over 300 s. Viewable at:

https://youtube.com/shorts/OqH0HE2rYIE?feature=share

Movie 11. Chemotactic response of *S.* Typhimurium IR715 to L-Ser and indole treatments.

Representative CIRA experiments with treatment sources as indicated, over 300 s. Viewable at:

https://www.youtube.com/watch?v=bNQMqF2QMek

720
721 **References**
722

1. Zhou, B., Szymanski, C. M. & Baylink, A. Bacterial chemotaxis in human diseases. *Trends*

in Microbiology **31**, 453–467 (2023).

2. Keegstra, J. M., Carrara, F. & Stocker, R. The ecological roles of bacterial chemotaxis.

Nature Reviews Microbiology (2022) doi:10.1038/s41579-022-00709-w.

3. Bi, S. & Sourjik, V. Stimulus sensing and signal processing in bacterial chemotaxis. *Current*

Opinion in Microbiology **45**, 22–29 (2018).

4. Matilla, M. A. & Krell, T. The effect of bacterial chemotaxis on host infection and

pathogenicity. *FEMS Microbiology Reviews* **42**, (2018).

5. Matilla, M. A., Gavira, J. A. & Krell, T. Accessing nutrients as the primary benefit arising

from chemotaxis. *Current Opinion in Microbiology* **75**, 102358 (2023).

6. Matilla, M. A. & Krell, T. Chemoreceptor-based signal sensing. *Current Opinion in*

Biotechnology **45**, 8–14 (2017).

7. Ortega, Á., Zhulin, I. B. & Krell, T. Sensory Repertoire of Bacterial Chemoreceptors.

Microbiol Mol Biol Rev **81**, e00033-17 (2017).

8. Yang, J. *et al.* Biphasic chemotaxis of *Escherichia coli* to the microbiota metabolite indole.

Proc. Natl. Acad. Sci. U.S.A. **117**, 6114–6120 (2020).

9. Livne, N. & Vaknin, A. Collective responses of bacteria to a local source of conflicting

effectors. *Sci Rep* **12**, 4928 (2022).

10. Keller, E. F. & Segel, L. A. Traveling bands of chemotactic bacteria: A theoretical analysis.

Journal of Theoretical Biology **30**, 235–248 (1971).

11. Adler, J. & Tso, W.-W. 'Decision'-Making in Bacteria: Chemotactic Response of

Escherichia coli to Conflicting Stimuli. *Science* **184**, 1292–1294 (1974).

12. Zhang, X. *et al.* Escape band in Escherichia coli chemotaxis in opposing attractant and

nutrient gradients. *Proc Natl Acad Sci U S A* **116**, 2253–2258 (2019).

13. Kalinin, Y., Neumann, S., Sourjik, V. & Wu, M. Responses of Escherichia coli Bacteria to

Two Opposing Chemoattractant Gradients Depend on the Chemoreceptor Ratio. *Journal of*

Bacteriology **192**, 1796 (2010).

14. Englert, D. L., Manson, M. D. & Jayaraman, A. Flow-Based Microfluidic Device for

Quantifying Bacterial Chemotaxis in Stable, Competing Gradients. *Appl Environ Microbiol*

75, 4557–4564 (2009).

15. Huang, J. Y., Sweeney, E. G., Guillemin, K. & Amieva, M. R. Multiple Acid Sensors

Control Helicobacter pylori Colonization of the Stomach. *PLOS Pathogens* **13**, e1006118

(2017).

16. Li, X., Zhang, B., Hu, Y. & Zhao, Y. New Insights Into Gut-Bacteria-Derived Indole and Its

Derivatives in Intestinal and Liver Diseases. *Front. Pharmacol.* **12**, 769501 (2021).

17. Lee, J.-H. & Lee, J. Indole as an intercellular signal in microbial communities. *FEMS*

Microbiol Rev **34**, 426–444 (2010).

18. Gupta, R. *et al.* Indole modulates cooperative protein–protein interactions in the flagellar

motor. *PNAS Nexus* **1**, pgac035 (2022).

19. Darkoh, C., Chappell, C., Gonzales, C. & Okhuysen, P. A Rapid and Specific Method for the

Detection of Indole in Complex Biological Samples. *Appl Environ Microbiol* **81**, 8093–8097

(2015).

20. Chappell, C. L. *et al.* Fecal Indole as a Biomarker of Susceptibility to Cryptosporidium

Infection. *Infect Immun* **84**, 2299–2306 (2016).

21. Ye, X. *et al.* Dual Role of Indoles Derived From Intestinal Microbiota on Human Health.

Front. Immunol. **13**, 903526 (2022).

22. Kumar, A. & Sperandio, V. Indole Signaling at the Host-Microbiota-Pathogen Interface.

mBio **10**, e01031-19 (2019).

23. Kohli, N. *et al.* The microbiota metabolite indole inhibits Salmonella virulence: Involvement

of the PhoPQ two-component system. *PLOS ONE* **13**, e0190613 (2018).

24. Nikaido, E. *et al.* Effects of indole on drug resistance and virulence of Salmonella enterica

serovar Typhimurium revealed by genome-wide analyses. *Gut Pathog* **4**, 5 (2012).

25. Bansal, T. *et al.* Differential Effects of Epinephrine, Norepinephrine, and Indole on

Escherichia coli O157:H7 Chemotaxis, Colonization, and Gene Expression. *Infect Immun*

75, 4597–4607 (2007).

26. Bai, X. *et al.* Engineering the gut microbiome. *Nat Rev Bioeng* **1**, 665–679 (2023).

27. Wolter, M. *et al.* Leveraging diet to engineer the gut microbiome. *Nat Rev Gastroenterol*

Hepatol **18**, 885–902 (2021).

28. Human Metabolome Database. https://hmdb.ca/.

29. Rogers, A. W. L., Tsolis, R. M. & Bäumler, A. J. Salmonella versus the Microbiome.

Microbiology and Molecular Biology Reviews **85**, e00027-19 (2020).

30. Rivera-Chávez, F. *et al.* Salmonella Uses Energy Taxis to Benefit from Intestinal

Inflammation. *PLoS Pathog* **9**, e1003267 (2013).

31. Rivera-Chávez, F. *et al.* Energy Taxis toward Host-Derived Nitrate Supports a Salmonella

Pathogenicity Island 1-Independent Mechanism of Invasion. *mBio* **7**, e00960-16 (2016).

32. Gül, E. *et al.* Salmonella T3SS-2 virulence enhances gut-luminal colonization by enabling

chemotaxis-dependent exploitation of intestinal inflammation. *Cell Rep* **43**, 113925 (2024).

33. Winter, S. E., Lopez, C. A. & Bäumler, A. J. The dynamics of gut‑associated microbial

communities during inflammation. *EMBO Reports* **14**, 319–327 (2013).

34. Winter, S. E. *et al.* Gut inflammation provides a respiratory electron acceptor for Salmonella.

Nature **467**, 426–429 (2010).

35. Cooper, K. G. *et al.* Regulatory protein HilD stimulates Salmonella Typhimurium

invasiveness by promoting smooth swimming via the methyl-accepting chemotaxis protein

McpC. *Nat Commun* **12**, 348 (2021).

36. Glenn, S. J. *et al.* Bacterial vampirism mediated through taxis to serum. *eLife* **12**, RP93178

(2024).

37. Piñas, G. E., DeSantis, M. D., Cassidy, C. K. & Parkinson, J. S. Hexameric rings of the

scaffolding protein CheW enhance response sensitivity and cooperativity in Escherichia coli

chemoreceptor arrays. *Science Signaling* **15**, eabj1737 (2022).

38. Burt, A. *et al.* Complete structure of the chemosensory array core signalling unit in an E. coli

minicell strain. *Nat Commun* **11**, 743 (2020).

39. Chen, X., Bi, S., Ma, X., Sourjik, V. & Lai, L. Discovery of a New Chemoeffector for

Escherichia coli Chemoreceptor Tsr and Identification of a Molecular Mechanism of

Repellent Sensing. *ACS Bio Med Chem Au* **2**, 386–394 (2022).

40. Vega, N. M., Allison, K. R., Samuels, A. N., Klempner, M. S. & Collins, J. J. *Salmonella*

typhimurium intercepts *Escherichia coli* signaling to enhance antibiotic tolerance. *Proc. Natl.*

Acad. Sci. U.S.A. **110**, 14420–14425 (2013).

41. Roager, H. M. & Licht, T. R. Microbial tryptophan catabolites in health and disease. *Nat*

Commun **9**, 3294 (2018).

42. Chowdhury, R., Pavinski Bitar, P. D., Chapman, H. M. & Altier, C. Salmonella Invasion Is

Controlled by Competition among Intestinal Chemical Signals. *mBio* **14**, e00012-23 (2023).

43. Ehrhardt, K. & Grassl, G. A. Mouse Model to Study Salmonella-Induced Colitis. *Methods*

Mol Biol **2427**, 201–213 (2022).

44. Grassl, G. A., Valdez, Y., Bergstrom, K. S. B., Vallance, B. A. & Finlay, B. B. Chronic

enteric salmonella infection in mice leads to severe and persistent intestinal fibrosis.

Gastroenterology **134**, 768–780 (2008).

45. Radlinski, L. C. *et al.* Salmonella virulence factors induce amino acid malabsorption in the

ileum to promote ecosystem invasion of the large intestine. *Proceedings of the National*

Academy of Sciences **121**, e2417232121 (2024).

46. Mandal, B. K. & Mani, V. COLONIC INVOLVEMENT IN SALMONELLOSIS. *The*

Lancet **307**, 887–888 (1976).

47. Kronsteiner, B. *et al.* Helicobacter pylori Infection in a Pig Model Is Dominated by Th1 and

Cytotoxic CD8 [+] T Cell Responses. *Infect Immun* **81**, 3803–3813 (2013).

48. Smith, A. C. & Swindle, M. M. Preparation of Swine for the Laboratory. *ILAR Journal* **47**,

358–363 (2006).

49. Poutahidis, T. *et al. Helicobacter pylori* -induced Gastritis in Experimentally Infected

Conventional Piglets. *Vet Pathol* **38**, 667–678 (2001).

50. Boyen, F. *et al.* Porcine *in vitro* and *in vivo* models to assess the virulence of *Salmonella*

enterica serovar Typhimurium for pigs. *Lab Anim* **43**, 46–52 (2009).

51. Boyen, F. *et al.* Coated fatty acids alter virulence properties of Salmonella Typhimurium and

decrease intestinal colonization of pigs. *Veterinary Microbiology* **132**, 319–327 (2008).

52. Köppen, K. *et al.* Ex vivo infection model for Francisella using human lung tissue. *Front.*

Cell. Infect. Microbiol. **13**, (2023).

53. Bomjan, R., Zhang, M. & Zhou, D. YshB Promotes Intracellular Replication and Is Required

for Salmonella Virulence. *Journal of Bacteriology* **201**, 10.1128/jb.00314-19 (2019).

54. Birhanu, B. T., Park, N.-H., Lee, S.-J., Hossain, M. A. & Park, S.-C. Inhibition of

Salmonella Typhimurium adhesion, invasion, and intracellular survival via treatment with

methyl gallate alone and in combination with marbofloxacin. *Veterinary Research* **49**, 101

(2018).

55. MacBeth, K. J. & Lee, C. A. Prolonged inhibition of bacterial protein synthesis abolishes

Salmonella invasion. *Infect Immun* **61**, 1544–1546 (1993).

56. Sharma, A. & Puhar, A. Gentamicin Protection Assay to Determine the Number of

Intracellular Bacteria during Infection of Human TC7 Intestinal Epithelial Cells by Shigella

flexneri. *BIO-PROTOCOL* **9**, (2019).

57. Thiennimitr, P. *et al.* Intestinal inflammation allows *Salmonella* to use ethanolamine to

compete with the microbiota. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 17480–17485 (2011).

58. Haneda, T. *et al.* The Capsule-Encoding viaB Locus Reduces Intestinal Inflammation by a

Salmonella Pathogenicity Island 1-Independent Mechanism. *Infection and Immunity* **77**,

2932–2942 (2009).

59. Kitamoto, S. *et al.* Dietary l-serine confers a competitive fitness advantage to

Enterobacteriaceae in the inflamed gut. *Nature Microbiology* **5**, 116–125 (2020).

60. Sugihara, K. & Kamada, N. THE ROLE OF DIETARY L-SERINE IN THE REGULATION

OF INTESTINAL MUCUS BARRIER DURING INFLAMMATION. *Inflammatory Bowel*

Diseases **26**, S42–S42 (2020).

61. Perkins, A., Tudorica, D. A., Amieva, M. R., Remington, S. J. & Guillemin, K. Helicobacter

pylori senses bleach (HOCl) as a chemoattractant using a cytosolic chemoreceptor. *PLoS*

Biol **17**, e3000395 (2019).

62. Ranque, C. L. *et al.* Examination of Carbohydrate Products in Feces Reveals Potential

Biomarkers Distinguishing Exclusive and Nonexclusive Breastfeeding Practices in Infants. *J*

Nutr **150**, 1051–1057 (2020).

63. Ke, Z., Lu, Z., Li, Q. & Tong, W. Intestinal glucose excretion: A potential mechanism for

glycemic control. *Metabolism* **152**, 155743 (2024).

64. Larke, J. A. *et al.* Dietary Intake of Monosaccharides from Foods is Associated with

Characteristics of the Gut Microbiota and Gastrointestinal Inflammation in Healthy US

Adults. *The Journal of Nutrition* **153**, 106–119 (2023).

65. Shen, X. *et al.* Asparagine, colorectal cancer, and the role of sex, genes, microbes, and diet:

A narrative review. *Front Mol Biosci* **9**, 958666 (2022).

66. Yoo, W. *et al. Salmonella* Typhimurium expansion in the inflamed murine gut is dependent

on aspartate derived from ROS-mediated microbiota lysis. *Cell Host & Microbe* **32**, 887-

899.e6 (2024).

67. Beltran, P. *et al.* Reference collection of strains of the Salmonella typhimurium complex

from natural populations. *Journal of General Microbiology* **137**, 601–606 (1991).

68. Shariat, N. *et al.* Subtyping of Salmonella enterica Serovar Newport Outbreak Isolates by

CRISPR-MVLST and Determination of the Relationship between CRISPR-MVLST and

PFGE Results. *J Clin Microbiol* **51**, 2328–2336 (2013).

69. Ferrari, R. G. *et al.* Worldwide Epidemiology of *Salmonella* Serovars in Animal-Based

Foods: a Meta-analysis. *Appl Environ Microbiol* **85**, e00591-19 (2019).

70. Wiles, T. J. *et al.* Modernized Tools for Streamlined Genetic Manipulation and Comparative

Study of Wild and Diverse Proteobacterial Lineages. *mBio* **9**, e01877-18 (2018).

71. Falke, J. J. & Hazelbauer, G. L. Transmembrane signaling in bacterial chemoreceptors.

Trends Biochem Sci **26**, 257–265 (2001).

72. Lai, R.-Z. *et al.* Cooperative Signaling among Bacterial Chemoreceptors. *Biochemistry* **44**,

14298–14307 (2005).

73. Gavira, J. A. *et al.* How Bacterial Chemoreceptors Evolve Novel Ligand Specificities. *mBio*

11, e03066-19 (2020).

74. Matilla, M. A., Ortega, Á. & Krell, T. The role of solute binding proteins in signal

transduction. *Computational and Structural Biotechnology Journal* **19**, 1786–1805 (2021).

75. Chimerel, C., Field, C. M., Piñero-Fernandez, S., Keyser, U. F. & Summers, D. K. Indole

prevents Escherichia coli cell division by modulating membrane potential. *Biochim Biophys*

Acta **1818**, 1590–1594 (2012).

76. Melander, R. J., Minvielle, M. J. & Melander, C. Controlling bacterial behavior with indole-

containing natural products and derivatives. *Tetrahedron* **70**, 6363–6372 (2014).

77. Moore, J. P., Kamino, K., Kottou, R., Shimizu, T. S. & Emonet, T. Signal integration and

adaptive sensory diversity tuning in *Escherichia coli* chemotaxis. *Cell Systems* **15**, 628-

638.e8 (2024).

78. Gül, E. *et al.* Intraluminal neutrophils limit epithelium damage by reducing pathogen assault

on intestinal epithelial cells during Salmonella gut infection. *PLOS Pathogens* **19**, e1011235

(2023).

79. Antunes, L. C. M. *et al.* Antivirulence activity of the human gut metabolome. *mBio* **5**,

e01183-01114 (2014).

80. Peixoto, R. J. M. *et al.* Repression of Salmonella Host Cell Invasion by Aromatic Small

Molecules from the Human Fecal Metabolome. *Appl Environ Microbiol* **83**, e01148-17

(2017).

81. Lamas, A., Regal, P., Vázquez, B., Cepeda, A. & Franco, C. M. Short Chain Fatty Acids

Commonly Produced by Gut Microbiota Influence Salmonella enterica Motility, Biofilm

Formation, and Gene Expression. *Antibiotics (Basel)* **8**, 265 (2019).

82. Huang, J. Y. *et al.* Chemodetection and Destruction of Host Urea Allows Helicobacter pylori

to Locate the Epithelium. *Cell Host & Microbe* **18**, 147–156 (2015).

83. Sugihara, K. *et al.* Mucolytic bacteria license pathobionts to acquire host-derived nutrients

during dietary nutrient restriction. *Cell Reports* **40**, 111093 (2022).

84. Chandra, K. Gentamicin Protection assay (Intracellular Survival Assay) for Salmonella

Typhimurium/Typhi v1. Preprint at https://doi.org/10.17504/protocols.io.b4zmqx46 (2022).

85. Rivera Calo, J., Rubinelli, P. M. & Ricke, S. C. Gentamicin Susceptibility and Comparison

of Adhesion and Invasion of Caco-2 and HD11 Cell Lines by Salmonella enterica Serotypes.

Applied Sciences **14**, 3305 (2024).

86. Ben-David, A. & Davidson, C. E. Estimation method for serial dilution experiments. *Journal*

of Microbiological Methods **107**, 214–221 (2014).

87. Macho, A. P., Zumaquero, A., Ortiz⬚Martín, I. & Beuzón, C. R. Competitive index in mixed

infections: a sensitive and accurate assay for the genetic analysis of *Pseudomonas syringae* –

plant interactions. *Molecular Plant Pathology* **8**, 437–450 (2007).

88. Auerbuch, V., Lenz, L. L. & Portnoy, D. A. Development of a competitive index assay to

evaluate the virulence of Listeria monocytogenes actA mutants during primary and

secondary infection of mice. *Infect Immun* **69**, 5953–5957 (2001).

Supplemental Information

Table S1. Summary of prior studies related to indole chemotaxis.

Species, Strain(s)	Indole Treatment [a]	Reported Response [b]	Experimental Method(s)	Chemotaxis & Motility Proteins Involved in Response	Ref.
E. coli, strain RP437	≤1 mM >1 mM	chemorepulsion chemoattraction	tethered cell assay [c]	Tsr, Tar	[8]
E. coli, strain MG1655	1 mM ± 0.5-0.1 mM L-Ser	"bet-hedging" [d]	diffusion-based assay into long channel with fluorescent imaging	Tar, Tsr	[9]
E. coli, strain RP437	500 µM	chemorepulsion, overridden in presence of 500 µM Autoinducer-2	flow-based microfluidic chemotaxis device coupled to a gradient generator	Tar, Tsr	[14]
E. coli, strain RP437	≤1 mM >1 mM	decreased motility slower flagellar rotation	tethered cell assay [c]	Non-CheY dependent	[18]

E. coli, strain O157:H7 CDC EDL933	500 μM	chemorepulsion	agarose plug chemotaxis assay imaged every 5 minutes for 30 minutes	MotB and FliD	[25]
S. Typhimurium, strain ATCC14028s	1 mM	decreased motility	swimming motility agar plates [c]	MotA	[23]
S. Typhimurium, strain ATCC14028s	1 mM	decreased motility	swimming motility agar plates [c]	FlhC	[24]

[a] Source concentration, or ranges, used in experiment.

[b] The motility or chemotactic response as reported by the study authors; note that some methods

employed may not be able to distinguish between responses as a consequence bacterial growth

versus chemotaxis.

[c] The assay may have limitations in its ability to report on either rapid temporal responses, or

localization to or from an effector source.

[d] In this work, we refer to behaviors of this type as "chemohalation" to be similar to the widely-

used terms chemoattraction and chemorepulsion.

Data S1. Enumeration of colony-forming units (CFUs) from explant studies.

Fig. S1. Competition of WT and *invA* in explant infections. A. Competitive indices (CI) of colony-forming units (CFUs) recovered from co-infected swine explant tissue, either from the total homogenate (total, open box and whiskers plots), or from tissue washed with gentamicin to kill extracellular and attached cells (invaded, checkered box and whisker plots), as indicated. Each data point represents a single experiment and the CI of CFUs recovered from that tissue (n=9-10). Boxes show median values (line) and upper and lower quartiles, and whiskers show max and min values. Effect size (Cohen's *d*) and statistical significance are noted for each experiment in relation to competitive advantage, i.e. deviation from a CI of 1 (not significant, ns; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). B. Disaggregated CFU enumerations for each experimental group prior to CI calculation.

Fig. S2. Colony-forming units (CFU) recovered from swine colonic explant infections. Shown are disaggregated CFU enumerations for each co-infection experiment with *S.* Typhimurium IR715 WT and mutant strains following tissue treatments, as indicated. CFUs extracted either using the entire tissue homogenate (total) or extracted from tissue treated with a gentamicin wash to kill exterior non-invaded cells (invaded) are noted separately (see Materials & Methods). Each data point represents CFUs recovered from a single explant experiment (n=8-10). Box and whisker plots represent the sample median (line), the edges are the upper and lower quartiles, and whiskers extend to the max and min values. The limit of detection was 1×10^5 CFUs, and experiments at or below this limit are not included here. See Data S1 for numeric values.

Fig. S3. CIRA design, diffusion modeling, and responses to L-Ser or indole. A. CIRA experimental design. B. CIRA microgradient diffusion model, simulated with a source of 1.13 mM A488 dye after 30 s of injection. Image rendered on a pink (100% source concentration)-blue-black (0%) color intensity scale, based on previous work [36]. C. CIRA quantification overview, with blue lines corresponding to chemoattraction, null chemotactic response corresponding to gray dashed lines, and chemorepulsion corresponding to brown lines. D-G. Microgradient models of the local effector concentration for treatments with 5 mM L-Ser or indole, respectively, at Δ300 s. Color gradient corresponds to gradually decreasing effector concentrations from yellow to blue. H-I. Representative 10 s max projections of *S.* Typhimurium IR715 response to sources of L-Ser (strong chemoattraction) or indole (strong chemorepulsion) using CIRA. The glass microcapillary that injects the treatment solution is centered within the field of view. Note that cells contain a plasmid expressing mPlum (see Materials & Methods), and fluorescence data are collected in the far-red channel, so the glass microcapillary is poorly visible.

990   **Fig. S4.** *Enterobacter cloacae* exhibits fecal attraction. A. Shown are max projections over 5 s
991   from a CIRA experiment from time -5 s before fecal treatment and after 5 minutes of treatment
992   with liquid human feces. Data were collected using phase-contrast.
993

Fig. S5. Model and summary for explant infection data. Based on analyses in this study, we provide this summary of the role of chemotaxis in mediating infection advantages for different tissue pretreatments. The strength of chemotactic advantage for transiting each chemical gradient and accessing the host tissue is indicated by the width of the solid black arrows. The baseline level of chemotactic advantage seen in buffer treatments may be from effectors emitted from the host tissue (gray gradient). Other gradients containing fecal chemoattractants show a similar level of chemotactic advantage, with the highest being for fecal treatment (purple). Only soaking the tissue in pure indole results in a chemical gradient for which chemotaxis and Tsr do not provide an infection advantage. Note that bacteria are exposed only to low concentrations of residual effectors that remain after the tissue is soaked and then transferred to 300 µl of buffer for infection; they are not immersed in the more concentrated effector solution. See also Fig. 1, Data S1, and Fig. S2.