Title

Systematic discovery of directional regulatory motifs associated with

human insulator sites

Naoki Osato[1]* and Michiaki Hamada [2,3,4]

[1] Department of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

[2] Graduate School of Advanced Science and Engineering, Waseda University, Tokyo,

Japan.

[3] Computational Bio Big-Data Open Innovation Laboratory (CBBD-OIL), National

Institute of Advanced Industrial Science and Technology (AIST), Tokyo, Japan.

[4] Graduate School of Medicine, Nippon Medical School, Tokyo, Japan.

* Corresponding author.

Tel/Fax: +81-3-5734-3777 Email: naokiosato11@gmail.com, osato.n.a71d@m.isct.ac.jp

Abstract

Insulator proteins function as barriers to enhancer–promoter interactions (EPIs), thereby regulating gene expression. The primary insulator protein in vertebrates is CTCF, a DNA-binding protein (DBP); however, the roles of other DBPs in EPI insulation are not fully understood. To address this, we developed a systematic and comprehensive deep learning–based approach to identify DNA motifs of DBPs associated with insulator function. Applying this method to human fibroblast cells, we identified 97 directional motifs and a smaller number of non-directional motifs. These motifs were mapped to 23 DBPs previously linked to insulator activity, CTCF, and/or other forms of chromosomal transcriptional regulation. We found that the estimated orientation bias of CTCF was consistently proportional to the orientation bias observed in chromatin interaction data. Furthermore, these motifs showed significant enrichment at insulator sites that separate repressive and active chromatin regions, at chromatin interaction–defined boundaries, and at splice sites, compared to motifs of other DBPs. For instance, we observed that the key regulator MyoD-binding site is located at an insulator site near a gene involved in skeletal muscle differentiation and function. Importantly, our findings support the previously proposed insulator-pairing model, which suggests that insulator–insulator interactions are orientation-dependent, and highlight the involvement of multiple DNA-binding proteins beyond CTCF. Together, these results provide new insights into transcriptional regulatory mechanisms mediated by insulator-associated DBPs.

Introduction

Enhancer‒promoter interactions (EPIs), which are organized by chromatin architecture, regulate gene transcription [1]. Chromatin interactions establish loop structures stabilized by transcription factors (TFs) and DNA-binding proteins (DBPs), such as CTCF and components of the cohesin complex (e.g., RAD21 and SMC3). CTCF typically acts as an insulator by anchoring chromatin loops and restricting enhancer activity within these loops [2]. Pairs of CTCF-binding sites display a characteristic directional bias, favoring a forward‒reverse (FR) orientation over forward ‒forward (FF), reverse‒reverse (RR), or reverse‒forward (RF) orientations [3]. The forward and reverse orientations of DNA-binding sites are complementary in sequence and can be readily distinguished based on reference genome sequences. Moreover, although FR and RF orientations are sequence complements, their frequencies at chromatin loop anchors differ significantly [3], suggesting that the directional bias of DNA-binding sites is associated with chromatin loop formation.

While CTCF and cohesin are recognized as the primary insulator proteins in vertebrates [4], regional micro-capture (Micro-C) analyses have demonstrated that microcompartment interactions are largely independent of cohesin and transcription [5]. These findings imply that other DBPs may function as insulators or be associated with insulator activity. However, experimentally identifying insulator-associated DBPs among the numerous DBPs present in the genome remains labor-intensive and technically challenging when relying on in situ assays.

One potential approach to identifying insulator-associated DBPs is to analyze the

enrichment of DNA-binding sites for chromatin- and insulator-associated proteins at

chromatin interaction sites. For example, DBPs such as ZNF143 and YY1 have been

shown to participate in chromatin interactions and enhancer−promoter interactions

(EPIs) [1,6]. However, while the binding sites of ZNF143 are indeed enriched at chromatin

interaction sites, those of YY1 are not significantly enriched [6]. Moreover, CTCF,

ZNF143, and YY1 participate in a variety of biological functions beyond chromatin

organization. This multifunctionality likely reduces motif enrichment at chromatin

interaction sites, complicating the identification of DBPs that are specifically associated

with insulator functions.

Another potential approach for predicting insulator-associated DBPs is to focus on

the directional bias of CTCF DNA-binding sites. The orientation of a DBP's binding

site may not be relevant when the protein binds to a single genomic location [7]. However,

when a DBP or a complex of DBPs binds to multiple genomic loci through chromatin

interactions, their binding is often structurally organized, contributing to the observed

directional bias of DNA-binding sites [8-10]. Some DBPs involved in chromatin

interactions, such as YY1, are multifunctional proteins, with certain binding sites

participating in both chromatin interactions and insulator functions. As a result, the

directional bias of DNA-binding sites for multifunctional DBPs may be diminished.

This multifunctionality complicates the identification of their specific

insulator-associated roles.

Computational methods have been developed to predict chromatin interactions

and their associated DBPs [11,12]. However, these methods are not designed to identify the

binding motifs of DBPs specifically associated with insulator function. Regarding DBP

binding motifs, Xie et al. noted that a key challenge is developing systematic

approaches for determining their specific functions on a genome-wide scale [13]. In terms

of gene regulation, insulators influence EPIs and gene expression; therefore,

incorporating gene expression data may improve the identification of

insulator-associated DBPs compared to approaches that rely solely on enrichment

analysis or the directional bias of DNA-binding sites. Recently, a novel deep

learning–based method was introduced to estimate gene expression levels in human

tissues based on DNA-binding sites within promoter regions [14]. However, this method

does not account for DNA-binding sites associated with EPIs as input features and was

not intended to identify DBPs linked to insulator function.

Therefore, we aimed to identify insulator-associated DBPs using a computational

approach that integrates and analyzes genome-wide experimental data. In this study, we

refined an existing deep learning method by incorporating EPI information based on

eQTL data and enhancer–promoter association rules (McLean et al., Supplementary Fig.

2) [15]. We analyzed the directional bias of DNA-binding sites for CTCF and

insulator-associated DBPs, including known factors such as RAD21 and SMC3, using

contribution scores generated by the deep learning model. Our analysis demonstrated

that the directional bias of CTCF binding sites closely matches the orientation bias

observed in chromatin interaction data. Furthermore, statistical testing of contribution

scores between directional and non-directional DNA-binding sites of

insulator-associated DBPs showed that directional sites contribute more significantly to

gene expression prediction than non-directional sites. Overall, we identified 23

insulator-associated DBPs related to insulator function, CTCF-associated activity,

and/or other transcriptional regulatory mechanisms affecting EPIs, as previously

reported. To validate these findings, we showed that the DNA-binding sites of the

identified DBPs are located within putative insulator regions, and that some of these

sites exist independently of nearby CTCF or cohesin binding sites. Finally, consistent

with the insulator-pairing model proposed from experimental studies in flies, we

confirmed that both homologous and heterologous insulator–insulator pairing

interactions are orientation-dependent. Our approach facilitates the identification of

insulator- and chromatin-associated DNA-binding sites that modulate EPIs and

uncovers novel functional roles and molecular mechanisms of DBPs involved in

transcriptional condensation, phase separation, and transcriptional regulation.

Results

Predictive modeling of gene expression from directional DBP binding

Accurate gene expression prediction can be achieved by integrating both distal

and proximal DNA-binding protein (DBP) binding sites as features within a machine

learning framework. By combining ChIP-seq data from the Gene Transcription

Regulation Database (GTRD v20.06) with expression quantitative trait locus (eQTL)

data from GTEx v8 [16], our model was able to account for enhancer–promoter

interactions (EPIs) by linking single nucleotide polymorphisms (SNPs) to gene

expression outcomes (**Fig. 1a-c**, **Fig. 2a**, **b**). Specifically, DBP binding sites within ±50

bp of an eQTL SNP were assumed to influence the associated transcript. Focusing on

four human cell types—HFFs, monocytes, HMECs, and neural progenitor cells—the

model achieved strong concordance with observed expression levels: Spearman's ρ

values reached 0.74, 0.71, 0.74, and 0.73 for all 27,428 genes, and 0.70, 0.67, 0.69, and

0.70 for 2,705 test genes (**Fig. 3a**, **Fig. S2a**). Importantly, this performance was

obtained without incorporating RNA-binding protein or miRNA data as required in

previous approaches [14].

To build a robust foundation for regulatory modeling, we comprehensively

identified binding motif sequences for 715 out of 1,310 DBPs using ChIP-seq peaks

from the Gene Transcription Regulation Database (GTRD, v20.06 release) and the

Protein Interaction Quantitation (PIQ) tool, which integrates DNase-seq data and a

curated library of motifs from TRANSFAC (2019.2), JASPAR (2018), UniPROBE

(2018), high-throughput SELEX, ENCODE ChIP-seq–derived motifs, and

HOCOMOCO (versions 9 and 11) [17,18-23]. Only motif occurrences overlapping ChIP-seq

peaks for each DBP were retained, thereby minimizing false positives and enabling the

high-resolution prediction of DBP binding in open chromatin regions.

We next examined how the directionality and genomic position of

insulator-associated DBP binding sites affect gene expression prediction. Using

DeepLIFT, which quantifies each feature's contribution to the model by measuring its

deviation from a reference, we demonstrated that both the orientation and the position

of DBP binding sites have substantial regulatory effects [24]. Directional differences in

putative insulator-associated DBP sites emerged as critical determinants of EPI strength

and gene expression outcomes (**Fig. 2d**). This systematic approach allowed us to dissect

how the regulatory effects of insulator-associated DBPs depend on both their chromatin

context and strand orientation.

In particular, we found that CTCF binding site orientation is a key determinant of

its regulatory impact on gene expression. CTCF sites in the forward–reverse (FR,

convergent) orientation—accounting for 76.4% of distal CTCF-mediated chromatin

interaction sites identified by ChIA-PET (**Fig. 3b**)—exhibited significantly different

DeepLIFT score distributions compared to the reverse–forward (RF) orientation

(Mann–Whitney U test, FDR-adjusted $p < 0.05$; $p < 10^{-3}$; **Fig. 3c**, **d**; **Table S1**) [3]. This

difference was robustly detected when comparing FR and RF CTCF-binding sites

located in predicted enhancer and open chromatin regions both upstream and

downstream of TSSs (**Fig. 2c**). Further analyses using alternative DNA-binding site

selection criteria supported these findings (**Supplementary Fig. 1** and **Note**),

demonstrating that our approach captures the regulatory significance of CTCF binding

site orientation.

Extending this analysis, we observed that directional orientation is a widespread

and biologically meaningful predictor of gene expression among putative

insulator-associated DBPs. DBPs such as RAD21, SMC3, and CTCF exhibited strong

differences in DeepLIFT score distributions between FR and RF orientations (**Fig. 3c**,

Table S1), and this effect was not limited to these proteins. Among 228 DBPs with at

least 1,000 binding sites in HFF cells, 92—including ZNF143 and YY1—showed

significant differences between sites with different orientations (e.g., FR vs. RF), while 79—including YY1—also showed differences between same-orientation sites (e.g., FF vs. RR) (**Fig. 3c**, **Fig. S2b**, **Table S1**, sheet **"DBP List 1A"**). Comparisons between directional (FR, RF, FF, RR) and non-directional (X) binding sites consistently produced higher z-scores for the former, highlighting the greater contribution of directional binding sites to gene expression prediction. The most pronounced effects were observed for established insulator- and chromatin-associated proteins, underlining the biological importance and sensitivity of our model. For increased stringency, we restricted the analysis to DBPs with $p < 10^{-3}$, confirming all such cases overlapped with those defined by the broader FDR-adjusted $p < 0.05$ threshold (**Table S1**). These findings prompted us to systematically classify insulator-associated DBPs by their directional or non-directional binding patterns for further analysis.

Identification and classification of insulator-associated DBPs

Insulator-associated DBPs demonstrate widespread and significant orientation-dependent as well as orientation-independent DNA-binding preferences, reflecting their diverse roles in chromatin regulation. Our analysis revealed that established insulator DBPs—including CTCF, RAD21, and SMC3—exhibit a predominant forward–reverse (FR) orientation in their binding site distributions. Pairwise comparisons of DeepLIFT scores across four orientations (FR, RF, FF, RR) and non-directional sites showed significant differences for 23 DBPs with different orientations and for 17 DBPs with the same orientation. Additionally, non-directional

binding sites showed significant score differences for six DBPs ($p < 10^{-3}$; **Table 1** and

Table S2, sheet "DBP List 1A"; FDR-adjusted $p < 0.05$; **Table S3**, "DBP List 1A").

Beyond these established factors, we identified several novel DBPs that displayed both

directional and non-directional binding preferences, expanding the pool of proteins

potentially involved in insulator function.

Importantly, these widespread orientation biases were supported by chromatin

interaction data. Our deep learning–based predictions accurately recapitulated the

characteristic orientation bias of CTCF and RAD21 DNA-binding sites at chromatin

interaction sites, as validated by ChIA-PET data. Specifically, among CTCF binding

sites overlapping distal interaction anchors, 76.4% displayed an FR orientation, with

substantially lower frequencies of FF (11.0%), RR (10.3%), and RF (2.3%) (**Fig. 3b**;

Guo Y et al., *Cell* 2015 [3], Fig. 4). DeepLIFT scores revealed that the difference between

FR and RF orientations was significantly greater than that between FR and FF or RR

(Mann–Whitney U test), consistent with experimental distributions (**Fig. 3c**, **d**).

Similarly, RAD21 ChIA-PET data revealed 76.3% FR, 3.4% RF, 9.3% FF, and 11.1%

RR orientations, and DeepLIFT scores for RAD21 binding sites closely mirrored these

results (**Fig. 3c**; **Table 1**). These findings confirm that our framework robustly predicts

the experimentally observed orientation biases of insulator-associated DBPs at

chromatin interaction sites.

Applying stringent orientation criteria, we identified a comprehensive set of 97

insulator-associated DBPs that directly interact with DNA and exhibit either strong

directional or non-directional binding preferences. Most of these DBPs exhibited clear

directional biases, while only six—including FOS, FOSL2, JUN, JUNB, JUND (all

with palindromic motifs), and SPI1 (their interacting partner)—were characterized by

non-directional binding. Notably, 23 of these DBPs have been previously implicated in

insulator activity, CTCF association, or chromatin regulatory functions affecting

enhancer–promoter interactions (EPIs), with 13 of them directly interacting with CTCF

(**Table S7**). These proteins are multifunctional, serving as transcriptional regulators,

repressors, chromatin regulators, or cofactors for epigenetic modifier enzymes.

Collectively, our findings indicate that CTCF coordinates EPI regulation through

a network of diverse DBPs and that both directional and non-directional binding

preferences underlie the regulatory capacity of these proteins. The involvement of these

multifunctional DBPs in chromatin architecture and long-range genomic interactions

emphasizes the complex and modular organization of chromatin regulatory networks.

Extending these analyses to the spatial organization of chromatin, we found that

insulator-associated DBPs are frequently co-localized at both ends of chromatin

interaction sites, supporting their role in mediating long-range chromatin architecture.

Across 30,369 chromatin interactions defined by 400-bp regions (separated by >5 kb

and excluding TSS-proximal areas), DNA-binding sites for 44 (45%) and 97 (100%)

insulator-associated DBPs were identified at both ends or at either end of an interaction

pair, respectively. For 10,384 interactions defined by 200-bp regions, binding sites for

30 (31%) and 97 (100%) DBPs were detected at both or either end. These analyses,

based on orientation-aware pairings (FR, RF, FF, RR) (**Fig. 3e**), likely underestimate the

full extent of insulator DBP involvement due to incomplete coverage of chromatin

interaction data [25]. Canonical insulator proteins such as CTCF, RAD21, and SMC3

exhibited strong directional binding within these regions, consistent with ChIA-PET

results. To further validate the predictions of candidate insulator-associated DBPs, we

next sought independent lines of evidence—including histone modification signatures

characteristic of insulator function—to corroborate their roles in chromatin boundary

and architecture.

Epigenomic validation of insulator activity

Epigenomic analyses provide independent support for the predicted regulatory

roles of insulator-associated DBPs. We identified insulator-associated DBP binding sites

by detecting asymmetry in H3K27me3 signal profiles across their upstream and

downstream regions, reflecting boundaries between active and repressed chromatin

domains. Specifically, loci where the mean H3K27me3 signal intensity differed by more

than twofold between upstream and downstream regions, with the higher value

exceeding the first quartile of the genome-wide distribution, were selected to ensure

robust discrimination and to filter out low-intensity noise. This approach builds on prior

observations that CTCF and USF1 occupy the edges of repressive chromatin domains

(marked by H3K27me3) versus active regions (marked by H2AK5ac) [26,27], making

H3K27me3 a suitable marker for broad transcriptional repression.

DNA-binding sites of predicted insulator-associated DBPs were significantly

enriched at chromatin boundaries defined by asymmetric H3K27me3 signals. Known

insulator-associated factors, such as CTCF, RAD21, and SMC3, showed the highest

number of such sites (**Table S4**). Quantitatively, the median and mean counts of binding

sites with greater than twofold H3K27me3 signal differences between upstream and

downstream regions were significantly higher for predicted insulator-associated DBPs

than for other DBPs (Mann–Whitney U test, $p < 0.05$; **Table 2**, **Table S4**, **Fig. 4a**, **b**).

These results demonstrate that insulator-associated DBPs preferentially occupy genomic

positions flanked by distinct chromatin states, providing further evidence for their

function in defining regulatory domains.

Furthermore, predicted insulator-associated DBPs are localized to transcriptional

transition regions and chromatin boundaries. Analysis of RNA-seq data from HFF cells

showed that these DBPs are significantly enriched at sites where transcript signal

intensities differ by more than 1.5-fold between upstream and downstream regions, with

the higher value exceeding the first quartile of the transcriptome-wide distribution

(**Table 2**, **Table S4**, **Fig. 4a**, **c**). Additionally, DNA-binding sites of these DBPs more

frequently overlapped with chromatin boundary regions, as defined by insulation scores

from chromatin interaction data [28], than those of non-insulator DBPs (**Table 2**, **Fig. 4d**).

These results underscore the role of insulator-associated DBPs in partitioning active and

inactive chromatin domains and functionally positioning at regulatory boundaries. To

explore the fine-scale genomic context of these sites, we next conducted genome

browser inspections.

Genomic inspection using the UCSC Genome Browser revealed that predicted

insulator-associated DBPs consistently occupy sites with insulator-like features,

including chromatin boundary regions identified by Micro-C data and transitional

regions between H3K27me3-marked and transcriptionally active domains. Many of

these binding sites were located at the edges of H3K27me3-enriched domains and were

frequently distant from canonical insulator proteins such as CTCF, RAD21, and SMC3

(**Fig. 4a**, **Fig. S3**). For each DBP, up to three representative binding sites in

insulator-associated contexts were manually selected (**Fig. S3a**, **S3f**). Notably, MyoD1

was found at a chromatin boundary near the SELENON (SEPN1) gene—both involved

in skeletal muscle differentiation [29] —suggesting a possible regulatory role for MyoD1

via insulator mechanisms (**Fig. 4a**).

Expanding this analysis to alternative repressive histone modifications,

insulator-associated DBPs were also enriched at the edges of H3K9me3-enriched

domains (**Fig. S3b**), supporting their ability to define chromatin boundaries beyond

those marked by H3K27me3. The distribution of H3K9me3 differed from that of

H3K27me3 in certain regions, highlighting the context-dependent nature of insulator

activity. Additionally, predicted DBP binding sites frequently formed spatial clusters at

insulator regions and were often associated with chromatin interaction anchors (**Fig. 4f**,

Fig. S3c, **S3d**). This clustering aligns with the role of architectural proteins in

organizing TAD boundaries, while discrete insulator motifs contribute to intra-TAD

looping [30]. To further clarify the relationship between these epigenetically defined

insulator sites and chromatin architecture, we compared their locations with chromatin

boundaries identified by chromatin interaction data.

Comparative analyses showed that histone- and transcription-defined insulator

sites often coincided with, or were located near, chromatin boundaries identified by

interaction data. In HFF cells, 29% of H3K27me3-based insulator sites (type 1) and

37% of H3K27me3 plus transcription-based sites (type 2) overlapped with 25-kb

boundary regions, whereas only 6% and 7% overlapped with 5-kb boundaries,

respectively, reflecting differences in boundary resolution (**Fig. 4f**, **Fig. S3**).

Genome-wide, DNA-binding sites of the 97 predicted insulator-associated DBPs were

found at type 1 insulator sites assigned to 3,170 genes, at type 2 sites assigned to 1,044

genes, and at chromatin interaction–defined boundaries assigned to 6,275 genes. Among

these, 1,212 genes (38%) harbored both type 1 insulator and boundary sites, and 389

genes (37%) contained both type 2 insulator and boundary sites. These data indicate that

while chromatin interaction boundaries and epigenetically or transcriptionally defined

insulator sites are not completely overlapping, they frequently target overlapping gene

sets and are co-located within the same regulatory landscapes, supporting their

complementary roles in chromatin compartmentalization.

Independent and cooperative clustering of insulator-associated DBPs

at chromatin regulatory landmarks

Predicted insulator-associated DBPs are enriched at critical transcript regulatory

regions, such as transcription start sites (TSSs), transcription termination sites (TTSs),

and splice sites, suggesting potential roles in shaping transcript structure. Statistical

analyses showed that these DBPs are significantly more abundant at splice sites than

other DBPs (Mann–Whitney U test, $p < 0.05$; **Fig. 4e**, **Table 2**), with pronounced

enrichment at the first and last introns compared to internal introns (**Table 2**, **Table S9**).

This finding is consistent with prior reports that variation in TSS/TTS usage is a key

driver of tissue-specific isoform diversity [31]. Examination of **Fig. 4a** and **Fig. S3** further

revealed clustering of multiple insulator-associated DBP binding sites within single loci,

suggesting potential cooperative functions in establishing insulator elements. Notably,

transcribed regions with overlapping H3K27me3 enrichment and alternative

transcriptional repression contained DBP sites between active and repressed domains

(**Fig. 5**, **Fig. S3e**), indicating that insulator-associated DBPs may delineate boundaries

between transcriptional states, thereby influencing selection of TSSs, TTSs, and exons

in a context-dependent manner.

Chromatin interaction–associated DBPs, including several predicted insulator

factors, are strongly enriched near chromatin interaction centers, emphasizing their

potential in supporting chromatin looping. High-resolution Micro-C data from HFF

cells identified DBP binding sites located within $\pm$500 bp of interaction anchors [32].

Canonical organizers CTCF, RAD21, and SMC3 exhibited sharp enrichment at these

sites (**Fig. 6a–c**), while YY1 and ZNF143 displayed broader or distinct distributions

(**Fig. 6d**, **e**), and MAZ—a DBP implicated in CTCF-independent insulator activity [33]—

demonstrated a unique peak (**Fig. 6f**). Even after removing binding sites overlapping

CTCF, RAD21, and SMC3, other predicted insulator-associated DBPs remained

consistently enriched at interaction centers, supporting their independent contributions

to 3D chromatin structure.

Analysis of 228 DBPs with over 1,000 binding sites in HFF cells revealed that

nearly 90% exhibited distinct enrichment near chromatin interaction sites (**Table S8**,

Fig. S4), displaying sharp, broad, or canonical CTCF-, RAD21-, or SMC3-like peak

profiles. Although DBPs such as MeCP2 and PRDM9, with fewer than 1,000 binding

sites, did not exhibit this pattern, these findings demonstrate that most DBPs—including

many predicted insulator-associated proteins—are spatially positioned to participate in

genome architecture regulation.

Co-localization analyses revealed that most predicted insulator-associated DBPs

are spatially segregated from canonical insulator proteins. Using 10-bp bins across ±500

bp windows centered at interaction anchors and ChIP-seq–validated binding sites, we

identified 245 DBP pairs (involving 135 DBPs) with ≥30% co-localization (Selection

Criteria S1 in Fig. S1; **Table S5**). CTCF was highly co-localized with RAD21 (39%)

and SMC3 (81%), whereas YY1/ZNF143 had much lower rates (<6%) with canonical

factors (**Table 3**). Some DBPs (CTCFL, RXRA, PPARG, NEUROD1, ZBTB7A)

exhibited moderate co-localization (>10%), but the majority of predicted

insulator-associated DBPs showed minimal overlap (<10%), suggesting functional

independence. Spatial clustering of multiple DBP binding sites was also observed at

putative insulator loci (**Fig. S3d**), and co-localization was especially strong at 5-kb

boundary-associated sites (197 pairs, 101 DBPs, ≥30% overlap; **Table S6**), suggesting

that these DBPs, like CTCF and cohesin, may assemble cooperatively at boundaries to

mediate domain partitioning.

Overlap analysis with ChIP-seq data showed that >98% of CTCF, RAD21, and

SMC3 binding sites overlapped their own ChIP-seq peaks under all conditions, thereby

validating site definitions. In contrast, nearly all other predicted insulator-associated

DBPs showed <50% overlap with CTCF, RAD21, or SMC3 peaks, even when data

from various cell types were considered. This low co-occupancy supports the notion

that predicted DBPs function largely independently of classical insulator complexes and

may establish chromatin boundaries by distinct mechanisms.

Interestingly, a distinct group of predicted insulator-associated DBPs—including

BACH2, FOS, ATF3, NFE2, and MAFK—exhibited binding site distributions around

chromatin interaction centers similar to canonical insulator proteins, but had less than

1% direct spatial overlap within 10-bp bins (**Fig. S4c**). However, these factors showed

strong mutual co-localization (**Table 3**), suggesting the existence of parallel,

functionally distinct DBP clusters that independently regulate chromatin structure [343536].

Thus, multiple DBP clusters with similar positional preferences may exert parallel

effects on chromatin remodeling or loop accessibility via independent regulatory

pathways.

Discussion

Here, we present the first, to our knowledge, computational method and

genome-wide analysis for the detection of insulator-associated DBPs. Transcriptional

regulation mediated by the binding of DBPs at sites located far from TSSs was modeled

using our deep learning approach based on eQTL data. Through this framework, the

effects of DNA-binding sites of insulator-associated DBPs on EPIs and gene expression

were estimated, and insulator-associated DBPs were predicted. We further identified

DBPs, including known insulator-associated proteins, that exhibit insulator-associated

functions and regulate transcription, thereby affecting EPIs as reported in previous

studies. The DNA-binding sites of the predicted insulator-associated DBPs

demonstrated orientation bias; unexpectedly, the significant differences in the

orientation bias of CTCF DNA-binding sites based on DeepLIFT scores (contribution

scores) were found to be consistent with the percent differences in the orientation of

CTCF DNA-binding sites at CTCF-mediated chromatin interaction sites. Furthermore,

statistical testing of the contribution scores revealed that the directional binding sites of

DBPs contributed more significantly to the prediction of gene expression levels than the

non-directional sites. The DNA-binding sites of the predicted insulator-associated DBPs

were located at potential insulator sites more frequently than those of other DBPs, as

indicated by differences in H3K27me3 signal intensities and transcript levels, and by

their overlap with boundary sites identified by chromatin interactions. We further

observed that only a small percentage of the DNA-binding sites of insulator-associated

DBPs co-localized with the DNA-binding sites of known insulator-associated proteins

such as CTCF, RAD21, and SMC3 around chromatin interaction sites. Finally, we

discovered that DBPs other than the well-established insulator proteins, as well as the

directionality of their DNA-binding sites, are associated with insulator function.

Our deep learning analysis accurately recapitulated the percent differences in the

FR, FF, RR, and RF orientations of CTCF DNA-binding sites at chromatin interaction

sites, by incorporating the genomic locations of DNA-binding sites for putative

insulator-associated DBPs. The other predicted insulator-associated DBPs also

demonstrated orientation bias in their DNA-binding sites. These results suggest that the

directional bias of DNA-binding sites of insulator-associated DBPs may contribute to

insulator function and chromatin regulation through structural interactions among DBPs,

other proteins, DNA, and RNA. For example, the N-terminal amino acids of CTCF

have been shown to interact with RAD21 in chromatin loops [8-10]. Cohesin complexes

that interact with convergent CTCF sites, that is, the N-terminus of CTCF, might be

protected from WAPL, whereas those that interact with divergent CTCF sites, i.e., the

C-terminus of CTCF, may not be protected from WAPL, which could release cohesin

from chromatin and thus disrupt cohesin-mediated chromatin loops [37]. As another

example, although sequence-based predictions of EPIs underestimated actual

interactions, an activity-by-contact (ABC) approach, based on a simple distance model,

performed well, suggesting that factors in addition to sequence influence EPIs [38,39].

Furthermore, CTCF-mediated chromatin looping induces the formation of

transcriptional condensates and phase separation between the inner and outer regions of

the loops, resulting in insulator function [40-42]. Therefore, structural features that drive

phase separation, rather than functional differences in DBPs themselves, may underlie

insulator activity. CEBPA, among the identified insulator-associated DNA-binding

proteins, has also been reported to participate in transcriptional condensates and phase

separation [43]. The FOXA1 pioneer factor functions as an initial chromatin-binding and

chromatin-remodeling factor and has also been shown to form biomolecular

condensates [44].

Other types of chromatin regulation are also likely to be related to the structural

interactions of molecules. In the boundary-pairing (insulator-pairing) model, molecules

bound to insulators physically pair with their partners, either head-to-head or

head-to-tail, with different degrees of specificity at the termini of TADs in flies (**Fig. 7**).

Although experiments have not clarified how partners locate each other, the mechanism

is unlikely to require loop extrusion, as is presumed in CTCF- and cohesin-mediated

chromatin looping. Both homologous and heterologous insulator–insulator pairing

interactions are thought to be central to the architectural functions of insulators.

Importantly, the nature of insulator–insulator interactions is orientation-dependent [45-47].

Regional micro-capture (Micro-C) analyses have revealed that microcompartment

interactions are largely independent of cohesin and transcription [5]. Depletion of cohesin

results in a subtle reduction in longer-range enhancer−promoter interactions, while

CTCF depletion can lead to rewiring of regulatory contacts [48]. Another group reported

that enhancer−promoter interactions and transcription are largely maintained upon

depletion of CTCF, cohesin, WAPL, or YY1; instead, cohesin depletion reduced

transcription factor binding to chromatin, suggesting that cohesin may enable

transcription factors to find and bind their targets more efficiently [49]. Furthermore, loop

extrusion is not essential for enhancer−promoter interactions but contributes to their

robustness, specificity, and the precise regulation of gene expressio [48]. The 'loop

extrusion' hypothesis is motivated by in vitro observations. Experiments in yeast, using

cohesin variants unable to extrude DNA loops but capable of topologically entrapping

DNA, suggested that in vivo chromatin loops can form independently of loop extrusion.

Instead, transcription was shown to promote loop formation and act as an extrinsic

motor that extends these loops and defines their final positions [50]. Maz and MyoD1,

among the identified insulator-associated DNA-binding proteins, have been shown to

form loops without CTCF [4,33,51]. Furthermore, protein post-translational modifications

(PTMs) facilitate control over the molecular valency and strength of protein−protein

interactions [52]. O-GlcNAcylation as a PTM has been reported to inhibit CTCF binding

to chromatin [53]. We found that the identified insulator-associated DNA-binding proteins

tend to cluster at potential insulator sites (Fig. 4f and Supplementary Fig. 3c). These

proteins may interact and actively regulate chromatin interactions, transcriptional

condensation, and phase separation via PTMs. Our analyses will contribute to revealing

novel functional and structural roles of DBPs in insulator functions, chromatin

regulation, and phase separation, and to advancing their experimental analysis.

High-resolution chromatin interaction data from Micro-C assays demonstrated

that most of the predicted insulator-associated DBPs exhibited peaks in the

DNA-binding site distribution around chromatin interaction sites, suggesting that these

DBPs are involved in chromatin interactions and highlighting the high resolution of

Micro-C data. Base pair−level resolution has been reported using Micro Capture-C [32].

However, due to the high resolution of Micro-C assays, it is challenging to obtain

genome-wide chromatin interaction data at a reasonable and accessible cost, whereas

Micro Capture-C and Region Capture Micro-C assays focus on specific genomic

regions to increase the coverage of chromatin interactions. The extent of chromatin

interaction coverage varies depending on the specific Micro-C experimental design. In

addition, chromatin interactions include types of interactions that are not associated

with EPIs and insulator functions [54-56]. Therefore, we analyzed and predicted

insulator-associated DBPs independently of chromatin interaction data. Our deep

learning analysis successfully learned enhancer–promoter interactions (EPIs) that affect

gene expression levels and accurately recapitulated the percent differences in the FR, FF,

RR, and RF orientations of CTCF DNA-binding sites at chromatin interaction sites.

When comparing insulator sites separated by active and repressive domains with

boundary sites identified by chromatin interaction data, we defined insulator sites based

on the total signal levels of H3K27me3 histone modification marks and transcript

expression levels within a region. Some insulator sites may not display smooth,

continuous distributions of histone marks and expression, which could account for

differences between insulator and boundary sites. Another possible explanation is that

boundary sites are more closely associated with topologically associating domains

(TADs) of chromosomes than are insulator sites [30]. Boundary sites represent regions

identified based on the separation of numerous chromatin interactions. In contrast, we

found that multiple DNA-binding sites of insulator-associated DNA-binding proteins

were frequently clustered at insulator sites and were associated with distinct nested and

focal chromatin interactions, as reported by Micro-C assays [5]. These interactions may

be transient and relatively weak, such as tissue-, cell type-, condition-, or

lineage-specific interactions [57].

AI (Artificial Intelligence) is often regarded as a black box, since the reasons and

causes of its predictions are difficult to determine. To address this issue, various tools

and methods have been developed to enhance model interpretability, collectively

referred to as Explainable AI. DeepLIFT is considered one such tool for Explainable AI.

However, DeepLIFT does not provide the specific reasons or causal mechanisms

underlying a prediction. Instead, it computes scores representing the contribution of

each input feature to the prediction outcome. In this study, analysis of numerous

contribution scores (i.e., DeepLIFT scores) for directional DNA-binding sites of

putative insulator-associated DBPs in the prediction of transcript expression levels

enabled accurate prediction of the percent differences in the FR, FF, RR, and RF

orientations of CTCF DNA-binding sites at chromatin interaction sites, based on the

contribution scores of DNA-binding sites in putative enhancer and open chromatin

regions. It should be noted that some contribution scores may not be accurate, and

predictions of insulator-associated DBPs or DNA-binding sites based on a small number

of scores may be unreliable. To enable comparison of contribution scores calculated

from different input–output pairs within our deep learning model, we normalized the

contribution scores for each input–output pair as previously described [58]. The analysis

of large sets of contribution scores will facilitate the estimation of the effects of other

molecules and functions involved in gene regulation.

One of the next computational challenges will be to predict and refine the

potential set of insulator-associated DNA-binding sites for multifunctional DBPs. To

validate machine learning predictions and identify insulator-associated DNA-binding

sites in the genome for subsequent experimental investigation, information on histone

modifications such as H3K27me3 and transcript levels will be valuable, and other types

of epigenomic data can also be utilized for this purpose. For example, as a reference for

an application of this approach, Liu et al. focused on CTCF-binding sites and performed

various assays in human and mouse cell cultures, as well as in mice, with the goal of

applying these sites in gene therapy [59]. The analysis of insulator-associated DBPs may

ultimately contribute to the development of technologies for precise gene expression

control. In summary, our novel computational method has identified

insulator-associated DBPs—including previously characterized factors—and revealed

that the directional bias of these DBPs is potentially involved in chromatin looping,

phase separation, transcriptional condensate formation, and insulator function through

the structural interactions of DBPs. These findings will contribute to a more

comprehensive and global understanding of transcriptional regulation and gene

expression.

Methods

Prediction of gene expression levels based on DNA-binding sites

Gene expression data as output data for machine learning analysis

To predict DNA-binding sites that may also function as insulator sites in the

human genome, we modified a previously developed deep learning method originally

designed to predict transcript expression levels across tissues and cell types based on

DNA-binding sites of DBPs in promoter regions and other genomic regions (Fig. 2a;

Fig. S4) [14].   To minimize the influence of cellular heterogeneity inherent in tissue

samples, we utilized gene expression data from four distinct human cell types for which

extensive experimental datasets are publicly available: human foreskin fibroblasts

(HFFs; GSM941744), monocytes (Blueprint RNA-seq FPKM data, file

'C0010KB1.transcript_quantification.rsem_grape2_crg.GRCh38.20150622.results',

available at http://dcc.blueprint-epigenome.eu/#/files), human mammary epithelial cells

(HMECs; 'wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam', ENCODE project

via UCSC Genome Browser http://genome.ucsc.edu/encode/), and neural progenitor

cells (NPCs; GSM915326, ENCFF529SIO).   FPKM (Fragments Per Kilobase of exon

per Million mapped reads) values representing gene expression levels were retrieved or

calculated based on GENCODE v19 annotations from the ENCODE portal [60]

(https://www.encodeproject.org/)   using   the   RSeQC   tool   [61]

(http://rseqc.sourceforge.net/).   Following the data processing pipeline described by

Tasaki et al. [14], we calculated $\log_2$-transformed FPKM values (adding 0.25 to avoid

negative infinity), determined median $\log_2$-FPKM values across the four cell types, and

computed $\log_2$-fold changes relative to the median expression for use as outputs in the

deep learning model.

DNA-binding sites as input data for machine learning analysis

To investigate the effect of insulator sites on EPIs and gene expression, the

DNA-binding sites in putative enhancer and open chromatin regions located away from

promoter regions were added as input data in the deep learning method. To identify the

DNA-binding sites of DBPs in promoter and enhancer regions, ChIP-seq data of the

DBPs were used and were updated to GTRD database release v20.06

(https://gtrd.biouml.org) [62]. The binding site of a DBP regulating the expression level of

each transcript was selected using eQTL data (GTEx_Analysis_v8_eQTL.tar) from

GTEx portal release v8 (https://gtexportal.org). If a DNA-binding site was located

within a 100-bp region around a single-nucleotide polymorphism (SNP) of an eQTL,

we assumed that the DBP regulated the expression of the transcript corresponding to the

eQTL. If the SNP of an eQTL was not located on the same chromosome as the

transcribed region of the eQTL (i.e., *trans*-eQTL), the eQTL data were not used. The

genomic positions of the eQTLs were converted to the corresponding positions in the

hg19 version of the human genome using the LiftOver tool

(http://hgdownload.soe.ucsc.edu/downloads.html#liftover).

The original model of the input data of DNA-binding sites in promoter regions

used for the deep learning analysis consisted of the lengths of a DBP ChIP-seq peak in

each 100-base bin in a region between 2 kbp upstream and 1 kbp downstream of the

transcription start site (TSS) of each gene or transcript [14]. To add the input data of

DNA-binding sites of DBPs in more upstream and downstream regions of a TSS and to

reduce the number of bins, we modified the model and first computed the length of a

DBP ChIP-seq peak in a 100-base bin and then selected the maximum length in each

interval of 100 bins in a region between 1 Mbp upstream and downstream of a TSS.

This step outputted the 200 longest DBP ChIP-seq peaks in the examined regions.

Training step for machine learning analysis

In the training of the modified deep learning model, we used the same test,

training, and validation data as in the original method: all 2,705 genes encoded on

chromosome 1 as test data, training data (22,251 genes), and validation data (2,472

genes) randomly selected from the other genes [14]. The input data for promoters and

putative enhancer regions were concatenated into a total of 230 bins of the lengths of

1,310 DBP ChIP-seq peaks, consisting of 30 bins for promoters and 200 bins for

putative enhancers because the DeepLIFT tool 'RevealCancel rule on dense layers and

the Rescale rule on conv layers (genomics default)', with high scoring accuracy, [24]

cannot be applied to a neural network with multiple input layers (Fig. S5). Ubuntu

(version 20.04), Python (version 3.8.10), TensorFlow (version 2.6.0), NVIDIA CUDA

toolkit (version 11.4.2), GeForce RTX 3080 GPU with 10 GB memory, and Intel Core

i7-10700 CPU with 128 GB memory were used for training and prediction, and the

original scripts were modified to run on newer versions of the software. The training

method was the same as the original, and the training time was two to four hours. In this

study, we focused on the analysis of HFF cells and assessed the training process using

all the input data as the test data, which slightly improved the prediction of gene

expression. Therefore, we used the trained model for the analysis of insulator-associated

DBPs.

Prediction of insulator-associated DBPs affecting EPIs

DNA-binding sites of putative insulator-associated DBPs

To identify the direction of DBP-binding motifs in the DNA-binding sites of

predicted insulator-associated DBPs, we searched for DBP-binding motifs in open

610 chromatin regions in HFF cells (GSE170964) using the PIQ tool [17] and the position

611 frequency matrices of the DBP-binding motifs. The motif sequences were collected

612 from databases and previous studies, specifically, TRANSFAC (2019.2), JASPAR

613 (2018), UniPROBE (2018), DNA motifs from high-throughput SELEX, DNA motifs

614 from ENCODE DBP ChIP-seq data, and HOCOMOCO versions 9 and 11 [18-23]. To

615 reduce the number of false-positive prediction of DNA-binding sites, only

616 DNA-binding sites that overlapped the ChIP-seq peaks of the same DBP (GTRD

617 database release v20.06 [https://gtrd.biouml.org]) were used for further analyses. The

618 genomic positions of ChIP-seq and DNase-seq data in public databases were converted

619 to those provided in the hg19 version of the human genome using the LiftOver tool.

620  To confirm that CTCF DNA-binding sites were located at chromatin interaction

621 sites, we examined the overlap between CTCF ChIP-seq peaks and the sites of *in situ*

622 ChIA-PET chromatin interactions mediated by CTCF in human foreskin fibroblast

623 (HFF) cells (4D Nucleome data portal,

624 https://data.4dnucleome.org/files-processed/4DNFI73AFN12/). The directional

625 difference between RAD21 DNA-binding sites located at chromatin interaction sites

626 was also examined using ChIA-PET chromatin interactions mediated by RAD21 in

627 human fibroblast from skin (ENCLB147AJZ). RAD21 DNA-binding sites within 100

628 bp of chromatin interaction sites were associated with chromatin interactions. To

629 eliminate DNA-binding sites that overlapped EPI, we obtained ChIP-seq peaks of

630 histone modifications, such as H3K4me3, H3K27ac, and H3K4me1, associated with

631 promoter and enhancer regions in HFF cells (4D Nucleome data portal,

4DNFIJOK1BWR, 4DNFI442BQLO, and 4DNFIAGV3X99).

Preparation of input data considering the DNA-binding sites of an

insulator-associated DBP

To prepare the input data of the DNA-binding sites of DBPs in enhancers and

promoters for the deep learning analysis, first, a transcript regulated by a DBP bound to

enhancers and promoters was predicted based on the single nearest gene association rule

(Fig. 2c; McLean et al. [15] Fig. S2c), as we found that the association rule predicted EPIs

more accurately than the other rules in our previous study [15]. The maximum length of

the regulatory domain for each transcript was set to 1 Mbp upstream and downstream of

the TSS. Then, considering the genomic locations and effects of the insulator sites of a

DBP on EPIs, the length of a DBP ChIP-seq peak in a 100-base bin was calculated from

a TSS to the DNA-binding site of a predicted insulator-associated DBP. To reduce the

computational time of the analysis, we first calculated the length of a DBP ChIP-seq

peak in the regulatory domains of all transcripts in a 100-base bin. Then, for each

interval of 100 bins between a TSS and a DNA-binding site of a predicted

insulator-associated DBP in the regulatory domain, we computed the maximum length

of each DBP ChIP-seq peak in 100 bins on the basis of the precomputed length of the

DBP ChIP-seq peak in each bin. In the case of a set of 100 bins containing the

DNA-binding site of a predicted insulator-associated DBP, we computed the maximum

lengths from the bin closest to a TSS to the first bin containing the DNA-binding site of

the insulator-associated DBP, taking into account the orientation of the DNA-binding

site.

Prediction of insulator-associated DBPs using DeepLIFT tool

The DeepLIFT tool (version 0.6.13.0) was used to calculate the contribution

(DeepLIFT score) of each DNA-binding site to the prediction of the expression level of

a transcript [24]. The contribution scores of DNA-binding sites of DBPs in putative

enhancers and promoters upstream and downstream of each transcript were normalized

by the sum of the absolute values of all contribution scores for the transcript, as the

range of contribution scores changes according to the range of the predicted output (i.e.,

the expression level of a transcript), as described in a previous study [58]. To estimate the

effect of the DNA-binding sites of a predicted insulator-associated DBP on the

prediction of transcript levels, positive normalized contribution scores for transcripts

with DNA-binding sites in upstream or downstream regions were subjected to

comparisons between all combinations of two different orientations (i.e., all

combinations of two orientations among FR, RF, forward-forward [FF], and

reverse-reverse [RR] orientations, and no orientation [X]: FR vs. RF, FR vs. X, RF vs.

X, FF vs. RR, FF vs. X, RR vs. X, FF vs. FR, FF vs. RF, RR vs. FR, and RR vs. RF)

among the DNA-binding sites of a predicted insulator-associated DBP by using a

statistical test (Mann–Whitney U test, FDR-adjusted p value < 0.05 and p value $< 10^{-3}$).

For EPIs that were common to the predictions based on two different orientations of

DNA-binding sites of a predicted insulator-associated DBP, the DNA-binding sites of

DBPs in putative enhancers and promoters among the common EPIs were omitted in

the comparison of positive normalized contribution scores between the two different

orientations. Based on the results of the statistical analysis and the mean differences in

contribution scores, we predicted an insulator-associated DBP on the basis of the

directional bias or lack of orientation bias of DNA-binding sites; for example, to predict

an insulator-associated DBP with a DNA-binding site in the FR orientation, we selected

a DBP for which the contribution scores among all combinations of FR vs. RF, FR vs.

FF, FR vs. RR, and FR vs. X were significant (Mann–Whitney U test, FDR-adjusted p

value < 0.05 and p value $< 10^{-3}$) and showed positive (high) mean differences among

the contribution scores of all the combinations.

Search for insulator-associated DNA-binding sites based on histone modifications

and gene expression levels

The genomic positions in the BAM files of H3K27me3 and H3K9me3 ChIP-seq

data (ENCFF267JBO, ENCFF528UZJ) from HFF cells were converted to the

corresponding positions in the hg19 version of the human genome using the CrossMap

tool [63] and then converted to WIG files using bam2wig.pl from Bio::ToolBox

(https://github.com/tjparnell/biotoolbox) to visualize the data on the UCSC Genome

Browser through custom tracks (http://www.genome.ucsc.edu/cgi-bin/hgCustom) and a

track hub (https://genome.ucsc.edu/cgi-bin/hgHubConnect). The BAM file of RNA-seq

data (GSM941744) of HFF cells was converted to a WIG file using bam2wig.pl in

Bio::ToolBox. The genomic positions in the BED file of the boundaries and the

distribution of insulation scores identified through a Micro-C assay (4D Nucleome data

portal,   https://data.4dnucleome.org/files-processed/4DNFI5OZ7TBE/   and

4DNFIPMHBA41) were converted to the corresponding positions in the hg19 version

of the human genome.

Distribution of DNA-binding sites around chromatin interaction sites

We examined the distribution of the DNA-binding sites of predicted

insulator-associated DBPs in regions within $\pm$500 bp of the center of a chromatin

interaction site using chromatin interaction data from HFF cells obtained via a Micro-C

assay   (4D   Nucleome   data   portal,

https://data.4dnucleome.org/files-processed/4DNFI1MCF9GZ/).   For   the

insulator-associated DBPs other than CTCF, RAD21, and SMC3, the DNA-binding

sites that do not overlap with those of CTCF, RND21, and SMC3 were used to examine

their distribution around interaction sites. The number of DNA-binding sites of

predicted insulator-associated DBPs that cooccurred with DNA-binding sites of known

insulator-associated DBPs, such as CTCF, RAD21, and SMC3, was counted in each

10-bp bin in the regions within ±500 bp of the center of a chromatin interaction site.

Insulator-associated DNA-binding sites near splice sites

Insulator-associated DNA-binding sites were searched within 200 bp of the center

of splice sites using the file of the genomic positions of alternative transcripts in the

UCSC   Genome   Browser

(https://hgdownload.soe.ucsc.edu/goldenPath/hg19/database/all_mrna.txt.gz   ;

https://genome.ucsc.edu/cgi-bin/hgTables?db=hg19&hgta_group=rna&hgta_track=mrn

a&hgta_table=all_mrna&hgta_doSchema=describe+table+schema). We performed the

statistical test to estimate the enrichment of insulator-associated DNA-binding proteins

at the splicing sites compared with the other DNA-binding proteins (Mann–Whitney U

test, p value < 0.05).

Statistical analysis

Comparison of the distribution of DeepLIFT scores between different orientations

of DNA-binding sites was performed using the Mann-Whitney U test with the

scipy.stats.mannwhitneyu library. Two-tailed p values were calculated. For multiple

comparisons, adjusted p values were calculated by false discovery rate (FDR) using the

Benjamini-Hochberg procedure. Z scores for the Mann-Whitney U test were not output

by the library. Therefore, we calculated z scores by z-standardizing the U values. The

code is made available on GitHub at https://github.com/reposit2/insulator/tree/main.

Data availability

Human foreskin fibroblast (HFF) cell RNA-seq data were obtained from GSM941744.

Human monocyte RNA-seq data were obtained from the Blueprint DCC portal

http://dcc.blueprint-epigenome.eu/#/files   at

'C0010KB1.transcript_quantification.rsem_grape2_crg.GRCh38.20150622.results'. The

RNA-seq data of human mammary epithelial cells (HMEC) were found on ENCODE at

UCSC   http://genome.ucsc.edu/encode/   at

'wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam'. Human neural progenitor cell

(NPC) RNA-seq data were obtained from GSM915326. The DNase-seq data of HFF

cells were obtained from GSE170964. The *in situ* ChIA-PET chromatin interaction data

mediated by CTCF in HFF cells were available at the 4D Nucleome data portal,

https://data.4dnucleome.org/files-processed/4DNFI73AFN12/. The ChIP-seq data of

H3K4me3, H3K27ac, and H3K4me1 of HFF cells were found at the 4D Nucleome data

portal, 4DNFIJOK1BWR, 4DNFI442BQLO, and 4DNFIAGV3X99. The ChIP-seq data

of H3K27me3 and H3K9me3 in HFF cells were obtained from the ENCODE portal,

ENCFF267JBO and ENCFF528UZJ. The chromatin interaction data and boundary data

of HFF cells by the Micro-C were obtained from the 4D Nucleome data portal,

https://data.4dnucleome.org/files-processed/4DNFI1MCF9GZ/   and

https://data.4dnucleome.org/files-processed/4DNFI5OZ7TBE/. ChIP-seq data of TFs

were downloaded from the v20.06 release of the GTRD database

(https://gtrd.biouml.org). eQTL data (GTEx_Analysis_v8_eQTL.tar) were downloaded

from the v8 release of the GTEx portal (https://gtexportal.org). DNA binding motifs of

TFs were collected from TRANSFAC (2019.2), JASPAR (2018)

(https://jaspar.genereg.net/),   UniPROBE   (2018)

(http://the_brain.bwh.harvard.edu/uniprobe/),   high-throughput   SELEX

(https://doi.org/10.1016/j.cell.2012.12.009), transcription factor binding sequences of

ENCODE TF ChIP-seq data (http://compbio.mit.edu/encode-motifs/), and

HOCOMOCO version 9 and 11 (https://autosome.org/HOCOMOCOS/)

(https://hocomoco11.autosome.org/). The genomic positions of alternative transcripts in

the UCSC Genome Browser

(https://hgdownload.soe.ucsc.edu/goldenPath/hg19/database/all_mrna.txt.gz).

767 Code availability

768 Demonstration codes are made available at

769 https://github.com/reposit2/insulator/tree/main and

770 https://doi.org/10.5281/zenodo.8216164.

772 Figures

Fig. 1. CTCF insulator function mediates enhancer–promoter interactions. a CTCF

DNA-binding sites with a forward–reverse (FR) orientation are frequently observed at

chromatin interaction sites. The forward and reverse orientations correspond to

complementary sequences, which are clearly distinguishable in the reference genome. **b**

Position weight matrices (PWMs) of CTCF binding motifs. The forward and reverse

motifs are complementary on opposite DNA strands. A sequence logo is a graphical

representation of nucleotide conservation, where the height of each letter reflects its

frequency at that position (adapted from Wikipedia 'Sequence logo'). **c** CTCF can block

interactions between an enhancer and a promoter (or the transcription start site [TSS] of

a gene), thereby restricting enhancer activity within a defined chromatin domain (region

between red and green arrows). The red and green arrows indicate the DNA-binding

sites of a putative insulator-associated DBP, such as the FR orientation of CTCF sites

illustrated here. The numbers of DNA-binding sites with FR, RF, FF, and RR

orientations differ, as reflected by the intensity of chromatin loops (right panel). Images

of double-stranded DNA loops were adapted from "Transcriptional Regulation by

CTCF and Cohesin" at BioRender.com (2023).

Fig. 2. Overview of the prediction pipeline for insulator-associated DBPs. a We employed a machine learning method, DEcode, with modifications. Gene expression levels in human cells were predicted using the genomic locations of DBP binding sites at promoters and putative enhancers as input features. **b** In the machine learning phase, a DBP bound to a putative enhancer overlapping an eSNP (expression quantitative trait locus SNP) was linked to a gene (eGene) potentially regulated by the DBP, based on eQTL data. For each gene, DBP ChIP-seq peak locations were binned (green boxes) and

summarized in a matrix as input: Promoters were represented by 30 bins of 100 bp each

spanning the region from -2 kbp to +1 kbp relative to the TSS. Enhancer–gene

regulatory regions were covered by 200 bins of 10 kbp each, spanning -1 Mbp to +1

Mbp from the TSS, excluding the promoter. **c** For prediction of putative

insulator-associated DBPs, a DBP bound to a putative enhancer was associated with a

gene based on a computationally defined regulatory domain, independent of eQTL data.

The single nearest gene association rule defines a regulatory domain as the region

between the TSS of a gene and the midpoints to the TSS of the nearest upstream and

downstream genes. The TSSs and regulatory domains for each gene are indicated with

purple, light blue, or blue arrows and bracketed lines, respectively. The regulatory

domain for enhancer–promoter associations was further constrained by DNA-binding

sites (red and green arrows) of putative insulator-associated DBPs. For each gene, DBP

ChIP-seq peak locations were binned (green boxes in non-gray regions) and

summarized in a matrix as input data. **d** Estimation of the directional effect of

DNA-binding sites (red and green arrows) of putative insulator-associated DBPs on

gene expression prediction. A DBP whose binding (orange box) is highly informative

for gene expression prediction receives a high contribution score.

Fig. 3. Estimation of the effect of directional regulatory motifs on gene expression.

a The predicted log₂-transcripts per million (log₂-TPM) values of genes were compared with actual expression levels in human foreskin fibroblast (HFF) cells using Spearman's rank correlation coefficient. **b** The forward–reverse (FR) orientation of CTCF-binding sites is predominant (76.4%) at chromatin interaction sites, while forward–forward (FF),

reverse–reverse (RR), and reverse–forward (RF) orientations are less frequent (11.0%,

10.3%, and 2.3%, respectively) (Guo Y et al., Cell 2015) [3]. **c** Distribution of z-scores

from the Mann–Whitney U test for DeepLIFT scores comparing FR and RF orientations

of protein DNA-binding sites. The right panel displays the distribution of z-scores

comparing non-directional (X) sites with directional orientations (FR, RF, FF, or RR).

Results are shown according to Selection Criteria S2 in Supplementary Fig. 1 and Note.

d *Z*-scores from the Mann–Whitney U test for DeepLIFT scores in pairwise

comparisons of CTCF-binding sites with four different orientations (FR, RF, FF, RR)

and non-directional (X) sites. The differences in *z*-scores mirrored the percentages of

CTCF-binding site orientations at chromatin interaction sites. **e** The DNA-binding sites

of insulator-associated proteins were identified at both sites or at either site within pairs

of long-range chromatin interactions.

a

 Fig. 4. DNA-binding sites of predicted insulator-associated DBPs at potential

insulator sites. a Distribution of histone modification marks H3K27me3 (green) and

H3K9me3 (turquoise), along with transcript levels (pink), in the upstream and

downstream regions of a potential insulator site (light orange). These histone

modification marks are characteristic of transcriptionally repressed regions. Insulator

sites are positioned at the boundary between transcriptionally active and repressed

regions (indicated by light orange vertical bars). Blue and green lines in the GENCODE

track denote gene transcripts. Insulator scores and boundary regions were estimated

from chromatin interaction data available via the 4D Nucleome Data Portal. Dark blue

arcs represent chromatin interactions within 10 bp, 20 bp, 50 bp, and 200 bp of the

binding sites of CTCF (upper left panel), MAZ (upper right panel), and MyoD1 (lower

left panel), respectively. The binding sites of CTCF, RAD21, SMC3 (upper left), MAZ

(upper right), and MyoD1 (lower left) are displayed as black bars. Analyses were

performed using the Homo sapiens genome assembly GRCh37 (hg19) from the

Genome Reference Consortium. **b** Number of DNA-binding sites of predicted

insulator-associated DBPs where H3K27me3 signal intensities differed between

upstream and downstream regions. * indicates $p < 0.01$ (Mann–Whitney U test) (Table

2). Boxplots show the median (centerline), interquartile range (boxes), whiskers (up to

1.5× IQR), and all outliers (points). All data points are shown. **c** Number of

DNA-binding sites of predicted insulator-associated DBPs where both H3K27me3

signal intensities and transcript levels differed between upstream and downstream

regions. **d** Number of DNA-binding sites of predicted insulator-associated DBPs located

within boundary regions assigned by chromatin interaction data from human foreskin

fibroblast (HFF) cells. **e** Number of DNA-binding sites of predicted insulator-associated

DBPs overlapping within ±200 bp of splice sites. Insulator-associated DBP binding sites

were found within this window significantly more frequently than for other

DNA-binding proteins. **f** Cluster of DNA-binding sites of insulator-associated DBPs.

Fig. 5. Insulator-associated DBPs potentially regulate alternative transcription.

Distribution of the H3K27me3 histone modification mark (green) and transcript levels

(pink) is displayed. The H3K27me3 mark is characteristic of transcriptionally repressed

regions. Insulator sites are positioned at the boundaries between transcriptionally active

and repressed regions (indicated by light orange vertical bars). Blue lines in the

GENCODE track indicate gene transcripts. Chromatin interactions overlapping a

200-bp window centered on the binding sites of RBPJ (left panel) and SP1 (right panel)

are shown as dark blue arcs. The binding sites of RBPJ (left), SP1 (right), and CTCF,

RAD21, SMC3 (both panels) are shown as black bars. The DNA-binding sites of

insulator-associated DBPs were frequently observed near transcription start and

termination sites. These results suggest that insulator-associated DBPs may

differentially regulate transcribed regions that are repressed in alternative transcripts.

Fig. 6. Distribution of DNA-binding sites around chromatin interaction sites. a-f

The x-axis represents the distance of a DNA-binding site from the center of a chromatin

interaction site, while the y-axis indicates the number of chromatin interaction sites with

DNA-binding sites located within a ±500 bp region. For the analyses of YY1, ZNF143,

and MAZ, DNA-binding sites overlapping those of CTCF, RAD21, and SMC3 were

excluded.

Fig. 7. Boundary-pairing (insulator-pairing) model for the regulation of enhancer

–promoter interactions. Molecules bound to insulators physically pair with their

partners, either head-to-head or head-to-tail, with varying specificity at the termini of

topologically associating domains (TADs) in flies [45-47]. Although the precise mechanism

by which partners locate each other remains unclear, it is unlikely to require loop

extrusion. Homologous and heterologous insulator–insulator pairing interactions are

central to the architectural functions of insulators, and these interactions are

orientation-dependent. Homologous and heterologous interactions between

insulator-associated DNA-binding proteins may similarly occur in humans. Chromatin

looping can induce the formation of transcriptional condensates and phase separation, in

addition to cohesin–CTCF-anchored chromatin looping [40-42]. Figure created with

BioRender.com.

Tables

916 **Table 1. Predicted insulator-associated DNA-binding proteins (DBPs) with**

917 **directional or nondirectional binding site preferences.**

FR orientation					Low			High
DBP	z score				Mean difference			
	RF	FF	RR	X	RF	FF	RR	X
CEBPA	5.64	3.99	5.17	13.79	1.92E-05	2.41E-05	1.33E-05	1.47E-04
CTCF	30.51	22.70	22.14	4.15	1.30E-04	1.07E-04	1.22E-04	1.54E-05
CTCFL	23.39	17.21	21.19	14.70	1.03E-04	7.86E-05	1.27E-04	5.62E-05
EGR1	9.17	7.11	10.13	12.76	4.71E-05	3.81E-05	6.96E-05	7.41E-05
EP300	6.81	4.23	3.94	3.49	5.24E-05	2.39E-05	5.02E-05	1.42E-05
KLF9	9.11	6.92	4.52	17.23	7.56E-05	5.10E-05	6.36E-05	1.57E-04
RAD21	47.70	34.64	37.59	24.35	1.97E-04	1.80E-04	1.93E-04	1.34E-04
RBPJ	7.52	4.04	7.55	5.77	9.29E-05	8.10E-05	1.12E-04	9.77E-05
RUNX1	9.17	6.61	4.30	14.81	5.91E-05	4.42E-05	5.40E-05	1.87E-04
SIN3A	10.27	7.14	3.49	3.62	7.42E-05	6.50E-05	4.33E-05	2.03E-05
SMC3	31.72	23.76	25.66	24.07	1.60E-04	1.36E-04	1.70E-04	1.67E-04
ZEB1	3.53	4.32	3.85	7.06	2.67E-05	3.51E-05	3.69E-05	1.64E-04
ZNF18	8.62	9.08	4.47	7.69	8.16E-05	1.17E-04	6.78E-05	3.27E-04

FR orientation					Low			High
DBP	z score				Mean difference			
	RF	FF	RR	X	RF	FF	RR	X
CTCF	29.35	20.05	20.07	10.91	1.21E-04	9.10E-05	1.11E-04	4.12E-05
CTCFL	23.14	17.60	19.38	19.81	1.04E-04	8.77E-05	1.20E-04	9.44E-05
EGR1	10.80	8.16	10.01	13.69	5.96E-05	3.84E-05	8.50E-05	1.08E-04
EP300	9.77	7.57	7.91	10.66	6.67E-05	4.17E-05	7.85E-05	6.42E-05
KLF1	8.29	5.23	10.08	10.85	8.32E-05	4.23E-05	1.39E-04	1.49E-04
KLF9	11.70	7.66	5.01	12.95	8.88E-05	5.83E-05	6.72E-05	1.17E-04
MAZ	7.05	4.79	3.94	5.37	6.59E-05	2.47E-05	8.60E-05	6.87E-05
RAD21	40.64	31.19	29.65	25.69	1.73E-04	1.64E-04	1.61E-04	1.47E-04
RUNX1	8.47	5.66	3.50	13.54	6.70E-05	4.61E-05	6.52E-05	2.01E-04
SIN3A	13.18	6.66	4.99	4.91	1.02E-04	6.78E-05	6.17E-05	2.81E-05
SMC3	29.76	22.20	24.11	29.30	1.56E-04	1.30E-04	1.66E-04	2.10E-04
SP1	8.58	5.67	8.32	19.04	5.39E-05	4.78E-05	7.58E-05	2.02E-04
SP2	8.80	7.55	3.56	7.37	7.07E-05	8.81E-05	3.55E-05	1.81E-04
TAF1	6.57	4.74	5.15	4.91	1.20E-04	8.27E-05	1.35E-04	8.90E-05

Predicted insulator-associated DBPs

AHR	CTCF	FOS	JUND	MYB	PRDM9	SMAD3	TBX21	ZBTB7A
ASCL1	CTCFL	FOSL2	KLF1	MYCN	RAD21	SMC3	TCF12	ZEB1
ATF2	DUX4	FOXA3	KLF16	MYOD1	RBPJ	SP1	TCF3	ZFP3
ATF7	EBF1	FOXH1	KLF5	MYOG	REST	SP2	TCF4	ZIC2
BATF	EGR1	FOXK1	KLF8	NFYC	RUNX1	SP5	TEAD3	ZNF143
BATF3	EGR2	GATA2	KLF9	NR2C2	RUNX3	SPI1	TFAP2C	ZNF18
BCL6	EGR3	GRHL1	MAFG	NR2F2	RXRA	SPIB	TFAP4	ZNF320
CDX2	ELK4	GRHL2	MAZ	NR3C1	SCRT1	SREBF2	TP53	ZNF329
CEBPA	EP300	HNF4A	MECP2	NR5A2	SIN3A	STAT1	TRIM28	ZNF35
CLOCK	EPAS1	JUN	MEF2B	PAX5	SMAD1	STAT3	USF1	ZNF449
CREB5	ETV1	JUNB	MEIS2	PBX2	SMAD2	TAF1	VDR	ZZZ3

Z-scores from the Mann–Whitney U test compare DeepLIFT scores between

forward–reverse (FR) and other orientations—reverse–forward (RF), forward–forward

(FF), reverse–reverse (RR), and non-oriented (X)—for protein DNA-binding sites.

Mean differences in DeepLIFT scores are also reported. Color scales represent values

for each DBP. Rows (Top, Middle, Bottom) correspond to: Top—Selection Criteria S1

(Supplementary Fig. 1); Middle—Selection Criteria S2 (Supplementary Fig. 1);

Bottom—List of predicted insulator-associated DBPs. DBPs highlighted in yellow are

associated with insulator function, CTCF, and/or other transcriptional regulatory

mechanisms known to affect enhancer–promoter interactions (EPIs) in previous studies.

Table 2. Number of DNA-binding sites for predicted insulator-associated DBPs at

insulator sites and splice sites.

		Selection Criteria		Median number of DNA sites	Mean number of DNA sites	Number of DBP	p value
Experimental data for identifying insulator sites	H3K27me3	S1	Insulator-associated DBPs	66	115.66	58	2.35E-08
			Other DBPs	16	38.14	354	
		S2	Insulator-associated DBPs	65	102.00	50	3.33E-08
			Other DBPs	14	31.73	346	
		S3	Insulator-associated DBPs	45	93.43	40	3.20E-08
			Other DBPs	12	25.98	298	
		S4	Insulator-associated DBPs	40	69.65	37	1.13E-06
			Other DBPs	11	22.45	274	
	H3K27me3,Gene expression	S1	Insulator-associated DBPs	12	27.95	58	4.26E-06
			Other DBPs	4	10.04	354	
		S2	Insulator-associated DBPs	16	27.30	50	2.14E-06
			Other DBPs	4	8.28	346	
		S3	Insulator-associated DBPs	8	18.05	40	9.89E-06
			Other DBPs	2	5.09	298	
		S4	Insulator-associated DBPs	7	13.97	37	9.17E-05
			Other DBPs	2	4.46	274	
	Boundary of interactions	S1	Insulator-associated DBPs	48	96.36	58	6.43E-07
			Other DBPs	14	30.88	354	
		S2	Insulator-associated DBPs	54	91.58	50	5.39E-07
			Other DBPs	13	26.70	346	
		S3	Insulator-associated DBPs	26	74.33	40	4.57E-07
			Other DBPs	7	15.17	298	
		S4	Insulator-associated DBPs	16	58.08	37	1.27E-05
			Other DBPs	7	13.75	274	
Overlap with splice sites		S1	Insulator-associated DBPs	535	774.45	58	1.02E-06
			Other DBPs	132	350.11	354	
		S2	Insulator-associated DBPs	472	748.72	50	6.28E-06
			Other DBPs	122	290.21	346	
		S3	Insulator-associated DBPs	120	305.7	40	2.13E-07
			Other DBPs	35	99.23	298	
		S4	Insulator-associated DBPs	103	198.24	37	1.45E-05
			Other DBPs	30	81.01	274	

Selection Criteria:

S1: DNA-binding sites of predicted insulator-associated DBPs.

S2: DNA-binding sites of predicted insulator-associated DBPs that overlap with a

chromatin interaction site located more than 5 kb upstream or downstream of a

transcribed region.

S3: DNA-binding sites of predicted insulator-associated DBPs that do not overlap with

ChIP-seq peaks for H3K4me3, H3K4me1, or H3K27ac.

S4: DNA-binding sites of predicted insulator-associated DBPs that both overlap a

chromatin interaction site (located more than 5 kb upstream or downstream of a

transcript) and do not overlap ChIP-seq peaks for H3K4me3, H3K4me1, or H3K27ac.

P-values were calculated using the Mann–Whitney U test.

Table 3. Pairs of DBPs that colocalize at equivalent distances from chromatin

interaction sites.

DBP1	DBP2	Distance to start (bp)	Distance to end (bp)	# chromatin interactions with DBP1 sites	# chromatin interactions with DBP2 sites	# chromatin interactions with DBP1 and 2 sites	Ratio (DBP1 and 2) / DBP1	Ratio (DBP1 and 2) / DBP2	Max ratio (DBP1 or DBP2)
CTCF	CTCFL	30	39	36229	27159	19743	0.54	0.73	0.73
CTCF	PPARG	30	39	36229	2577	528	0.01	0.20	0.20
CTCF	RAD21	-10	-1	23108	19746	7661	0.33	0.39	0.39
CTCF	RXRA	220	229	33327	13204	9102	0.27	0.69	0.69
CTCF	SMC3	-30	-21	35558	24371	19701	0.55	0.81	0.81
CTCF	ZBTB7A	30	39	36229	5190	1539	0.04	0.30	0.30
YY1	CTCF	-50	-41	12717	35247	291	0.02	0.01	0.02
YY1	RAD21	-30	-21	13931	32793	203	0.01	0.01	0.01
ZNF143	CTCF	20	29	8993	35212	430	0.05	0.01	0.05
ZNF143	RAD21	-30	-21	9064	32793	500	0.06	0.02	0.06
BACH2	ATF3	290	299	2496	20052	634	0.25	0.03	0.25
BACH2	BACH1	-280	-271	2471	2863	730	0.30	0.25	0.30
BACH2	BATF	-250	-241	2709	5924	486	0.18	0.08	0.18
BACH2	FOS	-240	-231	2769	39887	884	0.32	0.02	0.32
BACH2	FOSB	-70	-61	2774	16108	602	0.22	0.04	0.22
BACH2	FOSL1	240	249	2589	19417	687	0.27	0.04	0.27
BACH2	FOSL2	260	269	2649	37682	837	0.32	0.02	0.32
BACH2	JUN	260	269	2649	36745	783	0.30	0.02	0.30
BACH2	JUNB	440	449	2519	23889	535	0.21	0.02	0.21
BACH2	JUND	260	269	2649	40876	925	0.35	0.02	0.35
BACH2	MAF	430	439	2500	1520	367	0.15	0.24	0.24
BACH2	MAFF	-220	-211	2676	3302	295	0.11	0.09	0.11
BACH2	MAFG	180	189	2514	2738	385	0.15	0.14	0.15
BACH2	MAFK	-30	-21	2726	7561	896	0.33	0.12	0.33
BACH2	MYC	90	99	2674	6900	417	0.16	0.06	0.16
BACH2	NFE2	-10	-1	2399	8289	1284	0.54	0.15	0.54
BACH2	NFE2L2	440	449	2519	4908	928	0.37	0.19	0.37

Distance to start and end (bp) indicates the genomic distance (in base pairs) from the

center of a chromatin interaction site to the 10-bp bin with the highest number of DBP1

and DBP2 binding sites, among 100 bins (covering a 1-kbp region) surrounding the site.

A positive value denotes the distance to the downstream side of an insulator site, while a

negative value indicates the upstream side. # chromatin interactions with DBP1 and

DBP2 sites shows the number of chromatin interaction sites that contain both DBP1 and

DBP2 binding sites within ±500 bp of the interaction center. Ratio (DBP1 and 2)/DBP2

refers to the ratio of chromatin interaction sites containing both DBP1 and DBP2

binding sites to the total number of interaction sites with DBP2 binding sites.

References

1 Weintraub, A. S. *et al.* YY1 Is a Structural Regulator of Enhancer-Promoter
Loops. *Cell* **171**, 1573-1588 e1528, doi:10.1016/j.cell.2017.11.008 (2017).
2 Rao, S. S. *et al.* A 3D map of the human genome at kilobase resolution reveals
principles of chromatin looping. *Cell* **159**, 1665-1680,
doi:10.1016/j.cell.2014.11.021 (2014).
3 Guo, Y. *et al.* CRISPR Inversion of CTCF Sites Alters Genome Topology and
Enhancer/Promoter Function. *Cell* **162**, 900-910, doi:10.1016/j.cell.2015.07.038
(2015).
4 Ortabozkoyun, H. *et al.* CRISPR and biochemical screens identify MAZ as a
cofactor in CTCF-mediated insulation at Hox clusters. *Nature genetics* **54**,
202-212, doi:10.1038/s41588-021-01008-5 (2022).
5 Goel, V. Y., Huseyin, M. K. & Hansen, A. S. Region Capture Micro-C reveals
coalescence of enhancers and promoters into nested microcompartments. *Nature*
genetics, doi:10.1038/s41588-023-01391-1 (2023).
6 Bailey, S. D. *et al.* ZNF143 provides sequence specificity to secure chromatin

interactions at gene promoters. *Nat Commun* **2**, 6186, doi:10.1038/ncomms7186
(2015).

7   Lis, M. & Walther, D. The orientation of transcription factor binding site motifs
in gene promoter regions: does it matter? *BMC Genomics* **17**, 185,
doi:10.1186/s12864-016-2549-x (2016).

8   Nora, E. P. *et al.* Molecular basis of CTCF binding polarity in genome folding.
Nat Commun **11**, 5612, doi:10.1038/s41467-020-19283-x (2020).

9   Li, Y. *et al.* The structural basis for cohesin-CTCF-anchored loops. *Nature* **578**,
472-476, doi:10.1038/s41586-019-1910-z (2020).

10   Pugacheva, E. M. *et al.* CTCF mediates chromatin looping via N-terminal
domain-dependent cohesin retention. *Proc Natl Acad Sci U S A* **117**, 2020-2031,
doi:10.1073/pnas.1911708117 (2020).

11   Tao, H. *et al.* Computational methods for the prediction of chromatin interaction
and organization using sequence and epigenomic profiles. *Brief Bioinform* **22**,
doi:10.1093/bib/bbaa405 (2021).

12   Zhang, K., Li, N., Ainsworth, R. I. & Wang, W. Systematic identification of
protein combinations mediating chromatin looping. *Nat Commun* **7**, 12249,
doi:10.1038/ncomms12249 (2016).

13   Xie, X. *et al.* Systematic discovery of regulatory motifs in human promoters and
3' UTRs by comparison of several mammals. *Nature* **434**, 338-345,
doi:10.1038/nature03441 (2005).

14   Tasaki, S., Gaiteri, C., Mostafavi, S. & Wang, Y. Deep learning decodes the
principles of differential gene expression. *Nat Mach Intell* **2**, 376-386,
doi:10.1038/s42256-020-0201-6 (2020).

15   McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory
regions. *Nature biotechnology* **28**, 495-501, doi:10.1038/nbt.1630 (2010).

16   Consortium, G. T. The GTEx Consortium atlas of genetic regulatory effects
across human tissues. *Science (New York, N.Y.)* **369**, 1318-1330,
doi:10.1126/science.aaz1776 (2020).

17   Sherwood, R. I. *et al.* Discovery of directional and nondirectional pioneer
transcription factors by modeling DNase profile magnitude and shape. *Nature*
biotechnology **32**, 171-178, doi:10.1038/nbt.2798 (2014).

18   Wingender, E., Dietze, P., Karas, H. & Knuppel, R. TRANSFAC: a database on

transcription factors and their DNA binding sites. *Nucleic acids research* **24**,
238-241 (1996).

19 Newburger, D. E. & Bulyk, M. L. UniPROBE: an online database of protein
binding microarray data on protein-DNA interactions. *Nucleic acids research* **37**,
D77-82, doi:10.1093/nar/gkn660 (2009).

20 Portales-Casamar, E. *et al.* JASPAR 2010: the greatly expanded open-access
database of transcription factor binding profiles. *Nucleic acids research* **38**,
D105-110, doi:10.1093/nar/gkp950 (2010).

21 Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell*
152, 327-339, doi:10.1016/j.cell.2012.12.009 (2013).

22 Kheradpour, P. & Kellis, M. Systematic discovery and characterization of
regulatory motifs in ENCODE TF binding experiments. *Nucleic acids research*
42, 2976-2987, doi:10.1093/nar/gkt1249 (2014).

23 Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of
transcription factor binding models for human and mouse via large-scale
ChIP-Seq analysis. *Nucleic acids research* **46**, D252-D259,
doi:10.1093/nar/gkx1106 (2018).

24 Shrikumar, A., Greenside, P. & Kundaje, A. in *Proceedings of the 34th*
International Conference on Machine Learning Vol. 70   (eds Precup Doina &
Teh Yee Whye) 3145--3153 (PMLR, Proceedings of Machine Learning Research,
2017).

25 Lee, B. H., Wu, Z. & Rhie, S. K. Characterizing chromatin interactions of
regulatory elements and nucleosome positions, using Hi-C, Micro-C, and
promoter capture Micro-C. *Epigenetics Chromatin* **15**, 41,
doi:10.1186/s13072-022-00473-4 (2022).

26 Cuddapah, S. *et al.* Global analysis of the insulator binding protein CTCF in
chromatin barrier regions reveals demarcation of active and repressive domains.
Genome research **19**, 24-32, doi:10.1101/gr.082800.108 (2009).

27 Huang, S., Li, X., Yusufzai, T. M., Qiu, Y. & Felsenfeld, G. USF1 recruits
histone modification complexes and is critical for maintenance of a chromatin
barrier. *Mol Cell Biol* **27**, 7991-8002, doi:10.1128/MCB.01326-07 (2007).

28 Crane, E. *et al.* Condensin-driven remodelling of X chromosome topology
during dosage compensation. *Nature* **523**, 240-244, doi:10.1038/nature14450

(2015).

29   Noda, Y., Okada, S. & Suzuki, T. Regulation of A-to-I RNA editing and stop
codon recoding to control selenoprotein expression during skeletal myogenesis.
Nat Commun **13**, 2503, doi:10.1038/s41467-022-30181-2 (2022).

30   Matharu, N. K. & Ahanger, S. H. Chromatin Insulators and Topological
Domains: Adding New Dimensions to 3D Genome Architecture. *Genes (Basel)*
6, 790-811, doi:10.3390/genes6030790 (2015).

31   Reyes, A. & Huber, W. Alternative start and termination sites of transcription
drive most transcript isoform differences across human tissues. *Nucleic acids*
research **46**, 582-592, doi:10.1093/nar/gkx1165 (2018).

32   Hua, P. *et al.* Defining genome architecture at base-pair resolution. *Nature* **595**,
125-129, doi:10.1038/s41586-021-03639-4 (2021).

33   Xiao, T., Li, X. & Felsenfeld, G. The Myc-associated zinc finger protein (MAZ)
works together with CTCF to control cohesin positioning and genome
organization. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2023127118
(2021).

34   Ishihara, K., Oshimura, M. & Nakao, M. CTCF-dependent chromatin insulator
is linked to epigenetic remodeling. *Molecular cell* **23**, 733-742,
doi:10.1016/j.molcel.2006.08.008 (2006).

35   Vierbuchen, T. *et al.* AP-1 Transcription Factors and the BAF Complex Mediate
Signal-Dependent Enhancer Selection. *Molecular cell* **68**, 1067-1082 e1012,
doi:10.1016/j.molcel.2017.11.026 (2017).

36   Gong, Q. H., McDowell, J. C. & Dean, A. Essential role of NF-E2 in remodeling
of chromatin structure and transcriptional activation of the epsilon-globin gene
in vivo by 5' hypersensitive site 2 of the beta-globin locus control region. *Mol*
Cell Biol **16**, 6055-6064, doi:10.1128/MCB.16.11.6055 (1996).

37   Davidson, I. F. & Peters, J. M. Genome folding through loop extrusion by SMC
complexes. *Nat Rev Mol Cell Biol* **22**, 445-464,
doi:10.1038/s41580-021-00349-7 (2021).

38   Karollus, A., Mauermeier, T. & Gagneur, J. Current sequence-based models
capture gene expression determinants in promoters but mostly ignore distal
enhancers. *Genome Biol* **24**, 56, doi:10.1186/s13059-023-02899-9 (2023).

39   Fulco, C. P. *et al.* Activity-by-contact model of enhancer-promoter regulation

1076   from thousands of CRISPR perturbations. *Nature genetics* **51**, 1664-1669,
1077   doi:10.1038/s41588-019-0538-0 (2019).

1078   40   Lee, R. *et al.* CTCF-mediated chromatin looping provides a topological
1079   framework for the formation of phase-separated transcriptional condensates.
1080   *Nucleic acids research* **50**, 207-226, doi:10.1093/nar/gkab1242 (2022).

1081   41   Amankwaa, B., Schoborg, T. & Labrador, M. <i>Drosophila</i>
1082   insulator proteins exhibit in vivo liquid-liquid phase separation properties. *Life*
1083   *Sci Alliance* **5**, doi:10.26508/lsa.202201536 (2022).

1084   42   Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N. & Mirny, L. A.
1085   Chromatin organization by an interplay of loop extrusion and compartmental
1086   segregation. *Proc Natl Acad Sci U S A* **115**, E6697-E6706,
1087   doi:10.1073/pnas.1717730115 (2018).

1088   43   Christou-Kent, M. *et al.* CEBPA phase separation links transcriptional activity
1089   and 3D chromatin hubs. *Cell Rep* **42**, 112897, doi:10.1016/j.celrep.2023.112897
1090   (2023).

1091   44   Ji, D. *et al.* FOXA1 forms biomolecular condensates that unpack condensed
1092   chromatin to function as a pioneer factor. *Molecular cell* **84**, 244-260 e247,
1093   doi:10.1016/j.molcel.2023.11.020 (2024).

1094   45   Bing, X. *et al.* Chromosome structure in Drosophila is determined by boundary
1095   pairing not loop extrusion. *eLife* **13**, doi:10.7554/eLife.94070 (2024).

1096   46   Ke, W., Fujioka, M., Schedl, P. & Jaynes, J. B. Stem-loop and circle-loop TADs
1097   generated by directional pairing of boundary elements have distinct physical and
1098   regulatory properties. *eLife* **13**, doi:10.7554/eLife.94114 (2024).

1099   47   Fujioka, M., Mistry, H., Schedl, P. & Jaynes, J. B. Determinants of Chromosome
1100   Architecture: Insulator Pairing in cis and in trans. *PLoS Genet* **12**, e1005889,
1101   doi:10.1371/journal.pgen.1005889 (2016).

1102   48   Aljahani, A. *et al.* Analysis of sub-kilobase chromatin topology reveals
1103   nano-scale regulatory interactions with variable dependence on cohesin and
1104   CTCF. *Nat Commun* **13**, 2139, doi:10.1038/s41467-022-29696-5 (2022).

1105   49   Hsieh, T. S. *et al.* Enhancer-promoter interactions and transcription are largely
1106   maintained upon acute loss of CTCF, cohesin, WAPL or YY1. *Nature genetics*
1107   **54**, 1919-1932, doi:10.1038/s41588-022-01223-8 (2022).

1108   50   Guerin, T. M., Barrington, C., Pobegalov, G., Molodtsov, M. I. & Uhlmann, F.

An extrinsic motor directs chromatin loop formation by cohesin. *EMBO J* **43**, 4173-4196, doi:10.1038/s44318-024-00202-5 (2024).

Wang, R. *et al.* MyoD is a 3D genome structure organizer for muscle cell identity. *Nat Commun* **13**, 205, doi:10.1038/s41467-021-27865-6 (2022).

Snead, W. T. & Gladfelter, A. S. The Control Centers of Biomolecular Phase Separation: How Membrane Surfaces, PTMs, and Active Processes Regulate Condensation. *Molecular cell* **76**, 295-305, doi:10.1016/j.molcel.2019.09.016 (2019).

Tang, X. *et al.* The PTM profiling of CTCF reveals the regulation of 3D chromatin structure by O-GlcNAcylation. *Nat Commun* **15**, 2813, doi:10.1038/s41467-024-47048-3 (2024).

Phillips, J. E. & Corces, V. G. CTCF: master weaver of the genome. *Cell* **137**, 1194-1211, doi:10.1016/j.cell.2009.06.001 (2009).

Kumar, S., Kaur, S., Seem, K., Kumar, S. & Mohapatra, T. Understanding 3D Genome Organization and Its Effect on Transcriptional Gene Regulation Under Environmental Stress in Plant: A Chromatin Perspective. *Front Cell Dev Biol* **9**, 774719, doi:10.3389/fcell.2021.774719 (2021).

Dekker, J. & Misteli, T. Long-Range Chromatin Interactions. *Cold Spring Harb Perspect Biol* **7**, a019356, doi:10.1101/cshperspect.a019356 (2015).

Spurrell, C. H., Dickel, D. E. & Visel, A. The Ties That Bind: Mapping the Dynamic Enhancer-Promoter Interactome. *Cell* **167**, 1163-1166, doi:10.1016/j.cell.2016.10.054 (2016).

Kim, D. S. *et al.* The dynamic, combinatorial cis-regulatory lexicon of epidermal differentiation. *Nature genetics* **53**, 1564-1576, doi:10.1038/s41588-021-00947-3 (2021).

Liu, M. *et al.* Genomic discovery of potent chromatin insulators for human gene therapy. *Nature biotechnology* **33**, 198-203, doi:10.1038/nbt.3062 (2015).

Sloan, C. A. *et al.* ENCODE data at the ENCODE portal. *Nucleic acids research* **44**, D726-732, doi:10.1093/nar/gkv1160 (2016).

Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments. *Bioinformatics (Oxford, England)* **28**, 2184-2185, doi:10.1093/bioinformatics/bts356 (2012).

Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation.

1142   *Nucleic acids research* **49**, D104-D111, doi:10.1093/nar/gkaa1057 (2021).

1143   63   Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between

1144   genome assemblies. *Bioinformatics (Oxford, England)* **30**, 1006-1007,

1145   doi:10.1093/bioinformatics/btt730 (2014).

1146

1147

Acknowledgements

We would like to thank Prof. Martin Frith for reviewing the manuscript. Supercomputing resources were provided by the Human Genome Center of the Institute of Medical Science at the University of Tokyo and ROIS National Institute of Genetics. This work was partially supported by JSPS KAKENHI (Grant Number 16K00387), JST CREST (Grant Number JPMJCR15G1), Osaka University start-up research grant, KIOXIA young scientist research grant at Waseda University, Sumitomo basic science research projects to NO, AMED [grant nos. JP21ae0121049, 21479280], and JSPS KAKENHI [grant nos. 22H04925, 20H00624, 17K20032] to MH.

Author contributions

N.O. designed and performed the research, and wrote and revised the manuscript. M.H. revised the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material

available at https://doi.org/ .

Correspondence and requests for materials should be addressed to Naoki Osato.