1 **The blast effector Pwl2 is a virulence factor that modifies the cellular localisation of**
2 **host protein HIPP43 to suppress immunity.**
3

Vincent Were[1], Xia Yan[1], Andrew J. Foster[1], Jan Sklenar[1], Thorsten Langner[5], Amber Gentle[1], Neha Sahu[1], Adam Bentham[1], Rafał Zdrzałek[2], Lauren Ryder[1], Davies Kaimenyi[1], Diana Gomez De La Cruz[1], Yohan Petit-Houdenot[3], Alice Bisola Eseola[1], Matthew Smoker[1], Mark Jave Bautista[1], Weibin Ma[1], Jiorgos Kourelis[6], Dan Maclean[1], Mark J. Banfield[2], Sophien Kamoun[1], Frank L.H. Menke[1], Matthew J. Moscou[1,4] and Nicholas J. Talbot[1] *

[1] The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.

[2] Department of Biochemistry and Metabolism, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.

[3] Université Paris-Saclay, INRAE, UR BIOGER, 91120, Palaiseau, France.

[4] USDA-ARS Cereal Disease Laboratory, University of Minnesota, 1551 Lindig Street, Saint Paul, Minnesota, 55108 USA.

[5] Max-Planck-Institute for Biology, 72076, Tübingen, Germany.

[6] Imperial College London, Department of Life Sciences – Faculty of Natural Sciences, London, SW7 2AZ, United Kingdom

*Corresponding author: Nicholas J. Talbot (Nick.Talbot@tsl.ac.uk)

Abstract

The rice blast fungus *Magnaporthe oryzae* secretes a battery of effector proteins to facilitate

host infection. Among these effectors, Pwl2 was first identified as a host specificity

determinant for infection of weeping lovegrass (*Eragrostis curvula*) and is also recognised by

the barley Mla3 resistance gene. However, its biological activity is not known. Here we show

that *PWL2* expression is regulated by the Pmk1 MAP kinase during cell-to-cell movement by

M. oryzae at plasmodesmata (PD)-containing pit field sites. Consistent with its regulation, we

provide evidence that Pwl2 binds to a barley heavy metal-binding isoprenylated protein

HIPP43, which results in its displacement from plasmodesmata. Transgenic barley lines

overexpressing either *PWL2* or HIPP43 exhibit attenuated immune responses and increased

disease susceptibility. By contrast, a Pwl2^{SNDEYWY} mutant that does not interact with HIPP43,

fails to alter the PD localisation of HIPP43. Targeted deletion of three copies of *PWL2* in *M.*

oryzae results in a *Δpwl2* mutant showing gain-of-virulence to weeping lovegrass and barley

Mla3 lines, but also a reduction in severity of blast disease on susceptible host plants. Taken

together, our results provide evidence that Pwl2 is a virulence factor that acts by suppressing

host immunity through perturbing the plasmodesmatal deployment of HIPP43.

Introduction

Plant pathogens secrete virulence proteins called effectors during infection to suppress plant

immunity and facilitate infection (Jones and Dangl, 2006). Fungal pathogens, such as the

devastating blast fungus *Magnaporthe oryzae*, utilise an extensive battery of more than 500

effectors (Wilson and Talbot, 2009; Yan *et al.*, 2023), but very few have been functionally

characterised (Oliveira-Garcia *et al.*, 2023a). A subset of effectors are recognized by plant

intracellular nucleotide-binding leucine rich repeat (NLR) immune receptors to activate disease

resistance. In rice for example, the paired NLRs RGA5/RGA4, Pik-1/Pik-2, and Piks-1/Piks-2

confer resistance to *M. oryzae* strains that secrete AVR1-CO39/AVR-Pia, AVR-Pik or AVR-

Mgk1 effectors, respectively (Ashikawa *et al.*, 2008; Kanzaki *et al.*, 2012; Cesari *et al.*, 2013;

Ortiz *et al.*, 2017; Zhang *et al.*, 2018; Sugihara *et al.*, 2023). These effectors are members of

the *Magnaporthe* Avrs and ToxB-like (MAX) family which are sequence-unrelated,

structurally conserved (de Guillen *et al.*, 2015) and overrepresented among effectors

recognised by rice NLRs (Cesari *et al.*, 2013; Maqbool *et al.*, 2015). Pwl2 is a MAX effector

first identified as a host specificity determinant that controls pathogenicity towards weeping

lovegrass (*Eragrostis curvula*), a widely-grown forage grass (Sweigard *et al.*, 1995), and was

recently shown to be recognised in barley by the Mla3 immune receptor (Brabham *et al.*, 2023).

While Pwl2 has been widely studied to understand effector secretion and delivery (Mosquera

et al., 2009; Zhang and Xu, 2014; Oliveira-Garcia *et al.*, 2023b), the biological function of

Pwl2 is unknown.

Recent studies of MAX-effector perception by NLRs have implicated host small heavy metal-

associated (sHMA) domain containing proteins as potential targets of some of these effectors,

including Pwl2 (Zdrzałek *et al.*, 2024). Small HMAs are highly expanded across plant species

with putative functions including heavy metal detoxification and potential metallochaperones,

but in most cases their functions are not known (Dykema *et al.*, 1999; Suzuki *et al.*, 2002; Gao

et al., 2009). HMAs can be broadly classified into two families, heavy metal-associated plant

proteins (HPPs), and heavy metal-associated isoprenylated plant proteins (HIPPs), which

possess a C-terminal isoprenylation motif (CaaX, where 'a' represents an aliphatic residue and

'X' is any amino acid) important for membrane anchoring (Hála and Žárský, 2019). The rice

sHMA protein Pi21, for example, is a blast disease susceptibility factor (Fukuoka *et al.*, 2009),

which has led to deployment of loss-of-function alleles of *pi21*, as a recessive form of rice blast

resistance (Fukuoka *et al.*, 2009; Mutiga *et al.*, 2021). Furthermore, differences in host sHMA

protein repertoires are linked to host-specific, adaptive evolution of the APikL2 effector family

(Bentham *et al.*, 2021). Importantly, HMA protein domains have been identified as non-

canonical immune sensory domains in some NLR proteins, such as the rice Pik-1 of the Pik-

1/2 and RGA5 of the RGA5/4 pair receptors respectively, essential for recognition of cognate

M. oryzae effectors (Ashikawa *et al.*, 2008; Kanzaki *et al.*, 2012; Cesari *et al.*, 2013; Maqbool

et al., 2015). However, the function of sHMAs and their link to disease susceptibility is not

understood, limiting our understanding of the role of MAX effectors in plant disease.

In this study we set out to investigate the biological function of *PWL2*. We were motivated to

understand how a broadly distributed effector such as Pwl2 functions during a susceptible

interaction between *M. oryzae* and its host. We report that *PWL2* has undergone extensive

duplication and expansion in copy number with many rice blast isolates possessing 3-5 copies,

making its functional analysis challenging. Using CRISPR-Cas9 gene editing, however, we

have generated a *Δpwl2* mutant confirming that Pwl2 is a host specificity determinant but also

revealing a hitherto un-recognized virulence function. Furthermore, we demonstrate here that

PWL2 expression during infection is controlled by the Pmk1 MAP kinase, which regulates cell-

to-cell movement by the fungus at plasmodesmata-containing pit fields (Sakulkoo et al., 2018).

We reveal that Pwl2 targets the HIPP43 sHMA protein in barley, thereby displacing it from

plasmodesmata and show that transgenic plants overexpressing either *PWL2* or HIPP43 have

reduced immune responses and greater susceptibility to blast disease. Finally, we show that a

Pwl2[SNDEYWY] mutant unable to interact with HIPP43 fails to displace it from PDs and cannot

complement the reduced virulence of *Δpwl2* mutants. When considered together, our study

provides evidence that Pwl2 is a virulence factor in *M. oryzae* that suppresses host defense by

re-localising HIPP43 away from PDs to facilitate fungal invasion of plant tissue.

Results

Pwl2 is a cytoplasmic effector expressed during blast infection.

Expression of *PWL2* is specific to the initial biotrophic phase of plant infection, peaking at 48

hours post-infection (hpi) (Yan *et al.*, 2023) (**Fig. S1A - C**). The Pwl2 effector localises in the

biotrophic interfacial complex (BIC), a plant membrane-rich structure –that is clearly visible

as a single bright punctum, initially at the tip of a penetration hypha and then adjacent to

bulbous, branched invasive hyphae (Khang *et al.*, 2010). The BIC is the predicted site of

effector delivery (Kankanala *et al.*, 2007; Giraldo *et al.*, 2013; Oliveira-Garcia *et al.*, 2023b;

Were and Talbot, 2023), although experiments to date have not precluded that the BIC could

be a site of effector sequestration from the host plant. To investigate this possibility, we

generated a single *M. oryzae* strain expressing two BIC-localised effectors Pwl2-mRFP and

Bas1-GFP and visualised their localisation during infection of leaf sheath tissue in the

susceptible rice cultivar Moukoto. The two effectors were observed as small punctate signals

within the same BIC (**Fig. 1A and B**). By contrast, when two different *M. oryzae* strains,

Ina168 and Guy11, expressing Pwl2-GFP and Pwl2-mRFP respectively were used to

simultaneously infect rice tissue, we observed that the BIC formed by each individual invasive

hypha exclusively contained either Pwl2-GFP or Pwl2-mRFP respectively but never both

fluorescence signals (**Fig. 1C and D).** This is consistent with Pwl2 secretion by each fungal

strain into the BIC, because sequestration of previously secreted Pwl2 from the plant cytoplasm

to the BIC would result in a mixed GFP/mRFP fluorescence signal in the BIC. We conclude

that Pwl2 is expressed early during infection and secreted to the BIC from where it is delivered

into host cells.

***PWL2* expression is regulated by the Pmk1 MAP kinase during invasive growth.**

The blast fungus invades rice tissue by means of pit field sites containing plasmodesmata

(PDs), which enable it to move between rice cells while maintaining integrity of the rice plasma

membrane. PD conductance is also regulated by the fungus, enabling effectors like Pwl2 to be

deployed in adjacent uninfected cells (Kankanala et al., 2007; Sakulkoo et al., 2018). Cell-to-

cell movement by the fungus is regulated by the pathogenicity mitogen-activated protein kinase

1 (Pmk1 MAPK) pathway. An analogue-sensitive mutant of Pmk1 has been shown to be unable

to move through pit fields in the presence of the MAPK inhibitor 1NA-PP1 (Sakulkoo et al.,

2018). Given that *PWL2* is expressed during the initial stages of infection, we decided to test

whether it is regulated by Pmk1. We therefore re-analyzed RNA-seq data (**Fig. 1E**) (Sakulkoo

et al., 2018) by separating raw reads of *M. oryzae* and *O. sativa* and quantifying transcript

abundance using Kallisto, followed by determining differential expression using Sleuth. We

found that *PWL2* is significantly down-regulated in a *M. oryze pmk1^{AS}* mutant during cell-to-

cell movement in the presence of 1NA-PP1 together with a subset of known effector genes -

BAS83, *BAS52*, *BAS2, AVR-Pita-1, BAS3, BAS4, BAS162* and *AVR-Pik-C* (**Fig. 1F***)*. By

contrast, *MoNLP1* showed up-regulation, while expression of actin (MGG_03982) and the

RP27 40S 27a ribosomal subunit genes (MGG_02872) were not significantly affected (**Fig.**

1F*)*. We then used live cell imaging to investigate Pwl2-GFP expression in a *pmk1^{AS}* mutant

during rice leaf sheath infection ±1Na-PP1. We initially observed Pwl2-GFP in the BIC at early

stages of infection, unaffected (Stage 1) 30-36hpi (**Fig. 1G and H**), as previously reported

(Sakulkoo *et al.*, 2018)*,* but by later stages of infection Stage 2 (36-40hpi), (**Fig. 1I-J**), and

Stage 3 (40-48hpi) (**Fig. 1K-L**), 14-22h after 1NA-PP1 treatment, Pwl2-GFP fluorescence was

significantly reduced (**Fig. 1J, L and M**). The reduction in Pwl2-GFP fluorescence was

consistently associated with the stage at which *M. oryzae* traverses PD-containing pit field sites

and enters adjacent plant cells. Taken together, we conclude that Pwl2 is regulated by the

Pmk1 MAPK signalling pathway during cell-to-cell invasive growth by *M. oryzae*.

PWL2 is highly conserved in *M. oryzae.*

PWL2 belongs to a large gene family (Kang *et al.*, 1995; Sweigard *et al.*, 1995), but its

conservation in the global rice blast population is not known. We therefore investigated *PWL*

gene family distribution in isolates that infect a variety of different grass species (**Fig. 2A**) This

revealed that *PWL2* is found in the majority of host-limited forms of *M. oryzae* and related

Magnaporthe species, except for *Setaria* and closely-related *Panicum, Cynodon* and *Urochloa*-

infecting isolates (although these were less well represented than other isolates) as shown in

Fig. 2A. By contrast, *PWL1* is present in a sub-set (Asuke *et al.*, 2020) of *Eleusine*-infecting

isolates and some *Oryza*-infecting isolates, but largely absent from other host-specific lineages,

except one *Eragrostis*-infecting isolate (EtK19-1), and one *Cynodon*-infecting isolate

(Cd88215) (**Fig. 2A**). Similarly, *PWL3* is present in most *Oryza*-infecting isolates, some

Setaria, *Lolium*, and *Eleusine*-infecting isolates, but largely absent in *Eragrostis, Triticum*,

Digitaria and *Pennisetum* lineages, while *PWL4* is present in *Eleusine*, *Eragrostis, Lolium* and

Triticum-infecting isolates, *Pennisetum* and *Digitaria*-infecting isolates, but only two *Oryza*-

infecting isolates (**Fig. 2A**). We conclude that the *PWL* gene family (Kang *et al.*, 1995) is

broadly distributed among blast fungus isolates infecting numerous grasses but that *PWL2*, is

widespread across rice-infecting isolates and the majority of host-limited forms of *M. oryzae.*

Having determined that *PWL2* is broadly distributed in *M. oryzae*, we next sought to determine

its allelic variability. In addition to a loss of recognition allele *pwl2-2*, which contains a single

Asp-90-Asn substitution (Sweigard *et al.*, 1995), we identified 14 new alleles of *PWL2* (**Fig.**

S2A). Notably, most polymorphic residues occur between positions His87 to Ser92, suggesting

these residues might contribute to Pwl2 recognition by a cognate resistance gene (**Fig. S2A**).

Interestingly, we could only identify 1 variant of *PWL1*, 5 variants of *PWL3* and 1 variant of

PWL4, despite these effector genes occurring in finger millet, rice, wheat and ryegrass-

infecting lineages, respectively (**Fig. 2A and S2B - S2D**). It is possible that sample size for

some host-specific lineages may lead to an underestimate of allelic variability in *PWL1, PWL3,*

and *PWL4,* but from this analysis *PWL2* appears to be highly polymorphic and conserved,

compared to other members of the gene family (**Fig. S2E**). We also tested the ability of a subset
of *PWL2* alleles to be recognised by barley *Mla3*. In some cases, we found variant *PWL2* alleles
occurred in a *M. oryzae* isolate carrying *PWL1* (such as U34 and E39) or isolates with multiple
copies of *PWL2* (for example TH3), precluding functional analysis, and not all isolates were
available for testing. As expected, Guy11 did not produce lesions on cv. Baronesse, but was
able to infect cv. Nigrate (**Fig. S3A – C**). However, JUM1 (*pwl2-2*), BTJP4-16 (*pwl2-3)* and
BN0293 (variant *pwl2*) all caused blast disease on cv. Baronesse, confirming that they contain
loss of recognition *pwl2* alleles (**Fig. S3A - C**). Pwl2 has recently been structurally described
as a MAX effector, containing a core β-sandwich fold formed of two antiparallel β-sheets, as
well as a single α-helix, C-terminal to the β-sandwich fold (Zdrzałek *et al.*, 2024). We were
interested to understand how these variable residues affect Pwl2 recognition. We therefore
mapped a subset of the identified variants of Pwl2 to the effector crystal structure. We noted
that all polymorphic residues in *pwl2* alleles, such as the Glu-89-Gln, Lys-91-Gln and Ser-92-
Ile substitutions in *pwl2-3,* were present in the C-terminal α-helix, a distinct region from the
core MAX-fold (**Fig. 2B**). We conclude that this interface is likely used by immune receptors
present in resistant barley (Mla3) and weeping lovegrass to bind Pwl2 and is potentially
stabilized by the MAX-fold.

***PWL2* has undergone copy number expansion in *M. oryzae*.**
PWL2 copy number appears to vary significantly among *M. oryzae* isolates. The *M. oryzae*
reference genome assembly (strain 70-15) has two copies of *PWL2* on chromosomes 3 and 6,
annotated as MGG_04301 and MGG_13683, respectively (Dean *et al.*, 2005). Southern blot
hybridization (**Fig. S4**), however, and analysis of long-read assembled genome sequence of *M.*
oryzae Guy11 identified three copies of *PWL2* (**Fig. 3A)**. *PWL2* loci were associated with
MGR583, *POT2* and *MAGGY* transposable elements, suggesting their potential involvement in
genome rearrangements (**Fig. 3A**). To further assess *PWL2* copy number variation in *M. oryzae*
isolates, we employed k-mer analysis to determine copy number variation of *PWL2* in 286 *M.*
oryzae genomes. Copy number variation is common among effector-encoding genes, such as
AVR-Pik, *AVR-Pizt, BAS1, BAS4* and *SLP1*, but was particularly prevalent for *PWL2* with one
isolate, for instance, containing 9 copies (**Fig. 3B).** We conclude that *PWL2* has expanded in
copy number in *M. oryzae*.

Pwl2 is both a host range determinant and a virulence factor

We next set out to determine the function of *PWL2* in blast disease through gene functional analysis. Given that *PWL2* occurs in multiple copies, we used CRISPR/Cas9 genome editing (Foster *et al.*, 2018) to simultaneously delete all three copies of *PWL2* found in Guy11. We designed a sgRNA to target *PWL2* and introduced a hygromycin phosphotransferase (*HPH*) encoding gene cassette (**Fig. 4A**). Transformants with ectopic integrations, or where some *PWL2* had not been deleted, gave a predicted PCR amplicon of 645 bp in size. By contrast, *Δpwl2* mutants generated a larger amplicon of 1.5 kb (**Fig. 4A**). Four putative transformants were selected for whole genome sequencing. Raw reads were aligned to the *M. oryzae* reference genome using samtools v.15, before converting into BAM files and visualising using IGV viewer. No reads mapping to the *PWL2* locus were identified in mutants T6 and T12, whereas fewer reads mapped to the same locus from sequenced T4 and T5 (**Fig. S5A**). This confirmed that T6 and T12 are *Δpwl2* transformants. Transformant T4, T5 and T6 displayed vegetative growth that was identical to Guy11 with normal dark concentric rings and light growing edges, while T12 showed slight reduced growth, melanisation and conidiation (**Fig. S5B**). Given that *PWL2* is a host range determinant and avirulence gene, we reasoned that *M. oryzae Δpwl2* deletion mutants would gain virulence on weeping lovegrass (Kang *et al.*, 1995; Sweigard *et al.*, 1995) and barley cv. Baronesse expressing *Mla3* (Brabham *et al.*, 2023). The *Δpwl2* deletion transformants T5, T6 and T12 and Guy11 were therefore used to infect weeping lovegrass seedlings and barley Baronesse using spray inoculation. The *Δpwl2* mutant T6 produced disease symptoms identical to those produced by *Eragrostis*-infecting isolate G17 and *Oryza*-infecting isolate Ina168 which lacks *PWL2,* while Guy11 and the two complemented isogenic strains (T6+*PWL2*p:*PWL2*) and (T6+*RP27*p:*PWL2*) were unable to cause disease (**Fig. 4B**). Similarly, T5 and T12 also produced disease symptoms on weeping lovegrass seedlings and cv. Baronesse (**Fig. S5C**). We conclude that T5, T6 and T12 are loss of recognition mutants of *PWL2* (**Fig. 4B and S5C**)*,* and we can be confident that T6 and T12 have complete deletion of all three copies of the gene.

We next tested whether Pwl2 has a virulence function during a compatible interaction between the blast fungus and its host. Having observed that T12 had slight differences in vegetative growth, we decided to select T6 for virulence assays and as a genetic complementation background for *PWL2*. T6 was used to inoculate susceptible rice cultivar CO39 by spray infection, and on susceptible barley lines cv. Nigrate (-*Mla3)*, Siri (+*Mla8*), cv. Golden Promise

(+*Mla8*), and resistant cv. Baronesse (+*Mla3*) using leaf drop inoculation. In four independent

replicates, the *Δpwl2* mutant T6 showed a reproducible reduction in virulence on CO39

compared to wildtype Guy11 or a complemented isogenic strain (T6+*PWL2*p:*PWL2*), based

on lesion density and lesion length (**Fig. 4C - E)**. As expected, T6 was able to cause blast

disease on cv. Baronesse unlike Guy11 and produced statistically smaller lesion sizes on barley

lines Siri and Golden Promise, compared to Guy11. Both strains were able to infect Nigrate

(**Fig. 4F and G**). We conclude that *PWL2* contributes to the ability of *M. oryzae* to cause blast

disease.

Pwl2 suppresses host immunity

Having determined that the Pwl2 effector contributes to virulence, we decided to study its

biological function during host cell colonisation. To do this we first generated stable transgenic

barley lines (in cv. Golden Promise) expressing *PWL2-YFP* (without its signal peptide) under

control of the CaMV35S promoter (**Fig. 5A)**. We tested two independent *PWL2-YFP*

transgenic plants for their response to two elicitors of PAMP-triggered immunity (PTI), flg22

and chitin, compared to wild type cv. Golden Promise (**Fig. 5A**). Barley perceives flg22

through the pattern recognition receptor FLS2 and chitin through HvCEBiP and HvCERK1,

leading to immune responses such as generation of reactive oxygen species (ROS) (Yu *et al.*,

2023). We observed that Pwl2 expression abolished flg22 (**Fig. 5B - C)** and chitin-induced

ROS generation (**Fig. 5D - E)**. Pwl2 therefore contributes to virulence by suppressing PTI

during compatible interactions. To investigate the effect of elevated Pwl2 effector expression

on blast infection, we infected two independent transgenic barley plants expressing *PWL2* with

M. oryzae Guy11. *PWL2*-expressing barley lines were more susceptible and developed blast

disease symptoms earlier (2-3 dpi) compared to infection of isogenic cv. Golden Promise (**Fig.**

5F and G). We conclude that Pwl2 acts as a modulator of PTI that helps facilitate fungal

infection.

Pwl2 interacts with the heavy metal binding isoprenylated protein HIPP43.

To determine the likely target of Pwl2, transgenic barley lines expressing Pwl2-YFP were used

in co-immunoprecipitation coupled to mass spectrometry (IP-MS) analysis. This was followed

by spectral search in the *H. vulgare* (barley, cv Morex version 3) proteome database. We

identified a total of 282 frequently occurring proteins in the eight biological replicates

containing Pwl2, that were not identified in the six free-YFP control samples. To avoid

focusing on false positives in the form of sticky and abundant proteins, fold-changes were

calculated by first determining the average number of peptides per protein candidate, before

estimating $\log_2$ fold change compared to the control experiment. We further filtered for proteins

that produced at least two peptide hits in more than half of replicates ($\geq$4/8 biological

replicates). A total of 52 proteins met this criterion and were considered as enriched in samples

compared to controls and were therefore selected for further analysis. Furthermore, because

Pwl2 attenuates the ROS-burst in barley transgenic lines, we focused initially on protein

candidates previously reported to have potential roles in immunity or ROS generation (**Fig.**

6A). These were selected for one-to-one interactions using a yeast-two-hybrid (Y2H) assay by

co-transforming constructs expressing Pwl2 and P̲utative P̲wl2 I̲nteracting P̲roteins (PPIPs)

identified by IP-MS (**Fig. 6A)**. Pwl2 showed a strong interaction with PPIP4, a HMA-domain

protein (**Fig. 6B)**. We named this protein *Hv*HIPP43, for *Hordeum vulgare* heavy metal domain

containing isoprenylated plant protein 43 based on homology to *Os*HIPP43 (Zdrzałek *et al.*,

2024). We also tested to ensure that neither Pwl2 nor HIPP43 demonstrated auto-activity in

yeast-2-hybrid drop out media (**Fig. S6A)**. To independently verify the interaction between

Pwl2 and HIPP43, we carried out co-immunoprecipitation (co-IP) analysis using protein

extracts from *N. benthamiana* and confirmed that *Hv*HIPP43 interacts with Pwl2-YFP (**Fig.**

6C and D). Conversely, three other MAX-fold effectors, MEP3, AVR-PikE, AVR-Piz-t, or

free YFP did not interact with HIPP43 (**Fig. 6D)**. We next tested whether *PWL* gene family-

encoded proteins Pwl1, Pwl3, Pwl4 and the pwl2-3 variant could also interact with *Hv*HIPP43

and showed that they can all interact based on Y2H assays, suggesting that Pwl2 belongs to an

effector family that interacts with a host sHMA protein (**Fig. 6E)**. By contrast, three MAX-

fold effectors AVR-Pia, AVR-PikD and AVR-Mgk did not interact with HIPP43 (**Fig. 6E)**.

Barley and wheat have three copies of HIPP43 per haploid genome, but based on our IP-MS

results, we identified peptides that mapped to two copies of HIPP43, which both strongly

interacted with Pwl2 in Y2H assays HIPP43 (**Fig. S6B-C)**.

Pwl2 can interact with HIPP43 orthologs of diverse grass species.

HIPPs are expanded in plant genomes and to test whether HIPP43 has orthologs in other grass

species, we generated a maximum likelihood phylogenetic tree of HMA domains from diverse

grasses with high quality genomes and annotations (**Fig. S6D)**. The HIPP43 family formed a

distinct clade which includes orthologs from rice (*O. sativa)*, wheat (*T. aestivum)*, wild wheat

(*Haynaldia villosa*), foxtail millet (*S. italica)*, weeping lovegrass (*E. curvula)*, Sorghum

(*Sorghum bicolor*), *Oropetium thomaeum*, *Zea mays* and *Brachypodium distachyon* (**Fig.**

S6D). Furthermore, we could identify copy number variation ranging from one to five copies

in grasses, with barley and wheat (i.e. *Triticeae* lineage) having three or more paralogs of

HIPP43, one per haploid genome (**Fig. S7A**). We found that Pwl2 interacts with *Hv*HIPP43

orthologs from rice (*Os*HIPP43), *Triticum* (*Taes*HIPP43), *Setaria* (*SitiHIPP43*), *Eragrostis*

(*Ec*HIPP43) and *Haynaldia* (HIPP1-V) in Y2H assays (**Fig. 6F**). The interaction of Pwl2 with

rice *Os*HIPP43 was verified by *in vitro* biochemistry and a structure of the complex was

obtained by *X*-ray crystallography (Zdrzałek *et al.*, 2024). We also analyzed the transcriptional

profile of *Os*HIPP43 during infection of Guy11 in a susceptible rice line CO39 (Yan *et al.*,

2023) and found that *Os*HIPP43 expression is up-regulated during plant infection, consistent

with a role in host defense (**Fig. S7B**).

Overexpression of HIPP43 suppresses PTI and increases blast susceptibility.

To investigate the role of the Pwl2-HIPP43 interaction, we generated barley transgenic lines

overexpressing *Hv*HIPP43. Two independent transgenic lines were challenged with two PTI

elicitors, flg22 (**Fig. 7A-B)** and chitin (**Fig. 7C-D)**. Strikingly, we observed that both the chitin

and flg22-induced ROS burst was abolished in plants expressing *Hv*HIPP43, compared to wild

type (**Fig. 7A-D)**. Similarly, when we challenged transgenic barley plants expressing

*Hv*HIPP43 with *M. oryzae*, we observed increased susceptibility compared to the wildtype cv.

Golden Promise, and disease lesions appeared earlier than in wild type plants (**Fig. 7E and F)**.

These findings are consistent with over-expression of both *PWL2* and HIPP43 leading to

enhanced blast disease susceptibility.

Pwl2 prevents plasmodesmatal localisation of HIPP43.

To investigate the function of *Hv*HIPP43, we performed live cell imaging following transient

expression in *N. benthamiana* to determine its sub-cellular localisation. When tagged at the N-

terminus (mCherry-HIPP43), *Hv*HIPP43 localised predominantly to puncta at the plasma

membrane that co-localise with callose staining (aniline blue), consistent with sites of

plasmodesmata (PD) (**Fig. 7G and H**). Like all HIPPs, *Hv*HIPP43 has an isoprenylation motif

at its C-terminus (**Figure S8A and B**), implicated in membrane anchoring (Hála and Žárský,

2019), so we investigated whether PD localisation is affected by its deletion. We observed that

PD localisation was indeed abolished in the absence of the isoprenylation motif and -Iso-

mCherry-HIPP43 (*Hv*HIPP43 without isoprenylation motif) instead accumulated in the

nucleoplasm and at the plasma membrane. Moreover, co-expression with a plasma membrane

marker LTi6b-GFP and aniline blue staining confirmed its localisation at the plasma membrane

and loss of PD localisation (**Fig. S8C**) compared to the wild type (**Fig. S8D**).

Given the strong interaction between Pwl2 and HIPP43 and the localisation of HIPP43 to PD,

we were keen to see the effect of co-expressing the effector and putative host target in the same

cells. When we co-expressed YFP-Pwl2 with mCherry-HIPP43, they co-localised in the

cytosol as mobile puncta approximately 2 μm in diameter (**Fig. 7I**). In some cases, we observed

larger mobile structures ranging from 2-4 μm in diameter (**Fig. 7I**). We also tested whether

Pwl1, Pwl3 and Pwl4 (**Fig. S9A**) could also co-localise with HIPP43 when co-expressed in *N.*

benthamiana. In the presence of Pwl1, HIPP43 PD localization was abolished, and the two

proteins co-localise in the cytoplasm and partially in cytoplasmic mobile structures (**Fig. S9B**)

Conversely, Pwl3 did not affect mCherry-HIPP43 localisation to PD (**Fig. S9C**), while Pwl4

(**Fig. S9D**) only showed cytoplasmic co-localisation without larger mobile bodies forming. In

addition, because we had observed that *Hv*HIPP43 orthologs from wheat (*Taes*HIPP43), foxtail

millet (*Siti*HIPP43) and weeping lovegrass (*Ec*HIPP43) interact with Pwl2 in Y2H assays, we

also tested whether *Taes*HIPP43, *Siti*HIPP43 or *Ec*HIPP43 localise as PD puncta in the same

way as *Hv*HIPP43. Interestingly, *Taes*HIPP43 (wheat) mostly localized to the nucleus,

cytoplasm and PD, even though PD localization was reduced (**Fig. S9E**), whereas *Ec*HIPP43

(weeping lovegrass) (**Fig. S9F**) and *Siti*HIPP43 (foxtail millet) (**Fig. S9G**) localized to small

puncta equivalent to PD, like *Hv*-HIPP43. Because Pwl2 is able to alter sub-cellular

localisation of *Hv*HIPP43, we tested whether this also occurred when *Siti*HIPP43 and Pwl2

were co-expressed. We observed Siti-HIPP43/Pwl2 co-localisation as cytoplasmic mobile

bodies away from PD, mirroring the Pwl2/HvHIPP43 interaction (**Fig. S9H-J**). When

considered together, we conclude that Pwl2 and its family members can interact with

*Hv*HIPP43 and that Pwl2 can also interact with HIPP43 orthologs from other cereals, thereby

altering their deployment to PD.

Pwl2 consistently alters the plasmodesmatal localization of HIPP43.

To understand the effect of Pwl2 expression on the PD localisation of *Hv*HIPP43, we carried

out a detailed quantitative analysis. In control experiments we expressed *Hv*HIPP43 with free

cytoplasmic YFP, and two MAX effectors AVR-PikE and MEP3 (Yan *et al.*, 2023). In six

independent co-expression experiments, we observed co-localisation of Pwl2 and *Hv*HIPP43

as mobile cytoplasmic puncta or large cytoplasmic mobile structures, while PD localisation

was significantly reduced (**Fig. 7I and J**). By contrast, co-localisation of *Hv*HIPP43 with AVR-

PikE or MEP3 led to mCherry-*Hv*HIPP43 fluorescence remaining at PDs (**Fig. 7K-N**).

Moreover, mCherry-HIPP43 fluorescence showed greater intensity when co-expressed with

Pwl2 compared to when co-expressed with AVR-PikE or MEP3 **(Fig. 7K-N)**. It is therefore

possible that Pwl2 is able to stabilise *Hv*HIPP43 or increases its accumulation, because the

mCherry-*Hv*HIPP43 signal is barely detectable in the absence of Pwl2 **(Fig. 7K and M)**. Our

previous structural analysis demonstrated that Pwl2 and *Os*HIPP43 produce a robust binding

interface that requires up to seven mutations in Pwl2 (*Pwl2*^SNDEYWY^) to abolish (Zdrzałek *et*

al., 2024). To test whether binding of Pwl2 to HIPP43 is necessary for the alteration in its

cellular localisation, we co-infiltrated mCherry-HIPP43 and YFP-Pwl2^SNDEYWY^ in *N.*

benthamiana and carried out live cell imaging. We were able to observe YFP-Pwl2^SNDEYWY^

expression, but this did not alter mCherry-*Hv*HIPP43 localisation at PD **(Fig. 7O-P)**.

Furthermore, the fluorescence intensity of mCherry-*Hv*HIPP43 did not increase as it did upon

co-expression with YFP-Pwl2 **(Fig. 7I and O)**. To rule out the possibility that Pwl2 degrades

*Hv*HIPP43 upon interaction in plant cells, we carried out immuno-blot analysis using protein

extracts from *N. benthamiana* following co-infiltration of mCherry-HIPP43 with either Pwl2,

MEP3 or AVR-PikE. Strikingly, the presence of Pwl2 led to pronounced mCherry-*Hv*HIPP43

accumulation compared to when co-infiltrated with MEP3 or AVR-PikE **(Fig. 7Q)**. We were

also concerned that Pwl2/HIPP43 co-localisation as puncta could be mis-interpreted as also

occurring at PD. To rule out such a possibility, we stained PD with aniline blue after co-

infiltrating *Hv*HIPP43 with free-YFP, YFP-Pwl2, AVR-PikE-YFP and MEP3-YFP. In a

control experiment we expressed free-YFP, YFP-Pwl2, AVR-PikE-YFP and MEP3-YFP in the

absence of *Hv*HIPP43. We found that free-YFP, Pwl2, AVR-PikE-YFP and MEP3 show

localisation to the cytosol **(Fig. S10A-D)**, but MEP3-YFP also showed some additional PD

localisation. By contrast YFP-Pwl2 was always observed as cytoplasmic mobile puncta **(Fig.**

S10B). When cells expressing Pwl2 were stained with aniline blue there was no co-localization

of Pwl2 and the aniline blue signal **(Fig. S10E-G)**. In the presence of mCherry-HIPP43 and

YFP-Pwl2, co-localisation at cytoplasmic puncta/mobile bodies was observed and did not co-

localise with aniline blue staining **(Fig. S10H).** In limited cases where co-localisation was

observed, this was transient because the Pwl2/HIPP43 puncta are mobile. By contrast, when

expressed on its own, or in the presence of free-YFP or two MAX-fold effectors AVR-PikE,

free-YFP and MEP3, mCherry-HIPP43 co-localised with aniline blue to PD **(Fig. S10I-K)**. To

investigate the motility of HIPP43-Pwl2 puncta, we carried out time-lapse imaging using

confocal laser microscopy of aniline blue-stained *N. benthamiana* cells following co-

infiltration of mCherry-HIPP43 with YFP-Pwl2, MEP3-YFP, AVR-PikE-YFP or YFP-

Pwl2^SNDEYWY^. Blue stained puncta equivalent to PD remained immobile while mCherry-

HIPP43/YFP-Pwl2 fluorescence was observed at mobile cytoplasmic structures (see **Movie S1**

and individual frames shown in **Fig. S11A)**. By contrast, co-localisation of mCherry-HIPP43

at aniline blue-stained PD was clearly visible and not altered by the presence two MAX-fold

effectors MEP3-YFP and AVR-PikE-YFP, YFP-Pwl2^{SNDEYWY} or of free-YFP (see **Movie S2-**

S5) and frames in **Fig S11B-E)**. When considered together, these results provide evidence that

Pwl2 interacts with HIPP43 altering its cellular location away from plasmodesmata.

The ability of Pwl2 to bind HIPP43 is necessary for its avirulence and virulence functions.

As the interaction between Pwl2 and HIPP43 appears to be critical for altering its sub-cellular

localisation to PD, we decided to test if this was also necessary for the role of Pwl2 in blast

disease. We therefore set out to see whether the *pwl2*^{SNDEYWY} allele could complement the

mutant phenotypes of a *M. oryzae Δpwl2* mutant. To do this, we transformed *Δpwl2* mutant

with a construct expressing Pwl2^{SNDEYWY} under its native promoter (*PWL2p:pwl2*^{SNDEYWY}) and

successful transformants were selected (**Fig. 8A and B**). We then quantified *Pwl2*^{SNDEYWY} gene

expression in selected transformants using qRT-PCR (**Fig. 8C**). We reasoned that if recognition

of Pwl2 by Mla3 requires HIPP43 recognition (Gómez De La Cruz *et al.*, 2024), then

Pwl2^{SNDEYWY} should not complement the observed gain of virulence of *Δpwl2* mutants on

barley cv. Baronesse (+*Mla3*). Moreover, if the Pwl2/HIPP43 interaction is required for the

virulence function of Pwl2, then Pwl2^{SNDEYWY} should not restore virulence to *Δpwl2* mutants.

In three independent experiments of two independent strains, C6 (*Δpwl2+PWL2p:pwl2*^{SNDEYWY})

and C14 (*Δpwl2+PWL2p:pwl2*^{SNDEYWY}) showed a compatible interaction on cv. Baronesse (**Fig.**

8D) providing evidence that the Pwl2^{SNDEYWY} is not recognised by Mla3, because the strains

retained the ability to infect a susceptible barley cv. Nigrate (-*Mla3*) (**Fig. 8E**). We also

observed that *Δpwl2 +PWL2p:Pwl2*^{SNDEYWY} transformants C6 and C14 did not restore full

virulence on susceptible rice cv. CO39 compared to the wildtype Guy11 (**Fig. 8F**), based on

lesion density (**Fig. 8G)** and lesion length (**Fig. 8H**). We conclude that the interaction between

Pwl2 and HIPP43 is required for both its recognition as an avirulence factor and its function as

a virulence determinant.

Discussion

The Pwl2 effector was first identified as a host specificity determinant for infection of the forage grass species weeping lovegrass (Kang *et al.*, 1995; Sweigard *et al.*, 1995). Its initial identification provided evidence that host range in plant pathogenic fungi was conditioned in a similar way to cultivar specificity in a gene-for-gene manner, involving dominant pathogen genes recognised by the products of cognate disease resistance genes (Kang *et al.*, 1995). Furthermore, Pwl2 was found to belong to an expanded gene family, suggesting an important function in pathogenicity and fitness (Kang *et al.*, 1995). However, in the following two decades, the function of Pwl2 remained elusive despite its extensive use as a marker in cell biological studies for investigating effector regulation, secretion and delivery during plant infection (Kankanala *et al.*, 2007; Giraldo *et al.*, 2013; Oliveira-Garcia *et al.*, 2023b).

In this study, we set out to explore the function of Pwl2 and investigate why it is such a highly conserved effector in *M. oryzae*. We found that *PWL2* is highly conserved in *M. oryzae*, including each of its host-limited forms and even in sister *Magnaporthe* species infecting crabgrass and pearl millet. The observation that *PWL2* and the wider *PWL* family genes have been maintained in the global blast population and there are effector variants in certain host-adapted lineages, suggests that Pwl2 and members of this family serve an important function in pathogenesis. Furthermore, *PWL2* is present at high copy number in many *M. oryzae* isolates having undergone extensive gene duplication and is specifically expressed during the initial stages of blast infection, particularly at the stage when *M. oryzae* moves from an initially colonised epidermal cell, following appressorium penetration, to adjacent host cells. This cell-to-cell movement by the fungus utilises PD-containing pit fields and the fungus forms a transpressorium structure that undergoes severe hyphal constriction to traverse each pit field (Kankanala *et al.*, 2007; Cruz-Mireles *et al.*, 2021), a process that requires activity of the Pmk1 MAP kinase (Sakulkoo et al., 2018). We found that *PWL2* is expressed during this process in a Pmk1-dependent manner, forming part of a regulated set of effectors deployed by the fungus during cell-to-cell movement. The generation of a *Δpwl2* mutant, made possible by CRISPR/Cas9 gene editing to delete all three native copies of the gene, enabled us to confirm the role of Pwl2 in host specificity to weeping lovegrass (Sweigard et al., 1995), as an avirulence effector for barley Mla3 (Brabham et al., 2024), and also revealed its importance in blast disease. Pwl2 is therefore an important virulence determinant for blast disease, explaining its conservation and amplification.

To identify the likely target of Pwl2, we utilised discovery proteomics which revealed its
interaction with an isoprenylated small HMA protein, HIPP43. This is consistent with Pwl2
being a MAX effector – many of which have been shown to interact with sHMA protein
domains – although previous studies have focused on incorporation of sHMA domains into
paired NLR immune receptors leading to disease resistance (Ortiz *et al.*, 2017; Bentham *et al.*,
2021; Maidment *et al.*, 2021; Mukhi *et al.*, 2021). The crystal structure of the Pwl2/HIPP43
complex (Zdrzałek *et al.*, 2024) demonstrates that Pwl2 uses an expansive interface to mediate
binding to HIPP43, largely using elements of the MAX fold to interact with the β-sheet of its
host target. Indeed, when HIPP43 is incorporated into the Pik-1 NLR, in place of its naturally
occurring integrated HMA domain, this leads to an immune response to Pwl2 (Zdrzałek *et al.*,
2024). A recent study has also revealed that the barley resistance protein Mla3 acquired the
ability to bind Pwl2 by mimicking the HMA fold of its host target, HIPP43 (Gómez De La
Cruz *et al.*, 2024). Interestingly, polymorphic residues found in Pwl2 variants, such as pwl2-2
and pwl2-3 are not located at the HIPP43 binding interface but are located away from the MAX
fold in the C-terminal helix of Pwl2. It is possible that these polymorphic residues are essential
to stabilize the Pwl2, MAX-fold/HMA-like interface in Mla3. Consistent with this idea,
introducing mutations that disrupt the Pwl2/HIPP43 interaction also results in loss of
recognition by Mla3.
468
In spite of the importance of sHMA domains in plant immunity, little is known regarding their
actual function or why they are targeted by fungal effectors. HMA domains are known to be
involved in biotic/abiotic stress responses, transport of metals and metal detoxification, and
their expression can be organ-specific or tissue-specific within roots, leaves, or stems (Barth *et*
al., 2009; Zschiesche *et al.*, 2015; Zhang *et al.*, 2020). Moreover, sHMAs can localize to the
nucleus, plasma membrane, cytoplasm, or to plasmodesmata (Barth *et al.*, 2009; Cowan *et al.*,
2018; Barr *et al.*, 2023; Oikawa *et al.*, 2024). Furthermore, sHMAs are expanded in plant
genomes with more than 45, and more than 50, sHMA domain-encoding genes occurring in
Arabidopsis and rice, respectively (de Abreu-Neto *et al.*, 2013). Unique intragenic deletions in
OsHIPP05 (*Pi21*)*,* a proline rich HMA domain protein-encoding gene in rice leads to rice blast
resistance (Fukuoka *et al.*, 2009) and gene silencing or deletion mutants of *TaHIPP1* or
AtHMAD1 in wheat and *Arabidopsis* provide enhanced resistance against *Puccinia striiformis*
f. sp*. tritici* and *Pseudomonas syringae* DC3000, respectively (Imran *et al.*, 2016; Wang *et al.*,
2023). In addition, deletion mutants of *AtHIPP27* in *Arabidopsis* lead to increased resistance
against cyst nematodes (Radakovic *et al.*, 2018). However, it is not clear why deletion of sHMA

protein-encoding genes impacts immunity. Therefore, the observation that Pwl2 interacts with

HIPP43 is revealing, especially because over-expression of either Pwl2 or *Hv*HIPP43

suppresses PTI responses and enhances blast disease susceptibility. How this potential

enhancement of HIPP43 activity is stimulated by Pwl2 is not completely clear, but transient

co-expression of both Pwl2 and HIPP43 sequesters HIPP43 away from plasmodesmata (PD),

and the two proteins instead co-localize in large mobile structures within the cytoplasm. The

localisation of HIPP43 to PD requires its isoprenylation motif and may be associated with an

immune signaling role at this site. Given the Pmk1 MAP kinase-dependent expression of Pwl2

during PD traversal, it may be that sequestering HIPP43 away from these sites is critical for

the fungus to invade plant tissue efficiently. This is consistent with the reduced virulence

phenotype of *Δpwl2* mutants, which results in slower generation of disease symptoms and

reduced lesion size. A recent study has highlighted how a *P. infestans* effector PiE354 can

interfere with a host immune response by re-routing a plant Rab8a away from the

plant/pathogen interface, a similar potential effector function that alters the cellular location of

a target rather than impairing a particular function (Yuen *et al.*, 2024).

Localization of two HIPP proteins, HIPP7 and HIPP26, to PD has been previously reported,

which is also dependent on isoprenylation motifs like HIPP43 (Cowan *et al.*, 2018; Guo *et al.*,

2021). Although characterized HIPPs have been proposed to be metallochaperones, there are

no studies that link metal detoxification to immunity. Some studies suggest that there may be

a connection between concentration of metal ions, such as iron, copper, cadmium and zinc,

with plasmodesmatal permeability (O'Lexy *et al.*, 2018), which might explain the function of

a plasmodesmatal localized HIPP in regulating permeability, associated with their role in

immunity. Both the HMA domain and isoprenylation regions of HIPPs have been shown to be

important for plant immunity. For example, a mutation targeting the proline-rich region of *pi21*

is sufficient to lead to gain of resistance against *M. oryzae* (Fukuoka *et al.*, 2009), while

interfering with the isoprenylation of HIPP1-V from wild wheat (*H. villosa*) leads to loss of

resistance against *Blumeria graminis* f. sp. *tritici* accompanied by reduced HIPP1-V

localization to plasma membrane. Interestingly, HIPP1-V interacts with the E3-ligase CMPG1-

V at the plasma membrane leading to resistance to powdery mildew in an isoprenylation-

dependent manner (Wang *et al.*, 2023). This interaction has been reported to activate

expression of genes involved in ROS generation and salicylic acid biosynthesis, suggesting

that HIPP1-V is required for PTI regulation (Wang *et al.*, 2023). Interestingly, we found that

HIPP1-V is an ortholog of HvHIPP43 and can interact with Pwl2 in a Y2H assay. It is possible,

therefore, that the Pwl2 interaction with HvHIPP43 attenuates PTI through a similar

mechanism, which will require further investigation.

How the change in cellular localization of HIPP43 induced by the Pwl2 effector prevents its

function in immunity or leads to a new role that enhances disease susceptibility remains

unclear. A very recent study has provided evidence that *M. oryzae* AVR-Pik binding stabilizes

the rice small HMA (sHMA) proteins OsHIPP19 and OsHIPP20 (Oikawa *et al.*, 2024),

suggesting that the function of Pwl2 may be mirrored by other blast effectors, targeting a wider

pool of HIPPs. In this regard, future work will be necessary to determine whether HIPP43

directly regulates ROS generation, which might reduce permeability of plasmodesmata (Cui

and Lee, 2016), or acts in a more indirect manner through interaction with other signaling

components involved in PTI. Finally, Pwl2 is known to be a highly mobile effector and has

been shown to move into neighbouring rice cells ahead of *M. oryzae* hyphal growth (Giraldo

et al., 2013). This has been suggested to be a step to prepare un-invaded cells for fungal

colonisation, consistent with its Pmk1-dependent regulation. Pwl2 may therefore alter the sub-

cellular localisation and concentration of HIPP43, sequestering it away from PD that the fungus

uses for effector movement and hyphal invasion, thereby enabling more rapid tissue

colonisation by the blast fungus.

Materials and methods

Fungal strains, growth conditions and infection assays

Fungal isolates were routinely grown on complete medium (CM) at 24°C with a controlled 12 h light and dark cycle for up to 12 days (Talbot *et al.*, 1993). *H. vulgare*, *E. curvula* and *O. sativa* plants were grown for 7, 14 and 21 days respectively in 9 cm diameter plastic plant pots or seed trays. Conidia were recovered from 10-day old cultures using a sterile disposable plastic spreader in 3 mL sterile distilled water. The conidial suspension was filtered through sterile Miracloth and centrifuged at 5000 x g for 5 min at room temperature before adjusting to a final of concentration of 1×10^5 conidia mL^{-1} in 0.2 % gelatin. The spore suspension was used for spray or leaf drop infections assays. After spray inoculation, plants were placed in polythene bags and incubated in a controlled plant growth chamber at 24°C for 48 h with a 12 h light-dark cycle and 85% relative humidity, before removing polythene bags. Inoculated plants were incubated for 3 days before scoring lesions. For each treatment 10 leaves were collected before counting typical ellipsoid necrotic disease lesions with a grey centre (Valent *et al.*, 1991). Each experiment was repeated a minimum of three times, yielding consistent outcomes.

Leaf infection assay and live-cell imaging

Rice leaf sheath from 4 week old susceptible cultivars Moukoto or CO39 were inoculated with 4 mL of a suspension at 5×10^4 conidia mL^{-1} in ddH2O using a micropipette (Kankanala *et al.*, 2007). Inoculated leaf sheaths were incubated at 24°C for 24 h before a thin layer of inner leaf sheath was dissected and mounted on a glass slide. Treatment with INA-PP1 was carried as described previously (Sakulkoo *et al.*, 2018). Live cell imaging was carried out on an IX81 motorized inverted microscope (Olympus, Hamburg, Germany) for conventional and differential interference contrast (DIC) microscopy using Photometrics CoolSNAP HQ2 camera (Roper Scientific, Germany). Images were analyzed using ImageJ. For Leica SP8 laser confocal microscopy, settings were as follows; GFP, YFP and RFP/mCherry tagged proteins were excited using 488, 514 and 561 nm laser diodes and emitted fluorescence detected using 495-550, 525-565 and 570-620 nm respectively. Auto-fluorescence from chlorophyll was detected at 650-740 nm.

Generation of fungal transformation plasmids

Single or multiple DNA fragments were cloned into fungal transformation vectors using In-Fusion HD Cloning (Clontech, USA). Briefly, fragments from cDNA, genomic DNA or synthesized DNA were amplified using primers to introduce a 15bp overhang complementary

to sequences at restriction sites of a destination vector or adjacent insert fragments. This allows
the ends to fuse by homologous recombination. Positive transformants were selected by colony
PCR and constructs sequenced by GENEWIZ. A list of primers is provided in Table S3.

RNA isolation, RNA sequencing and analysis
To study *in-planta* gene expression of *PWL2* and other effectors, leaf drop infection assays
were carried out using susceptible rice cv. Moukoto and samples collected at 24 and 72 hpi.
Infected plant material was ground to a fine powder using a sterile nuclease-free mortar &
pestle containing liquid N_2. RNA was isolated from *M. oryzae* mycelium, or inoculated rice
leaves using QIAGEN RNeasy Plant Mini Kit. RNA quality was determined NanoDrop
spectrophotometry (Thermo Scientific, UK) and Agilent 2100 Bioanalyser (Agilent
Technologies, UK). Library preparation was carried out using Illumina® sequencing TruSeq
Stranded Total RNA Library Prep Kit before sequencing 100 bp paired ends reads using
Illumina Genome Analyser GXII platform by Exeter Sequencing Service (University of
Exeter). To determine differential gene expression, raw reads were separated by mapping to
both *M. oryzae* and *O. sativa* using kraken2. Reads specific to *M. oryzae* were used to quantify
transcript abundance using Kallisto. To quantify genes missing in 70-15, separated reads were
mapped to KE002. R package Sleuth was used to determine genes showing differential
expression with log2fold > 1 and P-adjust value < 0.05 defined as up-regulated and a log2fold
> 1 and P-adjust value < 0.05 as down-regulated. Southern blot analysis of *M. oryzae* genomic
DNA was carried out as described previously (Talbot *et al.*, 1993). Quantitative real-time PCR
was carried out using CFX OPUS 96 and CT values normalised to a house keeping gene, β-
Tubulin (MGG_00604). Fold change was determined using the formula $2^{-\Delta\Delta CT}$, where $\Delta\Delta Ct =$
((CtGOI in infected sample - CtNC in infected sample) – (CtNC in mycelia - CtNC in
mycelia)), and GOI is the gene of interest and NC (negative control) is β-Tubulin.

Generation of Cas9-sgRNA targeted gene deletion

A sgRNA was designed for CRISPR-Cas9 genome editing using online tool E-CRISP
http://www.e-crisp.org/E-CRISP/. A 20-nucleotide sequence was selected at the *PWL2* locus
(not including the PAM NGG-sequence). The sgRNA was first synthesised using the EnGen
sgRNA synthesis kit New England Biolabs (NEB #E3322) before mixing with Cas9-NLS to
form an RNP complex (Foster *et al.*, 2018). The mixture was incubated at room temperature
for 10 min before fungal transformation. *M. oryzae* protoplasts were generated as described
previously (Talbot *et al.*, 1993). The RNP complex together with donor template was mixed

with Guy11 protoplasts re-suspended in 150 µL STC to a concentration of 1×10^{8} mL^{-1}

incubated at room temperature for 25 min before adding 60% PEG. Successful transformants

were selected on complete medium (CM) agar containing 200 µg mL^{-1}Hygromycin B .

Whole genome sequencing

Purified RNA-free DNA was obtained using Hexadecyltrimethylammonium Bromide

(CTAB). Template quality was assessed by NanoDrop and Qubit spectrophotometry.

Sequencing was carried out at Exeter Sequencing services (University of Exeter, UK) and

Novogene (Cambridge, UK). NEXTflex^{TM} Rapid DNA-seq Library Prep Kit was used to

prepare and index libraries before sequencing on HiSeq 2500 (Illumina) with two lanes per

sample. Quality of sequencing reads was checked using FastQC

(http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/). From raw data (fastq files), adaptor

sequences were trimmed from sequences containing adaptors and low-quality reads removed

by fastq-mcf. Trimmed sequences were aligned to the reference genome (70-15) (Dean *et al.*,

2005) using BWA (Burrow Wheeler Aligner) https://github.com/lh3/bwa (Li and Durbin,

2009). Bam files were visualized by IGV genome viewer to determine CRISPR gene deletion.

Copy number variation of *PWL2*

A total of 286 *M. oryzae* isolates with raw Illumina-based sequencing information were

downloaded from NCBI (performed on October 16[th], 2019). Copy number variation was

assessed using a k-mer analysis approach using the k-mer analysis toolkit (KAT; v2.4.1).

Coding sequence information for *M. oryzae* isolate 70-15 was used as template for k-mer

analysis and raw Illumina reads were input. Default parameters were used including k-mer

length 27 nt. Copy number variation of individual effectors is based on average k-mer coverage

compared with the median coverage for all genes.

Phylogenetic analysis of *Magnaporthe* isolates

Genome sequences of diverse *M. oryzae* isolates were identified by literature review and

searches on NCBI (performed August 9[th], 2019). For isolates having only Illumina sequencing

data, raw paired reads were downloaded from NCBI, trimmed using Trimmomatic (v0.36)

using the parameters: removal of adapters with ILLUMINACLIP:TruSeq2-PE.fa:2:30:10,

remove leading low quality or N based with quality below 5 for leading (LEADING:5) and

trailing sequence (TRAILING:5), scan and cut reads with 4 bp sliding window below 10

(SLIDINGWINDOW:4:10), and minimum length 36 bp (MINLEN:36). KmerGenie (v1.7048)

was used to identify an optimal k-mer for genome assembly using default parameters. Genome

assembly was performed using minia (v0.0.102) with default parameters. kSNP3 (v.3.021) was

used to develop a phylogenetic tree of *M. oryzae* using assembled genomes as input with

parameters of k-mer of 29 bp and minimum fraction of 0.4. The phylogenetic tree was

generated using RAxML (v8.2.12) using the General Time Reversible model of nucleotide

substitution under the Gamma model of rate heterogeneity.

Generation of *Agrobacterium* transformation plasmids

Single or multiple DNA fragments were cloned into binary vectors using In-Fusion HD

Cloning (Clontech, USA). Briefly, fragments from cDNA were amplified to introduce a 15bp

overhang complementary to sequences at restriction sites of a destination vector or adjacent

insert fragments. Alternatively, synthesized DNA fragments were designed with a 15bp

overhang complementary to sequences at restriction sites of binary vector pG514 customized

for infusion cloning by digestion by either *Xba*I and *Pca*I or *Sac*I (New England Biolabs) to

remove the *ccDB* toxin encoding gene. Positive transformants were accessed by colony PCR

and constructs sequenced and analyzed on SNAP gene.

Transient expression in *Nicotiana benthamiana*

Agrobacterium tumefaciens strain *GV3101* (Holsters *et al.*, 1980) was used for transient

expression. Three-week-old *N. benthamiana* leaves were infiltrated with transformed

Agrobacterium carrying T-DNA constructs expressing the gene of interest. Bacterial cultures

were diluted to obtain a final OD_{600} of 0.4 in agroinfilitration buffer (10mM MES, 10 mM

$MgCl_2$, 150 µM acetosyringone, pH 5.6). Leaf discs were cut from agroinfiltrated tissue 48 hpi

and subjected for microscopy or used for co-IP.

Plant transformation and ROS measurement

Agrobacaterium tumefaciens strain *AGL1* (Holsters *et al.*, 1980) was used for plant

transformation (Hensel *et al.*, 2009). Positive transgenic plants were selected on HygromycinB

followed by confirmation using PCR. Alternatively, leaf discs were collected from transgenic

plants and analysed for copy number by iDNA Genetics Ltd (Norwich, UK). In addition,

expression of the protein was evaluated using SDS-PAGE. To measure the response to PTI

elicitors, a 4-mm diameter biopsy punch (Integra[TM] Miltex [TM]) was used to cut leaf discs from

5-week-old *H. vulgare* transgenic plants. Leaf discs were transferred to 96-well-plates (Greiner

Bio-One) containing 100 µL ddH_2O in each well and incubated overnight at room temperature.

The assay was carried out by replacing ddH_2O in wells by a 100 µL of 100 µM Luminol or L-

012 (Merck), 20 µg mL^{-1} horseradish peroxidase (Merck) together with elicitors, flg22 (1 µM

final concentration), chitin (1mg/mL final concentration) or without (mock). Photon count was

carried out using a HRPCS218 (Photek) equipped with a 20 mm F1.8 EX DG ASPHERICAL

RF WIDE LENS (Sigma Corp). Each experiment was repeated three times, yielding consistent

outcomes.

Co-immunoprecipitation and sample preparation for mass spectrometry

Leaves were harvested from 14-21 day old barley transgenic plants and rapidly frozen in liquid

N_2 before storage or immediately ground to fine powder using a GenoGrinder® tissue

homogenizer. Ground powder was quickly transferred to ice-cold 1.5 mL Eppendorf tubes and

2 mL of ice-cold extraction buffer (GTEN [10% (v/v) glycerol, 25 Mm Tris pH 7.5, 1 mM

EDTA 150 mM NaCl, 2% (w/v) PVPP, 10 Mm DTT, 1 X protease inhibitor cocktail (Sigma),

0.5% (v/v) IGEPAL added and mixed thoroughly. This was centrifuged at 3000 x g for 10 min

at 4°C to recover total protein in the supernatant and repeated twice. A volume of 2 mL of total

protein was mixed with 25 μL of GFP-TRAP agarose beads (50% slurry, ChromoTek) and

incubated shaking overnight at 4°C. The GFP-TRAP agarose beads were centrifuged for 2 min

at 3000 x g before washing three times with wash buffer (ChromoTek). Bound proteins were

recovered by resuspending beads in loading buffer (loading dye, 10 mM DTT, H2O) before

incubating at 70°C for 10 min. An aliquot was run by SDS-PAGE and used for immuno-blot

analysis (Burnette, 1981). Recovered proteins were fractionated by SDS-PAGE for

approximately 1 cm. The gel was washed for 1 h in ddH$_2$O followed by 1 incubation

SimplyBlue^{TM} Safe stain. The gel was then washed three times in ddH$_2$O and the region

containing proteins excised using a scalpel.

Mass spectrometry and data processing

Protein purification, immunoprecipitation, sample preparation, liquid chromatography

followed by tandem mass spectrometry (LC-MS/MS), and data analysis were carried out as

previously described (Li *et al.*, 2023). Proteins identified in immunoaffinity-enriched samples

were measured by high resolution LC-MS systems, Orbitrap Fusion (Thermo Ltd). Acquired

spectra were peak-picked and searched by Mascot (Matrix Science Ltd) to identify peptide

sequences from the search space defined by background proteome. Peptides were combined

into proteins based on the principle of parsimony by the search engine. Resulting proteins were

further described by quantitative values based on the number of spectra that identify them.

Individual runs were combined in Scaffold (Proteome Software Inc.), where the data were

evaluated and filtered to contain less than 1% false positives (FDR) and the resulting matrix

exported. The matrix of proteins detected in different samples serves as the input for an R script
for further processing and visualization.

Yeast two-hybrid analysis
To clone genes of interest into bait (pGBKT7 DNA-BD cloning) vector or prey (pGADT7
activating domain) vector, by In-Fusion HD Cloning (Clontech), bait and prey vectors were
digested by *Bam*H1 and *Eco*R1 while gene to be inserted was PCR-amplified using primers
containing 15 bp overhangs with homology to two ends of the digested bait vector. For
transformation, a single colony Y2H gold yeast strain was mixed in 1 mL of liquid yeast extract
peptone dextrose (YPD). Competent cells were prepared according to manufacturer
instructions (Zymo Research). Briefly, for transformation 700 ng – 1 µg of plasmids expressing
Pwl2 or other effectors in pGADT7 and *Hv*HIPP43 or plant proteins in pGBKT7 were co-
transformed in competent cells and Frozen-EZ Yeast Solution 3 added before incubation at
28ºC for 1 h and transformed cells plated on selection media lacking Leucine (L) and
Tryptophan (W) and incubated at 28º C for 3-5 days. To detect interactions colonies were
transferred to media lacking Leucine (L), Tryptophan (W), Adenine (A) Histidine (H), Bromo-
4-Chloro-3-Indolyl a-D-galactopyranoside (X-a-gal) and 10 mM 3-amino-1,2,4-triazole (3AT)
(Sigma). The plates were imaged after 60-72 incubation at 28ºC. Each experiment was
conducted a minimum of three times.

In-planta **co-immunoprecipitation (co-IP)**
To test interactions of proteins *in planta*, genes of interest tagged with either GFP or Myc were
cloned into vector pGW514, transformed into *Agrobacterium* strain *GV3101* and co-infiltrated
(OD_{600} = 0.4 for effectors and OD_{600} = 0.6 for HIPP43) into 3-4 week-old *N. benthamiana*
leaves and incubated for 48 h to allow protein expression. Total protein was isolated in ice-
cold extraction buffer (GTEN [10% (v/v) glycerol, 25 Mm Tris pH 7.5, 1 mM EDTA 150 mM
NaCl], 2% (w/v) PVPP, 10 Mm DTT, 1 X protease inhibitor cocktail (Sigma), 0.5% (v/v)
IGEPAL). Total protein was co-immunoprecipitated using anti-GFP M2 resin (Sigma-Aldrich,
St. Louis, MO) and washed 3 times using immunoprecipitation buffer before analyzing by
SDS-PAGE. Recovered proteins from co-immunoprecipitation were separated by SDS-PAGE
and transferred to a polyvinylidene diflouride (PVDF) membrane using a Trans-Blot turbo
transfer system (Bio841 Rad, Germany). Detection was performed using the appropriate
antibody either anti-GFP-HRP or anti-Myc-HRP. Imaging was carried out using an

ImageQuant LAS 4000 luminescent imager (GE Healthcare 844 Life Sciences, Piscataway,

NJ, U.S.A.) according to manufacturer's instructions.

Sub-cellular localization and plasmodesmata quantification

Leaf discs were obtained from *N. benthamiana* plants agroinfiltrated with different construct

combinations at 24 and 48 hpi and mounted on a slide immersed in perfluorodecalin and

observed using X63 oil immersion lens. For Leica SP8 laser confocal microscopy, settings

were as follows; GFP, YFP and RFP/mCherry tagged proteins were excited using 488, 514,

and 561 nm laser diodes and emitted fluorescence detected using 495-550, 525-565, and 570-

620 nm respectively. Auto-fluorescence from chlorophyll was detected at 650-740 nm. To stain

and quantify plasmodesmata (PD), agroinfiltrated leaves were further infiltrated with 0.1%

aniline (Sigma-Aldrich #415049, in PBS buffer, pH 7). Images were analyzed using ImageJ to

determine the number of PD. First, cell peripheries were divided into 20 µm sections and the

number of PD determined per 20 µm.

Phylogenetic analysis of grass HMA proteins

Proteins from *B. distachyon* (314; v3.1), *H. vulgare* (Barley cv. Morex V3, Jul 2020), *T.*

aestivum (RefSeqv2.1; 09-16-2020), *O. sativa* (323; v7.0), *Oropetium thomaeum* (386; v1.0),

Sorghum bicolor (454; v3.1.1), *Setaria italica* (312; v2.2), and *Zea mays* (RefGen_V4)

containing an HMA domain were identified using InterProScan (v5.59-91.0; Pfam PF00403).

The   HMA   domain   was   extracted   using   the   script   QKdomain_process.py

(https://github.com/matthewmoscou/QKdomain) including an additional 10 amino acid

sequences N and C-terminal of the Pfam boundaries (-n 10 -c 10). The non-redundant set of

HMA domains were identified using CD-HIT (v.4.8.1) with parameter -c 1.0. Structure-guided

multiple sequence alignment was performed using MAFFT with parameters dash, max iteration

of 1000, and globalpair. Maximum likelihood phylogenetic analysis was performed using

RAxML (v8.2.12) using the Gamma model of rate heterogeneity, JTT amino acid substitution

model,- and 1,000 bootstraps. Coding sequence was identified for the HIPP43 gene family and

aligned using MUSCLE (v5) translation alignment using default parameters. Maximum

likelihood phylogenetic analysis was performed using RAxML (v8.2.12) using General Time

Reversible model of nucleotide substitution under the Gamma model of rate heterogeneity and

1,000 bootstraps.

Statistical analysis and Protein Structure Prediction

Significance difference between groups of samples was performed using GraphPad Prism 10.

P values < 0.05 were considered significant, ****$P<0.0001$, ***$P<0.001$, **$P<0.01$ while

values *P* values > 0.05 were considered as non-significant in an unpaired Student's t-test.

Structure prediction was carried out using AlphaFold3 (Abramson *et al.*, 2024), the structure

analyzed and figures generated using ChimeraX (Meng *et al.*, 2023).

Acknowledgements

We thank current and past members of the "BLASTOFF" team from the Banfield, Kamoun,

Terauchi, Talbot, and Moscou Laboratories. For funding, we thank Gatsby Charitable

Foundation and Biotechnology and Biological Sciences Research Council (BBSRC)

BBS/E/J/000PR9797 and a BBSRC grant to NJT and FLHM (BB/V016342/1), United States

Department of Agriculture-Agricultural Research Service CRIS #5062-21220-025-000D

(MJM) and BB/X010996/1, funding for APH ISP. We acknowledge Y.K Gupta, J.C De la

Concepcion, J. Rhodes and J. Win for constructive discussions.

Author Contributions

Conceptualization: NJT and VW

Formal analysis: VW, XY, AF, JK, JS, DG, TL, YPH, ABE, MS, MJB, MM, WM and RZ

Funding acquisition: NJT

Investigation VW, XY, AF, JS, DG, AG, NS, DK, YPH, ABE, MJB, WM and RZ

Writing – original draft: VW and NJT

Writing – review & editing: VW, NJT, LR, AB, TL, MB, SK, MM, FM

ORCID IDs

Vincent Were: 0000-0002-9885-4877

Xia Yan: 0000-0003-4509-0137

Andrew J. Foster: 0000-0002-8459-7590

Jan Sklenar: 0000-0003-1858-2574

Thorsten Langner: 0000-0002-3401-8888

Amber Gentle: 0009-0006-1944-4754

Neha Sahu   0009-0008-0721-6337

Adam R. Bentham: 0000-0001-5906-0962

Rafał Zdrzałek: 0000-0003-3669-924X

Lauren Ryder: 0000-0003-0370-5746

Davies Kaimenyi: 0000-0002-6320-3740

Diana Gómez De La Cruz: 0000-0001-6683-9245

Yohann Petit-Houdenot: 0000-0003-4806-8349

Alice Bisola Eseola: 0000-0001-5698-8204

Mark Jave Bautista: 0000-0002-6016-5953

Weibin Ma: 0000-0002-0516-2151

Jiorgos Kourelis: 0000-0002-9007-1333

Dan MacLean: 0000-0003-1032-0887

Mark J. Banfield: 0000-0001-8921-3835

Sophien Kamoun: 0000-0002-0290-0315

Frank L.H.Menke: 0000-0003-2490-4824

Matthew J. Moscou: 0000-0003-2098-6818

Nicholas J Talbot: 0000-0001-6434-7757

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung,
C.C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A.,
Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve,
M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain,
R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S.,
Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski,
M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., and Jumper, J.M.
(2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630,** 493-500.

Ashikawa, I., Hayashi, N., Yamane, H., Kanamori, H., Wu, J., Matsumoto, T., Ono, K., and Yano, M. (2008). Two Adjacent Nucleotide-Binding Site–Leucine-Rich Repeat Class Genes Are Required to Confer Pikm-Specific Rice Blast Resistance. Genetics **180,** 2267-2276.

Asuke, S., Tanaka, M., Hyon, G.S., Inoue, Y., Vy, T.T.P., Niwamoto, D., Nakayashiki, H., and Tosa, Y. (2020). Evolution of an Eleusine-Specific Subgroup of Pyricularia oryzae Through a Gain of an Avirulence Gene. Mol Plant Microbe Interact **33,** 153-165.

Barr, Z.K., Werner, T., and Tilsner, J. (2023). Heavy Metal-Associated Isoprenylated Plant Proteins (HIPPs) at Plasmodesmata: Exploring the Link between Localization and Function. In Plants.

Barth, O., Vogt, S., Uhlemann, R., Zschiesche, W., and Humbeck, K. (2009). Stress induced and nuclear localized HIPP26 from Arabidopsis thaliana interacts via its heavy metal associated domain with the drought stress related zinc finger transcription factor ATHB29. Plant Molecular Biology **69,** 213-226.

Bentham, A.R., Petit-Houdenot, Y., Win, J., Chuma, I., Terauchi, R., Banfield, M.J., Kamoun, S., and Langner, T. (2021). A single amino acid polymorphism in a conserved effector of the multihost blast fungus pathogen expands host-target binding spectrum. PLoS Pathog **17,** e1009957.

Brabham, H.J., Gómez De La Cruz, D., Were, V., Shimizu, M., Saitoh, H., Hernández-Pinzón, I., Green, P., Lorang, J., Fujisaki, K., Sato, K., Molnár, I., Šimková, H., Doležel, J., Russell, J., Taylor, J., Smoker, M., Gupta, Y.K., Wolpert, T., Talbot, N.J., Terauchi, R., and Moscou, M.J. (2023). Barley MLA3 recognizes the host-specificity effector Pwl2 from Magnaporthe oryzae. The Plant Cell, koad266.

Burnette, W.N. (1981). Burnette WN. "Western Blotting": electrophoretic transfer of proteins from sodium dodecyl sulfate-polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein. Analytical Biochemistry **112,** 195 - 203.

Cesari, S., Thilliez, G., Ribot, C., Chalvon, V., Michel, C., Jauneau, A., Rivas, S., Alaux, L., Kanzaki, H., Okuyama, Y., Morel, J.-B., Fournier, E., Tharreau, D., Terauchi, R., and Kroj, T. (2013). The Rice Resistance Protein Pair RGA4/RGA5 Recognizes the Magnaporthe oryzae Effectors AVR-Pia and AVR1-CO39 by Direct Binding The Plant Cell **25,** 1463-1481.

Cowan, G.H., Roberts, A.G., Jones, S., Kumar, P., Kalyandurg, P.B., Gil, J.F., Savenkov, E.I., Hemsley, P.A., and Torrance, L. (2018). Potato Mop-Top Virus Co-Opts the Stress Sensor HIPP26 for Long-Distance Movement. Plant Physiol **176,** 2052-2070.

Cruz, C.D., and Valent, B. (2017). Wheat blast disease: danger on the move. Trop Plant Pathol **42,** 210-222.

Cruz-Mireles, N., Eseola, A.B., Osés-Ruiz, M., Ryder, L.S., and Talbot, N.J. (2021). From appressorium to transpressorium—Defining the morphogenetic basis of host cell invasion by the rice blast fungus. PLOS Pathogens **17,** e1009779.

Cui, W., and Lee, J.-Y. (2016). Arabidopsis callose synthases CalS1/8 regulate plasmodesmal
permeability during stress. Nature Plants **2,** 16034.

**de Abreu-Neto, J.B., Turchetto-Zolet, A.C., de Oliveira, L.F.V., Bodanese Zanettini,
M.H., and Margis-Pinheiro, M.** (2013). Heavy metal-associated isoprenylated plant
protein (HIPP): characterization of a family of proteins exclusive to plants. The FEBS
Journal **280,** 1604-1616.

de Guillen, K., Ortiz-Vallejo, D., Gracy, J., Fournier, E., Kroj, T., and Padilla, A. (2015).
Structure Analysis Uncovers a Highly Diverse but Structurally Conserved Effector
Family in Phytopathogenic Fungi. PLoS Pathog **11,** e1005228.

**Dean, R.A., Talbot, N.J., Ebbole, D.J., Farman, M.L., Mitchell, T.K., Orbach, M.J., Thon,
M., Kulkarni, R., Xu, J.R., Pan, H., Read, N.D., Lee, Y.H., Carbone, I., Brown, D.,
Oh, Y.Y., Donofrio, N., Jeong, J.S., Soanes, D.M., Djonovic, S., Kolomiets, E.,
Rehmeyer, C., Li, W., Harding, M., Kim, S., Lebrun, M.H., Bohnert, H.,
Coughlan, S., Butler, J., Calvo, S., Ma, L.J., Nicol, R., Purcell, S., Nusbaum, C.,
Galagan, J.E., and Birren, B.W.** (2005). The genome sequence of the rice blast fungus
Magnaporthe grisea. Nature **434,** 980-986.

Dykema, P.E., Sipes, P.R., Marie, A., Biermann, B.J., Crowell, D.N., and Randall, S.K.
(1999). A new class of proteins capable of binding transition metals. Plant Molecular
Biology **41,** 139-150.

Foster, A.J., Martin-Urdiroz, M., Yan, X., Wright, H.S., Soanes, D.M., and Talbot, N.J.
(2018). CRISPR-Cas9 ribonucleoprotein-mediated co-editing and counterselection in
the rice blast fungus. Scientific Reports **8,** 14355.

**Fukuoka, S., Saka, N., Koga, H., Ono, K., Shimizu, T., Ebana, K., Hayashi, N., Takahashi,
A., Hirochika, H., Okuno, K., and Yano, M.** (2009). Loss of function of a proline-
containing protein confers durable disease resistance in rice. Science **325,** 998-1001.

Gao, W., Xiao, S., Li, H.-Y., Tsao, S.-W., and Chye, M.-L. (2009). Arabidopsis thaliana
acyl-CoA-binding protein ACBP2 interacts with heavy-metal-binding farnesylated
protein AtFP6. New Phytologist **181,** 89-102.

**Giraldo, M.C., Dagdas, Y.F., Gupta, Y.K., Mentlak, T.A., Yi, M., Martinez-Rocha, A.L.,
Saitoh, H., Terauchi, R., Talbot, N.J., and Valent, B.** (2013). Two distinct secretion
systems facilitate tissue invasion by the rice blast fungus Magnaporthe oryzae. Nat
Commun **4,** 1996.

Gómez De La Cruz, D., Zdrzałek, R., Banfield, M.J., Talbot, N.J., and Moscou, M.J.
(2024). Molecular mimicry of a pathogen virulence target by a plant immune receptor.
bioRxiv**,** 2024.2007.2026.605320.

Guo, T., Weber, H., Niemann, M.C.E., Theisl, L., Leonte, G., Novák, O., and Werner, T.
(2021). Arabidopsis HIPP proteins regulate endoplasmic reticulum-associated
degradation of CKX proteins and cytokinin responses. Mol Plant **14,** 1918-1934.

Hála, M., and Žárský, V. (2019). Protein Prenylation in Plant Stress Responses. In Molecules.

Hensel, G., Kastner, C., Oleszczuk, S., Riechen, J., and Kumlehn, J. (2009).
Agrobacterium-mediated gene transfer to cereal crop plants: current protocols for
barley, wheat, triticale, and maize. Int J Plant Genomics **2009,** 835608.

Holsters, M., Silva, B., Van Vliet, F., Genetello, C., De Block, M., Dhaese, P., Depicker,
A., Inze, D., Engler, G., Villarroel, R., and et al. (1980). The functional organization
of the nopaline A. tumefaciens plasmid pTiC58. Plasmid **3,** 212-230.

Imran, Q.M., Falak, N., Hussain, A., Mun, B.G., Sharma, A., Lee, S.U., Kim, K.M., and
Yun, B.W. (2016). Nitric Oxide Responsive Heavy Metal-Associated Gene AtHMAD1
Contributes to Development and Disease Resistance in Arabidopsis thaliana. Front
Plant Sci **7,** 1712.

Inoue, Y., Vy, T.T.P., Yoshida, K., Asano, H., Mitsuoka, C., Asuke, S., Anh, V.L.,
Cumagun, C.J.R., Chuma, I., Terauchi, R., Kato, K., Mitchell, T., Valent, B.,
Farman, M., and Tosa, Y. (2017). Evolution of the wheat blast fungus through
functional losses in a host specificity determinant. Science **357,** 80-83.

Jones, J.D., and Dangl, J.L. (2006). The plant immune system. Nature **444,** 323-329.

Kang, S.C., Sweigard, J.A., and B, V. (1995). The PWL host specificity gene family in the
blast fungus Magnaporthe grisea. Molecular Plant-Microbe Interactions **8,** 939-948.

Kankanala, P., Czymmek, K., and Valent, B. (2007). Roles for rice membrane dynamics and
plasmodesmata during biotrophic invasion by the blast fungus. Plant Cell **19,** 706-724.

Kanzaki, H., Yoshida, K., Saitoh, H., Fujisaki, K., Hirabuchi, A., Alaux, L., Fournier, E.,
Tharreau, D., and Terauchi, R. (2012). Arms race co-evolution of Magnaporthe
oryzae AVR-Pik and rice Pik genes driven by their physical interactions. The Plant
Journal **72,** 894-907.

Khang, C.H., Berruyer, R., Giraldo, M.C., Kankanala, P., Park, S.-Y., Czymmek, K.,
Kang, S., and Valent, B. (2010). Translocation of Magnaporthe oryzae Effectors into
Rice Cells and Their Subsequent Cell-to-Cell Movement The Plant Cell **22,** 1388-
1403.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics **25,** 1754-1760.

Li, H., Wang, J., Kuan, T.A., Tang, B., Feng, L., Wang, J., Cheng, Z., Skłenar, J.,
Derbyshire, P., Hulin, M., Li, Y., Zhai, Y., Hou, Y., Menke, F.L.H., Wang, Y., and
Ma, W. (2023). Pathogen protein modularity enables elaborate mimicry of a host
phosphatase. Cell **186,** 3196-3207.e3117.

Maidment, J.H.R., Franceschetti, M., Maqbool, A., Saitoh, H., Jantasuriyarat, C.,
Kamoun, S., Terauchi, R., and Banfield, M.J. (2021). Multiple variants of the fungal
effector AVR-Pik bind the HMA domain of the rice protein OsHIPP19, providing a
foundation to engineer plant defense. Journal of Biological Chemistry **296**.

Maqbool, A., Saitoh, H., Franceschetti, M., Stevenson, C.E.M., Uemura, A., Kanzaki, H.,
Kamoun, S., Terauchi, R., and Banfield, M.J. (2015). Structural basis of pathogen

recognition by an integrated HMA domain in a plant NLR immune receptor. eLife **4,** e08709.

Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H., and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and analysis. Protein Sci **32,** e4792.

Mosquera, G., Giraldo, M.C., Khang, C.H., Coughlan, S., and Valenta, B. (2009). Interaction Transcriptome Analysis Identifies Magnaporthe oryzae BAS1-4 as Biotrophy-Associated Secreted Proteins in Rice Blast Disease. American Society of Plant Biologists **21,** 1273–1290.

Mukhi, N., Brown, H., Gorenkin, D., Ding, P., Bentham, A.R., Stevenson, C.E.M., Jones, J.D.G., and Banfield, M.J. (2021). Perception of structurally distinct effectors by the integrated WRKY domain of a plant immune receptor. Proc Natl Acad Sci U S A **118**.

Mutiga, S.K., Rotich, F., Were, V.M., Kimani, J.M., Mwongera, D.T., Mgonja, E., Onaga, G., Konaté, K., Razanaboahirana, C., Bigirimana, J., Ndayiragije, A., Gichuhi, E., Yanoria, M.J., Otipa, M., Wasilwa, L., Ouedraogo, I., Mitchell, T., Wang, G.-L., Correll, J.C., and Talbot, N.J. (2021). Integrated Strategies for Durable Rice Blast Resistance in Sub-Saharan Africa. Plant Disease **105,** 2749-2770.

O'Lexy, R., Kasai, K., Clark, N., Fujiwara, T., Sozzani, R., and Gallagher, K.L. (2018). Exposure to heavy metal stress triggers changes in plasmodesmatal permeability via deposition and breakdown of callose. Journal of Experimental Botany **69,** 3715-3728.

Oikawa, K., Fujisaki, K., Shimizu, M., Takeda, T., Nemoto, K., Saitoh, H., Hirabuchi, A., Hiraka, Y., Miyaji, N., Białas, A., Langner, T., Kellner, R., Bozkurt, T.O., Cesari, S., Kroj, T., Banfield, M.J., Kamoun, S., and Terauchi, R. (2024). The blast pathogen effector AVR-Pik binds and stabilizes rice heavy metal-associated (HMA) proteins to co-opt their function in immunity. PLOS Pathogens **20,** e1012647.

Oliveira-Garcia, E., Yan, X., Oses-Ruiz, M., de Paula, S., and Talbot, N.J. (2023a). Effector-triggered susceptibility by the rice blast fungus Magnaporthe oryzae. New Phytologist **n/a**.

Oliveira-Garcia, E., Tamang, T.M., Park, J., Dalby, M., Martin-Urdiroz, M., Rodriguez Herrero, C., Vu, A.H., Park, S., Talbot, N.J., and Valent, B. (2023b). Clathrin-mediated endocytosis facilitates the internalization of Magnaporthe oryzae effectors into rice cells. The Plant Cell **35,** 2527-2551.

Ortiz, D., de Guillen, K., Cesari, S., Chalvon, V., Gracy, J., Padilla, A., and Kroj, T. (2017). Recognition of the Magnaporthe oryzae Effector AVR-Pia by the Decoy Domain of the Rice NLR Immune Receptor RGA5. The Plant Cell **29,** 156-168.

Ou, S.H. (1980). Pathogen Variability and Host-Resistance in Rice Blast Disease. Annu Rev Phytopathol **18,** 167-187.

Radakovic, Z.S., Anjam, M.S., Escobar, E., Chopra, D., Cabrera, J., Silva, A.C., Escobar, C., Sobczak, M., Grundler, F.M.W., and Siddique, S. (2018). Arabidopsis HIPP27 is a host susceptibility gene for the beet cyst nematode Heterodera schachtii. Mol Plant Pathol **19,** 1917-1928.

Sakulkoo, W., Osés-Ruiz, M., Oliveira Garcia, E., Soanes, D.M., Littlejohn, G.R., Hacker,
C., Correia, A., Valent, B., and Talbot, N.J. (2018). A single fungal MAP kinase
controls plant cell-to-cell invasion by the rice blast fungus. Science **359,** 1399-1403.

Sugihara, Y., Abe, Y., Takagi, H., Abe, A., Shimizu, M., Ito, K., Kanzaki, E., Oikawa, K.,
Kourelis, J., Langner, T., Win, J., Białas, A., Lüdke, D., Contreras, M.P., Chuma,
I., Saitoh, H., Kobayashi, M., Zheng, S., Tosa, Y., Banfield, M.J., Kamoun, S.,
Terauchi, R., and Fujisaki, K. (2023). Disentangling the complex gene interaction
networks between rice and the blast fungus identifies a new pathogen effector. PLOS
Biology **21,** e3001945.

Suzuki, N., Yamaguchi, Y., Koizumi, N., and Sano, H. (2002). Functional characterization
of a heavy metal binding protein CdI19 from Arabidopsis. The Plant Journal **32,** 165-
173.

Sweigard, J.A., Carroll, A.M., Kang, S., Farrall, L., Chumley, F.G., and Valent, B. (1995).
Identification, cloning, and characterization of PWL2, a gene for host species
specificity in the rice blast fungus. Plant Cell **7,** 1221-1233.

Talbot, N.J. (2003). On the trail of a cereal killer: Exploring the biology of Magnaporthe
grisea. Annu Rev Microbiol **57,** 177-202.

Talbot, N.J., Ebbole, D.J., and Hamer, J.E. (1993). Identification and characterization of
MPG1, a gene involved in pathogenicity from the rice blast fungus Magnaporthe grisea.
Plant Cell **5,** 1575-1590.

Valent, B., Farrall, L., and Chumley, F.G. (1991). Magnaporthe-Grisea Genes for
Pathogenicity and Virulence Identified through a Series of Backcrosses. Genetics **127,**
87-101.

Wang, Z., Zhang, H., Li, Y., Chen, Y., Tang, X., Zhao, J., Yu, F., Wang, H., Xiao, J., Liu,
J., Zhang, X., Sun, L., Xie, Q., and Wang, X. (2023). Isoprenylation modification is
required for HIPP1-mediated powdery mildew resistance in wheat. Plant Cell Environ
46, 288-305.

Were, V., and Talbot, N.J. (2023). Breaking the biotrophic interfacial complex: How genome
editing can lead to rice blast resistance. Molecular Plant **16,** 1243-1245.

Wilson, R.A., and Talbot, N.J. (2009). Under pressure: investigating the biology of plant
infection by Magnaporthe oryzae. Nat Rev Microbiol **7,** 185-195.

Yan, X., Tang, B., Ryder, L.S., MacLean, D., Were, V.M., Eseola, A.B., Cruz-Mireles, N.,
Ma, W., Foster, A.J., Osés-Ruiz, M., and Talbot, N.J. (2023). The transcriptional
landscape of plant infection by the rice blast fungus Magnaporthe oryzae reveals
distinct families of temporally co-regulated and structurally conserved effectors. The
Plant Cell **35,** 1360-1385.

Yu, D., Fan, R., Zhang, L., Xue, P., Liao, L., Hu, M., Cheng, Y., Li, J., Qi, T., Jing, S.,
Wang, Q., Bhatt, A., and Shen, Q.-H. (2023). HvWRKY2 acts as an immunity
suppressor and targets HvCEBiP to regulate powdery mildew resistance in barley. The
Crop Journal **11,** 99-107.

Yuen, E.L.H., Tumtas, Y., King, F., Ibrahim, T., Chan, L.I., Evangelisti, E., Tulin, F.,
Skłenar, J., Menke, F.L.H., Kamoun, S., Bubeck, D., Schornack, S., and Bozkurt,
T.O. (2024). A pathogen effector co-opts a host RabGAP protein to remodel pathogen
interface and subvert defense-related secretion. Science Advances **10,** eado9516.

Zdrzałek, R., Xi, Y., Langner, T., Bentham, A.R., Petit-Houdenot, Y., De la Concepcion,
J.C., Harant, A., Shimizu, M., Were, V., Talbot, N.J., Terauchi, R., Kamoun, S.,
and Banfield, M.J. (2024). Bioengineering a plant NLR immune receptor with a robust
binding interface toward a conserved fungal pathogen effector. Proceedings of the
National Academy of Sciences **121,** e2402872121.

Zhang, H., Zhang, X., Liu, J., Niu, Y., Chen, Y., Hao, Y., Zhao, J., Sun, L., Wang, H.,
Xiao, J., and Wang, X. (2020). Characterization of the Heavy-Metal-Associated
Isoprenylated Plant Protein (HIPP) Gene Family from Triticeae Species. Int J Mol Sci
21.

Zhang, S., and Xu, J.R. (2014). Effectors and effector delivery in Magnaporthe oryzae. PLoS
Pathog **10,** e1003826.

Zhang, X., He, D., Zhao, Y., Cheng, X., Zhao, W., Taylor, I.A., Yang, J., Liu, J., and Peng,
Y.-L. (2018). A positive-charged patch and stabilized hydrophobic core are essential
for avirulence function of AvrPib in the rice blast fungus. The Plant Journal **96,** 133-
146.

Zschiesche, W., Barth, O., Daniel, K., Böhme, S., Rausche, J., and Humbeck, K. (2015).
The zinc-binding nuclear protein HIPP3 acts as an upstream regulator of the salicylate-
dependent plant immunity pathway and of flowering time in Arabidopsis thaliana. New
Phytologist **207,** 1084-1096.

1018

1019
Fig.1. *PWL2* **expression is regulated in a Pmk1-dependent manner during host infection.**
(**A+B**) Micrographs and line scan graph showing Pwl2 secreted through the Biotrophic
Interfacial Complex (BIC) in rice cells during early infection. Conidial suspension at 1x 10^5

mL^{-1} of *M. oryzae* strain co-expressing two BIC localised effectors, Bas1-GFP and Pwl2-mRFP

were inoculated onto a susceptible cultivar Moukoto rice leaf sheath and images captured at 26

hpi. Fluorescence of the two effectors was observed as small punctate signals in the same BIC.

(**C+D**) Co-infection assay of rice leaf sheath with two different *M. oryzae* strains, one

expressing Pwl2-mRFP and the other Pwl2-GFP at 30 hours post-infection (hpi). Micrographs

and line scan graph shows there is an absence of mixed fluorescence confirming the BIC does

not contain Pwl2 transferred from rice cells. BICs indicated by magenta arrows for mRFP and

green arrows for GFP. (**E**) Schematic illustration to describe the workflow used to test genes

regulated in a Pmk1-dependent manner. Rice leaf sheaths were infected with a *M. oryzae*

pmk1^{AS} mutant spores before mock or 1NA-PP1 treatment. Treated and mock treated, infected

leaf sheaths were trimmed and used for RNA isolation followed by sequencing. Figure created

with BioRender https://biorender.com/. (**F**) Bar charts to show that a subset of effectors is

regulated in a manner that requires Pmk1. Gene expression is shown as TPM (transcripts per

million) at 32 hpi. (**G-L**) Micrographs showing expression of Pwl2-GFP by *M. oryzae pmk1^{AS}*

in leaf sheaths of a susceptible rice line CO39 using conidial suspension at 1x 10^5 mL^{-1}.

Fluorescence of Pwl2 at different stages of infectious hyphal progression, starting with early

stage of infection, Stage 1 (30-36 hpi) where a newly differentiated bulbous hyphae is formed

as per (Sakulkoo *et al.*, 2018). Analysis was also carried out at later stages of infection

including Stage 2 (36-40 hpi) where a primary invaded cell is filled with differentiated bulbous

hyphae) and Stage 3 (40-48 hpi) where colonization of primary invaded cell is complete and

there is full invasion of secondary invaded cells. To test Pmk1 inhibition, inoculated leaf

sheaths were treated with 5 μM 1NA-PP1 at 26 hpi. (**M**) Inhibition of Pwl2 by 1NA-PP1 is

quantified as percentage of cells showing Pwl2 fluorescence in the BIC. Arrows indicate

fluorescence in the BIC. Scale bars represent 20 μm. The primary and secondary invaded cells

are indicated with 1 and 2 respectively. Significance between groups of samples was performed

using Unpaired Student's t-test. ****$P<0.0001$, **$P<0.01$, *$P<0.05$, NS indicates no

significant difference.

1050

1051

1052

1053 **Fig. 2**. **Pwl2 is highly conserved in isolates of *M. oryzae* and displays conserved structural**

1054 **features.**

(**A)** Phylogenetic analysis and *PWL* gene family distribution in *M. oryzae*. A maximum parsimony tree was generated using kSNP3 to include isolates from different host-limited forms of *M. oryzae* including isolates that infect *Oryza sativa* (rice), *Eleusine* spp. (finger millet), *Hordeum vulgare* (barley), *Setaria* spp. (foxtail millet), *Triticum aestivum* (wheat), *Lolium* spp. (rye grass), *Brachiaria* spp. (signal grass), *Panicum* spp. (torpedo grass), *Eragrostis* spp. (weeping lovegrass), *Stenotaphrum* spp. (buffalo grass), *Cynodon* spp. (Bermuda grass) and *Urochloa* spp. (signal grass) (Ou, 1980; Talbot, 2003; Cruz and Valent, 2017; Inoue *et al.*, 2017), as well as *Magnaporthe* species that infect *Digitaria sanguinalis* (crabgrass) and *Pennisetum* spp. (pearl millet). We used Pwl1 (BAH22184.1), Pwl2 (QNS36448.1), Pwl3 (AAA80240.1) and Pwl4 (AAA80241.1) protein sequences to query the presence or absence of each gene using tblastn. The heatmap indicates the presence/absence of genes in the *PWL* family. *PWL1* is predominantly present in group EC-1I (Asuke *et al.*, 2020) of *Eleusine*-infecting isolates and some *Oryza*-infecting isolates, but largely absent from other host-specific lineages, except one *Eragrostis*-infecting isolate, EtK19-1, and one *Cynodon*-infecting isolate, Cd88215, but not in *Digitaria* and *Pennisetum* spp lineages. *PWL3* is present in most *Oryza*-infecting isolates, some *Setaria*, *Lolium*, and *Eleusine*-infecting isolates but largely absent in *Eragrostis* and *Triticum* lineages, as well as *Digitaria* and *Pennisetum* lineages. *PWL4* is present in *Eleusine*, *Eragrostis, Lolium* and *Triticum*-infecting isolates, but only two *Oryza*-infecting isolates. Several *Pennisetum* and *Digitaria*-infecting isolates, however, carry *PWL4. PWL2* is found in most host-limited forms of *M. oryzae* and related *Magnaporthe* species but absent in *Brachiaria, Setaria, Panicum, Cynodon* and *Urochloa*. (**B**) Superimposition of different Pwl2 variants predicted using AlphaFold3 onto the resolved Pwl2 structure (magenta) indicating region of polymorphism. Different color represent different variant as follows, Pwl2-2 (teal), Pwl2-3 (brick red), Pwl2-TH3 (dark olive green), Pwl2-IN15 (dark teal), Pwl2-KE210 (blue), Pwl2-CKF3584 (dark grey), Pwl2-EtKY19-1 (orange), Pwl2-DS9461 (light grey). The superimposition shows overall structural conservation in MAX-fold, without the signal peptide and the C-terminus. The variants of Pwl2 are named with corresponding isolate name and grouped according to host species. Varying residues are colored according to the genome from which they were identified.

Fig. 3. *PWL2* has undergone copy number expansion in *M. oryzae* field isolates.

(A) Schematic diagram showing the estimated chromosomal location of *PWL2* on multiple loci
(Chr3 and 6) of the reference genome 70-15 and on equivalent region of assembled contigs of
laboratory strain Guy11 genome. *PWL2* is flanked by *POT2* and *MGR583* repeated sequences
suggesting a possible involvement in translocation of events into different loci in the genome.

Arrows indicate location of *POT2* and *MGR58* while *PWL2* is labelled in magenta. **(B)** A k-

mer analysis on sequenced raw reads was used to determine copy number variation of *PWL*

family genes in different *M. oryzae* isolates. Plot shows high copy number of *PWL2* in analyzed

genomes (n = 286) compared to the other *PWL* gene family members, another effector *SLP1,*

and selected control genes *MPG1* and *SEC5*. Similarly, *AVR-Pik* shows multiple copies in

different isolates. Copy number of *PWL2* in selected isolates Guy11, KE002, BF48, 70-15 is

indicated with Ina168 that lack *PWL2* used as a negative control.

1098

1099

1100

1101 **Fig. 4. CRISPR-Cas9 *Δpwl* mutants demonstrate Pwl2 as both host range and a**

1102 **virulence factor**

(A) Schematic illustration of the CRISPR-Cas9 mediated gene editing process for inserting the Hygromycin Phosphotransferase (*HPH*) gene cassette at the *PWL2* locus. The guided sequence (sgPWL2) directs Cas9 to introduce a double stranded break at the *PWL2* locus, with the DNA repair template containing the Hygromycin resistance gene cassette and flanking regions of the *PWL2* gene. This leads to mutations in the form of indels or gene replacements. Figure created with BioRender https://biorender.com/. Positive mutants were identified by amplifying the hygromycin cassette from transformants with the three copies deleted. (B) Comparison of disease symptoms in weeping lovegrass (*E. curvula*) infected with different *M. oryzae* isolates. In the top panel, typical disease lesions produced by *E. curvula*-infecting isolate G17 (-*PWL2*) and a rice infecting isolate Ina168 (-*PWL2*) are shown. In the middle panel, CRISPR-Cas9 deletion mutants exhibit gain of virulence on weeping lovegrass compared to Guy11. In the bottom panel, complemented strains with both native and constitutive RP27 promotors showed a loss of virulence on weeping lovegrass. (C) *Δpwl2* mutants display reduced pathogenicity on rice cultivar CO39. Conidial suspensions from Guy11, *Δpwl2* (T6) and complemented *Δpwl2*(T6) +*PWL2*p:*PWL2*) were used to inoculate 21-day-old seedlings of the blast-susceptible cultivar CO39, and disease symptoms recorded after 5 dpi. The box plots show the lesion density (D) and lesion size (E) of seedlings infected with Guy11, *Δpwl2* (T6) and complemented *Δpwl2* (T6 +*PWL2*p:*PWL2*). (F) The *Δpwl2* mutant exhibits enhanced pathogenicity on barley cultivar Baronesse (*Mla3*). Conidial suspensions from Guy11 and *Δpwl2* T6 were used to inoculate 10-day old seedlings of barley lines Golden Promise, Siri, Nigrate and Baronesse, and disease symptoms recorded after 5 dpi. Conidial suspensions at 1×10^5 mL^{-1} spores/mL was used for infection assays. (G) The box plot shows lesion size of barley seedlings infected with Guy11 and *Δpwl2* (T6). The lower horizontal line shows the minimum value, and the upper horizontal line shows the maximum value. The lower border and upper border of the box shows the lower quartile and upper quartile, respectively. The line in the box shows the median. Significance between groups of samples was performed using Unpaired Student's t-test. ****P<0.0001, **P<0.01, *P<0.05, NS indicates no significant difference.

Fig. 5. Pwl2 suppresses PAMP-induced ROS in transgenic barley lines

(**A**) Schematic illustration to describe the workflow used to generate transgenic plants, test for

PTI response and identify putative Pwl2 interactors using discovery proteomics. ROS

production in leaf discs collected from 4–week-old stable transgenic lines expressing Pwl2-

YFP compared to cv. Golden Promise induced by 1 uM flg22 (**B-C**) or chitin (**D-E**) (n > 8).

Points represent mean; error bars represent SEM. For ROS assay, line graphs (**B+D**) points

represent mean per time point and dot plots represent cumulative ROS (**C+E**) production over

60 min, error bars represent SEM. For dot plot, graphs show the average cumulative photon

count for cv. Golden Promise control and two independent Pwl2-YFP lines. The lower

horizontal line shows the minimum value, and the upper horizontal line shows the maximum

value, the middle line shows the mean value. (**F**) Leaf drop infection on two independent barley

transgenic lines expressing Pwl2-YFP compared to infection on wild type cv. Golden Promise

3-4 dpi. (**G**) The dot plot represents the disease lesion length on cv. Golden Promise compared

to two independent Pwl2-YFP transgenic lines. Unpaired Student's t-test was performed to

determine significant differences ****P<0.0001, ***P<0.001, **P<0.01. These experiments

were repeated three times to obtain consistent result.

1148

1149

1151   **Fig. 6. Pwl2 interacts with HIPP43 and its orthologs from other grass species.**

(**A**) Putative Pwl2 interacting peptides were immunoprecipitated from protein extracts of 3-week-old stable cv. Golden Promise transgenic lines expressing Pwl2-YFP or free cytoplasmic YFP using anti-GFP antibodies, and LC-MS/MS was performed to identify unique putatively interacting peptides. The scale 0-20 represents the Log2fold change in peptides when compared to the control. (**B**) One-to-one yeast two-hybrid between Pwl2 and selected top candidates from IP-MS analysis, Pwl2 and Putative Pwl2 Interacting Proteins (PPIPs). HMA integrated in rice NLR Pikm-1 and Pikp-1 were used as specificity controls. Simultaneous co-transformation of pGADT7-Pwl2 (prey vector) and pGBK-PPIPs (bait vector) and PGBKT7-53 and pGADT7-T (positive control) pGADTT7-T and pGBKT7-Lam (negative control) into Y2H gold strain. Positive interaction resulted in the activation of four reporter genes and growth on high-stringency medium (−Ade, −Leu, –Trp, _His +X-α-gal and 3-amino-1,2,4-triazole). Co-transformation also activates the expression of *MEL1*, which results in the secretion of α–galactosidase and the hydrolysis of X-α-gal in the medium, turning the yeast colonies blue. HIPP43 exclusively interacts with Pwl2 in SD/-L-W-H-A X-α-gal medium with 3AT added. These experiments were repeated several times over three years obtaining consistent result. (**C**) Co-immunoprecipitation (co-IP) of Pwl2-YFP and Myc-HIPP43 or (**D**) mCherry-HIPP43 in *N. benthamiana* leaves. C-or N-terminal GFP tagged Pwl2 and C-terminal Myc-HIPP43 was cloned into the vector pGW514 and transformed into *Agrobacterium* strain *GV3101* and co-infiltrated into *N. benthamiana* leaves and left to incubate for 48 h. Immunoprecipitates were obtained with anti-GFP affinity matrix beads and probed with anti-GFP-peroxidase, anti-mCherry-peroxidase and anti-Myc-peroxidase (HRP-conjugated) antibodies. Total protein extracts were also probed with appropriate (HRP-conjugated) antibodies. (**E**) Yeast-two-hybrid analysis shows; upper panel, *Hv*HIPP43 interacts with *PWL* gene family products Pwl1, Pwl3, Pwl4 and variant *pwl2-3*. (**F**) Pwl2 interacts with HIPP43 homologs from rice (*Os*HIPP43), wheat (*Taes*HIPP43), weeping love grass (*Ec*HIPP43), foxtail millet (*SitiHIPP43*) and wild wheat (HIPP1-V). These experiments were repeated several times over three years obtaining consistent result.

1180

Fig. 7. *Hv*HIPP43 suppresses PAMP-induced ROS in transgenic barley and is stabilized

by Pwl2.

(A-D) ROS production measured from leaf disks collected from 4–week-old stable transgenic

line expressing YFP-HIPP43 and cv. Golden promise (control) in the absence and presence of

1 uM flg22 or 1 mg/ml Chitin (n > 8). For ROS assay, line graphs **(A+C)** points represent mean

per time point and dot plots represent cumulative ROS **(B+D)** production over 60 min, error

bars represent SEM for cv. Golden Promise control and two independent YFP-HIPP43 lines.

For dot plot, graphs show the average cumulative photon count for cv. Golden Promise control

and two independent YFP-HIPP43 lines. **(E)** Leaf drop infection on barley transgenic lines

expressing YFP-HIPP43 compared to wild type Golden Promise. Conidial suspensions at 1

$\times 10^5$ mL^{-1} spores/mL from Guy11 were used for inoculation. Disease symptoms were recorded

after 4 dpi. **(F)** Dot plot showing lesion length on barley lines infected with Guy11. All

experiments were repeated three times giving consistent results. **(G+H)** Micrographs and line

scan graph showing mCherry-HIPP43 localizing as small puncta on the plasma membrane

when expressed in *N. benthamiana*. Staining of callose using aniline blue overlaps with

mCherry-HIPP43, confirming mCherry-HIPP43 localises exclusively at the plasmodesmata

(PD) localization and cytoplasmic mobile bodies (MB) localization is absent. **(I+J)**

Micrographs and line scan graph showing the presence of YFP-Pwl2, alters mCherry-HIPP43

localisation, which was observed to translocate to the cytoplasm, mobile cytoplasmic bodies.

(K+L) Micrographs and line scan graph showing mCherry-HIPP43 remains in the PD in the

presence of cytoplasmically localised AVR-PikE**. (M+N)** Micrographs and line scan graph

showing mCherry-HIPP43 remains in the PD in the presence of cytoplasmically and PD

localised MEP3**. (O+P)** Micrographs and line scan graph showing mCherry-HIPP43 remains

in the PD in the presence of Pwl2$^{\text{SNDEYWY}}$, a mutant that does not interact with HIPP43**.** Dotted

lines in the overlay panels correspond to distance and direction of line intensity plot. Scale bars

represent 20 μm**.** For dot plots, the lower horizontal line shows the minimum value, the upper

horizontal line shows the maximum value, the middle line shows the mean value. Unpaired

Student's t-test was performed to determine significant differences *****P*<0.0001,

****P*<0.001, ***P*<0.01. **(Q)** Western blot to show that Pwl2 does not degrade HIPP43. Total

protein extracts from *N. benthamiana* leaves co-infiltrated with YFP-Pwl2, MEP3 or AVR-

PikE and mCherry-HIPP43 were immunoblotted and probed with anti-GFP-peroxidase (left)

or anti-mCherry-peroxidase (right) (HRP-conjugated) antibodies. Microscope imaging

experiments were repeated several times over two years giving consistent result.

1214
1215 **Fig. 8. Pwl2^{SNDEYWY} does not complement Mla3 recognition and virulence on a blast-**
1216 **susceptible rice cultivar CO39.**
1217
(**A and B**) Complemented *Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}*) positive transformants were screened
using PCR to amplify *PWL2* coding sequence. (**C**) Bar charts showing relative expression as
log2 fold change of *Pwl2^{SNDEYWY}* in two selected transformants, C6 and C14 using qRT-PCR.
Detached leaves of 10-day old seedlings of barley were inoculated with
*Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}*) and infected tissue collected 40 hpi and used for RNA isolation
(3 biological replicates), cDNA synthesised and samples used for qRT-PCR. (**D and E**)
*Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}* complemented strains C6 and C14 produced compatible disease
lesions on barley cultivar Baronesse (*+Mla3*) (**D**) and Nigrate (*-Mla3*) (**E**). Conidial

suspensions at 1 ×10^5 mL^{-1} spores/mL from Guy11, *Δpwl2* and complemented

*Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}* were used to inoculate 10-day old seedlings of barley and disease

symptoms recorded after 5 dpi. (**F-H**) Complemented *Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}* display

reduced pathogenicity on rice cultivar CO39. Conidial suspensions at 1 ×10^5 mL^{-1} spores/mL

from Guy11, *Δpwl2* T6 and complemented *Δpwl2+PWL2*p:*Pwl2^{SNDEYWY}* were used to inoculate

21-day-old seedlings of the blast-susceptible cultivar CO39, and disease symptoms recorded

after 5 dpi. Significance between groups of samples was performed using Unpaired Student's

t-test. ***P<0.0001, **P<0.01, *P<0.05, NS indicates no significant difference.

1234
1235
1236

1237

Fig. S1. *M. oryzae* **Pwl2 is co-expressed during plant infection with other known effector genes.**

(**A**) Attached leaves of three-weeks old rice seedlings inoculated with KE002 conidia suspension at 0, 48, 72 and 120 hpi. Samples were collected at 24, 36, 48, 59 and 72 hpi and used for RNA isolation (3 biological replicates). (**B**) RNA isolated from infected rice leaves were sequenced and the reads normalized against *M. oryzae* mycelia KE002 RNA to determine differentially expressed genes. A weighted gene co-expression network analysis grouped gene expression profile into 10 co-expression modules. The line graph (WGCNA) shows the representative eigengene for module 4 in which *PWL2* is co-expressed with other known effector protein encoding genes, *AVR-Pi9*, *SLP1*, *BAS1-4* and suppressors of cell death effectors 3, 6, 7 and 8 that possibly serve related function during biotrophic phase of invasion. (**C***)* Heatmap showing relative expression levels (log2fold change) of *PWL2* and other known effectors during infection.

Fig. S2. *PWL2* is highly polymorphic compared to other members in the *PWL* family.

(**A**) Multiple amino acid sequence alignment of Pwl2 coding sequence and naturally occurring alleles. The region between His-87 and Ser-92 shows highest number of polymorphisms and is the region important for cognate resistance gene recognition. Multiple amino acid sequence alignment of *PWL1* (**B**), *PWL3* (**C**) and *PWL4* (**D**) Alleles identified from genomes of field isolates. Alignments were carried using ClustalW (Jalview). (**E**) Neighbor-joining tree based on amino acid sequence of alleles of the *PWL* gene family. Tree IDs contains isolate name from which allele was identified, and each dot represents presence in each host-limited form. Unlike other family members, *PWL2* alleles are present in host-limited forms apart from lineage 2 of *Brachiaria* and *Setaria. Panicum, Cynodon* and *Urochloa*-infecting isolates.

Fig. S3. Two *PWL2* alleles are not recognized by the barley resistance gene Mla3.

A subset of isolates carrying alleles of *PWL2* were used in leaf drop inoculation on (**A**) barley accessions cv. Nigrate (-*Mla3*) and (**B**) cv. Baronesse (+*Mla3*). Conidial suspensions at 1×10^5 mL^{-1} spores/mL were harvested from Guy11 and selected isolates with *PWL2* alleles and used to inoculate 10-day-old seedlings of cv. Nigrate (-*Mla3*) and cv. Baronesse (+*Mla3*). Disease lesions were scored 5 days post-infection. Isolates BF5 (variant *Pwl2*), BTJP4-16 (*pwl2-3)*, BN0293 (variant *Pwl2*), PM1 (variant *Pwl2*), Guy11 (+*PWL2*), JUM1 (*pwl2-2*), and P131 (-*PWL2*) were used for leaf drop infection assay. As expected Guy11 could infect cv. Nigrate but not cv. Baronesse. BF5 and PM1 did not produce blast disease on barley cv. Baronesse*,* suggesting they express *PWL2* alleles recognised by Mla3, (or express effectors recognised by additional R-genes contained (in barley cv. Baronesse). Notably, PM1 could also not infect cv. Nigrate (-*Mla3*). P131 does not carry *PWL2* and caused blast disease on cv. Baronesse as expected. JUM1 (*pwl2-2*), BTJP4-16 (*pwl2-3)* and BN0293 (*pwl2-BN0293*) all caused blast

disease on cv. Baronesse (**C**) Summary of disease reaction of isolates with distinct *PWL2* alleles on Nigrate (-*Mla3*) and Baronesse (+*Mla3*). Observations were consistent in three independent infection replicates.

1309

Fig. S4. Southern blot analysis to determine copy number variation of *PWL2* in selected

***M. oryzae* isolates.**

(**A**) Genomic DNA was digested using different enzyme combination *Bam*HI/*Pst*I (upper

panel), *Pst*I/*Afl*II (upper middle panel), *Afl*II single digest (lower middle panel) or *Pst*I single

digest (lower panel) and probed using the *PWL2* 438 bp probe. Size estimates are from the

PWL2 loci annotated *M. oryzae* 70-15 genome. (**B**) Left panel, the order of digested genomic

DNA from different isolates is indicated on the lanes. Right panel, when Guy11 genomic DNA

was digested using *Pst*1 and *Afl*II, we detected two bands of 3.5kb and 6kb. When digested

with *Afl*II *in a* single digest were detected two bands of 3.5 kb and 10 kb. While in a single

digest using *Pst*I only produce a single band of 6.5 kb.

1321

1322

1323

Fig. S5. *M. oryzae Δpwl2* deletion mutants show gain of virulence on weeping lovegrass and cultivar Baronesse (*Mla3*).

(**A**) Reads alignment display of *PWL2* locus on Chromosome 6 from four sequenced *pwl2* mutant strains, T12, T5, T4 and T6 (lower panel). No reads aligned to the MGG_04301T0 (*PWL2*) locus for T6 and T12 and partial alignment for T4 and T5. Read depth around *PWL2* locus is higher compared to the neighbouring gene MGG_17636T0 suggesting the *PWL2* locus occurs multiple times. Read coverage and locus length are indicated. (**B**) Guy11, T12, T5, T4,

T6 were grown at 25^0 C, on CM media for 10 days before imaging. (**C**) Disease symptoms on

weeping lovegrass (*E. curvula*) and barley cultivar Baronesse (*Mla3*) to show gain of virulence

for CRISPR-Cas9 deletion mutants T5 and T12. Conidial suspensions at 1×10^5 mL^{-1}

spores/mL from Guy11, *Δpwl2* (T5) and *Δpwl2* (T12) were used to inoculate 10-day-old

seedlings of barley cultivar Baronesse (*Mla3*) or weeping lovegrass, and disease symptoms

recorded after 5 dpi. Observations were consistent in three independent infection replicates.

1337

Fig. S6. Pwl2 interacts with two copies of *Hv*HIPP43 identified by IP-MS.

(**A**) Y2H assay to show that Pwl2 (in pGADT7 vector) and HIPP43 (in pGBKT7 vector) do

not show autoactivity (**B**) Table showing peptides identified by LC-MS that mapped to two

copies of HIPP43. (**C**) Pwl2 interacts with the two copies of HIPP43 in a yeast-two-hybrid

assay. (**D**) Maximum likelihood phylogenetic tree of HMA domains from diverse grass species.

HMA domains were identified from proteins from the grass species *B. distachyon*, *H. vulgare*,

T. aestivum, *O. sativa*, *Oropetium thomaeum*, *Sorghum bicolor*, *Setaria italica*, and *Zea mays*.

Structure guided multiple sequence alignment was performed using MAFFT-DASH and

phylogenetic analysis with RAxML (v8.2.12). Scale bar indicates substitutions per site. Dots

on branches indicate support of 80% or more based on 1,000 bootstraps. The HIPP43 family

formed a distinct clade with bootstrap support of 98%.

1351

1352

Fig. S7 Phylogenetic analysis of HIPP43 and expression profile during the rice blast infection on a susceptible rice cv. CO39.

(**A**) Full length coding sequence was identified for HIPP43 gene family members, aligned using MUSCLE translation alignment, and phylogenetic analysis with RAxML (v8.2.12). Barley and wheat (i.e. *Triticeae* lineage) have three copies of HIPP43 per haploid genome. Dots on branches indicate support of 80% or more based on 1,000 bootstraps. (**B**) RNA-seq analysis of

1359   *OsHIPP43* expression during infection of a susceptible rice cv. CO39 by Guy11. Hour post

1360   inoculation is shown on the X –axis. HIPP43 expression is shown as relative mean expression

1361   TPM (transcripts per million) at different stages of infection.

1362
1363

Fig. S8. HIPP43 isoprenylation motif is required for PD localisation.

(**A +B**) A schematic representation of domain structure of HIPP43 showing the HMA domain region and the isoprenylation motif at the C-terminus. Residues occurring at the C-terminus of HIPP43 orthologs in rice (Os-HIPP43), wheat (Taes-HIPP43), foxtail millet (Siti-HIPP43) and weeping lovegrass (Ec-HIPP43) are indicated. A multiple alignment was generated using Clustal X in Jalview and the conserved motif visualized on Weblogo. (**C-D**) Micrographs showing localization of HIPP43 without the isoprenylation motif (-Iso-mCherry-HIPP43) and of WT mCherry-HIPP43. (**C**) Deletion of the isoprenylation abolishes PD localisation but HIPP43 localises in the nucleus, the plasma membrane after 48h. Co-expression with a plasma membrane marker LTi6b-GFP and aniline blue staining confirmed localisation at the plasma membrane and lack of PD localisation. White arrows indicate regions of HIPP43 localisation away from aniline blue stained PD (**D**) Micrographs showing typical mCherry-HIPP43 localizing as small puncta on the plasma membrane when expressed in *N. benthamiana*. Staining of callose using aniline blue overlaps with mCherry-HIPP43, confirming mCherry-

1379   HIPP43 localises exclusively at PD. White arrows indicate regions of co-localisation between

1380   HIPP43, and aniline blue stained PD. Scale bars represent 20 µm.

Fig. S9. Pwl1, Pwl3, Pwl4 co-localise with HIPP43 and Pwl2 co-localises with HIPP43 orthologs from other grass species.

(**A**) Micrographs showing localization of YFP-Pwl1, YFP-Pwl3 and YFP-Pwl4. Like YFP-Pwl2, Pwl1, Pwl3 and Pwl4 showed localisation in cytoplasm and cytoplasmic mobile structures. (**B-D**) Micrographs showing co-localisation of Pwl1, Pwl3 and Pwl4 with m-Cherry-HIPP43. (**B**) In presence of Pwl1 mCherry-HIPP43 translocates to the cytoplasm and

partially in cytoplasmic mobile structures. **(C)** Pwl3 did not dislodge mCherry-HIPP43 from

the PD while Pwl4 **(D)** only showed cytoplasmic co-localisation. (**E-G**) Micrographs showing

HIPP43 orthologs fluorescently tagged to mCherry. (**E**) *Taes*HIPP43 (wheat) localizes in the

nucleus and cytoplasm while (**F**) *Ec*HIPP43 (weeping lovegrass) (**G**) *Siti*HIPP43 (foxtail

millet) localize as small puncta equivalent to PD. Images shown as maximum projections **(H-**

J) Micrographs showing co-localisation of YFP-Pwl2 with mCherry-*Siti*HIPP43. mCherry-

*Siti*HIPP43 localisation, was altered to the cytoplasm, cytoplasmic mobile structures. mCherry-

*Siti*HIPP43 fluorescence signal was higher in presence of YFP-Pwl2. White arrows indicate

regions of co-localisation and blue arrows show aniline blue stained PD. Scale bars represent

20 μm.

1398

1399

1400

Fig. S10. Pwl2 stabilises and promotes HIPP43 cytoplasmic localisation.

(A-D) Micrographs showing localization of free-YFP (**A**), AVR-PikE-YFP (**B**), YFP-Pwl2 (**C**)

and MEP3-YFP (**D**). Both free-YFP and AVR-PikE-YFP show localisation in the cytoplasm

while YFP-Pwl2 showed localisation in cytoplasm and cytoplasmic mobile structures. MEP3

localises in the cytosol and PD (**E-I**) Micrographs and line scan graphs showing the presence

of YFP-Pwl2 in the cytosol and puncta. Lack of co-localization with aniline blue stain confirms

that, without HIPP43, Pwl2 does not accumulate at PD but rather mobile puncta. (**H-K**)

Micrographs showing changes of mCherry-HIPP43 localisation when co-expressed with (**H**)

YFP-Pwl2, (**I**) AVR-PikE-YFP (**J**) EV-YFP and (**K**) MEP3-YFP. Staining of callose using

aniline blue is used to differentiate between immobile PD and mobile cytoplasmic mCherry-

HIPP43 co-localisation with YFP-Pwl2. In presence of AVR-PikE, EV and MEP3 aniline blue

staining co-localises with immobile mCherry-HIPP43. Scale bars represent 20 μm.

1413

1414

1415

1416

1417

1418

1419

1420

Fig. S11. Pwl2 alters HIPP43 cellular localisation from the plasmodesmata.

(**A**) A time-lapse confocal microscopy micrograph to shows changes of mCherry-HIPP43

localisation upon interaction with Pwl2 after co-expression in *N. benthamiana.* Pwl2 and

HIPP43 co-localise as small or large mobile puncta. White arrow heads indicate mobile

Pwl2/HIPP43 co-localisation puncta and blue arrow heads show immobile aniline blue stained

PD. Arrow 1 indicate large cytoplasmic mobile body, arrow 2 show small mobile puncta. (**B-**

E) Shows lack of co-localisation between MEP3 (**B**), AVR-PikE (**C**), Pwl2[SNDEYWY] (**D**) and

(**E**) free-YFP with HIPP43. HIPP43 co-localise with PD aniline blue stain as shown with cyan

arrow heads and remains immobile. Note, MEP3 localises at PD in presence or absence of

HIPP43. Green arrows show expression of AVR-PikE, MEP3 and free-YFP in the cytosol

without HIPP43 co-localisation. Scale bars represent 20 μm. Aniline blue is shown in blue,

free-YFP, Pwl2, MEP3, AVR-PikE and Pwl2[SNDEYWY] are shown as green while mCherry-

HIPP43 is shown as magenta.

1434

List of Movies

Movie S1

Mobility of Pwl2/HIPP43 sub-cellular co-localisation cytosolic puncta, 48 h after co-
expression in *N. benthamiana* cells and visualized using laser confocal microscopy. PD are
visualized as callose stained with aniline blue, Pwl2 as green and HIPP43 magenta fluorescence
respectively. Arrows are used to track Pwl2/HIPP43 co-localised mobile cytosolic puncta.

Movie S2

A laser confocal microscopy time-lapse video to show lack of co-localisation between AVR-
PikE and HIPP43, 48 h after co-expression in *N. benthamiana* cells. PD are visualized as
callose stained with aniline blue, AVR-PikE as green and HIPP43 magenta fluorescence
respectively. Cyan arrows show co-localisation of aniline blue stain and HIPP43 as immobile
signal at PD. Most of AVR-PikE expression occurred in the cytosol without HIPP43 co-
localisation.

Movie S3

A laser confocal microscopy time-lapse video to show lack of co-localisation between MEP3
and HIPP43, 48 h after co-expression in *N. benthamiana* cells. PD are visualized as callose
stained with aniline blue, MEP3 as green and HIPP43 magenta fluorescence respectively. Cyan
arrows show co-localisation of aniline blue stain, MEP3 and HIPP43 as immobile signal at PD.
Most of MEP3 expression occurred in the cytosol without HIPP43 co-localisation.

Movie S4

A laser confocal microscopy time-lapse video to show lack of co-localisation between a
septuple mutant of Pwl2, Pwl2^{SNDEYWY} and HIPP43, 48 h after co-expression in *N.*
benthamiana cells. PD are visualized as callose stained with aniline blue, Pwl2^{SNDEYWY} as
green and HIPP43 magenta fluorescence respectively. Cyan arrows show co-localisation of
aniline blue stain and HIPP43 as immobile signal at PD while expression of Pwl2^{SNDEYWY}
remained in the cytosol without HIPP43 co-localisation

Movie S5

A laser confocal microscopy time-lapse video to show lack of co-localisation between free-
YFP and HIPP43, 48 h after co-expression in *N. benthamiana* cells. laser confocal microscopy.
PD are visualized as callose stained with aniline blue, free-YFP as green and HIPP43 magenta
fluorescence respectively. Cyan arrows show co-localisation of aniline blue stain and HIPP43
as immobile signal at PD while expression of free-YFP remained in the cytosol without HIPP43
co-localisation.

Table S1 Primers used in this study.

Supplemental Data 1. Blast genomes used for phylogenetic analysis.

Supplemental Data 2. Blast genomes used for copy number variation analysis.

Supplemental Data 3. IP-MS spectrum report.

Supplemental Data 4. IP-MS samples report with clusters.

1473

1474

1475

1476   **Table S1 Primers used in this study.**

1477

Primer	Sequence
PWL2ORF	ATGAAATGCAACAACATCATCCTCCC
PWL2ORR	ACATAATATTGCAGCCCTCTTCTCGC
PWL2f	GGTTCTTATTATGGTCCCGGGTGA
PWL2r	GGTTCTTATTATGGTCCCGGGTGA
PWL2848F	CGCGGTGGCGGCCGCTCTAGAGACCGGGCACGAACCCGGCAGGC
PWL2-R	CATAATATTGCAGCCCTCTT
PWL2_GFPF	GGGCTGCAATATTATGATGGTGAGCAAGGGCGAGGAGCTG
KOLFF	ACAACGCGGTGTAAAGATTCGGGT
KOLFR	GTCGTGACTGGGAAAACCCTGGCGAGACCAAAGCAAAGGGAGGATGA
KORFF	TCCTGTGTGAAATTGTTATCCGCTGCGAGAAGAGGGCTGCAATATTA
PG514-myc-F	AACACGGGGGACTCTAGAGAACAAAAGTTGATCTCTGA
YFPPWL2-F	GACGAGCTGTACAAGGGTGGCGGGTGGACTAAC
PWL2-N-term-YFPR	GTTAGTCCACCCGCCACCCTTGTACAGCTCGTC
PWL2-R-p514	GAACGATCGGGGAAATTCGTTACATAATATTGCAGCCCT
HAPwl2F	AGAGAACACGGGGGACTCTAGAATGGGTGGCGGGTGGACTAAC
PG514-myc-F	AACACGGGGGACTCTAGAGAACAAAAGTTGATCTCTGA
myc-HIPP43-R	TCCAAGACGCCCATAAGGTCCTCTTCAGAAA
myc-HIPP43F	ATTTCTGAAGAGGACCTTATGGGCGTCTTGGA
HIPP43-NOS-R	ACGATCGGGGAAATTCGTCACATGATGGTGCAG
PG514-mCherry-F	AACACGGGGGACTCTAGAATGGTGAGCAAGGGC
mCherry-HIPP43F	GGACGAGCTGTACAAGATGGGCGTCTTGGA
mCherry-HIPP43R	TCCAAGACGCCCATCTTGTACAGCTCGTCC
KT7_HIPP34F	CATGGAGGCCGAATTCATGGGCGTCTTGGA
KT7_HIPP34R	GCAGGTCGACGGATCCTCACATGATGGTGCAG

1478