1   **The blast effector Pwl2 is a virulence factor that modifies the cellular localisation of host protein**

2   **HIPP43 to suppress immunity.**

Vincent Were[1], Xia Yan[1], Andrew J. Foster[1], Jan Sklenar[1], Thorsten Langner[1], Adam Bentham[1], Rafał Zdrzałek[2], Lauren Ryder[1], Davies Kaimenyi[1], Diana Gomez De La Cruz[1], Amber Gentle[1], Yohan Petit-Houdenot[4], Alice Bisola Eseola[1], Matthew Smoker[1], Mark Jave Bautista[1], Weibin Ma[1], Jiorgos Kourelis[1], Dan Maclean[1], Mark J. Banfield[2], Sophien Kamoun[1], Frank L.H. Menke[1], Matthew J. Moscou[1,3] and Nicholas J. Talbot[1] *

[1] The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.
[2] Department of Biochemistry and Metabolism, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.
[3] USDA-ARS Cereal Disease Laboratory, University of Minnesota, 1551 Lindig Street, Saint Paul, Minnesota, 55108 USA.
[4] Université Paris-Saclay, INRAE, UR BIOGER, 91120, Palaiseau, France.

*Corresponding author: Nicholas J. Talbot (Nick.Talbot@tsl.ac.uk)

ORCID IDs

Vincent Were:	0000-0002-9885-4877
Xia Yan:	0000-0003-4509-0137
Andrew J. Foster:	0000-0002-8459-7590
Jan Sklenar:	0000-0003-1858-2574
Thorsten Langner:	0000-0002-3401-8888
Adam R. Bentham:	0000-0001-5906-0962
Rafał Zdrzałek:	0000-0003-3669-924X
Lauren Ryder:	0000-0003-0370-5746
Davies Kaimenyi:	0000-0002-6320-3740
Diana Gómez De La Cruz:	0000-0001-6683-9245
Amber Gentle:	0009-0006-1944-4754
Yohann Petit-Houdenot:	0000-0003-4806-8349
Alice Bisola Eseola:	0000-0001-5698-8204
Mark Jave Bautista:	0000-0002-6016-5953
Weibin Ma:	0000-0002-0516-2151
Jiorgos Kourelis:	0000-0002-9007-1333
Dan MacLean:	0000-0003-1032-0887
Mark J. Banfield:	0000-0001-8921-3835
Sophien Kamoun:	0000-0002-0290-0315
Frank L.H.Menke:	0000-0003-2490-4824
Matthew J. Moscou:	0000-0003-2098-6818
Nicholas J Talbot:	0000-0001-6434-7757

Abstract

The rice blast fungus *Magnaporthe oryzae* secretes a battery of effector proteins during host
infection to suppress plant immunity and promote disease. Among these effectors, the MAX
effector-Pwl2 was first identified as a host specificity determinant for infection of weeping
lovegrass (*Eragrostis curvula*). However, its biological activity has remained unknown. Here
we show that the *PWL2* gene is ubiquitous in all host-limited forms of *M. oryzae* and has
undergone substantial copy number expansion indicating that it is a core effector of the blast
fungus. We used CRISPR/Cas9-mediated gene editing to delete all three copies of *PWL2* and
generate a *pwl2* null mutant in *M. oryzae* strain Guy11. This resulted in gain-of-virulence
towards weeping lovegrass, but a reduction in the severity of disease symptoms on rice. To
further investigate the virulence mechanism of Pwl2, we showed that transgenic rice and barley
lines constitutively expressing *PWL2* display suppression of reactive oxygen species
generation and increased susceptibility to infection by *M. oryzae*. Also, we identify the barley
and rice heavy metal-binding isoprenylated protein HIPP43 as a target of the Pwl2 effector.
Transgenic lines overexpressing HIPP43 exhibit attenuated defense responses and increased
susceptibility to blast infection, consistent with the hypothesis that Pwl2 binds HIPP43 to
modulate immunity. This binding is coupled to changes in cellular localisation of these
proteins. Taken together, our results provide evidence that Pwl2 is a virulence factor in *M.
oryzae* that acts by suppressing host immunity through binding and re-localising HIPP43.

Introduction

Plant pathogens secrete virulence proteins called effectors during infection to suppress the plant
immune system and facilitate invasion of host tissue. This process, called effector-triggered
susceptibility, enables pathogens to manipulate plant metabolic processes, overcome plant
immunity and successfully colonise host plants (1). Fungal pathogens, such as the devastating
blast fungus *Magnaporthe oryzae* (which destroys enough rice to feed 60 million people every
year) deploy an extensive battery of effectors, with more than 500 identified to date (2, 3), but
very few are functionally characterised (4). However, effectors are also recognised by plant
intracellular nucleotide-binding leucine rich repeat (NLR) immune receptors to activate disease
resistance. For example, in rice, the paired NLRs RGA5/RGA4, Pik-1/Pik-2, and Piks-1/Piks-
2 confer resistance to *M. oryzae* strains that secrete AVR1-CO39/AVR-Pia, AVR-Pik or AVR-
Mgk1 effectors, respectively (5-9). Each of these proteins are members of the *Magnaporthe*
Avrs and ToxB-like (MAX) effector family that are sequence unrelated, but structurally
conserved (10), and are overrepresented among effectors recognised by rice NLR immune
receptors (6, 7, 11, 12). Pwl2 was first identified as a host specificity determinant that controls
pathogenicity towards weeping lovegrass (*Eragrostis curvula*), a widely grown forage grass
and was recently shown to be recognised in barley by the Mla3 immune receptor (13, 14) While
PWL2 has been widely studied and used as an important tool to understand effector secretion
and delivery, the molecular mechanisms by which Pwl2 interacts with the host have remained
unknown, with little research into the virulence function of this effector.

A companion paper Zdrzałek *et al* (15) identifies Pwl2 as a MAX effector, and recent studies
of MAX-effector perception by NLRs have identified host small heavy metal-associated
(sHMA) domain containing proteins to be the likely targets of several of these effectors. Small
HMA proteins are highly expanded across plant species with putative functions ranging from
heavy metal detoxification, to potential metallochaperones and other yet to be characterised
functions (16-18). These proteins can be broadly separated into two families, heavy metal-
associated plant proteins (HPPs), and heavy-metal associated isoprenylated plant proteins
(HIPPs), of which the latter possess a C-terminal isoprenylation motif (CaaX, where 'a'
represents an aliphatic residue and 'X' is any amino acid) important for anchoring to

membranes (19). The rice sHMA protein Pi21 is a disease susceptibility factor, with rice lines
expressing this protein exhibiting delayed defence responses to rice blast (19, 20). This has led
to the use of loss-of-function alleles of *pi21*, as a recessive form of rice blast resistance in
breeding programs (20). Furthermore, differences in host sHMA protein repertoires have been
linked to host-specific adaptive evolution of the APikL2 effector family, while studies of AVR-
Pik and its target OsHIPP20 suggests that effector binding may stabilize the host protein (21,
22). Importantly, sHMA protein domains have also been found integrated as non-canonical
immune sensory domains in some NLR proteins, such as the rice Pik-1/2 and Pia (RGA5/4)
receptors, and are essential for recognition of their cognate *M. oryzae* effectors (6, 8, 9, 11).
The integration of sHMA proteins into immune receptors further underlines the relevance of
sHMA proteins in pathogen virulence. However, the exact function of sHMA proteins in plants
and their link to disease susceptibility in plant-pathogen interactions is still poorly understood,
obscuring the specific function of MAX effectors that target these proteins.
In this study, we investigate the biological function of *PWL2* and its role in fungal pathogenesis.
We were motivated to understand how a ubiquitous effector like Pwl2 functions during a
susceptible interaction between *M. oryzae* and its host. We demonstrate that the *PWL2* gene is
ubiquitous across *M. oryzae* lineages and has undergone significant gene duplication and
expansion in copy number, with many rice blast isolates possessing 3-5 copies, making its
functional analysis challenging. Using CRISPR-Cas9 gene editing, we generated a *pwl2* null
mutant and verified the role of Pwl2 as a host specificity determinant towards weeping
lovegrass and barley lines containing *Mla3*. We constructed transgenic rice and barley lines
constitutively expressing *PWL2* and show that Pwl2 suppresses PAMP-triggered immunity
through attenuation of the ROS burst, resulting in increased susceptibility to rice blast. Finally,
we reveal that Pwl2 targets the sHMA protein HIPP43 in both rice and barley and show that
transgenic plants overexpressing HIPP43 have reduced immune responses. We also show that
Pwl2 co-localizes and alters sub-cellular localisation of HIPP43 in *Nicotiana benthamiana*.
Taken together, our study provides evidence that Pwl2 plays a role in defence suppression by
binding and re-localising HIPP43 to facilitate colonization of plant tissue by the blast fungus.

Results

Pwl2 is an *in-planta* effector, expressed during infection and localizes to the BIC.

The *PWL2* gene is specifically expressed during the biotrophic phase of infection, peaking in
expression at 36 - 48 hours post-infection (hpi) (3) (**Fig. S1*A* and *B***). The Pwl2 effector
localises in the biotrophic interfacial complex (BIC) and is clearly visible as a single bright
punctum, initially at the tip of an invasive hypha and then adjacent to the bulbous, branched
hyphae as they develop (**Fig. 1*A* and *B***). The BIC is a plant membrane-rich structure and the
predicted site of effector delivery (23-25), although experiments to date have not precluded
that the BIC could be a site of effector sequestration by the host. To investigate Pwl2 secretion
at the BIC during infection, we generated a single *M. oryzae* strain expressing two BIC-
localised effectors Pwl2-mRFP and Bas1-GFP and visualised their localisation during infection
of leaf sheath tissue of the susceptible rice cultivar Moukoto. The two effectors were observed
as small punctate signals within the same BIC (**Fig. 1*A* and *C***). By contrast, when two different
M. oryzae strains, Ina168 and Guy11, expressing Pwl2-GFP and Pwl2-mRFP respectively were
used to simultaneously infect rice tissue, we observed that the BIC formed by each individual
invasive hypha harboured exclusively either Pwl2-GFP or Pwl2-mRFP but never both signals
(**Fig. 1*B* and *D***). This suggests that the Pwl2 effector is secreted from the BIC into host cells,
especially when considered with previous reports showing the BIC to be the site of
unconventional fungal secretion and clathrin-mediated endocytosis (24).

PWL2* is ubiquitous in isolates of *M. oryzae.

The *PWL2* effector belongs to a wider gene family (13, 26), but conservation of the *PWL* family
in the global rice blast population, and among different host-limited forms of *M. oryzae*, is not
fully known. Therefore, we investigated *PWL* gene family conservation in publicly available
genome sequences, including isolates that infect *Oryza sativa* (rice), *Eleusine* spp. (finger
millet), *Hordeum vulgare* (barley), *Setaria* spp. (foxtail millet), *Triticum aestivum* (wheat),
Lolium spp. (rye grass), *Brachiaria* spp. (signal grass), *Panicum repens* (torpedo grass) and
Eragrostis spp (weeping lovegrass) (27-30). We also included two close relatives of *M. oryzae*
that infect crabgrass (*Digitaria sanguinalis*) and pearl millet (*Pennisetum* spp.). We used Pwl1
(BAH22184.1), Pwl2 (QNS36448.1), Pwl3 (AAA80240.1) and Pwl4 (AAA80241.1) protein
sequences to query for the presence or absence of each protein-encoding gene. We observed
that *PWL1* is present in group EC-1I (31) of *Eleusine*-infecting isolates, one *Eragrostis*-
infecting isolate and some *Oryza*-infecting isolates but absent in all the other host-specific
lineages (**Fig. 1*F* and S2*B***). Similarly, *PWL3* is found in most *Oryza*-infecting isolates, but
almost completely lost in the other lineages, except for some *Setaria*, *Lolium*, and *Eleusine*-
infecting isolates. *PWL4* is present in *Eleusine*, *Lolium* and *Triticum*-infecting isolates, but
almost completely lost in *Oryza*-infecting isolates. We however found one *Digitaria*-infecting
isolate that carry *PWL4* (**Fig. S2*B***). In contrast to all other *PWL* family members, *PWL2* is
found in all host-limited forms of *M. oryzae* and related *Magnaporthe* species (**Fig. 1*E* and
S2*B***). We conclude that Pwl2 is a core effector of the blast fungus.

119
Having determined Pwl2 is ubiquitous in isolates of *M. oryzae*, we next sought to determine
how polymorphic the effector is, in different host-limited forms of *M. oryzae*. Apart from the
pwl2-2 variant that contains a single Asp-90-Asn substitution (13, 14), we found 14 new
variants of Pwl2 that can be renamed Pwl2-3 to Pwl2-14. Majority of the polymorphic residues
occurred at positions His87 to Ser92 (**Fig. S2*A***). In comparison, we identified 2 variants of
Pwl1 (renamed Pwl1-2 and Pwl1-3), 4 variants of Pwl3 and 1 variant of Pwl4 (**Fig. S3*A*, *B*
and *C***). From this analysis, we conclude that Pwl2 is highly polymorphic compared to other
members in the family. We next tested the ability of *PWL2* variant alleles to be recognised on

known resistant hosts. A subset of the isolates carrying variants of *PWL2* BTJP4-16 (*pwl2-3)*,
JUM1 (*pwl2-2*), BN0293 (variant *Pwl2*), BF5 (variant *Pwl2*), VO107 (variant *Pwl2*), and PM1
(variant *Pwl2*) were used to infect barley lines that carry *Mla3* or weeping lovegrass. Some of
the variants were identified in same background isolate carrying *PWL1,* one or multiple copies
of *PWL2*, and it was not possible functionally characterise them using infection assays. We
found BTJP4-16 (*pwl2-3)*, JUM1 (*pwl2-2*) and BN0293 to be virulent on these hosts, providing
evidence that these alleles are non-functional (**Fig. S2*C-D***). Isolates VO107, BF5 and PM1 did
not produce blast disease lesions in the presence of *Mla3* suggesting that alleles present in these
isolates were either recognised by the host, or the *M. oryzae* isolates carry other effectors
recognised by additional R-genes contained in barley cv. Baronesse or weeping lovegrass. (**Fig.**
S2*E*).

Pwl2 allelic variability affecting host specificity is not associated with the MAX effector

fold.

The majority of *Oryza-*, *Lolium*- and *Triticum*-infecting isolates of *M. oryzae* analysed have
PWL2 or the variant *pwl2-2* allele, which is associated with gain of virulence to *E. curvula* and
Mla3 (**Fig. S1*C***) (13, 14). Our analyses of the presence of *PWL2* in *Lolium* and *Triticum*-
infecting isolates from the pandemic wheat blast lineage (32) identified an allele of *PWL2,*
designated *pwl2-3*, with several polymorphisms in the core effector fold: Glu-89-Gln, Lys-91-
Gln and Ser-92-Ile (**Fig. 1*E*, S1*D* and S2*A***). Interestingly, *pwl2-3* was also identified from
Digitaria and *Pennisetum spp*-infecting isolates of *Magnaporthe* (**Fig. 1*F*, S2*A* and *B***). In an
accompanying paper by Zdrzalek *et al* (15) Pwl2 has been structurally verified as a predicted
MAX effector, with a core β-sandwich fold formed of two antiparallel β-sheets, as well as a
single α-helix, C-terminal to the β-sandwich fold. We therefore mapped a subset of the
identified polymorphic residues of Pwl2 including *pwl2-3* to the effector crystal structure. All
polymorphisms between *pwl2* alleles were found to reside in the C-terminal α-helix, a distinct
region from the core MAX-fold (**Fig. 1*E***). This suggests that the interface used to bind Pwl2
by the immune receptors present in resistant barley (Mla3) and weeping love grass, unlikely
involves the MAX-fold.

***PWL2* occurs in high copy number in *M. oryzae* field isolates.**

The *M. oryzae* reference genome assembly (strain 70-15) has two copies of *PWL2* on
chromosomes 3 and 6, annotated as MGG_04301 and MGG_13683, respectively. We
observed, however, that *PWL2* copy number varies significantly in different *M. oryzae* isolates.
Southern blot hybridization (**Fig. S4**), and analysis of long-read assembled genomes of the *M.*
oryzae rice pathogenic strain Guy11 identified three copies of *PWL2*. A duplication of *PWL2*
is present on a single contig in the Guy11 genome, whilst the third copy involves a translocation
resulting in a distinct *PWL2* allele in a separate contig (**Fig. 2*A***). We observed that the *PWL2*
loci were associated with the occurrence of *MGR583*, *POT2* and *MAGGY* transposable
elements, suggesting a possible involvement in genome translocation. To further assess *PWL2*
copy number in *M. oryzae* isolates, we employed a k-mer analysis approach to determine copy
number variation of *PWL2* in *M. oryzae* genomes (n = 286). Strikingly, *PWL2* copy number
was high in many *M. oryzae* strains, with one isolate showing 8 copies (**Fig. 2*B***). We conclude
that *PWL2* is ubiquitous in *Magnaporthe* and has expanded in copy number both in *M. oryzae*
and its sister species *M. grisea*.

Δpwl2 **mutants show altered host range and are reduced in ability to cause rice blast**

disease.

Having determined the copy number variation of *PWL2* in *M. oryzae*, we used a CRISPR/Cas9
genome editing approach to simultaneously delete the three copies of *PWL2* found in Guy11
(33). We designed a sgRNA to target *PWL2* and introduced a hygromycin phosphotransferase-
encoding gene cassette (*HPH*) (**Fig. 3*A***). To identify *Δpwl2* null mutants, transformants were
analysed by PCR using primers designed to amplify the *PWL2* coding sequence. Transformants
with ectopic integrations or where not all copies of *PWL2* were deleted, gave a predicted
amplicon size of 645 bp, consistent with untransformed Guy11. By contrast, in null mutants,
we amplified a larger amplicon of 1.5 kb, consistent with introgression of the hygromycin
resistance cassette into all three copies of *PWL2* (**Fig. 3*A***). Four putative transformants were
then selected for whole genome sequencing. Raw reads were aligned to the *M. oryzae* reference
genome using samtools v.15, before converting into BAM files and visualised using IGV
viewer. No reads mapping to the *PWL2* locus were identified in mutants T6 and T12 whereas
fewer reads mapped to the same locus from sequenced T5 and T4 (**Fig. S5**). Regions upstream
and downstream of the *PWL2* coding sequence were 100 % identical in both mutants and
reference genomes. This confirmed that T6 and T12 are *Δpwl2* gene deletion transformants and
they were therefore selected for further analysis. We reasoned that *M. oryzae Δpwl2* deletion
mutants would gain virulence on weeping lovegrass (13, 26) and the barley cultivar Baronesse
carrying *Mla3* cultivars of barley. The *Δpwl2* deletion transformants T5, T6 and T12 and
Guy11 were therefore used to inoculate weeping lovegrass seedlings. We observed that the
Δpwl2 mutants T6, T5 and T12 showed gain of virulence and symptoms of infection which
were identical to those exhibited by a virulent G17 isolate that lacks *PWL2* (**Fig. 3*B***). We
conclude that T5, T6 and T12 are loss of recognition mutants of *PWL2*, and we can be confident
that T6 and T12 have complete deletion of all three copies of the gene.
We next tested whether Pwl2 has a virulence function during a compatible interaction between
the blast fungus and its host. The *Δpwl2* mutant T6 was infected on the susceptible rice cultivar
CO39, susceptible barley lines Nigrate, Siri (*Mla8*), Golden Promise (*Mla8*), and resistant
cultivar Baronesse (*Mla3*). The *Δpwl2* mutants showed a small but reproducible reduction in
virulence on CO39 compared to wildtype Guy11, based on observed disease lesion size (**Fig.
3*C* and *E***). Additionally, *Δpwl2* produced statistically smaller lesion sizes on barley lines Siri
and Golden Promise (**Fig. 3*D* and *F***). We conclude that *PWL2* contributes to the ability of *M.
oryzae* to cause blast disease.

Pwl2 suppresses host immunity.

Having determined that the Pwl2 effector contributes to virulence, we aimed to gain insight
into its biological function during host cell colonisation. We generated stable transgenic barley
(Golden Promise) and rice (Kitaake) lines expressing *PWL2-YFP* (without its signal peptide)
under control of the CaMV35S promoter. Transgenic lines expressing cytoplasmic YFP were
used as controls. Transgenic plants, expressing Pwl2 and free YFP did not show any adverse
growth or fitness phenotypes (**Fig. 4*A***). We first assayed *PWL2-YFP* transgenics for their
response to two elicitors of PAMP-triggered immunity, flg22 and chitin. Rice and barley
perceive flg22 through the pattern recognition receptor FLS2 and perceive chitin through
CEBiP and CERK1 leading to immune response for example production of reactive oxygen
species (ROS) (34, 35). We observed that Pwl2 expression abolished chitin and flg22-induced
ROS generation (**Fig. 4*B* and *C***). Pwl2 therefore contributes to virulence by suppressing
PAMP-triggered immunity during compatible interactions. To investigate the effect of elevated
Pwl2 effector expression on blast infections, we infected barley plants expressing Pwl2 with

M. oryzae Guy11. Pwl2-expressing lines were significantly more susceptible and developed
blast disease symptoms earlier (2-3 dpi) compared to infection of isogenic Golden Promise
plants (**Fig. 4D and E**). We conclude that Pwl2 is a modulator of plant immunity.

Pwl2 binds to the heavy metal binding domain isoprenylated protein, HIPP43.

To determine the likely target of Pwl2, transgenic barley and rice plants expressing Pwl2-YFP
were used for protein extraction before performing anti-YFP co-immunoprecipitation coupled
to mass spectrometry (IP-MS). This revealed a total of 791 proteins after spectral search in the
H. vulgare (barley, cv Morex version 3) proteome database. We identified a total of 282
frequently occurring proteins in the biological replicates containing Pwl2, that were not
identified in the six free-YFP control samples. To avoid focusing on false positives in the form
of sticky and abundant proteins, fold-changes were calculated by first determining the average
number of peptides per protein candidate, before estimating $\log_2$ fold change compared to the
control experiment. We further filtered for proteins that produced at least two peptide hits in
more than half of replicates ($\geq$4/8 biological replicates). A total of 52 proteins met this criterion
and were considered as enriched in samples compared to controls and were therefore selected
for further analysis. Furthermore, because Pwl2 attenuates the ROS-burst in barley transgenics
lines, we focused initially on protein candidates previously reported to have roles in plant
immunity or ROS generation (**Fig. 4F**). These were selected for one-to-one interactions using
a yeast-two-hybrid (Y2H) assay by co-transforming constructs expressing Pwl2 and <u>P</u>utative
<u>P</u>wl2 <u>I</u>nteracting <u>P</u>roteins (PPIPs) identified by IP-MS (**Fig. 4F and Fig. S6**). Pwl2 showed
strong binding to PPIP4 (**Fig. 4G and H**). We named this protein HvHIPP43, for *Hordeum*
vulgare heavy metal binding domain containing isoprenylated plant protein 43 based on
homology to *Os*HIPP43 defined in a companion paper by Zdrzalek *et al* (15). To validate the
interaction between Pwl2 and HIPP43, we carried out co-immunoprecipitation (co-IP) analysis
in *N. benthamiana* and could detect HvHIPP43 interacting with Pwl2-YFP (**Fig. 4H**). Having
determined that Pwl2 binds to HvHIPP43, we investigated the *PWL* gene family-encoded
proteins Pwl1, Pwl3, Pwl4 and pwl2-3 variant and showed that they were all able to bind
HvHIPP43 in Y2H assays, confirming that Pwl2 belongs to an effector family that binds to a
host sHMA protein (**Fig. 4I**).

Pwl2 interacts with HIPP43 orthologs from other grass species.

HIPPs are expanded in plant genomes and to test whether HIPP43 has orthologs in other grass
species, we generated a maximum likelihood phylogenetic tree of HMA domains from diverse
grasses with high quality genomes and annotations. We found that the HIPP43 family formed
a distinct clade which included orthologs from rice (*O. sativa)*, wheat (*T. aestivum*)*,* wild wheat
(*Haynalidia villosa*), foxtail millet (*S. italica),* weeping lovegrass (*E. curvula),* Sorghum
(*Sorghum bicolor*)*, Oropetium thomaeum*, *Zea mays* and *Brachypodium distachyon* (**Figure**
S7 and S8). Furthermore, we could identify copy number variation ranging from one to five
copies in grasses, with barley and wheat (i.e. Triticeae lineage) having three paralogs of
HIPP43, one per haploid genome. We followed this by confirming that, Pwl2 interacts with
HvHIPP43 orthologs from *Triticum* (*Taes*HIPP43), *Setaria* (*Siti*HIPP43), and *Eragrostis*
(*Ec*HIPP43), in Y2H assays (**Fig. 4J**). Binding of Pwl2 to rice *Os*HIPP43 was verified by in
vitro biochemistry and a structure of the complex obtained from *X*-ray crystallography (see
companion paper by Zdrzalek *et al* (15). We also analyzed the transcriptional profile of
*Os*HIPP43 during infection of Guy11 on a susceptible rice line CO39 from dataset available
from Yan *et al* (3). Interesting we found that, *Os*HIPP43 expression was up regulated during
plant infection, suggesting it might play an important role in plant immune response.

Overexpression of HIPP43 suppresses PAMP-induced ROS in transgenic barley.

To investigate the biological relevance of the interaction between Pwl2 and HIPP43, we generated barley transgenic lines overexpressing HIPP43. These transgenic lines were challenged with two elicitors of PAMP-triggered immunity, flg22 and chitin. Strikingly, we observed that the chitin and flg22-induced ROS burst was abolished in plants expressing HIPP43 compared to wild type and plants expressing free YFP (**Fig. 5** *A, B, C* and *D*). Similarly, when we challenged transgenic barley plants expressing HIPP43 with *M. oryzae*, we observed increased susceptibility compared to the wildtype Golden Promise with disease lesions appearing earlier than in wild type plants. This observation is consistent with earlier results showing attenuated ROS burst in plants expressing Pwl2-YFP (**Fig. 5***E* and *F*).

Isoprenylation motif is required for HIPP43 plasmodesmata localization.

So far, there is no direct evidence for the involvement of HIPP proteins in host immunity and regulation of ROS so we reasoned that HIPP43 might bind to another host protein following interaction with Pwl2. HIPP43 possesses an isoprenylation motif at its C-terminus (**Figure S9***A*), and we performed live cell imaging of mCherry and YFP-tagged HIPP43 (at the N or C terminus, respectively), following transient expression in *N. benthamiana* to determine the localisation of the protein. We also tested localisation of HIPP43 without its predicted isoprenylation motif. When tagged at the N-terminus, HIPP43 mainly localises to puncta at the plasma membrane that co-localise with callose staining (aniline blue) consistent with these being sites of plasmodesmata (PD) (**Fig. 5***G* and *H*). However, without its isoprenylation motif, PD localisation was abolished, and we observed accumulation in the nucleoplasm and plasma membrane (**Fig. S9***B*). When tagged at its C-terminus (HIPP43-YFP), HIPP43 mainly localises to PD, cytoplasm, plasma membrane and chloroplasts (**Fig. S9***C*).

Pwl2 stabilises and alters plasmodesmata localization of HIPP43.

When we co-expressed YFP-Pwl2 with mCherry-HIPP43, we observed co-localisation in the cytoplasm with the appearance of cytoplasmic mobile bodies approximately 2 μm in diameter, likely to be very large vesicles or P-bodies (**Fig. 5***I*). In some cases, we also observed larger cytoplasmic mobile bodies ranging from 2-4 μm in diameter (**Fig. 5***I*). We quantified the number of times HIPP43 signal was observed at PD when co-expressed with free-YFP, AVR-PikE and the effector Mep3 (3) as controls. We observed that co-localisation of YFP-Pwl2 and mCherry-HIPP43 as cytoplasmic vesicles compromised PD localisation (**Fig. 5***I* and *J*).. In addition, localization of HIPP43 in large cytoplasmic mobile bodies was not observed when we co-localised HIPP43 with empty vector controls expressing YFP (**Fig. 5***K* and *L*) or two other MAX-fold effectors, AVR-PikE (**Fig. 5***M* and *N*) and MEP3 (**Fig. S10***E* and *F*), as it remained localised at PDs. Moreover, the intensity of mCherry-HIPP43 fluorescence was higher when co-expressed with Pwl2 compared to when expressed in the absence of Pwl2, or when co-expressed with free YFP, AVR-PikE or MEP3 (**Fig. 5***I* and **Fig. S10***D*). When considered together, these results suggest that Pwl2 binds to HIPP43, potentially stabilising the protein and altering its cellular location away from plasmodesmata.

Discussion

The Pwl2 effector was first identified as a host specificity determinant for infection of the forage grass species weeping lovegrass (13, 26). Its initial identification provided evidence that host range in plant pathogenic fungi was conditioned in a similar way to cultivar specificity in a gene-for-gene manner, involving dominant pathogen genes recognised by the products of cognate disease resistance genes (26). Furthermore, Pwl2 was found to belong to an expanded gene family, suggesting an important function in pathogenicity and fitness (26). However, in the following two decades, the function of Pwl2 has remained elusive despite its extensive use as a marker in cell biological studies for investigating effector regulation, secretion and delivery during plant infection (23-25).

In this study, we set out to explore the function of Pwl2 and investigate why it is such a ubiquitous effector in *Magnaporthe oryzae*. We found that the *PWL2* gene is present across all rice blast lineages, and in all publicly available genome sequences of *M. oryzae*, including each of its host-limited forms and even in sister species infecting crabgrass and pearl millet. The observation that *PWL2* and the wider *PWL* family genes have been maintained in the global blast population and occurrence of effector variants or loss-of-recognition alleles associated with certain host-adapted lineages, suggests that Pwl2 and members of this family serve an important function in pathogenesis. Furthermore, *PWL2* is present at high copy number in many *M. oryzae* isolates having undergone extensive gene duplication. This may have been facilitated by the presence of *MGR583*, *POT2* and *MAGGY* transposable elements (36, 37) flanking *PWL2* loci, providing the capacity to generate translocations and genomic instability. It is not clear whether the signatures of lineage-specific diversification are due to isolates infecting rice varieties carrying a yet unidentified resistance gene against Pwl2, or resistance genes that have evolved in different host species such as barley cultivars expressing Mla3 or *E. curvula* (14).

We were able to apply CRISPR/Cas9 gene editing in *M. oryzae* to functionally characterise genes occurring in multiple copies through the generation of *Δpwl2* mutants. Our analysis confirmed the previously reported host specificity function of Pwl2, but also revealed it is important for blast disease, with *Δpwl2* mutants showing reduced virulence compared to the isogenic wild type strain. Pwl2 therefore plays an important role during host tissue invasion, consistent with the observation of smaller and fewer disease lesions in susceptible interactions with *Δpwl2* mutants.

Using discovery proteomics, we identified putative Pwl2-interacting proteins, including an isoprenylated small HMA protein, HIPP43, which contains a predicted C-terminal isoprenylation motif. We found that all proteins encoded by the *PWL* gene family can bind HIPP43, including HIPP43 homologs from wheat, rice, weeping lovegrass and foxtail millet. Multiple MAX effectors bind sHMA protein domains, although most previous studies have focused on the incorporation of such domains into paired NLR immune receptors, leading to disease resistance (7, 22, 38, 39). HMA domains are known to be involved in biotic/abiotic stress responses, transport of metals and metal detoxification, and their expression can be organ-specific or tissue-specific within roots, leaves, or stems (40-42). Moreover, sHMAs can localise to the nucleus, plasma membrane, cytoplasm, or plasmodesmata (41, 43, 44), suggesting they might be involved in numerous cellular processes. Consistent with this, sHMAs are expanded in plant genomes with more than 45, and more than 50 sHMA domain-encoding genes occuring in *Arabidopsis* and rice respectively (45). However, little is known about the function of sHMA proteins in plants, especially during plant/microbe interactions, even though they have been reported to act as susceptibility factors (20, 21). For example, deletion mutants of OsHIPP05 (*Pi21*), a proline rich HMA domain protein-encoding gene in rice leads to rice blast resistance (20). Consequently, *pi21* is a recessive resistance gene that

has been successfully deployed to control blast disease (20). Similarly, gene silencing or
deletion mutants of *TaHIPP1* or *AtHMAD1* in wheat and *Arabidopsis* provide enhanced
resistance against *Puccinia striiformis* f. sp. *tritici* and *Pseudomonas syringae* DC3000,
respectively (46, 47). In addition, deletion mutants of *AtHIPP27* in *Arabidopsis* lead to
increased resistance against cyst nematode (48). However, it is not clear why deletion of sHMA
protein-encoding genes impacts immunity. Therefore, the observation that Pwl2 interacts with
HIPP43 is revealing, especially as Pwl2 binding appears to enhance its activity, or stability,
rather than inhibiting its function. This is consistent with the observation that over-expression
of HIPP43 enhances blast susceptibility in rice and barley, respectively.

The crystal structure of the Pwl2/HIPP43 complex (presented in the companion paper by
Zdrzałek *et al.* (15)) demonstrates that Pwl2 uses an expansive interface to mediate binding to
HIPP43, largely using elements of the MAX fold to interact with the β-sheet of the host target.
Interestingly, the polymorphic residues found between Pwl2 variants, Pwl2, pwl2-2 and pwl2-
3 are not located at the HIPP43 binding interface, but rather located away from the MAX fold
on the C-terminal helix of Pwl2. As these polymorphisms are determinants for Pwl2
recognition in resistance plants carrying functional cognate resistance gene, it suggests the
interface used for immune recognition of Pwl2 in both barley Mla3 lines and resistant weeping
love grass are likely spatially distinct from the interface used by Pwl2 to bind to HMA host
targets.

This observation contrasts with an emerging model where host immune receptors can utilize
binding interfaces spatially similar as to those found between effectors and their host targets
(38, 39). Indeed, Zdrzałek *et al.* (15) show that if HIPP43 is incorporated into the Pik-1 NLR,
in place of its naturally occurring integrated HMA domain, there is potential to gain immune
responses to Pwl2. Therefore, we hypothesize Pwl2 may have additional host targets with
spatially distinct interfaces from that observed with HIPP43. If true, this may explain
differences in Pwl2 HIPP43-binding and immune recognition sites. This suggests immune
recognition may have evolved to recognize Pwl2 through interfaces shared with the other
undetermined host targets. However, to test this hypothesis, structural information for the
natural recognition of Pwl2 by host resistance genes, such as Mla3, or genetic evidence for
additional Pwl2 host targets is required.

Finally, we show Pwl2 interaction with HIPP43 is important for fungal virulence. A *Δpwl2*
null mutant, lacking all three copies of the gene, produced smaller disease lesions that were
also fewer in number than caused by the isogenic wild type *M. oryzae* strain Guy11. Moreover,
when Pwl2 is over-expressed in transgenic rice or barley, it abolished chitin or flg22-elicited
generation of ROS. We also found that over-expression of HIPP43 in rice and barley had a
similar effect, suggesting that Pwl2-HIPP43 interaction results in attenuated generation or ROS
and reduced immune responses. Consistent with these observations, over-expression of Pwl2
or HIPP43 leads to increased susceptibility to *M. oryzae*. How this potential enhancement of
HIPP43 activity is stimulated by Pwl2 is not completely clear, but we found that transient co-
expression of both Pwl2 and HIPP43 sequestered HIPP43 away from plasmodesmata, with the
two proteins co-localizing in the cytoplasm and in large mobile vesicular bodies. The sub-
cellular localization of Pwl2 and HIPP43 was distinct when separately expressed, with Pwl2
accumulating in the cytoplasm and mobile bodies while HIPP43 localized to the membrane,
and particularly to plasmodesmata. Deletion of the isoprenylation motif abolishes PD
localization showing the C-terminal isoprenylation motif is important for plasmodesmatal (PD)
localization. This suggests that disrupting PD localization may lead to HIPP43 translocation to
other organelles, which may in-turn influence immune signaling.

Localization of HIPPs at PD has been previously reported. HIPP7 and HIPP26 localise at PD,
and this localisation can be altered by mutations in the isoprenylation motif (44, 49). Although
characterised HIPPs through their HMA domains are thought to be metallochaperones, there

are no studies that link metal detoxification and immunity. Some studies suggest that there may
be a connection between concentration of metal ions, such as iron, copper, cadmium and zinc
and plasmodesmatal permeability (50). This might explain why a plant pathogen effector
targets a plasmodesmatal localized HIPP. Both the HMA domain and the isoprenylation
regions of HIPPs have been shown to be important in plant immunity. For example, a mutation
targeting the proline-rich region of *pi21* is sufficient to lead to gain of resistance against *M.*
oryzae (20), while interfering with the isoprenylation of HIPP1-V from wild wheat (*H. villosa*)
leads to loss of resistance against *Blumeria graminis* f. sp. *tritici* accompanied by reduced
localization to PD. Interestingly, HIPP1-V interacts with the E3-ligase CMPG1 at the plasma
membrane leading to resistance against powdery mildew in an isoprenylation-dependent
manner (51). This interaction has been reported to activate expression of genes involved in
ROS generation and salicylic acid biosynthesis, suggesting that HIPP1-V is required for PTI
regulation (51) An *A. thaliana* HIPP3 has also been proposed to be involved in regulation of
the pathway involved in salicylic-dependent plant immunity via LSD1, PAD4 and SAG101
(40).
The change in cellular localization of HIPP43 induced by Pwl2 may therefore inhibit its
function in immunity or lead to a new role that enhances disease susceptibility. Future work
will be necessary to see if HIPP43 directly regulates ROS generation or acts in a more indirect
manner through binding other signaling components involved in PTI. By dislocating HIPP43
from the PD, Pwl2 may attenuate a role for HIPP43 in PD immunity. Pwl2 is a highly mobile
effector and has been shown to move into neighbouring rice cells ahead of *M. oryzae* hyphal
growth (23). This has been suggested to be a step to prepare un-invaded cells for fungal
colonisation. Its role may therefore be to alter the sub-cellular localisation, concentration and
stability of HIPP43, enabling more rapid tissue colonisation by the blast fungus.
426

Acknowledgements

We thank current and past members of the "BLASTOFF" team from the Banfield, Kamoun, Terauchi, Talbot, and Moscou Laboratories. For funding, we thank Gatsby Charitable Foundation and Biotechnology and Biological Sciences 462 Research Council (BBSRC) BBS/E/J/000PR9797 and a BBSRC grant to NJT and FLHM 463 (BB/V016342/1 and 101077853 "PANDEMIC"). United States Department of Agriculture-Agricultural Research Service CRIS #5062-21220-025-000D (MJM. We acknowledge Y.K Gupta, J.C De la Concepcion, J. Rhodes and J. Win for constructive discussions.

Author Contributions

Conceptualization: NJT and VW
Formal analysis: VW, XY, AF, JK, JS, DG, TL, YPH, ABE, MS, MJB, WM and RZ
Funding acquisition: NJT
Investigation VW, XY, AF, JS, DG, AG, DK, YPH, ABE, MJB, WM and RZ
Writing – original draft: VW and NJT
Writing – review & editing: VW, NJT, LR, AB, TL, MB, SK, MM, FM

Materials and methods

Fungal strains, growth conditions and infection assays

Fungal isolates were routinely grown on complete medium (CM) at 24°C with a controlled 12 h light and dark cycle for up to 12 days (52). *H. vulgare*, *E. curvula* and *O. sativa* plants were grown for 7, 14 and 21 days respectively in 9 cm diameter plastic plant pots or seed trays. Conidia were recovered from 10-day old cultures using a sterile disposable plastic spreader in 3 mL sterile distilled water. The conidial suspension was filtered through sterile Miracloth and centrifuged at 5000 x g for 5 min at room temperature and adjusted to a final of concentration of 1 x10^5 conidia mL^{-1} in 0.2 % gelatin. The spore suspension was used for spray or leaf drop infections assays. After spray inoculation, the plants were covered in polythene bags and incubated in a controlled plant growth chamber at 24°C for 48 h with a 12 h light and dark cycle, and 85% relative humidity, before removing the polythene bags. The inoculated plants were incubated for 3 more days before scoring the lesions. For each treatment 10 leaves were collected before counting typical disease lesions that are characterised by ellipsoid necrotic lesions with a grey centre, on the leaf surface (53). Each experiment was repeated minimum of three times, yielding consistent outcomes.

Leaf infection assay and live-cell imaging

Rice leaf sheaths from 4 weeks old susceptible cultivars Moukoto or CO39 were inoculated with 4 mL of a suspension at 5 x 10^4 of conidia mL^{-1} in distilled water using a micropipette as described in Kankanala et al 2007 (25). The inoculated leaf sheaths were incubated at 24°C for at least 24 h before a thin layer of the inner leaf sheath was dissected using a blade and mounted on a glass slide. Live cell imaging was carried out on an IX81 motorized inverted microscope (Olympus, Hamburg, Germany) for conventional and differential interference contrast (DIC) microscopy using the Photometrics CoolSNAP HQ2 camera (Roper Scientific, Germany). Images were analyzed using ImageJ. For Leica SP8 laser confocal microscopy, settings were as follows; GFP, YFP and RFP/mCherry tagged proteins were excited using 488, 514 and 561 nm laser diodes and the emitted fluorescence detected using 495-550, 525-565 and 570-620 nm respectively. The auto-fluorescence from chlorophyll was detected at 650-740 nm.

Generation of fungal transformation plasmids

Single or multiple DNA fragments were cloned into fungal transformation vector using In-Fusion HD Cloning Kit (Clontech, USA). Briefly, fragments from cDNA, genomic DNA or synthesized DNA were amplified using primers that introduce a 15bp overhang complementary to sequences at the restriction sites of the destination vector or adjacent insert fragments. This allows the ends to fuse by homologous recombination during cloning. Positive transformation was accessed using colony PCR and constructs sequenced then analysed on SnapGene. A list of primers is provided in Table S3.

RNA isolation, RNA sequencing and analysis

To study *in-planta* gene expression profile of *PWL2* and other effectors, leaf drop infection assays were carried out on a susceptible rice line Moukoto and samples collected at 24 and 72 hpi. Infected plant material was ground into fine powder using a sterile nuclease-free mortar & pestle containing liquid nitrogen. RNA was isolated from *M. oryzae* mycelia inoculated or rice leaves using QIAGEN RNeasy Plant Mini Kit according to manufacturer's instructions. The RNA quality was determined on a NanoDrop spectrophotometer (Thermo Scientific, UK) and Agilent 2100 Bioanalyser using an RNA nano-chip kit (Agilent Technologies, UK). Library preparation was carried out using Illumina® sequencing TruSeq Stranded Total RNA Library

Prep Kit before sequencing of 100 bp paired ends reads using Illumina Genome Analyser GXII
platform by the Exeter Sequencing Service (University of Exeter). *M. oryzae* specific reads
were recovered as unmapped reads after mapping to the *O. sativa* L. ssp. japonica genome
(IRGSP-1.0) (54). To determine differentially genes expression, reads were then mapped to
transcripts from the M. oryzae 70-15 reference genome (55) or a laboratory strain KE002 using
TopHat2 splice site-aware aligner (56). Differential expression analysis was carried out using
DESeq2 packages (57). .

Generation of Cas9-sgRNA targeted gene deletion

A sgRNA was designed for CRISPR-Cas9 genome editing using an online tool E-CRISP
http://www.e-crisp.org/E-CRISP/. A 20-nucleotide sequence was selected at the *PWL2* locus
(not including the PAM NGG-sequence). The sgRNA was first synthesised using the EnGen
sgRNA synthesis kit New England Biolabs (NEB #E3322) before mixing with the Cas9-NLS
to form the RNP complex. The mixture was incubated at room temperature for 10 min before
fungal transformation. *M. oryzae* protoplast were generated as described by Talbot *et al* (52).
The RNP complex together with the donor template was mixed with Guy11 protoplasts re-
suspended in 150 µL of STC to a concentration of 1 X 10^8 and incubated at room temperature
for 25 min before adding 60% PEG. Successful transformants were selected on complete media
(CM) agar containing Hygromycin B to a final concentration of 200 µg mL^{-1}.

Southern blot blotting

This procedure was adapted from the protocol of Southern (58) with adjustments as described
by Sambrook *et al* (59). Fungal genomic DNA (15 µg) was digested with the appropriate
restriction enzyme overnight at 37°C. The digested genomic DNA was then separated by gel
electrophoresis. The DNA embedded in the agarose gel was de-purinated by immersion in 0.25
M HCl before transferring the DNA onto Hybond-NX membrane (Amersham Biosciences).
The membrane was exposed to UV light in a BLX-254 cross linker (Bio-Link®) to fix and
immobilise the DNA. The membrane was then probed overnight using a DIG-labelled PCR
product in freshly prepared hybridization buffer (25-50ng mL) in a 50 mL falcon tube.
Chemiluminescent detection of DIG-labelled DNA was carried using CDP-Star solution
(Roche) before detection.

Whole genome sequencing

Purified RNA-free DNA was obtained using Hexadecyltrimethylammonium Bromide
(CTAB). Template quality was assessed using a NanoDrop spectrophotometer and Qubit.
Sequencing was carried out at Exeter Sequencing services (University of Exeter, UK) and
Novogene (Cambridge, UK). NEXTflex^{TM} Rapid DNA-seq Library Prep Kit was used to
prepare and index libraries before sequencing on HiSeq 2500 (Illumina) with two lanes per
sample. The quality of sequencing reads was first checked using the FastQC toolkit
(http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/). From the raw data (fastq files),
adaptor sequences were trimmed from sequences containing adaptors and low-quality reads
removed using the fastq-mcf program. The trimmed sequences were aligned to the reference
genome (70-15) (55) using BWA (Burrow Wheeler Aligner) https://github.com/lh3/bwa (60).
Bam files were visualized on IGV genome viewer to determine CRISPR gene deletion.

Copy number variation of *PWL2*

A total of 286 *M. oryzae* isolates with raw Illumina-based sequencing information were downloaded from NCBI (performed on October 16th, 2019). Copy number variation was assessed using a k-mer analysis approach using the k-mer analysis toolkit (KAT; v2.4.1). Coding sequence information for *M. oryzae* isolate 70-15 were used as template for k-mer analysis and raw Illumina reads were input. Default parameters were used including k-mer of length 27 nt. Copy number variation of individual effectors is based on average k-mer coverage as compared with the median coverage for all genes.

Phylogenetic analysis of *Magnaporthe*

Genome sequences of diverse *M. oryzae* isolates were identified based on literature review and search on NCBI (performed on August 9th, 2019). For isolates having only Illumina sequencing data, raw paired reads were downloaded from NCBI, trimmed using Trimmomatic (v0.36) using the parameters: removing of adapters with ILLUMINACLIP:TruSeq2-PE.fa:2:30:10, remove leading low quality or N based with quality below 5 for leading (LEADING:5) and trailing sequence (TRAILING:5), scan and cut reads with 4 bp sliding window below 10 (SLIDINGWINDOW:4:10), and minimum length of 36 bp (MINLEN:36). KmerGenie (v1.7048) was used to identify an optimal k-mer for genome assembly using default parameters. Genome assembly was performed using minia (v0.0.102) with default parameters. kSNP3 (v.3.021) was used to develop a phylogenetic tree of *M. oryzae* using assembled genomes as input with parameters of k-mer of 29 bp and minimum fraction of 0.4. The phylogenetic tree was generated using RAxML (v) using the General Time Reversible model of nucleotide substitution under the Gamma model of rate heterogeneity.

Generation of *Agrobacterium* transformation plasmids

Single or multiple DNA fragments were cloned into binary vector using In-Fusion HD Cloning Kit (Clontech, USA). Briefly, fragments from cDNA were amplified using primers that introduce a 15bp overhang complementary to sequences at the restriction sites of the destination vector or adjacent insert fragments. Alternatively synthesized DNA fragments were designed to have 15bp overhang complementary to sequences at the restriction sites of a binary vector pG514 customized for infusion cloning by digesting with either *Xba*1 and *Pca*1 or *Sac*1 (New England Biolabs) to remove the *ccDB* toxin encoding gene. Positive transformation was accessed using colony PCR and constructs sequenced and analyzed on SNAP gene.

Transient expression in *Nicotiana benthamiana*

Agrobacterium tumefaciens strain *GV3101* [61] was used for transient expression. Three-week-old *N. benthamiana* leaves were infiltrated with transformed *Agrobacterium* carrying T-DNA constructs, expressing the gene of interest. Bacterial cultures were diluted to obtain a final OD 600nm of 0.4 in agroinfiltration buffer (10mM MES, 10 mM $MgCl_2$, 150 μM acetosyringone, pH 5.6). Leaf discs were cut from agroinfiltrated tissue 48 hpi and subjected for microscopy or used co-IP.

Plant transformation and ROS measurement

Agrobacaterium tumefaciens strain *AGL1* [61] was used for plant transformation as described in [62]. Positive transgenics plants were first selected on hygromycin followed by confirmation using PCR. Alternatively, leaf discs were collected from transgenic plants and analysed for copy number at iDNA Genetics Ltd (Norwich, UK) to detect hygromycin insertion events. In addition, expression of the protein was evaluated using SDS-PAGE. To measure response to

PTI elicitors, a 4-mm diameter biopsy punch (Integra^TM Miltex ^TM) was used to cut leaf discs
from 5 weeks old *H. vulgare* and *O. sativa* transgenic plants. The leaf discs were transferred
into 96-well-plates (Greiner Bio-One) containing 100 µL of distilled H_2O in each well and
incubated overnight at room temperature. The assay was carried out by replacing the distilled
H_2O in the wells by a 100 µL reaction solution of 100 µM Luminol or L-012 (Merck), 20 µg
mL^{-1} horseradish peroxidase (Merck) together with elicitors, flg22 (1 µM final concentration),
chitin (1mg/mL final concentration) or without (mock). The photon count was carried out using
the (HRPCS218, Photek) equipped with a 20 mm F1.8 EX DG ASPHERICAL RF WIDE
LENS (Sigma Corp). Each experiment was repeated three times conducted a minimum of three
times, yielding consistent outcomes.

Co-immunoprecipitation and sample preparation for mass spectrometry

Leaves were harvested from two-three weeks old barley transgenic plants and rapidly frozen
in liquid nitrogen before storage or immediately ground into fine powder using a
GenoGrinder® tissue homogenizer. Ground powder was quickly transferred into ice-cold 1.5
mL Eppendorf tube and 2 mL of ice-cold extraction buffer (GTEN [10% (v/v) glycerol, 25 Mm
Tris pH 7.5, 1 mM EDTA 150 mM NaCl, 2% (w/v) PVPP, 10 Mm DTT, 1 X protease inhibitor
cocktail (Sigma), 0.5% (v/v) IGEPAL added and mixed thoroughly. This was centrifuged at
3000 x g for 10 min at 4°C to recover total proteins in the supernatant and repeated twice. A
volume of 2 mL of total protein was mixed with 25 µL of GFP-TRAP agarose beads (50%
slurry, ChromoTek) and incubated shaking overnight at 4°C. The GFP-TRAP agarose beads
were centrifuged for 2 min at 3000 x g before washing three times with wash buffer
(ChromoTek). Bound proteins were recovered by resuspending the beads in loading buffer
(loading dye, 10 mM DTT, H2O) and incubated at 70°C for 10 minutes. An aliquot of the
sample was run on SDS-PAGE gel and used for Western blotting (63). Recovered proteins
from co-immunoprecipitation were run on an SDS-PAGE gel for approximately 1 cm. The gel
was washed for 1 h in distilled water followed by 1 incubation SimplyBlue^TM Safe stain. The
gel was then washed three times in distilled water the region containing proteins excised using
a scalpel.

Mass spectrometry and data processing

Protein purification, immunoprecipitation, sample preparation, liquid chromatography
followed by tandem mass spectrometry (LC-MS/MS), and data analysis were previously
described in full detail (64). Briefly proteins identified in immunoaffinity-enriched samples
were measured with data dependent method on high resolution LC-MS systems, Orbitrap
Fusion (Thermo Ltd). The acquired spectra were peak-picked and searched by Mascot search
engine (Matrix Science Ltd) to identify the peptide sequences from the search space defined
by the background proteome. The peptides were combined into proteins based on the principle
of parsimony by the search engine. Resulting proteins were further described by quantitative
values based on the number of spectra that identify them. The individual runs were combined
in the Scaffold program (Proteome Software Inc.), where the data were evaluated and filtered
to contain less than 1% false positives (FDR) and resulted matrix exported as a spreadsheet.
The matrix of proteins detected in different samples serves as the input for an R script for
further processing and visualization.

Yeast two-hybrid analysis

To clone genes of interest into bait (pGBKT7 DNA-BD cloning) vector or prey (pGADT7
activating domain) vector, using the In-Fusion HD Cloning Kit (Clontech) kit. The bait and
prey vectors were digested using vector was digested using *Bam*H1 and *Eco*R1 while the gene

to be inserted was PCR-amplified using primers containing 15 bp overhangs with homology to
the two ends of the digested bait vector. For transformation, a single colony Y2H gold yeast
strain was mixed in 1 mL of liquid yeast extract peptone dextrose (YPD). Competent cells were
prepared as described in manufactures transformation protocol (Zymo Research). Briefly, for
transformation 700 ng – 1 µg of plasmids expressing Pwl2 or other effectors in pGADT7 and
*Hv*HIPP43 or plant proteins in pGBKT7 were co-transformed in competent cells and Frozen-
EZ Yeast Solution 3 added before incubation at 28°C for 1 h and transformed cells plated on
selection media lacking Leucine (L) and Tryptophan (W) and incubated at 28° C for 3-5 days.
To detect interactions the colonies were transferred to media lacking Leucine (L), Tryptophan
(W), Adenine (A) Histidine (H), Bromo-4-Chloro-3-Indolyl a-D-galactopyranoside (X-a-gal)
and 10 mM 3-amino-1,2,4-triazole (3AT) (Sigma). The plates were imaged after 60-72
incubation at 28°C. Each experiment was conducted a minimum of three times, yielding
consistent outcomes.

In-planta co-immunoprecipitation (co-IP)

To test interaction of protein *in planta*, genes of interest tagged with either GFP or Myc were
cloned into the vector pGW514 and transformed into *Agrobacterium* strain *GV301* and co-
infiltrated (OD_{600} = 0.4 for effectors and OD_{600} = 0.6 for HIPP43) into 3 to 4-week-old *N.*
benthamiana leaves and incubated for 48 h to allow protein expression. Total protein was
isolated in ice-cold extraction buffer (GTEN [10% (v/v) glycerol, 25 Mm Tris pH 7.5, 1 mM
EDTA 150 mM NaCl], 2% (w/v) PVPP, 10 Mm DTT, 1 X protease inhibitor cocktail (Sigma),
0.5% (v/v) IGEPAL). Total protein was co-immunoprecipitated using the anti-GFP M2 resin
(Sigma-Aldrich, St. Louis, MO) and washed 3 times using immunoprecipitation buffer before
analyzing on an SDS-PAGE gel. Recovered proteins from co-immunoprecipitation were
separated on an SDS-PAGE gel and transferred onto a polyvinylidene diflouride (PVDF)
membrane using a Trans-Blot turbo transfer system (Bio841 Rad, Germany). Detection was
performed using the appropriate antibody either anti-GFP-HRP or anti-Myc-HRP. Imaging
was carried out using an ImageQuant LAS 4000 luminescent imager (GE Healthcare 844 Life
Sciences, Piscataway, NJ, U.S.A.) according to manufacturer's instructions.

Sub-cellular localization and plasmodesmata quantification

Leaf discs were obtained from *N. benthamiana* plants agroinfiltrated with different construct
combinations at 24 and 48 hpi and mounted on a slide immersed in perflurodecalin and
observed using X63 oil immersion lens. For Leica SP8 laser confocal microscopy, settings
were as follows; GFP, YFP and RFP/mCherry tagged proteins were excited using 488, 514,
and 561 nm laser diodes and the emitted fluorescence detected using 495-550, 525-565, and
570-620 nm respectively. The auto-fluorescence from chlorophyll was detected at 650-740 nm.
To stain and quantify plasmodesmata, agroinfiltrated leaves were further infiltrated with 0.1%
aniline (Sigma-Aldrich #415049, in PBS buffer, pH 7). Images were analyzed using ImageJ to
determine the number of plasmodesmata. First, cell peripheries were divided into 20 µm
sections and the number of plasmodesmata determined per 20 µm.

Phylogenetic analysis of grass HMA proteins

Proteins from *B. distachyon* (314; v3.1), *H. vulgare* (Jul2020), *T. aestivum* (RefSeqv2.1; 09-
16-2020), *O. sativa* (323; v7.0), *Oropetium thomaeum* (386; v1.0), *Sorghum bicolor* (454;
v3.1.1), *Setaria italica* (312; v2.2), and *Zea mays* (RefGen_V4) containing an HMA domain
were identified using InterProScan (v5.59-91.0; Pfam PF00403). The HMA domain was
extracted using the script QKdomain_process.py including an additional 10 amino acid
sequences N and C-terminal of the Pfam boundaries (-n 10 -c 10). The non-redundant set of

HMA domains were identified using CD-HIT (v.4.8.1) with parameter -c 1.0. Structure guided
multiple sequence alignment was performed using MAFFT with parameters dash, max iteration
of 1000, and globalpair. Maximum likelihood phylogenetic analysis was performed using
RAxML (v8.2.12) using the Gamma model of rate heterogeneity, JTT amino acid substitution
model, and 1,000 bootstraps. Coding sequence was identified for the HIPP43 gene family and
aligned using MUSCLE (v5) translation alignment using default parameters. Maximum
likelihood phylogenetic analysis was performed using RAxML (v8.2.12) using the General
Time Reversible model of nucleotide substitution under the Gamma model of rate
heterogeneity and 1,000 bootstraps.

Statistical analysis and Protein structure prediction using AlphaFold2

Significance difference between groups of samples was performed using GraphPad Prism 10.
P values < 0.05 were considered significant, ****$P<0.0001$, ***$P<0.001$, **$P<0.01$ while
values *P* values > 0.05 were considered as non-significant in an unpaired Student's t-test.
Structure prediction was carried out using AlphaFold2 and the structure analyzed, and figures
made using ChimeraX

Fig. 1. **Pwl2 is a BIC localized effector that is ubiquitous in isolates of *M. oryzae* and**
displays conserved structural features.
(*A+C*) Micrographs showing Pwl2 secreted through the Biotrophic Interfacial Complex (BIC)
in rice cells during early infection. Conidial suspension at 1X 10^5 mL^{-1} of *M. oryzae* strain co-
expressing Bas1-GFP and Pwl2-mRFP were inoculated onto a susceptible cultivar Moukoto
rice leaf sheath and images captured at 26 hpi. Fluorescence of the two effectors was observed
as small punctate signals in the same BIC. (*C+D*) Co-infection assay of rice leaf sheath with
M. oryzae strains expressing Pwl2-mRFP and Pwl2-GFP at 30 hours post-infection (hpi).
Micrographs and line scan graph shows there is an absence of mixed fluorescence confirming
the BIC does not contain Pwl2 transferred from rice cells. BICs indicated by magenta arrows
for mRFP and green arrows for GFP. (*E*) Superimposition of different Pwl2 variants predicted
using AlphaFold2 onto the resolved Pwl2 structure (companion paper to this study by Zdrzalek
et al (15)) indicating region of polymorphism. (*F*) Phylogenetic analysis and *PWL* gene family
distribution in *M. oryzae*. A maximum parsimony tree including *M. oryzae* strains and isolates
from related species *Magnaporthe grisea* and *Pyricularia pennisetigena*. The heatmap
indicates the presence/absence of genes in the *PWL* family. *PWL1* is found in *Eleusine* and
Oryza lineages. *PWL2* is conserved across various *M. oryzae* lineages and isolates infecting
crabgrass and pearl millet. *PWL3* is specific to *Oryza*, *Lolium*, *Setaria*, and *Eleusine* lineages.
PWL4 is predominantly found in *Eleusine*, *Eragrostis*, *Lolium*, and *Triticum* lineages.
Maximum parsimony tree of *M. oryzae* isolates was generated using a standalone, whole
genome multiple alignment/SNP calling program, kSNP3. We used *Pwl1, Pwl2, Pwl3* and
Pwl4 protein sequence to query for presence/absence polymorphism of the gene families using
tblastn.
709

710
711
Fig. 2. *PWL2* occurs in multiple loci and shows high copy number variations in field
isolates.
(*A*) Schematic showing the estimated chromosomal location of *PWL2* on multiple loci (Chr3
and 6) of the reference genome 70-15 and on equivalent region of assembled contigs of
laboratory strain Guy11 genome. *PWL2* is flanked by *POT2* and *MGR583* repeated sequences
suggesting a possible involvement in translocation of events into different loci in the genome.
Arrows indicate location of *PWL2*. (*B*) A k-mer analysis on sequenced raw reads was used to

determine copy number variation of *PWL* family genes in different *M. oryzae* isolates. Plot
shows high copy number of *PWL2* in analyzed genomes (n = 286) compared to the other *PWL*
gene family members, another effector *SLP1,* and selected control genes *MPG1* and *SEC5*.
Similarly, *AVR-Pik* shows multiple copies in different isolates. Copy number of *PWL2* in
selected isolates Guy11, KE002, BF48, 70-15 is indicated with Ina168 that lack *PWL2* used as
a negative control.
725
726

Fig. 3. CRISPR-Cas9 *Δpwl2* mutants demonstrate gain of virulence on weeping lovegrass
and barley carrying *Mla3*.
(*A*) Schematic illustration of the CRISPR-Cas9 mediated gene editing process for inserting the
Hygromycin Phosphotransferase (*HPH*) gene cassette at the *PWL2* locus. The guided sequence
(sgPWL2) directs Cas9 to introduce a double stranded break at the *PWL2* locus, with the DNA
repair template containing the Hygromycin resistance gene cassette and flanking regions of the
PWL2 gene. This leads to mutations in the form of indels or gene replacements. Figure created
with BioRender https://biorender.com/. Positive transformants were identified by amplifying
the *PWL2* coding sequence from the wild type Guy11 and the hygromycin cassette from
transformants with the three copies deleted. (*B*) Comparison of disease symptoms in weeping
lovegrass (*Eragrostis curvula*) infected with different *M. oryzae* isolates. In the top panel,
typical disease lesions produced by *E. curvula*-infecting isolate G17 are shown. In the middle
panel, CRISPR-Cas9 deletion mutants exhibit gain of virulence on weeping lovegrass
compared to Guy11. In the bottom panel, complemented strains with both native and
constitutive RP27 promotors showed a loss of virulence on weeping lovegrass. (*C*) *Δpwl2*
mutants display reduced pathogenicity on rice cultivar CO39. Conidial suspensions at 1×10^5
mL^{-1} spores/mL from Guy11 and *Δpwl2* T6 were used to inoculate 21-d-old seedlings of the
blast-susceptible cultivar CO39, and disease symptoms recorded after 5 dpi. (*D*) The box plots
show the lesion size (top) and lesion density (bottom) of seedlings infected with Guy11 and
Δpwl2 T6. The lower horizontal line shows the minimum value, and the upper horizontal line
shows the maximum value. The lower border and upper border of the box show the lower
quartile and upper quartile, respectively. The line in the box shows the median. (*E*) The *Δpwl2*
mutant exhibits enhanced pathogenicity on barley cultivar Baronesse (*Mla3*). Conidial
suspensions at 1×10^5 mL^{-1} spores/mL from Guy11 and *Δpwl2* T6 were used to inoculate 10-
day old seedlings of barley lines Golden Promise, Siri, Nigrate and Baronesse, and disease
symptoms recorded after 5 dpi. (*F*) The box plot shows lesion size of barley seedlings infected
with Guy11 and *Δpwl2* T6. The lower horizontal line shows the minimum value, and the upper
horizontal line shows the maximum value. The lower border and upper border of the box shows
the lower quartile and upper quartile, respectively. The line in the box shows the median.
Significance between groups of samples was performed using Unpaired Student's t-test.
****$P<0.0001$, **$P<0.01$, $P<0.05$, NS indicates no significant difference.

Fig. 4. Pwl2 suppresses PAMP-induced ROS in transgenic barley lines and interacts with a host HIPP43.

(*A*) Transgenic lines expressing Pwl2-YFP, control (free YFP) did not show growth defect compared to the background line, Golden Promise. (*B*) ROS production in leaf discs collected from 4–week-old stable transgenic lines expressing Pwl2-YFP, control (free YFP) and Golden Promise induced by 1 uM chitin or flg22 (n > 8). Points represent mean; error bars represent SEM. (*C*) Dot plot shows cumulative ROS production over 60 min for Golden Promise control and Pwl2-YFP. The lower horizontal line shows the minimum value, and the upper horizontal line shows the maximum value, the middle line shows the mean value. Significance between groups of samples was performed using Unpaired Student's t-test. (*D*) Leaf drop infection on barley transgenic lines expressing free-YFP, Pwl2-YFP compared to infection on wild type Golden Promise 3-4 dpi. (*E*) The dot plot represents the number of rice blast disease lesions per 5 cm on Golden Promise, EV-YFP and Pwl2-YFP transgenic lines in three independent repetitions of the experiment. Unpaired Student's t-test was performed to determine significant differences. (*F*) Putative Pwl2 interacting peptides were immunoprecipitated from protein extracted from 3-week-old stable Golden Promise transgenic lines expressing Pwl2-YFP or free cytoplasmic YFP using anti-GFP antibodies, and LC-MS/MS was performed to identify unique putatively interacting peptides. The scale 0-20 represents the Log2fold change in peptides when compared to the control. (*G*) Pwl2 interacts with *Hv*HIPP43 in a yeast-two-hybrid assay. Simultaneous co-transformation of pGAD-Pwl2 (prey vector) and pGBK-HIPP43 (bait vector) and PGBKT7-53 and pGADT7-T (positive control) pGADTT7-T and pGBKT7-Lam (negative control) into Y2H gold strain resulted in the activation of four reporter genes and growth on high-stringency medium (−Ade, −Leu, –Trp, _His +X-α-gal and 3-amino-1,2,4-triazole). Co-transformation also activates the expression of *MEL1*, which results in the secretion of α–galactosidase and the hydrolysis of X-α-gal in the medium, turning the yeast colonies blue (*H*) Co-immunoprecipitation (co-IP) of Pwl2-YFP and Myc-HIPP43 in *N. benthamiana* leaves. C-terminal GFP tagged Pwl2 and C-terminal Myc-HIPP43 was cloned into the vector pGW514 and transformed into *Agrobacterium* strain *GV301* and co-infiltrated into *N. benthamiana* leaves and left to incubate for 48 h. Immunoprecipitates were obtained with anti-GFP or anti-Myc affinity matrix beads and probed with anti-GFP-peroxidase and anti-Myc-peroxidase (HRP-conjugated) antibodies. Total protein extracts were also probed with appropriate (HRP-conjugated) antibodies. (*I*) Yeast-two-hybrid analysis shows; upper panel, *Hv*HIPP43 interacts with *PWL* gene family products Pwl1, Pwl3, Pwl4 and variant *pwl2-3*. (*J*) Pwl2 binds to HIPP43 homologs from rice (*Os*HIPP43), wheat (*Taes*HIPP43), weeping love grass (*Ec*HIPP43), and foxtail millet (*Si*HIPP43).

Fig. 5. HvHIPP43 suppresses PAMP-induced ROS in transgenic barley.
(*A-D*) ROS production measured from leaf disks collected from 4–week-old stable transgenic
line expressing YFP-HIPP43 and Golden promise (control) in the absence and presence of 1
uM flg22 or 1 mg/ml Chitin (n > 8). For ROS assay, line graphs (upper panel) points represent
mean per time point and as cumulative ROS (*C+D*) production over 60 min, error bars
represent SEM. For dot plot graphs show the average cumulative photon count. The lower
horizontal line shows the minimum value, the upper horizontal line shows the maximum value,
the middle line shows the mean value. Significance between groups of samples was performed
using unpaired Student's t-test. ****$P<0.0001$, ***$P<0.001$, **$P<0.01$. (*E*) Leaf drop
infection on barley transgenic lines expressing YFP-HIPP43 compared to wild type Golden
Promise dpi. Conidial suspensions at 1×10^5 mL^{-1} spores/mL from Guy11 were used to
inoculate 21-d-old seedlings of the blast-susceptible Golden Promise control and Golden
Promise expressing YFP-HIPP43. Disease symptoms were recorded after 5 dpi. (*F*) Dot plot
showing the average lesion size on barley lines infected with Guy11. The lower horizontal line
shows the minimum value, the upper horizontal line shows the maximum value, the middle
line shows the mean value. Significance between groups of samples was performed using
unpaired Student's t-test. ***$P<0.001$. (*G+H*) Micrographs and line scan graph showing
mCherry-HIPP43 localizing as small puncta on the plasma membrane when expressed in *N.*
benthamiana. Staining of callose using aniline blue overlaps with mCherry-HIPP43,
confirming mCherry-HIPP43 localises exclusively at the plasmodesmata (PD) localization and
is absent from vesicular bodies (VB). (*I+J*) Micrographs and line scan graph showing the
presence of YFP-Pwl2, alters m-Cherry-HIPP43 localisation, where it was observed to
translocate to the cytoplasm, cytoplasmic and vesicular structures (*K+L*) Micrographs and line
scan graph showing free-YFP fluorescence is observed in the cytoplasm and does not overlap
with mCherry-HIPP43 PD localization. (*M+N*) Micrographs and line scan graph showing
mCherry-HIPP43 remains in the PD in the presence of cytoplasmically localised AVR-PikE**.**
Scale bars represent 20 μm**.** Significance between groups of samples was performed using
unpaired Student's t-test. ****$P<0.0001$.
835

836
837
838
839

Fig. S1. Pwl2 is co-expressed *in-planta* with other known effectors and is ubiquitous in
isolates of *M. oryzae* showing global distribution.
(*A*) Illustration of how *PWL2* has been described as a host range determinant gene in weeping
lovegrass and barley. *M. oryzae* strain carrying *PWL2* produces incompatible reaction on
weeping lovegrass and cv. Baronesse (*Mla3*). (*B*) A susceptible rice line Moukoto, was infected
by a *M. oryzae strain* KE002 using leaf drop infection method and samples collected at 24, 36,
48, 59 and 72 hpi. RNA isolated from infected rice leaves were sequenced and the reads
normalized against *M. oryzae* mycelia KE002 RNA to determine differentially expressed
genes. A weighted gene co-expression network analysis grouped gene expression profile into
10 co-expression modules. The line graph (WGCNA) showing the representative eigengene
for module 4 in *PWL2* is co-expressed with other known effector protein encoding, *AVR-Pi9*,
SLP1, *BAS1-4* and a necrosis inducing effector *NEP1*. (*C*) Heatmap showing relative
expression levels (log2fold change) of *PWL2* and other know effectors during infection. (*D*)
Global distribution of Pwl2 alleles in *M. oryzae* isolates with publicly available genomes.
Tblastn was used determine presence of Pwl2 in *M. oryzae* genomes and *R* package *rworldmap*
v.1.3.4 used to create the map and occurrence of different alleles according to isolate point of
collection. *pwl2-2* mostly present in isolates from Asia while *pwl2-3* present in wheat blast
isolates from South America.

Fig. S2. A variant of Pwl2, pwl2-3 is not recognized by Mla3 and cognate resistance

against Pwl2 in weeping lovegrass.

(*A*) Multiple amino acid sequence alignment of Pwl2 protein and naturally occurring variants.
Alignment was carried using ClustalW (Jalview). (*B*) Neighbor-joining tree based on amino
acid sequence of variants encoded by *PWL* gene family. Tree IDs contains isolate name from
which the variant was identified, and each dot represent presence in each host-limited forms.
(*C* and *D*) Wheat blast isolate BTJP-14-6 carrying a *PWL2* variant, *pwl2-3* with Glu-89-Gln,
Lys-91-Gln and Ser-92-Ile substitutions was used for spray inoculation on barley accessions
cv. Golden promise (-*Mla3*) and cv. Baronesse (+*Mla3*) or (*D*) weeping lovegrass seedlings.
Disease symptoms on barley and weeping lovegrass were scored 5 and 7 days respectively,
post-infection. Observations were consistent in three independent infection replicates. (*E*)
Summary of disease reaction of isolates carrying variants of Pwl2 on Nigrate (-*Mla3*) and
Baronesse (+*Mla3*). Lesions on barley were scored 5 days post-infection.
874

877 **Fig. S3. *M. oryzae* isolates carry variants of Pwl1, Pwl3 and Pwl4 and copy number**
878 **variation occurs in all rice blast genetic lineage groups.**
879 Multiple amino acid sequence alignment of Pwl1 (*A*), Pwl3 (*B*) and Pwl4 (*C*) protein and
880 naturally occurring variants. Alignment was carried using ClustalW (Jalview) regions or low
881 conservation are shown in magenta.
882
883

884
885
Fig. S4. Southern blot to determine copy number variation of *PWL2* in selected *M. oryzae*
isolates.
(*A*) Genomic DNA was digested using different enzyme combination *Bam*H1/*Pst*1 and probed
using the *PWL2* 438 bp probe. (*B*) The order of digested genomic DNA from different isolates
is indicated on the lanes. Size estimates are from the *PWL2* loci annotated *M. oryzae* 70-15
genome. When Guy11 genomic DNA was digested using *Pst*1 and *Afl*II or single digested
using *Afl*II, two bands were detected, a single digest using *Pst*1 only produce a single band.
893
894
895

896
897
Fig. S5. Analysis of CRISPR-Cas9 mutants using Integrative Genomics Viewer (IGV).
(A) Reads alignment display of *PWL2* locus on Chromosome 6 from four sequenced *pwl2*
mutant strains, T12, T5, T4 and T6 (lower panel). No reads aligned at *PWL2* (MGG_04301T0)
locus for T6 and T12 and partial alignment for T4 and T5. Read depth around *PWL2* locus is
higher compared to the neighbouring gene MGG_17636T0 suggesting the *PWL2* locus occurs
multiple times. Read coverage and locus length are indicated. (*B*) Sashimi plot showing depth
of reads from T12, T5, T4 and T6 that aligned to the region surrounding *PWL2* locus.
905

Fig. S6. Raw data results from Y2H assay with all other putative interactors from IP-MS.
One-to-one yeast two-hybrid between Pwl2 and selected top candidates from IP-MS analysis,
Pwl2 and Putative Pwl2 Interacting Proteins (PPIPs). HMA integrated rice NLR Pikm-1 and
Pikp-1 were used as specificity controls. Simultaneous co-transformation of pGADT7-Pwl2
(prey vector) and pGBK-HIPP43 (bait vector) and PGBKT7-53 and pGADT7-T (positive
control) pGADTT7-T and pGBKT7-Lam (negative control) into Y2H gold strain resulted in
the activation of four reporter genes and growth on high-stringency medium (−Ade, −Leu, −
Trp, _His +X-α-gal and 3-amino-1,2,4-triazole). Co-transformation also activates the
expression of *MEL1*, which results in the secretion of α–galactosidase and the hydrolysis of X-
α-gal in the medium, turning the yeast colonies blue. HIPP43 exclusively bind Pwl2 in SD/-L-
T-H-A X-a-gal medium with 3AT added.
918

1.0

HIPP43 family

919
920

Fig. S7. Maximum likelihood phylogenetic tree of HMA domains from diverse grass species.

HMA domains were identified from proteins from the grass species *B. distachyon*, *H. vulgare*, *T. aestivum*, *O. sativa*, *Oropetium thomaeum*, *Sorghum bicolor*, *Setaria italica*, and *Zea mays*. Structure guided multiple sequence alignment was performed using MAFFT-DASH and phylogenetic analysis with RAxML. Scale bar indicates substitutions per site. Dots on branches indicate support of 80% or more based on 1,000 bootstraps. The HIPP43 family formed a distinct clade with bootstrap support of 98%.

A

B

930
931
Fig. S8 Maximum likelihood phylogenetic tree for HIPP43 gene family.
(*A*) Full length coding sequence was identified for HIPP43 gene family members, aligned using
MUSCLE translation alignment, and phylogenetic analysis with RAxML. Barley and wheat
(i.e. *Triticeae* lineage) have three copies of HIPP43 per haploid genome. Dots on branches
indicate support of 80% or more based on 1,000 bootstraps. (*B*) Expression at transcript level

of *Os*HIPP43 during infection of a susceptible rice cv. CO39 by Guy11. Hour post inoculation
is shown on the X –axis.

945
946
Fig. S9. HIPP43 isoprenylation motif is required for PD localisation.
(*A*) A schematic representation of domain structure of HIPP43 showing the HMA domain
region and the isoprenylation motif at the C-terminus. (*B*+*C*) Micrographs showing
localization of HIPP43 without the isoprenylation motif (-Iso-mCherry-HIPP43) and C-
terminus tagged, HIPP43-YFP. (*B*) Deletion of the isoprenylation abolishes PD localisation
but HIPP43 localises in the nucleus and partially in the plasma membrane and cytoplasm after
48h. (*C*) When HIPP43 was tagged at the C-terminus, localisation was observed in the plasma
membrane, partially in the cytoplasm and PD, but predominantly in the chloroplast. Scale bars
represent 20 µm.
956

957
958
Fig. S10. Pwl2 promotes HIPP43 cytoplasmic localisation.
(*A-C*) Micrographs showing localization of free-YFP (A), AVR-PikE-YFP (B) and YFP-Pwl2
(C). Both free-YFP and AVR-PikE-YFP show localisation in the cytoplasm while YFP-Pwl2
showed localisation in cytoplasm and cytoplasmic mobile structures. (*D-F*) HIPP43
localisation in presence of AVR-PikE and Mep3 is unaffected but altered in the presence of
Pwl2. (*D*) Micrographs showing that the presence of YFP-Pwl2, alters m-Cherry-HIPP43
localisation, where it was observed to translocate to the cytoplasm and cytoplasmic vesicular
structures. Arrows indicated cytoplasmic localisation and cytoplasmic mobile bodies (*E+F*)

Micrographs showing HIPP43 is co-localisation with two different MAX-fold effectors AVR-
PikE and Mep3. mCherry-HIPP43 remains in the PD in the presence of cytoplasmically
localized AVR-PikE and Mep3. Arrows indicated PD. Scale bars represent 20 µm.
970
971

972 **Table S1 Primers used in this study.**

Primer	Sequence
PWL2ORF	ATGAAATGCAACAACATCATCCTCCC
PWL2ORR	ACATAATATTGCAGCCCTCTTCTCGC
PWL2f	GGTTCTTATTATGGTCCCGGGTGA
PWL2r	GGTTCTTATTATGGTCCCGGGTGA
PWL2848F	CGCGGTGGCGGCCGCTCTAGAGACCGGGCACGAACCCGGCAGGC
PWL2-R	CATAATATTGCAGCCCTCTT
PWL2_GFPF	GGGCTGCAATATTATGATGGTGAGCAAGGGCGAGGAGCTG
KOLFF	ACAACGCGGTGTAAAGATTCGGGT
KOLFR	GTCGTGACTGGGAAAACCCTGGCGAGACCAAAGCAAAGGGAGGATGA
KORFF	TCCTGTGTGAAATTGTTATCCGCTGCGAGAAGAGGGCTGCAATATTA
PG514-myc-F	AACACGGGGGACTCTAGAGAACAAAAGTTGATCTCTGA
YFPPWL2-F	GACGAGCTGTACAAGGGTGGCGGGTGGACTAAC
PWL2-N-term-YFPR	GTTAGTCCACCCGCCACCCTTGTACAGCTCGTC
PWL2-R-p514	GAACGATCGGGGAAATTCGTTACATAATATTGCAGCCCT
HAPwl2F	AGAGAACACGGGGGACTCTAGAATGGGTGGCGGGTGGACTAAC
PG514-myc-F	AACACGGGGGACTCTAGAGAACAAAAGTTGATCTCTGA
myc-HIPP43-R	TCCAAGACGCCCATAAGGTCCTCTTCAGAAA
myc-HIPP43F	ATTTCTGAAGAGGACCTTATGGGCGTCTTGGA
HIPP43-NOS-R	ACGATCGGGGAAATTCGTCACATGATGGTGCAG
PG514-mCherry-F	AACACGGGGGACTCTAGAATGGTGAGCAAGGGC
mCherry-HIPP43F	GGACGAGCTGTACAAGATGGGCGTCTTGGA
mCherry-HIPP43R	TCCAAGACGCCCATCTTGTACAGCTCGTCC
KT7_HIPP34F	CATGGAGGCCGAATTCATGGGCGTCTTGGA
KT7_HIPP34R	GCAGGTCGACGGATCCTCACATGATGGTGCAG

References

1. J. D. Jones, J. L. Dangl, The plant immune system. *Nature* **444**, 323-329 (2006).

2. R. A. Wilson, N. J. Talbot, Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*. *Nat Rev Microbiol* **7**, 185-195 (2009).

3. X. Yan *et al.*, The transcriptional landscape of plant infection by the rice blast fungus *Magnaporthe oryzae* reveals distinct families of temporally co-regulated and structurally conserved effectors. *The Plant Cell* **35**, 1360-1385 (2023).

4. E. Oliveira-Garcia, X. Yan, M. Oses-Ruiz, S. de Paula, N. J. Talbot, Effector-triggered susceptibility by the rice blast fungus *Magnaporthe oryzae*. *New Phytologist* **n/a** (2023).

5. Y. Sugihara *et al.*, Disentangling the complex gene interaction networks between rice and the blast fungus identifies a new pathogen effector. *PLOS Biology* **21**, e3001945 (2023).

6. S. Cesari *et al.*, The Rice Resistance Protein Pair RGA4/RGA5 Recognizes the *Magnaporthe oryzae* Effectors AVR-Pia and AVR1-CO39 by Direct Binding *The Plant Cell* **25**, 1463-1481 (2013).

7. D. Ortiz *et al.*, Recognition of the *Magnaporthe oryzae* Effector AVR-Pia by the Decoy Domain of the Rice NLR Immune Receptor RGA5. *The Plant Cell* **29**, 156-168 (2017).

8. I. Ashikawa *et al.*, Two Adjacent Nucleotide-Binding Site–Leucine-Rich Repeat Class Genes Are Required to Confer Pikm-Specific Rice Blast Resistance. *Genetics* **180**, 2267-2276 (2008).

9. H. Kanzaki *et al.*, Arms race co-evolution of *Magnaporthe oryzae* AVR-Pik and rice Pik genes driven by their physical interactions. *The Plant Journal* **72**, 894-907 (2012).

10. K. de Guillen *et al.*, Structure Analysis Uncovers a Highly Diverse but Structurally Conserved Effector Family in Phytopathogenic Fungi. *PLoS Pathog* **11**, e1005228 (2015).

11. A. Maqbool *et al.*, Structural basis of pathogen recognition by an integrated HMA domain in a plant NLR immune receptor. *eLife* **4**, e08709 (2015).

12. X. Zhang *et al.*, A positive-charged patch and stabilized hydrophobic core are essential for avirulence function of AvrPib in the rice blast fungus. *The Plant Journal* **96**, 133-146 (2018).

13. J. A. Sweigard *et al.*, Identification, cloning, and characterization of PWL2, a gene for host species specificity in the rice blast fungus. *Plant Cell* **7**, 1221-1233 (1995).

14. H. J. Brabham *et al.*, Barley MLA3 recognizes the host-specificity effector Pwl2 from *Magnaporthe oryzae*. *The Plant Cell* 10.1093/plcell/koad266, koad266 (2023).

15. Rafał Zdrzałek *et al.*, Bioengineering a plant NLR immune receptor with a robust recognition interface towards a conserved fungal pathogen effector. *bioRxiv* (2024).

16.   N. Suzuki, Y. Yamaguchi, N. Koizumi, H. Sano, Functional characterization of a heavy
metal binding protein CdI19 from *Arabidopsis*. *The Plant Journal* **32**, 165-173 (2002).

17.   P. E. Dykema *et al.*, A new class of proteins capable of binding transition metals. *Plant
Molecular Biology* **41**, 139-150 (1999).

18.   W. Gao, S. Xiao, H.-Y. Li, S.-W. Tsao, M.-L. Chye, *Arabidopsis thaliana* acyl-CoA-
binding protein ACBP2 interacts with heavy-metal-binding farnesylated protein AtFP6.
New Phytologist **181**, 89-102 (2009).

19.   M. Hála, V. Žárský, Protein Prenylation in Plant Stress Responses.
http://dx.doi.org/10.3390/molecules24213906.

20.   S. Fukuoka *et al.*, Loss of function of a proline-containing protein confers durable
disease resistance in rice. *Science* **325**, 998-1001 (2009).

21.   K. Oikawa *et al.*, The blast pathogen effector AVR-Pik binds and stabilizes rice heavy
metal-associated (HMA) proteins to co-opt their function in immunity. *bioRxiv*
10.1101/2020.12.01.406389, 2020.2012.2001.406389 (2020).

22.   A. R. Bentham *et al.*, A single amino acid polymorphism in a conserved effector of the
multihost blast fungus pathogen expands host-target binding spectrum. *PLoS Pathog*
17, e1009957 (2021).

23.   M. C. Giraldo *et al.*, Two distinct secretion systems facilitate tissue invasion by the rice
blast fungus *Magnaporthe oryzae*. *Nat Commun* **4**, 1996 (2013).

24.   E. Oliveira-Garcia *et al.*, Clathrin-mediated endocytosis facilitates the internalization
of *Magnaporthe oryzae* effectors into rice cells. *The Plant Cell* **35**, 2527-2551 (2023).

25.   P. Kankanala, K. Czymmek, B. Valent, Roles for rice membrane dynamics and
plasmodesmata during biotrophic invasion by the blast fungus. *Plant Cell* **19**, 706-724
(2007).

26.   Kang S.C., Sweigard J.A., V. B, The PWL host specificity gene family in the blast
fungus *Magnaporthe grisea*. *Molecular Plant-Microbe Interactions* **8**, 939-948 (1995).

27.   N. J. Talbot, On the trail of a cereal killer: Exploring the biology of *Magnaporthe
grisea*. *Annu Rev Microbiol* **57**, 177-202 (2003).

28.   S. H. Ou, Pathogen Variability and Host-Resistance in Rice Blast Disease. *Annu Rev
Phytopathol* **18**, 167-187 (1980).

29.   C. D. Cruz, B. Valent, Wheat blast disease: danger on the move. *Trop Plant Pathol* **42**,
210-222 (2017).

30.   Y. Inoue *et al.*, Evolution of the wheat blast fungus through functional losses in a host
specificity determinant. *Science* **357**, 80-83 (2017).

31.   S. Asuke *et al.*, Evolution of an Eleusine-Specific Subgroup of *Pyricularia oryzae*
Through a Gain of an Avirulence Gene. *Mol Plant Microbe Interact* **33**, 153-165
(2020).

32.   S. M. Latorre *et al.*, Genomic surveillance uncovers a pandemic clonal lineage of the
1056   wheat blast fungus. *PLOS Biology* **21**, e3002052 (2023).

1057   33.   A. J. Foster *et al.*, CRISPR-Cas9 ribonucleoprotein-mediated co-editing and
1058   counterselection in the rice blast fungus. *Scientific Reports* **8**, 14355 (2018).

1059   34.   A. Sánchez-Vallet, J. R. Mesters, B. P. H. J. Thomma, The battle for chitin recognition
1060   in plant-microbe interactions. *FEMS Microbiology Reviews* **39**, 171-183 (2015).

1061   35.   T. Shimizu *et al.*, Two LysM receptor molecules, CEBiP and OsCERK1, cooperatively
1062   regulate chitin elicitor signaling in rice. *The Plant Journal* **64**, 204-214 (2010).

1063   36.   R. Mostafa *et al.*, Recombination of standing variation in a multi-hybrid swarm drove
1064   adaptive radiation in a fungal pathogen and gave rise to two pandemic plant diseases.
1065   *bioRxiv* 10.1101/2021.11.24.469688, 2021.2011.2024.469688 (2021).

1066   37.   D. Pereira, U. Oggenfuss, B. A. McDonald, D. Croll, Population genomics of
1067   transposable element activation in the highly repressive genome of an agricultural
1068   pathogen. *Microb Genom* **7** (2021).

1069   38.   J. H. R. Maidment *et al.*, Multiple variants of the fungal effector AVR-Pik bind the
1070   HMA domain of the rice protein OsHIPP19, providing a foundation to engineer plant
1071   defense. *Journal of Biological Chemistry* **296** (2021).

1072   39.   N. Mukhi *et al.*, Perception of structurally distinct effectors by the integrated WRKY
1073   domain of a plant immune receptor. *Proc Natl Acad Sci U S A* **118** (2021).

1074   40.   W. Zschiesche *et al.*, The zinc-binding nuclear protein HIPP3 acts as an upstream
1075   regulator of the salicylate-dependent plant immunity pathway and of flowering time in
1076   *Arabidopsis thaliana*. *New Phytologist* **207**, 1084-1096 (2015).

1077   41.   O. Barth, S. Vogt, R. Uhlemann, W. Zschiesche, K. Humbeck, Stress induced and
1078   nuclear localized HIPP26 from *Arabidopsis thaliana* interacts via its heavy metal
1079   associated domain with the drought stress related zinc finger transcription factor
1080   ATHB29. *Plant Molecular Biology* **69**, 213-226 (2009).

1081   42.   H. Zhang *et al.*, Characterization of the Heavy-Metal-Associated Isoprenylated Plant
1082   Protein (HIPP) Gene Family from *Triticeae* Species. *Int J Mol Sci* **21** (2020).

1083   43.   Z. K. Barr, T. Werner, J. Tilsner, Heavy Metal-Associated Isoprenylated Plant Proteins
1084   (HIPPs) at Plasmodesmata: Exploring the Link between Localization and Function.
1085   http://dx.doi.org/10.3390/plants12163015.

1086   44.   G. H. Cowan *et al.*, Potato Mop-Top Virus Co-Opts the Stress Sensor HIPP26 for Long-
1087   Distance Movement. *Plant Physiol* **176**, 2052-2070 (2018).

1088   45.   J. B. de Abreu-Neto, A. C. Turchetto-Zolet, L. F. V. de Oliveira, M. H. Bodanese
1089   Zanettini, M. Margis-Pinheiro, Heavy metal-associated isoprenylated plant protein
1090   (HIPP): characterization of a family of proteins exclusive to plants. *The FEBS Journal*
1091   **280**, 1604-1616 (2013).

46. Z. Wang *et al.*, Isoprenylation modification is required for HIPP1-mediated powdery
mildew resistance in wheat. *Plant Cell Environ* **46**, 288-305 (2023).

47. Q. M. Imran *et al.*, Nitric Oxide Responsive Heavy Metal-Associated Gene AtHMAD1
Contributes to Development and Disease Resistance in *Arabidopsis thaliana*. *Front
Plant Sci* **7**, 1712 (2016).

48. Z. S. Radakovic *et al.*, Arabidopsis HIPP27 is a host susceptibility gene for the beet
cyst nematode *Heterodera schachtii*. *Mol Plant Pathol* **19**, 1917-1928 (2018).

49. T. Guo *et al.*, Arabidopsis HIPP proteins regulate endoplasmic reticulum-associated
degradation of CKX proteins and cytokinin responses. *Mol Plant* **14**, 1918-1934
(2021).

50. R. O'Lexy *et al.*, Exposure to heavy metal stress triggers changes in plasmodesmatal
permeability via deposition and breakdown of callose. *Journal of Experimental Botany*
69, 3715-3728 (2018).

51. Z. Wang *et al.*, Isoprenylation modification is required for HIPP1-mediated powdery
mildew resistance in wheat. *Plant, Cell & Environment* **46**, 288-305 (2023).

52. N. J. Talbot, D. J. Ebbole, J. E. Hamer, Identification and characterization of MPG1, a
gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant
Cell* **5**, 1575-1590 (1993).

53. B. Valent, L. Farrall, F. G. Chumley, *Magnaporthe grisea* Genes for Pathogenicity and
Virulence Identified through a Series of Backcrosses. *Genetics* **127**, 87-101 (1991).

54. H. Sakai *et al.*, Rice Annotation Project Database (RAP-DB): an integrative and
interactive database for rice genomics. *Plant Cell Physiol* **54**, e6 (2013).

55. R. A. Dean *et al.*, The genome sequence of the rice blast fungus *Magnaporthe grisea*.
Nature **434**, 980-986 (2005).

56. D. Kim *et al.*, TopHat2: accurate alignment of transcriptomes in the presence of
insertions, deletions and gene fusions. *Genome Biol* **14**, R36 (2013).

57. S. Anders, W. Huber, Differential expression analysis for sequence count data. *Genome
Biol* **11**, R106 (2010).

58. E. Southern, Southern blotting. *Nat Protoc* **1**, 518-525 (2006).

59. F. Sambrook J, E.F., and Maniatis, T Molecular Cloning - a laboratory manual, second
edition. . *Cold Spring Harbour Laboratory press* (1989).

60. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).

61. M. Holsters *et al.*, The functional organization of the nopaline *A. tumefaciens* plasmid
pTiC58. *Plasmid* **3**, 212-230 (1980).

62.   G. Hensel, C. Kastner, S. Oleszczuk, J. Riechen, J. Kumlehn, Agrobacterium-mediated
1128   gene transfer to cereal crop plants: current protocols for barley, wheat, triticale, and
1129   maize. *Int J Plant Genomics* **2009**, 835608 (2009).

1130   63.   B. WN, Burnette WN. "Western Blotting": electrophoretic transfer of proteins from
1131   sodium dodecyl sulfate-polyacrylamide gels to unmodified nitrocellulose and
1132   radiographic detection with antibody and radioiodinated protein. *Analytical*
1133   *Biochemistry* **112**, 195 - 203 (1981).

1134   64.   H. Li *et al.*, Pathogen protein modularity enables elaborate mimicry of a host
1135   phosphatase. *Cell* **186**, 3196-3207.e3117 (2023).

1136