Resistance patterns in drug-adapted cancer cell lines reflect the complex evolution in clinical tumours

Helen E. Grimsley[1,2], Magdalena Antczak[1,3], Ian G. Reddin[4], Katie-May McLaughlin[1], Andrea Nist[5], Marco Mernberger[6], Thorsten Stiewe[5,6], Tim R. Fenton[4], Daniel Speidel[7,8], Catherine Harper-Wynne[9], Karina Cox[10], Jindrich Cinatl jr.[11], Mark N. Wass[1]*, Michelle D. Garrett[1]*, Martin Michaelis[1,11]*

[1]School of Biosciences, Stacey Building, University of Kent, Canterbury, Kent, CT2 7NJ, UK

[2]Department of Radiation Oncology and the Molecular Biology Program, Memorial Sloan Kettering Cancer Center, New York, United States

[3]Current address: Queensland Cyber Infrastructure Foundation Ltd., Facility for Advanced Bioinformatics, Brisbane, QLD, 4072, Australia

[4]Cancer Sciences, Faculty of Medicine, University of Southampton, Southampton SO16 6YD, UK

[5]Genomics Core Facility, Philipps University, 35043 Marburg, Germany

[6]Institute of Molecular Oncology, Member of the German Center for Lung Research (DZL), Philipps University, 35032 Marburg, Germany

[7]Children's Medical Research Institute, The University of Sydney, 214 Hawkesbury Road, Westmead, New South Wales, Australia

[8]Current address: Breakpoint Therapeutics GmbH, Essener Bogen 7, 22419 Hamburg, Germany

[9]Kent Oncology Centre, Maidstone and Tunbridge Wells NHS Trust, Hermitage Lane, Maidstone, ME16 9QQ, UK

[10]Peggy Wood Breast Unit, Maidstone Hospital, Hermitage Lane, Maidstone, Kent,

ME16 9QQ, UK

[11]Dr Petra Joh-Research Institute, 60528 Frankfurt am Main, Germany

*Corresponding authors' emails: M.Michaelis@kent.ac.uk; M.D.Garrett@kent.ac.uk;

M.N.Wass@kent.ac.uk

Abstract

Background: Here, we introduce a novel set of triple-negative breast cancer

(TNBC) cell lines consisting of MDA-MB-468, HCC38, and HCC1806 and their

sublines adapted to cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, or 5-

fluorouracil.

Methods: The cell lines were characterized by whole exome sequencing and the

determination of drug-response profiles. Moreover, genes harbouring resistance-

associated mutations were investigated using TCGA data for potential clinical

relevance.

Results: Sequencing combined with TCGA-derived patient data resulted in the

identification of 682 biomarker candidates in the pan-cancer analysis. Thirty-five

genes were considered the most promising candidates because they harboured

resistance-associated variants in at least two resistant sublines, and their expression

correlated with TNBC patient survival. Exome sequencing and response profiles to

cytotoxic drugs and DNA damage response inhibitors identified revealed remarkably

little overlap between the resistant sublines, suggesting that each resistance

formation process follows a unique route. All of the drug-resistant TNBC sublines

remained sensitive or even displayed collateral sensitivity to a range of tested

compounds. Cross-resistance levels were lowest for the CHK2 inhibitor CCT241533,

the PLK1 inhibitor SBE13, and the RAD51 recombinase inhibitor B02, suggesting

that CHK2, PLK1, and RAD51 are potential drug targets for therapy-refractory

TNBC.

Conclusions: We present novel preclinical models of acquired drug resistance in

TNBC and many novel candidate biomarkers for further investigation. The finding

that each cancer cell line adaptation process follows an unpredictable route reflects

recent findings on cancer cell evolution in patients, supporting the relevance of drug-

adapted cancer cell lines as preclinical models of acquired resistance.

Key words

Triple Negative Breast Cancer, acquired drug resistance, exome sequencing DNA

repair, *de novo* variants, TCGA

Introduction

Triple-negative breast cancer (TNBC) is characterized by the absence of estrogen, progesterone, and HER2 receptors [1]. TNBC is responsible for approximately 15% of breast cancer cases and is associated with a poorer prognosis than hormone receptor- or HER2-positive breast cancers [1,2]. Current TNBC therapies are largely based on cytotoxic anticancer drugs, including platinum drugs, anthracyclins, eribulin, gemcitabine, paclitaxel, and 5-fluorouracil [1]. TNBC often responds well initially to cytotoxic chemotherapy, but recurrence and resistance are common, eventually leading to therapy failure. This combination of an initial high response rate followed by rapid resistance is referred to as the 'TNBC paradox' [1,3]. To improve TNBC therapy outcomes, new treatment approaches are needed, particularly those that are effective against treatment-refractory disease characterized by acquired resistance to cytotoxic chemotherapy.

In contrast to intrinsic drug resistance (which occurs independently of therapy and is a consequence of pre-existing often stochastic events in cancer cells), acquired resistance is the direct consequence of selection and adaptation processes caused by cancer treatment (directed tumor evolution) [4–8]. Understanding acquired resistance mechanisms is essential for optimizing cancer treatment for patients with therapy-refractory tumors.

Drug-adapted cancer cell lines are preclinical models that have been shown to reflect clinically relevant acquired drug resistance mechanisms in numerous studies [4,9–17]. Furthermore, drug-adapted cell lines enable detailed functional and systems-level studies that are not possible using clinical samples [4].

Here, we introduce a novel set of three parental TNBC cell lines and their 15 sublines adapted to cisplatin, doxorubicin, eribulin, gemcitabine, paclitaxel, or 5-

fluorouracil. These cell lines were characterized by whole exome sequencing and the

determination of response profiles to cytotoxic anti-cancer drugs and a panel of DNA

damage repair inhibitors. The resulting data showed that each resistance formation

process follows an individual and unpredictable route. The combined analysis of

resistance-associated mutations in combination with patient data from The Cancer

Genome Atlas (TCGA) [18] identified 35 novel candidate resistance biomarkers for

further investigation.

Results

Project cell line panel

Here, we characterized a cell line panel consisting of the parental TNBC cell

lines MDA-MB-468, HCC38, and HCC1806 and their sublines adapted to grow in the

presence of cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, or 5-fluorouracil,

which are all drugs used for the treatment of TNBC (Fig. 1A, Suppl. File 1) [19–25]. The

drug-resistant sublines were established by continuous exposure to stepwise

increasing drug concentrations as previously described [16]. All parental cell lines were

initially sensitive to therapeutic concentrations of the respective drugs, as indicated

by IC_{50} (concentration that reduces cell viability by 50%) values within the range of

clinical drug plasma concentrations (C_{max}) (Suppl. Fig. 1, Suppl. File 1) [26]. The

relative resistance factors (IC_{50} drug-adapted subline/ IC_{50} respective parental cell

line) ranged from 5.5-fold (HCC38^{r}PCL$^{2.5}$) to 5916.7-fold (HCC1806^{r}ERI^{50}) (Fig. 1B,

Suppl. File 1).

Characterization of the cell line panel by whole exome sequencing

The cell line panel was investigated by whole exome sequencing. Among the

identified variants, missense variants were most common, followed by synonymous

variants (Suppl. Fig. 2A). Insertions/deletions (INDELs), frameshift mutations, stop-

gain, stop-loss, and splice variants were identified at lower frequencies (Suppl. Fig.

2A). Between 217 (HCC38^{r}DOX^{40}) and 952 (HCC38^{r}GEM^{20}) variants differed in the

drug-adapted sublines relative to the respective parental cell lines (Suppl. Fig. 2B).

We grouped the resistance-associated variants into five categories (Fig. 2A,

see methods): 1. *Gained variants*, variants only called in the drug-adapted subline

but detectable at low confidence in the respective parental cell line; 2. *De novo*

variants, variants called in the drug-adapted subline but undetectable in the

respective parental cell line; 3. *Not-called variants*, variants only called in the

parental cell line but detectable with low confidence in the resistant subline; 4. *Lost*

variants; variants called in the parental cell line but undetectable in the drug-adapted

subline; and 5. *Shared variants*; variants called in both the parental and the

respective drug-adapted sublines (Fig. 2A).

The number of *gained* variants ranged from 44 (HCC38[r]DOX[40]) to 381

(HCC38[r]GEM[20]), the number of *de novo* variants ranged from 31 (HCC38[r]DOX[40]) to

225 (MDA-MB-468[r]PCL[20]), the number of *not-called* variants ranged from 88

(HCC38[r]GEM[20] and HCC1806[r]DOX[12.5]) to 345 (MDA-MB-468[r]PCL[20]), and the

number of *lost* variants ranged from 129 (HCC38[r]GEM[20]) to 398 (MDA-MB-

468[r]PCL[20]) (Fig. 2B, Fig. 2C, Suppl. File.2 and 3). The number of *shared* variants

that were both called in the parental cell lines and their sublines ranged from 128

(MDA-MB-468[r]PCL[20]) to 368 (HCC38[r]GEM[20]) (Fig. 2D, Suppl. File 2 and 3). The

number of *shared* variants that increased by at least two-fold in the resistant sub-

lines vs. the respective parental ranged from four (HCC1806[r]5-F[1500]) to 21 (MDA-

MB-468[r]CDDP[1000]), whilst the number of shared variants that decreased by at least

two-fold ranged from two (MDA-MB-468[r]PCL[20]) to 24 (HCC38[r]GEM[20]) (Fig. 2E,

Suppl. File 2).

Analysis of the distribution of *de novo* variants

To identify variants that may have a functional role in drug resistance, we

initially considered the 81 genes that harbored *de novo* variants in at least two

different sublines from more than one parental cell line (Fig. 3A, Suppl. File 4). This

list included 46 genes that have already been reported to be involved in drug

resistance in cancer and 33 new candidate genes with a possible role in drug

resistance (Fig. 3A, Suppl. File 4). Notably, 24 of the 33 new candidate genes are

reported to be relevant in cancer (Fig. 3A, Suppl. File 4).

Four of the five genes with the greatest number of *de novo* variants in the

drug-adapted sublines were mucin (MUC) genes. *MUC6* had *de novo* variants in 15

sublines, *MUC2* in 14 sublines, *MUC4* in 13 sublines, and MUC16 in nine sublines

(Fig. 3A, Suppl. File 4). The MUC genes are large genes that are known to be

commonly mutated in cancer and have been reported to be involved in cancer cell

drug resistance [27–31]. *De novo* mutations in CDC27, which has also been linked to

drug resistance in cancer, were also detected in nine drug-resistant sublines [32,33]

(Fig. 3A, Suppl. File 4).

GXYLT1, *KRTAP4-11*, and *RGPD4* were amongst those genes, which had

not previously been associated with drug resistance in cancer that displayed *de novo*

mutations in a high number (7) of drug-resistant sublines (Fig. 3A, Suppl. File 4). A

GXYLT1 mutation promoted metastasis in colorectal cancer through MAPK

signalling, a pathway known to confer resistance to a range of anti-cancer drugs [34–

37]. *RGPD4* mutations are correlated with vascular invasion in HBV-associated

hepatocellular carcinoma, and it is known that there is an overlap between pro-

angiogenic, pro-metastatic, and resistance-associated signalling in cancer [35,38].

There is no known link between KRTAP4-11 and cancer, but *KRTAP4-11* expression

levels have been reported to predict the methotrexate response in rheumatoid

arthritis patients [38]. Hence, it seems plausible that the products of these genes may

be involved in cancer cell drug resistance.

Taken together, our analysis identified 48 genes known to be involved in

cancer cell drug resistance alongside 33 novel candidates potentially contributing to

therapy failure. Further research will be required to characterize the roles of these

individual genes in detail.

When we compared the overlaps between *de novo* variants shared between

sublines adapted to the same drug, the numbers were too small to draw any

meaningful conclusions (Fig. 3B, Suppl. Fig. 3A).

Notably, *de novo* variants in drug-resistant sublines may not always represent

actual novel variants that are selected because they contribute to cancer cell

resistance. Many apparent *de novo* mutations may have already been present in a

small fraction of the cells of the parental cell line but may not have been detected

due to the sequencing depth. Hence, overlaps in *de novo* variants between sublines

of the same parental cell line can also be used to indicate the levels of relatedness

between the founding subpopulations of the different resistant sublines.

Analysis of the *de novo* variants shared between the sublines from the same

parental cell line indicated the largest overlap. On average, there was a 22.6%

overlap among the HCC1806 sublines, followed by a 15.0% overlap among the

HCC38 sublines and a 7.7% overlap among the MDA-MB-468 sublines (Fig. 3C).

However, there were also noticeable differences in the overlaps between *de novo*

variants identified in each of the sublines from the same parental cell line. For

example, only three *de novo* variants were shared between HCC38[r]CDDP[3000] (out of

98 in total, 3.1%) and HCC38[r]PCL[2.5] (out of 92 in total, 3.3%), while 53 variants were

shared between HCC38[r]ERI[10] (out of 131 in total, 40.5%) and HCC38[r]GEM[20] (out of

203 in total, 26.1%) (Fig. 3C, Suppl. Fig. 3B). These numbers suggest that there are

no pre-existing cell line subpopulations that are consistently selected in response to

anti-cancer drug treatment.

Protein functions related to variants that changed in drug-resistant sublines

Next, we used the Gene Ontology (GO) annotation to perform an analysis of

the protein functions associated with genes present in the *de novo*, *gained*, *not*

called, and *lost* variant sets as well as *shared* variants with a two-fold increase or

decrease in allele frequency (Suppl. Fig 4A, B).

There was limited overlap between the GO terms for the variants detected in

the sublines adapted to the same drug (Suppl. Fig. 4C, E). The extracellular matrix-

related GO terms 'extracellular matrix constituent lubricant activity', 'extracellular

matrix', and 'maintenance of gastrointestinal epithelium' were most common, which

reflects the high number of variants observed in the mucin genes (Suppl. Fig. 4C, E).

GO term analysis of the sublines from the same parental cell line revealed

very similar results, again revealing an overrepresentation of extracellular matrix-

related GO terms (Suppl. Fig. 4D, F). Further research will be required to investigate

the potential role of mucins and the extracellular matrix in acquired drug resistance in

TNBC cells.

Potential clinical relevance of selected variants

The potential clinical relevance of genes harboring *de novo*, *gained*, and

shared variants with a two-fold increase in allele frequency in the resistant subline as

well as genes harboring truncating variants was analysed using patient data derived

from The Cancer Genome Atlas (TCGA) [39]. Notably, there were only data available

from patients treated with cisplatin, doxorubicin, gemcitabine, paclitaxel, and/or 5-

fluorouracil, but no data on eribulin treatment were available.

We performed two analyses, one pan-cancer analysis, in which we

considered all patient survival data available for the drugs, and a second analysis, in

which we considered TNBC patients and for which only doxorubicin and paclitaxel

data were available (Fig. 4A). The pan-cancer analysis included data from 29 TCGA

cancer types for which mutation status and gene expression data were available (Fig

4B).

Six cases with at least two mutations in a resistance-associated gene were

associated with patient prognosis (Suppl. File 5). As the number of resistance-

associated genes with mutations in patients was low, we also considered gene

expression status associations with prognosis. For 1,018 cases there was a

significant association between gene expression and patient prognosis. This

included genes, whose products were known to play a role in cancer cell drug

resistance, such as CHEK2 [40–42] and APC [43–45] (Fig. 4C, Suppl. File 5). Moreover,

we also identified novel candidates, which had not previously been suggested to be

involved in cancer cell drug resistance, including KIAA2018, EYS, NBPF10, and

KIAA0586 (Fig. 4C, Suppl. File 5).

We further determined the association of the expression of genes harboring

de novo, *gained*, and *shared* variants with a two-fold increase as well as genes

harboring truncating variants with patient survival in response to treatment with

cisplatin, doxorubicin, gemcitabine, paclitaxel, and 5-fluorouracil (Fig. 4 D-E, Suppl.

File 5). In total, the expression of 682 genes was significantly correlated with patient

survival in response to at least one drug in the pan-cancer data. For 513 of these

682 genes, gene expression was associated with tumor response to the drug of the

respective resistant subline (Suppl. File 5). The expression of 91 genes was

associated with patient response to two drugs, the expression of 51 genes

associated with response to three drugs, the expression of 21 genes associated with

expression to four drugs, and the expression of 6 genes associated with response to

all five drugs (Fig. 4D, Suppl. File 5).

Considering the TNBC data alone, the expression of 165 genes was

significantly correlated with patient survival in response to either doxorubicin,

paclitaxel, or both drugs (Fig. 4E, Suppl. File 5). The expression of 141 of these 165

genes was associated with tumor response to the drug of the respective drug-

adapted subline. The expression of 22 genes was associated with patient response

to both doxorubicin and paclitaxel (Fig. 4E, Suppl. File 5).

Comparison of the analysis of the 165 genes identified in the TCGA analysis

with the 81 genes identified in the analysis of *de novo* variants (Suppl. File 4)

revealed 35 overlapping genes present in both datasets. This included 23 genes that

have already been associated with drug resistance and 12 genes (*ABCD1, AGAP6,*

CUBN, DNAJC13, FLG, GXYLT1, KIAA0586, PABPC3, RGPD3, RGPD4, SETX and

USP6) that are novel findings (Suppl. File 6).

Complex sensitivity patterns of drug-resistant sublines to cytotoxic drugs

Determining drug sensitivity profiles in the cell line panel against the drugs of

adaptation, i.e., cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, and 5-

fluorouracil (Fig. 5A, Suppl. File 1), revealed complex resistance patterns that did not

follow clear, predictable rules. For example, two of the three doxorubicin-adapted

sublines (HCC38^{r}DOX^{40} and HCC1806^{r}DOX$^{12.5}$) displayed increased (collateral)

sensitivity to cisplatin compared to the parental cell line, while MDA-MB-468^{r}DOX^{50}

displayed cross-resistance to cisplatin (Fig. 5A, Suppl. File 1). Moreover, all resistant

sublines remained sensitive to or showed collateral sensitivity to at least one of the

other chemotherapeutic agents (Fig. 5A, Suppl. File 1). The 5-fluorouracil-resistant

HCC1806^{r}5-F^{1500} subline was the only resistant subline that remained sensitive to all

other investigated cytotoxic drugs (Fig. 5A, Suppl. File 1).

The ATP-binding cassette (ABC) transporter ABCB1 (also known as P-

glycoprotein and MDR1) is an efflux transporter that mediates resistance to many

anti-cancer drugs, including doxorubicin, eribulin, and paclitaxel [46]. Only five of the

nine sublines adapted to the ABCB1 substrates doxorubicin, eribulin, and paclitaxel

(including all three eribulin-resistant sublines) displayed cross-resistance to all other

ABCB1 substrates. Among the ABCB1 substrate-adapted sublines, all the eribulin-

adapted sublines displayed cross-resistance to paclitaxel, and all the paclitaxel-

adapted sublines displayed cross-resistance to eribulin (Fig. 5A, Suppl. File 1).

Notably, eribulin and paclitaxel are both tubulin-binding agents but differ in their

mechanisms of interaction with tubulin. Eribulin is a destabilizing agent that binds to

the vinca binding site of tubulin and inhibits microtubule formation, while paclitaxel is

a stabilizing agent that binds to the taxane binding site that impairs microtubule

degradation [47–51]. Further research will be required to determine to what extent the

tubulin-binding agent cross-resistance profile of the tubulin-binding agent-adapted

sublines is the consequence of the expression of ABCB1 (and/or other transporters),

tubulin-related resistance mechanisms, or both.

Taken together, it is not possible to predict how resistance to a certain drug

will affect the sensitivity patterns of the resulting sublines to other cytotoxic agents.

However, all of the drug-resistant TNBC sublines remained sensitive and/or

displayed collateral sensitivity to at least one of the tested anti-cancer drugs. Future

research will be needed to elucidate the underlying mechanisms to identify

biomarkers for personalized therapy approaches that can guide effective drugs to the

right patients [4].

Complex sensitivity patterns of drug-resistant sublines to DNA damage

response (DDR) inhibitors

Triple-negative breast cancer cells have been shown to harbor defects in DNA

damage repair signalling, which can result in a dependence on the remaining intact

DNA damage repair (DDR) pathways and, in turn, in sensitivity to certain DDR

inhibitors [52]. Hence, we tested a panel of inhibitors targeting critical nodes of DDR

signalling in our novel resistant TNBC cell line panel (Fig. 5B).

All parental cell lines displayed sensitivity to the tested DDR inhibitors at

therapeutic concentrations, i.e., within the C_{max} values reported for these agents (for

which this information was available) (Suppl. Fig. 7). Similar to the results obtained

for the cytotoxic anti-cancer drugs, the DDR sensitivity profiles were complex and

unpredictable in the resistant sublines (Fig. 5C, Suppl. File 1). Relative to the

respective parental cell lines, the sensitivity remained unchanged for 128 DDR

inhibitor/ resistant subline combinations. Increased resistance (cross-resistance) was

detected in 96 DDR inhibitor/resistant subline combinations, and increased

sensitivity (collateral vulnerability) was recorded in 16 DDR inhibitor/resistant subline

combinations. Neither sublines of the same parental cell line nor sublines adapted to

the same drugs displayed substantial overlap in their DDR inhibitor sensitivity

profiles. Generally, cross-resistance levels were lowest for the CHK2 inhibitor

CCT241533, the PLK1 inhibitor SBE13, and the RAD51 recombinase inhibitor B02

among the investigated DDR inhibitors (Fig. 5C, Suppl. File 1).

Cross-resistance patterns were even inconsistent between DDR inhibitors

with the same targets. For example, different sensitivity patterns were observed

between the ATR inhibitors ceralasertib and berzosertib as well as the CHK1

inhibitors rabusertib, MK-8776, SRA737, and prexasertib (Fig. 5C, Suppl. File 1). The

reasons for these differences are unclear. Notably, the activity of the DDR inhibitors

may be modified by interactions with additional targets, and off-target resistance

mechanisms (e.g., processes associated with drug uptake or efflux) may contribute

to these differences [53].

In summary, and in line with the findings from the investigation of cytotoxic

anti-cancer drugs, the drug-adapted TNBC sublines displayed complex,

unpredictable sensitivity patterns against DDR inhibitors. This further demonstrates

that improved future therapies will depend on an advanced understanding of the

underlying molecular processes that enable the identification of biomarkers that can

guide effective therapies for individual patients after treatment failure[4]. Notably,

CHK2, PLK1, and RAD51 may have potential as new drug targets for the discovery

and development of next-line therapies for TNBC patients whose tumors have

stopped responding to chemotherapy.

Investigation of patterns in cell line drug response profiles

Finally, we used the delta (Δ) method to identify potential patterns in the

response of the cell lines to all investigated cytotoxic anti-cancer drugs and DDR

inhibitors [54]. The IC_{50} values were transformed to ΔIC_{50} values for each compound

(see methods) and correlated across the drug panel using linear regression analysis

and testing for statistical significance (Suppl. Table 1). Positive correlations indicate

that increased drug resistance is seen with both agents, whilst negative correlations

indicate that whilst increasing drug resistance is observed for one agent, collateral

sensitivity is observed for the other agent. In the MDA-MB-468, HCC38, and

HCC1806 sublines, we observed 19, 20, and 60 positive correlations and 2, 8, and 1

negative correlation, respectively (Suppl. Table 1).

We were most interested in the agents that demonstrated negative

correlations, as they may identify potential next-line treatments. However, among the

11 negative correlations, there were no consistent results across the cell line panel

(Fig. 6). This further confirms that acquired resistance mechanisms are complex,

individual, and unpredictable and that the identification of potential next-line

therapies after treatment failure will depend on an improved understanding of cancer

cell evolution enabling therapy monitoring and biomarker-guided treatment

adaptation.

Discussion

In this study, we introduced and characterized a novel set of 15 sublines

derived from the TNBC cell lines HCC38, HCC1806, and MDA-MB-468 that had

been adapted to cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, or 5-

fluorouracil.

We applied whole exome sequencing to identify biomarker candidates to

guide the use of anti-cancer therapies. In the first step, we focused on *de novo*

mutations, i.e., mutations found in a resistant subline but undetectable in the

respective parental cell line. Considering genes that displayed *de novo* mutations in

at least two sublines of two different parental cell lines resulted in 81 resistance-

associated variants, 48 of which were already known to be involved in cancer cell

drug resistance, while 33 variants were novel.

In a second approach, we used TCGA data to investigate the potential clinical

relevance of genes that harbored resistance-associated variants in the resistant

sublines. In the pan-cancer dataset, the expression of 682 of these genes was

correlated with patient survival in response to at least one of the investigated drugs.

Considering only TNBC, the expression of 165 genes was significantly correlated

with patient survival.

Comparison of the *de novo* variant analysis with the TNBC TCGA analysis

identified 35 overlapping genes. Twenty-three of these genes are known to be

associated with drug resistance. Twelve genes (*ABCD1, AGAP6, CUBN, DNAJC13,*

FLG, GXYLT1, KIAA0586, PABPC3, RGPD3, RGPD4, SETX and USP6) are novel

findings that may represent novel resistance biomarkers that have not been

previously associated with drug resistance in cancer. Further research will be

needed to investigate and define in more detail the role of these gene variants in

cancer therapy response and the expression of these genes as biomarkers for the

tailoring of personalized cancer therapies. Notably, numerous studies have shown

that drug-adapted cancer cell lines exhibit clinically relevant resistance mechanisms

[4,9–17].

Interestingly, the analysis of exome sequencing data revealed remarkably few

overlapping mutations between the investigated resistant sublines, including sublines

derived from the same parental cell line and sublines adapted to the same drug. This

suggests that resistance formation is the consequence of a complex, individual, and

unpredictable evolutionary process.

This complexity was confirmed by the determination of drug sensitivity profiles

to both cytotoxic anti-cancer drugs and DNA damage repair (DDR) inhibitors. Drug-

adapted sublines of the same parental cell line and sublines adapted to the same

drug displayed substantially different drug response patterns.

Notably, all the drug-adapted sublines remained sensitive and/or displayed

increased sensitivity (collateral vulnerability) to a range of tested compounds. This

suggests that it will be possible in the future to establish an improved understanding

of the processes underlying acquired resistance formation that result in the

identification of biomarkers that indicate effective next-line treatments for patients for

whom currently no effective treatment is available.

Among the investigated DDR inhibitors, the CHK2 inhibitor CCT241533, the

PLK1 inhibitor SBE13, and the RAD51 recombinase inhibitor B02 had the lowest

cross-resistance levels. Thus, CHK2, PLK1, and RAD51 are potential drug targets in

TNBC patients after failure of established therapies, particularly if reliable biomarkers

are found that identify cancer patients who are likely to benefit from such treatments.

Overall, the results from the characterization of the project cell line panel

indicated that cancer cell resistance is a complex, individual, and unpredictable

process. This finding is in agreement with data from studies in which cancer cell lines

were repeatedly adapted to the same drug in independent experiments [8,16,55–57] and

with recent findings from a comprehensive analysis of cancer cell evolution in lung

cancer patients [58–62].

In conclusion, we present a novel set of drug-adapted TNBC cell lines as

preclinical models of acquired drug resistance. Overlapping genes detected through

the characterization of *de novo* variants and patient-derived TCGA data identified 35

biomarker candidates for the guidance of personalized TNBC therapies for further

investigation, including 12 novel genes that have not been previously associated with

drug resistance in cancer. Finally, our results show that each cancer cell line

adaptation process follows an individual, unpredictable route, which reflects recent

clinical findings from the monitoring of cancer cell evolution in patients [58–62]. This

further supports the relevance of drug-adapted cancer cell lines as preclinical models

of acquired resistance that can be analysed and manipulated at a level of detail that

is impossible in the clinical setting.

Materials and Methods

Cell culture

MDA-MB-468, HCC38, and HCC1806 cells were obtained from the American Type Culture Collection (ATCC). The drug-adapted sublines (Fig. 1A, Suppl. File.1) were established by continuous exposure to stepwise increasing drug concentrations as previously described and derived from the Resistant Cancer Cell Line (RCCL) collection (https://research.kent.ac.uk/industrial-biotechnology-centre/the-resistant-cancer-cell-line-rccl-collection)[4,63]. All cell lines were cultured in Iscove's Modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (Sigma☐Aldrich, Germany), 2 mM L-glutamine, 25 mM HEPES (Fisher Scientific, UK), 100 IU/mL penicillin, and 100 µg/mL streptomycin (Life Technologies, UK) at 37 °C in a humidified atmosphere with 5% CO_2. Each drug-adapted subline was continuously cultured in the presence of the specific adaptation drug at a defined concentration, as indicated by the cell line name (ng/mL), e.g., MDA-MB-468[r]DOX[50], where r = the resistant subline, Dox = doxorubicin and 50 = 50 ng/ml.

Compounds

The following compounds were purchased from the indicated suppliers: Adavosertib, Alisertib, Berzosertib, Ceralasertib, MK-8776, Olaparib, Prexasertib, Rabusertib, Rucaparib, SBE13, Tozasertib (Adooq Bioscience), AZD0156, BI2536, Doxorubicin, Gemcitabine (Selleckchem), B02, Cisplatin, 5-Fluorouracil (Sigma☐Aldrich), CCT241533, SRA737 (a gift from the Institute of Cancer Research), Eribulin (Eisia), and Paclitaxel (Cayman Chemicals). All drug stocks were prepared in DMSO and stored at -20 °C, except for cisplatin, which was prepared in 0.9% NaCl solution and stored in the dark at room temperature.

Cell growth and viability assays

Cell viability was tested using the 3-(4,5-dimethylthiazol-2-yl)-2,5-

diphenyltetrazolium bromide (MTT) dye reduction assay after 120 hours of

incubation with each compound, modified as previously described [64,65].

Concentrations that reduced cell viability by 50% relative to an untreated control

(IC_{50}) were determined and used to calculate the resistance factor (RF; IC_{50} of drug-

adapted cell line/IC_{50} of respective parental cell line).

Whole exome sequencing

Whole exome sequencing (WES) libraries were prepared using the Nextera

Rapid Capture Exome Kit (Illumina). Sequencing was performed on a HiSeq 1500

platform in Rapid Run mode with 2 x 100 nucleotide paired-end reads. The two lanes

of the Rapid Run flow cell provided two sets of FASTQ data per cell line.

Variant calling

FASTQC was used to control the quality of the raw sequence data [66] prior to

the removal of sequencing adaptors. Trimmomatic (settings: NexteraPE-

PE.fa:2:30:10 LEADING:3 TRAILING:3 SLIDING WINDOW: 4:15 MILEN:36) [67]. Raw

FASTQ files were aligned to the human reference genome (GRCH37) using

Burrows☐Wheeler Alignment (v.0.7.17) with an output in sequence alignment map

(SAM) format applying the default settings -M -R [68–70]. Only paired reads were used,

and Samtools flagstat was used to print statistics throughout each of the subsequent

steps [68]. SAM files were input into Picard tools SortSam (v.2.17.10), where the read

alignments were sorted by coordinate and converted to a binary alignment map

(BAM) format (Picard Toolkit.2019. Broad Institute, GitHub Repository.

http://broadinstitute.github.io/picard/; Broad Institute). Picard Tools MarkDuplicates

(v2.17.10) was used for the removal of PCR duplicates (Picard Toolkit. 2019. Broad

Institute, GitHub Repository. http://broadinstitute.github.io/picard/; Broad Institute).

GenomeAnalysisTK-3.7.0 RealignerTargetCreator was used to perform base score

recalibration, and GenomeAnalysisTK-3.7.0 IndelRealigner was used for INDEL

realignment MAX_READS = 20000 [71]. SAMtools mpileup was used to generate

binary variant call format (BCF) files from the BAM files, which were then input into

BCFtools to call the SNVs and INDELS to generate a variant calling format (VCF) [72].

Variants were annotated with VEP [73].

Variant filtering

Only variants in coding regions of the genome were considered. To identify

high-confidence variants, variants with a Phred score < 30, variants with fewer than

10 reads supporting the base call, or variants with < 3 reads supporting the variant

were removed. Moreover, common variants with a frequency of ≥ 0.001% in the

genome aggregation database (gnomAD) were removed [74]; if not, ≥ 3 samples were

annotated in The Cancer Genome Atlas (TCGA), or ≥ 10 samples were annotated in

the Catalogue Of Somatic Mutations In Cancer (COSMIC) [39,75,76].

Definition of variants

Gained variants: variants that are called in the drug-resistant subline and are

called with low confidence in the parental cell line. *De novo* variants: variants that are

called in the drug-resistant subline but not called in the parental cell line. *Not called*

variants: variants that are called in the parental cell line but not called in the drug-

resistant subline, even at low confidence. *Lost* variants: variants that are called in the

parental cell line and are called in low confidence in the drug-resistant subline.

Shared variants: variants that are called in both the parental and drug-resistant

sublines.

Gene Ontology

Gene Ontology (GO) functional enrichment analysis was conducted using

G:profiler [77]. Gene lists were submitted as queries to the g:GOSt functional profiling

tool and run at a significance threshold of g:SCS and a user threshold of 0.05.

The Cancer Genome Atlas (TCGA) analysis

TCGA data retrieval

The data were collected from the UCSC Xena functional genomics browser

[https://xenabrowser.net]. Batch-corrected gene expression data (RNAseq,

log2(normalized value + 1)) for 11,060 patients (version 2016-12-29), clinical data for

12,591 patients (version 2018-09-13), and somatic mutation data (HG19) for 9,104

patients (version 2016-12-29) were downloaded for the TCGA pancancer (PANCAN)

cohort. Curated drug data were obtained from Moiso 2021 for 4,321 patients [78].

Final datasets

The 4 downloaded datasets were filtered to a final dataset for each drug for

which every data type was available (gene expression, somatic mutation, clinical,

and drug data). If a patient did not have at least 1 somatic mutation recorded, they

were excluded from further somatic mutation analyses. This resulted in final datasets

of 683 patients (23 cancer types) treated with cisplatin, 385 (17) with doxorubicin,

367 (11) with fluorouracil, 349 (20) with gemcitabine, and 544 (16) with paclitaxel for

which somatic mutation and clinical data were available (table –

"mutations/treatment_by_cancer_type_mutation_patients.tsv"). The gene expression

and treatment data included 765 patients (24 cancer types) treated with cisplatin,

571 (18) with doxorubicin, 452 (11) with fluorouracil, 438 (21) with gemcitabine, and

828 (15) with paclitaxel (table –

"expression/treatment_by_cancer_type_expression_patients.tsv"). Datasets including

only TNBC patients were also created for further analysis[79]. This was only completed

for those patients treated with doxorubicin (96 patients) and paclitaxel (63), as the

number of patients treated with cisplatin (2), gemcitabine (4), and fluorouracil (21)

was too low for meaningful analysis. For doxorubicin and paclitaxel treatments, gene

expression and clinical data were available for 93 and 62 TNBC patients,

respectively, while somatic mutation and clinical data were available for 74

doxorubicin- and 49 paclitaxel-treated patients. One TNBC patient (TCGA-AR-A256)

whose disease-specific survival (DSS) data were incomplete was excluded.

Survival analysis

Analysis was performed in R version 4.3.0. Kaplan-Meier (KM) plots were

generated for mutation status (mutated – MUT or wild type – WT) and for gene

expression status (high or low) using the survival (v3.5-5) and survminer (0.4.9)

packages. Somatic nonsynonymous mutations were considered in the genes of

interest. The cut-off for high/low gene expression was calculated using the

surv_cutpoint function in survminer, which makes use of the R package maxstat

(v0.7-25). This gave a threshold for high/low expression based on the most

significant relation with outcome, in this case, disease-specific survival. Any sample

with gene expression > the calculated threshold was considered to have "high

expression", and any sample with gene expression < the threshold was considered

to have "low expression". The p value displayed on the KM plots was calculated

using the log-rank test.

Statistical analysis and data manipulation

GraphPad Prism 6 (GraphPad Software, Inc., USA) was used to generate dose☐response curves and determine IC_{50} values via nonlinear regression (with variable slopes). Statistical significance was calculated using a two-tailed t-test, assuming unequal variance, in GraphPad Prism 6 (GraphPad Software, Inc., USA).

The delta method was used as described by Bracht *et al.*, 2006 [54]. IC_{50} values were transformed to ΔIC_{50} values: ΔIC_{50} = *log (average* IC_{50} *in drug over all cell lines) – log (individual* IC_{50} *in drug for each cell line)*. Linear regression analysis of ΔIC_{50X} versus ΔIC_{50Y}, where X and Y represent two different compounds from the panel, was performed. The Pearson correlation coefficient (*r*) was used to establish the level of significance in a two-tailed test with (n-2) degrees of freedom, where $p \leq 0.05$.

Availability of data and materials

Data generated or analyzed during this study are included in this published article and its supplementary information files. The exome sequencing datasets generated and analyzed during the current study are available at Gene Expression Omnibus (GEO) repository (#PRJNA1155201).

Competing interests

Nothing to declare.

Funding

This work was supported by grants from the Frankfurter Stiftung für krebskranke Kinder, Kent Health, Kent Cancer Trust, and the Rosetrees Trust.

Authors contribution

HEG, MMi, MDG, and MNW conceived and designed the study. HG, MA, IGR, KMM, AN, MMe, and JC acquired data. All authors analyzed and interpreted data. HEG, MM, MDG, and MNW drafted the work. All authors substantively revised the work and approved the submitted version.

Acknowledgements

We would like to thank the Institute of Cancer Research for their kind gift of CCT241533 and SRA737. Figures were created using BioRender.com.

References

1. Gupta, G.K., Collier, A.L., Lee, D., Hoefer, R.A., Zheleva, V., van Reesema,
L.L.S., Tang-Tan, A.M., Guye, M.L., Chang, D.Z., Winston, J.S., et al. (2020).
Perspectives on triple-negative breast cancer: Current treatment strategies,
unmet needs, and potential targets for future therapies. Cancers (Basel). *12*,
1–33. 10.3390/cancers12092392.

2. Hampton, J.M., Song, J., Jayasekera, J., Schechter, C., Alagoz, O., Stout,
N.K., Trentham-dietz, A., Lee, S.J., Huang, X., Mandelblatt, J.S., et al. (2024).
Analysis of Breast Cancer Mortality in the US—1975 to 2019. Jama *5405*,
233–241. 10.1001/jama.2023.25881.

3. Fornier, M., and Fumoleau, P. (2012). The paradox of triple negative breast
cancer: Novel approaches to treatment. Breast J. *18*, 41–51. 10.1111/j.1524-
4741.2011.01175.x.

4. Michaelis, M., Wass, M.N., and Cinatl, J. (2019). Drug-adapted cancer cell
lines as preclinical models of acquired resistance. Cancer Drug Resist. *2*, 447–
456. 10.20517/cdr.2019.005.

5. Oellerich, T., Schneider, C., Thomas, D., Knecht, K.M., Buzovetsky, O.,
Kaderali, L., Schliemann, C., Bohnenberger, H., Angenendt, L., Hartmann, W.,
et al. (2019). Selective inactivation of hypomethylating agents by SAMHD1
provides a rationale for therapeutic stratification in AML. Nat. Commun. *10*.
10.1038/s41467-019-11413-4.

6. Santoni-Rugiu, E., Melchior, L.C., Urbanska, E.M., Jakobsen, J.N., De
Stricker, K., Grauslund, M., and Sørensen, J.B. (2019). Intrinsic resistance to
EGFR-tyrosine kinase inhibitors in EGFR-mutant non-small cell lung cancer:
Differences and similarities with acquired resistance. Cancers (Basel). *11*, 1–
57. 10.3390/cancers11070923.

7. Touat, M., Li, Y.Y., Boynton, A.N., Spurr, L.F., Iorgulescu, J.B., Bohrson, C.L.,
Cortes-Ciriano, I., Birzu, C., Geduldig, J.E., Pelton, K., et al. (2020).
Mechanisms and therapeutic implications of hypermutation in gliomas. Nature
580, 517–523. 10.1038/s41586-020-2209-9.

8. Rothenburger, T., Thomas, D., Schreiber, Y., Wratil, P.R., Pflantz, T., Knecht,
K., Digianantonio, K., Temple, J., Schneider, C., Baldauf, H.M., et al. (2021).
Differences between intrinsic and acquired nucleoside analogue resistance in
acute myeloid leukaemia cells. J. Exp. Clin. Cancer Res. *40*, 1–19.
10.1186/s13046-021-02093-4.

9. Juliano, R.L., and Ling, V. (1976). A surface glycoprotein modulating drug
permeability in Chinese hamster ovary cell mutants. BBA - Biomembr. *455*,
152–162. 10.1016/0005-2736(76)90160-7.

10.   Cole, S.P.C., Bhardwaj, G., Gerlach, J.H., Mackie, J.E., Grant, C.E., Almquist,
618   K.C., Stewart, A.J., Kurz, E.U., Duncan, A.M.V., and Deeley, R.G. (1992).
619   Overexpression of a transporter gene in a multidrug-resistant human lung
620   cancer cell line. Science (80-.). *258*, 1650–1654. 10.1126/science.1360704.

621   11.   Engelman, J., Zejnullahu, K., Mitsudomi, T., Song, Y., and Hyland (2007). MET
622   amplification leads to gefitinib resistance in lung cancer by activating ERBB3
623   signaling. Science (80-.). *316*, 1039–1043.

624   12.   Sharma, S. V., Haber, D.A., and Settleman, J. (2010). Cell line-based
625   platforms to evaluate the therapeutic efficacy of candidate anticancer agents.
626   Nat. Rev. Cancer *10*, 241–253. 10.1038/nrc2820.

627   13.   Nazarian, R., Shi, H., Wang, Q., Kong, X., Koya, R.C., Lee, H., Chen, Z., Lee,
628   M.K., Attar, N., Sazegar, H., et al. (2010). Melanomas acquire resistance to B-
629   RAF(V600E) inhibition by RTK or N-RAS upregulation. Nature *468*, 973–977.
630   10.1038/nature09626.

631   14.   Crystal, A.S., Shaw, A.T., Sequist, L. V., Friboulet, L., Niederst, M.J.,
632   Lockerman, E.L., Frias, R.L., Gainor, J.F., Amzallag, A., Greninger, P., et al.
633   (2014). Patient-derived models of acquired resistance can identify effective
634   drug combinations for cancer. Science (80-.). *346*, 1480–1486.
635   10.1126/science.1254721.

636   15.   Schneider, C., Oellerich, T., Baldauf, H.M., Schwarz, S.M., Thomas, D., Flick,
637   R., Bohnenberger, H., Kaderali, L., Stegmann, L., Cremer, A., et al. (2017).
638   SAMHD1 is a biomarker for cytarabine response and a therapeutic target in
639   acute myeloid leukemia. Nat. Med. *23*, 250–255. 10.1038/nm.4255.

640   16.   Michaelis, M., Rothweiler, F., Barth, S., Cinat, J., Van Rikxoort, M.,
641   Löschmann, N., Voges, Y., Breitling, R., Von Deimling, A., Rödel, F., et al.
642   (2011). Adaptation of cancer cells from different entities to the MDM2 inhibitor
643   nutlin-3 results in the emergence of p53-mutated multi-drug-resistant cancer
644   cells. Cell Death Dis. *2*. 10.1038/cddis.2011.129.

645   17.   Berlak, M., Tucker, E., Dorel, M., Winkler, A., McGearey, A., Rodriguez-Fos,
646   E., da Costa, B.M., Barker, K., Fyle, E., Calton, E., et al. (2022). Mutations in
647   ALK signaling pathways conferring resistance to ALK inhibitor treatment lead
648   to collateral vulnerabilities in neuroblastoma cells. Mol. Cancer *21*, 1–19.
649   10.1186/s12943-022-01583-z.

650   18.   Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R.M., Ozenberger, B.A.,
651   Ellrott, K., Sander, C., Stuart, J.M., Chang, K., Creighton, C.J., et al. (2013).
652   The cancer genome atlas pan-cancer analysis project. Nat. Genet. *45*, 1113–
653   1120. 10.1038/ng.2764.

19.   Klaas, E., Sung, E., Azizi, E., Martinez, M., Barpujari, A., Roberts, J., and

Lucke-Wold, B. (2023). Advanced breast cancer metastasized in the brain:
treatment standards and innovations. J. Cancer Metastasis Treat. *9*.
10.20517/2394-4722.2022.125.

20. Wang, B., Sun, T., Zhao, Y., Wang, S., Zhang, J., Wang, Z., Teng, Y.E., Cai,
L., Yan, M., Wang, X., et al. (2022). A randomized phase 3 trial of Gemcitabine
or Nab-paclitaxel combined with cisPlatin as first-line treatment in patients with
metastatic triple-negative breast cancer. Nat. Commun. *13*. 10.1038/s41467-
022-31704-7.

21. Liu, Y., Fan, L., Wang, Z.H., and Shao, Z.M. (2023). Nab-paclitaxel Followed
by Dose-dense Epirubicin/Cyclophosphamide in Neoadjuvant Chemotherapy
for Triple-negative Breast Cancer: A Phase II Study. Oncologist *28*, 86-e76.
10.1093/oncolo/oyac223.

22. Gumusay, O., Huppert, L.A., Magbanua, M.J.M., Wabl, C.A., Assefa, M.,
Chien, A.J., Melisko, M.E., Majure, M.C., Moasser, M., Park, J., et al. (2023). A
phase Ib/II study of eribulin in combination with cyclophosphamide in patients
with advanced breast cancer. Breast Cancer Res. Treat. *203*, 197–204.
10.1007/s10549-023-07073-0.

23. Kim, S.H., Im, S.A., Suh, K.J., Lee, K.H., Kim, M.H., Sohn, J., Park, Y.H., Kim,
J.Y., Jeong, J.H., Lee, K.E., et al. (2023). Clinical activity of nivolumab in
combination with eribulin in HER2-negative metastatic breast cancer: A phase
IB/II study (KCSG BR18-16). Eur. J. Cancer *195*, 113386.
10.1016/j.ejca.2023.113386.

24. Velikova, G., Morden, J.P., Haviland, J.S., Emery, C., Barrett-Lee, P., Earl, H.,
Bloomfield, D., Brunt, A.M., Canney, P., Coleman, R., et al. (2023).
Accelerated versus standard epirubicin followed by cyclophosphamide,
methotrexate, and fluorouracil or capecitabine as adjuvant therapy for breast
cancer (UK TACT2; CRUK/05/19): quality of life results from a multicentre,
phase 3, open-label, randomised, . Lancet Oncol. *24*, 1359–1374.
10.1016/S1470-2045(23)00460-6.

25. Takahashi, M., Cortés, J., Dent, R., Pusztai, L., McArthur, H., Kümmel, S.,
Denkert, C., Park, Y.H., Im, S.A., Ahn, J.H., et al. (2023). Pembrolizumab Plus
Chemotherapy Followed by Pembrolizumab in Patients with Early Triple-
Negative Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial.
JAMA Netw. Open *6*, E2342107. 10.1001/jamanetworkopen.2023.42107.

26. Liston, D.R., and Davis, M. (2017). Clinically Relevant Concentrations of
Anticancer Drugs : A Guide for Nonclinical Studies. Clin. Cancer Res., 3489–
3498. 10.1158/1078-0432.CCR-16-3083.

27. Kim, N., Hong, Y., Kwon, D., and Yoon, S. (2013). Somatic Mutaome Profile in

Human Cancer Tissues. Genomics Inform. *11*, 239. 10.5808/gi.2013.11.4.239.

28. Reynolds, I.S., Fichtner, M., McNamara, D.A., Kay, E.W., Prehn, J.H.M., and
Burke, J.P. (2019). Mucin glycoproteins block apoptosis; promote invasion,
proliferation, and migration; and cause chemoresistance through diverse
pathways in epithelial cancers. Cancer Metastasis Rev. *38*, 237–257.
10.1007/s10555-019-09781-w.

29. Pandey, K., Lee, E., Park, N., Hur, J., Cho, Y. Bin, Katuwal, N.B., Kim, S.K.,
Lee, S.A., Kim, I., An, H.J., et al. (2021). Deregulated immune pathway
associated with palbociclib resistance in preclinical breast cancer models:
Integrative genomics and transcriptomics. Genes (Basel). *12*, 1–14.
10.3390/genes12020159.

30. Chang, Y., Wang, Y., Li, B., Lu, X., Wang, R., Li, H., Yan, B., Gu, A., Wang,
W., Huang, A., et al. (2021). Whole-Exome Sequencing on Circulating Tumor
Cells Explores Platinum-Drug Resistance Mutations in Advanced Non-small
Cell Lung Cancer. Front. Genet. *12*, 1–11. 10.3389/fgene.2021.722078.

31. Patel, N.M., Geropoulos, G., Patel, P.H., Bhogal, R.H., Harrington, K.J.,
Singanayagam, A., and Kumar, S. (2023). The Role of Mucin Expression in the
Diagnosis of Oesophago-Gastric Cancer: A Systematic Literature Review.
Cancers (Basel). *15*. 10.3390/cancers15215252.

32. Kim, S.H., Ho, J.N., Jin, H., Lee, S.C., Lee, S.E., Hong, S.K., Lee, J.W., Lee,
E.S., and Byun, S.S. (2016). Upregulated expression of BCL2, MCM7, and
CCNE1 indicate cisplatin-resistance in the set of two human bladder cancer
cell lines: T24 cisplatin sensitive and T24R2 cisplatin resistant bladder cancer
cell lines. Investig. Clin. Urol. *57*, 63–72. 10.4111/icu.2016.57.1.63.

33. Feng, Z., Zhang, L., Zhou, J., Zhou, S., Li, L., Guo, X., Feng, G., Ma, Z.,
Huang, W., and Huang, F. (2017). mir-218-2 promotes glioblastomas growth,
invasion and drug resistance by targeting CDC27. Oncotarget *8*, 6304–6318.
10.18632/oncotarget.13850.

34. Peng, L., Zhao, M., Liu, T., Chen, J., Gao, P., Chen, L., Xing, P., Wang, Z., Di,
J., Xu, Q., et al. (2022). A stop-gain mutation in GXYLT1 promotes metastasis
of colorectal cancer via the MAPK pathway. Cell Death Dis. *13*, 1–12.
10.1038/s41419-022-04844-3.

35. Michaelis, M., Klassert, D., Barth, S., Suhan, T., Breitling, R., Mayer, B.,
Hinsch, N., Doerr, H.W., Cinatl, J., and Cinatl, J. (2009). Chemoresistance
acquisition induces a global shift of expression of aniogenesis-associated
genes and increased pro-angogenic activity in neuroblastoma cells. Mol.
Cancer *8*, 80. 10.1186/1476-4598-8-80.

36. Bahar, M.E., Kim, H.J., and Kim, D.R. (2023). Targeting the RAS/RAF/MAPK

pathway for cancer therapy: from mechanism to clinical studies. Signal
Transduct. Target. Ther. *8*. 10.1038/s41392-023-01705-z.

37. Wang, P., Laster, K., Jia, X., Dong, Z., and Liu, K. (2023). Targeting CRAF
kinase in anti-cancer therapy: progress and opportunities. Mol. Cancer *22*, 1–
34. 10.1186/s12943-023-01903-x.

38. Xu, J., Zhou, Y., Dong, K., Gong, J., Xiong, W., Wang, X., Gu, C., Lu, X. yu,
Huang, D. pei, Shen, X. dong, et al. (2023). Gene variation profile and it's
potential correlation with clinical characteristics in HBV-associated HCC
patients of Sichuan Han nationality in China. Asian J. Surg. *46*, 4371–4377.
10.1016/j.asjsur.2023.02.056.

39. Weinstein, J.N. (2013). Cancer Genome Atlas Pan-cancer analysis project.
Nat Genet *45*, 113–1120. 10.3779/j.issn.1009-3419.2015.04.02.

40. Xu, H., Cheung, I.Y., Wei, X.X., Tran, H., Gao, X., and Cheung, N.K. V. (2011).
Checkpoint kinase inhibitor synergizes with DNA-damaging agents in G 1
checkpoint-defective neuroblastoma. Int. J. Cancer *129*, 1953–1962.
10.1002/ijc.25842.

41. Ling, V.Y., Straube, J., Godfrey, W., Haldar, R., Janardhanan, Y., Cooper, L.,
Bruedigam, C., Cooper, E., Tavakoli Shirazi, P., Jacquelin, S., et al. (2023).
Targeting cell cycle and apoptosis to overcome chemotherapy resistance in
acute myeloid leukemia. Leukemia *37*, 143–153. 10.1038/s41375-022-01755-
2.

42. Zeng, L., Nikolaev, A., Xing, C., della Manna, D.L., and Yang, E.S. (2020).
CHK1/2 inhibitor prexasertib suppresses notch signaling and enhances
cytotoxicity of cisplatin and radiation in head and neck squamous cell
carcinoma. Mol. Cancer Ther. *19*, 1279–1288. 10.1158/1535-7163.MCT-19-
0946.

43. VanKlompenberg, M.K., Bedalov, C.O., Soto, K.F., and Prosperi, J.R. (2015).
APC selectively mediates response to chemotherapeutic agents in breast
cancer. BMC Cancer *15*, 1–14. 10.1186/s12885-015-1456-x.

44. Stefanski, C.D., Keffler, K., McClintock, S., Milac, L., and Prosperi, J.R. (2019).
APC loss affects DNA damage repair causing doxorubicin resistance in breast
cancer cells. Neoplasia (United States) *21*, 1143–1150.
10.1016/j.neo.2019.09.002.

45. Astarita, E.M., Maloney, S.M., Hoover, C.A., Berkeley, B.J., Van Klompenberg,
M.K., Murlidharan Nair, T., and Prosperi, J.R. (2021). Adenomatous Polyposis
Coli loss controls cell cycle regulators and response to paclitaxel in MDA-MB-
157 metaplastic breast cancer cells. PLoS One *16*, 1–18.
10.1371/journal.pone.0255738.

46.   Szakács, G., Paterson, J.K., Ludwig, J.A., Booth-Genthe, C., and Gottesman,
M.M. (2006). Targeting multidrug resistance in cancer. Nat. Rev. Drug Discov.
5, 219–234. 10.1038/nrd1984.

47.   Lu, J.F., Pokharel, D., and Bebawy, M. (2015). MRP1 and its role in anticancer
drug   resistance.   Drug   Metab.   Rev.   *47*,   406–419.
10.3109/03602532.2015.1105253.

48.   Derry, W.B., Wilson, L., and Jordan, M.A. (1995). Substoichiometric Binding of
Taxol Suppresses Microtubule Dynamics. Biochemistry *34*, 2203–2211.
10.1021/bi00007a014.

49.   Snyder, J.P., Nettles, J.H., Cornett, B., Downing, K.H., and Nogales, E. (2001).
The binding conformation of Taxol in β-tubulin: A model based on electron
crystallographic density. Proc. Natl. Acad. Sci. U. S. A. *98*, 5312–5316.
10.1073/pnas.051309398.

50.   Jordan, M.A., and Wilson, L. (2004). Microtubules as a target for anticancer
drugs. Nat. Rev. Cancer *4*, 253–265. 10.1038/nrc1317.

51.   Smith, J.A., Wilson, L., Azarenko, O., Zhu, X., Lewis, B.M., Littlefield, B.A., and
Jordan, M.A. (2010). Eribulin binds at microtubule ends to a single site on
tubulin to suppress dynamic instability. Biochemistry *49*, 1331–1337.
10.1021/bi901810u.

52.   Jin, J., Tao, Z., Cao, J., Li, T., and Hu, X. (2021). DNA damage response
inhibitors: An avenue for TNBC treatment. Biochim. Biophys. Acta - Rev.
Cancer *1875*, 188521. 10.1016/j.bbcan.2021.188521.

53.   Baxter, J.S., Zatreanu, D., Pettitt, S.J., and Lord, C.J. (2022). Resistance to
DNA repair inhibitors in cancer. Mol. Oncol. *16*, 3811–3827. 10.1002/1878-
0261.13224.

54.   Bracht, K., Boubakari, Grünert, R., and Bednarski, P.J. (2006). Correlations
between the activities of 19 anti-tumor agents and the intracellular glutathione
concentrations in a panel of 14 human cancer cell lines: Comparisons with the
National   Cancer   Institute   data.   Anticancer.   Drugs   *17*,   41–51.
10.1097/01.cad.0000190280.60005.05.

55.   Michaelis, M., Schneider, C., Rothweiler, F., Rothenburger, T., Mernberger,
M., Nist, A., von Deimling, A., Speidel, D., Stiewe, T., and Cinatl, J. (2018).
TP53 mutations and drug sensitivity in acute myeloid leukaemia cells with
acquired MDM2 inhibitor resistance. bioRxiv, 404475.

56.   Michaelis, M., Wass, M.N., Reddin, I., Voges, Y., Rothweiler, F., Hehlgans, S.,
Cinatl, J., Mernberger, M., Nist, A., Stiewe, T., et al. (2020). YM155-adapted
cancer cell lines reveal drug- induced heterogeneity and enable the
identification of biomarker candidates for the acquired resistance setting.

Cancers (Basel). *12*, 1–17. 10.3390/cancers12051080.

57. Hata, A.N., Niederst, M.J., Archibald, H.L., Gomez-Caraballo, M., Siddiqui,
F.M., Mulvey, H.E., Maruvka, Y.E., Ji, F., Bhang, H.E.C., Radhakrishna, V.K.,
et al. (2016). Tumor cells can follow distinct evolutionary paths to become
resistant to epidermal growth factor receptor inhibition. Nat. Med. *22*, 262–269.
10.1038/nm.4040.

58. Karasaki, T., Moore, D.A., Veeriah, S., Naceur-Lombardelli, C., Toncheva, A.,
Magno, N., Ward, S., Bakir, M. Al, Watkins, T.B.K., Grigoriadis, K., et al.
(2023). Evolutionary characterization of lung adenocarcinoma morphology in
TRACERx. Nat. Med. *29*. 10.1038/s41591-023-02230-w.

59. Martínez-Ruiz, C., Black, J.R.M., Puttick, C., Hill, M.S., Demeulemeester, J.,
Larose Cadieux, E., Thol, K., Jones, T.P., Veeriah, S., Naceur-Lombardelli, C.,
et al. (2023). Genomic–transcriptomic evolution in lung cancer and metastasis.
Nature *616*, 543–552. 10.1038/s41586-023-05706-4.

60. Al Bakir, M., Huebner, A., Martínez-Ruiz, C., Grigoriadis, K., Watkins, T.B.K.,
Pich, O., Moore, D.A., Veeriah, S., Ward, S., Laycock, J., et al. (2023). The
evolution of non-small cell lung cancer metastases in TRACERx
10.1038/s41586-023-05729-x.

61. Frankell, A.M., Dietzen, M., Al Bakir, M., Lim, E.L., Karasaki, T., Ward, S.,
Veeriah, S., Colliver, E., Huebner, A., Bunkum, A., et al. (2023). The evolution
of lung cancer and impact of subclonal selection in TRACERx. Nature *616*,
525–533. 10.1038/s41586-023-05783-5.

62. Abbosh, C., Frankell, A.M., Harrison, T., Kisistok, J., Garnett, A., Johnson, L.,
Veeriah, S., Moreau, M., Chesh, A., Chaunzwa, T.L., et al. (2023). Tracking
early lung cancer metastatic dissemination in TRACERx using ctDNA. Nature
616, 553–562. 10.1038/s41586-023-05776-4.

63. Michaelis, M., Rothweiler, F., Barth, S., Cinat, J., Van Rikxoort, M.,
Löschmann, N., Voges, Y., Breitling, R., Von Deimling, A., Rödel, F., et al.
(2011). Adaptation of cancer cells from different entities to the MDM2 inhibitor
nutlin-3 results in the emergence of p53-mutated multi-drug-resistant cancer
cells. Cell Death Dis. *2*. 10.1038/cddis.2011.129.

64. Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival:
Application to proliferation and cytotoxicity assays. J. Immunol. Methods *65*,
55–63. 10.1016/0022-1759(83)90303-4.

65. Onafuye, H., Pieper, S., Mulac, D., Cinatl, J., Wass, M.N., Langer, K., and
Michaelis, M. (2019). Doxorubicin-loaded human serum albumin nanoparticles
overcome transporter-mediated drug resistance in drug-adapted cancer cells.
Beilstein J. Nanotechnol. *10*, 1707–1715. 10.3762/bjnano.10.166.

66.   Andrews S (2018). FastQC A Quality control tool for high throughput sequence
846   data. Babraham Bioinfo, 3–5.

847   67.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible
848   trimmer for Illumina sequence data. Bioinformatics *30*, 2114–2120.
849   10.1093/bioinformatics/btu170.

850   68.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
851   Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and
852   SAMtools. Bioinformatics *25*, 2078–2079. 10.1093/bioinformatics/btp352.

853   69.   Burrows, M., and Wheeler, D. (1994). A block-sorting lossless data
854   compression algorithm. Algorithm, Data Compression, 18. 10.1.1.37.6774.

855   70.   Church, D.M., Schneider, V.A., Graves, T., Auger, K., Cunningham, F., Bouk,
856   N., Chen, H.C., Agarwala, R., McLaren, W.M., Ritchie, G.R.S., et al. (2011).
857   Modernizing reference genome assemblies. PLoS Biol. *9*.
858   10.1371/journal.pbio.1001091.

859   71.   McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky,
860   A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The genome
861   analysis toolkit: A MapReduce framework for analyzing next-generation DNA
862   sequencing data. Genome Res. *20*, 1297–1303. 10.1101/gr.107524.110.

863   72.   Li, H. (2011). A statistical framework for SNP calling, mutation discovery,
864   association mapping and population genetical parameter estimation from
865   sequencing data. Bioinformatics *27*, 2987–2993.
866   10.1093/bioinformatics/btr509.

867   73.   McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A.,
868   Flicek, P., and Cunningham, F. (2016). The Ensembl Variant Effect Predictor.
869   Genome Biol. *17*, 1–14. 10.1186/s13059-016-0974-4.

870   74.   Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang,
871   Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2019).
872   Variation across 141,456 human exomes and genomes reveals the spectrum
873   of loss-of-function intolerance across human protein-coding genes. bioRxiv,
874   531210. 10.1101/531210.

875   75.   Ghandi, M., Huang, F.W., Jané-Valbuena, J., Kryukov, G. V., Lo, C.C.,
876   McDonald, E.R., Barretina, J., Gelfand, E.T., Bielski, C.M., Li, H., et al. (2019).
877   Next-generation characterization of the Cancer Cell Line Encyclopedia.
878   Nature. 10.1038/s41586-019-1186-3.

879   76.   Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A.,
880   Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., et al. (2004). The
881   COSMIC (Catalogue of Somatic Mutations in Cancer) database and website.
882   Br. J. Cancer *2*, 355–358. 10.1038/sj.bjc.6601894.

77.   Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and
Vilo, J. (2019). g:Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). Nucleic Acids Res. *47*, W191–W198.
10.1093/nar/gkz369.

78.   Moiso, E. (2021). Manual curation of TCGA treatment data and identification of
potential markers of therapy response. medRxiv, 2021.04.30.21251941.

79.   Lehmann, B.D., Colaprico, A., Silva, T.C., Chen, J., An, H., Ban, Y., Huang, H.,
Wang, L., James, J.L., Balko, J.M., et al. (2021). Multi-omics analysis identifies
therapeutic vulnerabilities in triple-negative breast cancer subtypes. Nat.
Commun. *12*, 1–18. 10.1038/s41467-021-26502-6.

Figure Legends

Figure 1. Confirmation of the resistance status of the project cell lines. A)
Panel of drug-naïve (MDA-MB-468, HCC38, HCC1806) and drug-adapted Triple
Negative Breast Cancer cell lines. B) Left: dose–response curve; bottom: IC_{50}
values; right: resistance factor ($IC_{50,}$ drug-adapted subline/$IC_{50,}$ respective parental
cell line)); when drug-naïve and drug-adapted cell lines are treated with the
respective agent: cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, or 5-
fluorouracil. Circles indicate drug-naïve cell lines, and crosses indicate drug-adapted
cell lines. Green, MDA-MB-468-derived; blue, HCC38-derived; orange, HCC1806-
derived. The data are from ≥ 3 independent experiments, and the statistics were
calculated using Student's t-test and are plotted as the means ± SDs.

Figure 2. Genomic characterization of drug-adapted cell lines. A) Diagram
illustrating the differences between *gained, de novo*, *not called*, *lost* and *shared*
variants. B) Count of *Gained* (blue) and *De novo* (green) variants, C) count of *Lost*
(orange) and *Not-called* (pink) variants, D) left panel; count of all *Shared* (purple)
variants, right panel; two-fold increase or decrease of shared variants.

Figure 3. Identification of novel candidates associated with therapy failure. A)
Flow chart of genes with *de novo* variants observed in two or more sublines from
more than one parental cell line. B) Venn diagrams of *de novo* variants shared
between sublines adapted to the same drug. C) Summary of relatedness between
sublines drug-adapted from the same parental cell line (%).

921
Figure 4. Tumor patient data available for mutations in resistance-associated
genes. (A) TCGA pan-cancer datasets for mutation status and gene expression.
Only patients for which clinical, drug and mutation status/gene expression data was
available for were considered in the TCGA pan-cancer analysis. (B) TCGA pan-
cancer mutation status and expression data available for chemotherapy drugs for 29
TCGA cancer classifications. (C) Kaplan-Meier plots for gene expression with most
significant association with prognosis in the pan-cancer dataset. Log-rank test was
the statistical test used with multiple test correction performed using Benjamini-
Hochberg method. (D-E) Genes for which expression is significantly associated with

patient prognosis. Upset plot showing the number of genes that are associated with
patient prognosis for (D) pan-cancer and (E) TNBC.

933
Figure 5. Complex sensitivity patterns to cytotoxic and DDR-targeted agents.
A) Heatmap of fold resistance and collateral sensitivity to cytotoxic agents. B)
Summary of pathways targeted by DNA damage response and repair (DDRR)
inhibitors used in screening. C) Heatmap of fold change resistance and collateral
sensitivity to DDRR inhibitors.

Figure 6. Lack of trends in drug or inhibitor sensitivity patterns. Graphs
demonstrating a negative correlation; collateral sensitivity to one agent but
resistance to the other (blue); positive correlation; resistance to both agents (red);
and no statistical correlation (black) for each set of sublines adapted from the MDA-
MB-468,   HCC38   or   HCC1806   TNBC   cell   lines.

Supplementary Figure 1. Chemo-naïve cell lines are clinically sensitive to
chemotherapy agents. IC_{50} values of drug-naïve parental cell lines treated with the
respective chemotherapy agents: cisplatin, doxorubicin, eribulin, paclitaxel,
gemcitabine or 5-fluorouracil. Green, MDA-MB-468 cells; blue, HCC38 cells; orange,
HCC1806 cells. The black line indicates known C_{max} values for each chemotherapy
agent. Data from n ≥ 3, statistics were calculated using Student's t-test and are
plotted as the mean ± SD.

951
952 **Supplementary Figure 2: Variant counts.** A) Total number of variants called for in
953 the panel of drug-naïve and drug-resistant cell lines. B) Types of variants called for in
954 the panel of drug-naïve and drug-resistant cell lines, including missense,
955 synonymous, frameshift, inframe insertion, inframe deletion, stop loss, stop gain,
956 splice acceptor and splice donor variants.

957
Supplementary Figure 3. *De novo* variant overlaps. The number of *de novo*
variants that overlap in A) drug-resistant cell lines adapted to the same
chemotherapy drug and B) drug-resistant cell lines adapted from the same parental
cell line but to different chemotherapy drugs.

Supplementary Figure 4. Gene ontology terms related to variants in drug-

resistant sublines. A) The number of variants increased in drug-resistant sublines

(*de novo* variants, *gained* variants and *shared* variants that demonstrated a ≥2

increase in variant allele frequency). B) The number of variants decreased in drug-

resistant sublines (*not-called* variants, *lost* variants and *shared* variants that

demonstrated ≤2 decreases in variant allele frequency). The number and

overlapping terms found in increased and decreased variants were compared

between cell lines adapted to the same chemotherapy drug (C, E) and sublines

derived from the same parental cell line but adapted to different chemotherapy drugs

(D, F). Green bars indicate increased variants (A, C, D), and red bars indicate

decreased variants (B, C, D).

Supplementary Figure 5. Chemo-naïve cell lines are clinically sensitive to DNA

damage response and repair (DDRR) inhibitors. IC_{50} values of drug-naïve cell lines

treated with the indicated drug. Green, MDA-MB-468-derived; blue, HCC38-derived;

orange, HCC1806-derived. The black line indicates known C_{max} values for each

DDRR agent. The data are from ≥ 3 independent experiments, and the statistics

were calculated using Student's t-test and are plotted as the means ± SDs.

Supplementary Table 1. Drug correlation of delta (Δ) values. The IC_{50} values

were transformed to ΔIC_{50} values for each drug (see methods) and correlated across

the drug panel, with linear regression analysis and statistical significance. The values

in the table indicate the r values of the correlations, where positive values indicate

positive correlations and negative values indicate negative correlations. P values of

the correlations are indicated in the blue color scheme, with light blue (p≤0.05)

indicating the lowest statistical significance and dark blue (p≤0.00001) indicating the

highest statistical significance.

Supplementary File 1. Mean IC_{50} values, SDs and resistance factors for the project

panel treated with chemotherapy drugs and DNA damage response inhibitors.

Supplementary File 2. Basic variant characterization of the cell line panel.

Supplementary File 3. Variants found to be *de novo*, *gained*, *not called*, *lost* and
shared in drug-resistant cell lines.

Supplementary File 4. List of genes with *de novo* variants in ≥2 drug-resistant cell
lines. The values in the table indicate the variant allele frequencies of the *de novo*
variants identified in the indicated genes. PMIDs for genes previously implicated in
cancer and drug resistance.

1003

1004 **Supplementary File 5.** Step-by-step analysis of both TNBC and pan-cancer patient

1005 data extracted from the TCGA.

1006 **Supplementary File 6.** Comparison of genes identified through *de novo* variant

1007 analysis   and   TCGA   analysis.

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

Figure 1

Figure 2

A

Parental → Drug-adapted

Gained Variants

De novo Variants

Not-called Variants

Lost Variants

Shared Variants

B

MDA-MB-468^{r}CDDP^{100}
MDA-MB-468^{r}DOX^{50}
MDA-MB-468^{r}ERI^{100}
MDA-MB-468^{r}PCL^{20}
HCC38^{r}CDDP^{3000}
HCC38^{r}DOX^{40}
HCC38^{r}ERI^{10}
HCC38^{r}GEM^{20}
HCC38^{r}PCL$^{2.5}$
HCC1806^{r}CDDP^{500}
HCC1806^{r}DOX$^{12.5}$
HCC1806^{r}ERI^{50}
HCC1806^{r}GEM^{20}
HCC1806^{r}PCL^{20}
HCC1806^{r}5-F^{1500}

Number of variants

C

Number of variants

D

DA-MB-468^{r}CDDP^{100}
MDA-MB-468^{r}DOX^{50}
MDA-MB-468^{r}ERI^{100}
MDA-MB-468^{r}PCL^{20}
HCC38^{r}CDDP^{3000}
HCC38^{r}DOX^{40}
HCC38^{r}ERI^{10}
HCC38^{r}GEM^{20}
HCC38^{r}PCL$^{2.5}$
HCC1806^{r}CDDP^{500}
HCC1806^{r}DOX$^{12.5}$
HCC1806^{r}ERI^{50}
HCC1806^{r}GEM^{20}
HCC1806^{r}PCL^{20}
HCC1806^{r}5-F^{1500}

Number of variants

E

Number of variants

- Gained
- De novo
- Lost
- Not-called
- Shared
- ≥ 2 fold Shared
- < 2 fold Shared

Figure 3

A

De novo variants in ≥2 sublines from more >1 parental cell line

81 genes

Involved in drug resistance
46 genes

Novel findings
33 genes

Related to cancer
24 genes

Not previously implicated in cancer
9 genes

ABCD1, ATAD2, ATXN1, CDC27, CDK2AP2, CSMD3, DDX11, DSPP, FBN2, FLG, FRG1, HSPD1, ITGB4, KMT2C, LAMA1, MAML2, MAML3, MMP16, MTCH2, MT-ND5, MUC16, MUC2, MUC3A, MUC4, MUC5B, MUC6, NBPF10, NCOA3, NCOR1, NCOR2, NEFH, NUP160, PABPC3, PHF2, PRSS3, PSPH, SCN1A, SLC2A12, SPTA1, TBP, TDG, TFAP2B, TTN, USP6, VCX, XDH

ACSM2A, CDON, CNTNAP3B, CRB1, CRIPAK, DENND4B, EP400, ERCC6-PGBD3, GALNT13, GOLGA6L2, GXYLT1, HERC2, IGFN1, INADL, MYH4, MYO5B, PGBD3, PRB4, RGPD3, RGPD4, RP11-830F9.6, RXFP2, TAS2R46, TMEM199, TRBV10-1, USH2A

C12orf55, C2orf16, CATSPER3, KRTAP4-11, KRTAP4-6, KRTAP4-7, TAS2R30, TRBV7-7, ZNF385C

C

B

Figure 4

Figure 5