Resistance patterns in drug-adapted cancer cell lines reflect complex evolution in clinical tumors

Helen E. Grimsley[1,2], Magdalena Antczak[1], Katie-May McLaughlin[1], Andrea Nist[3], Marco Mernberger[4], Thorsten Stiewe[3,4], Daniel Speidel[5,6], Catherine Harper-Wynne[7], Karina Cox[8], Jindrich Cinatl jr.[9,10], Mark Wass[1]*, Michelle D. Garrett[1]*, Martin Michaelis[1,9]*

[1]School of Biosciences, Stacey Building, University of Kent, Canterbury, Kent, CT2 7NJ, UK

[2]Department of Radiation Oncology and the Molecular Biology Program, Memorial Sloan Kettering Cancer Center, New York, United States

[3]Genomics Core Facility, Philipps-University, 35043 Marburg, Germany.

[4]Institute of Molecular Oncology, Member of the German Center for Lung Research (DZL), Philipps-University, 35032 Marburg, Germany.

[5]Children's Medical Research Institute, The University of Sydney, 214 Hawkesbury Road, Westmead, New South Wales, Australia

[6]Current address: Breakpoint Therapeutics GmbH, Essener Bogen 7, 22419 Hamburg, Germany

[7]Kent Oncology Centre, Maidstone and Tunbridge Wells NHS Trust, Hermitage Lane, Maidstone, ME16 9QQ, UK

[8]Peggy Wood Breast Unit, Maidstone Hospital, Hermitage Lane, Maidstone, Kent, ME16 9QQ, UK

[9]Dr Petra Joh-Research Institute, 60528 Frankfurt am Main, Germany

[10]Institute for Medical Virology, University Hospital, Goethe University Frankfurt am Main, 60596 Frankfurt, Germany.

*Corresponding authors emails: M.Michaelis@kent.ac.uk; M.D.Garrett@kent.ac.uk;

M.N.Wass@kent.ac.uk

Abstract

Here, we introduce a novel set of drug-adapted triple-negative breast cancer

(TNBC) cell lines consisting of the parental cell lines MDA-MB-468, HCC38, and

HCC1806 and their sublines adapted to cisplatin, doxorubicin, eribulin, paclitaxel,

gemcitabine, or 5-fluorouracil. Whole exome sequencing in combination with the

analysis of TCGA-derived patient data resulted in the identification of 135 biomarker

candidates for the guidance of personalized TNBC therapies for further investigation,

including 58 novel ones that had not been associated with drug resistance before.

The analysis of exome sequencing data showed remarkably few overlaps among the

resistant sublines, suggesting that each resistance formation process follows an

individual, unpredictable route. This complexity was confirmed by cancer cell line

drug sensitivity profiles to cytotoxic anti-cancer drugs and DNA damage repair

inhibitors. Drug-adapted sublines of the same parental cell line and sublines adapted

to the same drug substantially differed in their drug response patterns. Cross-

resistance levels were lowest for the CHK2 inhibitor CCT241533, the PLK1 inhibitor

SBE13, and the RAD51 recombinase inhibitor B02, making CHK2, PLK1, and

RAD51 promising drug targets for therapy-refractory TNBC. In conclusion, we

present novel preclinical models of acquired drug resistance in TNBC and 58 novel

candidate biomarkers for further investigation. Whole exome data and drug

sensitivity profiles showed that each cancer cell line adaptation process follows an

unpredictable route, which reflects recent findings on cancer cell evolution in

patients, supporting the relevance of drug-adapted cancer cell lines as preclinical

models of acquired resistance.

Key words

Triple Negative Breast Cancer, acquired drug resistance, exome sequencing DNA

repair, *de novo* variants, TCGA

Introduction

Triple negative breast cancer (TNBC) is characterized by the absence of estrogen, progesterone, and HER2 receptors [1]. It is responsible for about 15% of breast cancer cases and associated with a poorer prognosis than hormone receptor or HER2 positive breast cancers [1,2]. Current TNBC therapies are largely based on cytotoxic anti-cancer drugs with treatments including cisplatin, doxorubicin, eribulin, gemcitabine, paclitaxel, and 5-fluorouracil [1]. TNBC often responds initially well to cytotoxic chemotherapy, but recurrence and resistance formation are common, eventually leading to therapy failure. This combination of an initial high response rate followed by rapid resistance formation is referred to as the 'TNBC paradox' (Fornier and Fumoleau, 2012; Gupta *et al.*, 2020). To improve TNBC therapy outcomes, new treatment approaches are needed, in particular those that are effective against treatment-refractory disease characterized by acquired resistance to cytotoxic chemotherapy.

The processes underlying the formation of acquired drug resistance in cancer differ from those responsible for intrinsic resistance (Michaelis, Wass and Cinatl, 2019; Oellerich *et al.*, 2019; Santoni-Rugiu *et al.*, 2019; Touat *et al.*, 2020; Rothenburger *et al.*, 2021). In contrast to intrinsic drug resistance, that occurs independently of therapy, and is a consequence of pre-existing often stochastic events in the tumor, acquired resistance is the direct consequence of selection and adaptation processes caused by cancer treatment (directed tumor evolution). These discrepancies in origin can result in differences between the mechanisms underlying intrinsic and acquired drug resistance (Michaelis, Wass and Cinatl, 2019; Oellerich *et al.*, 2019; Santoni-Rugiu *et al.*, 2019; Touat *et al.*, 2020; Rothenburger *et al.*, 2021).

Identifying and understanding these mechanisms is therefore essential to optimize

cancer treatment for patients with therapy-refractory tumors.

Cancer cell line adaptation to anti-cancer drugs provides a preclinical platform

for the investigation of treatment-induced cancer cell evolution that has been shown

in numerous studies to reflect clinically relevant acquired drug resistance

mechanisms [5,10–18]. Furthermore, the resulting drug-resistant cell lines allow detailed

functional and systems level studies that are not possible using clinical samples

(Michaelis, Wass and Cinatl, 2019).

Here, we introduce a novel set of three parental TNBC sublines and their 15

sublines adapted to cisplatin, doxorubicin, eribulin, gemcitabine, paclitaxel, or 5-

fluorouracil. The project cell lines were characterized by whole exome sequencing

and the determination of response profiles to cytotoxic drugs and DNA damage

repair inhibitors. The resulting data showed that the resistance formation processes

are individual and unpredictable. The combined analysis of the resistance-

associated mutations in combination with patient data from The Cancer Genome

Atlas (TCGA) resulted in 58 candidate resistance biomarkers for further investigation

[19].

Results

Project cell line panel

The project cell line panel consists of the parental TNBC cell lines MDA-MB-

468, HCC38, and HCC1806 and their sublines adapted to grow in the presence of

cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, or 5-fluorouracil, drugs from

drug classes that are used for the treatment of this cancer type (Fig.1A, SupFile.1)

[20–26]. Drug-resistant sublines were established by continuous exposure to stepwise

increasing drug concentrations as previously described [17]. All parental cell lines were

initially sensitive to therapeutic concentrations of the respective drugs, as indicated

by IC_{50} (concentration that reduces cell viability by 50%) values within the range of

clinical drug plasma concentrations (C_{max}) (SupFig.1A) [27]. The relative resistance

factors (IC_{50} drug-adapted subline/ IC_{50} respective parental cell line) ranged from 5.5-

fold (HCC38^{r}PCL$^{2.5}$) to 5916.7-fold (HCC1806^{r}ERI^{50}) (Fig.1B, SupFile.1).

Characterization of the cell line panel by whole exome sequencing

We initially performed whole exome sequencing on the project cell line panel.

Between 186 (HCC38^{r}DOX^{40}) and 739 (HCC38^{r}GEM^{20}) DNA sequence variants

were detected in the drug-adapted sublines that differed from the respective parental

cell lines (SupFig.2A, SupFile.2). Missense variants were most common, followed by

synonymous variants (SupFig.2B). Insertions/ deletions (INDELs), frameshift

mutations, stop-gain, stop-loss, and splice variants were identified at lower

frequencies (SupFig.2B).

We grouped the resistance-associated variants into five categories (Fig.2A,

see methods): 1. *Gained variants*, variants only called in the drug-adapted subline,

but detectable at low confidence in the respective parental cell line; 2. *De novo*

variants, variants called in the drug-adapted subline but undetectable in the

respective parental cell line; 3. *Not-called variants*, variants only called in the

parental cell line, but detectable with low confidence in the resistant subline; 4. *Lost*

variants; variants called in the parental cell line, but undetectable in the drug-adapted

subline; and 5. *Shared variants*; variants called in both the parental and drug-

adapted cell lines that increased or decreased by at least two-fold are presented

(Fig.2A).

The number of *gained* variants ranged from 44 (HCC38[r]DOX[40]) to 381

(HCC38[r]GEM[20]), of *de novo* variants from 31 (HCC38[r]DOX[40]) to 225 (MDA-MB-

468[r]PCL[20]), of *not-called* variants from 88 (HCC38[r]GEM[20] and HCC1806[r]DOX[12.5]) to

345 (MDA-MB-468[r]PCL[20]), of *lost* variants from 129 (HCC38[r]GEM[20]) to 398 (MDA-

MB-468[r]PCL[20]), and of shared variants from 128 (MDA-MB-468[r]PCL[20]) to 368

(HCC38[r]GEM[20]) (Fig.2B-2D, SupFile.3).

Analysis of the distribution of *de novo* variants

To identify variants that may have a functional role in drug resistance, we

initially looked at genes that harbored *de novo* variants in at least two different

sublines from more than one parental cell line, resulting in a list of 81 genes (Fig.3A,

SupFile.4). This list includes 48 genes that had already been described to be

involved in drug resistance in cancer and 33 new candidate genes with a possible

role in drug resistance (Fig.3A, SupFile.4). Notably, 24 of the 33 new candidate

genes had already been reported to be of relevance in cancer (Fig.3A, SupFile.4).

Among the five genes with *de novo* variants in the most cell lines were the

mucin genes *MUC6* (15 cell lines), *MUC2* (14 cell lines), *MUC4* (13 cell lines), and

MUC16 (9 cell lines) (Fig.3A, SupFile.4). These are large genes that are known to be

commonly mutated and have been reported to be involved in cancer cell drug

resistance [28–32]. *De novo* mutations in CDC27, which has also been linked to drug

resistance in cancer, were also found in 9 resistant sublines [33,34] (Fig.3A, SupFile.4).

GXYLT1, *KRTAP4-11*, and *RGPD4* were the genes among those had not

previously been associated with drug resistance that displayed *de novo* mutations in

the most (7) resistant sublines (Fig.3A, SupFile.4). GXYLT1 promotes metastasis

formation in colorectal cancer through MAPK signaling, a pathway known to provide

resistance to a range of anti-cancer drugs [35–38]. RGPD4 was correlated with

vascular invasion in HBV-associated hepatocellular carcinoma, and it is known that

there is an overlap between pro-angiogenic, pro-metastatic, and resistance-

associated signaling in cancer [36,39]. There is no known link between KRTAP4-11 and

cancer, but KRTAP4-11 expression levels were reported to predict the methotrexate

response in rheumatoid arthritis patients [39]. Hence, it seems plausible that these

genes and their products may be involved in cancer cell drug resistance.

Taken together, our analysis identified many genes already known to be

involved in cancer cell drug resistance alongside a substantial number of novel

candidates potentially contributing to therapy failure. Further research will have to

characterize the roles of these individual genes in detail.

When we compared the overlaps between exactly the same *de novo* variants

in sublines adapted to the same drug the numbers were too small to draw any

meaningful conclusions (Fig.3B, Sup.Fig3A). Notably, *de novo* variants in drug-

resistant sublines may not always represent actual novel variants that are selected

because they contribute to cancer cell resistance. Many apparent *de novo* mutations

have probably already been present in a small fraction of the cells of the parental cell

line, but have not been detected due to limited sequencing depth. Hence, overlaps in

de novo variants between sublines of the same parental cell line can also be used to

indicate the levels of relatedness between the founding subpopulations of the

different resistant sublines.

Analysing *de novo* variants shared between the sublines of each resistant

sublines indicated the largest overlap (22.6% on average) and, thus, relatedness

among the HCC1806 sublines, followed by the HCC38 (15.0%), and the MDA-MB-

468 (7.7%) sublines (Fig.3C). There were noticeable differences in the overlaps

between *de novo* variants of the sublines of the individual parental cell lines. For

example, only three *de novo* variants were shared between HCC38[r]CDDP[3000] (out of

98 in total, 3.1%) and HCC38[r]PCL[2.5] (out of 92 in total, 3.3%), while 53 variants were

shared between HCC38[r]ERI[10] (out of 131 in total, 40.5%) and HCC38[r]GEM[20] (out of

203 in total, 26.1%) (Fig.3C, SupFig.3B). However, there were no patterns

suggesting consistent overlaps between sublines adapted to certain drugs.

Therefore, there is no indication that certain drugs may select certain pre-existing

cell line subpopulations.

Gene ontology (GO) terms related to gene variants that changed in drug-

resistant sublines

Next, we performed a gene ontology (GO) term analysis of *de novo*, *gained*,

not called, and *lost* variants as well as *shared* variants with a two-fold increase or

decrease in allele frequency (Sup.Fig4A, B).

There was a limited overlap between GO terms among sublines adapted to

the same drug (SupFig.4C, E).   The extracellular matrix-related GO terms

'extracellular matrix constituent lubricant activity', 'extracellular matrix', and

'maintenance of gastrointestinal epithelium' were most common which reflects the

high number of variants observed in the mucin genes (SupFig.4C, E).

A GO term analysis among the sublines of the same parental cell lines

resulted in very similar results, again revealing an overrepresentation of extracellular

matrix-related GO terms (SupFig.4D, F). Further research will have to investigate the

potential role of mucins and changes to the extracellular matrix in acquired drug

resistance in TNBC cells.

Potential clinical relevance of selected variants

The potential clinical relevance of *de novo*, *gained*, and *shared* variants with a

two-fold increase was analyzed using patient data derived from The Cancer Genome

Atlas (TCGA) data [40]. Initially, we investigated exact variants, i.e., variants that have

the same chromosomal position and base change in the TCGA data set. Next, we

analyzed variants in the same chromosomal position but with a different base

change that resulted in the same consequence i.e., a missense variant. Here, 27

Exact and 40 Same Consequence resistance-associated increased variants were

identified in TCGA-derived patient mutation data (Fig.4A, Fig.4B).

We also focused on all protein truncating variants in cells lines. Variants were

selected from the TCGA database if they were of a similar consequence (i.e., a

frameshift). Here, 65 resistance-associated variants were found to have protein

truncating variants of similar consequence identified in the sublines were also

identified in the TCGA-derived patient mutation data (Fig.4, Fig.4C).

The fraction of mutated tumors was too low for a meaningful analysis of the

potential role of the variants in clinical drug response. Hence, we used gene

expression data to evaluate further the role of the respective genes and their

products in response to the respective drugs. The following numbers of Exact and

Same Consequence variants could be examined for the respective drugs: cisplatin:

22 (9 Exact, 13 Same consequence), doxorubicin: 9 (3 Exact, 6 Same

Consequence), paclitaxel: 12 (5 Exact, 7 Same Consequence), gemcitabine: 16 (8

Exact and 8 Same Consequence), 5-fluorouacil: 8 (2 Exact, 6 Same Consequence)

(Fig.4B, Sup.File.5).

For the protein truncating variants, the number of analyzed genes was;

cisplatin: 16, doxorubicin: 14, paclitaxel: 7, gemcitabine: 22, 5-fluorouacil: 2. From

these genes, Kaplan Meier curves were plotted for high and low gene expression

and filtered for FDR and statistical significance (see methods). For the protein

truncating variant analysis, we focused on Kaplan Meier curves in which low gene

expression was associated with poor patient outcome, as truncations are most likely

to result in a loss of function (Fig.4C, SupFile.6).

In total, we identified 62 genes whose expression was associated with the

therapy response in cancer patients (Fig.4, Sup.File.5, SupFile.6). This included five

genes with a known role in resistance to the drug that the cell line, in which we

identified the variants, was adapted to (*COL22A1, FAT4, RGS9, SLC2A12, SLC4A8*)

(Fig.4, SupFile.7). Expression levels of 18 further genes have been reported to

mediate resistance to other drugs (*ABCB10, ADNP, C20orf27, CHST11, EXT1,*

FKBP7, NCOR1, PIK3C2B, ABCA8, ACIN1, DNAH5, MTCH2, INHBA, KLF11,

SLC22A23, SLC24A1, SMC1B, TYK2) (SupFile.7). For another 12 genes, there is

evidence that they contribute to both (*HUWE1, ITGB4, MSK1, PHF2, TRPM7,*

BRD7, CES2, IDO1, MSH2, PRLR, RPL14, TOP2A (Fig.4F) (SupFile.7).

In addition to these 35 genes and their products with a known role in drug

resistance, we identified 27 genes that have not previously been linked to drug

resistance (*ANK2, C11orf80, C5orf42, DNAJC13, EPB41, FGF14, FLG, GBGT1,*

GXYLT1, KCND2, OGN, RNF213, TBC1D9, USH2A, AGAP6, CDON, CEBPZ,

CNEP1R1, COG6, CUBN, EFCAB6, HSD17B3, KIAA0586, SETX, SYNGR1,

ZKSCAN3, ZNF442) (Fig.4E).

Eight of the genes and gene products (*CDON*, *FLG*, *GXYLT1*, *ITGB4*,

NCOR1, *PHF2*, *SLC2A12*, *USHS2A*) had already been identified in our analysis of

de novo variants (SupFile.4). Six of them have previously been associated with

cancer drug resistance (*ITGB4*, *NCOR1*, *PHF2*, *SLC2A12*, *USHS2A*) (Fig.3, Fig.4,

SupFile.4, SupFile.7. Further research will have to define in more detail the potential

use of the expression levels of and variants in these genes as biomarkers for the

direction of clinical therapies.

Complex sensitivity patterns of drug-resistant sublines against cytotoxic

drugs

Determining drug sensitivity profiles in the project cell line panel against the

drugs of adaptation, i.e., cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, and

5-fluorouracil (Fig.5A, SupFile.1), revealed complex resistance patterns that did not

follow clear, predictable rules. For example, two out of the three doxorubicin-adapted

sublines (HCC38^{r}DOX^{40}, HCC1806^{r}DOX$^{12.5}$) displayed increased (collateral)

sensitivity to cisplatin, while MDA-MB-468^{r}DOX^{50} displayed cross-resistance to

cisplatin (Fig.5A, SupFile.1). Moreover, all resistant sublines remained sensitive to or

showed collateral sensitivity against at least one of the other chemotherapeutic

agents (Fig.5A, SupFile.1). The 5-fluorouracil-resistant HCC1806^{r}5-F^{1500} subline was

the only resistant cell line that remained sensitive to all other investigated cytotoxic

drugs (Fig.5A, SupFile.1).

Only five of the nine sublines adapted to the ABCB1 substrates doxorubicin,

eribulin, and paclitaxel (including all three eribulin-resistant sublines) displayed

cross-resistance to all other ABCB1 substrates. Among the ABCB1 substrate-

adapted sublines, all eribulin and paclitaxel-adapted sublines displayed cross

resistance to the respective other drug (Fig.5A, SupFile.1). Notably, eribulin and

paclitaxel are both tubulin-binding agents, but differ in their mechanisms of

interaction with tubulin. Eribulin is a destabilizing agent that binds to the vinca

binding site of tubulin and inhibits microtubule formation, while paclitaxel is a

stabilizing agent that binds to the taxane-binding site that impairs microtubule

degradation [41–45]. Further research will have to show to which extent the tubulin-

binding agent cross-resistance profile of the tubulin-binding agent-adapted sublines

is the consequence of the expression of ABCB1 (and/ or other transporters), tubulin-

related resistance mechanisms, or both.

Taken together, it is not possible to predict how resistance formation to a

certain drug will affect the sensitivity patterns of the resulting sublines to other

cytotoxic agents. However, all of the drug-resistant triple-negative breast cancer

sublines remained sensitive and/ or displayed collateral sensitivity to at least one of

the tested chemotherapeutic drugs. Future research will have to elucidate the

underlying mechanisms to identify biomarkers for personalized therapy approaches

that can guide effective drugs to the right patients [5].

Complex sensitivity patterns of drug-resistant sublines against DNA damage

response inhibitors

Triple-negative breast cancer cells have been shown to harbor defects in DNA

damage repair signaling, which can result in a dependence on the remaining intact

DNA damage repair pathways and, in turn, in sensitivity to certain DNA damage

response inhibitors [4]. Hence, we tested a panel of inhibitors targeting critical nodes

of DNA damage repair signaling in the project cell lines (Fig.5B).

All parental cell lines displayed sensitivity to the tested DNA damage

response inhibitors in therapeutic concentrations, i.e., within the Cmax values

reported for these agents (if available) (SupFig.5). However and similarly to the

results obtained for cytotoxic anti-cancer drugs, the DNA damage response inhibitor

sensitivity profiles in the resistant sublines were complex and unpredictable (Fig.5C,

SupFile.1). Relative to the respective parental cell lines, the sensitivity remained

unchanged for 128 DNA damage response inhibitor/ resistant subline combinations.

Increased resistance (cross-resistance) was detected in 96 DNA damage response

inhibitor/ resistant subline combinations, and increased sensitivity (collateral

vulnerability) was recorded in 16 DNA damage response inhibitor/ resistant subline

combinations. Neither sublines of the same parental cell line nor sublines adapted to

the same drugs displayed substantial overlaps in their DNA damage response

inhibitor sensitivity profiles. Generally, cross-resistance levels were lowest for the

CHK2 inhibitor CCT241533, the PLK1 inhibitor SBE13, and the RAD51 recombinase

inhibitor B02 among the investigated DNA damage response inhibitors (Fig.5C,

SupFile.1).

Cross-resistance patterns were even inconsistent between DNA damage

repair inhibitors with the same targets. For example, different sensitivity patterns

were observed between the ATR inhibitors ceralasertib and berzosertib as well as

the CHK1 inhibitors rabusertib, MK-8776, SRA737, and prexasertib (Fig.5C,

SupFile.1). The reasons for these discrepancies are unclear. Notably, the activity of

the DNA damage repair inhibitors may be modified by interaction with additional

targets, and off-target resistance mechanisms (e.g., processes associated with drug

uptake or efflux) may contribute to these differences [46].

In summary and in line with the findings from the investigation of cytotoxic

anti-cancer drugs, the drug-adapted triple-negative breast cancer sublines displayed

complex, unpredictable sensitivity patterns against DNA damage agents, further

demonstrating that improved future therapies will depend on an improved

understanding of the underlying molecular processes resulting in the identification of

biomarkers that can guide effective therapies to individual patients after treatment

failure [5]. Notably, CHK2, PLK1, and RAD51 may have potential as next-line

therapies for triple-negative breast cancer patients, whose tumors have stopped

responding to chemotherapy.

334   Investigation of drug sensitivity patterns by the Delta (Δ) method

Finally, we used the delta (Δ) method to identify potential patterns in the

response of the project cell lines to all investigated cytotoxic drugs and DNA damage

response inhibitors [47]. The IC_{50} values were transformed to ΔIC_{50} values for each

drug (see methods) and correlated across the drug panel, with linear regression

analysis and statistical significance (Sup. Table1). Positive correlations indicate that

increased drug resistance is seen with both agents, whilst negative correlations

indicate that whilst increasing drug resistance is observed to one agent, acquired

vulnerability is observed in the other agent. In the MDA-MB-468, HCC38, and

HCC1806 sublines, we observed 19, 20, and 60 positive correlations and 2, 8, and 1

negative correlations, respectively (Sup. Table1).

We were most interested in the agents that demonstrate negative correlations

as they may identify potential next-line treatments. However, among the 11 negative

correlations, there were no consistent results across all three cell lines (Fig.6). This

further confirms that resistance mechanisms are complex, individual, and

unpredictable and that the identification of potential next-line therapies after

treatment failure will depend on an improved understanding enabling therapy

monitoring and biomarker-guided treatment adaptation.

Discussion

In this study, we introduce and characterize a novel set of 15 drug-adapted

cell lines derived from three parental cell lines that were sensitive to clinical plasma

concentrations of the respective drugs. Viability tests confirmed that all drug-adapted

sublines had developed substantial resistance to the respective drugs.

Next, we applied whole exome sequencing to identify biomarker candidates

for the guidance of anti-cancer therapies. In a first step, we focused on *de novo*

mutations, i.e. mutations found in a resistant subline but undetectable in the

respective parental cell line. Considering genes that displayed *de novo* mutations in

at least two sublines of two different parental cell lines resulted in 81 resistance-

associated variants, 48 of which were already known to be involved in cancer cell

drug resistance while 33 variants were novel.

In a second approach, we used TCGA data to investigate the potential clinical

relevance of genes that harbored resistance-associated variants in the resistant

sublines. This resulted in the identification of 64 genes, whose expression was

associated with drug response in cancer patients. This included 37 genes and gene

products with a known role in drug resistance and 27 genes that had not been linked

to drug resistance before.

Eight of the genes and gene products (*CDON*, *FLG*, *GXYLT1*, *ITGB4*,

NCOR1, *PHF2*, *SLC2A12*, *USHS2A*) were detected by both approaches, including

six ones with an already documented role in cancer drug resistance (*ITGB4*,

NCOR1, *PHF2*, *SLC2A12*, *USHS2A*). Hence, our study identified in total 135 genes

that may represent novel resistance biomarkers, 58 of which had not been

associated with drug resistance in cancer before. Further research will have to

investigate and define in more detail the role of variants in and the expression of

these genes as biomarkers for the tailoring of personalized cancer therapies.

Notably, drug-adapted cancer cell lines have already been shown to represent

clinically relevant resistance mechanisms in numerous studies [5,10–18].

Interestingly, the analysis of exome sequencing data resulted in remarkably

few overlaps between the investigated resistant sublines, including sublines derived

from the same parental cell line and sublines adapted to the same drug. This

suggests that resistance formation is each time the consequence of a complex,

individual, and unpredictable evolutionary process.

This complexity was confirmed by the determination of drug sensitivity

profiles, both to cytotoxic anti-cancer drugs and DNA damage repair inhibitors. Drug-

adapted sublines of the same parental cell line and sublines adapted to the same

drug displayed substantially different drug response patterns. Nevertheless and

notably, cross-resistance levels were lowest for the CHK2 inhibitor CCT241533, the

PLK1 inhibitor SBE13, and the RAD51 recombinase inhibitor B02 among the

investigated DNA damage response inhibitors. Thus, CHK2, PLK1, and RAD51 may

be promising drug targets in TNBC patients after failure of the established therapies,

in particular if reliable biomarkers are found that identify cancer patients that are

likely to benefit from the respective treatments.

Overall, the results from the characterization of the project cell line panel by

whole exome sequencing and from the determination of drug sensitivity profiles both

indicated that cancer cell resistance formation is a complex, individual, and

unpredictable process. This finding is in agreement with data from studies, in which

cancer cell lines were repeatedly adapted to the same drug in independent

experiments [17,48,49] and with recent findings from the comprehensive analysis of

cancer cell evolution in lung cancer patients [50–54].

In conclusion, we here present a novel set of drug-adapted TNBC cell lines as

preclinical models of acquired drug resistance. An initial characterization by whole

exome sequencing in combination with patient-derived TCGA data resulted in the

identification of 135 biomarker candidates for the guidance of personalized TNBC

therapies for further investigation, including 58 ones that are novel and had not been

associated with drug resistance in cancer before. Finally, whole exome data and

drug sensitivity profiles showed that each cancer cell line adaptation process follows

an individual, unpredictable route, which reflects recent clinical findings from the

monitoring of cancer cell evolution in patients [50–54]. This further supports the

relevance of drug-adapted cancer cell lines as preclinical models of acquired

resistance that can be analyzed and manipulated at a level of detail that is

impossible in the clinical setting.

Materials and Methods

Cell culture

MDA-MB-468, HCC38 and HCC1806 were obtained from ATCC. The drug-adapted sublines (Fig.1A, SupFile.1) were established by continuous exposure to stepwise increasing drug concentrations as previously described and derived from the Resistant Cancer Cell Line (RCCL) collection (https://research.kent.ac.uk/industrial-biotechnology-centre/the-resistant-cancer-cell-line-rccl-collection/)(Michaelis *et al.*, 2011; Michaelis, Wass and Cinatl, 2019). All cell lines were cultured in Iscove's Modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (Sigma-Aldrich, Germany), 2mM L-glutamine, 25mM HEPES (Fisher Scientific, UK), 100IU/mL penicillin, and 100µg/mL streptomycin (Life Technologies, UK) at 37 °C in a humidified atmosphere at 5 % CO_2. The drug-adapted sublines were continuously cultured in the presence of the respective concentrations of the drugs.

Compounds

Compounds were purchased from the indicated suppliers: Adavosertib, Alisertib, Berzosertib, Ceralasertib, MK-8776, Olaparib, Prexasertib, Rabusertib, Rucaparib, SBE13, Tozasertib (Adooq Bioscience), AZD0156, BI2536, Doxorubicin, Gemcitabine (Selleckchem), B02, Cisplatin, 5-Fluorouracil (Sigma-Aldrich), CCT241533, SRA737 (Institute of Cancer Research), Eribulin (Eisia), Paclitaxel (Cayman Chemicals). All drug stocks were prepared in DMSO, and stored at -20 °C, except cisplatin which was prepared in 0.9% NaCl solution and stored in the dark at room temperature.

Cell growth and viability assays

Cell viability was tested using the 3-(4,5-dimethylthiazol-2-yl)-2,5-
diphenyltetrazolium bromide (MTT) dye reduction assay after 120-hour incubation,
modified as previously described [55,56]. Concentrations that reduce cell viability by
50% relative to an untreated control (IC_{50}) were determined and used to calculate the
resistance factor (RF; IC_{50} of drug-adapted cell line / IC_{50} of drug-naive cell line).

Whole exome sequencing

Whole exome sequencing (WES) was performed using the Nextera Exome
Enrichment Kit (Illumina). 2 x 100 nucleotide paired end sequences were input into
Illumina HisSeq2000 with an output of 100 nucleotide paired end reads in FASTQ
format. The sequencing was performed in two lanes providing two sets of FASTQ
data per cell line.

Variant calling

FASTQC was used to control the quality of the raw sequence data [57], prior to
the removal of sequencing adaptors with parameters. Trimmomatic (settings:
NexteraPE-PE.fa:2:30:10 LEADING:3 TRAILING:3 SLIDING WINDOW: 4:15
MILEN:36) [58]. Raw FASTQ files were aligned to the human reference genome
(GRCH37) using the Burrows-Wheeler Alignment (v.0.7.17) with an output as
Sequence Alignment Map (SAM) format applying the default settings -M -R [59–61].
Only paired reads were used and Samtools flagstat used to print statistics
throughout each of the subsequent steps. SAM files were input into Picard tools
SortSam (v.2.17.10), where the read alignments were sorted by coordinate and
converted to a Binary Alignment Map (BAM) format output (Picard Toolkit.2019.
Broad Institute, GitHub Repository. http://broadinstitute.github.io/picard/; Broad

Institute). Picard Tools MarkDuplicates (v2.17.10) was used for the removal of PCR

duplicates (Picard Toolkit. 2019. Broad Institute, GitHub

Repository. http://broadinstitute.github.io/picard/; Broad Institute).

GenomeAnalysisTK-3.7.0 was used to perform base score recalibration [62]. SAMtools

mpileup was used to generate Binary Variant Call Format (BCF) files from the BAM

files, which were then input into BCFtools to call the SNVs and INDELS to generate

a Variant Calling Format (VCF) [63]. Variants were annotated with VEP [64].

Variant filtering

Only variants in the protein sequences were considered. To identify high

confidence variants, variants with a Phred score < 30, variants with less than 10

reads supporting the base call, or with < 3 reads supporting the variant were

removed. Moreover, common variants with a frequency of ≥ 0.001% in the genome

aggregation database (gnomAD) were removed [65], if not ≥ 3 samples were

annotated in The Cancer Genome Atlas (TCGA), or ≥ 10 samples in the Catalogue

Of Somatic Mutations In Cancer (COSMIC) [40,66,67].

Definition of variants

De novo variants: variants that are called in drug-resistant subline, but not

called in parental cell line, even at low confidence. *Gained* variants: variants that are

called in the drug-resistant subline and are called in low confidence in parental cell

line. *Not called* variants: variants that are called in the parental cell line, but not

called in the drug-resistant subline, even at low confidence. *Lost* variants: variants

that are called in the parental cell line and are called in low confidence in the drug-

resistant subline. *Shared* variants: variants that are called in both the parental and

drug-resistant subline.

Gene Ontology

495   Gene ontology (GO) functional enrichment analysis was conducted using
496   G:profiler [68].

TCGA analysis

Variant data was extracted via the GDSC Data portal and the Bioconductor R
package *TCGAbiolinks* was used to obtain clinical data [69,70]. Chromosomal locations
of patient variants were remapped from GRCh38 to GRCh37 using the NCBI
Genome Remapping service. Pan-cancer gene expression and survival data was
extracted for each chemotherapeutic agent. Survival analyses were conducted to
determine the response of the patient treated with the chemotherapeutic agent for
when the gene expression was high or low. Cox proportional hazards regression was
used to calculate the hazard ratio for cohorts expressing high vs low expression
levels of the given gene. The '*surv_*cutpoint' function of the package *survminer* in R
allowed for the identification of the optimal expression cut-ff point to give the lowest
p-value for high vs low expression. The cut-off selected was between the 20^{th} and
80^{th} percentiles of gene expression values as previously described by Uhlen *et al.*,
2017. The calculations used overall survival as the measure of clinical outcome.
Overall survival is defined as days to last medical follow up or death as was
previously described by Ng *et al.*, 2016. The calculations were performed using the
R *survminer* and *survival* packages. From this Kaplan-Meier survival curves were
generated using the R package *ggsurvplot*. Statistical analysis using the Wald test
(or log rank (Mantel-Cox)) test was performed to obtain the p-value of significance
for each Kaplan-Meier graph. It should be noted that eribulin variants were omitted

from this analysis as at the time of analysis, no eribulin patient data was available for

the further analysis steps.

Statistical analysis and data manipulation

GraphPad Prism 6 (GraphPad software Inc, USA) was used to generate

dose-response curves and determine GI_{50} values using non-linear regression (with

variable slope). Statistical significance was calculated using a two-tailed T-test,

assuming unequal variance in GraphPad Prism 6 (GraphPad software Inc, USA).

Delta method was used as described by Bracht *et al.*, 2006 . IC_{50} values were

transformed to Δ IC_{50} values: Δ IC_{50} = *log (average* IC_{50} *in drug over all cell lines)* –

log (individual IC_{50} *in drug for each cell line)*. Linear regression analysis of ΔIC_{50X}

versus ΔIC_{50Y} where X and Y represent two different drugs from the panel, were

performed. The Pearson correlation coefficient (*r*) was used to establish the level of

significance in a two tailed test with (n-2) degrees of freedom, where $p \leq 0.05$.

532
533

Acknowledgements

This work was supported by grants from the Frankfurter Stiftung für krebskranke Kinder, Kent Health, Kent Cancer Trust, and the Rosetrees Trust. We would like to thank Tim Fenton and lab members for experimental guidance and useful discussions and the Institute of Cancer Research for their kind gift of CCT241533 and SRA737. Figures were created using BioRender.com.

Conflict of Interests

Nothing to declare.

References

544   1.   Gupta, G.K., Collier, A.L., Lee, D., Hoefer, R.A., Zheleva, V., van Reesema,
545   L.L.S., Tang-Tan, A.M., Guye, M.L., Chang, D.Z., Winston, J.S., et al. (2020).
546   Perspectives on triple-negative breast cancer: Current treatment strategies,
547   unmet needs, and potential targets for future therapies. Cancers (Basel). *12*,
548   1–33. 10.3390/cancers12092392.

549   2.   Hampton, J.M., Song, J., Jayasekera, J., Schechter, C., Alagoz, O., Stout,
550   N.K., Trentham-dietz, A., Lee, S.J., Huang, X., Mandelblatt, J.S., et al. (2024).
551   Analysis of Breast Cancer Mortality in the US—1975 to 2019. Jama *5405*,
552   233–241. 10.1001/jama.2023.25881.

553   3.   Fornier, M., and Fumoleau, P. (2012). The paradox of triple negative breast
554   cancer: Novel approaches to treatment. Breast J. *18*, 41–51. 10.1111/j.1524-
555   4741.2011.01175.x.

556   4.   Jin, J., Tao, Z., Cao, J., Li, T., and Hu, X. (2021). DNA damage response
557   inhibitors: An avenue for TNBC treatment. Biochim. Biophys. Acta - Rev.
558   Cancer *1875*, 188521. 10.1016/j.bbcan.2021.188521.

559   5.   Michaelis, M., Wass, M.N., and Cinatl, J. (2019). Drug-adapted cancer cell
560   lines as preclinical models of acquired resistance. Cancer Drug Resist. *2*, 447–
561   456. 10.20517/cdr.2019.005.

562   6.   Oellerich, T., Schneider, C., Thomas, D., Knecht, K.M., Buzovetsky, O.,
563   Kaderali, L., Schliemann, C., Bohnenberger, H., Angenendt, L., Hartmann, W.,
564   et al. (2019). Selective inactivation of hypomethylating agents by SAMHD1
565   provides a rationale for therapeutic stratification in AML. Nat. Commun. *10*.
566   10.1038/s41467-019-11413-4.

567   7.   Santoni-Rugiu, E., Melchior, L.C., Urbanska, E.M., Jakobsen, J.N., De
568   Stricker, K., Grauslund, M., and Sørensen, J.B. (2019). Intrinsic resistance to
569   EGFR-tyrosine kinase inhibitors in EGFR-mutant non-small cell lung cancer:
570   Differences and similarities with acquired resistance. Cancers (Basel). *11*, 1–
571   57. 10.3390/cancers11070923.

572   8.   Touat, M., Li, Y.Y., Boynton, A.N., Spurr, L.F., Iorgulescu, J.B., Bohrson, C.L.,
573   Cortes-Ciriano, I., Birzu, C., Geduldig, J.E., Pelton, K., et al. (2020).

Mechanisms and therapeutic implications of hypermutation in gliomas. Nature
580, 517–523. 10.1038/s41586-020-2209-9.

9.   Rothenburger, T., Thomas, D., Schreiber, Y., Wratil, P.R., Pflantz, T., Knecht,
K., Digianantonio, K., Temple, J., Schneider, C., Baldauf, H.M., et al. (2021).
Differences between intrinsic and acquired nucleoside analogue resistance in
acute myeloid leukaemia cells. J. Exp. Clin. Cancer Res. *40*, 1–19.
10.1186/s13046-021-02093-4.

10.   Juliano, R.L., and Ling, V. (1976). A surface glycoprotein modulating drug
permeability in Chinese hamster ovary cell mutants. BBA - Biomembr. *455*,
152–162. 10.1016/0005-2736(76)90160-7.

11.   Cole, S.P.C., Bhardwaj, G., Gerlach, J.H., Mackie, J.E., Grant, C.E., Almquist,
K.C., Stewart, A.J., Kurz, E.U., Duncan, A.M.V., and Deeley, R.G. (1992).
Overexpression of a transporter gene in a multidrug-resistant human lung
cancer cell line. Science (80-.). *258*, 1650–1654. 10.1126/science.1360704.

12.   Engelman, J., Zejnullahu, K., Mitsudomi, T., Song, Y., and Hyland (2007). MET
amplification leads to gefitinib resistance in lung cancer by activating ERBB3
signaling. Science (80-.). *316*, 1039–1043.

13.   Sharma, S. V., Haber, D.A., and Settleman, J. (2010). Cell line-based
platforms to evaluate the therapeutic efficacy of candidate anticancer agents.
Nat. Rev. Cancer *10*, 241–253. 10.1038/nrc2820.

14.   Nazarian, R., Shi, H., Wang, Q., Kong, X., Koya, R.C., Lee, H., Chen, Z., Lee,
M.K., Attar, N., Sazegar, H., et al. (2010). Melanomas acquire resistance to B-
RAF(V600E) inhibition by RTK or N-RAS upregulation. Nature *468*, 973–977.
10.1038/nature09626.

15.   Crystal, A.S., Shaw, A.T., Sequist, L. V., Friboulet, L., Niederst, M.J.,
Lockerman, E.L., Frias, R.L., Gainor, J.F., Amzallag, A., Greninger, P., et al.
(2014). Patient-derived models of acquired resistance can identify effective
drug combinations for cancer. Science (80-.). *346*, 1480–1486.
10.1126/science.1254721.

16.   Schneider, C., Oellerich, T., Baldauf, H.M., Schwarz, S.M., Thomas, D., Flick,
R., Bohnenberger, H., Kaderali, L., Stegmann, L., Cremer, A., et al. (2017).

SAMHD1 is a biomarker for cytarabine response and a therapeutic target in
acute myeloid leukemia. Nat. Med. *23*, 250–255. 10.1038/nm.4255.

17. Michaelis, M., Rothweiler, F., Barth, S., Cinat, J., Van Rikxoort, M.,
Löschmann, N., Voges, Y., Breitling, R., Von Deimling, A., Rödel, F., et al.
(2011). Adaptation of cancer cells from different entities to the MDM2 inhibitor
nutlin-3 results in the emergence of p53-mutated multi-drug-resistant cancer
cells. Cell Death Dis. *2*. 10.1038/cddis.2011.129.

18. Berlak, M., Tucker, E., Dorel, M., Winkler, A., McGearey, A., Rodriguez-Fos,
E., da Costa, B.M., Barker, K., Fyle, E., Calton, E., et al. (2022). Mutations in
ALK signaling pathways conferring resistance to ALK inhibitor treatment lead
to collateral vulnerabilities in neuroblastoma cells. Mol. Cancer *21*, 1–19.
10.1186/s12943-022-01583-z.

19. Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R.M., Ozenberger, B.A.,
Ellrott, K., Sander, C., Stuart, J.M., Chang, K., Creighton, C.J., et al. (2013).
The cancer genome atlas pan-cancer analysis project. Nat. Genet. *45*, 1113–
1120. 10.1038/ng.2764.

20. Klaas, E., Sung, E., Azizi, E., Martinez, M., Barpujari, A., Roberts, J., and
Lucke-Wold, B. (2023). Advanced breast cancer metastasized in the brain:
treatment standards and innovations. J. Cancer Metastasis Treat. *9*.
10.20517/2394-4722.2022.125.

21. Wang, B., Sun, T., Zhao, Y., Wang, S., Zhang, J., Wang, Z., Teng, Y.E., Cai,
L., Yan, M., Wang, X., et al. (2022). A randomized phase 3 trial of Gemcitabine
or Nab-paclitaxel combined with cisPlatin as first-line treatment in patients with
metastatic triple-negative breast cancer. Nat. Commun. *13*. 10.1038/s41467-
022-31704-7.

22. Liu, Y., Fan, L., Wang, Z.H., and Shao, Z.M. (2023). Nab-paclitaxel Followed
by Dose-dense Epirubicin/Cyclophosphamide in Neoadjuvant Chemotherapy
for Triple-negative Breast Cancer: A Phase II Study. Oncologist *28*, 86-e76.
10.1093/oncolo/oyac223.

23. Gumusay, O., Huppert, L.A., Magbanua, M.J.M., Wabl, C.A., Assefa, M.,
Chien, A.J., Melisko, M.E., Majure, M.C., Moasser, M., Park, J., et al. (2023). A

636 phase Ib/II study of eribulin in combination with cyclophosphamide in patients
637 with advanced breast cancer. Breast Cancer Res. Treat. *203*, 197–204.
638 10.1007/s10549-023-07073-0.

639 24. Kim, S.H., Im, S.A., Suh, K.J., Lee, K.H., Kim, M.H., Sohn, J., Park, Y.H., Kim,
640 J.Y., Jeong, J.H., Lee, K.E., et al. (2023). Clinical activity of nivolumab in
641 combination with eribulin in HER2-negative metastatic breast cancer: A phase
642 IB/II study (KCSG BR18-16). Eur. J. Cancer *195*, 113386.
643 10.1016/j.ejca.2023.113386.

644 25. Velikova, G., Morden, J.P., Haviland, J.S., Emery, C., Barrett-Lee, P., Earl, H.,
645 Bloomfield, D., Brunt, A.M., Canney, P., Coleman, R., et al. (2023).
646 Accelerated versus standard epirubicin followed by cyclophosphamide,
647 methotrexate, and fluorouracil or capecitabine as adjuvant therapy for breast
648 cancer (UK TACT2; CRUK/05/19): quality of life results from a multicentre,
649 phase 3, open-label, randomised, . Lancet Oncol. *24*, 1359–1374.
650 10.1016/S1470-2045(23)00460-6.

651 26. Takahashi, M., Cortés, J., Dent, R., Pusztai, L., McArthur, H., Kümmel, S.,
652 Denkert, C., Park, Y.H., Im, S.A., Ahn, J.H., et al. (2023). Pembrolizumab Plus
653 Chemotherapy Followed by Pembrolizumab in Patients with Early Triple-
654 Negative Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial.
655 JAMA Netw. Open *6*, E2342107. 10.1001/jamanetworkopen.2023.42107.

656 27. Liston, D.R., and Davis, M. (2017). Clinically Relevant Concentrations of
657 Anticancer Drugs◻: A Guide for Nonclinical Studies. Clin. Cancer Res., 3489–
658 3498. 10.1158/1078-0432.CCR-16-3083.

659 28. Kim, N., Hong, Y., Kwon, D., and Yoon, S. (2013). Somatic Mutaome Profile in
660 Human Cancer Tissues. Genomics Inform. *11*, 239. 10.5808/gi.2013.11.4.239.

661 29. Reynolds, I.S., Fichtner, M., McNamara, D.A., Kay, E.W., Prehn, J.H.M., and
662 Burke, J.P. (2019). Mucin glycoproteins block apoptosis; promote invasion,
663 proliferation, and migration; and cause chemoresistance through diverse
664 pathways in epithelial cancers. Cancer Metastasis Rev. *38*, 237–257.
665 10.1007/s10555-019-09781-w.

666 30. Pandey, K., Lee, E., Park, N., Hur, J., Cho, Y. Bin, Katuwal, N.B., Kim, S.K.,

Lee, S.A., Kim, I., An, H.J., et al. (2021). Deregulated immune pathway
associated with palbociclib resistance in preclinical breast cancer models:
Integrative genomics and transcriptomics. Genes (Basel). *12*, 1–14.
10.3390/genes12020159.

31. Chang, Y., Wang, Y., Li, B., Lu, X., Wang, R., Li, H., Yan, B., Gu, A., Wang,
W., Huang, A., et al. (2021). Whole-Exome Sequencing on Circulating Tumor
Cells Explores Platinum-Drug Resistance Mutations in Advanced Non-small
Cell Lung Cancer. Front. Genet. *12*, 1–11. 10.3389/fgene.2021.722078.

32. Patel, N.M., Geropoulos, G., Patel, P.H., Bhogal, R.H., Harrington, K.J.,
Singanayagam, A., and Kumar, S. (2023). The Role of Mucin Expression in the
Diagnosis of Oesophago-Gastric Cancer: A Systematic Literature Review.
Cancers (Basel). *15*. 10.3390/cancers15215252.

33. Kim, S.H., Ho, J.N., Jin, H., Lee, S.C., Lee, S.E., Hong, S.K., Lee, J.W., Lee,
E.S., and Byun, S.S. (2016). Upregulated expression of BCL2, MCM7, and
CCNE1 indicate cisplatin-resistance in the set of two human bladder cancer
cell lines: T24 cisplatin sensitive and T24R2 cisplatin resistant bladder cancer
cell lines. Investig. Clin. Urol. *57*, 63–72. 10.4111/icu.2016.57.1.63.

34. Feng, Z., Zhang, L., Zhou, J., Zhou, S., Li, L., Guo, X., Feng, G., Ma, Z.,
Huang, W., and Huang, F. (2017). mir-218-2 promotes glioblastomas growth,
invasion and drug resistance by targeting CDC27. Oncotarget *8*, 6304–6318.
10.18632/oncotarget.13850.

35. Peng, L., Zhao, M., Liu, T., Chen, J., Gao, P., Chen, L., Xing, P., Wang, Z., Di,
J., Xu, Q., et al. (2022). A stop-gain mutation in GXYLT1 promotes metastasis
of colorectal cancer via the MAPK pathway. Cell Death Dis. *13*, 1–12.
10.1038/s41419-022-04844-3.

36. Michaelis, M., Klassert, D., Barth, S., Suhan, T., Breitling, R., Mayer, B.,
Hinsch, N., Doerr, H.W., Cinatl, J., and Cinatl, J. (2009). Chemoresistance
acquisition induces a global shift of expression of aniogenesis-associated
genes and increased pro-angogenic activity in neuroblastoma cells. Mol.
Cancer *8*, 80. 10.1186/1476-4598-8-80.

37. Bahar, M.E., Kim, H.J., and Kim, D.R. (2023). Targeting the RAS/RAF/MAPK

pathway for cancer therapy: from mechanism to clinical studies. Signal
Transduct. Target. Ther. *8*. 10.1038/s41392-023-01705-z.

38.   Wang, P., Laster, K., Jia, X., Dong, Z., and Liu, K. (2023). Targeting CRAF
kinase in anti-cancer therapy: progress and opportunities. Mol. Cancer *22*, 1–
34. 10.1186/s12943-023-01903-x.

39.   Xu, J., Zhou, Y., Dong, K., Gong, J., Xiong, W., Wang, X., Gu, C., Lu, X. yu,
Huang, D. pei, Shen, X. dong, et al. (2023). Gene variation profile and it's
potential correlation with clinical characteristics in HBV-associated HCC
patients of Sichuan Han nationality in China. Asian J. Surg. *46*, 4371–4377.
10.1016/j.asjsur.2023.02.056.

40.   Weinstein, J.N. (2013). Cancer Genome Atlas Pan-cancer analysis project.
Nat Genet *45*, 113–1120. 10.3779/j.issn.1009-3419.2015.04.02.

41.   Lu, J.F., Pokharel, D., and Bebawy, M. (2015). MRP1 and its role in anticancer
drug   resistance.   Drug   Metab.   Rev.   *47*,   406–419.
10.3109/03602532.2015.1105253.

42.   Derry, W.B., Wilson, L., and Jordan, M.A. (1995). Substoichiometric Binding of
Taxol Suppresses Microtubule Dynamics. Biochemistry *34*, 2203–2211.
10.1021/bi00007a014.

43.   Snyder, J.P., Nettles, J.H., Cornett, B., Downing, K.H., and Nogales, E. (2001).
The binding conformation of Taxol in β-tubulin: A model based on electron
crystallographic density. Proc. Natl. Acad. Sci. U. S. A. *98*, 5312–5316.
10.1073/pnas.051309398.

44.   Jordan, M.A., and Wilson, L. (2004). Microtubules as a target for anticancer
drugs. Nat. Rev. Cancer *4*, 253–265. 10.1038/nrc1317.

45.   Smith, J.A., Wilson, L., Azarenko, O., Zhu, X., Lewis, B.M., Littlefield, B.A., and
Jordan, M.A. (2010). Eribulin binds at microtubule ends to a single site on
tubulin to suppress dynamic instability. Biochemistry *49*, 1331–1337.
10.1021/bi901810u.

46.   Baxter, J.S., Zatreanu, D., Pettitt, S.J., and Lord, C.J. (2022). Resistance to
DNA repair inhibitors in cancer. Mol. Oncol. *16*, 3811–3827. 10.1002/1878-
0261.13224.

47.   Bracht, K., Boubakari, Grünert, R., and Bednarski, P.J. (2006). Correlations
730   between the activities of 19 anti-tumor agents and the intracellular glutathione
731   concentrations in a panel of 14 human cancer cell lines: Comparisons with the
732   National Cancer Institute data. Anticancer. Drugs *17*, 41–51.
733   10.1097/01.cad.0000190280.60005.05.

734   48.   Michaelis, M., Schneider, C., Rothweiler, F., Rothenburger, T., Mernberger,
735   M., Nist, A., von Deimling, A., Speidel, D., Stiewe, T., and Cinatl, J. (2018).
736   TP53 mutations and drug sensitivity in acute myeloid leukaemia cells with
737   acquired MDM2 inhibitor resistance. bioRxiv, 404475.

738   49.   Michaelis, M., Wass, M.N., Reddin, I., Voges, Y., Rothweiler, F., Hehlgans, S.,
739   Cinatl, J., Mernberger, M., Nist, A., Stiewe, T., et al. (2020). YM155-adapted
740   cancer cell lines reveal drug- induced heterogeneity and enable the
741   identification of biomarker candidates for the acquired resistance setting.
742   Cancers (Basel). *12*, 1–17. 10.3390/cancers12051080.

743   50.   Karasaki, T., Moore, D.A., Veeriah, S., Naceur-Lombardelli, C., Toncheva, A.,
744   Magno, N., Ward, S., Bakir, M. Al, Watkins, T.B.K., Grigoriadis, K., et al.
745   (2023). Evolutionary characterization of lung adenocarcinoma morphology in
746   TRACERx. Nat. Med. *29*. 10.1038/s41591-023-02230-w.

747   51.   Martínez-Ruiz, C., Black, J.R.M., Puttick, C., Hill, M.S., Demeulemeester, J.,
748   Larose Cadieux, E., Thol, K., Jones, T.P., Veeriah, S., Naceur-Lombardelli, C.,
749   et al. (2023). Genomic–transcriptomic evolution in lung cancer and metastasis.
750   Nature *616*, 543–552. 10.1038/s41586-023-05706-4.

751   52.   Al Bakir, M., Huebner, A., Martínez-Ruiz, C., Grigoriadis, K., Watkins, T.B.K.,
752   Pich, O., Moore, D.A., Veeriah, S., Ward, S., Laycock, J., et al. (2023). The
753   evolution of non-small cell lung cancer metastases in TRACERx
754   10.1038/s41586-023-05729-x.

755   53.   Frankell, A.M., Dietzen, M., Al Bakir, M., Lim, E.L., Karasaki, T., Ward, S.,
756   Veeriah, S., Colliver, E., Huebner, A., Bunkum, A., et al. (2023). The evolution
757   of lung cancer and impact of subclonal selection in TRACERx. Nature *616*,
758   525–533. 10.1038/s41586-023-05783-5.

54.   Abbosh, C., Frankell, A.M., Harrison, T., Kisistok, J., Garnett, A., Johnson, L.,

Veeriah, S., Moreau, M., Chesh, A., Chaunzwa, T.L., et al. (2023). Tracking
early lung cancer metastatic dissemination in TRACERx using ctDNA. Nature
616, 553–562. 10.1038/s41586-023-05776-4.

55.   Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival:
Application to proliferation and cytotoxicity assays. J. Immunol. Methods *65*,
55–63. 10.1016/0022-1759(83)90303-4.

56.   Onafuye, H., Pieper, S., Mulac, D., Cinatl, J., Wass, M.N., Langer, K., and
Michaelis, M. (2019). Doxorubicin-loaded human serum albumin nanoparticles
overcome transporter-mediated drug resistance in drug-adapted cancer cells.
Beilstein J. Nanotechnol. *10*, 1707–1715. 10.3762/bjnano.10.166.

57.   Andrews S (2018). FastQC A Quality control tool for high throughput sequence
data. Babraham Bioinfo, 3–5.

58.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible
trimmer for Illumina sequence data. Bioinformatics *30*, 2114–2120.
10.1093/bioinformatics/btu170.

59.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and
SAMtools. Bioinformatics *25*, 2078–2079. 10.1093/bioinformatics/btp352.

60.   Burrows, M., and Wheeler, D. (1994). A block-sorting lossless data
compression algorithm. Algorithm, Data Compression, 18. 10.1.1.37.6774.

61.   Church, D.M., Schneider, V.A., Graves, T., Auger, K., Cunningham, F., Bouk,
N., Chen, H.C., Agarwala, R., McLaren, W.M., Ritchie, G.R.S., et al. (2011).
Modernizing reference genome assemblies. PLoS Biol. *9*.
10.1371/journal.pbio.1001091.

62.   McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky,
A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The genome
analysis toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. *20*, 1297–1303. 10.1101/gr.107524.110.

63.   Li, H. (2011). A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation from
sequencing data. Bioinformatics *27*, 2987–2993.

10.1093/bioinformatics/btr509.

64.   McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A.,
Flicek, P., and Cunningham, F. (2016). The Ensembl Variant Effect Predictor.
Genome Biol. *17*, 1–14. 10.1186/s13059-016-0974-4.

65.   Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang,
Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2019).
Variation across 141,456 human exomes and genomes reveals the spectrum
of loss-of-function intolerance across human protein-coding genes. bioRxiv,
531210. 10.1101/531210.

66.   Ghandi, M., Huang, F.W., Jané-Valbuena, J., Kryukov, G. V., Lo, C.C.,
McDonald, E.R., Barretina, J., Gelfand, E.T., Bielski, C.M., Li, H., et al. (2019).
Next-generation characterization of the Cancer Cell Line Encyclopedia.
Nature. 10.1038/s41586-019-1186-3.

67.   Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A.,
Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., et al. (2004). The
COSMIC (Catalogue of Somatic Mutations in Cancer) database and website.
Br. J. Cancer *2*, 355–358. 10.1038/sj.bjc.6601894.

68.   Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and
Vilo, J. (2019). g:Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). Nucleic Acids Res. *47*, W191–W198.
10.1093/nar/gkz369.

69.   Colaprico, A., Silva, T.C., Olsen, C., Garofano, L., Cava, C., Garolini, D.,
Sabedot, T.S., Malta, T.M., Pagnotta, S.M., Castiglioni, I., et al. (2016).
TCGAbiolinks: An R/Bioconductor package for integrative analysis of TCGA
data. Nucleic Acids Res. *44*, e71. 10.1093/nar/gkv1507.

70.   Grossman, R.L., Heath, A.P., Ferretti, V., Varmus, H.E., Lowy, D.R., Kibbe,
W.A., and Staudt, L.M. (2016). Toward a shared vision for cancer genomic
data. N. Engl. J. Med. *375*, 1109–1112. 10.1056/NEJMp1607591.

71.   Uhlen, M., Zhang, C., Lee, S., Sjöstedt, E., Fagerberg, L., Bidkhori, G.,
Benfeitas, R., Arif, M., Liu, Z., Edfors, F., et al. (2017). A pathology atlas of the
human cancer transcriptome. Science (80-.). *357*. 10.1126/science.aan2507.

72.   Ng, S.W.K., Mitchell, A., Kennedy, J.A., Chen, W.C., McLeod, J., Ibrahimova,
N., Arruda, A., Popescu, A., Gupta, V., Schimmer, A.D., et al. (2016). A 17-
gene stemness score for rapid determination of risk in acute leukaemia. Nature
540, 433–437. 10.1038/nature20598.

 Figure Legends

Figure 1: Confirmation of the resistance status of the project cell lines. A)
Panel of drug-naive (MDA-MB-468, HCC38, HCC1806) and drug-adapted Triple
Negative Breast Cancer cell lines. B) Left; dose response curve, bottom; IC$_{50}$ values,
right; resistance factor (see methods); when drug-naive and drug-adapted cell lines
are treated with the respective agent; cisplatin, doxorubicin, eribulin, paclitaxel,
gemcitabine, 5-Fluorouracil. Circles indicate drug-naive cell lines, crosses indicate
drug-adapted cell lines. Green; MDA-MB-468-derived, blue; HCC38-derived, orange;
HCC1806-derived. Data from n ≥ 3, statistics calculated using student t-test and
plotted with mean ± SD.

Figure 2: Characterization of drug-adapted cell lines. A) Diagram illustrating the
difference between; *Gained*, *De novo*, *Not-called*, *Lost* and *Shared* variants. B)
Count of *Gained* (blue) and *De novo* (green) variants, C) count of *Lost* (orange) and
Not-called (pink) variants, D) left panel; count of *Shared* (purple) variants, right
panel; two-fold increase or decrease of shared variants.

Figure 3: Novel candidates with link to therapy failure identified. A) Flow chart
of genes that have *de novo* variants observed in ≥2 sublines from >1 parental cell
line. B) Venn diagrams of exact *de novo* variants shared between sublines adapted
to the same drug. C) Summary of relatedness between sublines adapted from the
same parental (%).

Figure 4: Novel gene identified with potential relevance to drug resistance in
clinical samples. A) Increased variants and *de novo* and *gained* protein truncating
variants as input and screened with known TCGA variants. B) Summary of variants
identified and subsequent genes where the Kaplan-Meier graph was statistically
significant for high and low gene expression (see methods). C) *ADNP* Kaplan-Meier
identified in the doxorubicin, paclitaxel, gemcitabine, and 5-fluorouracil cell lines. D)
Left; novel genes associated with drug resistance, right; Kaplan-Meier example of a
novel gene KIAA0588. E) Left; genes found to have a role in drug resistance
mechanisms, right; Kaplan-Meier example of known gene TOP2A.

Figure 5: Complex sensitivity patterns to cytotoxic and DDR targeted agents.
A) Heat map of fold-resistance and collateral sensitivity to cytotoxic agents. B)

Summary of DNA repair pathways targeted by agents used in screening. C) Heat
map of fold-resistance and collateral sensitivity to DDR agents.

Figure 6: No trend of sensitivity patterns by Delta (Δ) method. Graphs
demonstrating negative correlation; collateral sensitivity in one agent but resistance
in to the other (blue), positive correlation; resistance to both agents (red) and no
statistical correlation (black) for each group of cells lines belong to MDA-MB-468,
HCC38 and HCC1806. For calculation of delta method values see methods.

Supplementary Figure 1: Chemo-naïve cell lines are clinically sensitive to
chemotherapy agents. IC_{50} values, of drug-naïve cell lines treated with the
respective agent; cisplatin, doxorubicin, eribulin, paclitaxel, gemcitabine, 5-
Fluorouracil. Green; MDA-MB-468-derived, blue; HCC38-derived, orange;
HCC1806-derived. Black line indicates known Cmax values for chemotherapy agent.
Data from n ≥ 3, statistics calculated using student t-test and plotted with mean ±
SD.
873
874 **Supplementary Figure 2: Variant counts.** A) Total number of variants called for the
875 panel of drug-naïve and drug resistant cell lines. B) Different type of variants called
876 for the panel of drug-naïve and drug resistant cell line including; missense,
877 synonymous, frameshift, inframe insertion, inframe deletion, stop lost, stop gain,
878 splice acceptor and splice donor variants.
879
880 **Supplementary Figure 3: *De novo* variant overlaps.** The number of *de novo*
881 variants found overlapped in A) drug-resistant cell lines adapted to the same drug, B)
882 drug-resistant cell lines adapted from the same parental cell line.
883
Supplementary Figure 4: Gene ontology (GO) terms related to gene variants
that changed in drug-resistant sublines. A) Number of variants considered
increased in drug-resistant sublines (*de novo* variants, *gained* variants and *shared*
variants which demonstrated ≥2 increase in variant allele frequency). B) Number of
variants considered decreased in drug-resistant sublines (*not called* variants, *lost*
variants and *shared* variants which demonstrated ≤2 decrease in variant allele
frequency). The number and overlapping GO terms found in increased and

decreased variants were compared between cell lines adapted to the same drug (C,
E) and cell lines derived from the same parental cell line (D, F). Green bars indicate
increased variants (A, C, D) and red bars indicate decreased variants (B, C, D).

Supplementary Figure 5: Chemo-naïve cell lines are clinically sensitive to DNA
damage response (DNA damage response) inhibitors. IC_{50} values, of drug-naïve cell
lines treated with the stated drug. Green; MDA-MB-468-derived, blue; HCC38-
derived, orange; HCC1806-derived. Black line indicates known Cmax values for
DDR agent. Data from n ≥ 3, statistics calculated using student t-test and plotted with
mean ± SD

Supplementary Table 1: Drug correlation of delta (Δ) values. The IC_{50} values
were transformed to ΔIC_{50} values for each drug (see methods) and correlated across
the drug panel, with linear regression analysis and statistical significance. Values in
table indicate r value of correlation where positive values indicate positive correlation
and negative values indicate negative correlation. P values of the correlation are
indicated in the blue color scheme, with light blue (p≤0.05) being the lowest
statistical significance, and dark blue (p≤0.00001) the highest statistical significance.

Supplementary File 1: Mean IC_{50} values, S.D and resistance factor for project panel
treated with chemotherapy agents and DNA damage response inhibitors.

Supplementary File 2: Basic variant characterization of the project panel

Supplementary File 3: Variants found to be; *de novo*, *gained*, *not called*, *lost* and
shared in drug-resistant cell lines

Supplementary File 4: List of genes that have *de novo* variants in ≥2 drug resistant
cell lines. Values in the table indicate the variant allele frequency of *de novo* variants
identified in stated genes. PMID for genes identified to be previously implicated in
cancer and drug resistance.

Supplementary File 5: Exact and same consequence variants Kaplan-Meir graphs
when stated gene is expressed high and low and patient is treated with stated drug.
Data extracted from the TCGA.

Supplementary File 6: Protein truncating variants Kaplan-Meir graphs when stated
gene is expressed high and low and patient is treated with stated drug. Data
extracted from the TCGA.

Supplementary File 7: Genes identified in both (i) Exact and same consequence
variants and (ii) protein truncating variants analysis. Analysis identifies variant type,
cell line and references if gene has been indicated in resistance to the stated drug,
or other drug resistance. Green highlighted rows indicate novel resistant candidates.

935
936

945   **Figure**   **2**

 Figure 3

953 **Figure 4**

 Figure 5

980 **Figure**