Axonal distribution of mitochondria maintains neuronal autophagy during aging via eIF2β

Kanako Shinno[1†], Yuri Miura[2], Koichi M. Iijima[3,4], Emiko Suzuki[1,5], Kanae Ando[1,6*]

1. Department of Biological Sciences, Graduate School of Science, Tokyo Metropolitan University, Hachioji, Tokyo, 192-0397, Japan
2. Research Team for Mechanism of Aging, Tokyo Metropolitan Institute for Geriatrics and Gerontology, Itabashi, Tokyo, 173-0015, Japan
3. Department of Alzheimer's Disease Research, National Center for Geriatrics and Gerontology, Obu, Aichi, 474-8511, Japan
4. Department of Experimental Gerontology, Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Aichi, 467-8603, Japan
5. Gene Network Laboratory, National Institute of Genetics and Department of Genetics, SOKENDAI, Mishima, Shizuoka, 411-8540, Japan
6. Department of Biological Sciences, School of Science, Tokyo Metropolitan University, Hachioji, Tokyo, 192-0397, Japan

*Correspondence: k_ando@tmu.ac.jp
†Current address: Department of Neuroscience and Pathobiology, Research Institute of Environmental Medicine, Nagoya University, Chikusa-Ku, Nagoya, Aichi, 464-8601, Japan

Summary

Neuronal aging and neurodegenerative diseases are accompanied by proteostasis collapse, while the cellular factors that trigger it have not been identified. Impaired mitochondrial transport in the axon is another feature of aging and neurodegenerative diseases. Using *Drosophila*, we found that genetic depletion of axonal mitochondria causes dysregulation of protein degradation. Axons with mitochondrial depletion showed abnormal protein accumulation and autophagic defects. Lowering neuronal ATP levels by blocking glycolysis did not reduce autophagy, suggesting that autophagic defects are associated with mitochondrial distribution. We found that eIF2β was increased by the depletion of axonal mitochondria via proteome analysis.

Phosphorylation of eIF2α, another subunit of eIF2, was lowered, and global translation
was suppressed. Neuronal overexpression of *eIF2β* phenocopied the autophagic defects
and neuronal dysfunctions, and lowering *eIF2β* expression rescued those perturbations
caused by depletion of axonal mitochondria. These results indicate the
mitochondria-eIF2β axis maintains proteostasis in the axon, of which disruption may
underlie the onset and progression of age-related neurodegenerative diseases.
42
43

Highlights
• Loss of axonal mitochondria impairs autophagy and accumulates proteins in the
axon
• Loss of axonal mitochondria increases eIF2β and decreases p-eIF2α
• Neuronal upregulation of eIF2β induces autophagic defects and locomotor
dysfunction
• Lowering eIF2β rescues autophagic defects caused by loss of axonal mitochondria
51
52

53   **Keywords**
54   Mitochondria; axonal transport; aging; neuronal proteostasis; protein aggregation;
55   autophagy; eukaryotic Initiation Factor 2β (eIF2β); *Drosophila*; neurodegeneration;
56   proteome
57
58

Introduction
Neurons have a morphologically complex architecture composed of microcompartments
and require tight regulation of the abundance of proteins and organelles spatially and
temporally[1]. Such control of protein amounts, or proteostasis, is essential for neuronal
functions[2] and is achieved through the orchestration of protein expression, folding,
trafficking, and degradation controlled by intrinsic and environmental signals[3].
Translation is initiated by the eukaryotic initiation factor 2 (eIF2) complex[4]. eIF2, a
heterotrimer of α, β, and γ subunits, transports Met-tRNA to the ribosome in a
GTP-dependent manner[5]. Under stressed conditions, phosphorylation of eIF2α
attenuates global translation and initiates translation of mRNAs related to the integrated
stress response (ISR)[6]. As for protein degradation, autophagy and proteasome are major
systems that maintain proteostasis[7]. The proteasome degrades unnecessary proteins,

followed by regulated ubiquitination processes[8], and autophagy removes damaged or
harmful components, including large protein aggregates and organelles, through
catabolism (selective autophagy)[9]. In addition to autophagy induced by acute stressors,
a basal level of selective autophagy mediates the global turnover of damaged proteins[10].
75
Such constitutive autophagy decreases during aging, which may underlie declines in the
structural and functional integrity of neurons[11]. Decreased protein degradation and
accumulation of abnormal proteins also contribute to increased risks of
neurodegenerative diseases. Age-related neurodegenerative diseases such as
Alzheimer's disease and Parkinson's disease are often associated with the accumulation
of misfolded proteins such as amyloid-β, tau, and α-synuclein[12]. Enhancement of
autophagy mitigates age-related dysfunctions and neurodegeneration caused by
proteotoxic stress[13]. However, it is not fully understood how aging disrupts the
regulation of this constitutive autophagy.
85
Neurons are also highly energy-demanding cells. At nerve terminals, action potentials
trigger the release of neurotransmitters via exocytosis of synaptic vesicles, which
requires a constant supply of ATP and calcium buffering[14]. Such neuronal activity relies
on mitochondrial functions[15], and mitochondria are actively transported from their
major sites of biogenesis in soma to axons[16]. However, the axonal transport of
mitochondria declines during aging[17,18,19]. Reduced axonal transport of mitochondria is
thought to contribute to age-related declines in neuronal functions[17,19,20,21]. The number
of functional mitochondria in synapses is reduced in the brains of patients suffering
from age-related neurodegenerative diseases such as Alzheimer's disease[22], and
mutations in genes involved in mitochondrial dynamics are linked to neurodegenerative
diseases[23]. The mislocalization of mitochondria is sufficient to cause age-dependent
neurodegeneration in *Drosophila* and mice[24,25], indicating that the proper distribution of
mitochondria is essential to maintain neuronal functions. Thus, depletion of functional
mitochondria from axons and proteostasis collapse are common features of aging and
neurodegenerative diseases.
101
102 Mitochondrial transport is regulated by a series of molecular adaptors that mediate the
103 attachment of mitochondria to molecular motors[16]. In *Drosophila*, mitochondrial
104 transport is mediated by milton and Miro, which attaches mitochondria to microtubules
105 via kinesin heavy chain[26,27]. In the absence of milton or Miro, synaptic terminals and
106 axons lack mitochondria, although mitochondria are numerous in the neuronal cell

body[28]. We previously reported that RNAi-mediated knockdown of *milton* or *Miro* in neurons causes a reduction in axonal mitochondria, age-dependent locomotor defects[29], and age-dependent neurodegeneration in neuropile area starting around 30 days after eclosion (day-old)[24], and enhances axon degeneration caused by human tau proteins[24], suggesting that these flies can be used as a model to analyze the effect of depletion of axonal mitochondria during aging. In this study, we investigated a causal relationship between mitochondrial distribution and neuronal proteostasis by using neuronal knockdown of *milton*. We found that depletion of axonal mitochondria reduced autophagy and increased the accumulation of aggregated proteins in the axon prior to gross neurodegeneration. Proteome analysis and follow-up biochemical analyses revealed that neuronal knockdown of *milton* increased eIF2β levels and lowered phosphorylation of eIF2αin the axon. In addition, *milton* knockdown suppressed global translation. Overexpression of *eIF2β* was sufficient to decrease autophagy and induce neuronal dysfunction, and genetic suppression of *eIF2β* restored autophagy and improved neuronal function in the *milton* knockdown background. These findings suggest that loss of axonal mitochondria and elevated levels of eIF2β mediate proteostasis collapse and neuronal dysfunction during aging.

Results

Depletion of axonal mitochondria by knockdown of *milton* or *Miro* causes protein accumulation in the axon

In *Drosophila*, mitochondrial transport is mediated by milton and Miro, which attach mitochondria to microtubules via kinesin heavy chain[26,27] (Figure 1A). It has been reported that expression of *milton* RNAi in neurons via pan-neuronal elav-GAL4 driver reduced milton protein levels in *Drosophila* head lysate to 40% and mito-GFP signals in axons to 50%[24,29].

To test how loss of axonal mitochondria affects proteostasis in neurons, we first examined the accumulation of ubiquitinated proteins. At 14-day-old, more ubiquitinated proteins were deposited in the brains of *milton* knockdown flies than in those of age-matched control flies (Figure 1B, $p < 0.005$ between control RNAi and *milton* RNAi). There was no significant increase in ubiquitinated proteins in *milton* knockdown flies at 1-day old, suggesting that the accumulation of ubiquitinated proteins caused by *milton* knockdown is age-dependent (Figure S1). We also analyzed the effect of the neuronal knockdown of *Miro*, a partner of milton, on the accumulation of

ubiquitin-positive proteins. Since severe knockdown of *Miro* in neurons causes lethality, we used UAS-*Miro* RNAi strain with low knockdown efficiency, whose expression driven by elav-GAL4 caused 30% reduction of *Miro* mRNA in head extract[24]. Although there was a tendency for increased ubiquitin-positive puncta in *Miro* knockdown brains, the difference was not significant (Figure 1B, $p>0.05$ between control RNAi and *Miro* RNAi). These data suggest that the depletion of axonal mitochondria induced by *milton* knockdown leads to the accumulation of ubiquitinated proteins before neurodegeneration occurs.

It has been reported that ubiquitinated proteins accumulate with aging[30]; thus, we analyzed the accumulation of ubiquitinated proteins in aged brains (30-day-old) with *milton* knockdown. The number of puncta of ubiquitinated proteins did not significantly differ between control and *milton* knockdown flies or between control and *Miro* knockdown flies (Figure 1C, $p > 0.05$). These results suggest that depletion of axonal mitochondria may have more impact on proteostasis in young neurons than in old neurons.

We examined the ultrastructure of presynaptic terminals and cell bodies in photoreceptor neurons with *milton* knockdown by transmission electron microscopy in 27-day-old flies (Figure 1D). As previously reported[24], the number of mitochondria in presynaptic terminals decreased in *milton* knockdown (Figure 1E). The swelling of presynaptic terminals, characterized by the enlargement and roundness, was not reported at 3-day-old[24] but observed at this age with about 4% of total presynaptic terminals (Figure 1F, asterisks).

Some presynaptic terminals of *milton* knockdown neurons contained dense materials (Figure 1F and G, arrowheads). Dense materials are rarely found in age-matched control neurons, indicating that *milton* knockdown induces abnormal protein accumulation in the presynaptic terminals (Figure 1G and H). In *milton* knockdown neurons, dense materials are found in swollen presynaptic terminals more often than in presynaptic terminals without swelling, suggesting a positive correlation between the disruption of proteostasis and axonal damage (Figure 1G). In contrast, dense materials were not observed in cell bodies in the *milton* knockdown retina (Figure 1H). These results indicate that the depletion of axonal mitochondria induces protein accumulation in the axon.

Depletion of axonal mitochondria impairs protein degradation pathways

Since abnormal proteins were accumulated in *milton* knockdown brains, we next examined if protein degradation pathways were suppressed. We analyzed autophagy via

western blotting of the autophagy markers LC3 and p62[31]. During autophagy
progression, LC3 is conjugated with phosphatidylethanolamine to form LC3-II, which
localizes to isolation membranes and autophagosomes. LC3-I accumulation occurs
when autophagosome formation is impaired, and LC3-II accumulation is associated
with lysosomal defects[31,32]. p62 is an autophagy substrate, and its accumulation suggests
autophagic defects[31,32]. We found that *milton* knockdown increased LC3-I, and the
LC3-II/LC3-I ratio was lower in *milton* knockdown flies than in control flies at
14-day-old (Figure 2A). We also analyzed p62 levels in head lysates sequentially
extracted using detergents with different stringencies (1% Triton X-100 and 2% SDS).
Western blotting revealed that p62 levels were increased in the brains of 14-day-old of
milton knockdown flies (Figure 2B). The increase in the p62 level was significant in the
Triton X-100-soluble fraction but not in the SDS-soluble fraction (Figure 2B),
suggesting that depletion of axonal mitochondria impairs the degradation of
less-aggregated proteins. Proteasome activity was also significantly decreased in brains
with neuronal knockdown of *milton* (Figure 2C, $p < 0.005$).
At 30-day-old, LC3-I was still higher, and the LC3-II/LC3-I ratio was lower, in *milton*
knockdown compared to the control (Figure 2D). At this age, *milton* knockdown
increased p62 significantly in 1% Triton X-100 fraction and 2% SDS fraction (Figure
2E). Proteasome activities were also decreased in *milton* knockdown flies at 30-day-old
(Figure 2F). These results indicate that depletion of axonal mitochondria impairs protein
degradation pathways.
200

ATP deprivation does not impair autophagy
milton knockdown downregulates ATP in the axon[33]. To examine whether the disruption
of protein degradation pathways by *milton* knockdown is due to ATP deprivation, we
investigated the effects of knocking down phosphofructokinase (*Pfk*), a rate-limiting
enzyme in glycolysis, on protein degradation pathways. Neuronal knockdown of *Pfk*
was reported to lower ATP levels in brain neurons[33]. *Pfk* knockdown and *milton*
knockdown decreased ATP to similar levels (Figure 3A-C). However, in contrast with
milton knockdown, *Pfk* knockdown did not affect the levels of LC3-I, LC3-II or the
LC3-II/LC3-I ratio (Figure 3D). *Pfk* knockdown decreased p62 level (Figure 3E),
suggesting that autophagy is promoted. On the other hand, proteasome activity was
decreased by *Pfk* knockdown (Figure 3F). These results suggest that the downregulation
of axonal ATP upon depletion of axonal mitochondria decreases proteasome activity, but
not autophagy.
214

Proteome analysis suggests that depletion of axonal mitochondria causes disruption of autophagy and premature aging

To identify the pathways that mediate the decrease in autophagy in *milton* knockdown brains, we performed proteome analysis to systematically detect differentially expressed proteins upon neuronal knockdown of *milton*. We analyzed flies at 7- and 21-day-old, the age before autophagic defects are detected and the age just before the onset of neurodegeneration, respectively (Figure 4A). 1039 proteins were detected by liquid chromatography-tandem mass spectrometry (LC-MS/MS). Expression of 36 proteins was significantly increased (22 proteins) or decreased (14 proteins) by *milton* knockdown at 7-day-old (Figure 4B, Tables 1 and S1). At 21-day-old, the expression of 41 proteins (31 upregulated and ten downregulated proteins) was significantly altered in *milton* knockdown flies compared with control flies (Figure 4C, Tables 1 and S1). The "Interaction search" algorithm using KeyMolnet showed that proteins whose expression was significantly altered in the brains of *milton* knockdown flies at both 7- and 21-day-old were closely associated with the autophagic pathway (Table 2). Proteins involved in pathways characteristics of aging, such as the immune response (Transcriptional regulation by STAT), cancer (Transcriptional regulation by SMAD, Transcriptional regulation by myc), longevity (Transcriptional regulation by FOXO, Sirtuin signaling pathway), and stress responses (HSP90 signaling pathway, MAPK signaling pathway)[34,35], were enriched in the proteome profiles of *milton* knockdown flies compared with those of control flies at 7-day-old (Table 2), suggesting that depletion of axonal mitochondria accelerates aging in the brain.

Depletion of axonal mitochondria upregulates eIF2β and decreases phosphorylation of eIF2α

Differentially expressed proteins at 7-day-old flies may reflect alterations that are causal for autophagic defects. We noticed that the expression level of eIF2β was 2.465-fold higher in the brains of *milton* knockdown flies than in those of control flies (Figure 4B and D). Upregulation of eIF2β in the brains of *milton* knockdown flies was confirmed by western blotting. *milton* knockdown increased eIF2β protein levels more than twice (Figure 4E), but did not change the level of *eIF2β* mRNA (Figure 4F).

We also investigated age-dependent changes in eIF2β by western blotting of control flies at 7-, 21-, 35-, and 49-, and 63-day-old. eIF2β levels increased during aging until 49-day-old (Figure 4G). These results suggest that upregulation of eIF2β in *milton* knockdown fly brain reflects early an onset of age-dependent increase of eIF2β levels.

eIF2β is a subunit of the eukaryotic initiation factor 2 (eIF2) complex, which is critical

for translation initiation and the integrated stress response (ISR)[4]. eIF2 is a heterotrimer
of α, β, and γ subunits, and eIF2α is phosphorylated during the ISR[6]. As for the other
subunits of eIF2 complex, proteome analysis did not detect a significant difference in
the protein levels of eIF2α and eIF2γ between *milton* knockdown and control flies at 7-
and 21-day-old (Figure 4D). Western blotting of brain lysates showed that *milton*
knockdown reduced eIF2α levels (Figure 5A), while p-eIF2α levels were not
significantly affected (Figure 5B).
To analyze local changes of eIF2α and p-eIF2α, we carried out immunostaining. We
focused on the mushroom body, where axons, dendrites, and cell bodies can be easily
identified (Figure 5C). Both eIF2α and p-eIF2α were downregulated in the cell body
(Kenyon cells) and dendritic (Calyxes) regions of the brains of *milton* knockdown flies
(Figure 5D). In axons (lobe tips), *milton* knockdown did not affect eIF2α (Figure 5E, *p*
= 0.271) but significantly downregulated p-eIF2α (Figure 5E). The ratio of p-eIF2α to
eIF2α was lower in the axon but not in the soma or dendritic region. These results
suggest that axonal distribution of mitochondria regulate the level of overall eIF2α
protein and local p-eIF2α.
267
Depletion of axonal mitochondria suppressed global translation
Phosphorylation of eIF2α induces conformational changes in the eIF2 complex and
inhibits global translation[36]. To analyze the effects of *milton* knockdown on translation,
we performed polysome gradient centrifugation to examine the level of ribosome
binding to mRNA. Since p-eIF2α was downregulated, we hypothesized that *milton*
knockdown would enhance translation. However, unexpectedly, we found that *milton*
knockdown significantly reduced the level of mRNAs associated with polysomes
(Figure 6A and B). We also compared the level of translation between the brains of
control and *milton* knockdown flies by assessing the incorporation of puromycin (Figure
6C). Puromycin incorporation was lower in the brains of *milton* knockdown flies than in
those of control flies, while it was not statistically significant (Figure 6C, indicated by a
bracket). These data suggest that the depletion of axonal mitochondria suppresses global
translation.
281
eIF2β upregulation reduces the level of p-eIF2α, impairs autophagy, and decreases
locomotor function
We were motivated to ask if eIF2β upregulation mediates autophagic defects caused by
milton knockdown. If so, neuronal overexpression of *eIF2β* would also induce
autophagy impairment. Neuronal overexpression of *eIF2β* increased LC3-II, while the

LC3-II/LC3-I ratio was not significantly different (Figure 7A and B). Overexpression of

eIF2β significantly increased the p62 level in the Triton X-100-soluble fraction (Figure

7C, 4-fold vs. control, $p < 0.005$ (1% Triton X-100)) but not in the SDS-soluble fraction

(Figure 7C, 2-fold vs. control, $p = 0.062$ (2% SDS)), as observed in brains of *milton*

knockdown flies (Figure 2B). These data suggest that neuronal overexpression of *eIF2β*

accumulates autophagic substrates.

Since *milton* knockdown reduced the p-eIF2α level (Figure 5E), we asked whether an

increase in eIF2β affects p-eIF2α. Neuronal overexpression of *eIF2β* did not affect the

eIF2α level but significantly decreased the p-eIF2α level (Figure 7D and E).

Depletion of axonal mitochondria causes age-dependent decline in locomotor function[24].

We found that neuronal overexpression of *eIF2β* also caused locomotor dysfunction

(Figure 7F). Locomotor functions were significantly impaired in those flies at

20-day-old and worsened further during aging (Figure 7F, compare 4-, 20-, and

30-day-old). We asked if *eIF2β* overexpression causes neurodegeneration, as depletion

of axonal mitochondria in the photoreceptor neurons causes axon degeneration in an

age-dependent manner[24]. *eIF2β* overexpression in photoreceptor neurons tends to

increase neurodegeneration in aged flies, while it was not statistically significant

(p>0.05, Figure S2).

These data indicate that an increase of eIF2β in neurons phenocopies depletion of

axonal mitochondria, including suppression of autophagy and age-dependent locomotor

dysfunction, and suggest that increase of eIF2β mediates these phenotypes downstream

of loss of axonal mitochondria.

Lowering *eIF2β* rescues autophagic impairment and locomotor dysfunction

induced by *milton* knockdown

Finally, we investigated whether suppression of eIF2β rescues autophagy impairment

and locomotor dysfunction caused by neuronal knockdown of *milton*. Null mutants and

flies with RNAi-mediated knockdown of *eIF2β* in neurons did not survive. Flies lacking

one copy of the *eIF2β* gene survived without any gross abnormality, and the level of

eIF2β mRNA in these flies was about 80% of that in control flies (Figure 8A). *eIF2β*

heterozygosity did not affect the eIF2α and p-eIF2α levels (Figure S3A and B).

Neuronal knockdown of *milton* causes accumulation of autophagic substrate p62 in the

Triton X-100-soluble fraction (Figure 2B), and we tested if lowering eIF2β ameliorates

it. We found that *eIF2β* heterozygosity caused a mild increase in LC3-I levels and

decreases in LC3-II levels, resulting in a significantly lower LC3-II/LC3-I ratio in

milton knockdown flies (Figure 8B). *eIF2β* heterozygosity decreased the p62 level in

the Triton X-100-soluble fraction in the brains of *milton* knockdown flies (Figure 8C).

The p62 level in the SDS-soluble fraction, which is not sensitive to *milton* knockdown

(Figure 2B), was not affected (Figure 8C). These results suggest that suppression of

eIF2β ameliorates the impairment of autophagy caused by *milton* knockdown.

eIF2β heterozygosity also rescued locomotor dysfunction induced by *milton* knockdown.

milton knockdown flies with *eIF2β* heterozygosity exhibited better locomotor function

than *milton* knockdown alone (Figure 8D). The *milton* mRNA level was not increased in

these flies, indicating that the rescue effect in the *eIF2β* heterozygous background was

not mediated by an increase in the *milton* mRNA level (Figure S3C). These data suggest

that eIF2β upregulation mediates autophagy impairment and locomotor dysfunction

caused by the depletion of axonal mitochondria.

Discussion

The depletion of axonal mitochondria and accumulation of abnormal proteins are both

characteristics of aged brains[37,38]. Proteostasis perturbations trigger the formation of

pathological aggregates and increase the risks of neurodegenerative diseases during

aging. By using neuronal *milton* knockdown to deplete mitochondria from the axon, we

provide evidence that loss of axonal mitochondria drives age-related proteostasis

collapse via eIF2β (Figure 9). We observed declines in autophagy-mediated degradation

of less-aggregated proteins and proteasome activity in *milton* knockdown flies (Figure

2). Accumulation of ubiquitinated proteins and changes in age-related pathways started

prematurely in *milton* knockdown flies (Figure 1 and Table 2). *milton* knockdown

increased eIF2β and lowered eIF2α phosphorylation in young fly brain (Figure 4 and 5).

Overexpression of *eIF2β* phenocopied the effects of *milton* knockdown, including

reduced autophagy and accelerated age-related locomotor defects (Figure 7).

Furthermore, lowering *eIF2β* levels suppressed the impairment of autophagy and

locomotor dysfunction induced by *milton* knockdown (Figure 8). From these results, we

propose that upregulation of eIF2β downstream of depletion of axonal mitochondria

drives age-dependent collapse of proteostasis (Figure 9). Our results suggest that

mitochondrial distribution and eIF2β are part of the mechanisms constituting

proteostasis.

milton knockdown causes loss of mitochondria in the axon and accumulation of

mitochondria in the soma. Thus, the detrimental effects may be mediated by the

accumulation of mitochondria. However, degeneration induced by *milton* knockdown is

prominent in the axon and not detected in the cell body[24]. Furthermore, abnormal
protein accumulation was observed in the axon (Figure 1), and p-eIF2α/eIF2α was
decreased in the neurites but not in the soma (Figure 5), suggesting that proteostasis
defects studied in this work are caused by depletion of mitochondria rather than
accumulation of mitochondria. Further analyses to dissect the effects of *milton*
knockdown on proteostasis and translation in the cell body and axon by experiments
with spatial resolution would be needed.

Our results suggest that the loss of axonal mitochondria is an event upstream of
proteostasis collapse during aging. The number of puncta of ubiquitinated proteins was
higher in milton knockdown at 14-day-old, but there was no significant difference at
30-day-old (Figure 1). Proteome analyses also showed that age-related pathways, such
as immune responses, are enhanced in young flies with *milton* knockdown (Table 2). We
also found that eIF2β protein levels increase in an age-dependent manner until
49-day-old and reduces after that (Figure 4G). In the brains with neuronal knockdown
of *milton*, eIF2β levels were higher at 7-day-old than those in control and lower at the
21-day-old (Figure 4D and Supplementary table). These results suggest that *milton*
knockdown is likely accelerating age-dependent changes rather than increasing their
magnitude. Disruption of proteostasis is expected to contribute neurodegeneration[38],
and it would be interesting to analyze the sequence of protein accumulation and axonal
degeneration in *milton* knockdown ([24,29] and Figure 1) in detail with higher time
resolution.

Our results revealed that eIF2β regulates autophagy and maintains proteostasis during
aging. eIF2β is a component of eIF2, which meditates translational regulation and ISR
initiation. When ISR is activated, phosphorylated eIF2α suppresses global translation
and induces translation of ATF4, which mediates transcription of autophagy-related
genes[39,40]. Since ISR can positively regulate autophagy, we suspected that suppression
of ISR underlies a reduction in autophagic protein degradation. We found neuronal
knockdown of *milton* reduced phosphorylated eIF2α, suggesting that ISR is reduced
(Figure 5). However, we also found that global translation was reduced (Figure 6).
Increased levels of eIF2β might disrupt the eIF2 complex or alter its functions. The
stoichiometric mismatch caused by an imbalance of eIF2 components may inhibit ISR
induction. Supporting this model, we found that eIF2β upregulation reduced the levels
of p-eIF2α (Figure 7). It is also possible that eIF2β mediates autophagy defects via
mechanisms independent of ISR since eIF2β has functions independent of eIF2[41,42]. For

example, suppression of *eIF2β* has been reported to slow down cancer cell growth[41]. In

developing neurons, eIF2β can directly interact with the translational repressor Kra to

regulate midline axon guidance[42]. Our results also suggest that milton knockdown and

overexpression of eIF2β affect autophagy via increased LC3-I abundance (Figures 2 and

7), suggesting an unconventional mechanism of autophagy suppression. To our

knowledge, the roles of eIF2β in aging and autophagy independent of ISR have not been

reported. Our results revealed a novel function of eIF2β to maintain proteostasis during

aging, while further investigation is required to elucidate underlying mechanisms.

How depletion of axonal mitochondria upregulates eIF2β is currently under

investigation. A major mitochondrial function is ATP production, and depletion of

axonal mitochondria downregulates ATP in axons[33]. However, we found that ATP

deprivation did not always suppress autophagy (Figure 3), suggesting it is unlikely to be

involved in the mechanisms that induce eIF2β upregulation. Mitochondria also serve as

signaling hubs for translation and protein degradation. Mitochondrial proteins are

regulated by co-translational protein quality control, and mitochondrial damage induces

translational stalling of mitochondrial outer membrane-associated *complex-I 30 kD*

subunit (*C-I30*) mRNA[43]. Additionally, the mitochondrial outer membrane ubiquitin

ligase MITOL (also known as MARCHF5) ubiquitinates and regulates not only

mitochondrial proteins such as Mfn2[44] but also microtubule-associated[45] and

endoplasmic reticulum[46] proteins. These findings indicate that mitochondria serve as

local signaling centers for proteostasis maintenance, and eIF2β levels may also be

regulated by mechanisms related to mitochondria.

In conclusion, our results suggest that axonal mitochondria and eIF2β form an axis to

maintain constitutive autophagy. Suppression of *eIF2β* rescued autophagic defects and

neuronal dysfunction upon loss of axonal mitochondria. Since eIF2β is conserved across

many species, including *Drosophila* and humans, our results suggest that eIF2β may be

a possible therapeutic target for aging and diseases associated with mitochondrial

mislocalization.

Materials and methods

Fly stocks and husbandry

Flies were maintained in standard cornmeal medium (10% glucose, 0.7% agar, 9%

cornmeal, 4% yeast extract, 0.3% propionic acid, and 0.1% n-butyl p-hydroxybenzoate) at 25°C under light–dark cycles of 12:12 h. The flies were transferred to fresh food vials for every 2–3 days. UAS-*milton* RNAi (v41508) was from VDRC and outcrossed to [w1118] for five generations in our laboratory. Transgenic fly lines carrying UAS-*luciferase* RNAi control for *milton* RNAi) was reported previously[24]. GMR-gal4, Elav-gal4, UAS-*Pfk* RNAi (Bloomington stock center #36782), UAS-*luciferase* RNAi (Bloomington stock center #31603) (control for *Pfk* RNAi), UAS-*GFP* (used for control for UAS-*eIF2β*) and UAS-*eIF2β* (eIF2β^{EY08063}, Bloomington stock center #17425) were from the Bloomington stock center. *eIF2β* heterozygous strain (PBac{SAstopDsRed} LL07719, DGRC#142114) was from KYOTO *Drosophila* Stock Center. UAS-*mitoGFP* was a kind gift from Drs. W. M. Saxton (University of California, Santa Cruz).

Immunohistochemistry and image acquisition

Fly brains were dissected in PBS and fixed for 45 minutes in formaldehyde (4% v/v in PBS) at room temperature. After incubation in PBST containing 0.1% Triton X-100 for 10 min three times, samples were incubated for 1h at room temperature in PBST containing 1% normal goat serum (Wako, #143-06561) and then incubated overnight with the primary antibody (anti-ubiquitin antibody Ubi-1 (Thermo Fisher #13-1600) (1:50), anti-eIF2α (abcam #ab26197) (1:50) and anti-p-eIF2α (Cell signaling #3398S) (1:50)) diluted in 1% NGS/PBST at 4 °C. Samples were then washed for 10 min with PBST including 0.1% Triton X-100 three times and incubated with the secondary antibody for overnight at 4° C. Brains were mounted in Vectashield (Vectorlab Cat#H-1100) and analyzed under a confocal microscope (Nikon). Quantitative analysis was performed using ImageJ (National Institutes of Health) with maximum projection images derived from Z-stack images acquired with same settings. Puncta were identified with mean intensity and area using ImageJ. For eIF2α and p- eIF2α immunostaining, the mushroom body was detected by mitoGFP expression.

Electron microscopy

Proboscis was removed from decapitated heads, which were then incubated in primary fixative solution (2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium cacodylate buffer) at R.T. for 2 hours. After washing heads with 3% sucrose in 0.1 M sodium cacodylate buffer, fly heads were post-fixed for 1 hour in secondary fixation (1% osmium tetroxide in 0.1 M sodium cacodylate buffer) on ice. After washing with H$_2$O, heads were dehydrated with ethanol and infiltrated with propylene oxide and Epon

mixture (TAAB and Nissin EM) for 3 hours. After infiltration, specimens were

embedded with an Epon mixture at 70°C for 2-3 days. Thin sections (70 nm) of laminas

were collected on copper grids. The sections were stained with 5% uranyl acetate in

50% ethanol and Reynolds' lead citrate solution. Electron micrographs were obtained

with a CCD Camera mounted on a JEM-1400 plus electron microscope (Jeol Ltd.).

Quantitation was performed using ImageJ (National Institutes of Health).

SDS–PAGE and immunoblotting

Western blotting was performed as reported previously[24]. Briefly, heads of 10-20

Drosophila were homogenized with SDS-Tris-Glycine sample buffer (0.312M Tris, 5%

SDS, 8% glycerol, 0.0625% BPB, 10% β-mercaptoethanol, 10μg/mL leupeptin, 0.4 μM

Pefabloc, 10mM β-glycerphosphate, 10mM NaF) and after boiling at 95°C for 2 minutes,

it was centrifuged at 13,200 rpm, and the supernatant was used as a sample. For p62

western blot, fly heads were homogenized with 1% PBST and after centrifugation at

13,200 rpm, the supernatant was mixed 1:1 SDS-Tris-Glycine sample buffer, and boiled

at 95 °C for 2 minutes. The pellet was dissolved with 2% SDS in PBS, then centrifuged

again at 13,200 rpm. The supernatant was mixed 1:1 SDS-Tris-Glycine sample buffer

and then boiled at 95 °C for 2 minutes. SDS–PAGE for western blotting was performed

using 15%(w/v) (LC3), 7.5%(w/v) (p62), 10% (w/v) (eIF2α, β, and p-eIF2α)

polyacrylamide gels. After electrophoresis, they were transferred to PVDF membrane

(Merck Millipore) using a transfer device (BIO-RAD). After transfer, the membrane

was blocked with 5% skim milk/TBST (50 mM Tris (pH 7.5), 0.15 M NaCl, 0.05%

Tween20) for 1 hour and incubated with primary antibody listed below overnight at 4 °C.

Membranes were rinsed twice with TBST containing 0.65M NaCl and once with TBST

containing 0.15M NaCl. After incubation with the secondary antibody at room

temperature for 1 hour, membranes were rinsed twice with TBST containing 0.65M

NaCl and once with TBST containing 0.15M NaCl. After incubation with Immobilon

Western Chemiluminescent HRP Substrate (Merck Millipore), chemiluminescent

signals were detected with Fusion FX (Vilber). Experiments were repeated at least 3

times with independent cohorts of flies.

Primary antibodies: anti-LC3 antibody Atg8 (Merck Millipore #ABC975) (1:1000),

anti-p62 antibody Ref2P (abcam #ab178440) (1:750), anti-eIF2β antibody (1:1500),

anti-eIF2α antibody (abcam #ab26197) (1:1000), anti-p-eIF2α antibody (Cell signaling

#3398S) (1:2000), anti-actin antibody (SIGMA A2066) (1:3000) and anti-β tubulin

antibody (Sigma #T9026) (1:800,000). Polyclonal anti-eIF2β antibody was raised

against a synthetic peptide (CGLEDDTKKEDPQDEA) corresponding to the C-terminal

residues 29-43 of *Drosophila* eIF2β (1:1500).

Secondary antibodies: Peroxidase-conjugated goat anti-mouse IgG antibody (Dako

#P0447) (1:2000), peroxidase-conjugated pig anti-rabbit IgG antibody (Dako #P0399)

(1:2000)

Proteasome assay

Heads from ten flies were homogenized in 150 µl of buffer B (25 mM Tris-HCl [pH

7.5], 2 mM ATP, 5 mM MgCl2, and 1 mM dithiothreitol). Proteasome peptidase activity

in the lysates was measured with a synthetic peptide substrate,

succinyl-Leu-Leu-Val-Tyr-7-amino-4-methyl-coumarin (Suc-LLVY-AMC) (Cayman).

Luminescence was measured on a multimode plate reader 2300 Enspire (PerkinElmer).

Experiments were repeated at least 3 times with independent cohorts of flies.

ATP assay

Heads from the 10 flies were homogenized in 50 µl of 6 M guanidine-HCl in

extraction buffer (100 mM Tris and 4 mM EDTA, pH 7.8) to inhibit ATPases.

Samples were boiled for 5 min and centrifuged. The supernatant was diluted 4% with

extraction buffer and mixed with a reaction solution (ATP Determination kit,

Invitrogen). Luminescence was measured on a multimode plate reader 2300 Enspire

(PerkinElmer). The relative ATP levels were calculated by dividing the luminescence by

the total protein concentration, which was determined by the Bradford method.

Experiments were repeated at least 3 times with independent cohorts of flies.

Proteomic assay and pathway analysis

Sample preparation

Heads from the 35 flies were homogenized in 110 µl of extraction buffer (0.25%

RapiGest SF, 50mM ammonium bicarbonate, 10mM dithiothreitol, 10µg/mL leupeptin,

0.4 µM Pefabloc, 10mM β-glycerphosphate, 10mM NaF). Homogenized samples were

centrifuged and boiled for 5 min. After quantification of the protein concentration

using a Pierce® 660 nm Protein Assay (Thermo Fisher Scientific), 10 µg proteins from

each sample were reduced using 5 mM tris (2-carboxyethyl) phosphine hydrochloride

(TCEP-HCl; Thermo Fisher Scientific) at 60°C for 1 h, alkylated using 15 mM

iodoacetamide (Fujifilm Wako Pure Chemical, Osaka, Japan) at room temperature for

30 min, and then digested using 1.5 µg Trypsin Gold (Mass Spectrometry Grade;

Promega, Madison, WI, USA) at 37°C for 17 h. The digests were acidified by the

addition of trifluoroacetic acid (TFA), incubated at 37°C for 30 min, and then

centrifuged at 17,000 ×*g* for 10 min to remove the RapiGest SF. The supernatants were collected and desalted using MonoSpin™ C18 (GL Sciences, Tokyo, Japan). The resulting eluates were concentrated *in vacuo*, dissolved in 2% MeCN containing 0.1% formic acid (FA), and subjected to LC-MS/MS analysis.

LC-MS/MS analysis and database search

LC-MS/MS analyses were performed on an Ultimate 3000 RSLCnano system (Thermo Fisher Scientific) coupled to a Q Exactive hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher Scientific) equipped with a nano electron spray ionization (ESI) source. The LC system was equipped with a trap column (C18 PepMap 100, 0.3 × 5 mm, 5 µm, Thermo Fisher Scientific) and an analytical column (NTCC-360/75-3-125, Nikkyo Technos, Tokyo, Japan). Peptide separation was performed using a 90-min gradient of water/0.1% FA (mobile phase A) and MeCN/0.1% FA (mobile phase B) at a flow rate of 300 nL/min. Elution was performed as follows: 0–3 min, 2% B; 3–93 min, 2%–40% B; 93–95 min, 40%–95% B; 95–105 min, 95% B; 105–107 min, 95%–2% B; and 107–120 min, 2% B. The mass spectrometer was operated in data-dependent acquisition mode. The MS parameters were as follows: spray voltage, 2.0 kV; capillary temperature, 275°C; S-lens RF level, 50; scan type, full MS; scan range, *m/z* 350–1500; resolution, 70,000; polarity, positive; automatic gain control target, 3×10^{6}; and maximum injection time, 100 msec. The MS/MS parameters were as follows: resolution, 17,500; automatic gain control target, 1×10^{5}; maximum injection time, 60 msec; normalized collision energy (NCE), 27; dynamic exclusion, 15 sec; loop count, 10; isolation window, 1.6 *m/z*; charge exclusion: unassigned, 1 and ≥8; and injection volume, 1 µL (containing 0.5 µg protein). Measurements were made in duplicate for each sample.

The identification of proteins and label-free quantification (LFQ) of the detected peptides were performed using Proteome Discoverer software ver. 2.4 (Thermo Fisher Scientific). The analytical parameters used for the database search were as follows: parent mass error tolerance, 10.0 ppm; fragment mass error tolerance, 0.02 Da; search engine, sequest HT; protein database, Drosophila melanogaster (Fruit fly: SwissProt Tax ID=7227); enzyme name, trypsin (full); maximum number of missed cleavages, 2; dynamic modification, oxidation (methionine), phosphorylation (serine, threonine, tyrosine), acetyl (lysine), GG (lysine); N-terminal modification, Met-loss (methionine), and Met-loss+acetyl (methionine); static modification, carbamidomethylation (cysteine) and FDR confidence, High < 0.01, 0.01 ≤ Medium < 0.05, 0.05 ≤ Low. The parameters

for LFQ were as follows: precursor abundance, based on area; and normalization mode,
total peptide amount.
The abundance ratio of *milton* RNAi to control RNAi at 7- or 21-day-old was calculated.
We considered proteins with an abundance ratio of ≥2.0 or ≤ 0.5 and an ANOVA
P-value of < 0.05 based on volcano plots to be differentially expressed of *milton* RNAi.
To extract molecular networks biologically relevant to the proteins that are differentially
expressed in *milton* RNAi, pathway analysis was performed using KeyMolnet (KM
Data Inc., Tokyo, Japan).
582
RNA extraction and quantitative real-time PCR analysis
Heads from more than 25 flies were mechanically isolated, and total RNA was extracted
using ISOGEN (NipponGene) followed by reverse-transcription using PrimeScript RT
reagent kit (Takara). The resulting cDNA was used as a template for PCR with
THUNDERBIRD SYBR qPCR mix (TOYOBO) on a Thermal Cycler Dice real-time
system TP800 (Takara). Expression of genes of interest was standardized relative to
rp49. Relative expression values were determined by the ΔΔCT method (Livak and
Schmittgen, 2001). Experiments were repeated three times, and a representative result
was shown.
Primers were designed using DRSC FlyPrimerBank (Harvard Medical School). Primer
sequences are shown below;
eIF2β for 5′-GGACGACGACAAGAGCGAAG -3′
eIF2β rev 5′-CGGTCGCATCACGAACTTTG -3′
milton for 5′-GGCTTCAGGGCCAGGTATCT-3′
milton rev 5′-GCCGAACTTGGCTGACTTTG-3′
Actin for 5′-TGCACCGCAAGTGCTTCTAA-3′
Actin rev 5′-TGCTGCACTCCAAACTTCCA-3′
rp49 for 5′-GCTAAGCTGTCGCACAAATG-3′
rp49 rev 5′- GTTCGATCCGTAACCGATGT-3′
602
Polysome gradient centrifugation
30 heads were homogenized in 150 µl of lysis buffer (25 mM Tris pH 7.5, 50 mM
MgCl2, 250 mM NaCl, 1 mM DTT, 0.5 mg/ml cycloheximide, 0.1 mg/ml heparin). The
lysates were centrifuged at 13,200 rpm at 4 °C for 5 minutes, and the supernatant was
collected. The samples containing 38µg of RNA was layered gently on top of a 10–50%
w/w sucrose gradient (50 mM Tris pH 7.5, 50 mM MgCl2, 250 mM NaCl, 0.1 mg/ml
heparin, 0.5 mg/ml cycloheximide in 5 ml polyallomer tube) and centrifuged at 37,000

rpm at 4°C for 150 minutes in a himac CP-NX ultracentrifuge using a P50AT rotor. Samples were fractionated from top to bottom, and absorbance at OD260 nm was analyzed by a Plate reader (EnSpire). Experiments were repeated at least 3 times with independent cohorts of flies.

Puromycin analysis

13-day-old flies were starved for 6 hr and fed 600 μM puromycin (Sigma) or 600 μM puromycin/35 mM cycloheximide (Sigma) in 5% sucrose solution for 20 hr. Incorporated puromycin was quantified by western blot with anti-puromycin antibody and normalized with actin. Experiments were repeated at least 3 times with independent cohorts of flies.

Histological analysis

Fly heads were fixed in Bouin's fixative solution for 48 h at room temperature, incubated for 24 h in 50 mM Tris/150 mM NaCl, and embedded in paraffin. Serial sections (7 μm thickness) through the entire heads were stained with hematoxylin and eosin and examined by bright-field microscopy. Images of the sections that include the lamina were captured with Keyence microscope BZ-X700 (Keyence), and the vacuole area was measured using ImageJ (National Institutes of Health).

Climbing assay

The climbing assay was performed as previously described[24]. Flies were placed in an empty plastic vial (2.5 cm in diameter × 10 cm in length). The vial was gently tapped to knock the flies to the bottom, and the number of flies reached to the top, middle and bottom areas of the vials in 10 s was counted. Experiments were repeated 10 times, and the mean percentage of flies in each area and standard deviations were calculated. Experiments were repeated with independent cohorts more than three times, and a representative result was shown.

Statistics

The number of replicates, what n represents, precision measurements, and the meaning of error bars are indicated in Figure Legends. Data are shown as means ± SEM. For pairwise comparisons, Student's t-test was performed with Microsoft Excel (Microsoft). For multiple comparisons, data were analyzed using one-way ANOVA with Tukey's HSD multiple-comparisons test in the GraphPad Prism 6.0 software (GraphPad Software, Inc., La Jolla, CA). Results with a p-value of less than 0.05 were considered

646   to be statistically significant.

Acknowledgments

The authors thank the Bloomington stock center; TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947); the Kyoto Drosophila Stock Center and the Vienna Drosophila RNAi Center for fly stocks. The authors thank Dr. Masayuki Miura from Department of Pharmaceutical Science, University of Tokyo, for proteasome activity assay protocol; Dr. Shin-ichi Hisanaga from the Department of Biological Sciences, Tokyo Metropolitan University, for critical comments; Dr. Taro Saito, Dr. Akiko Asada from the Department of Biological Sciences, Tokyo Metropolitan University, Dr. Michiko Sekiya from Department of Alzheimer's Disease Research, National Center for Geriatrics and Gerontology, and Dr. Seiji Watanabe, Dr. Koji Yamanaka from Department of Neuroscience and Pathobiology, Research Institute of Environmental Medicine, Nagoya University for technical supports.

Funding

This work was supported by the Sasakawa Scientific Research Grant (2021-4087) (to K.S.), the Takeda Science Foundation (to KA), Hoansha foundation grant (to KA), a research award from the Japan Foundation for Aging and Health (to KA), the Novartis Foundation (Japan) for the promotion of Science (to KA), JSPS KAKENHI Grant-in-Aid for Scientific Research on Challenging Research (Exploratory) JP19K21593 (to KA), JSPS KAKENHI Grant-in-Aid for Scientific Research(B) JP24K02860 (to KA), NIG-JOINT (National Institute of Genetics, 71A2018, 25A2019) (to K.A.) and TMU strategic research fund for social engagement (to KA).

Author contributions

K.S. and K.A. designed the experiments, interpreted the data, and wrote the manuscript; K.S., Y.M., K.M.I and E.S. conducted the experiments, and K.S., Y.M., E.S., and K.A. analyzed the data. All authors read and approved the manuscript.

Declaration of interests

The authors declare no competing interests.

Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author upon request.

Figure legends

Figure 1. Knockdown of *milton* or *Miro* causes protein accumulation in the axon

(A) Schematic representation of the mitochondrial transport machinery. Knockdown of *milton*, an adapter protein for mitochondrial transport, depletes mitochondria in the axon. (B, C) Ubiquitinated proteins in brains with neuronal knockdown of *milton* or *Miro*. Brains dissected at 14-day-old (B) or 30-day-old (C) were immunostained with an antibody against ubiquitin. Firefly *luciferase* RNAi was used as a control. Representative images (left) and quantitation of the number of ubiquitin-positive puncta (right) are shown. Scale bars of hemibrains, 100 µm, Scale bars of high magnifications, 10 µm. Means ± SE, n = 8. *N.S.*, $p > 0.05$; ***$p < 0.005$ (one-way analysis of variance (ANOVA) followed by Tukey's honestly significant difference (HSD) *post hoc* test). (D) Cross sections in the lamina and in the retina were used to analyze the ultrastructure of synapses and cell bodies, respectively. *milton* RNAi was expressed in the retina and neurons via a combination of GAL4 drivers, a pan-retinal gmr-GAL4 and pan-neuronal elav-GAL4. (E) Quantitation of the number of mitochondria in a presynaptic terminal from transmission electron micrographs. 180 presynaptic terminals from cross-sections of the lamina from three brains were analyzed. ***$p < 0.005$ (Chi-square test). (F, G) Presynaptic terminals of photoreceptor neurons of control and *milton* knockdown flies. Photoreceptor neurons are highlighted in blue. Swollen presynaptic terminals (asterisks in (F)), characterized by the enlargement and higher circularity, were found more frequently in *milton* knockdown neurons. Arrowheads indicate presynaptic terminals with dense materials. Scale bars, 2 µm. Representative images (Left) and quantitation (Right) are shown. 918–1118 from three heads were quantified for the percentage of swollen presynaptic terminals, and 180 presynaptic terminals from three heads were quantified for the size of presynaptic terminals. Mean ± SE, **$p < 0.01$, ***$p < 0.005$ (Student's *t*-test). (G) Dense materials (arrowheads in (G)) in the presynaptic terminals of *milton* knockdown neurons. Scale bars, 2 µm. The ratio of presynaptic terminals containing dense materials were quantified from 918–1118 presynaptic terminals from

three heads. Mean ± SE, ***p < 0.005 (Student's t-test). (H) Cell bodies of
photoreceptor neurons of control and *milton* knockdown flies. Scale bars, 2 μm. Flies
were 27-day-old.

Figure 2. *milton* knockdown impairs protein degradation pathways
(A, B) Western blotting of head extracts of control and *milton* knockdown flies with
antibodies against LC3 (A) and Ref2P, the fly homolog of mammalian p62 (B). For the
analyses of p62 levels, heads were extracted with 1% Triton X-100 or 2% SDS (B).
Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown.
Actin was used as a loading control. Means ± SE, n = 6 (LC3), n = 3 (p62). (C)
Proteasome activity in head extracts of control and *milton* knockdown flies was
measured by hydrolysis of Suc-LLVY-AMC at 14-day-old. Means ± SE, n = 3. (D, E)
Western blotting of head extracts of 30-day-old control and *milton* knockdown flies.
Blotting was performed with anti-LC3 (D) and anti-p62 (E) antibodies. Representative
blots (left) and quantitation (right) are shown. Actin was used as a loading control.
Means ± SE, n = 6 (LC3), n = 3 (p62). (F) Proteasome activity in head extracts of
30-day-old control and *milton* knockdown flies. Means ± SE, n = 3. *N.S.*, $p > 0.05$; *$p <$
0.05; **$p < 0.01$; ***$p < 0.005$ (Student's t-test).

Figure 3. ATP deprivation does not impair autophagy
(A-C) ATP levels in brain extracts of control and *milton* knockdown flies (A) and
control and *Pfk* knockdown flies (B) and comparison of the effects of *milton*
knockdown and *Pfk* knockdown on ATP levels (C). Flies were 14-day-old. Means ± SE,
n = 3. (D, E) Western blotting of head extracts of flies with neuronal expression of
control or *Pfk* RNAi. Blotting was performed with anti-LC3 (D) and anti-p62 (E)
antibodies. For analyses of p62 levels, heads were extracted with 1% Triton X-100 or
2% SDS. Representative blots (left) and quantitation (right) are shown. Actin was used
as a loading control. Means ± SE, n = 6 (LC3), n = 3 (p62). (F) Proteasome activity in
head lysates of flies with neuronal expression of control or *Pfk* RNAi was measured by
hydrolysis of Suc-LLVY-AMC. Means ± SE, n = 3. *N.S.*, $p > 0.05$; *$p < 0.05$; **$p <$
0.01; ***$p < 0.005$ (Student's t-test). Flies were at 14-day-old.

Figure 4. *milton* knockdown upregulates eIF2β in young flies
(A) Timing of proteome analysis and phenotypes observed in *milton* knockdown flies.
(B, C) Scatter plots of the $\log_2$ abundance ratio (x-axis) against the $-\log_{10}$ p-value
(y-axis) of proteins at 7-day-old (B) and 21-day-old (C). (D) eIF2 subunit protein levels

from proteome analysis of *milton* knockdown flies compared to those of control flies. (E) Western blotting of head extracts of flies expressing control or *milton* RNAi in neurons with an anti-eIF2β antibody. Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown. Tubulin was used as a loading control. Means ± SE, n = 6. (F) *eIF2β* mRNA levels quantified by qRT-PCR. Means ± SE, n = 4. (G) Western blotting of head extracts of wild-type flies with an anti-eIF2β antibody. Flies were 7-, 21-, 35-, 49- and 63-day-old. Representative blots (left) and quantitation (right) are shown. Tubulin was used as a loading control. Means ± SE, n = 3, *$p < 0.05$ (one-way analysis of variance (ANOVA) followed by Dunnett's multiple comparison test).

Figure 5. *milton* knockdown decreases phosphorylation of eIF2α

(A, B) Western blotting of head extracts with anti-eIF2α (A) and anti-p-eIF2α (B) antibodies. Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown. Tubulin was used as a loading control. Means ± SE, n = 6. (C) A schematic representation of the axon (Lobe tips), the cell body region (Kenyon cells), and dendritic region (Calyxes) in the fly brain. Scale bars, 100µm. (D, E) Immunostaining with anti-eIF2α and anti-p-eIF2α antibodies. The mushroom body was identified by expression of mito-GFP. Scale bars, 20µm. The signal intensities of eIF2α and p-eIF2α in axons, dendrites, and cell bodies were quantified and are shown as ratios relative to the control. Means ± SE, n□=12. *N.S.*, $p > 0.05$; *$p < 0.05$; **$p < 0.01$; ***$p < 0.005$ (Student's *t*-test).

Figure 6. *milton* knockdown suppressed global translation

(A) Representative polysome traces of head lysates of control and *milton* knockdown flies. (B) Quantitation of polysome fraction. The relative ratio of area under the curve (AUC) of polysome fractions (sedimentation 28%-50%). Means ± SE, n = 3. ***$p < 0.005$ (Student's *t*-test) (C) Western blotting of head lysates of control and *milton* knockdown flies fed puromycin alone or puromycin and cycloheximide (CHX) with an anti-puromycin antibody. Flies were 14-day-old. Actin was used as a loading control. Representative blots (left) and quantitation (right) are shown. Means ± SE, n = 3. Student's *t*-test.

Figure 7. eIF2β upregulation impairs autophagy and decreases locomotor function

(A) *eIF2β* mRNA levels in head extracts of flies with UAS-*eIF2β* driven by elav-Gal4 (*eIF2β* OE) or UAS-GFP driven by elav-Gal4 (control) were quantified by qRT-PCR. Flies were 2-day-old. Means ± SE, n=4. (B, C) Western blotting of head extracts with

anti-LC3 (B) and anti-p62 (C) antibodies. Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown. Tubulin and actin were used as loading controls. Means ± SE, n = 3 (p62), n = 5 (LC3). (D, E) Western blotting of head extracts with anti-eIF2α (D) and anti-p-eIF2α (E) antibodies. Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown. Tubulin was used as a loading control. Means ± SE, n = 6. (F) Climbing assay revealed early-onset of age-dependent locomotor defects in *eIF2β*-overexpressing flies. Means ± SE, n = 5. *N.S.*, $p > 0.05$; ***$p < 0.005$ (Student's *t*-test).

Figure 8. Lowering *eIF2β* rescues autophagic impairment and locomotor dysfunction induced by *milton* knockdown

(A) *eIF2β* mRNA levels with one disrupted copy of the *eIF2β* gene (*eIF2β*SAstopDsRed/+ (*eIF2β* -/+)). Head extracts of flies 2-3 day-old were analyzed by qRT-PCR. Means ± SE, n = 3. (B, C) Western blotting of head extracts of flies with neuronal expression of *milton* RNAi with or without *eIF2β* heterozygosity with anti-LC3 (B) and anti-p62 (C) antibodies. Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown. Actin was used as a loading control. Means ± SE, n = 5 (LC3), n = 3 (p62). (D) The climbing ability of 20-day-old flies expressing *milton* RNAi with or without *eIF2β* heterozygosity. Means ± SE, n = 15. *N.S.*, $p > 0.05$; *$p < 0.05$; ***$p < 0.005$ (Student's *t*-test).

Figure 9. Graphical abstract: the mitochondria-eIF2β axis in the axon maintains neuronal proteostasis during aging

Supplemental Figure 1. Ubiquitinated proteins in brains with neuronal knockdown of *milton* at 1-day-old

Brains dissected at 1-day-old were immunostained with an antibody against ubiquitin. Representative images (left) and quantitation of the number of ubiquitin-positive puncta (right) are shown. Scale bars of hemibrains, 100 µm, Means ± SE, n = 8. *N.S.*, $p > 0.05$ (Student's *t*-test).

Supplemental Figure 2. Histology analysis of fly heads with *eIF2β* overexpression

The morphology of the eye with eIF2β overexpression. The dotted lines indicate the retina. Vacuoles are indicated by arrows. Representative images (left) and quantification of vacuole area (right). The flies were 40-day-old. Means ± SE, n = 5-13 *N.S.*, $p > 0.05$

(Student's *t*-test).

Supplemental Figure 3. **Lowering the eIF2β level does not affect the levels of eIF2α**

and p-eIF2α.

(A, B) Blotting was performed with anti-eIF2α (A) and anti-p-eIF2α (B) antibodies.

Flies were 14-day-old. Representative blots (left) and quantitation (right) are shown.

Tubulin was used as a loading control. Means ± SE, n = 6. (C) *eIF2β* gene disruption

does not affect the knockdown efficiency of *milton*. *milton* mRNA levels in head

extracts were quantified by qRT-PCR. Flies were 2-day-old. Means ± SE, n = 3.

N.S., $p > 0.05$; *$p < 0.05$ (Student's *t*-test).

References

1.   Yerbury, J.J., Ooi, L., Dillin, A., Saunders, D.N., Hatters, D.M., Beart, P.M.,

Cashman, N.R., Wilson, M.R., and Ecroyd, H. (2016). Walking the tightrope:

proteostasis and neurodegenerative disease. J. Neurochem. *137*, 489-505.

10.1111/jnc.13575.

2.   Hetz, C. (2021). Adapting the proteostasis capacity to sustain brain healthspan.

Cell *184*, 1545-1560. 10.1016/j.cell.2021.02.007.

3.   Balch, W.E., Morimoto, R.I., Dillin, A., and Kelly, J.W. (2008). Adapting

proteostasis for disease intervention. Science *319*, 916-919.

10.1126/science.1141448.

4.   Kimball, S.R. (1999). Eukaryotic initiation factor eIF2. Int J Biochem Cell Biol

31, 25-29. 10.1016/s1357-2725(98)00128-9.

5.   Jackson, R.J., Hellen, C.U., and Pestova, T.V. (2010). The mechanism of eukaryotic

translation initiation and principles of its regulation. Nat. Rev. Mol. Cell Biol.

11, 113-127. 10.1038/nrm2838.

6.   Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A., and Gorman,

A.M. (2016). The integrated stress response. EMBO Rep *17*, 1374-1395.

10.15252/embr.201642195.

7.   Kroemer, G., Marino, G., and Levine, B. (2010). Autophagy and the integrated stress

response. Mol. Cell *40*, 280-293. 10.1016/j.molcel.2010.09.023.

8.   Nandi, D., Tahiliani, P., Kumar, A., and Chandu, D. (2006). The

ubiquitin-proteasome system. J Biosci *31*, 137-155. 10.1007/BF02705243.

9.   Glick, D., Barth, S., and Macleod, K.F. (2010). Autophagy: cellular and molecular

mechanisms. J. Pathol. *221*, 3-12. 10.1002/path.2697.

10.   Vargas, J.N.S., Hamasaki, M., Kawabata, T., Youle, R.J., and Yoshimori, T. (2023).
The mechanisms and roles of selective autophagy in mammals. Nat. Rev. Mol. Cell
Biol. *24*, 167-185. 10.1038/s41580-022-00542-2.

11.   Aman, Y., Schmauck-Medina, T., Hansen, M., Morimoto, R.I., Simon, A.K., Bjedov,
I., Palikaras, K., Simonsen, A., Johansen, T., Tavernarakis, N., et al. (2021).
Autophagy in healthy aging and disease. Nat Aging *1*, 634-650.
10.1038/s43587-021-00098-4.

12.   Ross, C.A., and Poirier, M.A. (2004). Protein aggregation and neurodegenerative
disease. Nat. Med. *10 Suppl*, S10-17. 10.1038/nm1066.

13.   Rubinsztein, D.C., Marino, G., and Kroemer, G. (2011). Autophagy and aging. Cell
146, 682-695. 10.1016/j.cell.2011.07.030.

14.   Vos, M., Lauwers, E., and Verstreken, P. (2010). Synaptic mitochondria in synaptic
transmission and organization of vesicle pools in health and disease. Front
Synaptic Neurosci *2*, 139. 10.3389/fnsyn.2010.00139.

15.   Cheng, A., Hou, Y., and Mattson, M.P. (2010). Mitochondria and neuroplasticity.
ASN Neuro *2*, e00045. 10.1042/AN20100019.

16.   Hollenbeck, P.J., and Saxton, W.M. (2005). The axonal transport of mitochondria.
J. Cell Sci. *118*, 5411-5419. 10.1242/jcs.02745.

17.   Takihara, Y., Inatani, M., Eto, K., Inoue, T., Kreymerman, A., Miyake, S., Ueno,
S., Nagaya, M., Nakanishi, A., Iwao, K., et al. (2015). In vivo imaging of axonal
transport of mitochondria in the diseased and aged mammalian CNS. Proc Natl Acad
Sci U S A *112*, 10515-10520. 10.1073/pnas.1509879112.

18.   Milde, S., Adalbert, R., Elaman, M.H., and Coleman, M.P. (2015). Axonal transport
declines with age in two distinct phases separated by a period of relative stability.
Neurobiol. Aging *36*, 971-981. 10.1016/j.neurobiolaging.2014.09.018.

19.   Vagnoni, A., Hoffmann, P.C., and Bullock, S.L. (2016). Reducing Lissencephaly-1
levels augments mitochondrial transport and has a protective effect in adult
Drosophila neurons. J. Cell Sci. *129*, 178-190. 10.1242/jcs.179184.

20.   Morsci, N.S., Hall, D.H., Driscoll, M., and Sheng, Z.H. (2016). Age-Related Phasic
Patterns of Mitochondrial Maintenance in Adult Caenorhabditis elegans Neurons.
J. Neurosci. *36*, 1373-1385. 10.1523/JNEUROSCI.2799-15.2016.

21.   Adalbert, R., and Coleman, M.P. (2013). Review: Axon pathology in age-related
neurodegenerative disorders. Neuropathol. Appl. Neurobiol. *39*, 90-108.
10.1111/j.1365-2990.2012.01308.x.

22.   Duncan, J.E., and Goldstein, L.S. (2006). The genetics of axonal transport and

898   axonal transport disorders. PLoS Genet. *2*, e124. 10.1371/journal.pgen.0020124.

899   23.   Chen, H., and Chan, D.C. (2009). Mitochondrial dynamics--fusion, fission, movement,

900   and mitophagy--in neurodegenerative diseases. Hum. Mol. Genet. *18*, R169-176.

901   10.1093/hmg/ddp326.

902   24.   Iijima-Ando, K., Sekiya, M., Maruko-Otake, A., Ohtake, Y., Suzuki, E., Lu, B.,

903   and Iijima, K.M. (2012). Loss of axonal mitochondria promotes tau-mediated

904   neurodegeneration and Alzheimer's disease-related tau phosphorylation via PAR-1.

905   PLoS Genet. *8*, e1002918. 10.1371/journal.pgen.1002918.

906   25.   Lopez-Domenech, G., Higgs, N.F., Vaccaro, V., Ros, H., Arancibia-Carcamo, I.L.,

907   MacAskill, A.F., and Kittler, J.T. (2016). Loss of Dendritic Complexity Precedes

908   Neurodegeneration in a Mouse Model with Disrupted Mitochondrial Distribution in

909   Mature Dendrites. Cell Rep *17*, 317-327. 10.1016/j.celrep.2016.09.004.

910   26.   Guo, X., Macleod, G.T., Wellington, A., Hu, F., Panchumarthi, S., Schoenfield,

911   M., Marin, L., Charlton, M.P., Atwood, H.L., and Zinsmaier, K.E. (2005). The GTPase

912   dMiro is required for axonal transport of mitochondria to Drosophila synapses.

913   Neuron *47*, 379-393. 10.1016/j.neuron.2005.06.027.

914   27.   Glater, E.E., Megeath, L.J., Stowers, R.S., and Schwarz, T.L. (2006). Axonal

915   transport of mitochondria requires milton to recruit kinesin heavy chain and is

916   light chain independent. J. Cell Biol. *173*, 545-557. 10.1083/jcb.200601067.

917   28.   Stowers, R.S., Megeath, L.J., Gorska-Andrzejak, J., Meinertzhagen, I.A., and

918   Schwarz, T.L. (2002). Axonal transport of mitochondria to synapses depends on

919   milton, a novel Drosophila protein. Neuron *36*, 1063-1077.

920   10.1016/s0896-6273(02)01094-2.

921   29.   Iijima-Ando, K., Hearn, S.A., Shenton, C., Gatt, A., Zhao, L., and Iijima, K. (2009).

922   Mitochondrial mislocalization underlies Abeta42-induced neuronal dysfunction in

923   a Drosophila model of Alzheimer's disease. PLoS One *4*, e8310.

924   10.1371/journal.pone.0008310.

925   30.   Tonoki, A., Kuranaga, E., Tomioka, T., Hamazaki, J., Murata, S., Tanaka, K., and

926   Miura, M. (2009). Genetic evidence linking age-dependent attenuation of the 26S

927   proteasome with the aging process. Mol. Cell. Biol. *29*, 1095-1106.

928   10.1128/MCB.01227-08.

929   31.   Klionsky, D.J., Abdel-Aziz, A.K., Abdelfatah, S., Abdellatif, M., Abdoli, A., Abel,

930   S., Abeliovich, H., Abildgaard, M.H., Abudu, Y.P., Acevedo-Arozena, A., et al.

931   (2021). Guidelines for the use and interpretation of assays for monitoring

932   autophagy (4th edition)(1). Autophagy *17*, 1-382. 10.1080/15548627.2020.1797280.

32.   Bartlett, B.J., Isakson, P., Lewerenz, J., Sanchez, H., Kotzebue, R.W., Cumming,

R.C., Harris, G.L., Nezis, I.P., Schubert, D.R., Simonsen, A., and Finley, K.D.

(2011). p62, Ref(2)P and ubiquitinated proteins are conserved markers of neuronal

aging, aggregate formation and progressive autophagic defects. Autophagy *7*,

572-583. 10.4161/auto.7.6.14943.

33. Oka, M., Suzuki, E., Asada, A., Saito, T., Iijima, K.M., and Ando, K. (2021).

Increasing neuronal glucose uptake attenuates brain aging and promotes life span

under dietary restriction in Drosophila. iScience *24*, 101979.

10.1016/j.isci.2020.101979.

34. Zia, A., Pourbagher-Shahri, A.M., Farkhondeh, T., and Samarghandian, S. (2021).

Molecular and cellular pathways contributing to brain aging. Behav Brain Funct

17, 6. 10.1186/s12993-021-00179-9.

35. Haigis, M.C., and Yankner, B.A. (2010). The aging stress response. Mol. Cell *40*,

333-344. 10.1016/j.molcel.2010.10.002.

36. Wek, R.C. (2018). Role of eIF2alpha Kinases in Translational Control and Adaptation

to Cellular Stress. Cold Spring Harb Perspect Biol *10*.

10.1101/cshperspect.a032870.

37. Currais, A., Fischer, W., Maher, P., and Schubert, D. (2017). Intraneuronal protein

aggregation as a trigger for inflammation and neurodegeneration in the aging brain.

FASEB J. *31*, 5-10. 10.1096/fj.201601184.

38. Grimm, A., and Eckert, A. (2017). Brain aging and neurodegeneration: from a

mitochondrial point of view. J. Neurochem. *143*, 418-431. 10.1111/jnc.14037.

39. Bond, S., Lopez-Lloreda, C., Gannon, P.J., Akay-Espinoza, C., and Jordan-Sciutto,

K.L. (2020). The Integrated Stress Response and Phosphorylated Eukaryotic

Initiation Factor 2alpha in Neurodegeneration. J. Neuropathol. Exp. Neurol. *79*,

123-143. 10.1093/jnen/nlz129.

40. B'Chir, W., Maurin, A.C., Carraro, V., Averous, J., Jousse, C., Muranishi, Y.,

Parry, L., Stepien, G., Fafournoux, P., and Bruhat, A. (2013). The eIF2alpha/ATF4

pathway is essential for stress-induced autophagy gene expression. Nucleic Acids

Res. *41*, 7683-7699. 10.1093/nar/gkt563.

41. Salton, G.D., Laurino, C., Mega, N.O., Delgado-Canedo, A., Setterblad, N.,

Carmagnat, M., Xavier, R.M., Cirne-Lima, E., Lenz, G., Henriques, J.A.P., and

Laurino, J.P. (2017). Deletion of eIF2beta lysine stretches creates a dominant

negative that affects the translation and proliferation in human cell line: A tool

for arresting the cell growth. Cancer Biol Ther *18*, 560-570.

10.1080/15384047.2017.1345383.

42. Lee, S., Nahm, M., Lee, M., Kwon, M., Kim, E., Zadeh, A.D., Cao, H., Kim, H.J.,

970    Lee, Z.H., Oh, S.B., et al. (2007). The F-actin-microtubule crosslinker Shot is

971    a platform for Krasavietz-mediated translational regulation of midline axon

972    repulsion. Development *134*, 1767-1777. 10.1242/dev.02842.

973 43. Wu, Z., Tantray, I., Lim, J., Chen, S., Li, Y., Davis, Z., Sitron, C., Dong, J.,

974    Gispert, S., Auburger, G., et al. (2019). MISTERMINATE Mechanistically Links

975    Mitochondrial Dysfunction with Proteostasis Failure. Mol. Cell *75*, 835-848 e838.

976    10.1016/j.molcel.2019.06.031.

977 44. Sugiura, A., Nagashima, S., Tokuyama, T., Amo, T., Matsuki, Y., Ishido, S., Kudo,

978    Y., McBride, H.M., Fukuda, T., Matsushita, N., et al. (2013). MITOL regulates

979    endoplasmic reticulum-mitochondria contacts via Mitofusin2. Mol. Cell *51*, 20-34.

980    10.1016/j.molcel.2013.04.023.

981 45. Yonashiro, R., Kimijima, Y., Shimura, T., Kawaguchi, K., Fukuda, T., Inatome, R.,

982    and Yanagi, S. (2012). Mitochondrial ubiquitin ligase MITOL blocks S-nitrosylated

983    MAP1B-light chain 1-mediated mitochondrial dysfunction and neuronal cell death.

984    Proc Natl Acad Sci U S A *109*, 2382-2387. 10.1073/pnas.1114985109.

985 46. Takeda, K., Nagashima, S., Shiiba, I., Uda, A., Tokuyama, T., Ito, N., Fukuda,

986    T., Matsushita, N., Ishido, S., Iwawaki, T., et al. (2019). MITOL prevents ER

987    stress-induced apoptosis by IRE1alpha ubiquitylation at ER-mitochondria contact

988    sites. EMBO J. *38*, e100999. 10.15252/embj.2018100999.

Shinno et. al., Figure 1

A

milton knockdown

Miro

milton

kinesin

microtubule

X ▶ to the axon

B

14-day-old

control RNAi   *milton* RNAi   *Miro* RNAi

Anti-Ubiquitin

High magnification

14-day-old

puncta of ubiquitinated proteins / a whole brain

***

p=0.452

control RNAi   *milton* RNAi   *Miro* RNAi

C

30-day-old

control RNAi   *milton* RNAi   *Miro* RNAi

Anti-Ubiquitin

High magnification

30-day-old

puncta of ubiquitinated proteins / a whole brain

N.S.   *N.S.*

control RNAi   *milton* RNAi   *Miro* RNAi

D

Presynaptic terminals (Lamina)

Brain

Cell bodies (Retina)

E

No. of presynaptic terminals

□ control RNAi

■ *milton* RNAi

***

0   1   2   3

No. of mitochondria in a synaptic terminal

F

control RNAi   *milton* RNAi

2μm   2μm

presynaptic terminal size (μm²)

3.5

2.5

1.5

0.5

***

control RNAi   milton RNAi

swollen presynaptic terminals (%)

**

control RNAi   milton RNAi

G

milton RNAi

2μm   2μm

presynaptic terminals with dense materials / **presynaptic terminals** (%)

*

control RNAi   *milton* RNAi

presynaptic terminals with dense material / **swollen presynaptic terminals** (%)

***

control RNAi   *milton* RNAi

H

control RNAi   *milton* RNAi

2μm   2μm

Shinno et. al., Figure 2

Shinno et. al., Figure 3

Shinno et. al., Figure 4

A

Autophagic defects
Protein accumulation
Locomotor dysfunction

Neurodegeneration

eclosion ——— death

7   14   21   30
(day-old)

Proteome analysis   Proteome analysis

Interaction search   Interaction search

B   7-day-old

C   21-day-old

D

	Accession	Abundance Ratio : milton RNAi / control RNAi	
		7-day-old	21-day-old
eIF2α	P41374	1.106	0.961
eIF2β	P41375	2.465***	0.753*
eIF2γ	Q24208	0.952	0.952

E

F

G

Shinno et. al., Figure 6

A

B

C

A

B

C

D / E

A

B

N.S.

N.S.

***

C

1% Triton X-100

2% SDS

*

N.S.

D

Shinno et. al., Figure 9