The unseen invaders: tracking phylogeographic dynamics and genetic diversity of cryptic

***Pomacea canaliculata and P. maculata* (Golden Apple Snails) across Taiwan**

Pritam Banerjee[1,2], Gobinda Dey[2,3], Jyoti Prakash Maity[4], Kathryn A. Stewart[5], Raju Kumar

Sharma[6], Michael W.Y. Chan[2], Kuan Hsien Lee[7, *], Chien-Yen Chen[8, 9*]

[1]Department of Environmental Science, Policy, and Management, University of California,
University Avenue and Oxford St, Berkeley, CA 94720, USA

[2]Department of Biomedical Sciences, Graduate Institute of Molecular Biology, National Chung
Cheng University, 168 University Road, Min-Hsiung, Chiayi County 62102, Taiwan.

[3]Department of Agricultural Chemistry, National Taiwan University, Taipei, 106319, Taiwan

[4]Department of Chemistry, School of Applied Sciences, KIIT Deemed to be University,
Bhubaneswar, Odisha 751024, India.

[5]Institute of Environmental Sciences, Leiden University, 2333 CC Leiden, The Netherlands

[6]Doctoral Program in Science, Technology, Environment, and Mathematics, Department of
Earth and Environmental Sciences, National Chung Cheng University, 168 University Rd.,
Chiayi County, Min-Hsiung, 62102, Taiwan

[7]Department of Emergency Medicine, Ditmanson Medical Foundation Chia-Yi Christian
Hospital

[8]Department of Earth and Environmental Sciences, National Chung Cheng University, 168
University Road, Min-Hsiung, Chiayi County 62102, Taiwan.

[9]Center for Nano Bio-Detection, Center for Innovative Research on Aging Society, AIM-HI,
National Chung Cheng University, Chiayi 62102, Taiwan

Corresponding Author

Chien-Yen Chen[8, 9*] and Kuan Hsien Lee[7, *]

*Address for correspondence: Department of Earth and Environmental Sciences, National Chung

Cheng University, 168 University Road, Min-Hsiung, Chiayi County 62102, Taiwan. and

Department of Emergency Medicine, Ditmanson Medical Foundation Chia-Yi Christian Hospital

Email: chien-yen.chen@oriel.oxon.org; yen@eq.ccu.edu.tw (C.Y. Chen); seriex.lee@gmail.com

(K.H.Lee) Tel.: +886-5-2720411 (ext. 66220); Fax: +886-5-2720807.

Abstract

The cryptic invasion of golden apple snails (*Pomacea canaliculata and P. maculata*) in Taiwan

has caused significant ecological and economical damage over last few decades, however, their

management remains difficult due to inadequate taxonomic identification, complex phylogeny

and limited population genetic information. We aim to understand the current distribution,

putative population of origin, genetic diversity and potential path of cryptic invasion of *Pomacea*

canaliculata and P. maculata across Taiwan to aid in improved mitigation approaches. The

present investigation conducted a nationwide survey with 254 samples collected from 41

locations from 14 counties or cities across Taiwan. We identified *P. canaliculata* and *P.*

maculata based on mitochondrial COI and compared their genetic diversity across Taiwan, as

well as other introduced and native countries (based on publicly available COI data) to

understand the possible paths of invasion in Taiwan. Based on mitochondrial COI barcoding,

sympatric and heterogeneous distributions of invasive *P. canaliculata* and *P. maculata* were

noted. Our haplotype analysis and mismatch distribution suggested multiple introductions of *P.*

canaliculata in Taiwan was likely originated directly from Argentina, whereas *P. maculata* was

probably introduced from a single, or a few, introduction event(s) from Argentina and Brazil.

Our population genetic data further demonstrated a higher haplotype and genetic diversity for *P.*

canaliculata and *P. maculata* in Taiwan compared to other introduced regions. Based on our

current understanding, the establishment of *P. canaliculata* and *P. maculata* is alarming and

widespread beyond geopolitical borders, requiring a concerted and expedited national and

international invasive species mitigation program.

Keywords: agricultural pest, biological invasion, DNA barcoding, golden apple snails,

mitochondrial COI, *Pomacea*, population genetics,

INTRODUCTION

Invasive alien species are major drivers of biodiversity loss and have massive impact on the

economic status of invaded countries (Linders et al., 2019; McNeely, 2001), even considered the

second most economically destructive event after natural hazards such as earthquakes or floods

(Turbelin et al., 2023). Making matters worse, morphologically similar species may invade

together, causing a masking effect to species distribution and establishment, further hindering

effective management practices (Saltonstall et al., 2002). Without in-depth knowledge about the

origins and population genetics of invasive species, management of natural landscapes and

aquascapes is rendered speculative (Sakai et al., 2001).

Pomacea canaliculata native to South America, have been both intentionally and

unintentionally introduced in several countries (Lowe et al., 2000; Zhao et al., 2022), recently

becoming a serious threat to agricultural and economic development worldwide (e.g., Asia,

Europe, Africa, North America, and the Pacific Islands) (Ranamukhaarachchi &

Wickramasinghe, 2006; Hayes et al., 2008; Buddie et al., 2021). Several other species of

Pomacea (e.g., *P. maculata*, *P. diffusa*, *P. scalaris*) have also been introduced across numerous

countries, and among the introduced species of *Pomacea*, *P. canaliculata* and *P. maculata* are

the most destructive and morphologically cryptic (Rama Rao et al., 2018) causing frequent

misidentifications. This can then lead to false biodiversity information, potentially obstructing

effective invasion management (Cowie et al., 2006). Molecular method implementation has

recently been found to be the best option for distinguishing *P. canaliculata* and *P. maculata*

(Matsukura & Wada, 2017; Banerjee et al., 2022), with population genetics analysis recently

conducted to comprehend the invasive origin and distribution of *P. canaliculata* and *P. maculata*

across Asia (Hayes et al., 2008; Yang et al., 2018; Dumidae et al., 2021; Yang et al., 2022; Zhao

et al., 2022; Liu et al., 2023)

Taiwan encompasses a high percentage of the world's biodiversity compared to its

landmass and houses an extraordinary array of fauna that comprises evolutionary lineages from

both Palearctic and Indomalaya biogeographic realms (He et al., 2018; Päckert et al, 2009).

However, as an island, Taiwan is highly sensitive to invasive species (Lee et al., 2019). Indeed,

several invasive alien species have been introduced to Taiwan which has negatively affected its

local biodiversity. Specifically, the introduction and establishment of *P. canaliculata* in Taiwan

have resulted in serious ecological and economical destruction.

Historically, Taiwan was noted as the first Asian region to import *Pomacea* spp. for

commercial utilization (dietary protein supplementation and aquarium trades) during 1979-1982

(Joshi & Sebastian, 2003; Yang et al, 2006; Cheng and Kao, 2006; Wu et al, 2010).

Subsequently, *Pomacea* spp. were introduced into other Asian countries for economic trade

(food and aquarium trades). Initially in Taiwan, *Pomacea* spp. were considered a healthy diet

alternative that contained high protein, minerals, and vitamins, thus motivating farmers in its

cultivation and trade (Cheng and Kao, 2006; Hayes et al., 2008). However, the trade of *Pomacea*

spp. was soon terminated because of the snail's poor texture and high production cost.

Exacerbating *Pomacea* spp. invasion, farmers were not content with the low market value and

thereafter released many *Pomacea* spp. into the neighboring countryside (Cheng and Kao, 2006;

Chiu et al., 2014). Almost immediately after release, *Pomacea* spp. became a serious threat to the

local agricultural system possibly as a result of its high reproduction rate, tolerance to various

environmental stressors, and voracious appetite (Joshi & Sebastian, 2003). The problem with

Pomacea was first recognized in 1982, raising national-level attention due to numerous damages

to local agronomy; *Pomacea* spp. spread into various agricultural lands, as well as natural

wetlands impacting cultivated plants such as rice seedlings, taro, and lotus, even destroying

several cultivation fields (Cheng & Kao, 2006). Making matters worse, *Pomacea* spp. acts as a

vector for various zoonotic diseases (e.g., *Angiostrongylus cantonensis*). Several cases of human

eosinophilic meningitis have been reported in Taiwan through either direct (consumption of raw

snail meat) or indirect (contaminated food) involvement of *Pomacea* spp. (Tsai et al., 2001).

More specifically, enormous annual economic losses were reported due to these invasive

gastropods (Globally: 3.94 billion USD; in Asia: 3.71 billion since 1961; (Jiang et al., 2022)),

with Asian countries as the predominantly affected region (Jiang et al., 2022). Across all

reported gastropods, *Pomacea* spp. (particularly *P. canaliculata*) were noted to be the major

contributor. Despite this, little attention has thus far been paid to understanding the invasive

biology and population genetics of *P. canaliculata* in Taiwan (Cheng & Kao, 2006; Banerjee et

al., 2022), which restricted the understanding of their regional (Taiwan) and potentially global

spread (Sakai et al., 2001).

Originally, two species of *Pomacea, P. canaliculata* and *P. scalaris,* were reported from

Taiwan, where *P. canaliculata* was noted as a destructive, dominant species and distributed

nationwide. Conversely, *P. scalaris* was reported to be less invasive and limited in distribution to

few sampling locations in Southern Taiwan (Cheng et al., 2006; Yang et al., 2008; Wu et al.,

2010). However, a more recent molecular study revealed the presence of another species, *P.*

macualta, hiding with its cryptic relative *P. canaliculata* in Southern Taiwan (Banerjee et al.,

2022). They reported that misidentification is very common between *P. canaliculata* and *P.*

macualta based on morphological observation alone (Banerjee et al., 2022), suggesting limited

knowledge on the true distribution and invasion history of *P. maculata.* In fact, out of the three

Pomacea species currently in Taiwan, *P. scalaris* is considered less destructive due to its smaller

hatching size and inferior growth performance (Wu et al., 2010), and thus assumed

geographically restricted mostly in the South of the country (Wu et al., 2011). On the other hand,

P. canaliculata and *P. maculata* are highly invasive and eco-economically destructive in nature,

yet their actual distribution remains unknown across Taiwan (Banerjee et al., 2022).

Here, our study aims to generate accurate identification of cryptic invasive *P.*

canaliculata and *P. macualta*, including their nation-wide distribution and genetic diversity to

better aid in adaptive management strategies. By doing so, we aspire to elucidate their (i)

possible origin, thereby minimizing ongoing propagation events, and (ii) proper distribution and

putative invasion path within Taiwan to maximize future eradication measures. As subsequent

introductions of *P. canaliculata* and *P. maculata* may occur from trades between Taiwan into

other Asian countries (e.g., China, Japan, Thailand, Malaysia) (Joshi & Sebastian, 2003), the

current lack of molecular data makes global invasive management efforts especially difficult.

Here too our study may expand knowledge of *Pomacea* distributions in Taiwan. To accomplish

these aims, our survey conducted collections across 41 different sampling locations within 14

counties or cities across Taiwan to reveal the origin, distribution, and population genetic

structure of *P. canaliculata* and *P. maculata*. We then compared the genetic diversity of *P.*

canaliculata and *P. maculata* with other countries (invaded or native) to understand their

possible paths of invasion.

2. METHODS

2.1 Snail sample collection and preservation

In Taiwan, *Pomacea* spp. were presumably introduced through human activities and later

distributed nationwide. Since this distribution may be affected by agricultural land or other

(human-induced) distribution barriers (e.g., roads, mountains), sampling locations (populations)

were centered around or near cities to understand the current nationwide distribution. We then

further subdivided Taiwan into different regional parts coordinates (e.g., Southern, Northern,

Eastern, Western and Middle) to more generally understand the current distribution of *Pomacea*

for downstream management decisions. To accomplish this, distribution of *Pomacea* spp. were

surveyed across 41 locations across 14 counties or cities in Taiwan, including Hualien (HU) and

Taitung (TT) from the East, Yunlin (YL), Changhua (CH), Taichung (TC), and Miaoli (MIO)

from the Middle of island, Hsinchu (HS), Taoyuan (TY), Taipei (TP), and Yilan (YI) from the

North, and Pingtung (PT), Kaohsiung (KSH), Tainan (TN), and Chiayi (CY) from the South.

This was achieved via manual hand-collection from crop fields and drainage ditches from April

2020 to January 2023. Sampling locations and the number of samples are shown in Figure 1 and

APPENDIX 1. Collected samples were transported directly to the laboratory as soon as possible

(as per sampling protocols). In cases of remote sampling or locations away from the laboratory,

specimens were heat-shocked using hot water (100^{o} C) for 10-15 minutes and subsequently kept

in an ice box in separate sampling bags to avoid contamination. Once in the laboratory, samples

were washed thoroughly to remove soil and debris, and then if not previously applied, were heat-

shocked using a microwave to remove inner tissue from shells. After removing the inner tissue

from the shell, 20-50 mg of foot tissue was used to extract DNA to avoid parasite contamination,

and the rest of the tissue was stored in 95% ethanol.

2.2 Genomic DNA extraction and amplification

Genomic DNA was extracted from the 20-50 mg foot tissues of *Pomacea* spp. with the help of a

DNeasy Blood and Tissue Kits (Qiagen, Germany), similar to Banerjee et al. (2022). The

mitochondrial DNA barcode of *Pomacea* spp. was amplified using a conventional PCR

(CLUBIO PCR system, 96-well, CB series; Scientific Biotech Corp., Taiwan) with species-

specific primers: PcanCOI (F) (5'-TGG GGT ATG ATC AGG CC-3') and PinsCOI (F) (5'-ATC

TGC TGC TGT TGA AAG-3') and HCO2198 (R) (5'-TAA ACT TCA GGG TGA CCA AAA

AAT CA-3') for initial identification of *P. canaliculata* and *P. maculata* (Matsukura et al, 2008).

The double distilled water and DNA from Giant African land snail (*Lissachatina fulica*) were

used as the negative control followed by the previous study (Banerjee et al., 2022). Furthermore,

successfully identified specimens from initial screening were amplified using universal

mitochondrial COI primers: LCO1490 (5' GGT CAA CAA ATC ATA AAG ATA TTG G-3')

and HCO2198 (5' TAA ACT TCA GGG TGA CAA AAA AAT CA-3') for a 658bp fragment.

The final PCR reaction contained 5 µL 5x Fast-RunTM Taq Master Mix with Dye (Protech

Technology Enterprise, Taiwan), 0.5 µL of 10 pmol forward and reverse primers (Genomics,

Taiwan), 1 µL DNA template from the tissue sample, for a final volume of 25 µL using ultrapure

water (Thermo Fisher, USA). The PCR conditions were 5 min at 94°C, 35 cycles of 30s at 94°C

(denaturation), 30s at 55°C (annealing), and 1 min at 72°C (extension), then the final extension

was at 72°C for 5 min. All amplified products were sent to Genomics Co., Ltd, Taiwan for

sequencing in both directions. All sequences derived from the present study were aligned using

the MUSCLE algorithm (Edgar, 2004), and edited in Mega v11 (Tamura et al, 2021), and

Geneious v11.1.5 (Kearse et al., 2012). A total of 254 of *Pomacea* spp. were amplified from 41

sampling spots in Taiwan.

2.3 Mitochondrial COI

The identification of 254 *Pomacea* spp. were accomplished via sequence similarity based on the

NCBI database (https://blast.ncbi.nlm.nih.gov/Blast.cgi) with the highest level of query cover

and percentage identity. To clarify the origins of our *Pomacea* spp. samples across Taiwan,

previously identified and uploaded sequences of *P. canaliculata* and *P. maculata* were

downloaded from NCBI which represent different geographic locations (Hayes et al., 2008;

Kannan et al., 2021; Liu et al, 2019; Lv et al., 2013; Yang et al., 2018). Sequence quality was

examined in Geneious v11.1.5 (Kearse et al., 2012), BioEdit v7.2 (Hall, 1999), and DnaSP

v6.12.03 (Rozas et al., 2017) and filtering sequences published after 2007 to clearly distinguish

P. maculata from *P. canaliculata* (Yang et al., 2018). A total of 1173 (948 *P. canaliculata* and

225 *P. maculata*) sequences were selected for final evaluation (APPENDIX 2, 3)*,* and further

screening was performed after a test run of phylogenetic and haplotyping network analysis

(details in sections 2.5). Sequence length variation was adjusted and trimmed manually in Mega

v11 (Tamura et al., 2021), allowing us to prepare three datasets, (i) the sequences generated from

the present study (samples only from Taiwan) represented 611bp for *P. maculata* and *P.*

canaliculata as Dataset 1, and global sequences (Taiwan and other previously published

sequences collected from NCBI and/or Hayes et al., 2008; Kannan et al., 2021; Liu et al, 2019;

Lv et al., 2013; Yang et al., 2018) datasets represent (ii) 579bp for *P. maculata* as Dataset 2, and

(iii) 569bp for *P. canaliculata* as Dataset 3.

2.4 DNA polymorphism and phylogeographic distribution

All sequences were aligned in Mega v11 (Tamura et al., 2021) using the MUSCLE algorithm

(Edgar, 2004), and DnaSP v6.12.03 (Rozas et al., 2017) was used to calculate the number of

haplotypes (*H*) and polymorphic sites (*S*) as well as haplotype diversity (*Hd*), nucleotide

diversities (π) and average number of difference (*K*). Neutrality tests such as Tajima's *D* and Fu's

Fs (Fu, 1997) were performed to understand signatures of population expansion using Arlequin

v3.5.2.2 (Excoffier & Lischer, 2010). Furthermore, to understand the level of genetic difference

between the populations, pairwise F_{st} values were calculated using Arlequin v3.5.2.2 (Excoffier

& Lischer, 2010), and the level of gene flow/effective migration rate was calculated using the

following formula: $Nm=((1/F_{ST})$ -1)/2) (Hudson et al., 1992). The connections between

haplotypes and their phylogeographic distribution were analyzed using the median-joining

network implemented in PopArt v1.7 (Leigh & Bryant, 2015). Furthermore, the mismatch

distribution analyses were performed in DnaSP v6.12.03 (Rozas et al., 2017) to understand the

genetic variation for the population of *P. canaliculata* and *P. maculata*. The inter- and intra-

specific genetic distance (*p*-distance) of *P. canaliculata* and *P. maculata* were calculated using

Mega v11 (Tamura et al., 2021).

2.5 Phylogenetic analysis

The evolutionary model was first tested in jModelTest v2.1.10 for each population (Darriba et al., 2012). The maximum-likelihood (ML) method with the GTR+G model was used based on 1000 bootstrap replicates (Felsenstein, 1985) to build phylogenetic relations in Mega v11 (Tamura et al., 2021). A phylogenetic tree using Bayesian inference (BI) was produced using BEAUti v1.10.4 and BEAST v1.10.4 (Bouckaert et al., 2014) under GTR+G substitution model, strict clock, constant tree prior with random starting tree model. A Markov Chain Monte Carlo Analysis (MCMC) was run for 10 million generations and sampled in every 1000 generation as well as the final consensus tree was generated in TreeAnnotator v1.10.4 with 10% burn-in. The convergence states were visualized using Tracer v1.7.2 (Rambaut et al., 2018) with all the ESS values above 220. Both the maximum-likelihood (ML) and Bayesian inference (BI) analysis were run twice independently. The resulting consensus trees were visualized in FigTree v1.4.4 (Rambaut & Drummond, 2010). In all phylogenetic trees (ML and BI), *P. occulta* (FJ710311), *P. lineata* (FJ710311), *P. paludosa* (EF514960), *P. diffusa* (EU528547), *P. scalaris* (collected from TC by our team), *Marisa cornuarietis* (MK919685), *Asolene platae* (EU528547), and *Saulea vitrea* (FJ710301) were added as outgroups.

3. RESULTS

3.1 Species distribution

We used a quick identification primer (PanCOI, PinsCOI and HCO2198) and full former region (LCO1490 and HCO2198), and both identified *Pomacea* spp up to species level. Based on our collections across Taiwan, the distribution of *P. canaliculata* and *P. maculata* was observed to be sympatric, and heterogeneous. *P. canaliculata* was distributed throughout Taiwan and represented the dominant species ($\approx$ 97%), whereas *P. maculata* represented $\approx$ 3% of the total number of collected individuals (Fig. 1; Map created using "ArcGIS v10.7 software; ESRI Inc., Redlands, CA, USA"). The genetic distance (p-distance) within *P. canaliculata* and *P. maculata* were 0.02 and 0.01 respectively, and genetic distance (p-distance) in between them were 0.0928 (APPENDIX 4)

3.2 Phylogenic analysis

Phylogenetic relationships were congruent among four independent analysis (both ML and BI) with *P. canaliculata* and *P. maculata* clustered into two clades separately (Fig. 2). *P. maculata* grouped with *Pomacea occulta* and *Pomacea lineata* together and formed a sister group of *P. canaliculata.* although this separation was not strongly supported with Maximum likelihood analysis (<80, Fig. 2). Furthermore, there were three clades noted for *P. canaliculata* (Clade A, B, and C) and two clades for *P. maculata* (Clade D and E).

3.3 Haplotype distribution and network analysis

3.3.1 Phylogeography of *P. canaliculata* in Taiwan

A total number of 246 sequences of *P. canaliculata* from 14 different cities/counties revealed the

presence of 18 haplotypes (Pc_Hap1-18) and three different networks (Network Tw-A-C) across

Taiwan (Fig. 3, Table 1, 2, APPENDIX 5). The first network (Network Tw-A) was most

dominant, representing 69.90% of all sequences with seven haplotypes. Among the seven

haplotypes, Pc_Hap 3 was widespread and dominant; the other six haplotypes included five

unique haplotypes from Tainan (Pc_Hap 1, 6), Taitung (Pc_Hap 16), Chiayi (Pc_Hap 12),

Changhua (Pc_Hap 15), and one shared by Chiayi, Yilan, Changhua, and Taichung (Pc_Hap 11).

The Pc_Hap 3 demonstrated the highest number of connections which created a star-like

appearance, could be a potential founder haplotype of this network. The Network Tw-B

represented 30.49% of all sequences with four shared haplotypes including Pc_Hap 2

(widespread), Pc_Hap 4 (shared by Tainan, Kaohsiung, and Taichung), Pc_Hap 8 (shared by

Pingtung, Yunlin, Hualien), Pc_Hap 5 (shared by Tainan, Pingtung, Chiayi, Taipei, Taichung),

and five unique haplotypes from Hualien (Pc_Hap 17, 18), Yilan (Pc_Hap 13, 14) and Chiayi

(Pc_Hap 10). Here, Pc_Hap 2 assumed to be the founder haplotype (with most number of

connections and a star-like appearance around), The Network Tw-C was represented by only

1.62% of all sequences with one shared haplotype Pc_Hap 9 (Kaohsiung, Hsinchu, Changhua),

and one unique haplotype Pc_Hap 7 (Tainan). The distribution Network Tw-C was limited and

found from the South (Kaohsiung, Tainan), Middle (Changhua), and North part (Hsinchu).

Compared to other cities/counties Tainan, Chiayi, Yilan, and Hualien were reported to have a

higher number of shared and unique haplotypes. (Table 1, 2).

3.3.2 Phylogeography of *P. canaliculata* worldwide

The combined mitochondrial COI datasets of *P. canaliculata* with 948 sequences across Africa,

Asia, North America, Oceania, and South America were compared to one another and revealed

the presence of a total of 65 haplotypes (Wc_Hap1-65) and three distinct networks (Network

Wc-A-C) (Fig. 4, APPENDIX 6). The Network Wc-A, representing 32 haplotypes, including the

most widely shared dominant Wc_Hap 1, and 31 unique haplotypes from Argentina (Wc_Hap 3-

8, 10, 11, 12, 15, 17), Taiwan (Wc_Hap 24, 26, 30, 31, 34), Philippines (Wc_Hap 52, 53, 55, 57,

59, 61), Indonesia (Wc_Hap 48, 49), Thailand (Wc_Hap 62, 63), USA (Wc_Hap 64), China

(Wc_Hap 35), Myanmar (Wc_Hap 51), Chile (Wc_Hap 47), Malaysia (Wc_Hap 50). All of

those unique haplotypes were connected with Wc_Hap 1, thus inferred as the founder haplotype.

This haplotype was also found to be a widespread and dominant haplotype in Taiwan, as well as

shared by other Asian countries.

Network Wc-B represented 22 haplotypes, including five shared; Wc_Hap 16 (Taiwan,

China, Japan, Malaysia, and Kenya), Wc_Hap 20 (China, Taiwan, and Philippines), Wc_Hap 21

(China, Philippines), Wc_Hap 23 (Taiwan, China, Japan, Philippines), and Wc_Hap 36 (China

and Japan) and 17 unique haplotypes from Taiwan (Wc_Hap 25, 27, 29, 32, 33), China (Wc_Hap

22, 43-46), Uruguay (Wc_Hap 40, 41), Philippines (Wc_Hap 54, 56, 58, 60), and Argentina

(Wc_Hap 14). All the shared haplotypes in Network Wc-B were reported from Taiwan, except

Wc_Hap 21, and Wc_Hap 36, which were one to three mutational steps away from the shared

one. The Wc_Hap 27 (from Taiwan) was noted to be the common connection among all the

haplotypes in Network Wc-B.

Lastly, Network Wc-C represented 11 haplotypes, including 2 shared: Wc_Hap 28

(Taiwan, Japan, and Singapore), Wc_Hap 9 (Taiwan and Argentina), and 9 unique haplotypes

from Argentina (Wc_Hap 2, 13, 17, 18, 65), and Uruguay (Wc_Hap 37-39, 42). Among the

haplotypes in Network Wc-C, Wc_Hap 9 may be inferred as the common ancestor. Most of the

haplotypes from introduced regions (Asian countries and African countries) were found to be

associated with Network Wc-A and B, however, few representatives were noted in Network Wc-

C.

The comparison of representative sequence in Dataset 1 (representing only Taiwan

sequences) and Dataset 2 (sequence from Taiwan as well as other native and non-native

countries), revealed that haplotypes of Network Tw-A, Network Tw-B, and Network Tw-C,

represent in Network Wc-A, Network Wc-B, and Network Wc-C, respectively without any

interchange (APPENDIX 6). Taiwan was represented with 18 haplotypes in Dataset 1 (619 bp),

however in the global dataset (Dataset 3 = 569 bp; 16 haplotypes from Taiwan) a slight decrease

in the number of haplotypes was noted (APPENDIX 7). Across all countries with sequences, the

total number of unique haplotypes were found to be the most prevalent in Argentina, Uruguay,

Philippines, China, and Taiwan.

3.3.3 Phylogeography of *P. maculata* in Taiwan

In Taiwan, eight sequences of *P. maculata* represented four different haplotypes, where Pm_Hap

19 (62.50%) was the most common and the other three haplotypes (Pm_Hap 20-22) were

represented by single sequences (Table 3; APPENDIX 5). Haplotype network analysis of *P.*

maculata showed the presence of a single network, where Pm_Hap 1 was a few mutational steps

away; however, Pm_Hap 20-22 were more closely related (Fig. 3).

3.3.4 Phylogeography of *P. maculata* worldwide

A total of 226 sequences of worldwide *P. maculata* revealed 46 haplotypes and seven different

networks (Network Wm-A-G). Among the haplotype networks, Network Wm-A was

widespread, consisting of 29 haplotypes which include four shared haplotypes; Wm_Hap 6 was

the most widely distributed (representing 37.61% of all sequences; APPENDIX 8), and

Wm_Hap 10 (9.73%) was shared by Argentina and USA, Wm_Hap 9 was shared by USA,

Japan, and Argentina, and Wm_Hap 1 was shared by Taiwan, Argentina, and Uruguay, including

unique haplotypes from Argentina (Wm_Hap 13-18), Uruguay (Wm_Hap 2-4), Taiwan

(Wm_Hap 7, 8), Vietnam (Wm_Hap 5), USA (Wm_Hap 11), and 11 unique to Brazil (Fig. 5).

All the other Networks were unique to Brazil. Among the shared haplotypes in Network Wm-A,

all other Asian-originated haplotypes created a star-like structure around Wm_Hap 6 and

Wm_Hap 1 which suggests those two haplotypes may have been founders throughout Asia.

Wm_Hap 6 and Wm_Hap 1 were reported from Taiwan.

3.4 Population genetic structure *P. canaliculata* and *P. maculata*

The high value of haplotype diversity (Hd), nucleotide diversity (π), and an average number of

nucleotide differences (K) suggested higher genetic diversity of *P. canaliculata* in Taiwan (Table

1). Among all 14 sites, Taitung, Taichung, Pingtung, Yilan, Kaohsiung, and Chiayi showed the

highest genetic diversity; Tainan, Yunlin, Miaoli, and Taoyuan showed high to moderate genetic

diversity; Changhua, Hsinchu, Hualien, and Taipei were noted to have less diverse populations

(Table 1). The significantly positive value of Fu's Fs and Tajima's D ($D < 0$) were recorded for

populations Taitung, Miaoli, Yunlin, Taoyuan, Yilan, Kaohsiung, and Pingtung. Conversely,

significantly negative values of Fu's Fs and Tajima's D ($D > 0$) were observed for the population

in Changhua, Taipei, and Hsinchu (Table 1). Furthermore, *P. canaliculata* in Taiwan was noted

to significantly differ among some provinces, where a higher F_{st} value ($F_{st} > 0.2$) resulted in less

gene flow ($Nm < 1$) and vice versa (Table 4). However, the negative F_{st} values likely indicate

more variation within the focal populations than between populations. While *P. maculata* was

found to be less dominant in distribution compared to *P. canaliculata*, polymorphism in

populations were higher (Table 4).

The genetic diversity of *P. canaliculata* and *P. maculata* were considerably higher in

their native counties (Argentina, Brazil, and Uruguay) compared to the places in which they were

introduced. As an invaded region Taiwan demonstrated a higher genetic diversity of *P.*

canaliculata and *P. maculata*, constituting of 16 haplotypes for *P. canaliculata*, and 4 for *P.*

maculata (Table 5 and 6).

3.5 Mismatch distribution

The mismatch distribution results of *P. canaliculata* in Taiwan showed two bimodal well-

separated peaks with some small intermediate peaks (Fig. 6a). However, in the case of *P.*

maculata three distinct peaks were observed, including two major peaks and an intermediate

peak (Fig. 6b). The mismatch distribution of *P. canaliculata* and *P. maculata* from the world

dataset showed multimodal distribution (Fig. 6c and d).

4. **DISCUSSION**

Based on our mitochondrial COI barcoding data, we found a wide distribution of *P. canaliculata*

across Taiwan, whereas *P. maculata* was less abundant and restricted few counties (Pingtung,

Kaohsiung, Tainan, and Chiayi, Miaoli and Yilan) (Fig. 1). Moreover, the distribution of *P.*

maculata and *P. canaliculata* were noted to be sympatric and heterogenous across the country

(Fig. 1 and Table 1), a similar distribution pattern was noted in our previous study in southern

Taiwan (Banerjee et al., 2022). Our haplotype analysis further demonstrated the possible origin

of *P. maculata* from Argentina, although *P. maculata* may have been introduced independently

from Argentina and Brazil (Hayes et al., 2008; Yang et al., 2022) instead of being imported

directly from Brazil as reported in cases of *P. maculata* populations in China (Yang et al., 2018).

Thus, the results from Yang et al. (2018, 2022) and this present study showed that the origin of

P. maculata may be different in mainland China from Hong Kong and Taiwan.

Interestingly, the invasion of *P. canaliculata* and *P. maculata* has also been noted to be

sympatric and heterogenous in other nearby Asian countries, such as in China (Yang et al.,

2018), Hong Kong (Yang et al., 2022), Malaysia (Rama Rao et al., 2018), and Thailand

(Dumidae et al., 2021), where *P. canaliculata* is also reported to be more widely distributed

compared to *P. maculata*. This corroborates the dominant hypothesis of predominately *P.*

canaliculata in Asia regions (Hayes et al., 2008). Interestingly, this distribution pattern is not

reflected in their native ranges, where *P. maculata* is routinely found to be widely distributed

throughout Brazil, Uruguay, and Argentina, whereas *P. canaliculata* remains restricted to North

Argentina and South Uruguay (Hayes et al., 2008; Glasheen et al., 2020). The difference in the

distribution in Asian regions may reflect different physiological tolerance, or species-specific

biology (Hayes et al., 2008; Matsukura et al, 2016). Alternatively, it can be posited that these

phylogeographical patterns and unequal distribution may be due to complex introduction

histories. Notably, current distribution data is based on mitochondrial DNA, and *P. canaliculata*

and *P. maculata* are reported to hybridized both in their native and non-native ranges (Glasheen

et al., 2020; Yang et al., 2020; Kannan et al., 2021). Thus, hybridization could also play an

important role in their unequal phylogeographic distribution. Similar to *P. maculata*, another

invasive species, *P. scalaris,* was previously reported to be restricted to the southern part of

Taiwan (Wu et al., 2010), however, we recorded some individuals from the middle region of

Taiwan during our collection , possibly suggesting slow expansion.

Although the origin of *P. canaliculata* and *P. maculata* in Asia has been widely studied

by previous researchers from different countries (Hayes et al., 2008; Liu et al., 2019; Dumidae et

al., 2021; Yang et al., 2022), additional data from Taiwan specifically (and for the first time) will

contribute to our knowledge regarding the possible path of introduction and spread. Our

haplotype analysis revealed the presence of all three independent networks of *P. canaliculata*

across Taiwan, whereas only two networks of *P. canaliculata* were detected from other countries

of Asia (Yang et al., 2018; Yang et al., 2021). The occurrence of three largely different networks

across Taiwan suggesting introduction may have occurred multiple times and possibly

propagated from multiple locations (Fig. 3).

The combined mitochondrial datasets of *P. canaliculata* with 948 global sequences

(originating from Africa, Asia, North America, Oceania, and South America) exhibited three

different networks, mirroring previous studies (Yang et al., 2018; Yang et al., 2021), with

sequences from Taiwan represented across all networks (Fig. 4). The Wc_Hap 1, Wc_Hap 27,

and Wc_Hap 9 represented the founder haplotypes of Network Wc-A, Network Wc-B, and

Network Wc-C respectively. The common sharing of founder haplotypes between Taiwan and

Argentina reaffirms the historical assumption of the introduction of *P. canaliculata* from

Argentina to Taiwan (Joshi & Sebastian, 2003; Hayes et al., 2008). Although the absence of a

shared haplotype from Argentina was noted in the case of WC_Hap 27 in Network Wc-B, which

again may simply be due to a smaller number of represented sequences from Argentina in our

global dataset. Furthermore, Network Wc-A and Network Wc-B were widely distributed,

however the haplotypes from Network Wc-C were rare in all native and non-native countries, not

just those haplotypes representative in Asia. This also does not lend support for sequence

sampling bias (i.e. more representative samples in Taiwan compared to other sampled countries),

although we cannot preclude it as a possible contributing factor.

Interestingly, our haplotype analysis revealed the presence of a single Network for *P.*

maculata across Taiwan, echoing patterns across other Asian countries (Yang et al., 2018; Yang

et al., 2021). This is in contrast to *P. canaliculata* for which the haplotype distribution pattern

suggests multiple introduction events (Fig. 3 and 5). Corroborating this further and despite the

complex network of unique haplotypes, Taiwan shares the founder haplotype with Argentina

(Wm_Hap 1). Notably, our results demonstrated the presence of Brazilian haplotypes with only a

few mutational steps away from the founder haplotype (Wm_Hap 1), suggesting the introduction

of *P. maculata* possibly originated via Argentina and Brazil, rather than solely from Brazil (Fig.

5). Similar findings were also noted in Hong Kong, where results from Yang et al. (2022)

demonstrated *P. maculata* was possibly introduced from Argentina and Brazil. The presence of

only a single network of *P. maculata* may suggest less frequent introductions events in Taiwan.

Unsurprisingly, the populations of *P. canaliculata* and *P. maculata* in Taiwan showed

lower haplotype diversity compared to their native population counterparts (e.g., Argentina and

Brazil) which can be the result of founder effects. However, compared with other Asian

countries, Taiwan showed similar (e.g., China) or higher (e.g., Malaysia, Thailand, and

Philippines) genetic diversity for *P. canaliculata* and *P. maculata* (Table 5 and 6). These

observations insinuate that Taiwan may have faced multiple introductions during the past.

Furthermore, as per our mismatch distribution analysis, two distinct peaks with large pairwise

differences created a well-separated bimodal mismatch graph for *P. canaliculata* (Fig. 6a),

however mismatch distribution analysis produced an intermediate peak in between the bimodal

graph for *P. maculata* (Fig. 6b). Those patterns suggest two episodes of population expansion

which may coincide with multiple introductions for *P. canaliculata*, and possibly a single or

continuous source for *P. maculata*. Furthermore, the multimodal mismatch distribution of *P.*

canaliculata and *P. maculata* for the global dataset (Dataset 2 and 3) indicated their complex

population expansion dynamics, and possibly introduction history, around several countries (Fig.

6c and d).

Their genetic diversity (value of *Hd*, π, and *K*) varied among the counties/cities of

Taiwan which may be related to multiple sources of introductions at different times, across

different land-use patterns (e.g., agriculture), and human transportation. Moreover, the dispersal

ability of *Pomacea* spp. is observed to be low and thus the widespread distribution across Taiwan

within such a short invasion period was probably facilitated predominantly from human

transportation.

The neutrality test, such as the value of Fu's *Fs* and Tajima's *D* (D < 0), was not

observed as significant for the overall population *P. canaliculata* and *P. maculata* in Taiwan,

however, the higher value of Fu's *Fs* and Tajima's *D* (*D* < 0) for the population of *P.*

canaliculata in Taitung, Miaoli, Yunlin, Taoyuan, Yilan, Kaohsiung, and Pingtung did

demonstrate a high number of rare alleles. On the contrary, the lower value of Fu's *Fs* and

Tajima's *D* (*D* > 0) for the population of *P. canaliculata* in Changhua, Taipei, and Hsinchu

suggests a scarcity of rare alleles, however, these results may equally be due to sampling error.

Furthermore, the significant difference in F_{st} value (F_{st} > 0.2) and negative F_{st} values (suggest

more variation within the populations than in between) in Table 4 could be result of multiple

introductions, human transportation and population mixing.

Although the mitochondrial DNA haplotyping data is useful to understand the genetic

diversity, as *P. canaliculata* and *P. maculata* are reported to hybridize, implementing nuclear

genomic data or whole genomic sequencing data across invaded and native countries will be

more useful for understanding invasion dynamics. The current datasets were limited in number

and range of collections across different countries or regions, which could be solved by adding

more precisely identified sample sequences.

Conclusions

Our data suggested the Argentinian origin of *P. canaliculata* in Taiwan, whereas *P. maculata*

might have both Argentinian and Brazilian origin. Comparing their introduction histories, current

distribution, and cryptic nature of *P. canaliculata* and *P. maculata*, we assume that *P. maculata*

may have been introduced along with *P. canaliculata* (either intentionally or unintentionally) in

Taiwan, however, *P. canaliculata* likely has several introduction histories than *P. maculata*.

Furthermore, our sampling across Taiwan, also reported the presence of *P. scalaris* (non-target

for this study) within the middle region of Taiwan, which possibly suggest the expansion of

another invasive species. Thus, the establishment of *Pomacea* spp. across all of Taiwan is

alarming, and our current data suggests the urgent need for a nationwide mitigation program and

proper management policy. For early detection and to prevent further introductions and spread of

multiple *Pomacea* species, eDNA-based methods have been previously demonstrated to be

useful (Banerjee et al., 2022).

ACKNOWLEDGEMENTS

P.B. has been supported by Overseas Research Scholarships (ORS) from National Chung Cheng

University as well as Ministry of Education (MOE)- Industry-Academia project (Taiwan). The

authors would also like to thank the Ministry of Science and Technology (Taiwan) for financial

support (MOST 109-2811-M-194-502; MOST 108-2811-M-194-510).

AUTHORS' CONTRIBUTIONS

P.B. and C.Y.C. conceived of the idea; P.B., performed experiments, P.B., K.A.S., and C.Y.C.,

prepared the first draft and revised the manuscript, figures and tables. G.D., R.K.S, J.P.M., and

M.W.Y.C, K. H. L., gave extensive edits and revised the manuscript. We would like to thank

Prof. Qian-Qian Yang (College of Life Sciences, China Jiliang University, Hangzhou 310018,

China) for her valuable suggestions during revisions.

CONFLICTS OF INTERESTS

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

All data obtained in this study have been deposited to GeneBank under the accession number

OQ977961 - OQ978203 and OQ977020-OQ977027.

Captions of Figures

Figure 1. Geographical distribution and frequency haplotypes of *P. canaliculata* (Pc_Hap 1-18) and *P. maculata* (Pm_Hap 19-22) in Taiwan (Map created using ArcGIS v10.7 software)

Figure 2. Phylogenetic tree generated from 254 sequences mitochondrial COI sequence of *P. canaliculata* and *P. maculata* collected from 41 different location (14 counties/cities) in Taiwan. Node labels are maximum likelihood/Bayesian posterior probabilities. The sequences of *P. lineata*, *P. paludosa, P. diffusa, P. scalaris*, *Marisa cornuarietis*, *Asolene platae*, *Saulea vitrea* were used as outgroups.

Figure 3. Mitochondrial COI haplotypes networks revealing frequency and relationship of *P. canaliculata* (Pc_Hap 1-18) and *P. maculata* (Pm _Hap 19-22) across Taiwan using Dataset 1.

Figure 4. Mitochondrial COI haplotypes (Wc_Hap) revealing frequency and relationship of *P. canaliculata* from the global dataset (Dataset 3).

Figure 5. Mitochondrial COI haplotypes (Wm_Hap) revealing frequency and relationship of *P. maculata* from the global dataset (Dataset 2).

Figure 6. Mismatch distribution analysis of *P. canaliculata* from (a) Taiwan (Dataset 1) and (c) the global dataset (Dataset 3), and *P. maculata* (b) Taiwan (Dataset 1) and (d) the global dataset (Dataset 2)

Captions of Tables

Table Caption

Table 1. Population genetic diversity of *P. canaliculata* and *P. maculata* across Taiwan. The

significant value of Tajima's *D* and Fu's *Fs* are marked in bold.

Table 2. Haplotype distribution (Pc_Hap) of *Pomacea canaliculata* in 14 counties/cities across

Taiwan.

Table 3. Haplotype distribution (Pm_Hap) of *Pomacea maculata* in 14 counties/cities across

Taiwan.

Table 4. Genetic differentiation coefficient (F_{st}) below diagonal and gene flow (N_m) above

diagonal based on mitochondrial COI sequences of *P. canaliculata* from 14 counties or cities of

Taiwan. Abbreviations: HU: Hualien, and TT: Taitung from the East, YL: Yunlin, CH:

Changhua, TC: Taichung, and MIO: Miaoli from the Middle, HS: Hsinchu, TY: Taoyuan, TP:

Taipei, and YI: Yilan from the North, and PT: Pingtung, KSH: Kaohsiung, TN: Tainan, and CY:

Chiayi from the South. The higher F_{st} value (F_{st} >0.2) resulted in less gene flow (Nm <1)

between the counties or cities (marked in bold).

Table 5. Population genetic diversity of *Pomacea canaliculata* from the global mitochondrial

COI dataset (Dataset 3).

Table 6. Population genetic diversity of *Pomacea maculata* from the global mitochondrial COI

dataset (Dataset 2).

Reference

ArcGIS 10.7 software; ESRI Inc., Redlands, CA, USA. Retrieved from https://www.esri.com/en-

us/arcgis/products/arcgis-desktop/overview

Banerjee, P., Stewart, K. A., Dey, G., Sharma, R. K., Maity, J. P., Chan, M. W., . . . Chen, C.-Y.

(2022). When conventional methods fall short: identification of invasive cryptic Golden

Apple Snails (Pomacea canaliculata; P. maculata) using environmental DNA.

Hydrobiologia, 849(19), 4241-4257.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., . . . Drummond, A. J.

(2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS*

computational biology, 10(4), e1003537.

Buddie, A. G., Rwomushana, I., Offord, L. C., Kibet, S., Makale, F., Djeddour, D., . . . Chacha,

D. (2021). First report of the invasive snail Pomacea canaliculata in Kenya. *CABI*

Agriculture and Bioscience, 2(1), 1-10.

Cheng, E. Y., & Kao, C. (2006). Control of golden apple snail, Pomacea canaliculata (Lamarck),

in Taiwan. *Global advances in ecology and management of golden apple snails*, 155-167.

Chiu, Y.-W., Wu, J.-P., Hsieh, T.-C., Liang, S.-H., Chen, C.-M., & Huang, D.-J. (2014).

Alterations of biochemical indicators in hepatopancreas of the golden apple snail,

Pomacea canaliculata, from paddy fields in Taiwan. *Journal of environmental biology,*

35(4), 667.

Cowie, R., Hayes, K., & Thiengo, S. (2006). What Are Apple Snails? Confused Taxonomy and

Some Preliminary Resolution. Dalam Global Advances in Ecology and Management of

Golden Apple Snails. RC Joshi and LS Sebastian (Eds), 3-23. *PhilRice. Philippines*.

Darriba, D., Taboada, G.L., Doallo, R. and Posada, D., 2012. jModelTest 2: more models, new

heuristics and parallel computing. Nat Meth 9: 772. Go to original source.

Dumidae, A., Janthu, P., Subkrasae, C., Polseela, R., Mangkit, B., Thanwisai, A., & Vitta, A.

(2021). Population genetics analysis of a Pomacea snail (Gastropoda: Ampullariidae) in

Thailand and its low infection by Angiostrongylus cantonensis. *Zoological Studies, 60*.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic acids research, 32*(5), 1792-1797.

Excoffier, L., & Lischer, H. E. (2010). Arlequin suite ver 3.5: a new series of programs to

perform population genetics analyses under Linux and Windows. *Molecular ecology*

resources, 10(3), 564-567.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap.

evolution, 39(4), 783-791.

Fu, Y.-X. (1997). Statistical tests of neutrality of mutations against population growth,

hitchhiking and background selection. *Genetics, 147*(2), 915-925.

Glasheen, P. M., Burks, R. L., Campos, S. R., & Hayes, K. A. (2020). First evidence of

introgressive hybridization of apple snails (Pomacea spp.) in their native range. *Journal*

of Molluscan Studies, 86(2), 96-103.

Hall, T. A. (1999). *BioEdit: a user-friendly biological sequence alignment editor and analysis*

program for Windows 95/98/NT. Paper presented at the Nucleic acids symposium series.

Hayes, K. A., Joshi, R., Thiengo, S. C., & Cowie, R. (2008). Out of South America: multiple

origins of non□native apple snails in Asia. *Diversity and distributions, 14*(4), 701-712.

He, J., Gao, Z., Su, Y., Lin, S., & Jiang, H. (2018). Geographical and temporal origins of

terrestrial vertebrates endemic to Taiwan. *Journal of Biogeography, 45*(11), 2458-2470.

Hudson, R.R., Slatkin, M. and Maddison, W.P., 1992. Estimation of levels of gene flow from

DNA sequence data. Genetics, 132(2), pp.583-589.

Jiang, X., Zheng, P., Soto, I., Haubrock, P. J., Chen, J., & Ji, L. (2022). Global economic costs

and knowledge gaps of invasive gastropods. *Ecological Indicators, 145*, 109614.

Joshi, R. C., & Sebastian, L. S. (2003). Global advances in ecology and management of golden

apple snails.

Kannan, A., Ratnayeke, S. and Yow, Y.Y., 2021. Molecular evidence of hybridisation in two

invasive species of Pomacea (Gastropoda: Ampullariidae) in Peninsular Malaysia.

RAFFLES BULLETIN OF ZOOLOGY, 69, pp.570-585.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., . . . Duran, C.

(2012). Geneious Basic: an integrated and extendable desktop software platform for the

organization and analysis of sequence data. *Bioinformatics, 28*(12), 1647-1649.

Lee, K.-H., Chen, T.-H., Shang, G., Clulow, S., Yang, Y.-J., & Lin, S.-M. (2019). A check list

and population trends of invasive amphibians and reptiles in Taiwan. *ZooKeys, 829*, 85.

Leigh, J. W., & Bryant, D. (2015). POPART: full-feature software for haplotype network

construction. *Methods in ecology and evolution, 6*(9), 1110-1116.

Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J.C., Guirao-Rico, S., Librado, P., Ramos-

Onsins, S.E., Sánchez-Gracia, A. (2017). DnaSP 6: DNA Sequence Polymorphism

Analysis of Large Datasets. Mol. Biol. Evol. 34: 3299-3302.

Linders, T. E. W., Schaffner, U., Eschen, R., Abebe, A., Choge, S. K., Nigatu, L., . . . Allan, E.

(2019). Direct and indirect effects of invasive species: Biodiversity loss is a major

mechanism by which an invasive tree affects ecosystem functioning. *Journal of Ecology,*

107(6), 2660-2672.

Liu, X., Zhou, Y., Ouyang, S. and Wu, X., 2023. Environmental Correlates to Genetic Diversity

and Structure in Invasive Apple Snail (Pomacea canaliculata) Populations in China.

Diversity, 15(10), p.1048.

Liu, X., Zhou, Y., Ouyang, S., & Wu, X. (2019). Phylogeographic patterns and demographic

history of Pomacea canaliculata and Pomacea maculata from different countries

(Ampullariidae, Gastropoda, Mollusca). *Nature Conservation, 36*, 71-92.

Lowe, S., Browne, M., Boudjelas, S. and De Poorter, M., 2000. 100 of the world's worst invasive

alien species: a selection from the global invasive species database (Vol. 12). Auckland:

Invasive Species Specialist Group

Lv, S., Zhang, Y., Liu, H. X., Hu, L., Liu, Q., Wei, F. R., . . . Zhou, X. N. (2013). Phylogenetic

evidence for multiple and secondary introductions of invasive snails: Pomacea species in

the People's Republic of China. *Diversity and distributions, 19*(2), 147-156.

Matsukura, K., & Wada, T. (2017). Identification of Pomacea species using molecular

techniques. *Biology and management of invasive apple snails*, 33-43.

Matsukura, K., Izumi, Y., Yoshida, K., & Wada, T. (2016). Cold tolerance of invasive freshwater

snails, Pomacea canaliculata, P. maculata, and their hybrids helps explain their different

distributions. *Freshwater Biology, 61*(1), 80-87.

Matsukura, K., Okuda, M., Kubota, K., & Wada, T. (2008). Genetic divergence of the genus

Pomacea (Gastropoda: Ampullariidae) distributed in Japan, and a simple molecular

method to distinguish P. canaliculata and P. insularum. *Applied Entomology and Zoology,*

43(4), 535-540.

McNeely, J. (2001). Invasive species: a costly catastrophe for native biodiversity. *Land Use and*

Water Resources Research, 1(1732-2016-140260).

Päckert, M., Martens, J., & Severinghaus, L. L. (2009). The Taiwan Firecrest (Regulus

goodfellowi) belongs to the Goldcrest assemblage (Regulus regulus sl): evidence from

mitochondrial DNA and the territorial song of the Regulidae. *Journal of Ornithology,*

150, 205-220.

Rama Rao, S., Liew, T.-S., Yow, Y.-Y., & Ratnayeke, S. (2018). Cryptic diversity: Two

morphologically similar species of invasive apple snail in Peninsular Malaysia. *PLoS*

One, 13(5), e0196582.

Rambaut, A., & Drummond, A. (2010). FigTree v1. 4.4. Institute of Evolutionary Biology,

University of Edinburgh. In.

Rambaut A, Drummond AJ, Xie D, Baele G and Suchard MA (2018) Posterior summarisation in

Bayesian phylogenetics using Tracer 1.7. *Systematic Biology.*

Ranamukhaarachchi, S., & Wickramasinghe, S. (2006). Golden apple snails in the world:

introduction, impact, and control measures. *Global advances in ecology and management*

of golden apple snails. Philippine Rice Research Institute, Nueva Ecija, 133-152.

Sakai, A.K., Allendorf, F.W., Holt, J.S., Lodge, D.M., Molofsky, J., With, K.A., Baughman, S.,

Cabin, R.J., Cohen, J.E., Ellstrand, N.C. and McCauley, D.E., 2001. The population

biology of invasive species. Annual review of ecology and systematics, 32(1), pp.305-

332.

Saltonstall, K., 2002. Cryptic invasion by a non-native genotype of the common reed, Phragmites

australis, into North America. Proceedings of the National Academy of Sciences, 99(4),

pp.2445-2449.

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: molecular evolutionary genetics

analysis version 11. *Molecular biology and evolution, 38*(7), 3022-3027.

Tsai, H.-C., Liu, Y.-C., Kunin, C. M., Lee, S. S.-J., Chen, Y.-S., Lin, H.-H., . . . Yen, M.-Y.

(2001). Eosinophilic meningitis caused byAngiostrongylus cantonensis: report of 17

cases. *The American journal of medicine, 111*(2), 109-114.

Turbelin, A. J., Cuthbert, R. N., Essl, F., Haubrock, P. J., Ricciardi, A., & Courchamp, F. (2023).

Biological invasions are as costly as natural hazards. *Perspectives in Ecology and*

Conservation.

Wu, J.-Y., Meng, P.-J., Liu, M.-Y., Chiu, Y.-W., & Liu, L.-L. (2010). A high incidence of

imposex in Pomacea apple snails in Taiwan: a decade after triphenyltin was banned. *Zool*

Stud, 49, 85-93.

Wu, J.Y., Wu, Y.T., Li, M.C., Chiu, Y.W., Liu, M.Y. and Liu, L.L., 2011. Reproduction and

Juvenile Growth of the Invasive Apple Snails Pomacea canaliculata and P.

scalaris(Gastropoda: Ampullariidae) in Taiwan. Zoological Studies, 50(1), pp.61-68.

Yang, P., Chen, Y., Lee, W., & Chen, Y. (2006). Golden apple snail management and prevention

in Taiwan. *Global advances in ecology and management of golden apple snails*, 169-179.

Yang, Q. Q., Ip, J. C. H., Zhao, X. X., Li, J. N., Jin, Y. J., Yu, X. P., & Qiu, J. W. (2022).

Molecular analyses revealed three morphologically similar species of non☐native apple

snails and their patterns of distribution in freshwater wetlands of Hong Kong. *Diversity*

and distributions, 28(1), 97-111.

Yang, Q.Q., He, C., Liu, G.F., Yin, C.L., Xu, Y.P., Liu, S.W., Qiu, J.W. and Yu, X.P., 2020.

Introgressive hybridization between two non☐native apple snails in China: widespread

hybridization and homogenization in egg morphology. Pest management science, 76(12),

pp.4231-4239.

Yang, Q.-Q., Liu, S.-W., He, C., & Yu, X.-P. (2018). Distribution and the origin of invasive

apple snails, Pomacea canaliculata and P. maculata (Gastropoda: Ampullariidae) in

China. *Scientific Reports, 8*(1), 1-8.

Zhao, B., Luo, M., Zhang, J., Liu, Y., Deng, Z., & Gong, X. (2022). Genetic diversity of two

globally invasive snails in Asia and Americas in relation with agricultural habitats and

climate factors. Diversity, 14(12), 1069.

APPENDIX 1.

Details of sampling sites, number of samples (n), and sampling locations across Taiwan for
Pomacea spp,

Sample site (n)	Sample ID (n)	Location	Date of sampling
East (42)	ES (HU+TT)		
Hualien (29)	HU1 (5)	23°57'29.3"N 121°33'57.0"E	11/5/2022
	HU2 (10)	23°39'15.9"N 121°32'20.5"E	11/5/2022
	HU3 (14)	23°14'18.7"N 121°18'51.0"E	12/5/2022
Taitung (13)	TT1 (3)	23°07'34.6"N 121°12'42.5"E	13/5/2022
	TT2 (5)	22°57'45.7"N 121°07'52.0"E	13/5/2022
	TT3 (5)	22°46'05.7"N 121°07'23.7"E	14/5/2022
Middle (56)	MD (CH+MIO+TC+YL)		
Changhua (15)	CH1 (4)	24°03'37.1"N 120°25'09.5"E	2/9/2021
	CH2 (5)	24°05'18.5"N 120°35'15.3"E	2/9/2021
	CH3 (6)	24°09'58.5"N 120°29'36.1"E	2/9/2021
Miaoli (14)	MIO1 (5)	24°39'23.9"N 120°51'25.6"E	7/7/2022
	MIO2 (4)	24°29'32.3"N 120°41'45.3"E	7/7/2022
	MIO3 (5)	24°24'13.6"N 120°40'06.9"E	8/7/2022
Taichung (14)	TC1 (5)	24°12'46.9"N 120°32'33.8"E	9/7/2022
	TC2 (6)	24°21'36.9"N 120°38'14.0"E	9/7/2022
	TC3 (3)	24°19'10.2"N 120°43'00.6"E	10/7/2022
Yunlin (13)	YL1 (3)	23°38'05.5"N 120°12'29.9"E	2/8/2022
	YL2 (5)	23°43'36.2"N 120°15'42.9"E	2/8/2022
	YL3 (5)	23°46'34.8"N 120°26'04.7"E	2/8/2022
North (50)	NR (HS+TP+TY+YI)		
Hsinchu (14)	HS1 (5)	24°48'05.7"N 120°55'17.1"E	5/8/2022
	HS2 (5)	24°50'57.8"N 120°58'17.6"E	5/8/2022
	HS3 (4)	24°54'47.3"N 121°01'06.8"E	5/8/2022

Taipei (12)	TP1 (7)	25°07'29.0"N 121°29'15.1"E	12/7/2022
	TP2 (5)	25°13'34.6"N 121°37'21.8"E	12/7/2022
Taoyuan (11)	TY1 (3)	24°58'56.2"N 121°07'29.7"E	12/7/2022
	TY2 (7)	25°01'40.8"N 121°13'00.2"E	15/7/2022
	TY3 (4)		15/7/2022
Yilan (13)	YI1 (4)	24°38'20.1"N 121°49'10.9"E	4/9/2021
	YI2 (6)	24°42'01.3"N 121°44'10.6"E	4/9/2021
	YI3 (3)	24°48'27.4"N 121°46'51.5"E	4/9/2021
South (106)	SU (CH+KSH+PT+TN)		
Chiayi (24)	CH1 (8)	23°32'26.6"N 120°22'50.2"E	1/9/2021
	CH2 (7)	23°28'44.5"N 120°10'10.5"E	1/9/2021
	CH3 (9)	23°28'41.7"N 120°12'55.8"E	1/9/2021
Kaohsiung (22)	KSH1 (8)	22°30'42.0"N 120°23'10.7"E	12/4/2022
	KSH2 (8)	22°44'28.8"N 120°16'34.7"E	12/4/2022
	KSH3 (6)	22°49'10.7"N 120°17'53.1"E	12/4/2022
Pingtung (34)	PT1 (8)	22°00'40.2"N 120°44'13.2"E	05/01/2023
	PT2 (14)	22°49'56.2"N 120°36'23.7"E	05/01/2023
	PT3 (12)	22°28'18.4"N 120°31'56.4"E	05/01/2023
Tainan (26)	TN1 (8)	22°58'43.7"N 120°15'50.9"E	3/9/2021
	TN2 (10)	23°04'41.6"N 120°18'24.8"E	3/9/2021
	TN3 (8)	23°12'26.1"N 120°16'15.6"E	3/9/2021

689 **APPENDIX 2.**

690 List of GenBank accession numbers used to prepare *Pomacea canaliculata* global mitochondrial

691 COI dataset (Dataset 3)

Continent	Country	n	Sequence		
Africa	Kenya	3	KN_MW363565; KN_MW363566; KN_MW363568		
Asia	China	583	CHI_MK862299;	CHI_MK862295;	CHI_MK862294;
			CHI_MK862293;	CHI_MK189419;	CHI_MK189413;
			CHI_MK189403;	CHI_MK189394;	CHI_MK189393;
			CHI_MK189362;	CHI_MK189360;	CHI_MK189359;
			CHI_MK189356;	CHI_MK189354;	CHI_MK189350;
			CHI_MK189349;	CHI_MK189348;	CHI_MK189347;
			CHI_MK189345;	CHI_MK189344;	CHI_MK189342;
			CHI_MK189340;	CHI_MK189339;	CHI_MK189338;
			CHI_MK189332;	CHI_MK189331;	CHI_MK189330;
			CHI_MK189329;	CHI_MH293388;	CHI_MH293375;
			CHI_MH293373;	CHI_MH293372;	CHI_MH293370;
			CHI_MH293369;	CHI_MH293368;	CHI_MH293367;
			CHI_MH293366;	CHI_MH293364;	CHI_MH293363;
			CHI_MH293362;	CHI_MH293361;	CHI_MH293360;

CHI_MH293358;	CHI_MH293357;	CHI_MH293356;
CHI_MH293354;	CHI_MH293353;	CHI_MH293345;
CHI_MH293337;	CHI_MH293336;	CHI_MH293333;
CHI_MH211415;	CHI_MH211414;	CHI_MH211413;
CHI_MH211412;	CHI_MH211410;	CHI_MH211409;
CHI_MH211408;	CHI_MH211407;	CHI_MH211406;
CHI_MH211405;	CHI_MH211404;	CHI_MH211403;
CHI_MH211402;	CHI_KT884521;	CHI_KT884519;
CHI_KT852762;	CHI_KT852761;	CHI_KT852760;
CHI_KT852759;	CHI_KT852758;	CHI_KT852722;
CHI_KT852721;	CHI_KT852720;	CHI_KT852719;
CHI_KT852718;	CHI_KT852717;	CHI_KT852716;
CHI_KT852715;	CHI_KT852714;	CHI_KT852713;
CHI_KT852712;	CHI_KT852711;	CHI_KT852710;
CHI_KT852709;	CHI_KT852708;	CHI_KT852707;
CHI_KT693026;	CHI_KT693023;	CHI_KT693017;
CHI_KT693015;	CHI_KT693013;	CHI_KT693011;
CHI_KT693010;	CHI_KT693007;	CHI_KT693006;
CHI_KT693005;	CHI_KT693003;	CHI_KT693002;
CHI_KT693000;	CHI_KT692999;	CHI_KT692997;
CHI_KT692996;	CHI_KT692995;	CHI_KT692994;
CHI_KT692992;	CHI_KT692990;	CHI_KT692989;
CHI_KR021013;	CHI_KR021012;	CHI_KR021011;
CHI_KR021010;	CHI_KR021008;	CHI_KR021007;
CHI_KR021006;	CHI_KR021004;	CHI_KR021003;
CHI_KR021002;	CHI_KR021001;	CHI_KR021000;
CHI_KR020999;	CHI_KR020998;	CHI_KR020997;
CHI_KR020996;	CHI_KR020995;	CHI_KR020994;
CHI_KR020978;	CHI_KR020977;	CHI_KR020976;
CHI_KR020975;	CHI_KR020974;	CHI_KR020973;
CHI_KR020972;	CHI_KR020971;	CHI_KR020970;
CHI_KR020969;	CHI_KR020967;	CHI_KR020966;
CHI_KR020965;	CHI_KR020964;	CHI_KR020963;
CHI_KR020962;	CHI_KP310435;	CHI_KP310434;
CHI_KP310433;	CHI_KP310432;	CHI_KP310431;
CHI_KP310430;	CHI_KP310429;	CHI_KP310428;
CHI_KP310427;	CHI_KP310426;	CHI_KP310425;
CHI_KP310424;	CHI_KP310423;	CHI_KP310422;
CHI_KP310421;	CHI_KP310420;	CHI_KP310419;
CHI_KP310418;	CHI_KP310417;	CHI_KP310416;
CHI_KP310415;	CHI_KP310414;	CHI_KP310413;
CHI_KP310412;	CHI_KP310411;	CHI_KP310410;
CHI_KP310409;	CHI_KP310408;	CHI_KP310407;
CHI_KP310406;	CHI_KP310405;	CHI_KP310404;
CHI_KP310403;	CHI_KP310402;	CHI_KP310401;
CHI_KP310400;	CHI_KP310399;	CHI_KP310398;

CHI_KP310397;	CHI_KP310396;	cCHI_KP310395;
CHI_KP310394;	CHI_KP310393;	CHI_KP310392;
CHI_KP310391;	CHI_KP310390;	CHI_KP310389;
CHI_KP310388;	CHI_KP310387;	CHI_KP310386;
CHI_KP310385;	CHI_KP310384;	CHI_KP310383;
CHI_KP310382;	CHI_KP310381;	CHI_KP310380;
CHI_KP310379;	CHI_KP310378;	CHI_KP310377;
CHI_KP310376;	CHI_KP310375;	CHI_KP310374;
CHI_KP310373;	CHI_KP310372;	CHI_KP310371;
CHI_KP310370;	CHI_KP310369;	CHI_KP310368;
CHI_KP310367;	CHI_KP310366;	CHI_FJ946820;
CHI_KR021009;	CHI_KP310290;	CHI_MK189424;
CHI_MK189410;	CHI_MK189412;	CHI_MK189407;
CHI_MK189404;	CHI_MK189401;	CHI_MK189395;
CHI_MK189391;	CHI_MK189387;	CHI_MK189385;
CHI_MK189383;	CHI_MK189382;	CHI_MK189381;
CHI_MK189380;	CHI_MK189379;	CHI_MK189378;
CHI_MK189377;	CHI_MK189375;	CHI_MK189374;
CHI_MK189373;	CHI_MK189372;	CHI_MK189370;
CHI_MK189369;	CHI_MK189368;	CHI_MK189365;
CHI_MK189364;	CHI_MK189363;	CHI_MK189361;
CHI_MK189358;	CHI_MK189357;	CHI_MK189355;
CHI_MK189352;	CHI_MK189351;	CHI_MK189343;
CHI_MK189341;	CHI_MK189337;	CHI_MK189336;
CHI_MK189335;	CHI_MK189334;	CHI_MK189333;
CHI_MK189328;	CHI_MK189327;	CHI_MK189326;
CHI_MK189325;	CHI_MK189324;	CHI_MK189323;
CHI_MH293450;	CHI_MH293449;	CHI_MH293447;
CHI_MH293446;	CHI_MH293445;	CHI_MH293444;
CHI_MH293443;	CHI_MH293441;	CHI_MH293440;
CHI_MH293438;	CHI_MH293437;	CHI_MH293436;
CHI_MH293435;	CHI_MH293434;	CHI_MH293433;
CHI_MH293432;	CHI_MH293431;	CHI_MH293429;
CHI_MH293428;	CHI_MH293427;	CHI_MH293426;
CHI_MH293425;	CHI_MH293424;	CHI_MH293422;
CHI_MH293421;	CHI_MH293420;	CHI_MH293419;
CHI_MH293418;	CHI_MH293417;	CHI_MH293416;
CHI_MH293415;	CHI_MH293414;	CHI_MH293413;
CHI_MH293412;	CHI_MH293411;	CHI_MH293410;
CHI_MH293409;	CHI_MH293408;	CHI_MH293407;
CHI_MH293406;	CHI_MH293405;	CHI_MH293404;
CHI_MH293403;	CHI_MH293402;	CHI_MH293401;
CHI_MH293400;	CHI_MH293399;	CHI_MH293398;
CHI_MH293397;	CHI_MH293396;	CHI_MH293395;
CHI_MH293394;	CHI_MH293393;	CHI_MH293392;
CHI_MH293390;	CHI_MH293389;	CHI_MH293386;

CHI_MH293385;	CHI_MH293384;	CHI_MH293383;
CHI_MH293382;	CHI_MH293381;	CHI_MH293380;
CHI_MH293379;	CHI_MH293378;	CHI_MH293377;
CHI_MH293352;	CHI_MH293351;	CHI_MH293350;
CHI_MH293349;	CHI_MH293348;	CHI_MH293347;
CHI_MH293346;	CHI_MH293344;	CHI_MH293343;
CHI_MH293342;	CHI_MH293341;	CHI_MH293340;
CHI_MH293339;	CHI_MH293338;	CHI_MH293334;
CHI_MH293332;	CHI_MH293331;	CHI_MH293330;
CHI_MH293329;	CHI_MH293328;	CHI_MH293327;
CHI_MH293326;	CHI_MH293325;	CHI_MH293324;
CHI_MH293323;	CHI_MH293322;	CHI_MH211411;
CHI_MH211401;	CHI_MH211400;	CHI_KT884522;
CHI_KT884520;	CHI_KT884518;	CHI_KT852756;
CHI_KT852755;	CHI_KT852754;	CHI_KT852753;
CHI_KT852752;	CHI_KT852751;	CHI_KT852750;
CHI_KT852749;	CHI_KT852748;	CHI_KT852747;
CHI_KT852746;	CHI_KT852745;	CHI_KT852744;
CHI_KT852743;	CHI_KT852742;	CHI_KT852741;
CHI_KT852740;	CHI_KT852739;	CHI_KT852738;
CHI_KT852737;	CHI_KT852736;	CHI_KT852735;
CHI_KT852734;	CHI_KT852733;	CHI_KT852732;
CHI_KT852731;	CHI_KT852730;	CHI_KT852729;
CHI_KT852728;	CHI_KT852727;	CHI_KT693022;
CHI_KT693021;	CHI_KT693020;	CHI_KT693019;
CHI_KT693018;	CHI_KT693016;	CHI_KT693014;
CHI_KT693009;	CHI_KT692988;	CHI_KR020993;
CHI_KR020992;	CHI_KR020991;	CHI_KR020990;
CHI_KR020989;	CHI_KR020988;	CHI_KR020987;
CHI_KR020986;	CHI_KR020985;	CHI_KR020984;
CHI_KR020983;	CHI_KR020980;	CHI_KR020961;
CHI_KR020960;	CHI_KR020959;	CHI_KR020958;
CHI_KR020957;	CHI_KR020956;	CHI_KR020955;
CHI_KR020954;	CHI_KR020953;	CHI_KR020952;
CHI_KR020951;	CHI_KR020950;	CHI_KR020949;
CHI_KR020948;	CHI_KR020947;	CHI_KR020946;
CHI_KR020945;	CHI_KR020944;	CHI_KR020943;
CHI_KR020942;	CHI_KP310365;	CHI_KP310364;
CHI_KP310363;	CHI_KP310362;	CHI_KP310361;
CHI_KP310360;	CHI_MK189423;	CHI_KP310359;
CHI_KP310358;	CHI_KP310357;	CHI_KP310356;
CHI_KP310355;	CHI_KP310354;	CHI_KP310353;
CHI_KP310352;	CHI_KP310351;	CHI_KP310350;
CHI_KP310349;	CHI_KP310348;	CHI_KP310347;
CHI_KP310346;	CHI_KP310345;	CHI_KP310344;
CHI_KP310343;	CHI_KP310342;	CHI_KP310341;

			CHI_KP310340;	CHI_KP310339;	CHI_KP310338;
			CHI_KP310337;	CHI_KP310336;	CHI_KP310335;
			CHI_KP310334;	CHI_KP310333;	CHI_KP310332;
			CHI_KP310331;	CHI_KP310330;	CHI_KP310329;
			CHI_KP310328;	CHI_KP310327;	CHI_KP310326;
			CHI_KP310325;	CHI_KP310324;	CHI_KP310323;
			CHI_KP310322;	CHI_KP310321;	CHI_KP310320;
			CHI_KP310319;	CHI_KP310318;	CHI_KP310317;
			CHI_KP310316;	CHI_KP310315;	CHI_KP310314;
			CHI_KP310313;	CHI_KP310312;	CHI_KP310311;
			CHI_KP310310;	CHI_KP310309;	CHI_KP310308;
			CHI_KP310306;	CHI_KP310305;	CHI_KP310304;
			CHI_KP310303;	CHI_KP310302;	CHI_KP310301;
			CHI_KP310300;	CHI_KP310299;	CHI_KP310298;
			CHI_KP310297;	CHI_KP310296;	CHI_KP310295;
			CHI_KP310294;	CHI_KP310293;	CHI_KP310292;
			CHI_KP310291;	CHI_KP310289;	CHI_KP310288;
			CHI_KP310287;	CHI_KP310286;	CHI_KP310285;
			CHI_KP310284;	CHI_KP310283;	CHI_KP310282;
			CHI_KP310281;	CHI_KP310280;	CHI_KP310279;
			CHI_KP310278;	CHI_KP310277;	CHI_KP310276;
			CHI_KP310275;	CHI_KP310274;	CHI_KP310273;
			CHI_KP310272;	CHI_KP310271;	CHI_KP310270;
			CHI_KP310269;	CHI_KP310268;	CHI_KP310267;
			CHI_KP310266;	CHI_KP310265;	CHI_KP310264;
			CHI_FJ946824;	CHI_EU528579;	CHI_KP310307;
			CHI_MH293391;	CHI_MH293387;	CHI_MH293376;
			CHI_MH293374;	CHI_MH293371;	CHI_MH293365;
			CHI_MH293359;	CHI_MH293355;	CHI_KT852726;
			CHI_KT852725;	CHI_KP310444;	CHI_KP310443;
			CHI_KP310442;	CHI_FJ946823;	CHI_KT852757;
			CHI_KT852724;	CHI_KT852723;	CHI_KR021014;
			CHI_KR020981;	CHI_KR020979;	CHI_KP310445;
			CHI_KP310441;	CHI_KP310440;	CHI_KP310439;
			CHI_KP310436;	CHI_FJ946822;	CHI_KT852706;
			CHI_KR021005;	CHI_KT693029;	CHI_KT693028;
			CHI_KT693027;	CHI_KT693025;	CHI_KT693024;
			CHI_KR106989;	CHI_KR021020;	CHI_KR021019;
			CHI_KR021018;	CHI_KR021017;	CHI_KR021016;
			CHI_KR021015;	CHI_KR020982;	CHI_KP310438;
			CHI_KP310437;	CHI_EU528578;	CHI_FJ946827;
			CHI_FJ946826; CHI_FJ946825; CHI_FJ946821		
Asia	China-Hong Kong	2	HK_KT313034; HK_KT313035		
	Indonesia	3	IN_EU528580; IN_EU528581; IN_KY574007		

	Japan	18	JP_AB433757; JP_AB433758; JP_AB433759; JP_AB433760; JP_AB433761; JP_AB433762; JP_AB433763; JP_AB433764; JP_AB433765; JP_AB433766; JP_AB433767; JP_AB433768; JP_AB433769; JP_AB433770; JP_AB433771; JP_AB433772; JP_AB433773; JP_AB433774
	Malaysia	23	ML_MN623427; ML_MN623428; ML_MN623429; ML_MN623430; ML_MN623431; ML_MN623433; ML_MN623434; ML_MN623435; ML_MN623436; ML_MN623437; Ml_MN623438; ML_MN623439; ML_MN623440; ML_MN758725; ML_MN758726; ML_MN758727; ML_MN758728; ML_MN758729; ML_MN758730; ML_MN758731; ML_MN758732; ML_MW591143
	Myanmar	1	MY_EU528582
	Philippines	12	PH_EU528483; PH_EU528484; PH_EU528485; PH_EU528486; PH_EU528487; PH_EU528488; PH_EU528490; PH_EU528491; PH_EU528492; PH_EU528493; PH_EU528494
	Singapore	1	SN_KY081757
	Taiwan	249	TW_EU528584; TW_EU528586; TW_EU528587 + 246 Present study
	Thailand	2	TH_EU528497; TH_EU528498
North America	USA	19	US_EF514967; US_EF514968; US_EF514969; US_EF514970; US_EF514971; US_EF514972; US_EF514973; US_EF514974; US_EF514975; US_EF514976; US_EF514977; US_EF514978; US_EF514979; US_EF514980; US_EF514981; US_EF514982; US_EF514983; US_EF514984; US_EU523129
Oceania	Papua New Guinea	1	PNG_EU528499
South America	Argentina	32	ARG_AB728574; ARG_AB728575; ARG_AB728576; ARG_AB728577; ARG_AB728578; ARG_AB728585; ARG_AB728586; ARG_AB728587; ARG_AB728588; ARG_EF514961; ARG_EF514962; ARG_EF514963; ARG_EF514964; ARG_EF514965; ARG_EF514966; ARG_EU528482; ARG_EU528489; ARG_EU528508; ARG_EU528509; ARG_EU528510; ARG_EU528513; ARG_EU528515; ARG_EU528516; ARG_EU528520; ARG_EU528522; ARG_EU528523; ARG_EU528524; ARG_EU528525; ARG_EU528528; ARG_EU528529; ARG_FJ710314; ARG_FJ710315
	Chile	4	Cle_KX965671; Cle_KX965672; Cle_KX965673; Cle_KX965674
	Uruguay	8	UR_EU528593; UR_FJ710313; UR_MK992483; UR_MK992484; UR_MK992485; UR_MK992486; UR_MK992487; UR_MK992488

693　**APPENDIX 3.**

694 List of GenBank accession numbers used to prepare *Pomacea maculata* mitochondrial COI
695 dataset (Dataset 2)

Continent	Country	n	Sequences
Asia	China	44	CHI_FJ946828; CHI_KP310480; CHI_KP310481; CHI_KP310482; CHI_KP310483; CHI_KP310484; CHI_KP310485; CHI_KP310486; CHI_KP310487; CHI_KP310488; CHI_KP310489; CHI_KP310490; CHI_KP310491; CHI_KP310492; CHI_KP310493; CHI_KP310494; CHI_KP310495; CHI_KP310496; CHI_MK189346; CHI_MK189353; CHI_MK189388; CHI_MK189389; CHI_MK189390; CHI_MK189392; CHI_MK189396; CHI_MK189397; CHI_MK189398; CHI_MK189399; CHI_MK189400; CHI_MK189402; CHI_MK189405; CHI_MK189406; CHI_MK189408; CHI_MK189409; CHI_MK189411; CHI_MK189414; CHI_MK189415; CHI_MK189416; CHI_MK189417; CHI_MK189418; CHI_MK189420; CHI_MK189421; CHI_MK189422; CHI_MK862301
	Japan	6	JP_AB433776; JP_AB433777; JP_AB433778; JP_AB433779; JP_AB433780; JP_AB433781
	Malaysia	12	ML_MN623417; ML_MN623418; ML_MN623419; ML_MN623420; ML_MN623421; ML_MN623422; ML_MN623423; ML_MN623424; ML_MN623425; ML_MN623426; ML_MN623432
	Taiwan	8	Present Study
	Thailand	1	TH_EU528496
	Vietnam	1	VT_EU528502
Europe	Spain	8	SP_GU133205; SP_GU133206; SP_GU236486; SP_GU236487; SP_GU236488; SP_GU236489; SP_GU236490; SP_GU236491
North America	USA	66	US_EF514942; US_EF514943; US_EF514944; US_EF514985; US_EF515013; US_EF515014; US_EF515015; US_EF515016; US_EF515017; US_EF515018; US_EF515019; US_EF515020; US_EF515021; US_EF515022; US_EF515023; US_EF515024; US_EF515025; US_EF515026; US_EF515027; US_EF515028; US_EF515029; US_EF515030; US_EF515031; US_EF515032; US_EF515033; US_EF515034; US_EF515035; US_EF515036; US_EF515037; US_EF515038; US_EF515039; US_EF515040; US_EF515041; US_EF515042; US_EF515043; US_EF515044; US_EF515045; US_EF515046; US_EF515047; US_EF515048; US_EF515049; US_EF515050; US_EF515051; US_EF515052; US_EF515053; US_EF515054; US_EF515055; US_EF515056;

			US_EF515057; US_EF515058; US_JX845573; US_KM107825; US_KM107826; US_KM107827; US_KM107828; US_KM107829; US_KM107830; US_KR350494; US_KR350495; US_KR350496; US_KR350497; US_KR350498; US_KR350499; US_KX149128; US_KX810826; US_MK992475
Oceania	New Zealand	1	NZ_HQ908057
South America	Argentina	14	ARG_AB728579; ARG_AB728580; ARG_AB728582; ARG_AB728583; ARG_AB728584; ARG_EF514945; ARG_EU528481; ARG_EU528495; ARG_EU528514; ARG_EU528521; ARG_EU528526; ARG_EU528527; ARG_EU528530; ARG_EU528531
	Brazil	60	BR_EF514986; BR_EF514987; BR_EF514988; BR_EF514989; BR_EF514990; BR_EF514991; BR_EF514992; BR_EF514993; BR_EF514994; BR_EF514995; BR_EF514996; BR_EF514997; BR_EF514998; BR_EF514999; BR_EF515000; BR_EF515001; BR_EF515002; BR_EF515003; BR_EF515004; BR_EF515005; BR_EF515006; BR_EF515007; BR_EF515008; BR_EF515009; BR_EF515010; BR_EF515011; BR_EF515012; BR_EU528533; BR_EU528538; BR_EU528539; BR_EU528540; BR_EU528544; BR_EU528549; BR_EU528550; BR_EU528551; BR_EU528552; BR_EU528554; BR_EU528557; BR_EU528558; BR_EU528559; BR_EU528560; BR_EU528561; BR_EU528562; BR_EU528563; BR_EU528566; BR_EU528567; BR_EU528568; BR_EU528569; BR_EU528570; BR_EU528571; BR_EU528572; BR_EU528575; BR_EU528576; BR_MK992470; BR_MK992471; BR_MK992472 ; BR_MK992479 ; BR_MK992480; BR_MK992481; BR_MK992482
	Uruguay	5	UR_MK992473; UR_MK992474; UR_MK992476; UR_MK992477; UR_MK992478

696
697
698 **APPENDIX 4.**
699 **Genetic distance (p-distance) among the taxa used in the phylogenetic tree, Pcan:*P.***
700 ***canaliculate*; Pma: *P. maculata*; Plin: *P. lineata*, Ppal: *P. paludosa*, Pdif: *P. diffusa*, and PC:**
701 *P. scalaris*
702

	Pcan	Ppal	Pdif	Plin	Pma	PC

Pcan					
Ppal	0.1333				
Pdif	0.1353	0.1429			
Plin	0.1020	0.1252	0.1411		
Pma	0.0928	0.1246	0.1523	0.0852	
PC	0.1456	0.1658	0.1340	0.1517	0.1584

703
704
705 **APPENDIX 5.**
706 Haplotype distribution of *Pomacea* spp. in 14 counties/cities across Taiwan (full names of
707 abbreviation can be found in APPENDIX 1.)
708

Pomacea canaliculata

	HU	TT	CH	MIO	TC	YL	HS	TP	TY	YI	CY	KSH	PT	TN	Total	%
Hap_1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	0.407
Hap_2	1	6	-	5	3	-	1	-	3	2	4	5	10	4	44	17.886
Hap_3	24	4	12	8	7	6	12	11	8	7	14	11	15	16	155	63.008
Hap_4	-	2	-	-	-	-	-	-	-	-	-	2	-	1	5	2.033
Hap_5	-	-	-	-	1	-	-	1	-	-	1	-	1	1	5	2.033
Hap_6	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	0.407
Hap_7	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	0.407
Hap_8	2	-	-	-	-	7	-	-	-	-	-	-	7	-	16	6.504
Hap_9	-	-	1	-	-	-	1	-	-	-	-	1	-	-	3	1.220
Hap_10	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.407
Hap_11	-	-	1	-	3	-	-	-	-	1	2	-	-	-	7	2.846
Hap_12	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.407
Hap_13	-	-	-	-	-	-	-	-	-	1	-	-	-	-	1	0.407
Hap_14	-	-	-	-	-	-	-	-	-	1	-	-	-	-	1	0.407
Hap_15	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	0.407
Hap_16	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	0.407
Hap_17	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.407

															Total	%
Hap_18	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.407

Pomacea maculata

	HU	TT	CH	MIO	TC	YL	HS	TP	TY	YI	CY	KSH	PT	TN	Total	%
Hap_19	-	-	-	1	-	-	-	-	-	1	-	3	-	-	5	62.5
Hap_20	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	12.5
Hap_21	-	-	-	-	-	-	-	-	-	-	-	-	1	-	1	12.5
Hap_22	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	12.5

711 **APPENDIX 6.**
712 Haplotype distribution of *Pomacea canaliculata* from the global mitochondrial COI dataset
713 (Dataset 3)

	TW	Cle	AR	UR	US	PNG	HK	CH	JP	ML	TH	SN	PH	IN	MY	KN	Total	%
Hap_1	162	-	9	-	17	1	2	214	7	11	-	-	-	-	-	-	423	44.4
Hap_2	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.21
Hap_3	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_4	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_5	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_6	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_7	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_8	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_9	1	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	3	0.3
Hap_10	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_11	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_12	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_13	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_14	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_15	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.21
Hap_16	46	-	-	-	-	-	-	1	4	2	-	-	-	-	-	3	56	5.8
Hap_17	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_18	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	3	0.3
Hap_19	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_20	1	-	-	-	-	-	-	319	1	-	-	-	-	-	-	-	321	33.
Hap_21	-	-	-	-	-	-	-	14	-	-	-	-	1	-	-	-	15	1.5
Hap_22	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1	0.1
Hap_23	5	-	-	-	-	-	-	11	3	-	-	-	-	-	-	-	19	2.0
Hap_24	1	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	2	0.21
Hap_25	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	0.6
Hap_26	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1

																	Total	
Hap_27	16	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	16	1.6
Hap_28	4	-	-	-	-	-	-	-	2	-	-	1	-	-	-	-	7	0.74
Hap_29	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_30	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_31	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_32	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_33	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_34	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_35	-	-	-	-	-	-	-	2	-	-	-	-	-	-	-	-	2	0.2
Hap_36	-	-	-	-	-	-	-	16	1	-	-	-	-	-	-	-	17	1.7
Hap_37	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_38	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	2	0.2
Hap_39	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_40	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_41	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Hap_42	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	2	0.2
Hap_43	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1	0.1
Hap_44	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1	0.1
Hap_45	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1	0.1
Hap_46	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1	0.1
Hap_47	-	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	0.4
Hap_48	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	2	0.2
Hap_49	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	1	0.1
Hap_50	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1	0.1
Hap_51	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	1	0.1
Hap_52	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_53	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_54	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_55	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_56	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_57	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_58	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_59	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_60	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_61	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	0.1
Hap_62	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	0.1
Hap_63	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	0.1
Hap_64	-	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-	2	0.2
Hap_65	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.1
Total	249	4	32	8	19	1	2	582	18	14	2	1	12	3	1	3	951	

 APPENDIX 7.

719 Comparison of Taiwan haplotype positions in different datasets, for *Pomacea canaliculata*

720 Dataset 1 (Taiwan sequence) compared with Dataset 2 (global dataset of *Pomacea canaliculata*),

721 and for *Pomacea maculata* Dataset 1 (Taiwan sequences) compared with Dataset 3 (global

722 dataset of *Pomacea maculata*)

Dataset 1	Networks	Dataset 3 (*P. canaliculata*)	Network
P. canaliculata			
Hap 1	Network Tw-A	Hap 24	Network Wc-A
Hap 2	Network Tw-B	Hap 16	Network Wc-B
Hap 3	Network Tw-A	Hap 1	Network Wc-A
Hap 4	Network Tw-B	Hap 25	Network Wc-B
Hap 5	Network Tw-B	Hap 23	Network Wc-B
Hap 6	Network Tw-A	Hap 26	Network Wc-A
Hap 7	Network Tw-C	Hap 9	Network Wc-C
Hap 8	Network Tw-B	Hap 27	Network Wc-B
Hap 9	Network Tw-C	Hap 28	Network Wc-C
Hap 10	Network Tw-B	Hap 16	Network Wc-B
Hap 11	Network Tw-A	Hap 1	Network Wc-A
Hap 12	Network Tw-A	Hap 1	Network Wc-A
Hap 13	Network Tw-B	Hap 29	Network Wc-B
Hap 14	Network Tw-B	Hap 20	Network Wc-B
Hap 15	Network Tw-A	Hap 30	Network Wc-A
Hap 16	Network Tw-A	Hap 31	Network Wc-A
Hap 17	Network Tw-B	Hap 32	Network Wc-B
Hap 18	Network Tw-B	Hap 33	Network Wc-B
Dataset1	**Networks**	**Dataset 2 (*P. maculata*)**	**Networks**
P. maculata			
Hap 19 (exclude KSH 27)	Network Tw-A	Hap 1	Network Wm-A
Hap 19 (only **KSH 27**)	Network Tw-A	Hap 8	Network Wm-A
Hap 20	Network Tw-A	Hap 6	Network Wm-A
Hap 21	Network Tw-A	Hap 7	Network Wm-A

724 **APPENDIX 8.**

725 Haplotype distribution of *Pomacea maculata* from the global mitochondrial COI dataset (Dataset

726 2)

	TW	BR	AR	UR	US	NZ	SP	CH	JP	ML	TH	VT	Total	%
Hap_1	4	-	4	1	-	-	-	-	-	-	-	-	9	3.982
Hap_2	-	-	-	1	-	-	-	-	-	-	-	-	1	0.442
Hap_3	-	-	-	1	-	-	-	-	-	-	-	-	1	0.442

Hap												Total	%
Hap_4	-	-	-	2	-	-	-	-	-	-	-	-	2	0.885
Hap_5	-	-	-	-	-	-	-	-	-	-	-	1	1	0.442
Hap_6	2	1	-	-	14	1	8	44	3	11	1	-	85	37.611
Hap_7	1	-	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_8	1	-	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_9	-	-	1	-	26	-	-	-	3	-	-	-	30	13.274
Hap_1-	-	-	1	-	21	-	-	-	-	-	-	-	22	9.735
Hap_11	-	-	-	-	1	-	-	-	-	-	-	-	1	0.442
Hap_12	-	-	-	-	4	-	-	-	-	-	-	-	4	1.770
Hap_13	-	-	1	-	-	-	-	-	-	-	-	-	1	0.442
Hap_14	-	-	1	-	-	-	-	-	-	-	-	-	1	0.442
Hap_15	-	-	3	-	-	-	-	-	-	-	-	-	3	1.327
Hap_16	-	-	1	-	-	-	-	-	-	-	-	-	1	0.442
Hap_17	-	-	1	-	-	-	-	-	-	-	-	-	1	0.442
Hap_18	-	-	1	-	-	-	-	-	-	-	-	-	1	0.442
Hap_19	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_2-	-	6	-	-	-	-	-	-	-	-	-	-	6	2.655
Hap_21	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_22	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_23	-	7	-	-	-	-	-	-	-	-	-	-	7	3.097
Hap_24	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_25	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_26	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_27	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_28	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_29	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_3-	-	1	-	2	-	-	-	-	-	-	-	-	1	0.442
Hap_31	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_32	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_33	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_34	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_35	-	4	-	-	-	-	-	-	-	-	-	-	4	1.770
Hap_36	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_37	-	4	-	-	-	-	-	-	-	-	-	-	4	1.770
Hap_38	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442
Hap_39	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_4-	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_41	-	3	-	-	-	-	-	-	-	-	-	-	3	1.327
Hap_42	-	1	-	-	-	-	-	-	-	-	-	-	1	0.442

Hap_43	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_44	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_45	-	2	-	-	-	-	-	-	-	-	-	-	2	0.885
Hap_46	-	3	-	-	-	-	-	-	-	-	-	-	3	1.327

Table 2. Haplotype distribution (Pc_Hap) of *Pomacea canaliculata* in 14 counties/cities across Taiwan.

Sample sites (Hn)	Sample ID	Shared Haplotypes	Unique Haplotypes
East (7)	HU and TT	**Total 4**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 4**, **Hap 8**	**Total 3**: Network Tw-A: **Hap 16**; Network Tw-B: **Hap 17**, **Hap 18**
Hualien (5)	HU	**Total 3**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 8**	**Total 2**: Network Tw-B; **Hap 17**, **Hap 18**
Taitung (4)	TT	**Total 3**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 4**	Network Tw-A: **Hap 16**
Middle (7)	CH, MIO, TC, YL	**Total 6**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-B: **Hap 2**, **Hap 5**, **Hap 8**; Network Tw-C: **Hap 9**	Network Tw-A: **Hap 15**
Changhua (4)	CH	**Total 3**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-C: **Hap 9**	Network Tw-A: **Hap 15**
Miaoli (2)	MIO	**Total 2**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**	-
Taichung (4)	TC	**Total 4**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-B: **Hap 2**, **Hap 5**,	-
Yunlin (2)	YL	**Total 2**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 8**	-
North (7)	HS, TP, TY, YI	**Total 3**: Network Tw-A: **Hap 3**; **Hap 11**; Network Tw-B: **Hap 2**, **Hap 5**; Network Tw-C: **Hap 9**	**Total 2**: Network Tw-B: **Hap 14**; Network Tw-A: **Hap 15**
Hsinchu (3)	HS	**Total 3**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**; Network Tw-C: **Hap 9**	-
Taipei (2)	TP	**Total 2**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 5**	-
Taoyuan (2)	TY	**Total 2**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**	
Yilan (5)	YI	**Total 3**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-B: **Hap 2**	**Total 2**: Network Tw-B: **Hap 14**; Network Tw-A: **Hap 15**
South (9)	CY, KSH, PT, TN	**Total 4**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-B: Hap 2, Hap 4, Hap 5, Hap 8; Network Tw-C: Hap 9	**Total 5**: Network Tw-A: **Hap 1**, **Hap 6**, **Hap 12**; Network Tw-B: **Hap 10**; Network Tw-C: **Hap 7**
Chiayi (6)	CY	**Total 4**: Network Tw-A: **Hap 3**, **Hap 11**; Network Tw-B: **Hap 2**, **Hap 5**	**Total 2**: Network Tw-A: **Hap 12**; Network Tw-B: **Hap 10**
Kaohsiung (4)	KSH	**Total 4**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 4**; Network Tw-C: **Hap 9**	-
Pingtung (4)	PT	**Total 4**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 5**, **Hap 8**	-

Tainan (7)	TN	**Total 4**: Network Tw-A: **Hap 3**; Network Tw-B: **Hap 2**, **Hap 4**, **Hap 5**	Total 3: Network Tw-A: **Hap 1**, **Hap 6**; Network Tw-C: **Hap 7**

Table 6. Population genetic diversity of *Pomacea maculata* from the global mitochondrial COI dataset (Dataset 2)

Country*	n	Hn	S	Hd	π	K
Argentina	16	11	36	0.925	0.011	6.375
Uruguay	5	4	3	0.900	0.002	1.4
Brazil	61	29	69	0.964	0.029	16.833
USA	66	5	11	0.705	0.006	3.649
Spain	8	1	0	0	0	0
Taiwan	8	4	11	0.75	0.008	4.928
Malaysia	11	1	0	0	0	0
Japan	6	2	9	0.6	0.009	5.4
China	44	1	0	0	0	0
China- Hong Kong*	10	2	-	0.356	0.003	1.778

Note: n = Number of sequences; Hn = Number of Haplotype, S = Number of segregating sites; Hd = Haplotype diversity; π = Nucleotide diversity; K = Average number of differences

*Countries represented with 1 sequence were not shown here.

* 10 samples added from Yang et al., 2022, as their sequence are not available publicly yet.

Table 1. Population genetic diversity of *P. canaliculata* and *P. maculata* across Taiwan. The significant value of Tajima's D and Fu's

FS are marked in bold.

Sampling sites (n)	Haplotypes			Number of segregating sites (S)	Haplotype diversity (Hd)	Nucleotide diversity (π)	Average number of differences (K)	Fu's FS	P value	Tajima's D	P value
	Total	Shared	Unique								
P. canaliculata (246)				**52**	**0.539**	**0.018**	**9.899**				
HU (29)	5	3	2	31	0.318	0.0131	7.993	**8.9429**	0.996	0.046	0.558
TT (13)	4	3	1	27	0.718	0.02198	13.410	**9.0785**	0.999	**2.361**	1
CH (15)	4	3	1	26	0.371	0.00568	3.467	3.0052	0.916	**-2.362**	0
MIO (13)	2	2	0	26	0.513	0.02186	13.333	**15.674**	1	**2.574**	1
TC (14)	4	4	0	32	0.703	0.02012	12.275	**9.1311**	0.997	0.946	0.873
YL (13)	2	2	0	25	0.538	0.02207	13.460	**15.754**	1	**2.913**	1
HS (14)	3	3	0	36	0.275	0.01122	6.846	**7.977**	0.997	**-1.711**	0.024
TP (12)	2	2	0	28	0.167	0.00765	4.667	**8.009**	0.995	**-2.227**	0.001
TY (11)	2	2	0	26	0.436	0.0186	11.345	**12.987**	1	1.280	0.93
YI (12)	5	3	2	33	0.667	0.02228	13.590	**6.171**	0.994	1.101	0.911
CY (23)	6	4	2	33	0.613	0.01832	11.178	**8.137**	0.989	0.957	0.879
KSH (19)	4	4	0	37	0.614	0.02359	14.392	**13.236**	1	1.443	0.955
PT (33)	4	4	0	31	0.676	0.02218	13.530	**18.395**	1	**2.748**	1
TN (25)	7	4	3	42	0.580	0.01915	11.680	**7.375**	0.995	0.192	0.608
P. maculata (8)											
PT, KSH, TN, CY, MIO, YI	4			10	0.643	0.007	4.500			0.826	

Table 3. Haplotype distribution (Pm_Hap) of *Pomacea maculata* in 14 counties/cities across Taiwan.

Sample sites (Hn)	Sample ID	Shared Haplotypes	Unique Haplotypes
Middle (1)	MIO	**Hap 19**	-
Miaoli (1)	MIO	**Hap 19**	-
North (1)	YI	**Hap 19**	-
Yilan (1)	YI	**Hap 19**	-
South (4)	CY, KSH, PT, TN	**Hap 19**	Total 3: **Hap 20**, **Hap 21**, **Hap 22**
Chiayi (1)	CY	-	**Hap 22**
Kaohsiung (3)	KSH	**Hap 19**	
Pingtung (1)	PT	-	**Hap 21**
Tainan (1)	TN	-	**Hap 20**

Table 2. Genetic differentiation coefficient (F_{st}) below diagonal and gene flow (N_m) above diagonal based on mitochondrial COI sequences of *P. canaliculata* from 14 counties or cities of Taiwan. Abbreviations: HU: Hualien, and TT: Taitung from the East, YL: Yunlin, CH: Changhua, TC: Taichung, and MIO: Miaoli from the Middle, HS: Hsinchu, TY: Taoyuan, TP: Taipei, and YI: Yilan from the North, and PT: Pingtung, KSH: Kaohsiung, TN: Tainan, and CY: Chiayi from the South. The higher F_{st} value ($F_{st} > 0.2$) resulted in

	TN	PT	KSH	CY	YI	TY	TP	HS	YL	TC	MIO	CH	TT	HU
TN		4.040	NA	NA	NA	NA	13.27	73.904	4.788	NA	NA	4.904	2.13	NA
PT	0.110		261.28	3.906	13.778	6.224	1.148	**1.368**	NA	5.85	NA	**0.858**	NA	**1.752**
KSH	-0.007	0.002		NA	NA	NA	2.584	3.83	NA	NA	NA	**1.692**	12.49	5.622
CY	-0.038	0.113	0.000		NA	NA	13.724	35.758	4.478	NA	NA	4.464	2.026	NA
YI	-0.046	0.035	-0.051	-0.050		NA	4.816	8.668	36.4	NA	NA	2.332	4.618	42.198
TY	-0.065	0.074	-0.039	-0.067	-0.083		9.828	60.032	7.904	NA	NA	3.286	2.754	NA
TP	0.036	**0.304**	0.162	0.035	0.094	0.048		NA	**0.998**	9.844	2.51	NA	**0.63**	NA
HS	0.007	**0.268**	0.115	0.014	0.055	0.008	-0.053		**1.272**	21.712	3.744	NA	**0.782**	NA
YL	0.095	-0.049	-0.015	0.100	0.014	0.060	**0.334**	**0.282**		7.608	NA	**0.694**	NA	**1.71**
TC	-0.050	0.079	-0.026	-0.058	-0.070	-0.080	0.048	0.023	0.062		NA	3.434	2.714	NA
MIO	-0.022	-0.006	-0.064	-0.021	-0.072	-0.061	0.166	0.118	-0.022	-0.046		**1.488**	12.258	7.12
CH	0.093	**0.368**	**0.228**	0.101	0.177	0.132	-0.038	-0.041	**0.419**	0.127	**0.251**		**0.464**	13.7
TT	0.190	-0.030	0.038	0.198	0.098	0.154	**0.443**	**0.390**	-0.032	0.156	0.039	**0.519**		**0.974**
HU	-0.011	**0.222**	0.082	-0.014	0.012	-0.031	-0.017	-0.029	**0.226**	-0.011	0.066	0.035	**0.339**	

less gene flow (Nm <1) between the counties or cities (marked in bold).

Table 5. Population genetic diversity of *Pomacea canaliculata* from the global mitochondrial COI dataset (Dataset 3)

Country*	n	Hn	S	Hd	π	K
Argentina	32	19	49	0.915	0.023	12.599
Uruguay	8	6	35	0.928	0.027	15.107
USA	19	2	1	0.199	0.0003	0.199
Chile	4	1	0	0	0	0
China	582	12	57	0.564	0.0223	12.315
Taiwan	249	16	51	0.539	0.0179	9.868
Indonesia	3	2	1	0.667	0.0012	0.667
Japan	18	6	41	0.797	0.0277	15.274
Malaysia	14	3	22	0.385	0.0105	5.780
Philippines	12	12	39	1	0.029	16.045
Kenya	3	1	0	0	0	0
Thailand	2	2	1	1	0.002	1
China-Hongkong**	2+ 139	5	-	0.441	0.018	8.91

Note: n = Number of sequences; Hn = Number of Haplotype, S = Number of segregating sites; Hd = Haplotype diversity; π = Nucleotide diversity; K = Average number of differences

*Countries represented with 1 sequence were not shown here.

* 139 samples added from Yang et al., 2022, as their sequence are not available publicly yet.