MagIC-Cryo-EM: Structural determination on magnetic beads for scarce macromolecules in heterogeneous samples

Yasuhiro Arimura[1,2*], Hide A. Konishi[1], Hironori Funabiki[1*]

[1] Laboratory of Chromosome and Cell Biology, The Rockefeller University, New York, NY 10065

[2] Current address: Basic Sciences Division, Fred Hutchinson Cancer Center, Seattle, WA, USA, 98109-1024

[*] Correspondence: funabih@rockefeller.edu or yarimura@fredhutch.org

Summary

Cryo-EM single-particle analyses typically require target macromolecule concentration at 0.05~5.0 mg/ml, which is often difficult to achieve. Here, we devise Magnetic Isolation and Concentration (MagIC)-cryo-EM, a technique enabling direct structural analysis of targets captured on magnetic beads, thereby reducing the targets' concentration requirement to < 0.0005 mg/ml. Adapting MagIC-cryo-EM to a Chromatin Immunoprecipitation protocol, we characterized structural variations of the linker histone H1.8-associated nucleosomes that were isolated from interphase and metaphase chromosomes in *Xenopus* egg extract. Combining Duplicated Selection To Exclude Rubbish particles (DuSTER), a particle curation method that excludes low signal-to-noise ratio particles, we also resolved the 3D cryo-EM structures of nucleoplasmin NPM2 co-isolated with the linker histone H1.8 and revealed distinct open and closed structural variants. Our study demonstrates the utility of MagIC-cryo-EM for structural analysis of scarce macromolecules in heterogeneous samples and provides structural insights into the cell cycle-regulation of H1.8 association to nucleosomes.

Introduction

Recent advances in cryogenic electron microscopy (cryo-EM) technology have enabled the structural characterization of biomolecules isolated from their native conditions[1]. However, the necessity for high sample concentration restricts its applicability to abundant targets[2–5]. The vitrification step of cryo-EM is a major contributor to this bottleneck. In a conventional plunge vitrification method, 3 µL of aqueous samples greater than 1 mg/mL are typically required to acquire sufficient numbers of particle images on cryo-EM micrographs for 3D structure reconstruction (Table S1)[6]. This is because most of the target complexes in the sample solution applied on a cryo-EM grid

must be removed by a blotting paper to make a thin ice layer suitable for analysis.
Several methods are currently available to lower sample volume/concentration needed
(Table S1). Jet vitrification[7] and Spotiton[8] require sub-nanoliters of the sample volume,
but they still require high-concentration samples. Affinity grids, such as Ni-NTA lipid
monolayer grids[9], chemically functionalized grids[10], antibody-attached grids[11], and
streptavidin monolayer grids[12], are amenable for lower concentration samples (~0.05
mg/mL), but concentrating natively isolated targets to such a level and reproducibly
generating the affinity grids remains challenging.

Structural characterization of native chromatin-associated protein complexes is
particularly challenging due to their heterogeneity and scarcity: more than 300 proteins
directly bind to the histone core surface[13], while each of these proteins is targeted to
only a fraction of nucleosomes in chromatin. For their structural analysis, it is a common
practice to assemble nucleoprotein complexes using purified recombinant proteins and
a specific short (10 – 1000 bp) linear DNA. However, this reconstitution approach has a
limitation since the structure and function of chromatin proteins can be altered by
several variances under native conditions, such as DNA sequence, DNA and protein
modifications, and short- and long-scale DNA folding. Although isolation of the
endogenous chromatin-associated complexes can be achieved through chromatin
immunoprecipitation (ChIP) [14–16] to determine the associated DNA sequences and
proteins[17,18], the amount obtained by this method is too little to apply for conventional
structural analysis.

To obtain high-resolution cryo-EM structures of chromatin-associated protein
complexes while they are functioning on the native chromosomes, we previously
analyzed structural variation of nucleosomes isolated from interphase and metaphase
chromosomes formed in *Xenopus laevis* egg extracts [3]. We found that the averaged
structures of the nucleosome core particle (NCP) in interphase and metaphase
chromosomes are essentially identical to the NCP crystal structure assembled with
histone proteins and DNA with strong nucleosome positioning sequences [19,20]. We also
observed that the major structural variation of the nucleosome structures between
interphase and metaphase chromosomes was attributable to the binding status of the
oocyte-specific linker histone H1.8. We were able to resolve the 3D structure of the
H1.8-bound nucleosome isolated from metaphase chromosomes but not from
interphase chromosomes[3]. The resolved structure indicated that H1.8 in metaphase is
most stably bound to the nucleosome at the on-dyad position, in which H1 interacts with
both the entry and exit linker DNAs [21–24]. This stable H1 association to the nucleosome
in metaphase likely reflects its role in controlling the size and the shape of mitotic
chromosomes through limiting chromatin accessibility of condensins [25], but it remains
unclear why H1.8 binding to the nucleosome in interphase is less stable. Since the low
abundance of H1.8-bound nucleosomes in interphase might have prevented us from

determining their structure, we sought to solve this issue by enriching H1.8-bound
nucleoprotein complexes through adapting ChIP-based methods.

Aiming to reduce sample requirements for single particle cryo-EM analyses to
levels lower than those widely used for ChIP-seq (10-50 ng DNA, Table S1)[17], here we
developed Magnetic Isolation and Concentration (MagIC)-cryo-EM, which enables direct
cryo-EM analysis of target molecules enriched on superparamagnetic nanobeads. By
adapting the ChIP protocol to MagIC-cryo-EM, we successfully determine the ~4 Å
resolution structures of H1.8-GFP-bound nucleosomes using highly heterogeneous
dilute fractions isolated from metaphase and interphase chromosomes. In addition, by
combining the particle curation method, Duplicated Selection To Exclude Rubbish
particles (DuSTER), which effectively removes particles with a low signal-to-noise ratio
(S/N), we revealed structural variations of the H1.8-bound chaperone NPM2 isolated
from interphase chromosomes, providing structural insights into the cell cycle regulation
of H1.8 stabilization on nucleosomes.

Results

<u>Development and optimization of MagIC-cryo-EM using nucleosomes</u>

Inspired by a report using 200-300 nm superparamagnetic beads directly loaded onto a
cryo-EM grid to image viral particles[26], we examined the feasibility of 50 nm streptavidin
nanobeads for cryo-EM single-particle analysis using poly-nucleosome arrays as pilot
targets (Figure 1A). Nanobeads were easily identified on the grid as black dots in the
intermediate-magnification montage map (Figure 1B), facilitating target identification for
subsequent high-magnification data collection. In the high-magnification micrographs,
poly-nucleosome fibers were observed around the nanobeads as expected (Figure 1C).
Using nucleosome-like particles selected from 550 micrographs by the machine-
learning-based software Topaz[27], we successfully determined the 3D structure of the
nucleosome at sub-nanometer resolution (Figure 1D). This result, however, revealed a
notable issue; an intense halo-like scattering covered a ~30 nm radius around the
nanobeads (Figure 1D, blue areas), interfering with the signal from particles that were
proximal to the beads.

To reduce the effect of the halo-like scattering surrounding the nanobeads, a
protein spacer module was attached to the beads so that the target biomolecules are
placed outside the reach of the halo (Figure 2A and 2B). After several rounds of
optimization using the *in vitro* reconstituted H1.8-bound nucleosome as a model target,
we chose a spacer module comprising an 11-nm triple helical bundle (3HB) protein[28]
and four copies of a 60-nm single alpha helix (SAH) protein[29] for its effectiveness and
reasonable production yield (Figure 2B, Figure S1). The distal end of the spacer module

was engineered to allow for exchangeable target-capturing modules by SPYcatcher-
SPYtag conjugation (Figure 2B)[30]. We hereon refer to these magnetic nanoparticles
coated with the spacer and target-capturing modules as MagIC-cryo-EM beads.

To assess the feasibility of the MagIC-cryo-EM beads for structural analysis of a
low-concentration target in heterogeneous samples, we isolated H1.8-GFP-bound
nucleosomes by anti-GFP nanobody coupled to the MagIC-cryo-EM beads from a
mixture of H1.8-GFP nucleosomes (1.7 nM, or 0.00047 mg/mL) and a large excess of
unbound mono-nucleosomes (53 nM, or 0.012 mg/mL) (Figure 2C and 2D). This target
concentration was approximately 100 to 1000 times lower than the concentration
required for conventional cryo-EM methods, including affinity grid approaches [9–11]. The
magnetic beads were captured on a cryo-EM grid by neodymium magnets for 5 min in a
humidified chamber (Figure 2E). This magnetic capture step significantly increased the
number of beads that were found in the sample holes of the grid (Figure 2F-I), thereby
mitigating the sample loss caused by filter paper blotting to generate a thin ice layer.

High-magnification micrographs of MagIC-cryo-EM beads show that the spacer
module successfully placed nucleosome-like particles outside the halo-like scattering
surrounding the nanobeads (Figure 2J). The local enrichment of target molecules
around MagIC-cryo-EM beads offers a substantial advantage in data collection
efficiency over available cryo-EM methods[9–11], in which target molecules are
disseminated across the grids and are difficult to identify. In contrast, the magnetic
beads are easily identified in the Medium-Magnification Montage (MMM) map (Figure
2G), enabling the selection of target-rich areas prior to high-magnification data
collection. Indeed, approximately 100 H1.8-GFP nucleosome particle images per bead
were efficiently collected even with a sample concentration as low as 0.00047 mg/mL of
H1.8-GFP nucleosomes in the heterogeneous sample (Figure 2J right panel).

After removing junk particles using decoy classification[3,31–33] (Figure S2), an H1.8
density-containing nucleosome class was isolated via *ab initio* reconstruction and
heterogeneous refinement using cryoSPARC[34]. Among the nucleosome-containing
particles, 55.7 % of them were classified as a nucleosome with H1.8 at the on-dyad
position (Figure S2), yielding a final 3D structure at 3.6 Å resolution (Figure 2K). This
high fraction of H1.8-bound nucleosome particles indicated that the MagIC-cryo-EM
beads efficiently isolated the target molecules. Notably, this method only required 5 ng
of H1.8-GFP-bound nucleosomes (including 2 ng of DNA) per cryo-EM grid, which is
comparable to or even lower than the requirements of widely used ChIP-seq[17].

<u>MagIC-cryo-EM application for ChIP to assess structural features of H1.8 in</u>
<u>chromosomes</u>

We next adapted MagIC cryo-EM to ChIP protocols to elucidate the cell-cycle-specific
mechanism that controls H1.8 stability on interphase and metaphase nucleosomes. We
previously reported the cryo-EM structure of *Xenopus* H1.8 bound to the metaphase
nucleosome at the on-dyad position, whereas no H1.8-containing structures were
reconstructed from interphase chromosomes[3] (Figure 3A, left). Despite the high
accumulation of H1.8 in the nucleus[35] (Figure 3B), the amount of nucleosome-
associated H1.8 in interphase is reduced to approximately 30% of that in metaphase[3].
Given the high mobility of the linker histone H1 on chromatin[25,36,37], we hypothesized
that H1.8 on nucleosome is destabilized by an interphase-specific mechanism. By
enriching H1.8-bound nucleosomes from interphase and metaphase chromosomes
using MagIC-cryo-EM, we intended to examine if H1.8 in interphase preferentially
associates with nucleosomes at more unstable binding positions, such as at off-dyad
positions[38,39] (Figure 3A, positioning model), or if there is an interphase-specific
mechanism (by chaperones, for example) that dissociates H1.8 from nucleosomes
(Figure 3A, chaperone model), although the amount H1.8-bound NAP1, the known
histone H1.8 chaperone [40], did not differ between metaphase and interphase egg
extracts (Fig. S3A).

To distinguish between these models, we applied MagIC-cryo-EM to enrich H1.8
bound nucleosomes from chromosomes assembled in interphase and metaphase
Xenopus egg extracts. Sperm nuclei were incubated in egg extracts supplemented with
H1.8-GFP to obtain replicated interphase chromosomes and metaphase chromosomes,
which were crosslinked and fragmented to generate soluble nucleoprotein complexes
(Figure 3B). We confirmed that H1.8-GFP is functional as it rescued the chromosome
elongation phenotype caused by H1.8 immunodepletion[25,35] (Figure S3B-D). Sucrose
density gradient centrifugation was conducted to separate different H1.8-containing
complexes, including mono-nucleosome fractions and oligo-nucleosome fractions, as
previously described[3] (Figure 3C and S4). As we had predicted that more H1.8 proteins
would associate with nucleosomes in metaphase than in interphase[3], we increased the
quantities of egg extract and sperm nuclei by 2.5 fold to prepare comparable amounts of
H1.8-bound interphase nucleosomes as compared to metaphase (Figure 3C, fractions
4-11). To prevent the dissociation of H1.8 from nucleosomes during DNA fragmentation,
the MNase concentration and the reaction time were optimized to generate DNA
fragment lengths with 180–200 bp (Fig. S4B), which is adequate for linker histone
association[22]. To ensure that most nucleosomes isolated through MagIC-cryo-EM were
bound by H1.8, we selected the fractions enriched with H1.8-bound mono-nucleosomes
(fraction 5 in Figure 3C and 3D), as oligo-nucleosomes (abundant in fractions 6-11)
might include H1.8-free nucleosomes. These fractions contain highly heterogeneous
protein mixtures (Figure 3E), in which H1.8-GFP is a minor constituent with an
estimated concentration at 1-2 nM (corresponding to 0.00025-0.0005 mg/ml of H1.8-
bound mono-nucleosomes) (Figure S4C). Mass spectrometry analysis of these fractions

also showed heterogeneity as they included several DNA-binding proteins, such as
PCNA (Table S2 and Table S5).

H1.8-GFP-bound mono-nucleosomes in fraction 5 (from metaphase and
interphase chromosomes) were captured by GFP nanobody-MagIC-cryo-EM beads and
applied to grids for cryo-EM analysis. Mass spectrometry analysis of the captured
MagIC-cryo-EM beads confirmed selective enrichment of H1.8 over other nonhistone
proteins found in fraction 5 (Table S2). To quantitatively assess the population of the H1-
bound structural modes of interphase and metaphase nucleosomes, we employed *in*
silico mixing 3D classification[3,41]. Micrographs of interphase and metaphase MagIC-
cryo-EM were mixed and used for particle picking and decoy classification to isolate the
nucleosome-containing classes (Figure S5). Subsequently, particles were classified into
three nucleosome-containing 3D models (A, B, C), which were generated by *ab initio*
reconstruction (Figure 3F and S5A). Further 3D classification on the class A, which has
weak H1.8 density, yielded three new nucleosome-containing structures, A1, A2, and A3
(Figure 3F and S5A). Then, the populations of interphase and metaphase particles in
each class were assessed (Figure 3F). Only class A1 had an apparent H1.8 density at
the on-dyad position of the nucleosome, with 27% and 23% of the nucleosome particles
assigned to this class coming from interphase and metaphase conditions, respectively.
Although class A2 had linker DNA densities on both sides of the entry/exit sites of the
nucleosome in a closed conformation, it did not have a clear H1.8 density. This
suggested that the structures of H1.8 in the particles assigned to this class were not
uniform, and that the H1.8 density was averaged out during the cryo-EM processing.
Class A3, to which 3-4 % of the nucleosome particles were assigned, had ambiguous
extra densities outside of the on-dyad position (Figure 3F, red arrows), possibly
representing H1.8 bound to non-dyad positions. Overall, the relative distributions of
these 5 classes were largely similar between interphase and metaphase (Figure 3F),
and the structures of H1.8-bound nucleosomes in interphase and metaphase were
indistinguishable (Figure 3G). The structures of GFP-tagged H1.8-bound nucleosomes
isolated from *Xenopus* egg extract chromosomes are essentially identical to the
endogenous H1.8-bound nucleosome structure we previously determined [3]. Therefore,
although the usage of GFP-tagged H1.8 and MagIC-cryo-EM potentially affect the
structure of the H1.8-bound nucleosome, we consider these influences to be minimal.
Altogether, the results suggest that differential positional preferences of H1.8 on the
nucleosome (Figure 3A, positioning model) are unlikely to drive the reduced H1.8
association to interphase nucleosomes.

MagIC-cryo-EM and DuSTER reconstructed cryo-EM structure of interphase-specific
H1.8-containing complex, NPM2

Although we could not discern structural differences of H1.8-bound mono-nucleosomes
from metaphase and interphase samples, we noticed that substantial portions of H1.8
were enriched in sucrose fractions 3 and 4 isolated from interphase chromosomes but
not from metaphase chromosomes (Figure 3C). As these interphase-specific H1.8
fractions were lighter than mono-nucleosome-containing fractions, we thought that they
may contain regulatory proteins that preferentially dissociate H1.8 from nucleosomes in
interphase, in line with the chaperone model (Figure 3A).

To characterize these interphase-specific fractions, we sought to determine their
structural features using MagIC-cryo-EM. However, our initial attempt failed to
reconstitute any reasonable 2D classes of the interphase-specific H1.8-containing
complex (Figure S6A), even though Topaz successfully picked most of the 60~80 Å
particles that are visible on motion-corrected micrographs and enriched around the
MagIC-cryo-EM beads (Figure S6A). This was likely due to their small size; most of the
particles did not have a high enough S/N to be properly classified during the 2D
classifications as they were masked by background noise from the ice and/or spacer
proteins (Figure S6B).

To solve this issue, we devised the particle curation method DuSTER that does
not requires the successful 2D classifications (Figure 4A). The principle of DuSTER is
based on our realization that low S/N ratio particles were not reproducibly recentered
during 2D classification (Figure S7). On the particles that were successfully recognized
during 2D classification, picked points were shifted to the center of the particles (Figure
4A, black arrows). However, on the low S/N ratio particles that could not be recognized
during 2D classification, picked points were shifted outside the center of the particles
(Figure 4A, green arrows). To assess the reproducibility of the particle recentering
during 2D classification, two independent particle pickings were conducted by Topaz so
that each particle on the grid has up to two picked points (Figure 4A, second left panel).
Some particles that only have one picked point will be removed in a later step. These
picked points were independently subjected to 2D classification. After recentering the
picked points by 2D classification, distances (D) between recentered points from the
first picking process and other recentered points from the second picking process were
measured. DuSTER keeps recentered points whose D are shorter than a threshold
distance (D_{TH}). By setting D_{TH} = 20 Å, 2D classification results were dramatically
improved in this sample; a five-petal flower-shaped 2D class was reconstructed (Figure
4B). This step also removes the particles that only have one picked point. Although
approaches to utilize the reproducibility of 2D class assignments have been proposed[42],
the advantage of DuSTER is that it can be applied to small particles that cannot even be
properly classified in 2D classification.

Repetitive rounds of particle curation using the picked point locations recentered
by 2D classification (referred to as 2D DuSTER) successfully reconstituted 2D classes

of 60~80 Å complexes (Figure 4B, and S8). As expected, the particles rejected by
DuSTER have a generally weak contrast (Fig S9A). Although higher contrast images
can be generated by increasing the defocus (the distance between the target particles
and the lens focus), the selected particles were evenly distributed in all defocus ranges
between 1.5 ~ 3.5 μm (Fig S9B), demonstrating that DuSTER did not merely select any
random high contrast particles. By selecting these 2D classes, an initial 3D model was
built (Figure S8, and S10). Using this 3D model, particle curation was revised with 3D
DuSTER. In the 3D DuSTER, three 3D maps were used as the initial models for the
cryoSPARC heterogenous refinement to centering the particles accurately (D_{TH} = 15 Å)
(Figure S10A). 3D DuSTER enabled the reconstruction of 3D structure of the
interphase-specific H1.8-containing complex, a pentameric macromolecule with a
diameter of approximately 60 Å (Figure 4C and S12).

To determine the identity of this complex, MagIC-cryo-EM beads used for
isolating the complex were analyzed by mass spectrometry (MS) (Figure 4D). Among
the proteins detected by MS, NPM2 aligned well with the MagIC-cryo-EM result.
Western blotting confirmed that NPM2 was preferentially enriched in interphase
chromatin fractions compared to metaphase (Figure 4E), while NPM2 interacts with
H1.8 in chromosome-free egg extracts both in interphase and metaphase (Fig. S3A).
The native PAGE of the chromatin fractions indicated that NPM2 forms various
complexes, including NPM2-H1.8, on the interphase chromatin fractions (Fig. S4D). In
addition, the crystal structure and AlphaFold2 (AF2)-predicted models of *Xenopus*
NPM2 matched the MagIC-cryo-EM structure of the interphase-specific H1.8-bound
complex (Figure 4F)[43].

<u>Structural variations of NPM2 bound to H1.8</u>

In *Xenopus* eggs, NPM2 replaces sperm protamines with core histones upon
fertilization, thereby promoting nucleosome assembly on sperm DNA[44–46]. NPM2 can
also extract out somatic linker histones from chromatin [47–49]. X-ray crystallography
suggested that recombinant *Xenopus* NPM2 forms a pentamer and a decamer (a dimer
of pentamers)[43]. The acidic tracts in the C-terminal tail of NPM2 binds H2A-H2B,
histone octamers, and the linker histone H5[50–52], while poly-glutamylation and
hyperphosphorylation of NPM2 promote its substrate sequestration[53,54]. In addition,
NPM1 (nucleophosmin), a paralog of NPM2, interacts with H1[49,55]. However, no
subnanometer-resolution structure of NPM2 or NPM1 with post-translational
modifications or with substrates is currently available.

By further analyzing our cryo-EM structure representing the H1.8-bound state of
NPM2, we identified two structural variants, classified as open and closed forms (Figure
5A, S11, and S12J-K). Due to its structural similarity to a flower, we call the highly acidic

putative substrate-binding surface the petal side, whereas the other more charge
neutral surface the sepal side (Figure 5A and S13). The major structural differences
between the two forms are found at C-terminal and N-terminal segments of NPM2 core
and at the A1 loop (Figure 5A, 6B, and S13). In the closed form, β8 runs straight from
the sepal to the petal sides of each pentamer and has an extended C-terminal segment
that protrudes past the petal side of the pentamer. In the open form, however, the C-
terminal portion of β8 is bent outward to the rim (Figure 5A). Along with this β8 bending,
C-terminal segment, N-terminal segment, and A1 loop are also positioned outward in
the open form. The configuration of β1, β8, and A1 loop in the crystal structure of
Xenopus NPM2 [43], the AF2-predicted structure of *Xenopus* NPM2 [56–58], and the cryo-
EM structure of the bacterially expressed human NPM1[59], which were all determined in
the absence of their target proteins, is similar to the closed form (Figure S13B-D).
Notably, extra cryo-EM densities, which may represent H1.8, are clearly observed in the
open form but much less in the closed form near the acidic surface regions proximal to
the N terminus of β1 and the C terminus of β8 (Figure 5A and 5B). Supporting this idea,
the acidic tract A1 (aa 36-40) and A2 (aa 120-140), which are both implicated in the
recognition of basic substrates such as core histones [43,50], respectively interact with and
are adjacent to the putative H1.8 density (Figure 5B). In addition, the NPM2 surface that
is in direct contact with the putative H1.8 density is accessible in the open form while it
is internalized in the closed form (Figure 5C). This structural change of NPM2 may
support more rigid binding of H1.8 to the open NPM2, whereas H1.8 binding to the
closed form is less stable and likely occurs through interactions with the C-terminal A2
and A3 tracts, which are not visible in our cryo-EM structures.

In the aforementioned NPM2-H1.8 structures, for which we applied C5 symmetry
during the 3D structure reconstruction, only a partial H1.8 density could be seen (Figure
5B). One possibility is that the H1.8 structure in NPM2-H1.8 does not follow C5
symmetry. As the size of the NPM2-H1.8 complex estimated from sucrose gradient
elution volume is consistent with pentameric NPM2 binding to a single H1.8 (Figure 3C
and Table S3), applying C5 symmetry during structural reconstruction likely blurred the
density of the monomeric H1.8 that binds to the NPM2 pentamer. The structural
determination of NPM2-H1.8 without applying C5 symmetry lowered the overall
resolution but visualized multiple structural variants of the NPM2 protomer with different
degrees of openness co-existing within an NPM2-H1.8 complex (Figure S14), raising a
possibility that opening of a portion of the NPM2 pentamer may affect modes of H1.8
binding. Although more detailed structural analyses of the NPM2-substrate complex are
the subject of future studies, MagIC-cryo-EM and DuSTER revealed structural changes
of NPM2 that was co-isolated H1.8 on interphase chromosomes.

Discussion

MagIC-cryo-EM offers sub-nanometer resolution structural determination using a heterogeneous sample that contains the target molecule at 1~2 nM, which is approximately 100 to 1000 times lower than the concentration required for conventional cryo-EM methods, including affinity grid approach [9–11]. This significant improvement was achieved through the four unique benefits of MagIC-cryo-EM (Figure 6). First, the on-bead-cryo-EM approach minimizes preparation steps, which can lead to sample loss, such as target isolation, enrichment, and buffer exchange (Figure 6A). Second, sample loss during the grid-freezing process is reduced by magnet-based enrichment of the targets on cryo-EM grids (Figures 2E-2I and 6B). Third, magnetic beads are easily identifiable on the grid (Figures 2G and 6C). Fourth, the target molecules are accumulated around magnetic beads, ensuring that each micrograph contains more than 100 usable particles independent of input sample concentration (Figure 2J and 6D). Adapting the ChIP-based method to MagIC cryo-EM, we successfully isolated and reconstructed the H1.8-bound nucleosome and the H1.8-bound NPM2 structures from interphase chromosomes, which have never been accomplished before.

To reconstitute the structure of H1.8-bound NPM2, we needed to devise the particle curation method DuSTER, which greatly helped the structural reconstitution of small particles with low S/N (Figure 4). By combining MagIC-cryo-EM and DuSTER, we were able to determine the sub-nanometer structure and structural variations of the NPM2-H1.8-GFP complex, in which the mass of the ordered region is only 60 kDa. Notably, particle curation by DuSTER does not require human supervision or machine learning, except for determining the distance threshold between repeatedly picked particles. This feature may allow for automating particle curation via DuSTER in the future.

MagIC-cryo-EM and DuSTER approaches hold the potential for targeting a wide range of biomolecules, including small ones, for two main reasons. First, the target-capturing module could be replaced with various other proteins, such as different nanobodies, single-chain variable fragments (scFv), protein A, dCas9, or avidin, to capture a wide range of biomolecules. Second, the sample requirement for MagIC-cryo-EM is a mere 5 ng per grid, which is comparable to or even lower than the requirements of widely used ChIP-seq [17]. Coupling next-generation sequencing with MagIC-cryo-EM beads would help the field determine structural features of functionally distinct chromatin regions, such as heterochromatin, euchromatin, transcription start sites, telomeres, and centromeres. The low sample requirement of MagIC-cryo-EM also opens the door to structural analysis using limited specimens, including patient tissues.

Combining MS, MagIC-cryo-EM and DuSTER, we found that the majority of chromatin-bound H1.8 in interphase existed as a complex with NPM2 rather than with

nucleosomes (Figure 5C and 5D). This contrasts to the reports suggesting that NAP1 is
the major H1.8-bound chaperone in *Xenopus* egg extracts [60,61], while it is consistent
with our previous MS analysis that also detected NPM2, but not NAP1, in fractions
enriched with nucleosomes in interphase [3]. Our observation is also in line with a
previous report that NPM2 is able to remove linker histones but not core histones from
somatic nuclei that are introduced to *Xenopus* egg extracts [47]. Since the amounts of
H1.8-associated NAP1 or NPM2 in the egg cytoplasm did not change between
interphase and metaphase (Figure S3A), a mechanism must exist such that NPM2
interacts with H1.8 on chromatin specifically in interphase and suppresses H1.8-
nucleosome interaction (Figure 5D). Two basic patches at the C-terminal tail of NPM2
may contribute to cell cycle-dependent DNA binding as they are flanked with potential
Cdk1 phosphorylation sites. NPM2 may maintain nucleosome-bound H1.8 at a low level
in interphase during early developmental cell cycles to support rapid DNA replication,
while mitotic induction of H1.8 association with nucleosomes tunes condensin loading
on chromosomes and ensures proper chromosome size to facilitate chromosome
segregation [25] (Figure 5D).

395   Structural studies based on *in vitro* reconstitution previously suggested that
NPM2 binds to its substrate as a homo-decamer [43,50], or a homo-pentamer [51,52]. Our
cryo-EM structure strongly suggests that the NPM2 binds to H1.8 as a homo-pentamer.
Structure variation analyses suggest that NPM2 subunits can exhibit two structural
configurations, open and closed forms, of which H1.8 is stably associated with only the
open form. Since the closed form is more similar to the reported crystal structure and
AF2-predicted structures (Figure S14B-D), both of which are determined in the absence
of the substrates, our analysis points toward a possibility that substrate binding induces
the structural transition of NPM2 to the open form. The conformational changes of the
NPM family have been proposed in other studies, such as NMR and negative stain-EM
[54,62,63]. Our cryo-EM structures of NPM2 indicate the potential mechanisms of NPM2
conformational changes and potential substrate binding sites. Among NPM2 acidic
tracts A1, A2 and A3, which are important for substrate recognition, our atomic models
visualize A1 and the edge of A2 at the petal side of the structure, where the density
corresponding to the predicted H1.8 can be found (Figure 5B). As the A2 and A3 belong
to the disordered C-terminal tail that extends from the petal side of the NPM2 complex,
our data suggest that the open form provides a stable association platform by exposing
the acidic surface at the petal side for the substrate recognition, while the C-terminal A2
and A3 at the flexible tail may facilitate recruitment and possibly also entrapment of the
substrate. Since our structural analysis further suggests that each NPM2 subunit may
independently adapt open and closed form within a pentamer, this flexibility in the core
domain may enable the association of substrates with diverse sizes and structures to
support its molecular chaperone functionality.

<u>Limitations of the study</u>

While MagIC-cryo-EM is envisioned as a versatile approach suitable for various

biomolecules from diverse sources, including cultured cells and tissues, it has thus far

been tested only with H1.8-bound nucleosome and H1.8-bound NPM2, both using anti-

GFP nanobodies to isolate GFP-tagged H1.8 from chromosomes assembled in

Xenopus egg extracts after pre-fractionation of chromatin. To apply MagIC-cryo-EM for

the other targets, the following factors must be considered: *1) Pre-fractionation.* This

step (e.g., density gradient or gel filtration) may be necessary to enrich the target

protein in a specific complex from other diverse forms (such as monomeric forms,

subcomplexes, and protein aggregates). *2) Avoiding bead aggregation.* Beads may be

clustered by targets (if the target complex contains multiple affinity tags or is

aggregated), nonspecific binders, and target capture modules. To directly apply

antibodies that recognize the native targets and specific modifications, optimization to

avoid bead aggregation will be important. *3) Stabilizing complexes.* The target

complexes must be stable during the sample preparation. Crosslink was necessary for

the H1.8-GFP-bound nucleosome. *4) Loading the optimum number of targets on the*

bead. The optimal number of particles per bead differs depending on target sizes, as

larger targets are more likely to overlap. For H1.8-GFP-bound nucleosomes, 500 to

2,000 particles per bead were optimal. We expect that fewer particles should be coated

for larger targets.

Regarding the cryo-EM data acquisition, the selection of data collection points is

currently performed through the manual picking of magnetic beads on the MMM map.

This method does not support image-shift-based data collection and serves as a

bottleneck for data collection speed, limiting throughput to approximately 500–1000

micrographs per day. The development of machine learning-based software to

automatically identify magnetic beads on MMM maps and establish parameters for

image-shift-based multiple shots could substantially enhance data collection efficiency.

The efficiency of magnetic bead capture can be further improved. In the current

MagIC-cryo-EM workflow, the cryo-EM grid is incubated on a magnet before being

transferred to the Vitrobot for vitrification. However, since the Vitrobot cannot

accommodate a strong magnet, the vitrification step occurs without the magnetic force,

potentially resulting in bead loss. This limitation could be addressed by developing a

new plunge freezer capable of maintaining magnetic force during vitrification.

While DuSTER enables the structural analysis of NPM2 co-isolated with H1.8-

GFP, the resulting map quality is modest, and the reported numerical resolution may be

overestimated. Furthermore, only partial density for H1.8 is observed. Although

structural analysis of small proteins is inherently challenging, it is possible that halo-like

scattering further hinders high-resolution structural determination by reducing the S/N
ratio. More detailed structural analyses of the NPM2-substrate complex will be
addressed in future studies.

<u>Acknowledgments</u>

This research was supported by a National Institutes of Health Grants (R35GM132111)
to HF, Japan Society for the Promotion of Science Overseas Research Fellowships to
HAK, and Osamu Hayaishi Memorial Scholarship for Study Abroad to YA. This research
was also supported by the Stavros Niarchos Foundation (SNF) as part of its grant to the
SNF Institute for Global Infectious Disease Research at The Rockefeller University.

We are grateful to Mark Ebrahim, Johanna Sotiris, and Honkit Ng for their
technical advice and assistance for the Cryo-EM and Soeren Heissel and Henrik Molina
for MS analysis, Amalia Pasolli assistance for EM, David Shechter for providing NPM2
and NAP1 antibodies, Genzhe Lu and Daniil Tagaev for their contributions to the
optimization of MagIC-cryo-EM, and Rochelle Shih, Nick Prescott, Yiming Niu, and
Isabel Wassing for comments on the manuscript. We also thank Seth Darst, Elizabeth
Campbell, Thomas Huber, Michael Rout, Peter Fridy, Christopher Caffalette, Trevor Van
Eeuwen, Hiro Furukawa, Takashi Onikubo, Sue Biggins, Daniel Barrero, and Mengqiu
Jiang for consulting on the project. This work was conducted with the help of the High-
Performance Computing Resource Center, Proteomics Resource Center, the Evelyn
Gruss Lipper Cryo-Electron Microscopy Resource Center, Electron Microscopy
Resource Center, and Bio-Imaging Resource Center at the Rockefeller University.

<u>Author contributions</u>

Y.A. conceived and designed the study. Y.A. and H.A.K. conducted experiments. Y.A.
performed cryo-EM analyses. Y.A. and H.F. supervised the study. Y.A., H.A.K., and H.F.
wrote the manuscript.

<u>Declaration of interests</u>

YA, HAK, and HF have filed a patent application encompassing aspects of MagIC-cryo-
EM (PCT/US2023/03315). HF is affiliated with the Graduate School of Medical
Sciences, Weill Cornell Medicine, and Cell Biology Program at the Sloan Kettering
Institute.

<u>Supplemental information</u>

Document S1. Figures S1–S14 and Table S1-S4

Table S5. Excel file containing additional data too large to fit in a PDF, related to Figure
4.

<u>Figure titles and legends</u>

Figure 1. Single particle cryo-EM analysis of poly-nucleosomes attached to
magnetic beads (A) Schematic of a pilot cryo-EM experiment on magnetic beads.
Biotin-labeled 19-mer nucleosome arrays attached to 50 nm streptavidin-coated
magnetic nanobeads were loaded onto the cryo-EM grid. **(B)** Representative medium
magnification micrographs. The magnetic beads are seen as black dots (red arrows).
(C) Left; a representative high magnification micrograph. The micrograph was motion-
corrected and low-pass filtered to 5 Å resolution. Right; green circles indicate the
nucleosome-like particles selected by Topaz, and the blue areas indicate the halo-like
scattering. **(D)** The 3D structure of the nucleosome bound on magnetic beads.

Figure 2. MagIC-Cryo-EM structural determination of low-quantity and low-purity
targets (A) Schematic depicting the principle steps of MagIC-cryo-EM. **(B)** Graphical
representation of the MagIC-cryo-EM beads with 3HB and SAH spacers and GFP
nanobody target capture module. **(C)** Schematic of MagIC-cryo-EM for *in vitro*
reconstituted H1.8-GFP bound nucleosomes isolated from an excess of H1.8-free
nucleosomes. **(D)** Native PAGE analysis of H1.8-GFP bound nucleosomes and
unbound nucleosomes in the input. DNA staining by SYTO-60 is shown. **(E)** A
handmade humidity chamber used for the 5 min incubation of the cryo-EM grids on the
magnet. The humidity chamber was assembled using a plastic drawer. Wet tissues are
attached to the side walls of the chamber, which is sealed with a plastic cover to
maintain high humidity. Two pieces of neodymium magnets are stacked. A graphene
grid is held by a non-magnetic vitrobot tweezer and placed on the magnets. 4 μL of
sample is applied on the grid and incubated for 5 min. **(F)** Micrograph montage of the
grids without using magnetic concentration. The GFP-nanobody-MagIC-cryo-EM beads
(4 μL of 12.5 pM beads) were applied on the graphene-coated Quantifoil R 1.2/1.3 grid
and vitrified without incubation on a magnet. **(G)** Micrograph montage of the grids
without using magnetic concentration. The GFP-nanobody-MagIC-cryo-EM beads (4 μL
of 12.5 pM beads) were applied on the graphene-coated Quantifoil R 1.2/1.3 grid and
vitrified with 5 min incubation on two pieces of 40 x 20 mm N52 neodymium disc
magnets. **(H)** Quantitative analysis of the percentage of holes containing MagIC-cryo-
EM beads. Each data point represents the percentage of holes containing MagIC-cryo-

EM beads on each square mesh. **(I)** Quantitative analysis of the average number of
MagIC-cryo-EM beads per hole. Each data point represents the average number of
MagIC-cryo-EM beads per hole on each square mesh. The edges of the boxes and the
midline indicates the 25th, 50th, and 75th percentiles. Whiskers indicate the maximum
and lowest values in the dataset, excluding outliers. For the quantification, 11 square
meshes with 470 holes without magnetic concentration and 11 square meshes with 508
holes with 5 min incubation on magnets were used. **(J)** Representative motion corrected
micrographs of *in vitro* reconstituted H1.8-GFP nucleosomes captured by MagIC-cryo-
EM beads. The micrographs were low-pass filtered to 10 Å resolution. Green circles
indicate the nucleosome-like particles picked by Topaz. **(K)** 3D structure of the *in vitro*
reconstituted H1.8-GFP-bound nucleosome determined through MagIC-cryo-EM. The
pipeline for structural analysis is shown in Figure S2.

Figure 3. MagIC-Cryo-EM structural determination of H1.8-bound nucleosomes
from interphase and metaphase chromosomes in *Xenopus* egg extract. (A) Models
of potential cell cycle-dependent H1.8 dynamic binding mechanisms **(B)** Experimental
flow of MagIC-cryo-EM analysis for GFP-H1.8 containing complexes isolated from
chromosomes assembled in interphase and metaphase *Xenopus* egg extract.
Fluorescence microscopy images indicate localization of GFP-H1.8 to interphase and
metaphase chromosomes. DNA and GFP-H1.8 were detected either by staining with
Hoechst 33342 or GFP fluorescence, respectively. **(C)** Native PAGE of fragmented
interphase and metaphase chromosome sucrose gradient fractions. GFP-H1.8 and DNA
were detected with either GFP fluorescence or SYTO-60 staining, respectively. **(D)**
Western blot of GFP-H1.8 in interphase and metaphase chromosome sucrose gradient
fractions. GFP-H1.8 was detected using anti-GFP antibodies. **(E)** SDS-PAGE of the
sucrose gradient fractions 4 and 5 shown in (C), demonstrating heterogeneity of the
samples. Proteins were stained by gel code blue. Red arrows indicate the H1.8-GFP
bands. The full gel image is shown in Figure S4A. **(F)** *In silico* 3D classification of
interphase and metaphase H1.8-bound nucleosomes isolated from chromosomes in
Xenopus egg extract. To assess the structural variations and their population of H1.8-
bound nucleosomes, *ab initio* reconstruction and heterogenous reconstruction were
employed twice for the nucleosome-like particles isolated by the decoy classification.
The initial round of *ab initio* reconstruction and heterogenous reconstruction classified
the particles into three nucleosome-containing 3D models (A, B, C). Subsequent *ab*
initio reconstruction and heterogenous reconstruction on the class A, which has weak
H1.8 density, yielded three new nucleosome-containing structures, A1, A2, and A3. 3D
maps represent the structural variants of GFP-H1.8-bound nucleosomes. Red arrows
indicate extra densities that may represent H1.8. Green densities indicate on-dyad
H1.8. The bar graphs indicate the population of the particles assigned to each 3D class

in both interphase and metaphase particles (gray), interphase particles (blue), and
metaphase particles (red). The pipeline for structural analysis is shown in Figure S5A.
(G) Structures of H1.8-bound nucleosomes isolated from interphase and metaphase
chromosomes.

Figure 4. MagIC-cryo-EM and DuSTER reconstructed cryo-EM structures of
interphase-specific H1.8-bound NPM2. (A) Schematic of DuSTER workflow. **(B)** 2D
classes before and after particle curation with DuSTER. More 2D classes are shown in
Figure S10B-S10E. **(C)** 3D cryo-EM structure of interphase-specific H1.8-containing
complex. C5 symmetry was applied during structural reconstruction. The complete
pipeline is shown in Figures S8, S10, and S11. **(D)** MS identification of proteins that
cofractionated with H1.8 in sucrose gradient fraction 4 from interphase chromosomes
shown in Figure 3C. Portions of MagIC-cryo-EM beads prepared for cryo-EM were
subjected to MS. Proteins shown in red are the proteins that comprise the GPF
nanobody-MagIC-cryo-EM beads. Proteins shown in blue represent signals from H1.8-
GFP. **(E)** Western blot of NPM2 in the sucrose gradient fractions of interphase and
metaphase chromosome fragments. **(F)** The structural comparison of the crystal
structure of the pentameric NPM2 core (PDB ID: 1K5J), and AF2 predicted structure of
the pentameric NPM2 core, and MagIC-cryo-EM structures of NPM2-H1.8. The MagIC-
cryo-EM structures indicate NPM2 in the NPM2-H1.8 complex forms pentamer.

Figure 5. Structural variations of NPM2 bound to H1.8. (A) Structural differences
between the opened and closed forms of NPM2. Left panels show cryo-EM maps of the
opened and closed forms of NPM2 with H1.8. Middle panels show the atomic models.
The right panel shows the zoomed-in view of the open form (green) and closed form
(gray) of the NPM2 protomer. In the closed form, β8 runs straight from the sepal side to
the petal side. In the open form, the C-terminal portion of β8 is bent outward to the rim.
(B) Putative H1.8 density (red arrow) in the averaged NPM2-H1.8 structure. **(C)** The
NPM2 surface that contacts the putative H1.8 density (corresponding to aa 42-44) is
shown in orange. The H1.8-binding sites are accessible in the open form while they are
internalized in the closed form. Note that C-terminal acidic tracts A2 and A3 (Figure
S13A) are not visible in the cryo-EM structure but are likely to contribute to H1.8 binding
as well in both open and closed forms. **(D)** Model of the mechanism that regulates the
amount of the H1.8 in interphase and metaphase nucleosome.

Figure 6. Advantages of MagIC-cryo-EM over conventional cryo-EM methods. (A)
The on-bead-cryo-EM approach reduces preparation steps (for example, target

isolation, enrichment, and buffer exchange), which can lead to sample loss. (B) Sample
loss during the grid-freezing process is reduced by magnet-based enrichment of the
targets on cryo-EM grids. (C) The magnetic beads are easily identified in medium -
magnification montage maps, enabling the selection of areas where targets exist prior to
high-magnification data collection. (D) Targets are highly concentrated around the
beads, ensuring that each micrograph contains more than 100 usable particles for 3D
structure determination.

STAR Methods

Xenopus laevis

Xenopus laevis was purchased from Xenopus 1 (female, 4270; male, 4235). Vertebrate
animal protocols (20031 and 23020) approved by the Rockefeller University Institutional
Animal Care and Use Committee were followed.

Purification of Biotin-3HB-SPYcatcher003

Biotin-3HB-SPYcatcher003 was bacterially expressed and purified using pQE80-His$_{14}$-
bdSUMO-Cys-3HB-SPYcatcher003. To build the plasmid, a pQE80 derivative vector
encoding an N-terminal His-tag was amplified by PCR from pSF1389 [Addgene plasmid
104962] [64]. gBlock DNAs encoding *Brachypodium distachyon* SUMO (bdSUMO) [64]
and a computationally designed monomeric three-helix bundle [28] were synthesized by
IDT and used as a PCR template. DNA encoding SPYcatcher003 was amplified using
pSpyCatcher003 [Addgene plasmid # 133447] [65] as a PCR template. DNA fragments
were assembled by the Gibson assembly method [66]. *E. coli* Rosetta (DE3) cells
expressing His$_{14}$-bdSUMO-Cys-3HB-SPYcatcher003 were induced with 1 mM
isopropyl-β-D-thiogalactopyranoside (IPTG) at 25 ºC and then resuspended in 100 mL
buffer A (8 mM Na$_2$HPO$_4$, 2 mM KH$_2$PO$_4$, 537 mM NaCl, 2.7 mM KCl, 10 % glycerol, 2
mM β-mercaptoethanol, 1 mM PMSF, 20 mM imidazole with 1x cOmplete Protease
Inhibitor Cocktail EDTA-free [Roche]). The cells were disrupted by sonication, and the
soluble fraction was collected by centrifugation at 20,000 rpm (46,502 rcf) at 4 ºC for 30
min using a 45Ti rotor in Optima L80 (Beckman Coulter). This fraction was then mixed
with Ni-NTA agarose beads (Qiagen). Protein-bound Ni-NTA agarose beads were
packed into an Econo-column (bio-rad) and washed with 170 column volumes (CV) of
buffer B (8 mM Na$_2$HPO$_4$, 2 mM KH$_2$PO$_4$, 937 mM NaCl, 2.7 mM KCl, 10 % glycerol, 2
mM β-mercaptoethanol, 1 mM PMSF, 40 mM imidazole with 1x cOmplete EDTA-free
Protease Inhibitor Cocktail [Roche], pH 7.4). The beads were further washed with 33 CV
of Phosphate-Buffered Saline (PBS: 8 mM Na$_2$HPO$_4$, 2 mM KH$_2$PO$_4$, 137 mM NaCl, 2.7
mM KCl, pH 7.4) containing additional 5 % glycerol to remove β-mercaptoethanol. The

His$_{14}$-SUMO-tag was cleaved by incubating overnight at 4 ºC with N-terminal His-tagged SENP1 protease, which was expressed and purified using the previously described method with pSF1389 [Addgene plasmid # 104962] [64]. Ni-NTA agarose beads that bound the cleaved His$_{14}$-bdSUMO-tag and His$_{14}$-SENP1 were filtered out using an Econo-column (bio-rad). The cleaved 3HB-SPYcatcher003 with a cysteine residue at the N-terminal was concentrated using Amicon 30K (Millipore), mixed with EZ-link Maleimide-PEG2-Biotin (Thermo A39261), and left at 4 ºC overnight. Biotinylated 3HB-SPYcatcher003 was dialyzed overnight against PBS at 4 ºC. The dialyzed Biotin-3HB-SPYcatcher003 was further purified through a Hi-load Superdex75 16/600 column (Cytiva) and stored at -20 ºC in PBS containing 47.5 % glycerol.

Purification of Biotin-60 nm-SAH-SPYcatcher003 and Biotin-90 nm-SAH-SPYcatcher003

Biotin-30 nm-SAH-SPYcatcher003 and Biotin-60 nm-SAH-SPYcatcher003 were bacterially expressed and purified using pQE80-His$_{14}$-bdSUMO-Cys-30nm-SAH-SPYcatcher003 and pQE80-His$_{14}$-bdSUMO-Cys-60 nm-SAH-SPYcatcher003. DNA encoding 30 nm single alpha-helix (SAH) from *Trichomonas vaginalis* was amplified using pCDNA-FRT-FAK30 [Addgene plasmid # 59121] [29] as a PCR template. To extend the repeat to the desired length, MluI and AscI sites were inserted at the top and bottom of the DNA segment encoding 30 nm SAH, respectively. Although the target sequences for AscI (GG/CGCGCC) and MluI (A/CGCGT) are distinct, the DNA overhangs formed after the DNA digestion are identical. In addition, the DNA sequence formed by ligating these DNA overhangs translated into Lys-Ala-Arg, which does not disrupt a single alpha-helix. To generate pQE80-His$_{14}$-bdSUMO-Cys-60 nm-SAH-SPYcatcher3, two DNA fragments were prepared. The longer fragment was prepared by digesting pQE80-His$_{14}$-bdSUMO-Cys-30 nm-SAH-SPYcatcher003 with XhoI and MluI. The shorter fragment was prepared by digesting pQE80-His$_{14}$-bdSUMO-Cys-30 nm-SAH-SPYcatcher003 with XhoI and AscI. Target fragments were isolated by agarose gel extraction and ligated to form pQE80-His$_{14}$-bdSUMO-Cys-60nm-SAH-SPYcatcher003. Repeating these steps, pQE80-His$_{14}$-bdSUMO-Cys-90 nm-SAH-SPYcatcher003 was also generated.

E. coli Rosetta (DE3) cells expressing His$_{14}$-bdSUMO-Cys-SAH-SPYcatcher003 were induced with 1 mM IPTG at 18 ºC and then resuspended in 100 mL of buffer A before being disrupted by sonication. The soluble fraction was collected by centrifugation at 20,000 rpm (46,502 rcf) at 4 ºC for 30 min using a 45Ti rotor in Optima L80 (Beckman Coulter) and applied to a HisTrap HP column (Cytiva). The column was washed with 4 column volumes (CV) of buffer B. His$_{14}$-bdSUMO-Cys-SAH-SPYcatcher003 was eluted from the HisTrap column with buffer D (8 mM Na$_2$HPO$_4$, 2 mM KH$_2$PO$_4$, 137 mM NaCl,

2.7 mM KCl, 5 % glycerol, 200 mM imidazole [pH 7.4]). The eluted His$_{14}$-bdSUMO-Cys-
SAH-SPYcatcher003 was mixed with His$_{14}$-SENP1 and dialyzed against PBS containing
5 % glycerol at 4 ºC overnight. The dialyzed protein was applied to the HisTrap HP
column (Cytiva) to remove the cleaved His$_{14}$-bdSUMO-tag and His$_{14}$-SENP1. The
cleaved SAH-SPYcatcher003 was further purified through a MonoQ 5/50 column
(Cytiva). The purified SAH-SPYcatcher003 with a cysteine residue at the N-terminus
was concentrated with Amicon 10K (Millipore), mixed with EZ-link Maleimide-PEG2-
Biotin (Thermo A39261), and placed overnight at 4 ºC. The biotinylated SAH-
SPYcatcher003 was dialyzed against PBS at 4 ºC overnight. The dialyzed Biotin-SAH-
SPYcatcher003 was purified through a Hi-load Superdex200 16/600 column (Cytiva)
and stored at -20 ºC in PBS containing 47.5 % glycerol.

<u>Purification of Mono-SPYtag-avidin tetramer</u>

Mono-SPYtag-avidin tetramer was purified using a modified version of the method
described by Howarth et al. [67]. pET21-SPY-His$_6$-tag streptavidin and pET21-streptavidin
were generated by using pET21a-Streptavidin-Alive [Addgene plasmid # 20860] [67] as a
PCR template. SPY-His$_6$-tag streptavidin and untagged avidin were expressed
individually in *E. Coli* BL21(DE3) as inclusion bodies by inducing with 1 mM IPTG at 37
ºC. The cells expressing the proteins were resuspended in 100 mL of buffer E (50 mM
Tris-HCl, 1 mM EDTA) and disrupted by sonication. Insoluble fractions were collected by
centrifugation at 20,000 rpm at 4 ºC for 30 min using a 45Ti rotor in Optima L80
(Beckman Coulter). The insoluble pellets were washed by resuspending them in 50 ml
of buffer E and re-collecting them through centrifugation at 20,000 rpm at 4 ºC for 30
min using a 45Ti rotor in Optima L80 (Beckman Coulter). The washed insoluble pellets
were resuspended in 8 mL of 6 M guanidine HCl (pH 1.5) and dialyzed against 200 ml
of 6 M guanidine HCl (pH 1.5) overnight at 4 ºC. The denatured proteins were collected
by centrifugation at 20,000 rpm at 4 ºC for 30 min using a 45Ti rotor in Optima L80
(Beckman Coulter). Protein concentrations in soluble fractions were estimated based on
the absorbance at 260 nm. Denatured SPY-His$_6$-tag streptavidin and untagged
streptavidin were mixed at a 1:2.9 molar ratio and rapidly refolded by diluting them with
250 mL of PBS at 4 ºC. After 6 h of stirring at 4 ºC, aggregated proteins were removed
by centrifugation at 20,000 rpm at 4 ºC for 30 min using a 45Ti rotor in Optima L80
(Beckman Coulter). The supernatant was mixed with 62.7 g of solid ammonium sulfate
and stirred overnight at 4 ºC. Insolubilized proteins were removed with centrifugation at
20,000 rpm at 4 ºC for 30 min using a 45Ti rotor in Optima L80 (Beckman Coulter). The
supernatant was loaded into the HisTrap HP column (Cytiva). Refolded avidin tetramers
were eluted from the column by a linear gradient of imidazole (10 mM to 500 mM) in
PBS. The peak corresponding to mono-SPY-His-tagged streptavidin tetramer was
collected and concentrated using Amicon 10K (Millipore). The concentrated mono-SPY-

His$_6$-tagged streptavidin tetramer was further purified through Hiload superdex75
(Cytiva) and stored at -20 ºC in PBS containing 47.5 % glycerol.

Purification of SPYtag-GFP nanobody

MagIC-cryo-EM beads were optimized by testing three different GFP nanobodies:
tandem GFP nanobody, GFP enhancer nanobody, and LaG (llama antibody against
GFP)-10 (Figure S1). To express SPYtag-GFP nanobodies, plasmids pSPY-GFP
nanobody were built. The plasmid has a pQE80 backbone, and the DNA sequences that
encode His$_{14}$-bdSUMO-SPYtag-GFP nanobody were inserted into the multiple cloning
sites of the backbone. DNA encoding tandem GFP nanobody was amplified from
pN8his-GFPenhancer-GGGGS4-LaG16 [Addgene plasmid # 140442][68]. DNA encoding
GFP enhancer nanobody [69] was amplified from pN8his-GFPenhancer-GGGGS4-
LaG16. DNA encoding the LaG10 nanobody was amplified from a plasmid provided by
Dr. Michael Rout [70]. GFP nanobodies were expressed at 16 ºC in *E. coli* Rosetta (DE3)
by IPTG induction. The cells expressing His$_{14}$-bdSUMO-SPYtag-GFP nanobody were
resuspended with 100 mL buffer A and disrupted by sonication. The soluble fraction was
collected with centrifugation at 20,000 rpm (46,502 rcf) at 4 ºC for 30 min using a 45Ti
rotor in Optima L80 (Beckman Coulter) and applied to the HisTrap HP column (Cytiva).
The protein was eluted from the column with a step gradient of imidazole (50, 200, 400
mM) in buffer F (50 mM Tris-HCl (pH 8), 100 mM NaCl, 800 mM Imidazole, 5 %
Glycerol). The eluted His$_{14}$-bdSUMO-SPYtag-GFP nanobody was mixed with His$_{14}$-
SENP1 and dialyzed against PBS containing 5 % glycerol at 4 ºC overnight. The
dialyzed protein was applied to the HisTrap HP column (Cytiva) to remove the cleaved
His$_{14}$-bdSUMO-tag and His$_{14}$-SENP1. The cleaved SPYtag-GFP-nanobody was
concentrated with Amicon 10K (Millipore). The concentrated SPYtag-singular GFP
nanobody was further purified through Hiload superdex75 (Cytiva) and stored at -20 ºC
in PBS containing 47.5 % glycerol.

Purification of H1.8-GFP

To purify *Xenopus laevis* H1.8-superfolder GFP (sfGFP, hereafter GFP), pQE80-His$_{14}$-
bdSUMO-H1.8-GFP was generated by replacing bdSENP1 in pSF1389 vector to H1.8-
GFP. Using this plasmid, His$_{14}$-bdSUMO-H1.8-GFP was expressed in *E. Coli* Rosetta
(DE3) at 18 ºC with 1 mM IPTG induction. The soluble fraction was collected through
centrifugation at 20,000 rpm (46,502 rcf) at 4 ºC for 30 min using a 45Ti rotor in Optima
L80 (Beckman Coulter) and applied to the HisTrap HP column (Cytiva). His$_{14}$-bdSUMO-
H1.8-GFP was eluted from the column with a linear gradient of imidazole (100 mM to
800 mM) in PBS. The fractions containing His$_{14}$-bdSUMO-H1.8-GFP were collected,

mixed with SENP1 protease, and dialyzed overnight against PBS containing 5 %
glycerol at 4 ºC. The SENP1-treated sample was then applied to a Heparin HP column
(Cytiva) and eluted with a linear gradient of NaCl (137 mM to 937 mM) in PBS
containing 5 % glycerol. The fractions containing H1.8-GFP were collected and
concentrated using Amicon 30K (Millipore) before being applied to a Hiload
Superdex200 16/600 column (Cytiva) in PBS containing 5 % glycerol. The fractions
containing H1.8-GFP were collected, concentrated using Amicon 30K (Millipore), flash-
frozen, and stored at -80 ºC.

Purification of MNase

To purify MNase, pK19-His-bdSUMO-MNase was generated. Using this plasmid, His14-
bdSUMO-MNase was expressed in E. Coli JM101 at 18 °C with 2 mM IPTG induction.
The soluble fraction was collected through centrifugation at 20,000 rpm (46,502 rcf) at 4
°C for 30 min using a 45Ti rotor in Optima L80 (Beckman Coulter) and applied to the
HisTrap HP column (Cytiva). His14-bdSUMO-MNase was eluted from the column with a
linear gradient of imidazole (100 mM to 500 mM) in PBS. The fractions containing
His14-bdSUMO-MNase were collected, mixed with SENP1 protease, and dialyzed
overnight against PBS containing 5 % glycerol at 4 °C. The dialyzed protein was applied
to the HisTrap HP column (Cytiva) to remove the cleaved His14-bdSUMO-tag and
His14-SENP1. The cleaved MNase was concentrated with Amicon 3K (Millipore). The
concentrated MNase was further purified through Hiload superdex75 (Cytiva) and
stored at -80 °C in PBS containing 60 % glycerol.

Purification of X. laevis histones

All histones were purified using the method described previously [71]. Bacterially
expressed X. laevis H2A, H2B, H3.2, and H4 were purified from inclusion bodies. His-
tagged histones (H2A, H3.2, and H4) or untagged H2B expressed in bacteria were
resolubilized from the inclusion bodies by incubation with 6 M guanidine HCl. For His-
tagged histones, the solubilized His-tagged histones were purified using Ni-NTA beads
(Qiagen). For untagged H2B, the resolubilized histones were purified using a MonoS
column (Cytiva) under denaturing conditions before H2A-H2B dimer formation. To
reconstitute the H3–H4 tetramer and H2A–H2B dimer, the denatured histones were
mixed at an equal molar ratio and dialyzed to refold the histones by removing the
guanidine. His-tags were removed by TEV protease treatment, and the H3–H4 tetramer
and H2A–H2B dimer were isolated through a HiLoad 16/600 Superdex 75 column

(Cytiva). The fractions containing the H3–H4 tetramer and H2A–H2B dimer were
concentrated using Amicon 10K, flash-frozen, and stored at −80 °C.

<u>Preparation of *in vitro* reconstituted poly-nucleosome</u>

pAS696 containing the 19-mer of the 200 bp 601 nucleosome positioning sequence was
digested using HaeII, DraI, EcoRI, and XbaI. Both ends of the 19-mer of the 200 bp 601
DNA were labeled with biotin by Klenow fragment (NEB) with biotin-14-dATP [72]. The
nucleosomes were assembled with the salt dialysis method [72]. Purified DNAs were
mixed with H3-H4 and H2A-H2B, transferred into a dialysis cassette, and placed into a
high salt buffer (10 mM Tris-HCl [pH 7.5], 1 mM EDTA, 2 M NaCl, 5 mM β-
mercaptoethanol, and 0.01 % Triton X-100). Using a peristaltic pump, the high salt
buffer was gradually exchanged with a low salt buffer (10 mM Tris-HCl [pH 7.5], 1 mM
EDTA, 50 mM NaCl, 5 mM β-mercaptoethanol, 0.01 % Triton X-100) at roughly 2 ml/min
overnight at 4 °C. In preparation for cryo-EM image collection, the dialysis cassette
containing the sample was then placed in a buffer containing 10 mM HEPES-HCl (pH
7.4) and 30 mM KCl and dialyzed for 48 h at 4 °C.

<u>Native PAGE and SDS-PAGE</u>

For the native PAGE for nucleosome (Figure 3C), 15 µL of nucleosome fractions were
loaded onto a 0.5x TBE 6 % native PAGE gel. For the native PAGE for nucleosomal
DNA (Figure S4B), 15 µL of nucleosome fractions were mixed with 1 µL of 10 mg/mL
RNaseA (Thermo Scientific) and incubated at 55 °C for 30 min. To deproteinize and
reverse-crosslink DNA, RNaseA treated samples were then mixed with 1 µL of 19 mg/ml
Proteinase K solution (Roche) and incubated at 55 °C for overnight. Samples were
loaded to 0.5x TBE 6 % native PAGE. Native PAGE gels were stained by SYTO-60 to
detect DNA. SYTO-60 and GFP signals were scanned on a LI-COR Odyssey. For SDS-
PAGE analysis (Figure S4B), 20 µL of nucleosome fractions were mixed with 5 µL of 4x
SDS-PAGE sample buffer (200 mM Tris- HCl pH 6.8, 8 % SDS, 40 % glycerol, 10% β-
mercaptoethanol) and boiled for 10 min at 96 °C. Samples were loaded to a 4 %–20 %
gradient gel (Bio-Rad, # 5671095).

<u>Western blot</u>

For the western blot of nucleosome fractions (Figure 3D), 20 µL of nucleosome fractions
were mixed with 5 µL of 4x SDS-PAGE sample buffer and boiled for 10 min at 96 °C.
Samples were loaded to a 4 %–20 % gradient gel (Bio-Rad, # 5671095).

For the H1.8-GFP complementation assay (Figure S3), 2 µL egg extract samples were
added to 38 µL of 1x SDS-PAGE sample buffer (50 mM Tris- HCl pH 6.8, 2 % SDS,
10 % glycerol, 2.5 % β-mercaptoethanol) and boiled for 5 min at 96°C. Samples were
mixed by vortex and spun at 13,200 rpm for 1 min before gel electrophoresis. 10 µL out
of 40 µL samples were separated in 4–20 % gradient gel (Bio-Rad, # 5671095).

The SDS-PAGE gels were transferred into the western blot cassette and transferred to
a nitrocellulose membrane (Cytiva, # 10600000) with 15 V at 4 °C overnight. The
transferred membranes were blocked with Intercept TBS Blocking Buffer (LI-COR
Biosciences, # 927-60001). Primary and secondary antibodies were diluted in Intercept
TBS Antibody Diluent (LI-COR Biosciences, #927-65001). For Figure S3A, as primary
antibodies, mouse monoclonal antibody against GFP (Santa Cruz Biotechnology, # sc-
9996, 1:1000 dilution) and rabbit polyclonal antibody against *X. laevis* H1.8 [73] (final: 1
µg/mL) were used. For Figure S14, as primary antibodies, rabbit polyclonal antibody
against *X. laevis* H1.8 [73], rabbit polyclonal antisera against *X. laevis* NAP1 (1:500
dilution) [54], NPM2 (1:500 dilution) [53], and rabbit polyclonal antibody against
phosphorylated histone H3 Thr3 (MilliporeSigma, # 07-424, 1:5000 dilution) were used.
NAP1 and NPM2 antibody are kind gifts of David Shechter. As secondary antibodies,
IRDye 800CW goat anti-rabbit IgG (LI-COR, # 926-32211; 1:10,000) and IRDye 680RD
goat anti-mouse IgG (LI-COR, # 926-68070; 1:15,000) were used. The images were
taken with Odyssey M Infrared Imaging System (LI-COR Biosciences).

<u>Immunoprecipitation (IP) assay in *Xenopus* egg extract</u>

For the IP assay (Figure S14), antibody against rabbit IgG, in-house purified from pre-
immune rabbit serum by HiTrap Protein A HP (# 17040301), and antibody against *X.*
laevis H1.8 (# RU2130) were conjugated to Protein-A coupled Dynabeads (Thermo
Fisher Scientific, # 10001D) at 20 µg/mL beads at 4 °C for overnight on a rotator. rIgG
and H1.8 antibody beads were crosslinked using 5 mM BS3 (Thermo Fisher Scientific, #
A39266) resuspended in PBS (pH 7.4) at room temperature for 30 min and quenched
by 50 mM Tris-HCl (pH 7.4) resuspended in PBS (pH 7.4) at room temperature for 20-
30 min on a rotator. All antibody beads were washed extensi vely using wash/coupling
buffer (10 mM K-HEPES (pH 8.0) and 150 mM KCl), followed by sperm dilution buffer
(10 mM K-HEPES (pH 8.0), 1 mM $MgCl_2$, 100 mM KCl, 150 mM sucrose). The beads
were left on ice until use.

Interphase egg extract (30 µL) was prepared by incubating at 20 °C for 60 min after
adding $CaCl_2$ (final: 0.4 mM) and cycloheximide (final: 100 µg/mL) to fresh CSF egg
extract. Mitotic egg extract (CSF egg extract, 30 µL) was also incubated at 20 °C for 60
min without any additives. After 60 min incubation, each mitotic and interphase egg
extract was transferred to antibody-conjugated beads (10 µL) after removing sperm

dilution buffer on a magnet stand (Sergi Lab Supplies, Cat# 1005). Beads-extract
mixtures were mixed and incubated on ice for 45 min with flicking tubes every 15 min.
After 45 min, beads were collected using a magnet stand at 4 °C and washed 3 times
with beads wash buffer (sperm dilution buffer supplemented 1x cOmplete EDTA-free
protease inhibitor cocktail (Roche, # 4693132001), 1x PhosSTOP (Roche, #
4906845001), and 0.1 % (v/v) Triton-X (BIO-RAD, # 1610407)). Beads are resuspended
in 20 µL of 1x SDS sample buffer and loaded 10 µL out of 20 µL to a SDS-PAGE gel.
Methods for SDS-PAGE and western blot are described above.

<u>Trial MagIC-cryo-EM with poly-nucleosome (used in Figure 1)</u>

A total of 60 fmol of Absolute Mag streptavidin nano-magnetic beads (CD bioparticles:
WHM-X047, 50 nM size) were mixed with 100 µL of EM buffer A (10 mM HEPES-KOH
[pH 7.4], 30 mM KCl, 1 mM EGTA, 0.3 ng/µL leupeptin, 0.3 ng/µL pepstatin, 0.3 ng/µL
chymostatin, 1 mM Sodium Butyrate, 1 mM beta-glycerophosphate, 1 mM $MgCl_2$, 2%
trehalose, 0.2 % 1,6-hexanediol). The beads were collected by incubation on two pieces
of 40 x 20 mm N52 neodymium disc magnets (DIYMAG: D40x20-2P-NEW) at 4 ºC for
30 min and then resuspended in 120 µL of EM buffer A. The two pieces of strong
neodymium magnets have to be handled carefully as magnets can leap and slam
together from several feet apart. Next, 60 µL of 34 nM nucleosome arrays formed on the
biotinylated 19-mer 200 bp 601 DNA were mixed with the beads and rotated at 20 ºC for
2 h. To remove unbound nucleosomes, the biotin-poly-nucleosome-bound nano-
magnetic beads were collected after 40 min of incubation on the N52 neodymium disc
magnets and then resuspended in 300 µL EM buffer containing 10 µM biotin. A 100 µL
portion of the biotin-poly-nucleosome-bound nano-magnetic beads solution was
incubated on the N52 neodymium disc magnets for 30 min and then resuspended in 20
µL EM buffer A. Finally, 3 µL of biotin-poly-nucleosome-bound nano-magnetic beads
solution was added onto a glow-discharged Quantifoil Gold R 1.2/1.3 300 mesh grid
(Quantifoil). The samples were vitrified under 100% humidity, with a 20-sec incubation
and 5-sec blotting time using the Vitrobot Mark IV (FEI).

The grid was imaged on a Talos Arctica (Thermo Fisher) equipped with a 200 kV field
emission gun and K2 camera. A total of 657 movies were collected at a magnification of
x 72,000 (1.5 Å/pixel) using super-resolution mode, as managed by SerialEM [74]. Movie
frames are motion-corrected and dose-weighted patch motion correction in CryoSPARC
v3 with output Fourier cropping fac½ 1/2 [34]. Particles were picked by Topaz v0.2.3 with
around 2000 manually picked nucleosome-like particles as training models [27]. Picked
particles were extracted using CryoSPARC v3 (extraction box size = 200 pixel). 2D
classification of extracted particles was done using 100 classes in CryoSPARC v3.
Using 2D classification results, particles were split into the nucleosome-like groups and

the non-nucleosome-like groups. Four 3D initial models were generated for both groups
with ab initio reconstruction in CryoSPARC v3 (Class similarity = 0). One nucleosome-
like model was selected and used as a given model of heterogeneous reconstruction
with all four of the "decoy" classes generated from the non-nucleosome-like group. After
the first round of 3D classification, the particles assigned to the "decoy" classes were
removed, and the remaining particles used for a second round of 3D classification using
the same settings as the first round. These steps were repeated until more than 90 % of
particles wer classified as a nucleosome-like class. To isolate the nucleosome class that
has visible H1.8 density, four to six 3D references were generated with ab initio
reconstruction of CryoSPARC v3 using purified nucleosome-like particles (Class
similarity = 0.9). Refined particles were further purified with the heterogeneous
refinement using an H1.8-visible class and an H1.8-invisible class as decoys. The
classes with reasonable extra density were selected and refined with homogeneous
refinement. The final resolution was determined with the gold stand FSC threshold (FSC
= 0.143).

Preparation of in vitro reconstituted mono-nucleosome and H1.8-GFP bound mono-
nucleosome

The 193 bp 601 DNA fragment was amplified by a PCR reaction [75,76]. The nucleosomes
were assembled with the salt dialysis method described above. The reconstituted
nucleosome was dialyzed into buffer XL (80 mM PIPES-KOH [pH 6.8], 15 mM NaCl, 60
mM KCl, 30 % glycerol, 1 mM EGTA, 1 mM $MgCl_2$, 10 mM β-glycerophosphate, 10 mM
sodium butyrate). H1.8-GFP was mixed with nucleosome with a 1.25 molar ratio in the
presence of 0.001 % poly L-glutamic acid (wt 3,000-15,000) (Sigma-Aldrich) and
incubated at 37 ºC for 30 min. As a control nucleosome sample without H1.8-GFP, the
sample without H1.8-GFP was also prepared. The samples were then crosslinked
adding a 0.5-time volume of buffer XL containing 3 % formaldehyde and incubating for
90 min on ice. The crosslink reaction was quenched by adding 1.7 volume of quench
buffer (30 mM HEPES-KOH (pH 7.4), 150 mM KCl, 1 mM EGTA, 10 ng/μL leupeptin, 10
ng/μL pepstatin, 10 ng/μL chymostatin, 10 mM sodium butyrate, 10 mM β-
glycerophosphate, 400 mM glycine, 1 mM $MgCl_2$, 5 mM DTT). The quenched sample
was layered onto the 10-25 % linear sucrose gradient solution with buffer SG (15 mM
HEPES-KOH [pH 7.4], 50 mM KCl, 10-22 % sucrose, 10 μg/ml leupeptin, 10 μg/ml
pepstatin, 10 μg/ml chymostatin, 10 mM sodium butyrate, 10 mM β-glycerophosphate, 1
mM EGTA, 20 mM glycine) and spun at 32,000 rpm (max 124,436 rcf) and 4 ºC for 13 h
using SW55Ti rotor in Optima L80 (Beckman Coulter). The centrifuged samples were
fractionated from the top of the sucrose gradient. The concertation of H1.8-GFP bound
nucleosome in each fraction is calculated based on the 260 nm light absorbance
detected by Nanodrop 2000 (Thermo Scientific).

Preparation of GFP nanobody attached MagIC-cryo-EM beads

A total of 25 fmol of Absolute Mag streptavidin nanomagnetic beads (CD Bioparticles: WHM-X047) were transferred to a 0.5 mL protein LoBind tube (Eppendorf) and mixed with 200 pmol of inner spacer module protein (biotin-3HB-SPYcatcher003 or biotin-60nm-SAH-SPYcatcher003) in 200 µL of EM buffer A (10 mM HEPES-KOH [pH 7.4], 30 mM KCl, 1 mM EGTA, 10 ng/µL leupeptin, 10 ng/µL pepstatin, 10 ng/µL chymostatin, 1 mM Sodium Butyrate, and 1 mM beta-glycerophosphate) and the mixture was incubated at 4 ºC for 10 h. To wash the beads, the mixture was spun at 13,894 rpm (16,000 rcf) at 4 ºC for 10 min using the SX241.5 rotor in an Allegron X-30R centrifuge (Beckman Coulter). The beads that accumulated at the bottom of the tube were resuspended in 200 µL of EM buffer A. Subsequently, 200 pmol of mono-SPYtag-avidin tetramer was added to the beads in 200 µL of EM buffer A, and the mixture was incubated at 4 ºC for 10 h. Again, the beads were washed by collecting them via centrifugation and resuspending them in 200 µL of EM buffer A. This washing step was repeated once more, and 800 pmol of outer spacer module protein (biotin-30 nm-SAH-SPYcatcher003, biotin-60 nm-SAH-SPYcatcher003 or biotin-90 nm-SAH-SPYcatcher003) were added and incubated at 4 ºC for 10 h. The beads were washed twice and resuspended with 25 µL of EM buffer A. 20 µL of this mixture was transferred to a 0.5 ml protein LoBind tube and mixed with 640 pmol of SPYtag-GFP nanobody and incubated at 4 ºC for 10 h. The beads were washed twice and resuspended with 25 µL of EM buffer A. The assembled GFP nanobody attached MagIC-cryo-EM beads can be stored in EM buffer A containing 50 % glycerol at -20ºC for several weeks.

Graphene grids preparation

Graphene grids were prepared using the method established by Han et al. [77] with minor modifications. Briefly, monolayer graphene grown on the copper foil (Grolltex) was coated by polymethyl methacrylate (Micro chem, EL6) with the spin coat method. The copper foil was removed by 1 M of ammonium persulfate. The graphene monolayer coated by polymethyl methacrylate was attached to gold or copper grids with carbon support film (Quantifoil) and baked for 30 min at 130 ºC. The polymethyl methacrylate was removed by washing with 2-butanone, water, and 2-propanol on a hotplate.

Optimization of the spacer module length by the MagIC-cryo-EM of *in vitro* reconstituted H1.8-GFP bound nucleosome (used in Figure S1)

To prepare the MagIC-cryo-EM beads capturing H1.8-GFP bound mono-nucleosome, 4
fmol of GFP nanobody-attached MagIC-cryo-EM beads with different spacer lengths
were mixed with 100 nM (28 ng/µL) of *in vitro* reconstituted crosslinked H1.8-GFP
bound mono-nucleosome in 100 µL of PBS containing 15~30 % glycerol and incubated
at 4 ºC for 12 h. To wash the beads, the beads were collected with centrifugation at
13,894 rpm (16,000 rcf) at 4 ºC for 20 min using SX241.5 rotor in Allegron X-30R
(Beckman Coulter) and resuspended with 200 µL of PBS containing 15~30 % glycerol.
This washing step was repeated once again, and the beads were resuspended with 100
µL of EM buffer C (10 mM HEPES-KOH [pH 7.4], 30 mM KCl, 1 mM EGTA, 10 ng/µL
leupeptin, 10 ng/µL pepstatin, 10 ng/µL chymostatin, 1 mM sodium butyrate, 1 mM β-
glycerophosphate, 1.2 % trehalose, and 0.12 % 1,6-hexanediol). This washing step was
repeated once again, and the beads were resuspended with 100~200 µL of EM buffer C
(theoretical beads concentration: 20~40 pM).

To vitrify the grids, a plasma-cleaned graphene-coated Quantifoil gold R1.2/1.3 400
mesh grid (Quantifoil) featuring a monolayer graphene coating [77] was held using a pair
of sharp non-magnetic tweezers (SubAngstrom, RVT-X). The two pieces of strong
neodymium magnets have to be handled carefully as magnets can leap and slam
together from several feet apart. Subsequently, 4 µL of MagIC-cryo-EM beads capturing
H1.8-GFP-nucleosomes were applied to the grid. The grid was then incubated on the 40
x 20 mm N52 neodymium disc magnets for 5 min within an in-house high-humidity
chamber to facilitate magnetic bead capture. Once the capture was complete, the
tweezers anchoring the grid were transferred and attached to the Vitrobot Mark IV (FEI),
and the grid was vitrified by employing a 2-second blotting time at room temperature
under conditions of 100% humidity.

We found that gold grids are suitable for MagIC-cryo-EM, whereas copper grids
worsened the final resolution of the structures presumably due to magnetization of the
copper grids during the concentration process which then interfered with the electron
beam and caused the grid to vibrate during data collection (Figure S1, Test 7).

The vitrified grids were loaded onto the Titan Krios (ThermoFisher), equipped with a 300
kV field emission gun and a K3 direct electron detector (Gatan). A total of 1890 movies
were collected at a magnification of x 64,000 (1.33 Å/pixel) using super-resolution
mode, as managed by SerialEM [74].

Movie frames were corrected for motion using MotionCor2 [78] installed in Relion v4 [79] or
patch motion correction implemented in CryoSPARC v4. Particles were picked with
Topaz v0.2 [80], using approximately 2000 manually picked nucleosome-like particles as
training models. The picked particles were then extracted using CryoSPARC v4
(extraction box size = 256 pixels) [34]. Nucleosome-containing particles were isolated
through decoy classification using heterogeneous reconstruction with one nucleosome-

like model and four decoy classes generated through ab initio reconstruction in
CryoSPARC v4. CTF refinement and Bayesian polishing were applied to the
nucleosome-containing particles in the Relion v4 [79,81]. To isolate the nucleosome class
with visible H1.8 density, four 3D references were generated through ab initio
reconstruction in CryoSPARC v4 using purified nucleosome-like particles (Class
similarity = 0.9). These four 3D references were used for heterogeneous reconstruction.
Two of the classes had strong H1.8 density. Using the particles assigned in these
classes, non-uniform refinement was performed in CryoSPARC v4. The final resolution
was determined using the gold standard FSC threshold (FSC = 0.143).

1021

MagIC-cryo-EM of *in vitro* reconstituted H1.8-GFP bound nucleosome using the mixture
of the H1.8-GFP bound and unbound nucleosomes (shown in Figure 2)

A total of 0.5 fmol of GFP-singular nanobodies conjugated to 3HB-60nm-SAH magnetic
beads were mixed with 1.7 nM (0.5 ng/µL) of H1.8-GFP bound nucleosome and 53 nM
(12 ng/µL) of H1.8-free nucleosome in 100 µL of buffer SG (15 mM HEPES-KOH [pH
7.4], 50 mM KCl, 12% sucrose, 1x LPC, 10 mM Sodium Butyrate, 10 mM β-
glycerophosphate, 1 mM EGTA) containing approximately 17 % sucrose. The mixture
was then incubated at 4 ºC for 10 h. To wash the beads, they were collected by
centrifugation at 13,894 rpm (16,000 rcf) at 4 ºC for 20 min using the SX241.5 rotor in
an Allegron X-30R centrifuge (Beckman Coulter). Subsequently, the beads were
resuspended in 200 µL of EM buffer C. This washing step was repeated twice, and the
beads were finally resuspended in approximately 80 µL of EM buffer C, resulting in a
theoretical bead concentration of 6.25 pM.

To vitrify the grids, 4 µL of the samples were applied to plasma-cleaned graphene-
coated Quantifoil gold R1.2/1.3 300-mesh grids (Quantifoil). The grid was then
incubated on the 40 x 20 mm N52 neodymium disc magnets for 5 minutes and vitrified
using the Vitrobot Mark IV (FEI) with a 2-sec blotting time at room temperature under
100 % humidity. The vitrified grids were loaded onto the Titan Krios (ThermoFisher),
equipped with a 300 kV field emission gun and a K3 direct electron detector (Gatan). A
total of 1890 movies were collected at a magnification of x 64,000 (1.33 Å/pixel) using
super-resolution mode, as managed by SerialEM [74].

The analysis pipeline is described in Figure S2. Movie frames were corrected for motion
using MotionCor2 [78], which was installed in Relion v4 [79]. Particles were picked with
Topaz v0.2.3 [80], using approximately 2000 manually picked nucleosome-like particles as
training models. The picked particles were then extracted using CryoSPARC v4
(extraction box size = 256 pixels) [34]. Nucleosome-containing particles were isolated
through decoy classification using heterogeneous reconstruction with one nucleosome-
like model and four decoy classes generated through ab initio reconstruction in

CryoSPARC v3.3. CTF refinement and Bayesian polishing were applied to the

nucleosome-containing particles in Relion v4 [79,81]. To isolate the nucleosome class with

visible H1.8 density, four 3D references were generated through ab initio reconstruction

in CryoSPARC v3.3 using purified nucleosome-like particles (Class similarity = 0.9).

These four 3D references were used for heterogeneous reconstruction. Two of the

classes had strong H1.8 density. Using the particles assigned in these classes, non-

uniform refinement was performed in CryoSPARC v3.3. The final resolution was

determined using the gold standard FSC threshold (FSC = 0.143).

1058

<u>Assessment of the efficiency of the magnetic concentration of the MagIC-cryo-EM on</u>

<u>cryo-EM grid (shown in Figure 2)</u>

A plasma-cleaned graphene-coated Quantifoil copper R1.2/1.3 400 mesh grid

(Quantifoil) was held using non-magnetic Vitrobot tweezers (SubAngstrom).

Subsequently, 4 µL of 12.5 pM GFP-nanobody attached MagIC-cryo-EM beads were

applied to the grid. The grid was then incubated on the 40 x 20 mm N52 neodymium

disc magnets for 5 min within a high-humidity chamber. As a control experiment, several

grids were frozen by omitting the magnetic incubation steps. Once the capture was

complete, the tweezers anchoring the grid were attached to the Vitrobot Mark IV (FEI),

and the grid was vitrified by employing a 2-sec blotting time at room temperature under

conditions of 100% humidity. The vitrified grids were subjected to cryo-EM to collect 8 x

8 or 9 x 9 montage maps at x2,600 magnification on Talos Arctica to capture the whole

area of each square mesh. The efficiency of the magnetic concentration of the MagIC-

cryo-EM beads was quantitatively assessed by counting the percentage of holes

containing MagIC-cryo-EM beads and counting the average number of MagIC-cryo-EM

beads per hole. For the quantification, 11 square meshes with 470 holes were used for

the condition without magnetic concentration. For the condition with 5 min incubation on

magnets, 11 square meshes with 508 holes were used. The boxplots and the scatter

plots were calculated by the seaborn.boxplot and seaborn.stripplot tools in the Seaborn

package [82] and visualized by Matplotlib [83]. Outlier data points that are not in 1.5 times of

the interquartile range, the range between the 25th and 75th percentile, were excluded.

1080

<u>Functional assessment of H1.8-GFP in *Xenopus* egg extract (Shown in Figure S3)</u>

The functional replaceability of H1.8-GFP in *Xenopus* egg extracts was assessed

through whether H1.8-GFP could rescue the chromosome morphological defect caused

by depletion of endogenous H1.8. Mitotic chromosome morphology and length were

assessed through the previously described method (23) with some modifications.

The cytostatic factor (CSF)-arrested metaphase *Xenopus laevis* egg extracts were
prepared using the method as described [84]. Anti-rabbit IgG (SIGMA, Cat# I5006) and
rabbit anti-H1.8 custom antibodies [25] (Identification# RU2130) were conjugated to
Protein-A coupled Dynabeads (Thermo Fisher Scientific, # 10001D) at 250 µg/ml beads
at 4 °C for overnight on a rotator. IgG and H1.8 antibody beads were crosslinked using
4 mM BS3 (Thermo Fisher Scientific, # A39266) resuspended in PBS (pH 7.4) at room
temperature for 45 min and quenched by 50 mM Tris-HCl (pH 7.4) resuspended in PBS
(pH 7.4) at room temperature for 20-30 min on a rotator. All antibody beads were
washed extensively using wash/coupling buffer (10 mM K-HEPES (pH 8.0) and 150 mM
KCl), followed by sperm dilution buffer (10 mM K-HEPES (pH 8.0), 1 mM $MgCl_2$, 100
mM KCl, 150 mM sucrose). After the two rounds of depletion at 4 °C for 45 min using 2
volumes of antibody-coupled beads on a rotator, the beads were separated using a
magnet (Sergi Lab Supplies, Cat# 1005). For the complementation of H1.8, 1.5 µM of
recombinantly purified H1.8 or H1.8-GFP was supplemented into H1.8-depleted CSF
egg extract.

To assess chromosome morphology in the metaphase chromosomes with spindles, 0.4
mM $CaCl_2$ was added to CSF-arrested egg extracts containing *X. laevis* sperm (final
concentration 2000/µL) to cycle the extracts into interphase at 20 °C for 90 min. To
induce mitotic entry, half the volume of fresh CSF extract and 40 nM of the non-
degradable cyclin BΔ90 fragment were added after 90 min and incubated at 20 °C for
60 min.

Metaphase spindles for fluorescent imaging were collected by a published method [85]. 15
µL metaphase extracts containing mitotic chromosomes were diluted into 2 mL of fixing
buffer (80 mM K-PIPES pH 6.8, 1 mM $MgCl_2$, 1 mM EGTA, 30 % (v/v) glycerol, 0.1 %
(v/v) Triton X-100, 2 % (v/v) formaldehyde) and incubated at room temperature for 5
min. The fixed samples were layered onto a cushion buffer (80 mM K-PIPES pH 6.8, 1
mM $MgCl_2$, 1 mM EGTA, 50 % (v/v) glycerol) with a coverslip (Fisher Scientific, Cat#
12CIR-1.5) placed at the bottom of the tube and centrifuged at 5,000x *g* for 15 min at
16 °C in a swinging bucket rotor (Beckman Coulter, JS-5.3 or JS-7.5). The coverslips
were recovered and fixed with pre-chilled methanol (–20 °C) for 5 min. The coverslips
were extensively washed with TBST (TBS supplemented 0.05% Tween-20) and then
blocked with antibody dilution buffer (AbDil; 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 %
BSA, 0.02 % NaN_3) at 4 °C for overnight.

Individualized mitotic chromosome samples were prepared as described previously [25].
10 µL of metaphase extracts containing mitotic chromosomes were diluted into 60 µL of
chromosome dilution buffer (10 mM K-HEPES pH 8, 200 mM KCl, 0.5 mM EGTA, 0.5
mM $MgCl_2$, 250 mM sucrose), mixed by gentle flicking, and incubated at room
temperature for 8 min. Diluted samples were transferred into 3 mL of fixing buffer (80
mM K-PIPES pH 6.8, 1 mM $MgCl_2$, 1 mM EGTA, 30 % (v/v) glycerol, 0.1 % (v/v) Triton

X-100, 2 % (v/v) formaldehyde), mixed by inverting tubes, and incubated for total 6 min

at room temperature. Similar to mitotic chromosome preparation, the fixed samples

were subjected to glycerol cushion centrifugation (7,000x *g* for 20 min at 16 °C) using a

swinging bucket rotor (Beckman, JS-7.5). Coverslips were recovered, fixed with pre-

chilled methanol (–20 °C) for 5 min, extensively washed with TBST, and then blocked

with AbDil buffer at 4 °C overnight.

For immunofluorescence microscopy, primary and secondary antibodies were diluted in

AbDil buffer. Coverslips were incubated in primary antibody solution at room

temperature for 60 min and secondary antibody at room temperature for 45 min. DNA

was stained using NucBlue™ Fixed Cell ReadyProbes™ Reagent (Thermo Fisher

Scientific, Cat# R37606) following manufacture's protocol. Coverslips were extensively

washed using TBST between each incubation and sealed on the slide glass using

ProLong™ Diamond Antifade Mountant (Thermo Fisher Scientific, Cat# P36965). For

primary antibodies, mouse monoclonal antibody against α-tubulin (MilliporeSigma, Cat#

T9026, 1:1000 dilution) and rabbit polyclonal antibody against *X. laevis* CENP-A [86]

(Identification# RU1286), 1:1000 dilution). For secondary antibodies, mouse IgG was

detected using Cy™3 AffiniPure F(ab')$_2$ Fragment Donkey Anti-Mouse IgG (H+L)

(Jackson ImmunoResearch, Cat# 715-166-150; 1:500 dilution) and rabbit IgG was

detected using Cy™5 AffiniPure Donkey Anti-Rabbit IgG (H+L) (Jackson

ImmunoResearch, Cat# 711-175-152; 1:500 dilution).

The immunofluorescence imaging was performed on a DeltaVision Image

Restoration microscope (Applied Precision), which is a widefield inverted microscope

equipped with a pco. edge sCMOS camera (pco). Immunofluorescence samples were

imaged with 1 µm z-sections using a 60× Olympus UPlan XApo (1.42 NA) oil objective,

and were processed with a iterative processive deconvolution algorithm using the Soft-

WoRx (Applied Precision).

For chromosome length measurements, the length of individualized mitotic

chromosomes were manually traced on a single maximum intensity slice using

segmented line tool in Fiji software (ver. 2.9.0). Data was summarized using R (ver.

4.2.2) and visualized as SuperPlots [87] using ggplot2 package in R and RStudio (ver.

RSTUDIO-2023.09.1-494). For the representative images in Figure S3, max projection

images were prepared in Fiji using z-stuck function. For the visibility, the brightness and

contrast of representative images were adjusted using GIMP software (ver. 4.2.2).

Adjustment was done using a same setting among all images.

1159

<u>Fractionation of chromosomes isolated from *Xenopus* egg extracts (Used for Figure 3)</u>

Nucleosomes were isolated from *Xenopus* egg extract chromosomes using the
previously described method [3]. To prevent the spontaneous cycling of egg extracts, 0.1
mg/ml cycloheximide was added to the CSF extract. H1.8-GFP was added to the CSF
extract at a final concentration of 650 nM, equivalent to the concentration of
endogenous H1.8 [88]. For interphase chromosome preparation, *Xenopus laevis* sperm
nuclei (final concentration 2000/µL) were added to 5 mL of CSF extracts, which were
then incubated for 90 min at 20 ºC after adding 0.3 mM $CaCl_2$ to release the CSF
extracts into interphase. For metaphase sperm chromosome preparation, cyclin B ∆90
(final concentration 24 µg/mL) and 1 mL of fresh CSF extract were added to 2 ml of the
extract containing interphase sperm nuclei prepared using the method described above.
To make up for the reduced H1.8-nucleosome formation in interphase, we used 5 ml of
egg extracts for preparing interphase chromosomes and 2 mL of extracts for metaphase
chromosomes. The extracts were incubated for 60 min at 20 ºC, with gentle mixing
every 10 min. To crosslink the *Xenopus* egg extracts chromosomes, nine times the
volume of ice-cold buffer XL (80 mM PIPES-KOH [pH 6.8], 15 mM NaCl, 60 mM KCl,
30 % glycerol, 1 mM EGTA, 1 mM $MgCl_2$, 10 mM β-glycerophosphate, 10 mM sodium
butyrate, 2.67 % formaldehyde, 0.001% digitonin) was added to the interphase or
metaphase extract containing chromosomes, which was further incubated for 60 min on
ice. These fixed chromosomes were then layered on 3 mL of fresh buffer SC (80 mM
HEPES-KOH [pH 7.4], 15 mM NaCl, 60 mM KCl, 1.17 M sucrose, 50 mM glycine, 0.15
mM spermidine, 0.5 mM spermine, 1.25x cOmplete EDTA-free Protease Inhibitor
Cocktail (Roche), 10 mM beta-glycerophosphate, 10 mM sodium butyrate, 1 mM EGTA,
1 mM $MgCl_2$) in 50 ml centrifuge tubes (Falcon, #352070). The tubes were spun at
3,300 (2,647 rcf) rpm at 4 ºC for 40 min using a JS 5.3 rotor in an Avanti J-26S
centrifuge (Beckman Coulter). Pellets containing fixed chromosomes were resuspended
with 10 mL of buffer SC, layered on 3 ml of fresh buffer SC in 14 mL centrifuge tubes
(Falcon, #352059), and spun at 3,300 (2,647 rcf) rpm at 4 ºC for 40 min using a JS 5.3
rotor in an Avanti J-26S centrifuge (Beckman Coulter). The chromosomes were
collected from the bottom of the centrifuge tube and resuspended with buffer SC.
Chromosomes were pelleted by centrifugation at 5,492 rpm (2,500 rcf) using an
SX241.5 rotor in an Allegron X-30R centrifuge (Beckman Coulter). The chromosome
pellets were resuspended with 200 µL of buffer SC. To digest chromatin, MNase
concentration and reaction time were tested on a small scale and optimized to the
condition that produce 180-200 bp DNA fragments. After the optimization, 0.6 and 0.3
U/µL of MNase were added to interphase and metaphase chromosomes, respectively.
Then, $CaCl_2$ was added to a final concentration of 7.4 mM, and the mixture was
incubated at 4 ºC for 4 h. The MNase reaction was stopped by adding 100 µL MNase
stop buffer B (80 mM PIPES-KOH (pH 6.8), 15 mM NaCl, 60 mM KCl, 30% glycerol, 20
mM EGTA, 1 mM $MgCl_2$, 10 mM β-glycerophosphate, 10 mM sodium butyrate, 3.00 %
formaldehyde). The mixtures were incubated on ice for 1 h and then diluted with 700 µL

of quench buffer (30 mM HEPES-KOH (pH 7.4), 150 mM KCl, 1 mM EGTA 1x LPC, 10
mM sodium butyrate, 10 mM β-glycerophosphate, 400 mM glycine, 1 mM MgCl$_2$, 5 mM
DTT). The soluble fractions released by MNase were isolated by taking supernatants
after centrifugation at 13,894 rpm (16,000 rcf) at 4 ºC for 30 min using an SX241.5 rotor
in an Allegron X-30R centrifuge (Beckman Coulter). The supernatants were collected
and layered onto a 10-22 % linear sucrose gradient solution with buffer SG (15 mM
HEPES-KOH [pH 7.4], 50 mM KCl, 10-22 % sucrose, 10 µg/mL leupeptin, 10 µg/mL
pepstatin, 10 µg/mL chymostatin, 10 mM sodium butyrate, 10 mM β-glycerophosphate,
1 mM EGTA, 20 mM glycine) and spun at 32,000 rpm (max 124,436 rcf) and 4 ºC for 13
h using an SW55Ti rotor in an Optima L80 centrifuge (Beckman Coulter). The samples
were fractionated from the top of the sucrose gradient. The concentration of H1.8 in
each fraction was determined by western blot. 15 µL of each sucrose gradient fraction
was incubated at 95 ºC with 1 % sodium dodecyl sulfate (SDS) and applied for SDS-
PAGE with a 4-20 % gradient SDS-PAGE gel (Bio-rad). The proteins were transferred to
a nitrocellulose membrane (Cytiva) from the SDS-PAGE gel using TE42 Tank Blotting
Units (Hoefer) at 15 V, 4 ºC for 4 h. As primary antibodies, 1 µg/mL of mouse
monoclonal Anti-GFP Antibody sc-9996 (Santa Cruz Biotechnology) and as secondary
antibodies, IR Dye 800CW goat anti-mouse IgG (Li-Cor 926-32210; 1:15,000) were
used. The images were taken with an Odyssey Infrared Imaging System (Li-Cor). The
existence of the H1.8-GFP bound nucleosomes was confirmed by native PAGE. 15 µL
of each sucrose gradient fraction was applied for a 6 % x0.5 TEB native PAGE gel. The
DNA was stained with SYTO-60 (Invitrogen S11342: 1:10,000). The images of SYTO-60
signal and GFP signal were taken with an Odyssey Infrared Imaging System (Li-Cor).

1224

<u>MagIC-cryo-EM of H1.8-GFP bound nucleosomes isolated from chromosomes</u>
<u>assembled in *Xenopus* egg extract (used in Figure 3)</u>

Tween 20 was added to a final concentration of 0.01% to the 350 µL of fraction 5 from
the interphase or metaphase sucrose gradient fractions shown in Figure 3 and S4.
These samples were then mixed with 1 fmol of GFP nanobody-conjugated MagIC-cryo-
EM beads. The mixture was incubated at 4 ºC for 10 h. The beads were washed four
times with EM buffer C containing 0.01 % Tween 20, as described above. Finally, the
beads were resuspended in approximately 80 µL of EM buffer C containing 0.001 %
Tween 20.

To vitrify the grids, 4 µL of the samples were applied to plasma-cleaned graphene-
coated Quantifoil gold R1.2/1.3 300-mesh grids (Quantifoil). The grid was then
incubated on the 40 x 20 mm N52 neodymium disc magnets for 5 minutes and vitrified
using the Vitrobot Mark IV (FEI) with a 2-second blotting time at room temperature
under 100 % humidity. The vitrified grids were loaded onto the Titan Krios

(ThermoFisher), equipped with a 300 kV field emission gun and a K3 direct electron
detector (Gatan). A total of 677 movies for the interphase and 965 movies for the
metaphase were collected at a magnification of x 64,000 (1.33 Å/pixel) using super-
resolution mode, as managed by SerialEM [74].

The analysis pipeline is described in Figure S5. Movie frames were corrected for motion
using MotionCor2 [78], which was installed in Relion v4 [79]. The micrographs for interphase
and metaphase MagIC-cryo-EM were combined and subjected to particle picking.
Particles were picked with Topaz v0.2.3 [80], using approximately 2000 manually picked
nucleosome-like particles as training models. The picked particles were then extracted
using CryoSPARC v4 (extraction box size = 256 pixels) [34]. Nucleosome-containing
particles were isolated through decoy classification using heterogeneous reconstruction
with one nucleosome-like model and four decoy classes generated through ab initio
reconstruction in CryoSPARC v4. CTF refinement and Bayesian polishing were applied
to the nucleosome-containing particles in Relion v4 [79,81]. To isolate the nucleosome
class with visible H1.8 density, three 3D references were generated through ab initio
reconstruction in CryoSPARC v4 using purified nucleosome-like particles (Class
similarity = 0.9). This step was repeated for the class with weak H1.8 density (Class A).
Non-uniform refinement was performed in CryoSPARC v4 for each class. Subsequently,
to isolate the H1.8-bound nucleosome structures in interphase and metaphase, the
particles were separated based on their original movies. Using these particle sets, the
3D maps of the interphase and metaphase H1.8-bound nucleosomes were refined
individually through non-uniform refinement in CryoSPARC v4. The final resolution was
determined using the gold standard FSC threshold (FSC = 0.143).

1262

<u>Isolation of interphase-specific H1.8-GFP containing complex by MagIC-cryo-EM (used</u>
<u>in Figure 4)</u>

Tween20 was added to a final concentration of 0.01% to 350 µL of fraction 4 from the
interphase sucrose gradient fractions shown in Figure 3C. The sample was then mixed
with 1 fmol of GFP nanobody-conjugated MagIC-cryo-EM beads. The mixture was
incubated at 4 °C for 10 h. The beads were washed four times with EM buffer C
containing 0.01% Tween 20, as described above. Finally, the beads were resuspended
in approximately 80 µL of EM buffer C containing 0.001 % Tween 20. The resuspended
MagIC-cryo-EM beads solution was subjected to the MS and cryo-EM.

1272

<u>Mass spectrometry</u>

For the MS analysis, 20 µL of the resuspended solution containing the MagIC-cryo-EM
beads isolating interphase-specific H1.8-GFP containing complex was incubated at

95 °C for 10 minutes to reverse the crosslink. The 20 µL each of the sucrose gradient
fractions 4 and 5 (interphase and metaphase) was also incubated at 95 °C. The
samples were then applied to an SDS-PAGE (4 %–20 % gradient gel, Bio-Rad). The gel
was stained with Coomassie Brilliant Blue G-250 (Thermo Fisher). The corresponding
lane was cut into pieces approximately 2 mm x 2 mm in size. The subsequent
destaining, in-gel digestion, and extraction steps were carried out as described [89]. In
brief, the cut gel was destained using a solution of 30 % acetonitrile and 100 mM
ammonium bicarbonate in water. Gel pieces were then dehydrated using 100 %
acetonitrile. Disulfide bonds were reduced with dithiothreitol, and cysteines were
alkylated using iodoacetamide. Proteins were digested by hydrating the gel pieces in a
solution containing sequencing-grade trypsin and endopeptidase LysC in 50 mM
ammonium bicarbonate. Digestion proceeded overnight at 37 °C. The resulting peptides
were extracted three times with a solution of 70 % acetonitrile and 0.1 % formic acid.
These extracted peptides were then purified using in-house constructed
micropurification C18 tips. The purified peptides were subsequently analyzed by LC-
MS/MS using a Dionex 3000 HPLC system equipped with an NCS3500RS nano- and
microflow pump, coupled to an Orbitrap ASCEND mass spectrometer from Thermo
Scientific. Peptides were separated by reversed-phase chromatography using solvent A
(0.1 % formic acid in water) and solvent B (80 % acetonitrile, 0.1 % formic acid in water)
across a 70-min gradient. Spectra were recorded in positive ion data-dependent
acquisition mode, with fragmentation of the 20 most abundant ions within each duty
cycle. MS1 spectra were recorded with a resolution of 120,000 and an AGC target of
2e5. MS2 spectra were recorded with a resolution of 30,000 and an AGC target of 2e5.
The spectra were then queried against a *Xenopus laevis* database [88,90], concatenated
with common contaminants, using MASCOT through Proteome Discoverer v.1.4 from
Thermo Scientific. The abundance value for each protein is calculated as the average of
the 3 most abundant peptides belonging to each protein [91]. All detected proteins are
listed in Table S5. The keratin-related proteins that were considered to be contaminated
during sample preparation steps and the proteins with less than 5% coverage that were
considered to be misannotation were not shown in Figure 4D and Supplementary Table
2.

1307

<u>Cryo-EM data collection of interphase-specific H1.8-GFP containing complex isolated by</u>
<u>MagIC-cryo-EM beads (used in Figure 4)</u>

To vitrify the grids, 4 µL of the resuspended solution containing the MagIC-cryo-EM
beads isolated interphase-specific H1.8-GFP containing complex were applied to
plasma-cleaned in-house graphene attached Quantifoil gold R1.2/1.3 300-mesh grids
(Quantifoil). The grid was then incubated on the 40 x 20 mm N52 neodymium disc
magnets for 5 min and vitrified using the Vitrobot Mark IV (FEI) with a 2-sec blotting time

at room temperature under 100 % humidity. The vitrified grids were loaded onto the
Titan Krios (ThermoFisher), equipped with a 300 kV field emission gun and a K3 direct
electron detector (Gatan). At a magnification of x 105,000 (0.86 Å/pixel), 4,543 movies
were collected. At a magnification of x 105,000 (1.08 Å/pixel), 1,807 movies were
collected.

1320

<u>Application of DuSTER for Cryo-EM analysis of interphase-specific H1.8-GFP</u>
<u>containing complex isolated by MagIC-cryo-EM beads (used in Figure 4)</u>

The pipeline to generate the initial 3D model is described in Figure S8. Movie frames
are motion-corrected and dose-weighted patch motion correction in CryoSPARC v4 with
output Fourier cropping factor 1/2 [34]. To remove low S/N ratio particles that are not
reproducibly recentered during 2D classification, through DuSTER, particles picking with
Topaz v0.2 [80] were repeated twice to assign two picked points for each protein particle
on micrographs. Training of Topaz was performed individually for each picked particle
set using the same approximately 2000 manually picked particles as training models.
The particles in these two picked particle sets were then extracted using CryoSPARC v4
(extraction box size = 185.8 Å) [34]. These two extracted particle sets were individually
applied to 2D classification in CryoSPARC v4 (600 classes). These 2D classifications
did not generate any reasonable 2D classes of interphase-specific H1.8-GFP containing
complex that was expected from the particle images on the original motion-corrected
micrographs. The reproducibility of the particles recentering can be assessed by the D.
Smaller value of D indicates that two pick points on each particle are reproducibly
recentered during 2D classification. To remove duplicate particles at closed distances,
we used this tool to keep the recentered points whose D are shorter than D_{TH}. The
DuSTER curation can be achieved by using the 'Remove Duplicate Particles' tool in
CryoSPARC. Although the tool was originally designed to remove duplicate particles at
closed distances, we used this tool to keep the recentered points whose D are shorter
than D_{TH}. All particles from two individual particle sets after the 2D classification were
applied to the 'Remove Duplicate Particles' tool in CryoSPARC v4 using the 'Remove
Duplicates Entirely' option (Minimum separation distance: 20Å). Although the tool was
originally designed to remove duplicate particles at closed distances, we used this tool
to keep the recentered points whose D are shorter than D_{TH}. The particles whose
recentered points whose D are shorter than D_{TH} and were the particles used in further
downstream processing, were sorted as 'rejected particles'. These particles were
applied to the Particle Sets Tool in CryoSPARC v4 to split them into two individual
particle sets. 2D DuSTER, including particle re-centering, particle extraction, and
particle splitting steps, was repeated seven times. After seven rounds of 2D DuSTER,
the particles were manually curated by removing the 2D classes with unreasonable
sizes or shapes for the interphase-specific H1.8-GFP containing complex. The 2D

images of removed classes are shown in Figure S8. After manual curation, the particles
were further cleaned by an additional four rounds of 2D DuSTER. The particles were
further cleaned by the Class Probability Filtering Tool in CryoSPARC v4. 2D
classification was performed twice for one of the cleaned particle sets. The particles
whose 2D class probability scores were lower than 0.3 in both replicates of 2D
classification were removed. The redundant 2D classifications were necessary to
prevent unintentional loss of high S/N particles.The duplicated class probability filtering
was repeated six times. Using the filtered particles, 2D classification was performed
twice. The high-resolution classes with reasonable protein-like features were manually
selected from both 2D classification results. To prevent unintentional contamination of
low S/N particles, the 92,382 particles that were selected in both 2D classification runs
were used for ab initio 3D reconstruction (C5 symmetry applied). The 3D structure was
highly similar to NPM2, and we were convinced that the interphase-specific H1.8-GFP
containing complex is NPM2-H1.8-GFP complex.

The pipeline for the particle cleaning using 3D DuSTER is described in Figure S10. After
seven rounds of 2D DuSTER for the particles picked by Topaz, decoy 3D classification
was employed to remove nucleosomes and GFP complexed with GFP-nanobody. The
nucleosome 3D model was generated by ab initio 3D reconstruction using the particles
assigned to nucleosome-like 2D classes. The 3D model of GFP complexed with GFP-
nanobody was modeled from the crystal structure of the complex (PDB ID: 3k1k) [69]
using EMAN2 [92]. Noise 3D models were generated by ab initio 3D reconstruction using
the low S/N particles that were removed during 2D DuSTER. Using these models and
the initial 3D model of NPM2-H1.8-GFP, heterogeneous 3D refinement was performed
twice in CryoSPARC v4. To prevent unintentional loss of high S/N particles, particles
that were assigned to the nucleosome and GFP complexed with GFP-nanobody class in
both heterogeneous 3D refinement results were removed. By using the Remove
Duplicate Particles and Particle Sets tools in CryoSPARC v4, the particles in picked
particle set 2 that corresponded to the particles cleaned by decoy classification were
selected. Using both picked particle sets, heterogeneous 3D refinement of CryoSPARC
v4 was performed individually. Using the same procedure as 2D DuSTER, the particles
that were reproducibly centered in each particle set were selected (Minimum separation
distance: 15 Å). 3D DuSTER was repeated six times. To conduct 3D DuSTER more
comprehensively, 3D refinements were performed for each picked particle set three
times. Particle curation based on the distance was performed for all nine combinations
of these 3D refinement results, and this comprehensive 3D DuSTER was repeated once
again. Using the particles in picked particle set 1 after 3D DuSTER, 2D classification
was performed twice. The noise classes were manually selected from both 2D
classification results. To prevent unintentional loss of high S/N particles, particles that
were assigned to the noise class in both 2D classification runs were removed. This
duplicated 2D classification and manual selection was repeated twice. During the 2D

classification, 2D classes that represent GFP-nanobody were found. To remove the
particles, duplicated decoy 3D classification was employed once again. The remaining
162,995 particles were used for the 3D structure reconstruction.

The pipeline for 3D structure reconstruction using the particle curated by 3D DuSTER is
described in Figure S11. Using the 162,995 particles after the 3D DuSTER, *ab initio* 3D
reconstruction (5 classes, C5) was performed five times. The particles assigned to the
NPM2-like classes were selected. To prevent unintentional loss of high S/N particles,
particles that were assigned to the noise class in all five *ab initio* 3D reconstruction runs
were removed. For the 'averaged' NPM2 structure, a single 3D map was built by ab
initio 3D reconstruction (1 class, C5) using the remaining 92,428 particles. The 3D map
was refined by local refinement using the particles after symmetry expansion. For the
structural variants of the NPM2, particles were split into the 2 classes by ab initio 3D
reconstruction (2 class, C5). The ab initio 3D reconstruction (3 class, C5) was
performed again for each class, and the particles were manually split into the 3 groups
to generate 'open,' 'half-open,' and 'closed' NPM2 structures.

The initial atomic model of *Xenopus laevis* NPM2 pentamer was built by ColabFold
v1.5.5, which implements AlphaFold2 and MMseqs2 [56–58]. The full-length *Xenopus*
laevis NPM2 pentamer structure was docked on the cryo-EM maps by the Dock-in-map
tool in Phenix v1.21[93]. The atomic coordinates of the disordered regions were removed.
The atomic model was refined using the Starmap v1.2.15 [94]. The refined models were
further refined using the real-space refinement in Phenix v1.21[93].

For reconstituting the 3D maps without applying symmetry, the particles used for
reconstituting 'open,' 'half-open,' and 'closed' NPM2 structures were applied to the
manual picking tool in cryoSPARC to remove the 3D alignment information attached to
the particle images. The particle images were extracted and applied to the ab initio 3D
reconstruction (1 class, C1).

3D FSC was plotted by the Orientation Diagnostics tool integrated in the cryoSPARC
v4.4.

1422

1423 <u>AlphaFold2 prediction of the NPM2-H1.8 complex structure</u>

1424 The AF2 models of the *Xenopus laevis* NPM2-H1.8 complex were built by ColabFold
1425 v1.5.5, by submitting five NPM2 and one H1.8 amino acid sequence as input [56–58].

1426

1427 <u>3D structure visualization</u>

1428 Local resolution was estimated by cryoSPARC v4.4. All 3D structures, including cryo-
1429 EM density maps, cartoon depictions, and surface depictions with electrostatic potential,
1430 were visualized by the UCSF ChimeraX software [95].

1431

<u>Data and materials availability</u>

Cryo-EM density maps have been deposited in the EM Data Resource under accession
codes EMD-42599 (*in vitro* reconstituted poly-nucleosome), EMD-42598 (*in vitro*
reconstituted H1-GFP bound nucleosome), EMD-42594 (*Xenopus* egg extract H1-GFP
bound nucleosome structure containing both interphase and metaphase particles),
EMD-42596 (interphase *Xenopus* egg extract H1-GFP bound nucleosome), EMD-42597
(metaphase *Xenopus* egg extract H1-GFP bound nucleosome), EMD-43238 (Averaged
NPM2-H1.8-GFP structure), EMD- 43239 (open NPM2-H1.8-GFP structure), and EMD-
43240 (closed NPM2-H1.8-GFP structure). The atomic coordinates have been
deposited in the Protein Data Bank under accession codes PDB 8VHI (averaged NPM2-
H1.8-GFP structure), PDB 8VHJ (open NPM2-H1.8-GFP structure), and PDB 8VHK
(closed NPM2-H1.8-GFP structure). The cryo-EM data will be disclosed upon the
publication of this manuscript. The plasmids for generating MagIC-cryo-EM beads were
deposited to Addgene under accession codes #214835 (Non tagged Avidin), #214836
(SPYtag-Histag-Avidin), #214837 (SPYtag-GFPnanobody), #214838 (Cys-3HB-
SPYcatcher), #214839 (Cys-30nmSAH-SPYcatcher), and #214840 (Cys-60nmSAH-
SPYcatcher).

1449

References

1. Azinas, S., and Carroni, M. (2023). Cryo-EM uniqueness in structure
determination of macromolecular complexes: A selected structural anthology.
Curr. Opin. Struct. Biol. *81*, 102621. 10.1016/j.sbi.2023.102621.

2. Natchiar, S.K., Myasnikov, A.G., Kratzat, H., Hazemann, I., and Klaholz, B.P.
(2017). Visualization of chemical modifications in the human 80S ribosome
structure. Nature *551*, 472–477. 10.1038/nature24482.

3. Arimura, Y., Shih, R.M., Froom, R., and Funabiki, H. (2021). Structural features of
nucleosomes in interphase and metaphase chromosomes. Mol. Cell *81*, 4377-
4397. 10.1016/j.molcel.2021.08.010.

4. Arimura, Y., and Funabiki, H. (2022). Structural Mechanics of the Alpha-2-
Macroglobulin Transformation. J. Mol. Biol. *434*, 167413.
https://doi.org/10.1016/j.jmb.2021.167413.

5. Leesch, F., Lorenzo-Orts, L., Pribitzer, C., Grishkovskaya, I., Roehsner, J.,
Chugunova, A., Matzinger, M., Roitinger, E., Belačić, K., Kandolf, S., et al. (2023).

A molecular network of conserved factors keeps ribosomes dormant in the egg.
Nature *613*, 712–720. 10.1038/s41586-022-05623-y.

6.　　Bhella, D. (2019). Cryo-electron microscopy: an introduction to the technique, and
considerations when working to establish a national facility. Biophys. Rev. *11*,
515–519. 10.1007/s12551-019-00571-w.

7.　　Ravelli, R.B.G., Nijpels, F.J.T., Henderikx, R.J.M., Weissenberger, G.,
Thewessem, S., Gijsbers, A., Beulen, B.W.A.M.M., López-Iglesias, C., and
Peters, P.J. (2020). Cryo-EM structures from sub-nl volumes using pin-printing
and jet vitrification. Nat. Commun. *11*, 1–9. 10.1038/s41467-020-16392-5.

8.　　Dandey, V.P., Wei, H., Zhang, Z., Tan, Y.Z., Acharya, P., Eng, E.T., Rice, W.J.,
Kahn, P.A., Potter, C.S., and Carragher, B. (2018). Spotiton: New features and
applications. J. Struct. Biol. *202*, 161–169. 10.1016/j.jsb.2018.01.002.

9.　　Kelly, D.F., Abeyrathne, P.D., Dukovski, D., and Walz, T. (2008). The Affinity Grid:
A Pre-fabricated EM Grid for Monolayer Purification. J. Mol. Biol. *382*, 423–433.
10.1016/j.jmb.2008.07.023.

10.　Llaguno, M.C., Xu, H., Shi, L., Huang, N., Zhang, H., Liu, Q., and Jiang, Q.X.
(2014). Chemically functionalized carbon films for single molecule imaging. J.
Struct. Biol. *185*, 405–417. 10.1016/j.jsb.2014.01.006.

11.　Yu, G., Li, K., and Jiang, W. (2016). Antibody-based affinity cryo-EM grid.
Methods *100*, 16–24. 10.1016/j.ymeth.2016.01.010.

12.　Wang, L., Ounjai, P., and Sigworth, F.J. (2008). Streptavidin crystals as
nanostructured supports and image-calibration references for cryo-EM data
collection. J. Struct. Biol. *164*, 190–198. 10.1016/j.jsb.2008.07.008.

13.　Skrajna, A., Goldfarb, D., Kedziora, K.M., Cousins, E.M., Grant, G.D., Spangler,
C.J., Barbour, E.H., Yan, X., Hathaway, N.A., Brown, N.G., et al. (2020).
Comprehensive nucleosome interactome screen establishes fundamental
principles of nucleosome binding. Nucleic Acids Res., 1–18.
10.1093/nar/gkaa544.

14.　Hebbes, T.R., Thorne, A.W., and Crane-Robinson, C. (1988). A direct link
between core histone acetylation and transcriptionally active chromatin. EMBO J.
7, 1395–1402. 10.1002/j.1460-2075.1988.tb02956.x.

15.　Solomon, M.J., Larsen, P.L., and Varshavsky, A. (1988). Mapping protein-DNA
interactions in vivo with formaldehyde: evidence that histone H4 is retained on a
highly transcribed gene. Cell *53*, 937–947. 10.1016/s0092-8674(88)90469-2.

16.　Gilmour, D.S., and Lis, J.T. (1986). RNA polymerase II interacts with the promoter
region of the noninduced hsp70 gene in Drosophila melanogaster cells. Mol. Cell.
Biol. *6*, 3984–3989. 10.1128/mcb.6.11.3984-3989.1986.

17.　Zou, Z., Ohta, T., Miura, F., and Oki, S. (2022). ChIP-Atlas 2021 update: a data-
mining suite for exploring epigenomic landscapes by fully integrating ChIP-seq,
ATAC-seq and Bisulfite-seq data. Nucleic Acids Res. *50*, W175–W182.

10.1093/nar/gkac199.

18. Wang, C.I., Alekseyenko, A.A., Leroy, G., Elia, A.E.H., Gorchakov, A.A., Britton, L.M.P., Elledge, S.J., Kharchenko, P. V., Garcia, B.A., and Kuroda, M.I. (2013). Chromatin proteins captured by ChIP-mass spectrometry are linked to dosage compensation in Drosophila. Nat. Struct. Mol. Biol. *20*, 202–209. 10.1038/nsmb.2477.

19. Luger, K., Mäder, A.W., Richmond, R.K., Sargent, D.F., and Richmond, T.J. (1997). Crystal structure of the nucleosome core particle at 2.8 Å resolution. Nature *389*, 251–260. 10.1038/38444.

20. Chua, E.Y.D., Vasudevan, D., Davey, G.E., Wu, B., and Davey, C.A. (2012). The mechanics behind DNA sequence-dependent properties of the nucleosome. Nucleic Acids Res. *40*, 6338–6352. 10.1093/nar/gks261.

21. Bednar, J., Garcia-Saez, I., Boopathi, R., Cutter, A.R., Papai, G., Reymer, A., Syed, S.H., Lone, I.N., Tonchev, O., Crucifix, C., et al. (2017). Structure and Dynamics of a 197 bp Nucleosome in Complex with Linker Histone H1. Mol. Cell *66*, 384-397.e8. 10.1016/j.molcel.2017.04.012.

22. Zhou, B.R., Jiang, J., Feng, H., Ghirlando, R., Xiao, T.S., and Bai, Y. (2015). Structural Mechanisms of Nucleosome Recognition by Linker Histones. Mol. Cell *59*, 628–638. 10.1016/j.molcel.2015.06.025.

23. Zhou, B.R., Feng, H., Kale, S., Fox, T., Khant, H., de Val, N., Ghirlando, R., Panchenko, A.R., and Bai, Y. (2021). Distinct Structures and Dynamics of Chromatosomes with Different Human Linker Histone Isoforms. Mol. Cell *81*, 166-182.e6. 10.1016/j.molcel.2020.10.038.

24. Dombrowski, M., Engeholm, M., Dienemann, C., Dodonova, S., and Cramer, P. (2022). Histone H1 binding to nucleosome arrays depends on linker DNA length and trajectory. Nat. Struct. Mol. Biol. *29*, 493–501. 10.1038/s41594-022-00768-w.

25. Choppakatla, P., Dekker, B., Cutts, E.E., Vannini, A., Dekker, J., and Funabiki, H. (2021). Linker histone h1.8 inhibits chromatin-binding of condensins and dna topoisomerase ii to tune chromosome length and individualization. Elife *10*, 2020.12.20.423657. 10.7554/eLife.68918.

26. Bonnafous, P., Perrault, M., Le Bihan, O., Bartosch, B., Lavillette, D., Penin, F., Lambert, O., and Pécheur, E.I. (2010). Characterization of hepatitis C virus pseudoparticles by cryo-transmission electron microscopy using functionalized magnetic nanobeads. J. Gen. Virol. *91*, 1919–1930. 10.1099/vir.0.021071-0.

27. Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A.J., and Berger, B. (2019). Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. Nat. Methods *16*, 1153–1160. 10.1038/s41592-019-0575-8.

28. Huang, P.S., Oberdorfer, G., Xu, C., Pei, X.Y., Nannenga, B.L., Rogers, J.M., DiMaio, F., Gonen, T., Luisi, B., and Baker, D. (2014). High thermodynamic

stability of parametrically designed helical bundles. Science *346*, 481–485.
10.1126/science.1257481.

29.   Sivaramakrishnan, S., and Spudich, J.A. (2011). Systematic control of protein
interaction using a modular ER/K α-helix linker. Proc. Natl. Acad. Sci. U. S. A.
108, 20467–20472. 10.1073/pnas.1116066108.

30.   Zakeri, B., Fierer, J.O., Celik, E., Chittock, E.C., Schwarz-Linek, U., Moy, V.T.,
and Howarth, M. (2012). Peptide tag forming a rapid covalent bond to a protein,
through engineering a bacterial adhesin. Proc. Natl. Acad. Sci. U. S. A. *109*.
10.1073/pnas.1115485109.

31.   Gong, X., Qian, H., Zhou, X., Wu, J., Wan, T., Cao, P., Huang, W., Zhao, X.,
Wang, X., Wang, P., et al. (2016). Structural insights into the Niemann-Pick C1
(NPC1)-mediated cholesterol transfer and ebola infection. Cell *165*, 1467–1478.
10.1016/j.cell.2016.05.022.

32.   Nguyen, A.H., Thomsen, A.R.B., Cahill, T.J., Huang, R., Huang, L.Y., Marcink, T.,
Clarke, O.B., Heissel, S., Masoudi, A., Ben-Hail, D., et al. (2019). Structure of an
endosomal signaling GPCR–G protein–β-arrestin megacomplex. Nat. Struct. Mol.
Biol. *26*, 1123–1131. 10.1038/s41594-019-0330-y.

33.   Lilic, M., Chen, J., Boyaci, H., Braffman, N., Hubin, E.A., Herrmann, J., Müller, R.,
Mooney, R., Landick, R., Darst, S.A., et al. (2020). The antibiotic sorangicin A
inhibits promoter DNA unwinding in a Mycobacterium
tuberculosis rifampicin-resistant RNA polymerase. Proc. Natl. Acad.
Sci. *117*, 30423 LP – 30432. 10.1073/pnas.2013706117.

34.   Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). CryoSPARC:
Algorithms for rapid unsupervised cryo-EM structure determination. Nat. Methods
14, 290–296. 10.1038/nmeth.4169.

35.   Maresca, T.J., Freedman, B.S., and Heald, R. (2005). Histone H1 is essential for
mitotic chromosome architecture and segregation in Xenopus laevis egg extracts.
J. Cell Biol. *169*, 859–869. 10.1083/jcb.200503031.

36.   Willcockson, M.A., Healton, S.E., Weiss, C.N., Bartholdy, B.A., Botbol, Y., Mishra,
L.N., Sidhwani, D.S., Wilson, T.J., Pinto, H.B., Maron, M.I., et al. (2021). H1
histones control the epigenetic landscape by local chromatin compaction. Nature
589, 293–298. 10.1038/s41586-020-3032-z.

37.   Yusufova, N., Kloetgen, A., Teater, M., Osunsade, A., Camarillo, J.M., Chin, C.R.,
Doane, A.S., Venters, B.J., Portillo-Ledesma, S., Conway, J., et al. (2021).
Histone H1 loss drives lymphoma by disrupting 3D chromatin architecture. Nature
589, 299–305. 10.1038/s41586-020-3017-y.

38.   Zhou, B.-R., Feng, H., Kato, H., Dai, L., Yang, Y., Zhou, Y., and Bai, Y. (2013).
Structural insights into the histone H1-nucleosome complex. Proc. Natl. Acad. Sci.
110, 19390 LP – 19395. 10.1073/pnas.1314905110.

39.   Song, F., Chen, P., Sun, D., Wang, M., Dong, L., Liang, D., Xu, R.M., Zhu, P., and

Li, G. (2014). Cryo-EM study of the chromatin fiber reveals a double helix twisted
by tetranucleosomal units. Science *344*, 376–380. 10.1126/science.1251413.

40.   Miller, K.E., and Heald, R. (2015). Glutamylation of Nap1 modulates histone H1
dynamics and chromosome condensation in Xenopus. J. Cell Biol. *209*, 211–220.
10.1083/jcb.201412097.

41.   Hite, R.K., and MacKinnon, R. (2017). Structural Titration of Slo2.2, a Na+-
Dependent K+ Channel. Cell *168*, 390-399.e11. 10.1016/j.cell.2016.12.030.

42.   Yang, Z., Fang, J., Chittuluru, J., Asturias, F.J., and Penczek, P.A. (2012).
Iterative stable alignment and clustering of 2D transmission electron microscope
images. Structure *20*, 237–247. 10.1016/j.str.2011.12.007.

43.   Dutta, S., Akey, I. V., Dingwall, C., Hartman, K.L., Laue, T., Nolte, R.T., Head,
J.F., and Akey, C.W. (2001). The crystal structure of nucleoplasmin-core:
Implications for histone binding and nucleosome assembly. Mol. Cell *8*, 841–853.
10.1016/S1097-2765(01)00354-9.

44.   Laskey, R.A., Honda, B.M., Mills, A.D., and Finch, J.T. (1978). Nucleosomes are
assembled by an acidic protein which binds histones and transfers them to DNA.
Nature *275*, 416–420. 10.1038/275416a0.

45.   Philpott, A., Leno, G.H., and Laskey, R.A. (1991). Sperm decondensation in
Xenopus egg cytoplasm is mediated by nucleoplasmin. Cell *65*, 569–578.
10.1016/0092-8674(91)90089-h.

46.   Ohsumi, K., and Katagiri, C. (1991). Characterization of the ooplasmic factor
inducing decondensation of and protamine removal from toad sperm nuclei:
involvement of nucleoplasmin. Dev. Biol. *148*, 295–305. 10.1016/0012-
1606(91)90338-4.

47.   Dimitrov, S., and Wolffe, A.P. (1996). Remodeling somatic nuclei in Xenopus
laevis egg extracts: molecular mechanisms for the selective release of histones
H1 and H1(0) from chromatin and the acquisition of transcriptional competence.
EMBO J. *15*, 5897–5906. 10.1002/j.1460-2075.1996.tb00976.x.

48.   Ramos, I., Prado, A., Finn, R.M., Muga, A., and Ausió, J. (2005). Nucleoplasmin-
mediated unfolding of chromatin involves the displacement of linker-associated
chromatin proteins. Biochemistry *44*, 8274–8281. 10.1021/bi050386w.

49.   Bañuelos, S., Omaetxebarria, M.J., Ramos, I., Larsen, M.R., Arregi, I., Jensen,
O.N., Arizmendi, J.M., Prado, A., and Muga, A. (2007). Phosphorylation of both
nucleoplasmin domains is required for activation of its chromatin decondensation
activity. J. Biol. Chem. *282*, 21213–21221. 10.1074/jbc.M702842200.

50.   Platonova, O., Akey, I. V., Head, J.F., and Akey, C.W. (2011). Crystal structure
and function of human nucleoplasmin (Npm2): A histone chaperone in oocytes
and embryos. Biochemistry *50*, 8078–8089. 10.1021/bi2006652.

51.   Ramos, I., Martín-Benito, J., Finn, R., Bretaña, L., Aloria, K., Arizmendi, J.M.,
Ausió, J., Muga, A., Valpuesta, J.M., and Prado, A. (2010). Nucleoplasmin binds

histone H2A-H2B dimers through its distal face. J. Biol. Chem. *285*, 33771–
33778. 10.1074/jbc.M110.150664.

52.   Taneva, S.G., Bañuelos, S., Falces, J., Arregi, I., Muga, A., Konarev, P. V,
Svergun, D.I., Velázquez-Campoy, A., and Urbaneja, M.A. (2009). A mechanism
for histone chaperoning activity of nucleoplasmin: thermodynamic and structural
models. J. Mol. Biol. *393*, 448–463. 10.1016/j.jmb.2009.08.005.

53.   Onikubo, T., Nicklay, J.J., Xing, L., Warren, C., Anson, B., Wang, W.L., Burgos,
E.S., Ruff, S.E., Shabanowitz, J., Cheng, R.H., et al. (2015). Developmentally
regulated post-translational modification of nucleoplasmin controls histone
sequestration and deposition. Cell Rep. *10*, 1735–1748.
10.1016/j.celrep.2015.02.038.

54.   Lorton, B.M., Warren, C., Ilyas, H., Nandigrami, P., Hegde, S., Cahill, S., Lehman,
S.M., Shabanowitz, J., Hunt, D.F., Fiser, A., et al. (2023). Glutamylation of Npm2
and Nap1 acidic disordered regions increases DNA charge mimicry to enhance
chaperone efficiency., 10.1101/2023.09.18.558337 10.1101/2023.09.18.558337.

55.   Gadad, S.S., Senapati, P., Syed, S.H., Rajan, R.E., Shandilya, J., Swaminathan,
V., Chatterjee, S., Colombo, E., Dimitrov, S., Pelicci, P.G., et al. (2011). The
multifunctional protein nucleophosmin (NPM1) is a human linker histone H1
chaperone. Biochemistry *50*, 2780–2789. 10.1021/bi101835j.

56.   Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger,
M. (2022). ColabFold: making protein folding accessible to all. Nat. Methods *19*,
679–682. 10.1038/s41592-022-01488-1.

57.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.
10.1038/s41586-021-03819-2.

58.   Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein
sequence searching for the analysis of massive data sets. Nat. Biotechnol. *35*,
1026–1028. 10.1038/nbt.3988.

59.   Saluri, M., Leppert, A., Gese, G.V., Sahin, C., Lama, D., Kaldmäe, M., Chen, G.,
Elofsson, A., Allison, T.M., Arsenian-Henriksson, M., et al. (2023). A "grappling
hook" interaction connects self-assembly and chaperone activity of
Nucleophosmin 1. PNAS Nexus *2*, 1–9. 10.1093/pnasnexus/pgac303.

60.   Shintomi, K., Iwabuchi, M., Saeki, H., Ura, K., Kishimoto, T., and Ohsumi, K.
(2005). Nucleosome assembly protein-1 is a linker histone chaperone in Xenopus
eggs. Proc. Natl. Acad. Sci. U. S. A. *102*, 8210–8215. 10.1073/pnas.0500822102.

61.   Freedman, B.S., Miller, K.E., and Heald, R. (2010). Xenopus egg extracts
increase dynamics of histone H1 on sperm chromatin. PLoS One *5*, 1–10.
10.1371/journal.pone.0013111.

62.   Onikubo, T., Nicklay, J.J., Xing, L., Warren, C., Anson, B., Wang, W.-L., Burgos,

E.S., Ruff, S.E., Shabanowitz, J., Cheng, R.H., et al. (2015). Developmentally
Regulated Post-translational Modification of Nucleoplasmin Controls Histone
Sequestration and Deposition. Cell Rep. *10*, 1735–1748.
10.1016/j.celrep.2015.02.038.

63.   González-Arzola, K., Díaz-Quintana, A., Bernardo-García, N., Martínez-Fábregas,
J., Rivero-Rodríguez, F., Casado-Combreras, M.Á., Elena-Real, C.A., Velázquez-
Cruz, A., Gil-Caballero, S., Velázquez-Campoy, A., et al. (2022). Nucleus-
translocated mitochondrial cytochrome c liberates nucleophosmin-sequestered
ARF tumor suppressor by changing nucleolar liquid-liquid phase separation. Nat.
Struct. Mol. Biol. *29*, 1024–1036. 10.1038/s41594-022-00842-3.

64.   Frey, S., and Görlich, D. (2014). A new set of highly efficient, tag-cleaving
proteases for purifying recombinant proteins. J. Chromatogr. A *1337*, 95–105.
10.1016/j.chroma.2014.02.029.

65.   Keeble, A.H., Turkki, P., Stokes, S., Anuar, I.N.A.K., Rahikainen, R., Hytönen,
V.P., and Howarth, M. (2019). Approaching infinite affinity through engineering of
peptide-protein interaction. Proc. Natl. Acad. Sci. U. S. A. *116*, 26523–26533.
10.1073/pnas.1909653116.

66.   Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A., and Smith,
H.O. (2009). Enzymatic assembly of DNA molecules up to several hundred
kilobases. Nat. Methods *6*, 343–345. 10.1038/nmeth.1318.

67.   Howarth, M., Chinnapen, D.J.F., Gerrow, K., Dorrestein, P.C., Grandy, M.R.,
Kelleher, N.L., El-Husseini, A., and Ting, A.Y. (2006). A monovalent streptavidin
with a single femtomolar biotin binding site. Nat. Methods *3*, 267–273.
10.1038/nmeth861.

68.   Zhang, Z., Wang, Y., Ding, Y., and Hattori, M. (2020). Structure-based
engineering of anti-GFP nanobody tandems as ultra-high-affinity reagents for
purification. Sci. Rep. *10*, 1–10. 10.1038/s41598-020-62606-7.

69.   Kirchhofer, A., Helma, J., Schmidthals, K., Frauer, C., Cui, S., Karcher, A., Pellis,
M., Muyldermans, S., Casas-Delucchi, C.S., Cardoso, M.C., et al. (2010).
Modulation of protein properties in living cells using nanobodies. Nat. Struct. Mol.
Biol. *17*, 133–139. 10.1038/nsmb.1727.

70.   Fridy, P.C., Li, Y., Keegan, S., Thompson, M.K., Nudelman, I., Scheid, J.F.,
Oeffinger, M., Nussenzweig, M.C., Fenyö, D., Chait, B.T., et al. (2014). A robust
pipeline for rapid production of versatile nanobody repertoires. Nat. Methods *11*,
1253–1260. 10.1038/nmeth.3170.

71.   Zierhut, C., Jenness, C., Kimura, H., and Funabiki, H. (2014). Nucleosomal
regulation of chromatin composition and nuclear assembly revealed by histone
depletion. Nat. Struct. Mol. Biol. *21*, 617–625. 10.1038/nsmb.2845.

72.   Guse, A., Fuller, C.J., and Straight, A.F. (2012). A cell-free system for functional
centromere and kinetochore assembly. Nat. Protoc. *7*, 1847–1869.
10.1038/nprot.2012.112.

73.   Jenness, C., Giunta, S., Müller, M.M., Kimura, H., Muir, T.W., and Funabiki, H.
(2018). HELLS and CDCA7 comprise a bipartite nucleosome remodeling complex
defective in ICF syndrome. Proc. Natl. Acad. Sci. U. S. A. *115*, E876–E885.
10.1073/pnas.1717509115.

74.   Mastronarde, D.N. (2003). SerialEM: A Program for Automated Tilt Series
Acquisition on Tecnai Microscopes Using Prediction of Specimen Position.
Microsc. Microanal. *9*, 1182–1183. DOI: 10.1017/S1431927603445911.

75.   Arimura, Y., Tachiwana, H., Oda, T., Sato, M., and Kurumizaka, H. (2012).
Structural analysis of the hexasome, lacking one histone H2A/H2B dimer from the
conventional nucleosome. Biochemistry *51*. 10.1021/bi300129b.

76.   Lowary, P.T., and Widom, J. (1998). New DNA sequence rules for high affinity
binding to histone octamer and sequence-directed nucleosome positioning. J.
Mol. Biol. *276*, 19–42. 10.1006/jmbi.1997.1494.

77.   Han, Y., Fan, X., Wang, H., Zhao, F., Tully, C.G., Kong, J., Yao, N., and Yan, N.
(2020). High-yield monolayer graphene grids for near-atomic resolution
cryoelectron microscopy. Proc. Natl. Acad. Sci. U. S. A. *117*, 1009–1014.
10.1073/pnas.1919114117.

78.   Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard,
D.A. (2017). MotionCor2: Anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. Nat. Methods *14*, 331–332.
10.1038/nmeth.4193.

79.   Scheres, S.H.W. (2012). RELION: Implementation of a Bayesian approach to
cryo-EM structure determination. J. Struct. Biol. *180*, 519–530.
10.1016/j.jsb.2012.09.006.

80.   Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A.J., and Berger,
B. (2019). Positive-unlabeled convolutional neural networks for particle picking in
cryo-electron micrographs. Nat. Methods *16*, 1153–1160. 10.1038/s41592-019-
0575-8.

81.   Zivanov, J., Nakane, T., and Scheres, S.H.W. (2019). A Bayesian approach to
beam-induced motion correction in cryo-EM single-particle analysis. IUCrJ *6*, 5–
17. 10.1107/S205225251801463X.

82.   Waskom, M. (2021). Seaborn: Statistical Data Visualization. J. Open Source
Softw. *6*, 3021. 10.21105/joss.03021.

83.   Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. *9*,
90–95. 10.1109/MCSE.2007.55.

84.   Murray, A.W. (1991). Cell cycle extracts. Methods Cell Biol. *36*, 581–605.

85.   Desai, A., Murray, A., Mitchison, T.J., and Walczak, C.E. (1998). Chapter 20 The
Use of Xenopus Egg Extracts to Study Mitotic Spindle Assembly and Function in
Vitro. Methods Cell Biol. *61*, 385–412. 10.1016/S0091-679X(08)61991-3.

86.　Wynne, D.J., and Funabiki, H. (2015). Kinetochore function is controlled by a
phosphodependent coexpansion of inner and outer components. J. Cell Biol. *210*,
899–916. 10.1083/jcb.201506020.

87.　Lord, S.J., Velle, K.B., Dyche Mullins, R., and Fritz-Laylin, L.K. (2020).
SuperPlots: Communicating reproducibility and variability in cell biology. J. Cell
Biol. *219*. 10.1083/JCB.202001064.

88.　Wühr, M., Freeman, R.M., Presler, M., Horb, M.E., Peshkin, L., Gygi, S.P., and
Kirschner, M.W. (2014). Deep proteomics of the xenopus laevis egg using an
mRNA-derived reference database. Curr. Biol. *24*, 1467–1475.
10.1016/j.cub.2014.05.044.

89.　Shevchenko, A., Tomas, H., Havliš, J., Olsen, J. V., and Mann, M. (2007). In-gel
digestion for mass spectrometric characterization of proteins and proteomes. Nat.
Protoc. *1*, 2856–2860. 10.1038/nprot.2006.468.

90.　Peshkin, L., Lukyanov, A., Kalocsay, M., Gage, R.M., Wang, D., Pells, T.J.,
Karimi, K., Vize, P.D., Wühr, M., and Kirschner, M.W. (2019). The protein
repertoire in early vertebrate embryogenesis. bioRxiv, 571174. 10.1101/571174.

91.　Silva, J.C., Gorenstein, M. V., Li, G.Z., Vissers, J.P.C., and Geromanos, S.J.
(2006). Absolute quantification of proteins by LCMSE: A virtue of parallel MS
acquisition. Mol. Cell. Proteomics *5*, 144–156. 10.1074/mcp.M500230-MCP200.

92.　Tang, G., Peng, L., Baldwin, P.R., Mann, D.S., Jiang, W., Rees, I., and Ludtke,
S.J. (2007). EMAN2: an extensible image processing suite for electron
microscopy. J. Struct. Biol. *157*, 38–46. 10.1016/j.jsb.2006.05.009.

93.　Afonine, P. V., Poon, B.K., Read, R.J., Sobolev, O. V., Terwilliger, T.C.,
Urzhumtsev, A., and Adams, P.D. (2018). Real-space refinement in PHENIX for
cryo-EM and crystallography. Acta Crystallogr. Sect. D Struct. Biol. *74*, 531–544.
10.1107/S2059798318006551.

94.　Lugmayr, W., Kotov, V., Goessweiner-Mohr, N., Wald, J., DiMaio, F., and
Marlovits, T.C. (2023). StarMap: a user-friendly workflow for Rosetta-driven
molecular structure refinement. Nat. Protoc. *18*, 239–264. 10.1038/s41596-022-
00757-9.

95.　Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris,
J.H., and Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in
visualization and analysis. Protein Sci. *27*, 14–25. 10.1002/pro.3235.

1778

1779

Figure 1. Single particle cryo-EM analysis of poly-nucleosomes attached to magnetic beads (A) Schematic of a pilot cryo-EM experiment on magnetic beads. Biotin-labeled 19-mer nucleosome arrays attached to 50 nm streptavidin-coated magnetic nanobeads were loaded onto the cryo-EM grid. **(B)** Representative medium magnification micrographs. The magnetic beads are seen as black dots (red arrows). **(C)** Left; a representative high magnification micrograph. The micrograph was motion-corrected and low-pass filtered to 5 Å resolution. Right; green circles indicate the nucleosome-like particles selected by Topaz, and the blue areas indicate the halo-like scattering. **(D)** The 3D structure of the nucleosome bound on magnetic beads.

1789

1790

Figure 2. MagIC-Cryo-EM structural determination of low-quantity and low-purity targets (A) Schematic depicting the principle steps of MagIC-cryo-EM. **(B)** Graphical representation of the MagIC-cryo-EM beads with 3HB and SAH spacers and GFP nanobody target capture module. **(C)** Schematic of MagIC-cryo-EM for *in vitro* reconstituted H1.8-GFP bound nucleosomes isolated from an excess of H1.8-free nucleosomes. **(D)** Native PAGE analysis of H1.8-GFP bound nucleosomes and unbound nucleosomes in the input. DNA staining by SYTO-60 is shown. **(E)** A handmade humidity chamber used for the 5 min incubation of the cryo-EM grids on the magnet. The humidity chamber was assembled using a plastic drawer. Wet tissues are

attached to the side walls of the chamber, which is sealed with a plastic cover to

maintain high humidity. Two pieces of neodymium magnets are stacked. A graphene

grid is held by a non-magnetic vitrobot tweezer and placed on the magnets. 4 µL of

sample is applied on the grid and incubated for 5 min. **(F)** Micrograph montage of the

grids without using magnetic concentration. The GFP-nanobody-MagIC-cryo-EM beads

(4 µL of 12.5 pM beads) were applied on the graphene-coated Quantifoil R 1.2/1.3 grid

and vitrified without incubation on a magnet. **(G)** Micrograph montage of the grids

without using magnetic concentration. The GFP-nanobody-MagIC-cryo-EM beads (4 µL

of 12.5 pM beads) were applied on the graphene-coated Quantifoil R 1.2/1.3 grid and

vitrified with 5 min incubation on two pieces of 40 x 20 mm N52 neodymium disc

magnets. **(H)** Quantitative analysis of the percentage of holes containing MagIC-cryo-

EM beads. Each data point represents the percentage of holes containing MagIC-cryo-

EM beads on each square mesh. **(I)** Quantitative analysis of the average number of

MagIC-cryo-EM beads per hole. Each data point represents the average number of

MagIC-cryo-EM beads per hole on each square mesh. The edges of the boxes and the

midline indicates the 25th, 50th, and 75th percentiles. Whiskers indicate the maximum

and lowest values in the dataset, excluding outliers. For the quantification, 11 square

meshes with 470 holes without magnetic concentration and 11 square meshes with 508

holes with 5 min incubation on magnets were used. **(J)** Representative motion corrected

micrographs of *in vitro* reconstituted H1.8-GFP nucleosomes captured by MagIC-cryo-

EM beads. The micrographs were low-pass filtered to 10 Å resolution. Green circles

indicate the nucleosome-like particles picked by Topaz. **(K)** 3D structure of the *in vitro*

reconstituted H1.8-GFP-bound nucleosome determined through MagIC-cryo-EM. The

pipeline for structural analysis is shown in Figure S2.

1824

Figure 3. MagIC-Cryo-EM structural determination of H1.8-bound nucleosomes from interphase and metaphase chromosomes in *Xenopus* egg extract. (A) Models of potential cell cycle-dependent H1.8 dynamic binding mechanisms **(B)** Experimental flow of MagIC-cryo-EM analysis for GFP-H1.8 containing complexes isolated from chromosomes assembled in interphase and metaphase *Xenopus* egg extract. Fluorescence microscopy images indicate localization of GFP-H1.8 to interphase and metaphase chromosomes. DNA and GFP-H1.8 were detected either by staining with Hoechst 33342 or GFP fluorescence, respectively. **(C)** Native PAGE of fragmented interphase and metaphase chromosome sucrose gradient fractions. GFP-H1.8 and DNA were detected with either GFP fluorescence or SYTO-60 staining, respectively. **(D)** Western blot of GFP-H1.8 in interphase and metaphase chromosome sucrose gradient fractions. GFP-H1.8 was detected using anti-GFP antibodies. **(E)** SDS-PAGE of the sucrose gradient fractions 4 and 5 shown in (C), demonstrating heterogeneity of the samples. Proteins were stained by gel code blue. Red arrows indicate the H1.8-GFP

bands. The full gel image is shown in Figure S4A. **(F)** *In silico* 3D classification of
interphase and metaphase H1.8-bound nucleosomes isolated from chromosomes in
Xenopus egg extract. To assess the structural variations and their population of H1.8-
bound nucleosomes, *ab initio* reconstruction and heterogenous reconstruction were
employed twice for the nucleosome-like particles isolated by the decoy classification.
The initial round of *ab initio* reconstruction and heterogenous reconstruction classified
the particles into three nucleosome-containing 3D models (A, B, C). Subsequent *ab*
initio reconstruction and heterogenous reconstruction on the class A, which has weak
H1.8 density, yielded three new nucleosome-containing structures, A1, A2, and A3. 3D
maps represent the structural variants of GFP-H1.8-bound nucleosomes. Red arrows
indicate extra densities that may represent H1.8. Green densities indicate on-dyad
H1.8. The bar graphs indicate the population of the particles assigned to each 3D class
in both interphase and metaphase particles (gray), interphase particles (blue), and
metaphase particles (red). The pipeline for structural analysis is shown in Figure S5A.
(G) Structures of H1.8-bound nucleosomes isolated from interphase and metaphase
chromosomes.

1856

Figure 4. MagIC-cryo-EM and DuSTER reconstructed cryo-EM structures of interphase-specific H1.8-bound NPM2. (A) Schematic of DuSTER workflow. **(B)** 2D classes before and after particle curation with DuSTER. More 2D classes are shown in Figure S10B-S10E. **(C)** 3D cryo-EM structure of interphase-specific H1.8-containing complex. C5 symmetry was applied during structural reconstruction. The complete pipeline is shown in Figures S8, S10, and S11. **(D)** MS identification of proteins that cofractionated with H1.8 in sucrose gradient fraction 4 from interphase chromosomes shown in Figure 3C. Portions of MagIC-cryo-EM beads prepared for cryo-EM were subjected to MS. Proteins shown in red are the proteins that comprise the GPF nanobody-MagIC-cryo-EM beads. Proteins shown in blue represent signals from H1.8-GFP. **(E)** Western blot of NPM2 in the sucrose gradient fractions of interphase and

1869   metaphase chromosome fragments. **(F)** The structural comparison of the crystal

1870   structure of the pentameric NPM2 core (PDB ID: 1K5J), and AF2 predicted structure of

1871   the pentameric NPM2 core, and MagIC-cryo-EM structures of NPM2-H1.8. The MagIC-

1872   cryo-EM structures indicate NPM2 in the NPM2-H1.8 complex forms pentamer.

1873

1874

Figure 5. Structural variations of NPM2 bound to H1.8. (A) Structural differences between the opened and closed forms of NPM2. Left panels show cryo-EM maps of the opened and closed forms of NPM2 with H1.8. Middle panels show the atomic models. The right panel shows the zoomed-in view of the open form (green) and closed form (gray) of the NPM2 protomer. In the closed form, β8 runs straight from the sepal side to the petal side. In the open form, the C-terminal portion of β8 is bent outward to the rim. **(B)** Putative H1.8 density (red arrow) in the averaged NPM2-H1.8 structure. **(C)** The NPM2 surface that contacts the putative H1.8 density (corresponding to aa 42-44) is shown in orange. The H1.8-binding sites are accessible in the open form while they are internalized in the closed form. Note that C-terminal acidic tracts A2 and A3 (Figure S13A) are not visible in the cryo-EM structure but are likely to contribute to H1.8 binding as well in both open and closed forms. **(D)** Model of the mechanism that regulates the amount of the H1.8 in interphase and metaphase nucleosome.

1888

1889

Figure 6. Advantages of MagIC-cryo-EM over conventional cryo-EM methods. (A) The on-bead-cryo-EM approach reduces preparation steps (for example, target isolation, enrichment, and buffer exchange), which can lead to sample loss. (B) Sample loss during the grid-freezing process is reduced by magnet-based enrichment of the targets on cryo-EM grids. (C) The magnetic beads are easily identified in medium - magnification montage maps, enabling the selection of areas where targets exist prior to high-magnification data collection. (D) Targets are highly concentrated around the beads, ensuring that each micrograph contains more than 100 usable particles for 3D structure determination.

1899