The Notch1/Delta-like-4 axis is crucial for the initiation and progression of oral

squamous cell carcinoma

Christian T. Meisel[1], Riccardo Destefani[1], Ilaria J. Valookkaran[1], Niels Rupp[2], Aashil

Batavia[2], Francesca Catto[3], Jacqueline Hofmann-Lobsiger[1], Thimios A. Mitsiadis[1], Cristina

Porcheri[1*]

[1] Institute of Oral Biology, Faculty of Medicine, University of Zurich, Plattenstrasse 11, 8032 Zurich, Switzerland

[2] Department of Pathology and Molecular Pathology, University Hospital Zurich, Rämistrasse 100, 8091 Zurich,

Switzerland

[3] IMAI-MedTech, Bio-technopark, Wagistrasse 18, 8952 Schlieren, Switzerland

* Correspondence: cristina.porcheri@zzm.uzh.ch

Abstract

The Notch signaling pathway is frequently altered in Oral Squamous Cell Carcinoma (OSCC), the

most common malignant neoplasm of the oral mucosa. In this study, we aimed to elucidate the

functional role of this pathway in OSCC initiation and progression. Using transgenic animal

models, advanced imaging, and next-generation-sequencing techniques, we analyzed Notch-

dependent changes driving OSCC. We found specific expression patterns of Notch1 and Delta-

like-4 confined to the malignant tissue, while Jagged1 was downregulated in OSCC. This mutually

exclusive expression of Delta-like-4 and Jagged1 occurs at the early hyperplastic stage and persists

until more advanced stages of the tumor. Transcriptomic analyses confirmed the dysregulation of

the Notch pathway circuitry associated with a genetic signature for undifferentiation in OSCC.

Furthermore, pharmacological Notch inhibition significantly impaired cancer cell motility. Taken

together, these results reveal the pivotal importance of the Notch1/Delta-like-4 signaling axis as a

central oncogenic driver in OSCC.

<u>Keywords</u>

Notch signaling pathway, oral cancers, squamous cell carcinoma, OSCC, keratinocyte

differentiation, Notch inhibitors, Notch ligands, Delta-like-4, Notch1, Jagged1

34 **List of abbreviations**

4NQO   – 4-nitroquinoline-1-oxide

CSL   – CBF, Suppressor of Hairless, Lag-1

Dll4   – Delta-like-4

DMSO   – Dimethyl sulfoxide

E-Cad   – E-cadherin

EMT   – Epithelial-to-mesenchymal-transition

GFP   – Green fluorescent protein

GO   – Gene Ontology

GSEA   – Gene set enrichment analysis

Hes   – Hairy and enhancer of split-1

Hey   – Hairy and enhancer of split related with YPRW motif protein

HNSCC   – Head and neck squamous cell carcinoma

Jag1   – Jagged1

K5/10/14   – Keratin 5/10/14

MesoSPIM   – Mesoscale selective plane illumination microscopy

OSCC   – Oral squamous cell carcinoma

RBPj   – Recombining binding protein suppressor of hairless

Runx1   – Runt-related transcription factor 1

Sox2   – SRY-Box Transcription Factor 2

TAM   – Tamoxifen

Main

Head and neck squamous cell carcinoma (HNSCC) is the sixth most common type of cancer,

annually affecting 850.000 people worldwide [1–3]. Although modern multimodal therapy has

improved clinical management of this disease, tumor recurrence and distal metastasis correlate

with a poor prognosis, with a median overall survival of ten months [4,5]. HNSCC regroups a variety

of cancers arising from the oral mucosa, nasal cavities, pharynx, and larynx, which have their own

unique carcinogenic evolution and clinical score [6].

In normal mucosal tissues, such as the oral and respiratory epithelia, keratinocytes undergo a well-

defined differentiation process as they move from the basal layer to the surface. In the basal layer,

stem-cell-like cells express undifferentiated markers such as SRY-box transcription factor 2

(SOX2), P63, and Keratin 5 and 14 (K5, K14). As these cells progress upward they begin to

differentiate, producing keratins and junctional and structural proteins essential for the formation

of the mucosal barrier [7]. In SCC, the normal keratinocyte differentiation process is disrupted,

leading to uncontrolled proliferation and impaired differentiation [8]. Basal keratinocytes bypass

typical differentiation stages and continue to proliferate in an undifferentiated or partially

differentiated state. This results in the accumulation of abnormal, poorly differentiated cells that

resist cell death and shedding.

Within tumors, cancer-initiating cells retain self-renewal capabilities, driving tumor initiation and

exhibiting resistance to conventional therapies [9]. These cells share undifferentiated markers with

epithelial stem cells, while the expression of keratins indicative of terminally differentiated

squamous cells (e.g., K10) is reduced or altered, reflecting the dedifferentiation character of tumor

cells.

Oral Squamous Cell Carcinoma (OSCC) originates from the mucosa of the oral cavity, with the

tongue being the most commonly affected site. OSCC develops progressively, starting with local

hyperplasia of squamous epithelial cells and advancing through dysplastic and invasive stages [10,3].

In squamous dysplasia, clusters of basaloid nonkeratinizing and poorly differentiated cells grow

on the epithelial surface as exophytic growth (e.i. Carcinoma in Situ) [11]. Aberrant epithelial cells

may infiltrate the neighboring muscles and spread along the mesenchymal lines to metastasize to

cervical lymph nodes [4,8]. Proximal metastases occur in up to 50% of patients and are associated

with the recurrence and development of distant metastases [6,12,13]. OSCC onset often relates to

chronic exposure to nitrosamines, polycyclic aromatic hydrocarbons, and other genotoxic

compounds (e.g. quinoline derivatives, arylamines), which in turn leads to genetic instability and

the dysregulation of the DNA repair process[10]. Clones with acquired genetic alterations have an

initial growth advantage and can potentially transform into aggressive subclones in the more

advanced stages of the disease [14,15,16]. High-risk premalignant lesions can be subtle, and timely

recognition relies on the detection of stage-specific molecular biomarkers.

Mutational profiling consistently identified chromosomal alterations resulting in the amplification

of *cyclin D1* and EGFR regions, while genetic mutations affecting tongue epithelium have been

mapped in cyclin-dependent kinase inhibitor 2A (CDKN2A), tumor protein 53 (TP53) [3,17,18].

Notably, the Notch signaling pathway is one of the major mutation hotspots in OSCC[19–21]. Exome

sequencing screenings revealed that approximately 15% of tumors harbor mutations in this

pathway, including truncations, insertions-deletions, and missense variants.

Notch signaling is an evolutionary conserved pathway involved in cell fate specification and

differentiation [22]. It plays a crucial role in maintaining epithelial integrity, particularly in the oral

cavity, where it regulates mucosal formation and homeostasis from embryonic development

onward [22–27]. In mammals, four Notch receptors (Notch1-4) and five Notch-ligands (Jagged1,2 and

Delta-like-1,3,4) have been identified [28,29]. Upon receptor-ligand binding, a series of cleavages

take place to ultimately release the Notch intracellular domain (NICD), which then translocates

into the nucleus and acts as a transcription factor in a complex with CSL (also known as RBPj)

[22,30]. The activity of Notch is strictly context-dependent, therefore spatial distribution of each

receptor, ligand and downstream targets reflects in a distinct modulation of the pathway, hence

influencing tissue homeostasis [31]. Dysregulation of Notch signaling is found in several human

pathologies, and novel therapeutic approaches have been developed in recent years to specifically

target Notch components [31,32]. Although frequently mutated in OSCC[19–21], the precise role of

Notch in the initiation and development of this malignancy remains largely unknown, with

contrasting results that are supportive of either an oncogenic or a tumor-suppressive role [33–37].

Using *ad hoc* murine models, we have explored the functional role of Notch signaling in the

progression of OSCC, from initiation to the most advanced aggressive stages. Our findings reveal

a mutually exclusive expression of Dll4 and Jag1 that is consistently maintained across all stages

of tumor growth, from precancerous lesions to advanced invasive forms. Transcriptomic analysis

further confirmed the downregulation of Notch-specific target genes, alongside dysregulation of

genes involved in keratinocyte differentiation and cell-to-cell and cell-to-extracellular matrix

(ECM) communication. In parallel, pharmacological modulation of Notch-specific transcription

in human cancer cells led to a reduction in the malignant phenotype. Together, our data underscore

the critical role of the Notch1/Dll4 signaling axis in OSCC initiation and progression and

substantiate its suitability as both an early diagnostic tool and a functional target for OSCC

treatment.

Materials and Methods

Animals

C57Bl/6 and $Notch1^{CreERT}$:$R26mTmG^{fl/fl}$ mice were used for this study [38,39],The mice were

maintained and handled according to the Swiss Animal Welfare Law. This study was approved by

the Cantonal Veterinary office, Zurich (License ZH197/2017 and ZH086/2021).

4NQO treatment of mice – OSCC murine model

For the induction of OSCC, treated mice received drinking water containing 50μg/mL 4NQO

(#N8141, Sigma-Aldrich) starting at week 6 until week 22 of age. Control mice were littermates

receiving drinking water containing the same volume of DMSO (#D4540, Sigma-Aldrich). The

drinking solution was renewed twice a week. Females and males were equally represented in the

experimental set-up. After 16 weeks mice were either euthanized (time point 22 weeks) or given

normal drinking water until euthanasia (time point 32 weeks or 38 weeks). For lineage tracing

Notch1-driven expression, Cre was activated by the intraperitoneal injection of tamoxifen

50mg/kg dissolved in corn oil/10% EtOH (Sigma-Aldrich Cat#T5648) for two consecutive days.

All analyses were conducted in groups of 3 to 5 mice (n ≥ 3).

Sample embedding and cryo-sectioning

After tissue isolation, samples were post fixed in 4% Paraformaldehyde (PFA) for 1h at 4°C and

cryopreserved in 30% sucrose. Tissues were embedded in a cryo-embedding matrix (Tissue-Tek,

O.C.T Compound, Sakura) and stored at -80°C until sectioning. Samples were sectioned (10μm)

in the cryostat (Leica CM3050S) and stored at -80°C until staining.

Hematoxylin & Eosin staining

Sections were washed for 3x5min in 1x PBS. Nuclei were stained with Hematoxylin for 20 seconds

(Sigma-Aldrich #51275). Cytoplasm was then stained with Eosin for 10 seconds (Sigma-Aldrich

#HT110216), thoroughly washed and mounted for imaging. Images were acquired using the Leica

DMC5400 microscope.

Immunostaining of cryosections

Immunofluorescence staining of cryosections was performed as previously described[40]. Briefly,

Cryosections (10–25 μm thick) were post-fixed with 4% PFA and permeabilized with a 0.5%

Triton/PBS solution for 30 minutes at room temperature. To minimize nonspecific antibody

binding, slides were then incubated for 1 hour at RT in a blocking buffer containing species-

specific serum and 0.1% Triton in PBS. Primary antibodies were diluted in the blocking buffer at

the following concentrations: polyclonal rabbit anti-Keratin14 (1:100; Poly19053, Biolegend),

polyclonal goat anti-E-Cadherin (1:200, AF748, R&D Systems), polyclonal goat anti-GFP (1:100;

ab6673; Abcam), polyclonal rabbit anti-DLL4 (1:100; ab7280; Abcam), polyclonal goat anti-

DLL4 (1:50; AF1389; R&D) polyclonal rabbit anti-Jag1 (1:100; ab7771; Abcam), polyclonal

rabbit anti-Keratin10 (1:100; PRB-159P; Biolegend), polyclonal Rabbit anti-Keratin5 (dilution

1:100; ab64081; Abcam) polyclonal rabbit anti-Notch1 (D1E11) (dilution 1:100; 3608S; Cell

Signaling), polyclonal rabbit anti-Runx1 (dilution 1:100; HPA004176; Atlas Antibodies),

polyclonal rabbit anti-p63 (dilution 1:100; ab53039; Abcam), monoclonal rabbit anti-Vimentin

(dilution 1:100; ab92547; Abcam), mouse anti-Snail (1:100, #14-9859-80 Invitrogen). Used

secondary antibodies: Goat anti-Rabbit IgG 488, Alexa Fluor Plus 488 (dilution 1:1000; A32731,

Invitrogen), goat anti-mouse IgG 647, Alexa Fluor Plus 647 (dilution 1:1000; A32728, Invitrogen),

donkey anti-Goat IgG 488, Alexa Fluor Plus 488 (dilution 1:1000; A32814; Invitrogen); donkey

anti-Rabbit IgG 647, Alexa Fluor 647 (dilution 1:1000; A32795: Invitrogen), streptavidin Alexa

Fluor 488 (dilution 1:100; S11223; Invitrogen). Nuclear counterstaining was performed using

DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride, #D1306, Invitrogen).

Image acquisition – Confocal microscopy

Immunofluorescence images of the tongue tissue were produced using a Leica DM6000B

microscope, or Confocal Leica SP5/ SP8 with the following settings: 1024x1024 pixel dimension;

400Hz laser; frame average 3 or 4. photomultipliers PMT or Hyd. Fluorochromes used: Alexa 488,

546, 594, 647. A range of 10μm (standard) to 70μm (whole mount) thick Z-stack was analyzed

collecting images every 2μm (standard) or 10μm (whole mount). Images were processed using

Imaris 9.9 software (Bitplane). Photomerge Script from Photoshop (Adobe, Version 23.0.2) was

used for overview images reconstruction.

Tissue clearance

The separated epithelial sheet was embedded in 2% low-melting agarose. The tissue was then

transferred into Histodenz (#D2158, Sigma-Aldrich), dissolved in PBS1X and diluted until a

measurable refractive index of 1.457 for tissue clearance.

Whole-organ 3D imaging

Whole tongue images were recorded with a custom-made selective plane illumination microscope

(mesoSPIM) [41]. SPIM imaging was done after clearing and refractive index matching. The

laser/filter combinations for mesoSPIM imaging were as follows: for Notch1-GFP at 488 nm

excitation, a 405-488-561-640 nm QuadrupleBlock bandpass filter was used as the emission filter;

for membrane localized dt-Tomato at 561 nm excitation, a 561 nm long pass filter was used.

Transparent tongue epithelial whole-mount was imaged at a voxel size of 3.26 × 3.26 × 3 μm (X

× Y × Z). Further technical details of the mesoSPIM have been previously reported [41]

(*mesoSPIM.org*).

Enzymatic separation of keratinized epithelium from mesenchyme of the murine tongue

Separation was performed as previously described [42]. In brief, an enzymatic cocktail of collagenase

A (1mg/mL; Roche) and Dispase II (2mg/mL; #D4693 Sigma-Aldrich;) in 0.1 M PBS1X was

injected into the subepithelial space from the posterior cutting end of the tongue. Upon incubation

at 37°C for 30 min the mesenchyme and epithelium were pulled in opposing directions until

complete separation is achieved.

Oral mucosal organoid establishment and treatment

Oral mucosal organoids were established and propagated as previously described [43]. Briefly,

following enzymatic separation of the tongue epithelium, a keratinocyte cell suspension was

obtained and cultured in Advanced DMEM^{+++} supplemented with B27 (Life Technologies,

#17504-044), 1.25 mmol/L N-acetyl-L-cysteine and 10 mmol/L nicotinamide. Every three days

freshly prepared mRSPO1 (100 ng/mL, PeproTech), mEGF (50 ng/mL,) , and 10 ng/mL FGF10

was added. Organoids were passaged every 8 days by dissociating them into single cells using a

combination of trypsin digestion and mechanical disruption via pipetting. Trypsin was inactivated

using 10% FBS for 10 minutes at 37 °C, after which cells were centrifuged and resuspended in

complete organoid medium with growth factors. For treatment, either CB103 (100 μM;

MedChemExpress, #HY-135145) or an equivalent volume of DMSO (Sigma-Aldrich, #D8418)

was added to the organoid medium. Treated single cells were imaged every day, and organoid size

was quantified using FIJI/ImageJ software. Quantitative analyses were performed on 40 organoids

per condition.

RNA isolation epithelial tissue

Separated epithelial sheets were snap-frozen with liquid nitrogen and homogenized on ice for 3min

using a motor driven grinder (Pellet Pestle Motor, Kontes). Further isolation according to

manufacturer's instructions (RNeasy Mini Kit; Qiagen). Quantification of the total RNA extracted

by NanoDrop (Thermo Fischer Scientific, Waltham, USA). RNA was retrotranscribed using the

iScript™cDNA synthesis kit (Bio-Rad Laboratories AG, Cressier FR, Switzerland) according to

manufacturer's instructions.

Quantitative real-time PCR

The quantitative real-time PCR was performed using the CFX Connect Real-Time System and the

software CFX Manager Version 3.1. Expression level analysis of *Actin* (housekeeping gene), and

all candidate genes were performed using the SYBR® Green PCR Master Mix (Applied

Biosystems, Carlsbad CA, USA) in combination with corresponding oligonucleotide primers

(Supplementary Table 1 and 2).

All amplifications were done in technical duplicates of 5 biological replicates. Using the following

thermocycling conditions: 95°C for 10min; 40 cycles of 95°C for 15sec; 60°C for 60sec. Melting

curve analysis was performed at 95°C for 15sec, 60°C for 5sec and 95°C for 30sec. Expression

levels were calculated using the ΔΔCt method ($2^{-\Delta\Delta Ct}$ formula), after being normalized to the Ct-

value of the *Actin* housekeeping gene.

Bioinformatic analysis

Library for bulk seqRNA was prepared using Illumina NovaSeq 6000. Differentially expressed

genes were obtained from the DEseq2 output (p.adj <= 0.05) (Supplementary Method 1 and

Supplementary Method 2). We determine pathway enrichment using the ReactomePA v1.38.0

package [44]. We produce a similarity matrix using the Jaccard distance between the enriched terms

needed to produce the enrichment map using the R package enrichplot v1.14.1.

Raw data available on NCBI/SRA database (https://www.ncbi.nlm.nih.gov/sra/PRJNA1067290

and https://www.ncbi.nlm.nih.gov/sra/PRJNA1187878).

Patients' samples

Samples from patients diagnosed with HPV-negative HNSCC were used under the license BASEC

2018-01729. Tissues from patients 18-90 years old with primary tumor occurrence were collected

upon primary surgical therapy performed at the University Hospital Zurich (USZ). Tissues were

processed, embedded in OCT and sectioned according to USZ SOPs. For immunohistochemistry

staining, primary antibody α-Jagged1 (1:100; Abcam, ab7771) and α-Dll4 (1:100; R&D Systems,

AF1389) were used.

Cell culture

The SCC-25 epithelial cell line, derived from the tongue of a 70 years-old OSCC patient, was

purchased from the ATCC/LGC company (SCC-25CRL-1628™) and maintained accordingly to

producer indications. Briefly, cells were cultured in DMEM/F12 medium containing 1.2 g/L

sodium bicarbonate, 2.5 mM L-glutamine, 15 mM HEPES, 0.5 mM sodium pyruvate (# 11330032,

Gibco-ThermoFisher) and 10% fetal bovine serum (#16000044, Gibco-ThermoFisher). Cells were

sub-cultured when 80% confluent using trypsin/EDTA (#15400054, Gibco-ThermoFisher) and

frozen at 10^6 cells/vial in 10% DMSO (#D4540, Sigma-Aldrich). All cells used in our experiments

are mycoplasma-free.

Scratch assay

For scratch assay, SCC25 cells were grown to confluence. A central scratch of 1mm of distance

was produced using a silicon scraper. Medium was changed immediately with conditioned

medium: Control group medium contained the anti-mitotic Aphidicolin (1ug/ml; # A4487 Sigma-

Aldrich) and DMSO (#D8418; Sigma-Aldrich); Treated group contained Aphidicolin (1ug/ml; #

A4487 Sigma-Aldrich) and CB103 (100μM; # HY-135145 MedChemExpress). Fresh conditioned

medium was used for medium changes every two days for 10 days. Images were taken at 20X

every day at the same time of the day using a Leica DMIL LED inverted microscope mounting a

Leica camera DFC295. Four wells per conditions were analyzed. Quantification of cell-free areas

was performed using FIJI/ImageJ free software.

Cell immunofluorescence

Treated SCC25 cells were plated on hatched coverslips at 20.000cells/well. Cells were fixed with

PFA4% for 15min at 4°C. After PBS washing, cells were permeabilized 30 min in 0.1% triton and

then blocked using a solution of 10% FBS, 0,1% triton in PBS. Primary antibodies were diluted in

blocking buffer and incubated overnight. Primary antibodies a-Vinculin 1:500 (#90227 Millipore)

was used for immunofluorescence on cells and detected via secondary antibody donkey anti-mouse

IgG 546, Alexa Fluor 546 (dilution 1:1000; Invitrogen). Nuclear counterstaining was performed

using DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride, #D1306, Invitrogen).

TRITC-conjugated Phalloidin staining of actin filaments was performed as indicated by the

manufacturer (1:1000 for 1h incubation at room temperature; Millipore # 90228). Mounting in

Mowiol 4-88/Glycerol (#81381 and #G-6279, Sigma-Aldrich).

Time-lapse imaging

Low passage SCC25 cells were treated with CB103 (100μM; # HY-135145 MedChemExpress) or

DMSO (#D8418; Sigma-Aldrich) for 4 days and then plated at 10000 cells/well in an 8-wells μ-

Slide high glass bottom (#80807, Ibidi). Cells were observed under an Olympus ScanR HCS at

20X with controlled environmental conditions (CO_2 5% and 37°C) for 3h, taking an image every

3 minutes. For analysis, Imaris 9.9 (Bitplane) spot and tracking function was used to automatically

quantify the displacement length.

297
298

Results

Physiological Distribution and Localization of Notch1, Dll4, and Jag1 in the Tongue

Epithelium

To investigate how the Notch signalling pathway is activated in OSCC, we initially sought to

examine receptor-ligand interactions in the keratinized epithelium of the tongue, the primary site

of OSCC development. Since different receptor-ligand combinations can either activate or inhibit

Notch signaling, identifying the specific interactions present in normal tissue is a critical first step

in understanding how the pathway is regulated in the oral mucosa.

To map Notch1 receptor activity in vivo, we used an inducible transgenic reporter mouse model,

Notch1^{CreERT}:R26mTmG$^{fl/fl}$, which allows for mGFP labeling of Notch1-expressing cells following

tamoxifen administration (Figure 1A). This system enabled us to monitor Notch1 activation over

time and to track the behavior and contribution of labeled cells as the tissue transitions from a

healthy state to advanced stages of OSCC.

Using light-sheet microscopy (mesoSPIM), we generated a 3D reconstruction of the entire tongue

epithelium and observed Notch1-driven mGFP expression in both ventral and dorsal regions

(Figure 1B, Movie 1). Cross-sections of adult tongues provided more detailed views of Notch1

localization within the stratified epithelium, confirming its broad distribution across both surfaces

(Figure 1C). In *Notch1^{CreERT}:R26mTmG$^{fl/fl}$* mice, Notch1:mGFP-positive cells were restricted to

the multilayered tongue epithelium, while unrecombined regions retained expression of the

mTomato cassette (Figure 1D). Both endogenous Notch1 protein and Notch1-driven mGFP were

detected from the basal layer up to the stratum granulosum (Figures 1C, 1D).

We then examine the spatial distribution of Notch-ligands Jagged1 (Jag1) and Delta-like-4 (Dll4)

in the tongue epithelium. We identified a compartmentalized expression pattern with mutually

exclusive localization (Figures 1E and 1F). Jag1 was present on both the dorsal and ventral sides

of the tongue, with staining in the spaces between filiform papillae of the dorsal region (Figure

1E). In contrast, Dll4 was exclusively localized in the epithelium of the dorsal tongue, specifically

on the outer surface of the filiform papillae (Figure 1F). This complementary expression pattern

of the two ligands suggests a potentially distinct role for Notch1-Jagged1 versus Notch1-Dll4

interactions in the physiology of tongue keratinocytes.

Mutually exclusive distribution patterns of Dll4 and Jag1 in the tongue epithelium of OSCC

To model the progressive stages of tongue squamous cell carcinoma, we administered the

carcinogenic substance 4-Nitroquinoline N-oxide (4NQO) in the drinking water of 6-weeks-old

mice for a duration of 16 weeks (Supplementary Figure 1A) [45]. As previously reported, chronic

exposure to 4NQO for 16 weeks induces hyperplasia and dysplasia in the tongue epithelium [46].

After an additional 10-week washout period, the tissue progresses to carcinoma with 100%

penetrance. Histological analysis of 22-week-old mice revealed epithelial hyperplasia,

leukoplakia, and hyperkeratinization, primarily on the dorsal side of the tongue and occasionally

on the ventral side (Supplementary Figures 1A, 1B, 1C). By 32 weeks of age, histological evidence

of carcinoma in situ was observed on the dorsal epithelium of the tongue (Supplementary Figures

1B, 1C) [46]

To study the role of Notch in OSCC initiation, we firstly treated $Notch1^{CreERT}$:$R26mTmG^{fl/fl}$ mice

with 4NQO or DMSO and analyzed the tissue at 22 weeks of age (Figures 2A). In 4NQO-treated

mice, we observed specific Notch-driven mGFP expression localized to the hyperplastic

epithelium (Figure 2B), indicating active Notch signaling within the cell population driving

malignant transformation. Next, we assessed the distribution of the Notch ligands Dll4 and Jag1

at this early stage. Dll4 staining was restricted to the hyperplastic epithelium (Figure 2C), while

Jag1 immunolabelling was notably absent in this same region (Figure 2D, white arrows and dashed

area). Interestingly, Jag1 expression displayed a gradual decline: it remained at physiological

levels in unaffected tongue regions distant from the epithelial overgrowth (Figure 2D, red arrows),

decreased progressively in the tissue bordering the hyperplastic area (Figure 2D, yellow arrows),

and became undetectable within the tumor's central core (Figure 2D, white arrows).

To determine whether the exophytic growth exhibited undifferentiated characteristics at this early

stage of the disease, we performed labeling for the undifferentiated markers Sox2 and K5,

alongside the mature keratinocyte marker K10 [47,48]. Sox2 and K5 were highly expressed in the

exophytic mass, whereas K10 was absent, indicating that the growing mass predominantly consists

of undifferentiated progenitor cells (Figure 2E).

We next aimed to determine whether the Notch1-Dll4 signaling axis remains active throughout

OSCC progression or is restricted to the tumor initiation phase. To address this, we analyzed

4NQO-treated *Notch1CreERT:R26mTmGfl/fl* mice at a later time point (32-weeks-old; Figure

3A). While DMSO-treated mice showed a similar pattern to the one observed in 22-weeks-old

tissue (Figure 3B, upper panel), carcinoma in situ lesions exhibited widespread Notch1-driven

mGFP labeling (Figure 3B, lower panel). Consistent with our previous observations, Jag1

expression was largely absent from the tumor mass but persisted in the adjacent tissue (Figures

3C), whereas strong Dll4 staining was detected within the exophytic tumor growth (Figure 3D).

Thus, the Notch1-Dll4 interaction appears to be a constant throughout the course of OSCC

development, from initiation to advanced lesions.

367 **Notch1-Dll4 signature correlate with maintenance of undifferentiation and early epithelial**

368 **to mesenchymal transition.**

As the Notch signaling have been implicated in the maintenance of stem cell identity in a variety

of tissues[49], we explored whether the presence of Notch1-Dll4 signaling correlates with

undifferentiation in OSCC lesions. To this end, 32-week-old mice tissue was stained for the

progenitor markers Sox2 and K5. In 4NQO-treated mice, these markers label the majority of

malignant cells, while in DMSO-treated controls, Sox2 and K5 staining was restricted to the basal

epithelial layer of the tongue (Supplementary Figures 2A and 2B) [50,51]. Conversely, the

differentiation marker K10 was confined in the cornified layer of the healthy tongue epithelium

(Supplementary Figure 2C), while remaining undetectable in the 4NQO-induced exophytic

growth. Finally, malignant epithelial cells retain the expression of the pan-epithelial marker E-

Cadherin (E-Cad)[52] (Supplementary Figure 2D). Taken together, these results suggest impaired

differentiation and an expansion of progenitor-like cells within the 4NQO-induced tumor

(Supplementary Figure 2C).

Next, we investigated whether inhibiting Notch in tongue keratinocytes directly affects stem cell

behavior [67]. To assess this, we performed an organoid formation assay using epithelial stem cells

isolated from the tongue. In control cultures treated with DMSO, organoids consistently formed

within 5 days, with an average diameter of 122.43 μm ± 36.9. However, in cultures treated with

the Notch inhibitor CB103, organoid formation was completely disrupted: Cells remained as single

cells or small clusters, with an average diameter of 20.21 μm ± 7.19 (DMSO vs. CB103,

$p \leq 0.0001$). These results suggest that Notch signaling is essential for maintaining the stem cell

functions required for organoids initiation and growth (Figure 4A).

To gain insight into the molecular changes driving malignant transformation, we performed further

analyses at the 32-weeks-old stage. After separating the tongue's outer epithelium from the

underlying connective tissue [42] total RNA was extracted for downstream analyses (Figure 4B).

qRT-PCR on 4NQO-treated epithelium showed an important increase in *Dll4* expression compared

to DMSO-treated epithelium (Figure 4C). Interestingly, the *Hes*-family of target genes remained

largely unaffected, whereas *Hey1* and *Hey2* (Hairy and enhancer of split-related with YPRW motif

protein) were significantly reduced in 4NQO-treated epithelia (Figure 4C). To obtain a

comprehensive view of the transcriptional changes, we additionally performed bulk RNA

sequencing analyses. Gene Set Enrichment Analysis (GSEA) revealed a significant enrichment of

the Notch signaling signature (Figure 4D), which was further supported by the volcano plot

highlighting a pronounced upregulation of *Dll4* in 4NQO-treated samples (Figure 4F). We

identified 3331 differentially expressed genes in 4NQO-treated versus DMSO-treated epithelia (p-

value threshold p≤ 0.01; Supplementary Method 1). The heatmap shows changes in genes mainly

involved in cell differentiation processes (Cluster 2 and Cluster 3), molecular pathways regulating

cell fate determination (Cluster 1 and Cluster 5), extracellular matrix rearrangement (Clusters

1,2,3,4), and cell motility (Cluster 3) (Figure 4E, Supplementary Table 3). Additional GSEA

showed changes in keratinocyte differentiation and cornification (Figure 4G and Supplementary

Figure 3). These results align with our phenotyping data, which show that the Notch1-Dll4 axis is

associated with the maintenance of an undifferentiated state.

We then conducted interactome analyses to examine changes in the tongue's epithelia upon 4NQO

and DMSO treatments. Independent *Enrichr (https://maayanlab.cloud/Enrichr/)* analyses

identified alteration of *Runt-related transcription factor 1 (Runx1)* amongst others (e.g., *p21, p300*

and *Stat3*) (Supplementary Figure 4). Importantly, this alteration is specific to *Runx1*, as other

members of the *Runx* family were not significantly altered (Supplementary Figure 4B). As *Runx1*

has been previously linked to Notch pathway activation[53–55], we wanted to demonstrate the

distribution of the Runx1 protein in the tumorigenic epithelium of the tongue. Indeed, a strong

Runx1 staining was detected in the tumor mass of the 4NQO-treated mice (Supplementary Figure

5A), while the Runx1 labelling was confined to the stratum basale and stratum spinosum of the

DMSO-treated mice. Furthermore, strong immunostaining against p63, a marker of

undifferentiated keratinocytes[56], was detected in the tumor mass of 4NQO-treated mice

(Supplementary Figure 5B) [53,57–62] . Together, these findings suggest that Notch1-Dll4 signaling

is closely linked to the expansion of progenitor-like cell populations in OSCC lesions, supporting

its role in maintaining an undifferentiated state during malignant transformation.

At more advanced stages of OSCC progression, malignant cells residing within the exophytic mass

may become invasive by undergoing epithelial-to-mesenchymal transition (EMT) [63]. Using light-

sheet microscopy (SPIM), we analyzed 4NQO-treated mice to identify the time window when

basal lamina dissolution begins. In 22-week-old mice, digital sectioning revealed an uninterrupted

basal lamina underneath the epithelial overgrowth (Supplementary Figure 6A, yellow arrows;

Movie 2). In 32-week-old mice, we observed a localized area of basal lamina degradation, allowing

the malignant overgrowth to directly interface with the underlying mesenchyme (Supplementary

Figure 3B, white arrows; Movie 3).

To investigate the molecular changes that may contribute to the dissolution of the basal lamina

observed in SPIM imaging, we examined our sequencing data. We identified an EMT signature

through GSEA, along with changes in anchoring junctions (Figure 4H). Gene Ontology analysis

for biological processes and cellular components further confirmed alterations in cell-cell and cell-

to-ECM communication, as well as cell migration (Figure 4I). To validate EMT onset at this stage,

we assessed the expression of the mesenchymal marker Vimentin (*VIM*), components of the

extracellular matrix Tenascin (*TNC*), and members of the zinc-finger transcription factor family

Snail. Transcriptomic analyses confirmed their upregulation, together with the early EMT markers

ZEB1, ZEB2 (Figure 4L and 4M).

While technical limitations of the 4NQO model prevent us from capturing the full extent of the

EMT process and metastasis formation, our data indicate that EMT initiation occurs at the 32-

week-old stage following 4NQO exposure, with changes in tissue architecture, molecular

signatures, and the upregulation of key markers of cellular transformation.

Invasive squamous cells conserve their dependency from the Notch1-Dll4 signaling axis

To confirm the progression toward an invasive phenotype, we analyzed a more advanced stage of

our OSCC model at 38-weeks-old. Tongue sections from *R26:mTomato* mice were stained with

the epithelial markers K14 and P63. This labeling revealed clusters of epithelial cells embedded

within the stromal tissue of the tongue (Figures 5B and 5C). These findings led us to hypothesize

that carcinoma in situ cells had breached the basal lamina and begun invading the underlying

connective tissue. As confirmation that EMT was occurring at this stage, we examined the

expression patterns of Snai1 and the adhesion molecule regulator E-cadherin [64,65]. Snail-positive

staining was detected in cells within the invasive tumor mass of 4NQO-treated mice (Figure 5D,

white arrows), whereas such labeling was absent in DMSO-treated controls. We then assessed the

distribution of Vimentin and E-cadherin, markers of mesenchymal and epithelial identity,

respectively [65,66]. A subset of cells within the tumor mass coexpressed vimentin and E-cadherin

(Figure 5E, yellow arrows), thus confirming an ongoing EMT process.

We therefore sought to clarify if the Notch1-Dll4 signaling axis is also involved in the process of

tumor invasion observed in 38-weeks-old mice. *Notch1^{CreERT}:R26mTmG$^{fl/fl}$* mice were used to trace

Notch1-expressing cells and their progeny (Figure 6A). In DMSO-treated mice, *mGFP* expression

remained restricted to the epithelial layers (Figure 6B). Conversely, in 4NQO-treated mice, mGFP-

labeled cells were observed within the carcinoma mass invading the mesenchyme (Figure 6C). We

next performed immunolabeling for the Notch1 protein. A widespread Notch1 signal labeled the

infiltrating mass of 4NQO-treated mice, while in DMSO-treated mice Notch1 expression remained

confined to the healthy tongue epithelium (Figure 6D). Finally, we examined the expression of

Dll4 in the invasive aberrant epithelium using *R26:mTomato* mice. Analysis of DMSO-treated

mice confirmed the expression of Dll4 in the dorsal portion of the filiform papillae, as previously

observed (Figure 6E and 1F). In 4NQO-treated mice, labeling for Dll4 was broadly present in the

infiltrating mass (Figure 6E). Our findings suggest a conserved paradigm governing tumor

progression, where the Notch1-Dll4 axis remains central throughout the phases of hyperplastic

growth, carcinoma in situ and the most advanced, invasive stages of the disease.

The Notch signaling pathway regulates malignancy in human OSCC

To evaluate the clinical relevance of the Notch pathway in OSCC, we examined the expression of

JAG1 and DLL4 in human tissue samples. We first performed Hematoxylin and Eosin (H&E)

staining on tissues from HPV-negative adult patients with primary tumors. This allowed us to

identify characteristic SCC features, including dysplastic epithelium, mitotic figures, and

abnormal keratin accumulation (Figure 7, overview). Based on these histological criteria, we

distinguished healthy tissue (Figure 7A) from hyperplastic (Figure 7D) and carcinoma samples

(Figure 7G).

Jag1 and Dll4 expression were then compared across these different tissue types. DLL4 showed

diffuse expression in the healthy epithelium (Figure 7B), with increased levels observed in

hyperplastic tissue (Figure 7E) and consistently high expression in carcinoma nests (Figure 7H).

In contrast, JAG1 staining in healthy tissue was limited to small, localized pockets, and its

expression varied across different regions of the carcinoma samples (Figure 7C and Figure 7F,7I).

To further investigate the molecular regulation of the Notch pathway in the human context, we

treated hSCC25 cell line (#CRL-1628, ATCC/LGC) with CB103, a pan-blocker of the Notch-

transcriptional activation complex[67] (Figure 8A). Interestingly, quantitative RT-PCR revealed a

specific downregulation of the *DLL4* expression, while *JAG1* levels remain largely unchanged

(Figure 8B). Additionally, CB103 treatment resulted in decreased expression of *SOX2* and *P63*

(Figure 8B), suggesting a direct link between Notch activity and the maintenance of an

undifferentiated state.

To further investigate the molecular impact of CB103 treatment, we performed bulk seqRNA

comparing DMSO- versus CB103-treated hSCC25 cells. This analysis identified 3329

differentially expressed genes (p≤ 0.01; Supplementary Method 2), with the top 2000 genes

visualized in a hierarchical clustering heatmap (Figure 8D). The heatmap highlights alterations in

genes involved in epithelial differentiation (Cluster 3, 4 and Cluster 5), cell-to-cell adhesion

(Cluster 3), cytoskeleton organization (Cluster 4, Cluster 5) and chemotaxis (Cluster 1) (Figure

8D, Supplementary Table 4). GO analysis for cellular components and biological processes, along

with *Enrichr* analyses, further supported these findings by identifying alterations in intermediate

filaments, cell-to-cell adhesion, and cell-to-ECM interactions (Figure 8C, Supplementary Figure

7A and Supplementary Figure 7B). Paralleling our findings in the mouse model, GSEA analysis

revealed significant changes in keratinization processes and keratinocytes differentiation (Figure

8E). Additionally, MA and volcano plot analysis indicated an upregulation in the expression of

keratins (Figure 8F), as well as a change in the expression of integrins and ECM molecules

(Supplementary Figure 7C). Together with GSEA results (Figure 8G-I), these data suggest that

CB103-treated cells undergo structural modifications and remodeling of anchoring molecules.

We hypothesized that such changes could affect cell motility, a crucial factor in acquiring an

invasive phenotype during cancer progression. We therefore aimed to explore the functional

consequences of CB103 treatment. Among the differentially expressed signatures, we identified

enrichment in genes associated with desmosomal complexes and apical junctions (Figures 9A, 9B

and 9D). qRT-PCR validation confirmed a significant upregulation of desmosome-associated

proteins upon CB103 treatment, including desmoplakin, desmoglein, and desmocollin (Figure 9C).

Consistently, CB103-treated cells exhibited a marked increase in the number of adhesive

complexes compared to DMSO-treated controls (Figures 9E and 9F)[68,69], reinforcing the role of

Notch inhibition in enhancing cell–cell adhesion. Finally, we conducted scratch assays and

monitored the progression of migrating cells over a period of 10 days (Figures 9G and 9H).

DMSO-treated cells were able to fully reform the monolayer within a ten-day timeframe, in

contrast to CB103-treated cells, which were unable to recolonize the cell-free zone (Figure 9G).

To further confirm the reduced motility of CB103-treated cells, we performed time-lapse analysis.

CB103-treated cells exhibited significantly reduced motility when compared to cells treated with

DMSO (Figures 9I and 9J; Movies 4 and 5). These results indicate that selective Notch inhibition

by CB103 markedly alters the expression of structural components within adhesive complexes that

anchor epithelial cells to the extracellular matrix, ultimately reducing the migration of human

carcinoma cells.

Overall, our findings from both murine and human models highlight the critical role of the Notch1-

Dll4 axis in maintaining undifferentiation and promoting motility. Disrupting this pathway

effectively impairs key drivers of OSCC malignancy.

Discussion

The Notch signaling pathway is a central regulator of carcinogenic transformation in a variety of

epithelial tumors, including lung, cervical, colon, and pancreatic cancers [70–73]. The dysregulation

of Notch is context-dependent and can shift from an activator to an inhibitor, depending on the

organ and the type of tissue affected [74–76]. Currently, a debate exists on whether the Notch pathway

acts as an oncogene or tumor suppressor in OSCC, and its exact role in this type of cancer remains

largely unknown [19,77,78]. Since Notch activity is strongly influenced by the specific receptor-ligand

interactions within a tissue, our first goal was to identify how these combinations evolve across

healthy tissue, initiation, and advanced stages of OSCC. To achieve this, we used the

Notch1CreERT:R26mTmGfl/fl transgenic line and studied Notch1-expressing cells and their

progeny throughout the progressive stages of malignant transformation. Short tamoxifen pulses

allowed precise control of the reporter activation, providing information on which cells had active

Notch1 at specific time points. Pulse-and-chase experiments instead, tracked labelled cells over

extended periods, revealing their movements and dynamic behavior as the cancer progresses and

its structure evolves. Through detailed phenotypic characterization, our findings suggest that

Notch signaling contributes to OSCC development and progression by preferentially interacting

with the Dll4 ligand rather than Jag1. Gene expression analyses confirmed that *Dll4* is the most

significant gene of the Notch pathway to be differentially regulated in OSCC condition. The

preferential binding of Notch1 to Dll4 in the tumorigenic epithelium ultimately leads to the

downregulation of *Hey1* and *Hey2*, while the expression of other Notch-specific target genes, such

as *Hes1* and *Hes2,* remains largely unaffected. The mutually exclusive expression of the Jag1 and

Dll4 ligands regulates fate definition and undifferentiation in a variety of tissues (e.g., during

angiogenesis and hematopoiesis)[40,79,80]. Similarly in our system, the selection between the Notch1-

Dll4 versus Notch1-Jag1 interaction observed in OSCC coincides with sustained keratinocyte

undifferentiation. The Notch1/Dll4-expressing keratinocyte progenitors retain their immature

state, resulting in the expansion of the undifferentiated portion of the tongue epithelium.

Important downstream changes occur during Notch pathway dysregulation in OSCC. Our

transcriptomic analyses reveal a complex scenario of crosstalk between pathways sustaining oral

carcinoma. Interestingly, p63 and Runx1 cooperate with the Notch pathway, similarly to what has

been found in other tissues. In skin keratinocytes a bidirectional crosstalk between Notch and p63

suggest a complex regulation of differentiation occurring at various levels (from regulation of

transcription to indirect regulation of the Notch pathway). A high expression of P63 correlates

with poor prognosis in patients with OSCC of the tongue[53,54,64,81–83][75]. Our model showed a similar

correlation, with an undifferentiated mass coherently expressing P63 and Notch1-Dll4. In HNSCC

screenings, the transcription factor RUNX1 was found to be significantly increased and thus, has

been suggested as a prognostic marker [83,84]. RUNX1 upregulation drives tumor progression and

metastasis, while suppression of RUNX1 inhibits HNSCC cell proliferation and migration [84,84,85].

In other settings, such as mammalian hematopoiesis, Runx1 controls the expansion, specification

and self-renewal of stem cells in a Notch-dependent manner [86] [87,88]. We have found a similar

scenario in our system, where both transcriptomic approaches and immunostainings show a

widespread expression of Runx1 and Notch1-Dll4 with an analogous expression pattern.

Continuous epithelial growth gives rise to a malignant mass with the potential of invading the

underling stroma[11]. In the invasive growth, we identified typical features of EMT, such as basal

lamina dissolution, overexpression of ZEB1, ZEB2 and Snail1 and cells co-expressing Vimentin

and E-cadherin. Transcriptomic analysis confirmed a specific hallmark signature for EMT in the

epithelium of 4NQO-treated mice. Importantly, cells of epithelial origin growing within the

mesenchyme maintain the expression of Dll4. Consequently, we identified the Notch-Dll4 axis

throughout all phases of OSCC progression, spanning from the initial hyperplastic phase to the

most advanced invasive stages, highlighting its promising potential as a diagnostic tool.

To strengthen the translational impact of our findings, we analyzed human malignant tissue and

confirmed elevated expression of DLL4. Dll4 blockers have been proposed for the treatment of

other epithelial tumors (e.g., intestinal tumors) primarily aiming to limit angiogenesis [89]. Since

OSCC growth is not strictly dependent on vascularization, a locally administered blocker could

directly target malignant epithelial cells in this tumor. Immunotherapy specifically targeting Dll4

combined to chemotherapy has been applied to treat solid tumors (e.g., pancreatic and lung

cancers) [90–93], with positive results spanning from a reduction of tumor growth to a decrease in the

reoccurrence rate [94–96]. Given the redundancy of the Notch pathway and the frequent activation of

compensatory mechanisms when a single component is targeted, we hereby adopted a broader

strategy to test the therapeutic potential of pan-Notch blockers in OSCC. The Notch

pharmacological inhibitor CB103 is increasingly gathering attention in cancer treatments, in view

of its high stability and unbeaten specificity in targeting the Notch pathway [67,97,98]. Several clinical

trials are currently in place to determine its usage as a therapeutic agent in carcinomas (clinical

trials: NCT03422679, NCT04714619) as well as leukemias and lymphomas (clinical trials:

NCT03422679, NCT05464836). In our system, SCC25 cells treated with CB103 displayed a

significant reduction in *DLL4* expression, with no observable change in *JAG1* expression,

confirming the crucial role of the NOTCH-DLL4 signaling axis in OSCC. Importantly, the

expression of undifferentiation markers SOX2 and P63 is significantly reduced in CB103-treated

cells, while transcriptomic analysis reveals induction of keratinization. These *in vitro* results

validate the *in vivo* findings in mice and denote the connection between an active NOTCH-Dll4

signaling axis and the prolonged undifferentiation cell state within OSCC. We finally tested the

functional effects of Notch blockage in OSCC cell behavior. We recorded a dramatic reduction in

cell migration when SCC25 cells were treated with CB103. This is accompanied by an increase in

adhesive complexes (focal adhesions and desmosomes), suggesting that NOTCH regulates cell

movement and is essential for the acquisition of tumor's invasive properties.

In conclusion, the present findings show that the Notch1-Dll4 signaling axis plays a crucial role in

the initiation and development of OSCC, contributing to the acquisition of critical oncogenic

features.

Acknowledgements

611 This work was supported by the University of Zurich, the Krebsliga (KFS-4890-08-2019-R) (T.A.M.), the Julius Klaus-Stiftung (C.P.), Stiftung Julius Müller zur Unterstützung der Krebsforschung (C.P.) and the Olga Mayenfisch Stiftung (C.P.). We thank Prof. Spyros Artavanis-Tsakonas (Harvard Medical School, Harvard University) for providing *Notch1^{creER}* mice.

616 Imaging was performed with equipment maintained by the Center for Microscopy and Image Analysis, University of Zurich. The authors gratefully acknowledge the Functional Genomics Center Zurich (FGCZ) of University of Zurich and ETH Zurich for the support on analyses.

References

References

1. Ferlay, J. *et al.* Estimating the global cancer incidence and mortality in 2018: GLOBOCAN

sources and methods. *Int J Cancer* **144**, 1941–1953 (2019).

2. Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and

mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*

68, 394–424 (2018).

3. Johnson, D. E. *et al.* Head and neck squamous cell carcinoma. *Nat Rev Dis Primers* **6**, 92

(2020).

4. Cramer, J. D., Burtness, B., Le, Q. T. & Ferris, R. L. The changing therapeutic landscape of

head and neck cancer. *Nat Rev Clin Oncol* **16**, 669–683 (2019).

5. Duprez, F. *et al.* Distant metastases in head and neck cancer: Distant metastases in head and

neck cancer. *Head & Neck* **39**, 1733–1743 (2017).

6. Pisani, P. *et al.* Metastatic disease in head & neck oncology. *Acta Otorhinolaryngol Ital* **40**,

S1–S86 (2020).

7. Eckert, R. L. & Rorke, E. A. Molecular biology of keratinocyte differentiation. *Environ*

Health Perspect **80**, 109–116 (1989).

8. Franchi, A. & Skalova, A. Undifferentiated and dedifferentiated head and neck carcinomas.

Semin Diagn Pathol **38**, 127–136 (2021).

9. Zhou, H.-M., Zhang, J.-G., Zhang, X. & Li, Q. Targeting cancer stem cells for reversing

therapy resistance: mechanism, signaling, and prospective agents. *Signal Transduct Target*

Ther **6**, 62 (2021).

10. Müller, S. Update from the 4th Edition of the World Health Organization of Head and Neck

Tumours: Tumours of the Oral Cavity and Mobile Tongue. *Head Neck Pathol* **11**, 33–40

(2017).

11. Pai, S. I. & Westra, W. H. Molecular pathology of head and neck cancer: implications for

diagnosis, prognosis, and treatment. *Annu Rev Pathol* **4**, 49–70 (2009).

12. Byers, R. M. *et al.* Frequency and therapeutic implications of 'skip metastases' in the neck

from squamous carcinoma of the oral tongue. *Head Neck* **19**, 14–19 (1997).

13. Kuperman, D. I. *et al.* Squamous cell cancer of the head and neck with distant metastasis at

presentation. *Head Neck* **33**, 714–718 (2011).

14. Califano, J. *et al.* Genetic progression and clonal relationship of recurrent premalignant head

and neck lesions. *Clin Cancer Res* **6**, 347–352 (2000).

15. Puram, S. V. *et al.* Cellular states are coupled to genomic and viral heterogeneity in HPV-

related oropharyngeal carcinoma. *Nat Genet* **55**, 640–650 (2023).

16. Bai, Y. *et al.* YBX1 integration of oncogenic PI3K/mTOR signalling regulates the fitness of

malignant epithelial cells. *Nat Commun* **14**, 1591 (2023).

17. Stransky, N. *et al.* The Mutational Landscape of Head and Neck Squamous Cell Carcinoma.

Science **333**, 1157–1160 (2011).

18. Sequeira, I. *et al.* Genomic landscape and clonal architecture of mouse oral squamous cell

carcinomas dictate tumour ecology. *Nat Commun* **11**, 5671 (2020).

19. Agrawal, N. *et al.* Exome sequencing of head and neck squamous cell carcinoma reveals

inactivating mutations in NOTCH1. *Science* **333**, 1154–1157 (2011).

20. Fan, W.-L. *et al.* Prognostic Genetic Biomarkers Based on Oncogenic Signaling Pathways

for Outcome Prediction in Patients with Oral Cavity Squamous Cell Carcinoma. *Cancers*

(Basel) **13**, 2709 (2021).

21. Loganathan, S. K. *et al.* Rare driver mutations in head and neck squamous cell carcinomas

converge on NOTCH signaling. *Science* **367**, 1264–1269 (2020).

22. Artavanis-Tsakonas, S. Notch Signaling: Cell Fate Control and Signal Integration in

Development. *Science* **284**, 770–776 (1999).

23. Casey, L. M. *et al.* Jag2-Notch1 signaling regulates oral epithelial differentiation and palate

development. *Dev. Dyn.* **235**, 1830–1844 (2006).

24. Mustonen, T. *et al.* Lunatic fringe, FGF, and BMP regulate the Notch pathway during

epithelial morphogenesis of teeth. *Dev Biol* **248**, 281–293 (2002).

25. Mitsiadis, T. A., Henrique, D., Thesleff, I. & Lendahl, U. Mouse Serrate-1 (Jagged-1):

expression in the developing tooth is regulated by epithelial-mesenchymal interactions and

fibroblast growth factor-4. *Development* **124**, 1473–1483 (1997).

26. Mitsiadis, T. A., Lardelli, M., Lendahl, U. & Thesleff, I. Expression of Notch 1, 2 and 3 is

regulated by epithelial-mesenchymal interactions and retinoic acid in the developing mouse

tooth and associated with determination of ameloblast cell fate. *Journal of Cell Biology* **130**,

407–418 (1995).

27. Porcheri, C. & Mitsiadis, T. A. Notch in Head and Neck Cancer. *Adv Exp Med Biol* **1287**,

81–103 (2021).

28. Artavanis-Tsakonas, S., Rand, M. D. & Lake, R. J. Notch signaling: cell fate control and

signal integration in development. *Science* **284**, 770–776 (1999).

29. Kopan, R. & Ilagan, M. X. G. The canonical Notch signaling pathway: unfolding the

activation mechanism. *Cell* **137**, 216–233 (2009).

30. Gordon, W. R., Arnett, K. L. & Blacklow, S. C. The molecular logic of Notch signaling--a

structural and biochemical perspective. *J Cell Sci* **121**, 3109–3119 (2008).

31. Siebel, C. & Lendahl, U. Notch Signaling in Development, Tissue Homeostasis, and Disease.

Physiological Reviews **97**, 1235–1294 (2017).

32. Louvi, A. & Artavanis-Tsakonas, S. Notch and disease: A growing field. *Seminars in Cell &*

Developmental Biology **23**, 473–480 (2012).

33. Yoshida, R. *et al.* The pathological significance of Notch1 in oral squamous cell carcinoma.

Lab. Invest. **93**, 1068–1081 (2013).

34. Setoguchi. Upregulation of Notch pathway molecules in oral squamous cell carcinoma. *Int J*

Oncol **36**, (2010).

35. Sakamoto, K. *et al.* Reduction of NOTCH1 expression pertains to maturation abnormalities

of keratinocytes in squamous neoplasms. *Laboratory Investigation* **92**, 688–702 (2012).

36. Sawangarun, W. *et al.* Loss of Notch1 predisposes oro-esophageal epithelium to

tumorigenesis. *Experimental Cell Research* **372**, 129–140 (2018).

37. Natsuizaka, M. *et al.* Interplay between Notch1 and Notch3 promotes EMT and tumor

initiation in squamous cell carcinoma. *Nat Commun* **8**, 1758 (2017).

38. Fre, S. *et al.* Notch lineages and activity in intestinal stem cells determined by a new set of

knock-in mice. *PLoS One* **6**, e25785 (2011).

39. Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-fluorescent

Cre reporter mouse. *Genesis* **45**, 593–605 (2007).

40. Porcheri, C. *et al.* Notch ligand Dll4 impairs cell recruitment to aortic clusters and limits

blood stem cell generation. *EMBO J.* **39**, e104270 (2020).

41. Voigt, F. F. *et al.* The mesoSPIM initiative: open-source light-sheet microscopes for imaging

cleared tissue. *Nat Methods* **16**, 1105–1108 (2019).

42. Meisel, C. T., Pagella, P., Porcheri, C. & Mitsiadis, T. A. Three-Dimensional Imaging and

Gene Expression Analysis Upon Enzymatic Isolation of the Tongue Epithelium. *Front*

Physiol **11**, 825 (2020).

43. Driehuis, E. *et al.* Oral Mucosal Organoids as a Potential Platform for Personalized Cancer

Therapy. *Cancer Discov* **9**, 852–871 (2019).

44. Yu, G. & He, Q.-Y. ReactomePA: an R/Bioconductor package for reactome pathway

analysis and visualization. *Mol. BioSyst.* **12**, 477–479 (2016).

45. Ishida, K. *et al.* Current mouse models of oral squamous cell carcinoma: Genetic and

chemically induced models. *Oral Oncol* **73**, 16–20 (2017).

46. Bouaoud, J. *et al.* The 4-NQO mouse model: An update on a well-established in vivo model

of oral carcinogenesis. *Methods Cell Biol* **163**, 197–229 (2021).

47. Sanz-Gómez, N. *et al.* Squamous differentiation requires G2/mitosis slippage to avoid

apoptosis. *Cell Death Differ* **27**, 2451–2467 (2020).

48. Sanz-Gómez, N. *et al.* Squamous differentiation requires G2/mitosis slippage to avoid

apoptosis. *Cell Death Differ* **27**, 2451–2467 (2020).

49. Bigas, A. & Porcheri, C. Notch and Stem Cells. *Adv Exp Med Biol* **1066**, 235–263 (2018).

50. Vasca, V. *et al.* Keratin 5 expression in squamocellular carcinoma of the head and neck.

Oncol Lett **8**, 2501–2504 (2014).

51. Alam, H., Sehgal, L., Kundu, S. T., Dalal, S. N. & Vaidya, M. M. Novel function of keratins

5 and 14 in proliferation and differentiation of stratified epithelial cells. *Mol Biol Cell* **22**,

4068–4078 (2011).

52. Huntsman, D. G. & Caldas, C. Assignment1 of the E-cadherin gene (CDH1) to chromosome

16q22.1 by radiation hybrid mapping. *Cytogenet Cell Genet* **83**, 82–83 (1998).

53. Burns, C. E., Traver, D., Mayhall, E., Shepard, J. L. & Zon, L. I. Hematopoietic stem cell

fate is established by the Notch-Runx pathway. *Genes Dev.* **19**, 2331–2342 (2005).

54. Burns, C. E. & Zon, L. I. Notch Signaling Is Necessary and Sufficient for Runx1-Dependent

Stem Cell Induction in the Embryonic Aorta-Gonad-Mesonephros (AGM) Region. *Blood*

104, 4161–4161 (2004).

55. Islam, R. *et al.* RUNX1 colludes with NOTCH1 to reprogram chromatin in T cell acute

lymphoblastic leukemia. *iScience* **26**, 106795 (2023).

56. Crum, C. P. & McKeon, F. D. p63 in epithelial survival, germ cell surveillance, and

neoplasia. *Annu Rev Pathol* **5**, 349–371 (2010).

57. Srivastava, S. S. *et al.* Keratin 5/14-mediated cell differentiation and transformation are

regulated by TAp63 and Notch-1 in oral squamous cell carcinoma-derived cells. *Oncol Rep*

39, 2393–2401 (2018).

58. Dotto, G. P. Crosstalk of Notch with p53 and p63 in cancer growth control. *Nat Rev Cancer*

9, 587–595 (2009).

59. Moses, M. A. *et al.* Molecular Mechanisms of p63-Mediated Squamous Cancer

Pathogenesis. *Int J Mol Sci* **20**, E3590 (2019).

60. Ortt, K., Raveh, E., Gat, U. & Sinha, S. A chromatin immunoprecipation screen in mouse

keratinocytes reveals Runx1 as a direct transcriptional target of DeltaNp63. *J Cell Biochem*

104, 1204–1219 (2008).

61. Nakagawa, M. *et al.* AML1/Runx1 rescues Notch1-null mutation-induced deficiency of para-

aortic splanchnopleural hematopoiesis. *Blood* **108**, 3329–3334 (2006).

62. Kouwenhoven, E. N. *et al.* Transcription factor p63 bookmarks and regulates dynamic

enhancers during epidermal differentiation. *EMBO Rep* **16**, 863–878 (2015).

63. González, M. M. *et al.* Is cancer a pure growth curve or does it follow a kinetics of

dynamical structural transformation? *BMC Cancer* **17**, 174 (2017).

64. Villarejo, A., Cortés-Cabrera, A., Molina-Ortíz, P., Portillo, F. & Cano, A. Differential role

of Snail1 and Snail2 zinc fingers in E-cadherin repression and epithelial to mesenchymal

transition. *J Biol Chem* **289**, 930–941 (2014).

65. Thiery, J. P. Epithelial–mesenchymal transitions in tumour progression. *Nat Rev Cancer* **2**,

442–454 (2002).

66. Dongre, A. & Weinberg, R. A. New insights into the mechanisms of epithelial–mesenchymal

transition and implications for cancer. *Nat Rev Mol Cell Biol* **20**, 69–84 (2019).

67. Lehal, R. *et al.* Pharmacological disruption of the Notch transcription factor complex. *Proc*

Natl Acad Sci U S A **117**, 16292–16301 (2020).

68. Ilić, D. *et al.* Reduced cell motility and enhanced focal adhesion contact formation in cells

from FAK-deficient mice. *Nature* **377**, 539–544 (1995).

69. Geiger, B., Tokuyasu, K. T., Dutton, A. H. & Singer, S. J. Vinculin, an intracellular protein

localized at specialized sites where microfilament bundles terminate at cell membranes. *Proc*

Natl Acad Sci U S A **77**, 4127–4131 (1980).

70. Avila, J. L. & Kissil, J. L. Notch signaling in pancreatic cancer: oncogene or tumor

suppressor? *Trends Mol Med* **19**, 320–327 (2013).

71. Brzozowa-Zasada, M. *et al.* Notch and its oncogenic activity in human malignancies. *Eur*

Surg **49**, 199–209 (2017).

72. Tyagi, A., Sharma, A. K. & Damodaran, C. A Review on Notch Signaling and Colorectal

Cancer. *Cells* **9**, E1549 (2020).

73. Rodrigues, C., Joy, L. R., Sachithanandan, S. P. & Krishna, S. Notch signalling in cervical

cancer. *Exp Cell Res* **385**, 111682 (2019).

74. Dotto, G. P. Notch tumor suppressor function. *Oncogene* **27**, 5115–5123 (2008).

75. Yap, L. *et al.* The opposing roles of NOTCH signalling in head and neck cancer: a mini

review. *Oral Dis* **21**, 850–857 (2015).

76. Fukusumi, T. & Califano, J. A. The NOTCH Pathway in Head and Neck Squamous Cell

Carcinoma. *J Dent Res* **97**, 645–653 (2018).

77. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck

squamous cell carcinomas. *Nature* **517**, 576–582 (2015).

78. Johnson, D. E. *et al.* Head and neck squamous cell carcinoma. *Nat Rev Dis Primers* **6**, 92

(2020).

79. Benedito, R. *et al.* The notch ligands Dll4 and Jagged1 have opposing effects on

angiogenesis. *Cell* **137**, 1124–1135 (2009).

80. Gama-Norton, L. *et al.* Notch signal strength controls cell fate in the haemogenic

endothelium. *Nat Commun* **6**, 8510 (2015).

81. Nguyen, B.-C. *et al.* Cross-regulation between Notch and p63 in keratinocyte commitment to

differentiation. *Genes Dev* **20**, 1028–1042 (2006).

82. Ichikawa, M. *et al.* A role for RUNX1 in hematopoiesis and myeloid leukemia. *Int J*

Hematol **97**, 726–734 (2013).

83. Gao, L. *et al.* RUNX1 and the endothelial origin of blood. *Exp Hematol* **68**, 2–9 (2018).

84. Feng, X. *et al.* Elevated RUNX1 is a prognostic biomarker for human head and neck

squamous cell carcinoma. *Exp Biol Med (Maywood)* **246**, 538–546 (2021).

85. Loljung, L. *et al.* High expression of p63 is correlated to poor prognosis in squamous cell

carcinoma of the tongue. *J Oral Pathol Med* **43**, 14–19 (2014).

86. Loljung, L. *et al.* High expression of p63 is correlated to poor prognosis in squamous cell

carcinoma of the tongue. *J Oral Pathol Med* **43**, 14–19 (2014).

87. Liu, K. *et al.* RUNX1 promotes MAPK signaling to increase tumor progression and

metastasis via OPN in head and neck cancer. *Carcinogenesis* **42**, 414–422 (2021).

88. Feng, X. *et al.* Elevated RUNX1 is a prognostic biomarker for human head and neck

squamous cell carcinoma. *Exp Biol Med (Maywood)* **246**, 538–546 (2021).

89. Ridgway, J. *et al.* Inhibition of Dll4 signalling inhibits tumour growth by deregulating

angiogenesis. *Nature* **444**, 1083–1087 (2006).

90. Hidalgo, M. *et al.* A phase 1b study of the anti-cancer stem cell agent demcizumab (DEM)

and gemcitabine (GEM) +/- nab-paclitaxel in patients with pancreatic cancer. *JCO* **34**, 341–

341 (2016).

91. McKeage, M. J. *et al.* Phase IB Trial of the Anti-Cancer Stem Cell DLL4-Binding Agent

Demcizumab with Pemetrexed and Carboplatin as First-Line Treatment of Metastatic Non-

Squamous NSCLC. *Target Oncol* **13**, 89–98 (2018).

92. Coleman, M. L. *et al.* Asparaginyl hydroxylation of the Notch ankyrin repeat domain by

factor inhibiting hypoxia-inducible factor. *J Biol Chem* **282**, 24027–24038 (2007).

93. Johnson, M. *et al.* Abstract A081: A phase 1b, open-label, dose escalation and expansion

study of demcizumab plus pembrolizumab in patients with locally advanced or metastatic

solid tumors. in *Clinical Trials* A081–A081 (American Association for Cancer Research,

2018). doi:10.1158/1535-7163.TARG-17-A081.

94. Fischer, M. *et al.* Anti-DLL4 inhibits growth and reduces tumor-initiating cell frequency in

colorectal tumors with oncogenic KRAS mutations. *Cancer Res.* **71**, 1520–1525 (2011).

95. Yen, W.-C. *et al.* Anti-DLL4 Has Broad Spectrum Activity in Pancreatic Cancer Dependent

on Targeting DLL4-Notch Signaling in Both Tumor and Vasculature Cells. *Clin Cancer Res*

18, 5374–5386 (2012).

96. Smith, D. C. *et al.* A Phase I Dose Escalation and Expansion Study of the Anticancer Stem

Cell Agent Demcizumab (Anti-DLL4) in Patients with Previously Treated Solid Tumors.

Clinical Cancer Research **20**, 6295–6303 (2014).

97. Fabbro, D., Bauer, M., Murone, M. & Lehal, R. Notch Inhibition in Cancer: Challenges and

Opportunities. *Chimia (Aarau)* **74**, 779–783 (2020).

98. Perez Garcia, J. M. *et al.* First-in-human phase 1-2A study of CB-103, an oral Protein-

Protein Interaction Inhibitor targeting pan-NOTCH signalling in advanced solid tumors and

blood malignancies. *JCO* **36**, TPS2619–TPS2619 (2018).

837
838

Figures legends

Figure 1: Notch1, Dll4 and Jag1 expression in the epithelium of the mouse tongue.

A) Schematic representation illustrating the murine tongue of a $Notch1^{CreERT}:R26mTmG^{fl/fl}$ mouse

after tamoxifen induction and subsequent $mGFP$ expression. B) MesoSPIM imaging and related

three-dimensional reconstruction of a whole-mount $Notch1^{CreERT}:R26mTmG^{fl/fl}$ mouse tongue.

Notch1-driven $mGFP$ expression (green) in the raw reconstruction (left panels) and relative

surface rendering (right panels). C) Cross-sections of the dorsal (left panels) and ventral (right

panels) part of the tongue stained against the full length Notch1 protein (green). Cell nuclei are

stained with DAPI (blue). Asterisks indicate the magnified areas in the corresponding region of

the overview. D) Cross-sections showing the dorsal (left panel) and ventral (right panel) parts of

the tongue of a $Notch1^{CreERT}:R26mTmG^{fl/fl}$ mouse. Both the dorsal and the ventral epithelium

express the Notch1-driven mGFP (green). Unrecombined tissue expresses $mTomato$ (red). Cell

nuclei stained with DAPI (blue). Asterisks indicates magnified area in the corresponding region of

the overview. E) Cross-sections of the dorsal (left panel) and ventral (right panel) parts of the

tongue showing Jag1 staining (green). Cell nuclei are stained with DAPI (blue). F) Cross-sections

of the dorsal (left panel) and ventral (right panel) parts of the tongue showing Dll4 staining (green).

Nuclei are stained with DAPI (blue). Scale bars in overviews100µm. Scale bars in magnification

50µm.

Figure 2: The Notch ligand Jagged1 and Dll4 expression is mutually exclusive in the early stage of OSCC

(A) Schematic representation of the tamoxifen induction and the 4NQO or DMSO treatment in 22-week-old *Notch1CreERT:R26mTmGfl/fl* mice. (B) Cross-sections of the dorsal part of the tongue in DMSO-treated (upper panels) and 4NQO-treated (lower panels) mice. mTomato expression (red) and mGFP expression (green) in the healthy (DMSO) and aberrant (4NQO) epithelium of the tongue. Cell nuclei are stained with DAPI (blue). (a-c) indicate the magnified areas in the corresponding regions of the overview (a-c'). (C) Cross-sections of the dorsal part of the tongue stained against Dll4 (green) in DMSO- (upper panels) and 4NQO-treated (lower panels) mice. Cell nuclei are stained with DAPI (blue). (D) Cross-sections of the dorsal part of the tongue stained against Jag1 (green) in DMSO-treated (left panels) and 4NQO-treated (right panels) mice. Cell nuclei are stained with DAPI (blue). Note the gradient of the Jag1 staining in the 4NQO-treated epithelium: the epithelium distant from the lesion is Jag1-positive (red arrows), the staining is reduced in the epithelium close to the hyperplasia (yellow arrows) and is absent in the hyperplastic tissue (white arrows). The dotted line indicates the borders of the epithelial hyperplasia. (E) Cross-sections of the dorsal part of the tongue showing Sox2 staining (green, upper panels), K5 (green, middle panels) and K10 (green, lower panels) in the healthy (DMSO) and aberrant (4NQO) epithelium of the tongue. Cell nuclei are stained with DAPI (blue). (d-i) indicate the magnified areas in the corresponding regions of the overview (d'-i'). Scale bars in overviews100µm. Scale bars in magnification 50µm.

Figure 3: Dll4 is specifically expressed in the OSCC exophytic outgrowth

(A) Schematic representation of the 4NQO or DMSO treatment in 32-week-old *R26:mTomatofl/fl*

mice. B) Cross-sections of the dorsal part of the tongue in DMSO-treated (upper panels) and

4NQO-treated (lower panels) mice. *mTomato* expression (red) and *mGFP* expression (green) in

the healthy (DMSO) and aberrant (4NQO) epithelium of the tongue. Cell nuclei are stained with

DAPI (blue). Asterisks indicate the magnified areas in the corresponding regions of the overview.

(C) Cross-sections of the dorsal part of the tongue stained against Jag1(green) in DMSO- (left

panels) and 4NQO-treated (right panels) mice. Cell nuclei are stained with DAPI (blue). (D) Cross-

sections of the dorsal part of the tongue stained against Dll4 (green) in DMSO-treated (left panels)

and 4NQO-treated (right panels) mice. Cell nuclei are stained with DAPI (blue). The dotted line

indicates the borders of the OSCC overgrowth. Asterisks indicate the magnified areas in the

corresponding regions of the overview. Scale bars in overviews100μm. Scale bars in magnification

50μm.

Figure 4: Transcriptome analysis of differential gene expression in the tongue epithelium of

4NQO-treated mice

(A) Organoid formation assay using single epithelial stem cells derived from the tongue.

Treatment with CB103 inhibits organoid formation from single-cell suspensions, indicating

impaired stem cell function. Scale bar: 50 μm. (B) Schematic representation of the tongue's

epithelium separation for RNA extraction and follow up analysis. (C) qRT-PCR shows

upregulation of Dll4 expression and downregulation of Hey1 and Hey2 expression in 4NQO-

treated mice. Error bars indicate standard error. *=p<0.05; **=p<0.01; ***=p<0.001 (D) Gene Set

Enrichment Analysis (GSEA) plot for the Notch-signaling signature shows changes in the Notch

pathway in 4NQO condition when compared to DMSO. (E) Heatmap visualization of 2000

differentially expressed genes with hierarchical clustering. (F) Volcano plot confirm higher

expression of Dll4 in 4NQO over DMSO conditions. (G) Gene Set Enrichment Analysis (GSEA)

shows alterations in the process of keratinocytes differentiation and keratinization in 4NQO

condition. (H) GSEA identify alterations in the apical junction complex and EMT signatures (I)

Gene Ontology (GO) classification analysis for categories of biological processes (left panel) and

cellular components (right panel) comparing DMSO- versus 4NQO-treated epithelium. (L)

Volcano plot identified an increased expression of EMT markers Vimentin, ZEB1, ZEB2,

Tenascin C and Snail3. (M) qRT-PCR confirmation of EMT markers ZEB1 and ZEB2 upregulated

in 4NQO versus DMSO condition. Error bars indicate standard error. $*=p<0.05$; $**=p<0.01$;

$***=p<0.001$

Figure 5: Epithelial cells invade the tongue's mesenchyme in 4NQO-treated mice

(A) Schematic representation of the 4NQO or DMSO treatment in 38-week-old *R26:mTomatofl/fl*

mice. (B) Cross-sections displaying the dorsal part of the tongue labelled with K14 (cyan) in

DMSO-treated (left panels) and 4NQO-treated mice (right panels). Background tissue expressed

mTomato (red). Asterisks indicate the magnified areas in the corresponding regions of the

overview. (C) Cross-sections displaying the dorsal part of the tongue labelled with P63 (cyan) in

DMSO-treated (left panels) and 4NQO-treated (right panels) mice. Background tissue expressed

mTomato (red). Asterisks indicate the magnified areas in the corresponding regions of the

overview. (D) Cross-sections of the dorsal part of the tongue stained with Snail (cyan) in DMSO-

treated (left panel) and 4NQO-treated condition (right panels) mice. Cell nuclei are stained with

DAPI (blue). Asterisk indicates the magnified area in the corresponding region of the overview.

White arrows indicate Snail-positive cells. (E) Cross-sections of the dorsal part of the tongue

stained with Vimentin (cyan) and E-cadherin (green) in DMSO-treated (left panel) and 4NQO-

treated (right panels) mice. Cell nuclei are stained with DAPI (blue). Asterisks indicate the

magnified areas in the corresponding regions of the overview. White arrows indicate sole Vimentin

staining, while yellow arrows indicate staining for both Vimentin and E-cadherin. Scale bars in

overviews 100μm. Scale bars in magnification 50μm.

Figure 6: Detection of Notch1-expressing epithelial cells and their progeny within the

infiltrated mesenchyme of the tongue.

(A) Schematic representation illustrating the tamoxifen induction and 4NQO or DMSO treatment

of 38-week-old *Notch1^CreER^T:R26mTmG^fl/fl* mice. (B) Cross-sections displaying the Notch1-driven

expression of *mGFP* in DMSO-treated epithelium. Unrecombined tissue expressed *mTomato*

(red). Cell nuclei are stained with DAPI (blue). Asterisk indicates the magnified area in the

corresponding region of the overview. (C) Tongue cross-sections displaying the Notch1-driven

expression of *mGFP* in 4NQO-treated mice after ten days of tamoxifen induction. Unrecombined

tissue expressed *mTomato* (red). Cell nuclei are stained with DAPI (blue). Asterisks indicate the

magnified areas in the corresponding regions of the overview. (D) Cross-sections displaying the

endogenous Notch1-full length expression (cyan) in the tongue epithelium of DMSO-treated (left

panels) and 4NQO-treated (right panels) mice. Unrecombined tissue expressed *mTomato* (red).

Asterisks indicate the magnified areas in the corresponding regions of the overview. (E) Cross-

sections showing the distribution of Dll4 (green) in the tongue epithelium of DMSO-treated (left

panels) and 4NQO-treated (right panels) mice. Asterisk indicates magnified area in the

corresponding region of the overview. Dashed lines indicate the extent of the epithelial infiltration

in the musculature of the tongue. Scale bars in overviews 100μm. Scale bars in magnification

50μm.

Figure 7: Notch signaling in human squamous cell carcinoma

(A) Histological examination of human healthy epithelium. Hematoxylin/Eosin staining is used to

identify the histological condition of the samples analyzed. Corresponding areas are then stained

for Dll4 (B) and Jag1(C). Overview of expression for Dll4 (upper panel) and magnifications (lower

panels). Scale bar in overview 500μm. Scale bar in magnification 50μm. (D) Histological

examination of human hyperplastic epithelium. Hematoxylin/Eosin staining is used to identify the

histological condition of the samples analyzed. Corresponding areas are then stained for Dll4 (E)

and Jag1(F). Overview of expression (upper panel) and magnifications (lower panels). Scale bar

in overview 500μm. Scale bar in magnification 50μm. (G) Histological examination of human

squamous cell carcinoma tissue. Hematoxylin/Eosin staining is used to identify aberrant histology

in two representative patients (Patient#1 with hyperplastic epithelium: middle panel, Patient#2

with squamous carcinoma: right panel). (H) Distribution of Dll4 in Patient#1 and Patient#2.

Magnified areas show expression of Dll4 protein. Scale bar in magnification: 100μm. Scale bar in

overview: 1mm. (I) Distribution of Jag1 protein in Patient#1 and Patient#2 in overview and

magnified areas. Scale bar in magnification: 100μm. Scale bar in overview: 1mm.

Figure 8: CB103 treatment leads to broad transcriptional reprogramming in human

squamous cell carcinoma cells

(A) Schematic overview of the experimental setup: SCC25 human squamous cell carcinoma cells

were treated with the pan-Notch inhibitor CB103 or DMSO (control) for five days prior to

downstream analyses. (B) qRT-PCR analysis shows that CB103 treatment significantly decreases

expression of the Notch ligand DLL4 and the stemness-associated markers SOX2 and P63. (C)

Bar plot summarizing Gene Ontology (GO) classification of cellular components, comparing gene

expression profiles between CB103- and DMSO-treated SCC25 cells.

(D) Heatmap with hierarchical clustering of the top 500 differentially expressed genes identified

via bulk RNA sequencing of SCC25 cells treated with CB103 or DMSO. (E) Gene Set Enrichment

Analysis (GSEA) plots show enrichment of keratinization (left) and keratinocyte differentiation

(right) signatures following CB103 treatment. (F) Volcano plot highlighting upregulation of

keratin genes (e.g., KRT1, KRT10) and downregulation of the stemness marker SOX2 after CB103

treatment.

(G) GSEA plot showing increased expression of genes involved in keratin filament organization

in CB103-treated cells. (H) GSEA plot illustrating changes in genes associated with cytoskeletal

components following CB103 exposure. (I) GSEA plot showing modulation of the epithelial-to-

mesenchymal transition (EMT) gene signature in response to CB103 treatment.

Figure 9: Blockage of Notch transcription results in disruption of SCC cells migratory

activity

A) Volcano plot showing upregulation of transcripts associated with desmosomes and cell

adhesion molecules in CB103-treated cells (e.g., plakoglobin (JUP), desmocollins (DSC1/2), and

desmogleins (DSG1/2). (B) GSEA plot highlighting enrichment of gene ontology cellular

components related to desmosomes in CB103- versus DMSO-treated conditions. (C) qRT-PCR

analysis of SCC25 cells treated with DMSO or CB103 demonstrates increased expression of

desmosomal structural components following Notch inhibition (e.g. desmoplakin (DSP),

desmoglein 1 (DSG1), and desmocollins (DSC1 and DSC2)). (D) GSEA plot showing changes in

apical junction-related gene signatures in SCC25 cells treated with CB103. (E) Representative

immunofluorescence images of actin filaments and focal adhesions in SCC25 cells treated with

DMSO or CB103. Green: phalloidin. Red: vinculin. Blue: DAPI. Scale bar: 10 μm. (F)

Quantification of focal adhesion complexes per cell. *** $p < 0.001$. (G) Scratch wound assay of

SCC25 cells treated with DMSO or CB103. Scale bar 100μm. (H) Quantification of the cell-free

area in the scratch assay. Values normalized to the initial scratched area (day 0). Error bars

represent standard error. **=$p<0.01$; ***=$p<0.001$; ****=$p<0.0001$ (I) Time-lapse imaging of

SCC25 cells treated with DMSO or CB103, showing single frames from representative recordings

over a 2-hour 25-minute period (related to Movies 3 and 4). Scale bar 50μm. (J) Quantification of

single-cell displacement lengths using Imaris spot tracking.***=$p<0.001$

1013

1014

A

N1^CreERT:mTomato_mGFP

6 weeks   22 weeks
                 TAM
4NQO/DMSO   tissue collection

TAM
DMSO
4NQO

B N1:mGFP/mTOMATO/DAPI

DMSO
a   a'

4NQO
c   b'
b   c'

C Dll4/DAPI

DMSO

4NQO

D DMSO   4NQO-22 weeks

Jag1/DAPI

E DMSO   4NQO-22 weeks

Sox2/DAPI
d   d'   e   e'

K5/DAPI
f   f'   g   g'

K10/DAPI
h   h'   i   i'