Differential growth and transcriptomic profile in stem cell-derived midbrain astrocytes

Zongze Li[1&2], Lucia Fernandez Cardo[1], Michal Rokicki[1], Jimena Monzon Sandoval[1], Viola Volpato[1], Frank Wessely[1], Caleb Webber[1*], Meng Li[2*]

[1] Dementia Research Institute, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, UK

[2] Neuroscience and Mental Health Innovation Institute, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, UK

*Corresponding authors:

Meng Li, lim26@cardiff.ac.uk,

Caleb Webber, webberc4@cardiff.ac.uk

Key words: astrocyte, regional heterogeneity, ventral midbrain, dopaminergic, single cell RNA-sequencing

Abstract

Regional specificity of stem cell-derived astrocytes is believed to be an important prerequisite for their applications in disease modelling and cell-based therapies. The regional identity of these astrocytes is often defined by the positional characteristics of their antecedent, stem cell-derived neural progenitors patterned to a fate of interest, with the assumption that the positional specification is to be preserved by the derived astrocytes. Using a human induced pluripotent stem cell line designed for tracing midbrain floor plate derivatives, here we show that lineage composition of the derived astrocytes is not a faithful recapitulation of the founder progenitor population, as demonstrated by the loss of floor plate differentiated progeny in the final astrocyte products. Using deep single cell RNA sequencing, we identified distinct transcriptomic signatures of midbrain floor plate-derived astrocytes. Our study highlights the need for rigorous characterisation of pluripotent stem cell-derived regional astrocytes and provides a valuable resource for assessing midbrain floor plate-derived human astrocytes.

Introduction

Astrocytes are the most abundant cells in the brain. They play important roles in the central nervous system in supporting neuronal survival and synaptic activities, including the regulation of ionic homeostasis, providing energetic support, elimination of oxidative stress, and neurotransmitter removal and recycle (Verkhratsky and Nedergaard 2018). Abnormalities in astrocytes have been linked to various neurodegenerative and neurodevelopmental disorders, such as Parkinson's disease, Alzheimer's disease, Huntington's disease, autism spectrum disorders and Alexander's disease (Molofsky et al. 2012; Phatnani and Maniatis 2015; Booth et al. 2017). There is therefore a growing interest in using human pluripotent stem cell (PSC)-derived astrocytes for disease modelling *in vitro* (Chandrasekaran et al. 2016).

Contrary to the widely held belief that astrocytes in the brain are largely identical, recent studies revealed diversity in their transcriptomic profile, physiological properties and function (Oberheim et al. 2009; Schober et al. 2022). For example, single cell and spatial transcriptomic studies have identified several astrocyte subpopulations in the mouse cortex (Zhu et al. 2018; Batiuk et al. 2020; Bayraktar et al. 2020). In human, although astrocyte heterogeneity remains largely elusive, heterogeneity in radial glia across brain regions and within midbrain has been reported (La Manno et al. 2016; Li et al. 2023). Furthermore, different molecular and physiological features and distinct responses to stimuli were also observed in astrocytes from different mouse brain regions (Takata and Hirase 2008; Chai et al. 2017; Morel et al. 2017; Itoh et al. 2018; Kostuk et al. 2019; Makarava et al. 2019; Xin et al. 2019; Lozzi et al. 2020). Indeed, astrocyte heterogeneity has been suggested to underly the regional susceptibility to human diseases (Schober et al. 2022). Therefore, recapitulating astrocyte regional specificity in PSC-derived astrocytes is generally accepted as an important prerequisite.

Several reports have described the generation of regional astrocytes from human embryonic stem cells or induced pluripotent stem cells (iPSCs) which include forebrain (Krencik et al. 2011; Zhou et al. 2016; Tcw et al. 2017; Lin et al. 2018; Bradley et al. 2019; Hedegaard et al. 2020; Peteri et al. 2021), ventral midbrain (Booth et al. 2019; Barbuti et al. 2020; Crompton et al. 2021; de Rus Jacquet et al. 2021), hindbrain and

spinal cord (Roybon et al. 2013; Serio et al. 2013; Holmqvist et al. 2015; Bradley et al.

2019; di Domenico et al. 2019; Yun et al. 2019). The regional identity of these

astrocytes is typically evaluated at the stage of early neural progenitors generated via

cell type- or region-directed neural patterning protocols, with the assumption that the

positional characteristics will be faithfully preserved in the final astrocyte products.

However, astrocyte production in vitro involves an extended period of astrocytic fate

induction and progenitor expansion using FGF and EGF while substantial literature

reported alterations in region-specific gene expression and/or neurogenic competence

in expanded neural progenitors (Jain et al. 2003; Sun et al. 2008; Koch et al. 2009;

Falk et al. 2012). Therefore, better characterisations of PSC-derived astrocytes and

their lineage-specific features are needed to advance our knowledge about the

molecular heterogeneity of human astrocytes.

Using a human iPSC line that allows the tracing of LMX1A expressing midbrain floor

plate neural progenitors and their differentiated progeny (Cardo et al. 2023), we

discovered an unexpected gradual depletion of LMX1A$^+$ progenitor progeny during

astrocyte induction from a bulk population of ventral midbrain patterned progenitors

despite LMX1A$^+$ progenitors being the predominant starting population. LMX1A$^+$

progenitor derived astrocytes can however be generated if astrocytic induction is

initiated from purified LMX1A$^+$ progenitors, indicating that the positional constituents

of the founding cell population may not be preserved faithfully in the derived

astrocytes. Single cell RNA sequencing (scRNAseq) of astrocytes derived from both

parental populations identified distinct transcriptomic signatures, providing a useful

resource for assessment of human PSC-derived midbrain astrocytes.

Results

Depletion of LMX1A$^+$ progenitors and/or derivatives in ventral midbrain patterned neural progenitor cultures during astrogenic induction

89 To investigate whether regionally patterned neural progenitors retain their lineage

90 identity during astrogenic induction and glial progenitor expansion, we made use of

the LMX1A-Cre/AAVS1-BFP iPSCs tracer line, which enables the tracking of LMX1A[+] midbrain floor plate progenitors and their differentiated progeny(Cardo et al. 2023)⌨. We differentiated the LMX1A-Cre/AAVS1-BFP iPSCs towards the ventral midbrain fate following a modified protocol based on Jaeger *et al.* and Nolbrant *et al.*(Jaeger et al. 2011; Nolbrant et al. 2017)⌨(Figure 1A). Immunocytochemistry of day (d)19 cultures confirmed a high proportion of cells expressing BFP (96.01±0.42%) and ventral midbrain progenitor markers LMX1A (92.94±0.91%), FOXA2 (94.76±0.57%) and OTX2 (97.82±0.28%; Figure 1B-D with the original images shown in Figure S1A). Most cells (91.26±1.64%) co-expressed LMX1A and FOXA2 (Figure 1B and 1D). At this stage, all BFP[+] cells also stained positive to a pan neural progenitor marker NESTIN (Figure 1C). We detected a small proportion of cells expressing midbrain basal plate marker NKX6.1 (2.03±0.47%, Figure 1B-C) whose expression domain in the early developing ventral midbrain partially overlap with that of (Andersson et al. 2006)(Andersson et al. 2006)⌨. However, few PAX6[+] cells were present (Figure 1C), which marks the forebrain and (Duan et al. 2013)(Duan et al. 2013)⌨. Immunocytochemical analysis of FACS-sorted BFP[+] cells confirmed highly enriched expression of LMX1A (95.51±0.09%) and FOXA2 (95.35±0.14%), and co-expression of both markers (94.68±0.10%; Figure 1B and 1D). In contrast, only a small number of LMX1A[+] cells (4.96±0.70%) were present in the sorted BFP[-] population (Figure 1B and1D). These findings provide further support that BFP expression faithfully recapitulate LMX1A expression and that LMX1A[+] ventral midbrain progenitors represent the major cell population in d19(Cardo et al. 2023)(Cardo et al. 2023)⌨.

The d19 cells were then induced to undergo astrogenic switch in media containing FGF2 and EGF with the BFP content monitored by flow cytometry at each weekly passaging (Figure 1A, representative gating strategy is shown in Figure S1B-E). We found, unexpectedly, a dramatic decrease of BFP[+] cell proportion from the starting point of 88.20±1.10% to only 27.59±8.28% at d43 and to nearly absent as the differentiation continued (Figure 1E). We did not observe evident cell death during culture and replating; thus, the absence of BFP could be either due to the silencing of BFP expression in the derivatives of LMX1A[+] progenitors or the loss of these cells through growth competition. To address this question, we performed progenitor expansion and astrogenic induction under the same culture condition with purified d19

BFP$^+$ progenitors isolated using fluorescence-activated cell sorting (FACS).
Interestingly, the sorted BFP$^+$ cells exhibited similar population growth rate to that of
unsorted cultures, and the proportion of BFP$^+$ cells remained at approximately 90%
throughout the astrogenic induction and glial progenitor expansion period (Figure 1E).
This observation demonstrates that BFP expression can be maintained in the
derivatives of LMX1A$^+$ midbrain floorplate progenitors, and that the loss of BFP$^+$ cells
in the unsorted culture is likely due to their growth disadvantage compared to the
derivatives of LMX1A$^-$ progenitors.

Our finding is unexpected and demonstrates that the regional or lineage identity of
PSC-derived astrocytic cells should not be assumed merely based on the dominant
regional property of their cellular origin, given that no *in vitro* fate specification
paradigm is 100% efficient.

Figure 1. Depletion of LMX1A+ progenitors and their derivatives during astrogenic induction in ventral midbrain patterned neural progenitor cultures.

A, Schematic diagram of ventral midbrain neural differentiation and astrogenic induction. B-C, Representative view of immunocytochemistry of ventral midbrain neural progenitor markers and other regional markers in d19 unsorted, sorted BFP+, and sorted BFP- population. Scale bar represents 100 μm. Images shown were cropped to 300 μm×300 μm by randomly selecting the region of interest in the nuclei-

only channel (uncropped greyscale images are shown in Figure S1A). D,
Quantification of marker expression in unsorted, sorted BFP$^+$, and sorted BFP$^-$
population. Error bars represent the standard error of means (SEM) of three
independent experiments. E, Flow cytometry quantification of unsorted and BFP+
population during astrogenic induction and progenitor expansion. Each data point
represents one biological replicate. The gating strategy used is shown in Figure S1B.

Astrogenic switch occurred earlier in derivatives of LMX1A$^+$ midbrain progenitors.

Since sorted BFP$^+$ (LMX1A$^+$ or their derivatives) and unsorted astrocytic cultures differ
distinctively in BFP expression soon after the initiation of astrogenic induction, for
simplicity, these cultures are hereafter referred to as the BFP$^+$ and BFP$^-$ cultures,
respectively. Wondering whether the two cell populations behave differently in the
process of astrogenic switch, we examined astrocytic marker expression in these
cultures at d45 and d98 by immunocytochemistry. SOX9 and NFIA are transcription
factors crucial for the initiation of astrogenesis and acquisition of astrogenic
competence in the developing central nervous system (Stolt et al. 2003; Deneen et al.
2006), while CD44 identifies astrocyte-restricted precursors (Liu et al. 2004). We found
that all these markers were more abundantly detected in the BFP$^+$ cultures (NFIA:
65.89±2.81%; SOX9: 57.19±4.25%) than the BFP$^-$ cultures (NFIA: 4.26±1.28%; SOX9:
8.88±1.82%) at d45 (Two-way ANOVA with post-hoc Turkey test, NFIA: p=2.52×10^{-5},
SOX9: p=9.21×10^{-6}; Figure 2A-B with the original images shown in Figure S2).
Although the number of NFIA$^+$ and SOX9$^+$ cells significantly increased in the BFP$^-$
cultures by d98 (NFIA: 44.07±4.56% on d98, p=3.58×10^{-4}; SOX9: 44.28±2.84% on
d98, p=1.21×10^{-4}), the BFP$^+$ cultures still contained more cells expressing NFIA
(65.71±4.25%; p=2.06×10^{-2}) and SOX9 (73.25±2.12%; p=5.51×10^{-4}) than BFP$^-$
cultures (Figure 2A and 2C).

To investigate whether the temporal difference in astrocytic switch between the BFP$^+$
and BFP$^-$ cultures affects maturation and functionality of the derived astrocytes, we
initiated astrocyte terminal differentiation by exposing the BFP$^+$ and BFP$^-$ astrocyte
precursors to CNTF and BMP4 (Krencik et al. 2011; Bradley et al. 2019) from d87
(referred to as early astrocytes) and d136 (late astrocytes). Both BFP$^+$ and BFP$^-$

cultures exhibited a similar expression profile of classic astrocyte markers, including AQP4, EAAT2, and S100B, but few GFAP$^+$ cells at both time points (Figure 2B and 2D with the original images shown in Figure S3).

As a reference, we also generated neural progenitors without employing any patterning cues and induced astrogenic switch and astrocyte differentiation from these non-patterned neural progenitors (Figure S4A-B). We found that, while the astrocyte cultures derived from the non-patterned progenitors contained a similar proportion of cells expressing AQP4, EAAT2 and S100B compared to the BFP$^+$ and BFP$^-$ astrocyte cultures, there are more GFAP$^+$ cells in the non-patterned astrocyte preparations (17.89±5.4%, Figure S4A-B).

Functional astrocytes exhibit transient calcium (Ca^{2+}) spikes upon chemical stimulation, such as ATP (Zhang et al. 2016). Using a FLIPR Ca^{2+} release assay, we observed a sharp increase in the intracellular Ca^{2+} concentration upon ATP administration in both the early and late BFP$^+$ astrocyte populations (early BFP$^+$: $p=5.37\times10^{-12}$; late BFP$^+$: $p<2.2\times10^{-16}$; Figure 2E-F). ATP induced Ca2$^+$ release is partially mediated by inositol trisphosphate. Indeed, addition of an inositol trisphosphate receptor antagonist 2-aminoethoxydiphenylborate (2-APB) reduced the amplitude (early BFP$^+$: $p=5.88\times10^{-5}$; late BFP$^+$: $p=9.22\times10^{-10}$; Figure 2F) and rise time (early BFP$^+$: $p=2.63\times10^{-5}$; late BFP$^+$: $p=3.59\times10^{-4}$; late BFP$^-$: $p=0.028$; Figure 2G) of ATP-induced Ca2$^+$ response in both the early and late BFP$^+$ astrocytes. Interestingly, the early BFP$^+$ astrocytes had significantly lower peak amplitude than that observed in late BFP$^+$ astrocytes ($p=6.92\times10^{-9}$; Figure 2F), despite their similar level of astrocyte marker expression, suggesting a difference in maturity at the functional level. Early and late BFP$^-$ astrocytes exhibited a similar profile of time-dependent increase in the amplitude of ATP-induced Ca^{2+} response but did not reach statistical significance (Figure 2E-F). However, late BFP$^-$ astrocytes showed a significantly lower peak amplitude than late BFP$^+$ astrocytes ($p<2.2\times10^{-16}$; Figure 2E-F).

Taken together, our data demonstrate that astrogenesis occurred earlier in BFP$^+$ cultures than BFP$^-$ cells. This temporal difference is also reflected in functional maturity of the derived astrocytes, despite a similar expression profile of classic astroglial markers.

Figure 2. Early astrogenic switch and astrocyte maturation in derivatives of LMX1A[+] midbrain progenitors.

A, Representative view of immunocytochemistry of astrogenic marker expression in BFP[+] and unsorted progenitors at day 45 and 98. Scale bar represents 100 μm. Images shown were cropped to 462 μm $\times$ 462 μm by randomly selecting the region of interest in the nuclei-only channel (uncropped greyscale images are shown in Figure S2). B, Representative view of immunocytochemistry of astrocyte marker expression in early and late, BFP[+] and BFP[-] (unsorted) astrocytes. Scale bar represents 100 μm. Images shown were cropped to 300 μm $\times$ 300 μm by randomly selecting the region of interest in the nuclei-only channel (uncropped greyscale images are shown in Figure S3). C, Quantification of immunocytochemistry of astrogenic marker in BFP[+] and BFP[-] progenitors at day 45 and 98 shown in Panel A. Error bars represent the standard error of means (SEM) of three independent experiments. Two-way ANOVA was

performed to compare between lineages (NFIA: p=5.389×10^{-6}, df=1, effect size=3.62;
SOX9: p=1.96×10^{-6}, df=1, effect size=4.77) and days of differentiation (NFIA:
p=7.82×10^{-5}, df=1, effect size=1.99; SOX9: p=2.62×10^{-5}, df=1, effect size=2.99) D,
Quantification of immunocytochemistry of astrocyte marker expression in astrocytes.
Error bars represent SEM of three independent experiments. Kruskal-Wallis test
results following Bonferroni correction are shown on the top of the figure (AQP4:
p.adjust=0.12, df=3, H=8.95; EAAT2: p.adjust=1.00, df=3, H=0.95; GFAP:
p.adjust=0.06, df=3, H=10.38; S100B: p.adjust=0.11, df=3, H=9.05). E, Averaged
trace of ATP-induced Ca^{2+} response assayed using FLIPR. Drugs or DMSO were
applied at 1 minute of the assay. The line represents the average fluorescence change
(ΔF/F0) in at least three independent experiments each with at least three replicate
wells. The shaded area represents the SEM across at least three independent
experiments. F, Quantitative comparison of the peak amplitude of ATP-induced Ca^{2+}
response among conditions (two-way ANOVA, p<2.2×10^{-16}, df=2, effect size=2.54)
and samples (p=2.87×10^{-14}, df=3, effect size=2.17). Error bars represent the SEM
across at least independent experiments. G: Quantitative comparison of the rise time
of ATP-induced Ca^{2+} response among conditions (two-way ANOVA, p=2.19×10^{-13},
df=2, effect size =1.958) and samples (p=0.064, df=3, effect size=0.76). Intergroup
comparison was performed using Post-hoc Tukey test. Error bars represent the SEM
across at least three independent experiments. (****: p<0.0001, ***: p<0.001, **:
p<0.01, *: p<0.05, ns: not significant).

Single cell RNA sequencing confirms the authenticity of PSC-derived astrocytes.

To further characterize the PSC-derived astrocytes, we performed full-length
scRNAseq on early and late BFP^{+} and BFP^{-} astrocytes using the iCELL8 platform and
SMART-seq technology, with non-patterned astrocytes derived from the LMX1A-
Cre/AAVS1-BFP tracer line as a control. A sample of iPSC-derived neurons was
included to facilitate downstream cell type identification. We profiled FACS purified
astrocytes expressing CD49f as well as unsorted cultures for comparison of all three
astrocyte populations (Barbar et al. 2020); Figure S4C-H). After stringent filtering
(Figure S5A-C, see Methods and Materials on filtering), we obtained a total of 17478
protein-coding genes in 1786 qualifying cells, with an average of 6326 protein-coding
genes detected per cell.

Unsupervised Louvain clustering identified 12 cell clusters (Figure 3A). Cells clustered
mainly based on sample type (astrocytes and neurons; Figure S6A) and the estimated
cell cycle phase (Figure S6B), while sorted CD49f^{+} and unsorted astrocytes were

clustered together (Figure S6A). *TagBFP* was detected at a higher level in BFP$^+$
astrocyte samples than in BFP$^-$ and NP samples, while *TagBFP* expression was
negligible in the neuronal samples derived from an iPSC line without BFP transgene
(Figure S6C-D). Using a set of known astrocyte and neuronal signature genes (Figure
3B), we identified cells in clusters 0, 1, and 5-11 as astrocytes (Figure 3B), which were
enriched in expression of *SOX9, NFIA, NFIB, HES1, S100A13, S100A16, EGFR,*
CD44 and *GJA1* (Figure 3B). These transcripts were also detected at high levels in
clusters 2 and 4, which were mostly estimated to be in cell cycle phases G2, M and S
(Figure S6B). In addition, clusters 2 and 4 showed high levels of proliferation-related
transcripts, such as *TOP2A, MKI67, CDK1* and *AURKA* (Figure 3B), and are thus
defined as astrocyte precursors. In contrast, Cluster 3 contains mostly cells from the
neuronal sample (Figure 3A and Figure S6A) and indeed expressed high levels of
genes closely related to neuronal structure and function (such as *STMN2*, *SYT1*, *DCX*,
MAPT, and *SNAP25*; Figure 3B). We did not detect transcripts indicative of endoderm
(*GATA4*), mesoderm (*TBXT* and *TBX6*), and oligodendrocyte progenitors (*SOX10* and
PDGFRA) in any of these clusters (Figure S6C).

To determine the authenticity of these PSC-derived astrocytes, we mapped our data
to published scRNAseq datasets obtained from five foetal, an adult human brain and
a PSC-derived astrocyte study using Seurat integration (La Manno et al. 2016; Sloan
et al. 2017; Zhong et al. 2018; Polioudakis et al. 2019; Agarwal et al. 2020; Fan et al.
2020; Bhaduri et al. 2021; Eze et al. 2021). We found that cells in clusters annotated
as astrocytes (clusters 0, 1, and 5-11) were indeed predominantly mapped to the
reference astrocyte or astrocyte precursor populations with high confidence
(prediction score over 0.5; Figure 3C and Figure S7). In contrast, neuronal cluster 2
was mapped to neurons of the foetal reference datasets, while the astrocyte precursor
clusters (2 and 4) were mapped to progenitor populations in the foetal reference
datasets (Figure 3C and Figure S7). These findings demonstrate that our iPSC-
derived astrocytes closely resemble those in the human brains.

Figure 3. Single cell RNA sequencing confirms the authenticity of PSC-derived astrocytes.

A, Uniform manifold approximation and projection plot of unbiased clustering, coloured by clusters. B, Heatmap of the normalised expression of selected markers in different clusters. The assigned identity to each cluster is shown at the top of the plot. C, Sankey plot summarising the result of reference mapping of cells in different clusters to eight published reference human brain scRNAseq datasets. The thickness of the thread is proportional to the number of cells mapped to the same identity in the reference datasets (predicted identity). Detailed results of referencing mapping to each reference datasets are shown in Figure S7A-H and prediction score shown in Figure S7I. (PRE: precursors; N: neurons)

Distinct transcriptome fingerprints of LMX1A⁺ midbrain floor plate-derived astrocytes

Significant advance has been made recently in understanding the molecular profiles
of midbrain dopamine neurons. However, our knowledge about midbrain astrocytes in
this regard remains limited and does not inform anatomic or lineage origin of the cells.

In this regard, the BFP$^+$ astrocytes provide a unique resource to determine the
transcriptomic characteristics of human midbrain floor plate derived astrocytes. By
performing pairwise differential gene expression (details described in Methods and
Materials), we identified 1153 genes differentially expressed (DEGs; adjusted p values
less than 0.05 and log2 fold change over 0.25) in BFP$^+$ astrocytes when compared to
either BFP$^-$ or non-patterned astrocyte populations (Supplementary Data 1). Of these,
159 were unique to BFP$^+$ astrocytes (BFP$^+$ enriched, Figure 4A), which include genes
associated with midbrain dopamine neuron development such as *SULF1*, *LMO3*,
NELL2, and *RCAN2* (Figure 4B) (Strelau et al. 2000; La Manno et al. 2016; Bifsha et
al. 2017; Ahmed et al. 2021). Interestingly, *LMX1A* and *FOXA2*, which was used to
evaluate PSC-derived midbrain astrocytes in previous studies, were not detected in
BFP$^+$ astrocytes (Figure S6D). We also identified 530 DEGs enriched in BFP$^-$
astrocytes only (BFP$^-$ enriched, Figure 4A and Supplementary Data 1), which include
those known to be expressed in the ventrolateral - dorsal domain of the midbrain and
hindbrain, such as *IRX3*, *IRX5*, *PAX3*, and *PAX7* (Figure 4B)(Houweling et al. 2001;
Matsunaga et al. 2001). This transcription profile supports the notion that the BFP$^-$
astrocytes are descendants of the initial minor populations of lateral midbrain
progenitors. Moreover, 72 DEGs were shared by BFP$^+$ and BFP$^-$ astrocytes against
the non-patterned astrocytes (midbrain enriched, Figure 4A and Supplementary Data
1). This set of genes includes *NR2F1*, *NR2F2, ZEB2, KCNJ6* and *SRPX* (Figure 4B)*,*
which have been reported to be signatures of mouse midbrain astrocytes (Endo et al.
2022). Together, our findings provide a new entry to transcriptomic characteristics of
midbrain astrocytes and specifically a gene expression map of midbrain floor plate
derived human astrocyte lineage.

Gene ontology (GO) enrichment analysis was performed on the 1153 DEGs enriched
in BFP$^+$ astrocytes (Supplementary Data 2). The significantly enriched GO terms were
mainly related to various aspects of metabolism, stress response, biosynthesis,
lysosomal activity, and cellular respiration (Figure 4C and Supplementary Data 3).
These biological processes have previously been shown to be disrupted by several
mutations causing familial Parkinson's disease (di Domenico et al. 2019; Barbuti et al.
2020; Sonninen et al. 2020). In contrast, the GO terms associated with 530 BFP$^-$
astrocyte enriched DEGs were mostly related to formation of extracellular matrix and

tissue development (Figure 4D and Supplementary Data 2-3). The differential enrichment of GO terms implicates functional difference between the BFP$^+$ and BFP$^-$ astrocytes and supports the need for generating regional specific astrocytes for disease modelling.

Figure 4. Distinct transcriptome fingerprints of LMX1A⁺ midbrain floor plate-derived astrocytes

A, Heatmap of the normalised expression of population-specific genes in different populations of astrocytes. B, Violin plots of the normalised expression of selected candidate markers for BFP⁺, BFP⁻, and non-patterned (NP) astrocytes. C-D, Representative GO terms significantly enriched in BFP⁺ (C) and BFP⁻ (D) enriched genes. Semantically similar representative terms were shown with the same colour.

Discussion

Despite the general belief that recapitulating astrocyte lineage heterogeneity is necessary for stem cell-based disease modelling and cell transplantation, the extent of astrocyte heterogeneity in different brain regions and subregions remains largely elusive. By harnessing an LMX1A based lineage tracing human iPSC line and cutting edge scRNAseq technology, we show that astrocytes derived from the LMX1A⁺ midbrain floor plate progenitors, the same cells giving rise to midbrain dopaminergic neurons, possess distinct transcriptional landmarks from those derived from non-midbrain patterned neural progenitors as well as other midbrain patterned progenitors. Moreover, we discovered unexpected negative selection against derivatives of LMX1A⁺ progenitors during astrocyte induction and progenitor expansion. Our study highlights the need for careful characterisation of PSC-derived astrocytes and provides a transcriptomic fingerprint for midbrain floor plate derived astrocytes.

Using a popular astrocyte in vitro differentiation paradigm, we found that astrocytes descended from the LMX1A⁺ midbrain progenitors could only be obtained from purified progenitors. In contrast, astrocytes derived from bulk midbrain patterned progenitors exhibits transcriptomic profiles of the lateral-dorsal midbrain, despite LMX1A⁺ midbrain progenitors being the predominant starting population. Our findings demonstrate that the lineage composition of the parent progenitors may not be faithfully preserved during astrocyte induction and progenitor expansion. FGF is the most popular inductive molecule used for astrocyte differentiation from stem cells (Chandrasekaran et al. 2016). It is evident however that FGF expanded neural progenitors, originated either from the brain or neutralized PSCs, exhibit restricted regional competence and

positional gene expression. For example, bulk expanded human ventral midbrain

neural progenitors (Jain et al. 2003), and fetal forebrain or spinal cord derived neural

stem (NS) cells only give rise to GABAergic neurons (Sun et al. 2008); likewise, lt-

NES cells display anterior hindbrain-like positional profile (Falk et al. 2012), while their

antecedents, PSC-derived neural rosettes and early passage derivatives, express

anterior forebrain markers (Koch et al. 2009). It is not clear whether this is due to

deregulation of the original patterning at the level of gene expression or the loss of

associated cell population (Gabay et al. 2003). In this study, since BFP$^+$ astrocytes

can be generated under the same culture condition with purified LMX1A$^+$ progenitors,

we reasoned that the loss of their derivatives in unsorted cultures was possibly due to

differential growth capacity.

Our study highlights the need for careful assessment of astrocyte positional identity.

A common practice in this regard is to confirm the regional characteristics of the

founder progenitors following fate directed neural induction, with the assumption that

the dominant positional features will be maintained by the astrocyte progeny (Krencik

et al. 2011). This strategy is at least partly dictated by our limited knowledge in gene

expression signatures of regional- and/or lineage-specific astrocytes. Hence, an end

point evaluation of PSC-derived astrocytes often relies on region-specific markers

defined in the developmental brain during the neurogenic period. For example, LMX1A

and FOXA2 expression were used as criteria for midbrain astrocytes in previous

studies (Barbuti et al. 2020; Crompton et al. 2023). However, scRNAseq of human

fetal ventral midbrain and adult substantia nigra revealed negligible expression of

these transcripts in astrocytes (La Manno et al. 2016; Agarwal et al. 2020; Kamath et

al. 2022). Consistent with these findings, we did not detect LMX1A or FOXA2 in neither

the BFP$^+$ nor BFP$^-$ astrocytes. Our analysis however identified new positive and

negative markers that could be applied to confirm the regional identity of ventral

midbrain astrocytes.

In addition to distinct transcriptomic profile, BFP$^+$ and BFP$^-$ astrocytes may also be

functionally different. Astrocytes generated from progenitors broadly patterned to the

dorsal forebrain, ventral forebrain and spinal cord have been shown to exhibiting

different GO enrichment profile as well as different physiological and functional

properties (Bradley et al. 2019). In comparison to the BFP$^-$ and non-patterned

astrocytes, the current study revealed that GO terms enriched in BFP⁺ astrocytes,
which originated from the same progenitor giving rise to midbrain dopaminergic
neurons, were closely related to various biological processes disrupted in astrocytes
carrying familial Parkinson's disease mutations (di Domenico et al. 2019; Barbuti et al.
2020; Sonninen et al. 2020). Such a distinct enrichment profile implicates BFP⁺
astrocytes being functionally adapted to supporting midbrain dopaminergic neurons
compared to BFP⁻ and non-patterned astrocytes.

In conclusion, this study lends further support on regional diversity of astrocytes and
identified a set of midbrain enriched genes. Crucially, the transcriptomic fingerprint of
human midbrain floor plate-derived astrocytes described here offers a much-needed
resource for assessing the authenticity of stem cell derived astrocytes in studies
associated with Parkinson's disease.

Methods and Materials

Stem cell culture and astrocyte differentiation

KOLF2 human iPSCs were maintained in E8 flex media (ThermoFisher) and manually
dissociated using Gentle Cell Dissociation Reagent (STEMCELL Technologies) as
previously described (Cardo et al. 2023). Astrocytes were differentiated using a three-
stage stepwise strategy consisting neural induction and regional patterning, astrogenic
switch and progenitor expansion, and astrocyte terminal differentiation. Midbrain floor
plate progenitors were generated as previously described (Cardo et al. 2023). At day
19, cells were replated as single cells onto poly-D-lysine-laminin-coated plates at
1×10^6 cells/cm^2 for astrogenic switch and progenitor expansion in N2B27 media
supplemented with 10 ng/mL FGF2 (Peprotech) and 10 ng/mL Human EGF
(Peprotech) and replated every 6-8 days. For astrocyte terminal differentiation,
expanded neural progenitors were re-plated at a density of 3×10^4 cells/cm^2 in
expansion media and 24 hours later switched to N2B27 supplemented with 10 ng/mL
human recombinant CNTF (Peprotech) and 10 ng/mL human recombinant BMP4
(Peprotech) for 7 days followed by media containing CNTF alone for another 13 days.
10 µM Y-27632 was used for 24 hours before and after each replating. The protocol

for generating non-patterned astrocytes was the same as for floor plate-derived
astrocyte except the neural progenitors were derived with duo-SMAD inhibitors only
without ventral patterning reagents.

Flow cytometry analysis and cell isolation

Cells were dissociated in Accutase as described above and washed twice with DPBS
by centrifugation for 5 minutes at 200 rcf. For evaluating BFP expression, dissociated
cells were resuspended in 0.5 mM EDTA in DPBS (Sigma-Aldrich) and analysed on a
BD LSRFortessa cell analyser (BD Biosciences). For purifying BFP$^+$ cells, dissociated
cells were resuspended in the same cell culture media. Background autofluorescence
was compensated for using KOLF2 parental cell line at a similar stage of differentiation
to define BFP$^-$ gating. For purifying CD49f$^+$ astrocytes, dissociated cells were stained
with Alexa Fluor 647-conjugated rat anti-CD49f antibody (5% v/v in a 100 μL reaction;
BD Biosciences) for 25 minutes at 37°C on an orbital shaker at 200 rcf and
resuspended in DPBS containing 0.5% bovine serum albumin and 50 units/mL DNase
I (Sigma Aldrich). Background autofluorescence was compensated for using KOLF2
parental cell line at a similar stage of differentiation to define BFP$^-$ gating and unstained
astrocytes to define CD49f$^-$ gating. Cell sorting was performed on a BD FACSAria III
(BD Biosciences) using an 80 μm nozzle. Sorted cells were collected in the same
media as for resuspension. Flow cytometry data were analysed in FlowJo v10.8.1 (BD
Biosciences) as shown in Figure S1B-E. Briefly, non-debris events were selected
using the eclipse gates on dot graphs of SSC-A versus FSC-A. Singlet events were
sequentially gated using polygonal gates on dot graphs of FSC-H versus FSC-A and
SSC-H versus SSC-A by selecting the events in the diagonal region. The positive and
negative gates in the fluorescence channel were set as bifurcate gates at a minimum
of 99.9% percentile (usually at 99.99% percentile) on the histogram of the
fluorescence intensity of the negative control sample of the same flow cytometry
experiment and applied to all samples of the same flow cytometry experiment.

Immunocytochemistry

Cultures were fixed with 3.7% PFA for 15-20 min at 4 ºC. For nuclear antigen
detection, an additional fixation with methanol gradient was performed, which include

5 mins each in 33% and 66% methanol at room temperature followed by 100%
methanol for 20 min at -20°C. Cultures were then returned to PBST via inverse
gradient and were then permeabilized with three 10-minute washes in 0.3% Triton-X-
100 in PBS (PBS-T) and then blocked in PBS-T containing 1% BSA and 3% donkey
serum. Cells were incubated with primary antibodies in blocking solution overnight at
4°C. Following three PBS-T washes, Alexa-Fluor secondary antibodies (Thermo
Fisher Scientific) were added at 1:1000 PBS-T for 1 hour at ambient temperature in
the dark. Three PBS-T washes were then performed that included once with DAPI
(Molecular Probes). Images were taken on a Leica DMI6000B inverted microscope.
Quantification was carried out in Cell Profiler (Stirling et al. 2021) or manually using
ImageJ (Schindelin et al. 2012) by examining at least four randomly selected fields
from three independent experiments. The antibodies used are provided in the
Supplementary Table 1. Representative images shown in main figures were cropped
by randomly selecting the region of interest in the DAPI-stained channel only, with the
original unedited images shown in Figure S1A, S2 and S3.

Single-cell RNA-sequencing

Cells were dissociated with Accutase with 10 units/mL of papain (Sigma-Aldrich) for
10 minutes at 37°C and resuspended in 0.5% bovine serum albumin (Sigma Aldrich)
with 50 units/mL DNase I in DPBS without calcium or magnesium (Gibco) and passed
through a cell strainer with 35 μm mesh. Cells were stained with 1 μM SYTO16
(Invitrogen) and 0.08% (v/v) propidium iodide (part of the Invitrogen ReadyProbes™
Cell Viability Imaging Kit, Blue/Red) for 20 minutes on ice and then dispensed into the
nano-well plates of the ICELL8® cx Single-Cell System (Takara). Wells containing
single viable cells were automatically selected using the ICELL8 cx CellSelect v2.5
Software (Takara) with the green and not red logic. Manual triage was performed to
recover additional candidate wells that contain viable single cells. The library for
sequencing was then prepared using the SMART-Seq ICELL8 application kit (Takara)
following the manufacturer's recommended protocol. Next-generation sequencing was
performed using the NovaSeq6000 and the Xp Workflow on a S4 flow cell for 200
cycles of pair-end sequencing.

Single-cell RNA-sequencing analysis

Using the Cogent NGS Analysis Pipeline software v 1.5.1 (Takara), FASTQ files containing all indices for each chip were demultiplexed to FASTQ files containing one index per file. Adaptor sequences were removed using cutadapt 3.2 with the following settings: -m 15 --trim-n --max-n 0.7 -q 20. Trimmed FASTQ files were aligned to the *Homo sapiens* GRCh38.106 primary assembly with the BFP reporter gene attached to the end of the genome, using STAR 2.7.9a (Dobin et al. 2013) with the following settings: --outSAMtype BAM Unsorted --quantMode TranscriptomeSAM --outReadsUnmapped Fastx --outSAMstrandField intronMotif --chimSegmentMin 12 --chimJunctionOverhangMin 8 --chimOutJunctionFormat 1 --alignSJDBoverhangMin 10 --alignMatesGapMax 100000 --alignIntronMax 100000 --alignSJstitchMismatchNmax 5 -1 5 5 --chimMultimapScoreRange 3 --chimScoreJunctionNonGTAG -4 --chimMultimapNmax 20 --chimNonchimScoreDropMin 10 --peOverlapNbasesMin 12 --peOverlapMMp 0.1 –alignInsertionFlush Right --alignSplicedMateMapLminOverLmate 0 --alignSplicedMateMapLmin 30. After alignment, gene-level quantification was performed using featureCounts from subread 2.0.0 (Liao et al. 2013) with the following settings: -t exon --primary -R CORE -F GTF -Q 0 -B -g gene_id. The count matrix of each index was combined in R 4.2.0 (Team 2023).

All downstream analysis was performed in R 4.3.0 using Seurat 4.3.0 (Stuart et al. 2019). Gene-level filtering was applied by including only protein-coding genes with at least five total counts across all cells and being expressed in at least 1% of all cells. Poor quality cells were then identified using the *is.outlier* function from the scater 1.28.0 (McCarthy et al. 2017). Poor quality cells were defined as having a high percentage of mitochondrial gene count, or high or low the total number of genes detected, or high total gene counts. The thresholds of each metrics for each sample were determined as twice the median absolute deviation of the sample. Raw gene counts were log normalised with a scale.factor setting of 1×10^5. Data from the two batches of experiments were integrated using the *FindIntegrationAnchors* and *IntegrateData* based on the common top 2000 highly variable genes and the first 30 dimensions of principal components. The percentage of mitochondrial gene count and

total gene count were regressed out using the *ScaleData* function. Principal
component analysis (PCA) was performed on the top 2000 high variable genes and
the number of principal components used for uniform manifold approximation and
projection (UMAP) was determined using the JackStraw method (Chung and Storey
2015). UMAP and unbiased Louvain clustering was performed on the first 33 principal
components. Pairwise differential gene expression analysis was performed using the
MAST method (Finak et al. 2015) with "Chip" as the latent variable. Gene ontology
enrichment analysis was performed using *enrichGO* function in the clusterProfiler
4.10.0 package (Stirling et al. 2021) with all genes in the filtered dataset as the
background. GO term database was downloaded using the org.Hs.eg.db 3.18.0
package (Carlson 2019). Revigo v1.8.1 was used to group the representative GO
terms based on semantic similarity using a size setting of 0.5, Homo sapiens
database, and SimRel method for semantic similarity (Schlicker et al. 2006; Supek et
al. 2011).

Published datasets were downloaded from NCBI's Gene Expression Omnibus
(Clough and Barrett 2016) and processed in R 4.2.0. Gene level filtering was
performed by retaining only protein-coding genes with more than five total counts
across all cells. Gene counts were normalised using the NormalizeData function
(scale.factor settings are listed in Supplementary Table 2). PCA was performed based
on the top 2000 highly variable genes to obtain the first 50 PCs. Visual inspection of
the elbow plot was used to determine the number of PCs for downstream analysis.
Batch effect between subjects was evaluated on the two-dimensional PC2~PC1 plot.
Where inter-subject batch effect was observed, Harmony integration was performed
based on the PCs selected in the previous step (Supplementary Table 2). UMAP was
performed based on either the PCA or Harmony reduction (using the top 30
dimensions), and Louvain clustering was performed (settings shown in Supplementary
Table 2). Cluster identities were verified against the reported annotation where
possible. For datasets without detailed annotation published or astrocyte lineage
reported (Supplementary Table 2), reannotation was performed based on the
expression of known markers (Figure S7A-E). Reference mapping was performed
using *FindTransferAnchors* and *TransferData* function in Seurat based on the first 30
dimensions of either the PCA or Harmony loadings of the reference dataset.

Statistical analyses

All data were collected from at least three independent experiments and presented as mean ± standard error of means unless otherwise specified. Data were tested for normality with the Shapiro-Wilk test and for equal variance with Levene test before performing statistical analyses by two-way ANOVA with post-hoc Tukey test for multiple comparisons where relevant. Kruskal-Wallis test with post-hoc Dunn's test for pairwise comparison was used where parametric test was not suitable. Effect size was calculated as Cohen's f for ANOVA or eta squared based on the H-statistic for Kruskal-Wallis test. All statistical tests were performed in R4.3.0.

Supplementary Information

Figure S1. Original images of immunocytochemistry of d19 progenitors and gating strategy of BFP flow cytometry analysis.

A, the original images shown in Figure 1B-C. The gating strategy used for BFP flow cytometry analysis are shown in B-E and described in Methods and Materials. B,

570 scatter plot of SSC-A versus FSC-A. C, scatter plot of FSC-H versus FSC-A. D-E,

571 histogram of BFP fluorescence in the negative control (D) and BFP-expressing (E)

572 samples.

575 **Figure S2. Original images of immunocytochemistry of astrogenic markers**

576 **shown in Figure 2A.**

Figure S3. Original images of immunocytochemistry of astrocyte markers shown in Figure 2B.

Figure S4. Non-patterned astrocytes and fluorescence-activated cell sorting of astrocytes for single cell RNA sequencing.

A, Representative view of immunocytochemistry of astrocyte marker expression in non-patterned astrocytes (scale bar represents 100 μm). B, Quantification of astrocyte marker expression in astrocytes. Error bars represent SEM of three independent experiments. C-E, negative control samples; F-H, one sample of BFP+ astrocytes). C and F shows the dot plot of SSC-A versus FSC-A. D and G shows the dot plot of FSC-H versus FSC-A. E and H shows the dot plot of Alexa Fluor-647-A (labelling CD49f) versus DAPI-A (labelling BFP). P3 was used to isolate CD49f+ population (including both BFP+ and BFP-), while P4 was used to isolate CD49f+/BFP+ population.

Figure S5. Processing of single cell RNA sequencing data.

A-C, Violin plots showing the results of adaptive cell-level filtering based on the number of genes detected per cell (A), percentage of mitochondrial gene counts per cell (B), and total gene count per cell (C). D-E, UMAP plot of all filtered cells before (D) and after integration (E) coloured by chip. F-G, UMAP plot of the subset of sorted BFP+ astrocytes on Chip A and B derived from the same astrocyte differentiation before (F) and after integration (G) coloured by chip. (N: neuron; SBN: sorted BFP-; SBP: sorted BFP+; SNP: sorted non-patterned; UBN: unsorted BFP-; UBP: unsorted BFP+; UNP: unsorted non-patterned; numbers represent independent samples)

Figure S6. Marker expression in scRNAseq.

A-B, UMAP plot of all filtered cells after integration coloured by sample type and sorting status (A) and estimated cell cycle phase (B). C, Raincloud plots of normalised expression of *BFP*. D, Histogram of the distribution of raw gene count of *BFP*. E, Violin plots of the normalised expression of endoderm, mesoderm, neuroectoderm, and oligodendrocyte progenitor markers. F, Violin plots of the normalised expression of classic ventral midbrain markers.

Figure S7. Reference mapping to human brain single cell RNA sequencing

datasets.

A-H, Sankey plot summarising the result of reference mapping of cells in different

clusters to eight published reference human brain scRNAseq datasets. The thickness

of the thread is proportional to the number of cells mapped to the same identity in the

reference datasets (predicted identity). Cluster IDs in this study are shown on the left. Heatmap in Panel A-E shows the expression of marker genes in different clusters in the re-annotated reference datasets. I, Heatmap of prediction score from Seurat integration.

Supplementary Table 1. Antibodies used in this study.

Target	Species	Producer	Catalogue number	Dilution for use
ALDH1L1	MS	Abcam	ab56777	1:500
AQP4	RB	Abcam	ab259318	1:200
BFP	rabbit	Evrogen	AB233	1:3000
CD44	mouse	Cell Signalling Technology	3750	1:400
EAAT2	goat	Abcam	ab235202	1:200
FOXA2	goat	R&D system	AF2400	1:1000
GFAP	rat	Life Tech	AB10533	1:1000
LMX1A	rabbit	Millipore	HPA030088	1:2000
NESTIN	mouse	BD	BD611659	1:300
NFIA	rabbit	Abcam	AB228897	1:1000
NKX6.1	mouse	DSHB	F55A10	1:100
OTX2	goat	R&D system	AF1979	1:200
PAX6	mouse	DSHB	PAX6	1:1000
S100B	rabbit	Abcam	ab52642	1:1000
SOX9	goat	R&D system	AF3075	1:300
Anti-Goat IgG, Alexa Fluor 488	donkey	Invitrogen	A32814	1:1000
Anti-Goat IgG, Alexa Fluor 555	donkey	Invitrogen	A32816	1:1000
Anti-Goat IgG, Alexa Fluor 647	donkey	Invitrogen	A32849	1:1000

Anti-Mouse IgG, Alexa Fluor 488	donkey	Invitrogen	A21202	1:1000
Anti-Mouse IgG, Alexa Fluor 555	donkey	Invitrogen	A31570	1:1000
Anti-Mouse IgG, Alexa Fluor 647	donkey	Invitrogen	A31571	1:1000
Anti-Rabbit IgG, Alexa Fluor 488	donkey	Invitrogen	A21206	1:1000
Anti-Rabbit IgG, Alexa Fluor 555	donkey	Invitrogen	A31572	1:1000
Anti-Rabbit IgG, Alexa Fluor 647	donkey	Invitrogen	A31573	1:1000
Anti-Rat IgG, Alexa Fluor 488	donkey	Invitrogen	A21208	1:1000

622 **Supplementary Table 2. Settings used for processing published datasets.**

PMID	Tissue	Age	Scale factor	PCs used	Harmony integration	Clustering resolution	Re-annotation
3282 6893	SN and cortex	Adult	10000	50	No	0.5	No
3372 3434	multiple	CS22	10000	50	Yes	0.3	Yes
3461 6070	multiple	GW25	10000	50	Yes	0.4	Yes
3292 3614	Neocortex	GW9-28	100000	30	Yes	0.9	Yes
3130 3374	Neocortex	GW17-18	10000	30	Yes	0.5	Yes
2953 9641	Prefrontal cortex	GW8-26	100000	30	Yes	0.9	Yes
2881 7799	PSC-derived astrocytes	-	10000	30	No	0.3	No
2771 6510	Ventral midbrain	GW7-11	10000	30	No	0.5	No

624 Supplementary Data 1: Pairwise DEGs of three astrocyte and one neuron populations.

625 Supplementary Data 2: GO enrichment of population enriched DEGs.

626 Supplementary Data 3: Representative enriched GO terms of DEGs for BFP+ and
627 BFP- astrocytes.

628 Supplementary Data 4: DEGs comparing sorted and unsorted populations.

Acknowledgements

630 We would like to thank Mark Bishop and Joanne Morgan for conducting FACS and
631 next generation sequencing, respectively. We thank Kathryn Peall and Laura Abram
632 for providing iPSC-derived neurons for scRNAseq. We also thank the support of the
633 Supercomputing Wales project, which is part-funded by the European Regional
634 Development Fund (ERDF) via the Welsh Government.

Author Contribution

Z.L. and M.L. conceived the study and designed the experiments. Z.L. performed all
cell experiments and analysed all data. L.C. and M.L. designed the lineage tracing
system, and L.C. generated the cell line with assistance from Z.L. M.R. performed
iCELL8 library preparation. J.M.S., F.W., and V.V. provided guidance and contributed
to general discussions on scRNAseq data analysis. C.W. contributed to the design of
scRNAseq and general discussions throughout the work. Z.L. and M.L wrote the
paper. All authors edited and approved the paper.

Funding

644 This work was supported by the UK Dementia Research Institute, jointly funded by the
645 UK Medical Research Council, Alzheimer's Society and Alzheimer's Research UK, to
646 C.W. (MC_PC_17112) and Z.L. (DRI-TRA2021-02), and a seed corn fund to Z.L. from
647 the Neuroscience and Mental Health Innovation Institute, Cardiff University. Z.L. was
648 funded by a UK Dementia Research Institute PhD studentship.

Data Availability

The sequencing data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Clough and Barrett 2016) and are accessible through a GEO Series accession number which will be made available upon final publication.

Declaration of interests

The authors declare no competing interests.

References

Agarwal, D. et al. 2020. A single-cell atlas of the human substantia nigra reveals cell-specific pathways associated with neurological disorders. *Nat Commun* 11(1), p. 4183. doi: 10.1038/s41467-020-17876-0

Ahmed, M., Owens, M. J. S., Toledo, E. M., Arenas, E., Bradley, M. and Ffrench-Constant, C. 2021. Combinatorial ECM Arrays Identify Cooperative Roles for Matricellular Proteins in Enhancing the Generation of TH+ Neurons From Human Pluripotent Cells. *Front Cell Dev Biol* 9, p. 755406. doi: 10.3389/fcell.2021.755406

Andersson, E. et al. 2006. Identification of intrinsic determinants of midbrain dopamine neurons. *Cell* 124(2), pp. 393-405. doi: 10.1016/j.cell.2005.10.037

Barbar, L. et al. 2020. CD49f Is a Novel Marker of Functional and Reactive Human iPSC-Derived Astrocytes. *Neuron* 107(3), pp. 436-453 e412. doi: 10.1016/j.neuron.2020.05.014

Barbuti, P. A. et al. 2020. IPSC-derived midbrain astrocytes from Parkinson's disease patients carrying pathogenic SNCA mutations exhibit alpha-synuclein aggregation, mitochondrial fragmentation and excess calcium release. *bioRxiv*, p. 2020.2004.2027.053470. doi: 10.1101/2020.04.27.053470

Batiuk, M. Y. et al. 2020. Identification of region-specific astrocyte subtypes at single
cell resolution. *Nature Communications* 11(1), doi: 10.1038/s41467-019-14198-8

Bayraktar, O. A. et al. 2020. Astrocyte layers in the mammalian cerebral cortex
revealed by a single-cell in situ transcriptomic map. *Nature Neuroscience* 23(4), pp.
500-509. doi: 10.1038/s41593-020-0602-1

Bhaduri, A. et al. 2021. An atlas of cortical arealization identifies dynamic molecular
signatures. *Nature* 598(7879), pp. 200-204. doi: 10.1038/s41586-021-03910-8

Bifsha, P., Balsalobre, A. and Drouin, J. 2017. Specificity of Pitx3-Dependent Gene
Regulatory Networks in Subsets of Midbrain Dopamine Neurons. *Molecular*
Neurobiology 54(7), pp. 4921-4935. doi: 10.1007/s12035-016-0040-y

Booth, H. D. E., Hirst, W. D. and Wade-Martins, R. 2017. The Role of Astrocyte
Dysfunction in Parkinson's Disease Pathogenesis. *Trends in neurosciences* 40(6), pp.
358-370. doi: 10.1016/j.tins.2017.04.001

Booth, H. D. E. et al. 2019. RNA sequencing reveals MMP2 and TGFB1
downregulation in LRRK2 G2019S Parkinson's iPSC-derived astrocytes. *Neurobiol*
Dis 129, pp. 56-66. doi: 10.1016/j.nbd.2019.05.006

Bradley, R. A. et al. 2019. Regionally specified human pluripotent stem cell-derived
astrocytes exhibit different molecular signatures and functional properties.
Development 146(13), p. dev170910. doi: 10.1242/dev.170910

Cardo, L. F., Monzón-Sandoval, J., Li, Z., Webber, C. and Li, M. 2023. Single-Cell
Transcriptomics and In Vitro Lineage Tracing Reveals Differential Susceptibility of
Human iPSC-Derived Midbrain Dopaminergic Neurons in a Cellular Model of
Parkinson's Disease. *Cells* 12(24), doi: 10.3390/cells12242860

Carlson, M. 2019. org.Hs.eg.db: Genome wide annotation for Human. R package
version 3.8.2.

Chai, H. et al. 2017. Neural Circuit-Specialized Astrocytes: Transcriptomic, Proteomic, Morphological, and Functional Evidence. *Neuron* 95(3), pp. 531-549 e539. doi: 10.1016/j.neuron.2017.06.029

Chandrasekaran, A., Avci, H. X., Leist, M., Kobolák, J. and Dinnyés, A. 2016. Astrocyte Differentiation of Human Pluripotent Stem Cells: New Tools for Neurological Disorder Research. *Frontiers in cellular neuroscience* 10, pp. 215-215. doi: 10.3389/fncel.2016.00215

Chung, N. C. and Storey, J. D. 2015. Statistical significance of variables driving systematic variation in high-dimensional data. *Bioinformatics* 31(4), pp. 545-554. doi: 10.1093/bioinformatics/btu674

Clough, E. and Barrett, T. 2016. The Gene Expression Omnibus Database. *Methods Mol Biol* 1418, pp. 93-110. doi: 10.1007/978-1-4939-3578-9_5

Crompton, L. A., McComish, S. F., Stathakos, P., Cordero-Llana, O., Lane, J. D. and Caldwell, M. A. 2021. Efficient and Scalable Generation of Human Ventral Midbrain Astrocytes from Human-Induced Pluripotent Stem Cells. *Journal of Visualized Experiments* (176), doi: 10.3791/62095

Crompton, L. A., McComish, S. F., Steward, T. G. J., Whitcomb, D. J., Lane, J. D. and Caldwell, M. A. 2023. Human stem cell-derived ventral midbrain astrocytes exhibit a region-specific secretory profile. *Brain Commun* 5(2), p. fcad114. doi: 10.1093/braincomms/fcad114

de Rus Jacquet, A., Tancredi, J. L., Lemire, A. L., DeSantis, M. C., Li, W. P. and O'Shea, E. K. 2021. The LRRK2 G2019S mutation alters astrocyte-to-neuron communication via extracellular vesicles and induces neuron atrophy in a human iPSC-derived model of Parkinson's disease. *Elife* 10, p. 2020.2007.2002.178574. doi: 10.7554/eLife.73062

Deneen, B., Ho, R., Lukaszewicz, A., Hochstim, C. J., Gronostajski, R. M. and Anderson, D. J. 2006. The Transcription Factor NFIA Controls the Onset of Gliogenesis in the Developing Spinal Cord. *Neuron* 52(6), pp. 953-968. doi: 10.1016/j.neuron.2006.11.019

di Domenico, A. et al. 2019. Patient-Specific iPSC-Derived Astrocytes Contribute to
Non-Cell-Autonomous Neurodegeneration in Parkinson's Disease. *Stem cell reports*
12(2), pp. 213-229. doi: 10.1016/j.stemcr.2018.12.011

Dobin, A. et al. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1),
pp. 15-21. doi: 10.1093/bioinformatics/bts635

Duan, D., Fu, Y., Paxinos, G. and Watson, C. 2013. Spatiotemporal expression
patterns of Pax6 in the brain of embryonic, newborn, and adult mice. *Brain Structure*
and Function 218(2), pp. 353-372. doi: 10.1007/s00429-012-0397-2

Endo, F. et al. 2022. Molecular basis of astrocyte diversity and morphology across the
CNS in health and disease. *Science* 378(6619), p. eadc9020. doi:
doi:10.1126/science.adc9020

Eze, U. C., Bhaduri, A., Haeussler, M., Nowakowski, T. J. and Kriegstein, A. R. 2021.
Single-cell atlas of early human brain development highlights heterogeneity of human
neuroepithelial cells and early radial glia. *Nature Neuroscience* 24(4), pp. 584-594.
doi: 10.1038/s41593-020-00794-1

Falk, A. et al. 2012. Capture of neuroepithelial-like stem cells from pluripotent stem
cells provides a versatile system for in vitro production of human neurons. *PLOS ONE*
7(1), p. e29597. doi: 10.1371/journal.pone.0029597

Fan, X. et al. 2020. Single-cell transcriptome analysis reveals cell lineage specification
in temporal-spatial patterns in human cortical development. *Science Advances* 6(34),
p. eaaz2978. doi: doi:10.1126/sciadv.aaz2978

Finak, G. et al. 2015. MAST: a flexible statistical framework for assessing
transcriptional changes and characterizing heterogeneity in single-cell RNA
sequencing data. *Genome Biology* 16(1), p. 278. doi: 10.1186/s13059-015-0844-5

Gabay, L., Lowell, S., Rubin, L. L. and Anderson, D. J. 2003. Deregulation of
dorsoventral patterning by FGF confers trilineage differentiation capacity on CNS stem
cells in vitro. *Neuron* 40(3), pp. 485-499. doi: 10.1016/s0896-6273(03)00637-8

Hedegaard, A., Monzón-Sandoval, J., Newey, S. E., Whiteley, E. S., Webber, C. and
Akerman, C. J. 2020. Pro-maturational Effects of Human iPSC-Derived Cortical
Astrocytes upon iPSC-Derived Cortical Neurons. *Stem cell reports* 15(1), pp. 38-51.
doi: https://doi.org/10.1016/j.stemcr.2020.05.003

Holmqvist, S. et al. 2015. Generation of human pluripotent stem cell reporter lines for
the isolation of and reporting on astrocytes generated from ventral midbrain and
ventral spinal cord neural progenitors. *Stem Cell Research* 15(1), pp. 203-220. doi:
https://doi.org/10.1016/j.scr.2015.05.014

Houweling, A. C., Dildrop, R., Peters, T., Mummenhoff, J., Moorman, A. F. M., Rüther,
U. and Christoffels, V. M. 2001. Gene and cluster-specific expression of the Iroquois
family members during mouse development. *Mechanisms of Development* 107(1), pp.
169-174. doi: https://doi.org/10.1016/S0925-4773(01)00451-8

Itoh, N. et al. 2018. Cell-specific and region-specific transcriptomics in the multiple
sclerosis model: Focus on astrocytes. *Proceedings of the National Academy of*
Sciences 115(2), pp. E302-E309. doi: doi:10.1073/pnas.1716032115

Jaeger, I. et al. 2011. Temporally controlled modulation of FGF/ERK signaling directs
midbrain dopaminergic neural progenitor fate in mouse and human pluripotent stem
cells. *Development* 138(20), pp. 4363-4374. doi: 10.1242/dev.066746

Jain, M., Armstrong, R. J., Tyers, P., Barker, R. A. and Rosser, A. E. 2003. GABAergic
immunoreactivity is predominant in neurons derived from expanded human neural
precursor cells in vitro. *Exp Neurol* 182(1), pp. 113-123. doi: 10.1016/s0014-
4886(03)00055-4

Kamath, T. et al. 2022. Single-cell genomic profiling of human dopamine neurons
identifies a population that selectively degenerates in Parkinson's disease. *Nature*
Neuroscience 25(5), pp. 588-595. doi: 10.1038/s41593-022-01061-1

Koch, P., Opitz, T., Steinbeck, J. A., Ladewig, J. and Brüstle, O. 2009. A rosette-type,
self-renewing human ES cell-derived neural stem cell with potential for in vitro
instruction and synaptic integration. *Proc Natl Acad Sci U S A* 106(9), pp. 3225-3230.
doi: 10.1073/pnas.0808387106

Kostuk, E. W., Cai, J. and Iacovitti, L. 2019. Subregional differences in astrocytes
underlie selective neurodegeneration or protection in Parkinson's disease models in
culture. *Glia* 67(8), pp. 1542-1557. doi: 10.1002/glia.23627

Krencik, R., Weick, J. P., Liu, Y., Zhang, Z.-J. and Zhang, S.-C. 2011. Specification of
transplantable astroglial subtypes from human pluripotent stem cells. *Nature*
Biotechnology 29, p. 528. doi: 10.1038/nbt.1877

https://www.nature.com/articles/nbt.1877#supplementary-information

La Manno, G. et al. 2016. Molecular Diversity of Midbrain Development in Mouse,
Human, and Stem Cells. *Cell* 167(2), pp. 566-580.e519. doi:
10.1016/j.cell.2016.09.027

Li, Y. et al. 2023. Spatiotemporal transcriptome atlas reveals the regional specification
of the developing human brain. *Cell* 186(26), pp. 5892-5909.e5822. doi:
https://doi.org/10.1016/j.cell.2023.11.016

Liao, Y., Smyth, G. K. and Shi, W. 2013. The Subread aligner: fast, accurate and
scalable read mapping by seed-and-vote. *Nucleic Acids Res* 41(10), p. e108. doi:
10.1093/nar/gkt214

Lin, Y.-T. et al. 2018. APOE4 Causes Widespread Molecular and Cellular Alterations
Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell
Types. *Neuron* 98(6), pp. 1141-1154.e1147. doi:
https://doi.org/10.1016/j.neuron.2018.05.008

Liu, Y. et al. 2004. CD44 expression identifies astrocyte-restricted precursor cells.
Developmental Biology 276(1), pp. 31-46. doi:
https://doi.org/10.1016/j.ydbio.2004.08.018

Lozzi, B., Huang, T. W., Sardar, D., Huang, A. Y. S. and Deneen, B. 2020. Regionally
Distinct Astrocytes Display Unique Transcription Factor Profiles in the Adult Brain.
Frontiers in Neuroscience 14, doi: 10.3389/fnins.2020.00061

Makarava, N., Chang, J. C.-Y., Kushwaha, R. and Baskakov, I. V. 2019. Region-
Specific Response of Astrocytes to Prion Infection. *Frontiers in Neuroscience* 13, doi:
10.3389/fnins.2019.01048

Matsunaga, E., Araki, I. and Nakamura, H. 2001. Role of Pax3/7 in the tectum
regionalization. *Development* 128(20), pp. 4069-4077. doi: 10.1242/dev.128.20.4069

McCarthy, D. J., Campbell, K. R., Lun, A. T. L. and Wills, Q. F. 2017. Scater: pre-
processing, quality control, normalization and visualization of single-cell RNA-seq data
in R. *Bioinformatics* 33(8), pp. 1179-1186. doi: 10.1093/bioinformatics/btw777

Molofsky, A. V. et al. 2012. Astrocytes and disease: a neurodevelopmental
perspective. *Genes & Development* 26(9), pp. 891-907. doi: 10.1101/gad.188326.112

Morel, L. et al. 2017. Molecular and Functional Properties of Regional Astrocytes in
the Adult Brain. *J Neurosci* 37(36), pp. 8706-8717. doi: 10.1523/JNEUROSCI.3956-
16.2017

Nolbrant, S., Heuer, A., Parmar, M. and Kirkeby, A. 2017. Generation of high-purity
human ventral midbrain dopaminergic progenitors for in vitro maturation and
intracerebral transplantation. *Nature Protocols* 12(9), pp. 1962-1979. doi:
10.1038/nprot.2017.078

Oberheim, N. A. et al. 2009. Uniquely hominid features of adult human astrocytes. *J
Neurosci* 29(10), pp. 3276-3287. doi: 10.1523/jneurosci.4707-08.2009

Peteri, U.-K., Pitkonen, J., Utami, K. H., Paavola, J., Roybon, L., Pouladi, M. A. and
Castrén, M. L. 2021. Generation of the Human Pluripotent Stem-Cell-Derived
Astrocyte Model with Forebrain Identity. *Brain sciences* 11(2), p. 209. doi:
10.3390/brainsci11020209

Phatnani, H. and Maniatis, T. 2015. Astrocytes in neurodegenerative disease. *Cold*
Spring Harbor perspectives in biology 7(6), p. a020628. doi:
10.1101/cshperspect.a020628

Polioudakis, D. et al. 2019. A Single-Cell Transcriptomic Atlas of Human Neocortical
Development during Mid-gestation. *Neuron* 103(5), pp. 785-801.e788. doi:
10.1016/j.neuron.2019.06.011

Roybon, L. et al. 2013. Human stem cell-derived spinal cord astrocytes with defined
mature or reactive phenotypes. *Cell Rep* 4(5), pp. 1035-1048. doi:
10.1016/j.celrep.2013.06.021

Schindelin, J. et al. 2012. Fiji: an open-source platform for biological-image analysis.
Nature Methods 9(7), pp. 676-682. doi: 10.1038/nmeth.2019

Schlicker, A., Domingues, F. S., Rahnenführer, J. and Lengauer, T. 2006. A new
measure for functional similarity of gene products based on Gene Ontology. *BMC*
Bioinformatics 7(1), p. 302. doi: 10.1186/1471-2105-7-302

Schober, A. L., Wicki-Stordeur, L. E., Murai, K. K. and Swayne, L. A. 2022.
Foundations and implications of astrocyte heterogeneity during brain development
and disease. *Trends in neurosciences* 45(9), pp. 692-703. doi:
10.1016/j.tins.2022.06.009

Serio, A. et al. 2013. Astrocyte pathology and the absence of non-cell autonomy in an
induced pluripotent stem cell model of TDP-43 proteinopathy. *Proceedings of the*
National Academy of Sciences of the United States of America 110(12), pp. 4697-
4702. doi: 10.1073/pnas.1300398110

Sloan, S. A. et al. 2017. Human Astrocyte Maturation Captured in 3D Cerebral Cortical
Spheroids Derived from Pluripotent Stem Cells. *Neuron* 95(4), pp. 779-790 e776. doi:
10.1016/j.neuron.2017.07.035

Sonninen, T.-M. et al. 2020. Metabolic alterations in Parkinson's disease astrocytes.
Scientific Reports 10(1), p. 14474. doi: 10.1038/s41598-020-71329-8

Stirling, D. R., Swain-Bowden, M. J., Lucas, A. M., Carpenter, A. E., Cimini, B. A. and
Goodman, A. 2021. CellProfiler 4: improvements in speed, utility and usability. *BMC*
Bioinformatics 22(1), p. 433. doi: 10.1186/s12859-021-04344-9

Stolt, C. C., Lommes, P., Sock, E., Chaboissier, M.-C., Schedl, A. and Wegner, M.
2003. The Sox9 transcription factor determines glial fate choice in the developing
spinal cord. *Genes & Development* 17(13), pp. 1677-1689. doi: 10.1101/gad.259003

Strelau, J. et al. 2000. Growth/differentiation factor-15/macrophage inhibitory cytokine-
1 is a novel trophic factor for midbrain dopaminergic neurons in vivo. *J Neurosci*
20(23), pp. 8597-8603. doi: 10.1523/jneurosci.20-23-08597.2000

Stuart, T. et al. 2019. Comprehensive Integration of Single-Cell Data. *Cell* 177(7), pp.
1888-1902.e1821. doi: 10.1016/j.cell.2019.05.031

Sun, Y. et al. 2008. Long-term tripotent differentiation capacity of human neural stem
(NS) cells in adherent culture. *Mol Cell Neurosci* 38(2), pp. 245-258. doi:
10.1016/j.mcn.2008.02.014

Supek, F., Bošnjak, M., Škunca, N. and Šmuc, T. 2011. REVIGO Summarizes and
Visualizes Long Lists of Gene Ontology Terms. *PLOS ONE* 6(7), p. e21800. doi:
10.1371/journal.pone.0021800

Takata, N. and Hirase, H. 2008. Cortical Layer 1 and Layer 2/3 Astrocytes Exhibit
Distinct Calcium Dynamics In Vivo. *PLOS ONE* 3(6), p. e2525. doi:
10.1371/journal.pone.0002525

Tcw, J. et al. 2017. An Efficient Platform for Astrocyte Differentiation from Human
Induced Pluripotent Stem Cells. *Stem cell reports* 9(2), pp. 600-614. doi:
10.1016/j.stemcr.2017.06.018

Team, R. C. 2023. : A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing, .

Verkhratsky, A. and Nedergaard, M. 2018. Physiology of astroglia. *Physiological*
Reviews 98(1), pp. 239-389. doi: 10.1152/physrev.00042.2016

Xin, W., Schuebel, K. E., Jair, K.-w., Cimbro, R., De Biase, L. M., Goldman, D. and
Bonci, A. 2019. Ventral midbrain astrocytes display unique physiological features and
sensitivity to dopamine D2 receptor signaling. *Neuropsychopharmacology* 44(2), pp.
344-355. doi: 10.1038/s41386-018-0151-4

Yun, W. et al. 2019. Generation of Anterior Hindbrain-Specific, Glial-Restricted
Progenitor-Like Cells from Human Pluripotent Stem Cells. *Stem Cells and*
Development 28(10), pp. 633-648. doi: 10.1089/scd.2019.0033

Zhong, S. et al. 2018. A single-cell RNA-seq survey of the developmental landscape
of the human prefrontal cortex. *Nature* 555(7697), pp. 524-528. doi:
10.1038/nature25980

Zhou, S. et al. 2016. Neurosphere Based Differentiation of Human iPSC Improves
Astrocyte Differentiation. *Stem Cells Int* 2016, p. 4937689. doi: 10.1155/2016/4937689

Zhu, Q., Shah, S., Dries, R., Cai, L. and Yuan, G. C. 2018. Identification of spatially
associated subpopulations by combining scRNAseq and sequential fluorescence in
situ hybridization data. *Nat Biotechnol* 36(12), pp. 1183-1190. doi: 10.1038/nbt.4260