1

Sociality shapes the life histories of animals along the fast-slow continuum, but does not affect their senescence

4　　　　　　　　　　　　　　Roberto Salguero-Gómez[1,*]

1 Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RB, United

Kingdom, ORCID ID 0000-0002-6085-4433

* Author for correspondence: rob.salguero@biology.ox.ac.uk

Total word count: 5,247

Total figures: 5

Total tables: 1

Total references: 116

SOMs: 1

Abstract (199/200 words)

The role of sociality on the demography of animal species has become an intense focus of research in recent decades. However, efforts to understand the sociality-demography nexus have focused on single species or isolated taxonomic groups. Consequently, we lack generality regarding how sociality shapes demographic traits along the Animal Kingdom. Here, I propose a continuum of sociality, from solitary to tightly social, and test whether it predicts key demographic properties of 152 species, from jellyfish to humans. After correction for body mass and phylogenetic relationships, I show that the sociality continuum predicts key life history traits: more social species live longer, postpone maturity, have greater generation time, and greater probability of achieving reproduction than solitary, gregarious, communal, or colonial species. The sociality continuum is positively correlated with the fast-slow continuum. However, contrary to predictions, sociality does not result in more buffered populations: more social species have a lower ability to benefit from disturbances, although display greater resistance than more solitary species. Finally, I also show that sociality does not shape reproductive or actuarial senescence rates. This cross-taxonomic examination of sociality across the demography of 13 taxonomic classes highlights keyways in which individual interactions shape most aspects of animal demography.

Keywords: eusocial, buffering, generation time, gregarious, solitary, colonial, communal, resilience, senescence.

Introduction

Quantifying the links between sociality and demography is crucial to understand and forecast the dynamics of animal wildlife populations (1,2). Indeed, social structures can influence various aspects of an organism's life, including its survival (3,4) and reproduction (5,6). The role of sociality in shaping the demography of animal species has gained increasing attention in recent years (7–9), as we strive to unravel the mechanisms underlying population fluctuations and viability (10,11).

Social interactions within a group can significantly influence the founding blocks of organismal fitness: the survival and reproduction of individuals in the group. For instance, cooperative breeding, where individuals assist in the care and rearing of offspring, occurs in many animal species, from meerkats (*Suricata suricatta*; (12) to fairy-wrens (*Malurus cyaneus;* (13). This behaviour may enhance reproductive success by optimising resource allocation (14) and providing protection against predators (15). Conversely, in some species, competition for mates within social groups can result in skewed reproductive success (16,17). Survival patterns within a population too are intricately linked to sociality. Indeed, social animals often experience reduced predation risk through group vigilance (18) and cooperative defence mechanisms (19). However, social hierarchies can also increase competition (20), and lower survival (21,22). As such, sociality has the potential to drastically affect the demography of animal species. Understanding how these contrasting factors interact is essential for predicting and managing population dynamics in socially structured species (23).

Our understanding of how sociality shapes the demography of animals lacks generality. Specifically, we lack a comprehensive overview of how sociality associates with various demographic attributes of animal populations across a wide range of taxonomic classes -

but see (24) for a cross-taxonomic examination of reproductive monopolisation across

ca. 300 species of wasps, ants, birds, and mammals. Most social comparative studies

have focused only on birds and mammals (25–28). In these studies, sociality has been

linked to enhanced reproductive success (29), and only weakly with reproductive

senescence (27). Within primates, the impacts of sociality are modulated via complex

group hierarchies, where dominance influences access to resources, mating

opportunities, and reproductive success (30–32). Among the carnivora, social species

like lions (*Panthera leo*) and spotted hyenas (*Crocuta crocuta*) display complex group

structures that influence demographic patterns. In them, cooperative hunting and

communal care of young contribute to the survival and overall fitness of individuals within

the social group (33). In eusocial insects (*e.g.*, ants, bees, and termites), the within-

colony division of labour affects demographic traits via resource allocation, colony

defence, and reproduction (34,35). In certain fish species, such as the clownfish

(*Amphiprion* spp.) and cleaner wrasses (*Labroides* spp.), sociality is linked to improved

foraging efficiency, reduced predation risk, and increased reproductive success (36).

Sociality is multifaceted. As such, different authors define the degree of sociality

depending on the taxonomic group under examination. The lack of cross-taxonomic

categorisation of sociality, together with the historical scarcity of demographic data

across the Tree of Life, I argue, is why a broader examination of sociality across the

demographies of the Animal Kingdom does not exist. However, taking inspiration from

the Plant Kingdom, one could think about the individuals in a group of animals as an

analogous organisation to the xylem vessels that connect the roots and shoots in an

individual vascular plant. The degree of integration-modularity of these individuals

(vessels) within the animal group (individual plant) provides a framework to categorise

sociality across the Animal Kingdom. Indeed, the degree of integration-modularity across

138 plant species strongly correlates with their longevity and rate of actuarial
senescence (37).

Here, I propose a continuum of animal sociality to examine whether sociality, the way
individual organisms organise themselves within a population and interact, shapes their
demography. My sociality continuum contains five categories: (1) solitary: individuals
spend their time alone, except to breed (*e.g.*, tigers (*Panthera tigris*), cheetahs (*Acinonyx*
jubatus), some wasps); (2) gregarious: individuals spend time in groups but social
interactions are loose (*e.g.*, wildebeests, zebras, flock-forming birds); (3) communal:
individuals live together in close proximity and often share a common nesting or dwelling
area, but do not engage in cooperative breeding (*e.g.*, purple martin (*Progne subis*)); (4)
colonial: individuals live in close proximity and always share a common nesting or living
area (*e.g.*, nesting birds, some wasps, coral polyps); and (5) social: individuals live in
close proximity and form stable, organised groups, engaging in social behaviours such
as cooperative breeding and hierarchical structures (*e.g.*, African elephant (*Loxodonta*
africana), most primates, dolphins, meerkat, honeybee (*Apis mellifera*)).

I apply this sociality continuum to 152 species across 13 taxonomic classes, from
cnidarians to mammals (Fig. 1), to carry out phylogenetic comparative analyses to
examine the demographic correlates of sociality across their life history traits (38), vital
rates and impacts on population growth rate (39), and metrics of long-term and short-
term performance (40). My multi-level examination of the demographic traits of these
species allows me to test expectations regarding the impacts of sociality on longevity and
senescence (41) and the 'social buffer' hypothesis (42), whereby more social organisms
should be able to better mitigate environmental stochasticity while also displaying lower
temporal oscillations in population growth rate.

Methods

To test the aforementioned hypotheses, I ran a series of comparative phylogenetic analyses based on high resolution demographic, body mass, sociality, and phylogenetic data of animal species across multiple taxonomic classes.

Demographic data

For the demographic data, I used the COMADRE Animal Matrix Database (43). COMADRE is an open-access database that contains high-resolution demographic information of animal species, mostly from peer-reviewed publications. In its current version, COMADRE (v. 4.32.1) contains 3,488 matrix population models (MPMs, hereafter) from 429 animal studies across 415 studies. An MPM is a numerical array that describes the demography of a species at a given location, time period, and environmental conditions in discrete time and along discrete life cycle stages (44). Specifically, in an MPM, the life cycle of a given population is discretised into stages (*e.g.*, age, discrete size ranges, developmental stages), such that the rates that are then stored in the MPM describe stage-specific survival (σ), growth/development (γ), shrinkage/de-development (ρ), and reproduction (φ) from time t to $t+1$.

To ensure comparability across the thousands of MPMs available in COMADRE, I imposed the following series of selection criteria using the *Rcompadre* R package (45): (1) Unmanipulated MPMs from wild populations, so that the derived demographic traits reflect the dynamics of the population under natural conditions, rather than under experimental treatments or in captivity; (2) MPMs with dimension equal to or greater than 4 × 4 (*i.e.*, 4 or more stages in the life cycle), so as to allowing for enough resolution in the life cycle to successfully discern age-specific patterns of survivorship and reproduction (46,47) and avoid mortality and fertility plateaus (48); (3) MPMs where the

rates of survival and reproduction had been separated into sub-matrices (such that the

overall MPM $A = U + F$, where U and F represent the sub-matrices of survival and

reproduction, respectively), to facilitate the calculation of life history traits and vital rates

and their elasticities to population growth rate; (4) MPMs whose stage-specific survival

values ≤ 1, to prevent erroneous calculations of inflated longevity and actuarial

senescence (47); and (5) MPMs corresponding to extant species (*e.g.*, a study

successfully passed through selection criteria 1-4, above, but in fact corresponds to an

extinct porcupine, *Hystrix refossa (49)*). These criteria resulted in 154 animal species, for

which 23 species had at least two available studies in COMADRE.

Next, for those 23 species with multiple studies available, I manually chose the single

study per species that provides higher representativity of the demography of the species

of interest by retaining the longer-term study, the MPM with greater dimensionality,

and/or the study where the rates of survival and/or reproduction in adults were not

assumed constant. For each of the resulting 154 animal species, I retained the 'grand

mean' MPM that described the dynamics of the population across the length of the study

by calculating the element-by-element arithmetic mean MPM, thus resulting in one MPM

per species. In addition, I supplemented my dataset with a recent study on meerkat

demography (50), not yet incorporated in COMADRE. For the 10 years of seasonal

MPMs available in this study, I retained the MPMs corresponding to 1998-2002,

corresponding to normal environmental conditions (M. Paniw, pers. comm.), and then

again estimated the element-by-element arithmetic grand mean MPM. This addition

increased the count to 155 animal species. The vast majority of MPMs were

parameterised with data collected once per year (n = 130, 84%; Fig. S1). However, to

154 make sure that all demographic traits derived from the MPMs (below) are on the same

155 time units, MPMs not on an annual periodicity were rescaled accordingly by elevating

156 each MPM element $a_{i,j}$ in **A** to the power of $1/P$, where P is the projection interval of the

157 MPM, which is also archived in COMADRE (43). The original sources of the individual

158 demographic studies used in this paper are shown in Table S1.

159 For each species' MPM, I derived a series of demographic traits that inform on the life

160 history traits, vital rates and their relative impacts on population growth rate, as well as

161 metrics of asymptotic (*i.e.*, long-term) and transient (*i.e.*, short-term) performance. To do

162 so, I used the R packages *Rage* (45), *popdemo* (51) and *popbio* (52).

163 Life history traits are key metrics that define specific moments and periods of time along

164 the life cycle of a species (38). The chosen life history traits describe ~70-80% of the

165 variation in life history strategies of animals (53–57). These are: generation time (T; (55),

166 the average age of reproductive individuals in a population; age at maturity (L_a; (58);

167 mean (η_e; (44)) and variance in life expectancy ($\Delta\eta_e$; (44); actuarial senescence,

168 quantified via the rate (shape) of age-specific survivorship (s_{lx}; (59,60); probability of

169 achieving reproduction before dying (p_R; (44); reproductive senescence, quantified via

170 the rate (shape) of age-specific reproduction (s_{mx}; (61); and Demetrius' entropy (S; (62),

171 which quantifies the degree of semelparity (individuals in the population reproduce once;

172 S = 1) *vs.* iteroparity (reproduce multiple times; S>>1) (63).

173 To further characterise potential differences that sociality might have on the demography

174 of species, I examined vital rates and their relative impacts on population growth rate.

175 Using the Rage R package (45), I estimated an average value of survival (σ), growth (γ),

176 shrinkage (ρ), and reproduction (φ), weighted by the stationary distribution of individuals

177 in the population (63). Next, I calculated the elasticities of population growth rate (λ) to

178 these vital rates. These elasticities quantify the effect that a relative, infinitesimally small

change in a vital rate would have on the long-term, asymptotics of the population (39,64),

quantified as the dominant eigenvector of the MPM **A**, λ (44) (not to confuse with Pagel's

λ, an estimate of phylogenetic signal - below). Next, I also quantified the population

growth rate (λ), and the deviance of population growth rate from $\lambda=1$ as $|1-\lambda|$, at which

point the population is neither increasing ($\lambda>1$) nor declining ($\lambda>1$). Deviance of $\lambda=1$ was

of particular interest here, as the social buffer hypothesis (42) predicts the canalisation of

λ towards demographic stability (λ close to 1) with increases in the degree of sociality.

Finally, I also calculated metrics of transient dynamics. Transient dynamics in an

structured population model, such as an MPM, quantify the short-term responses of a

population to potential disturbances across the structure of said population (40,65). The

full range of potential responses to disturbances creates a so-called transient envelope

(40), from which one can calculate the rate of recovery to the stationary equilibrium

(damping ratio, ζ), the number of oscillations that the population undergoes before

achieving said equilibrium (period of oscillation, P_i), the maximum increase in size that

the population can undertake following a disturbances (reactivity, $\overline{\rho}_1$), and the maximum

decrease in size that the population can display after a disturbance (first step

attenuation, $\underline{\rho}_1$). I note that many other transient metrics can be used to define the

transient envelope of an MPM (40), but four I chose here tend to be uncorrelated (66,67),

and together they inform on the inherent ability of a population to respond to

disturbances, that is, their demographic resilience (68). The vital rate elasticities, and

metrics of asymptotic and transient performance were only calculated for MPMs that are

reducible, primitive, and ergodic, thus guaranteeing the existence of a single dominant

eigenvalue, λ_1.

<u>Adult body mass data</u>

Most life history traits scale with adult body mass (53–56). As such, to explore the

relationships between sociality and life history traits beyond the blurring effect of body

mass, I first obtained adult body mass data for each species. For mammals, I used

information archived in (69); for birds, AVONET (70); for fish, FishBase (71) via

rFishBase (72); for mammals, birds, and reptiles and amphibians, AMNIOTE (73); and

for remaining groups, MOSAIC (74), and data from (54). For 18 remaining species for

which, adult body mass data were not available from these online databases (mostly

insects, bivalves, and corals), I obtained body mass information from peer-review

publications using the keywords "body mass" OR "weight" and the name of the species in

ISI WoS. When pertinent, all data were transformed to grams across all sources. The

final variable was log-transformed to fulfil assumptions of the statistical analyses

employed below.

Sociality data

No centralised, open-access database exists to my knowledge offering a

comprehensive categorisation of the sociality of species across kingdom Animalia. As

such, I classified sociality following the continuum described in the introduction by first

contrasting information from the following online resources for each species in my

dataset: Animal Diversity Web (75), Smithsonian's National Zoo and Conservation

Biology   Institute   (https://nationalzoo.si.edu/),   FishBase

(https://www.fishbase.se/search.php), IUCN Red List of Threatened Species (76), and

chatGPT (https://chat.openai.com/) - for the latter, suggested references were cross-

checked and validated via ISI WoK. Next, I double-checked existing classifications in my

data and complemented missing information from peer-review publications for each of

the 154 animal species. Once the list was completed, I discussed it with experts in each

of the taxonomic classes, and made the appropriate corrections (Table S1).

Phylogenetic data

The analyses below explicitly consider how our species are related phylogenetic. To that end, I obtained a phylogeny from the Open Tree of Life (77) via the R package *ROTL* (78). To do so, I first matched the scientific names of my subset of animal species from COMADRE to the Taxonomic Name Resolution Service via the function *tnrs_match_names*. All species matched successfully, except the two species from class Demospongiae (*Spongia graminea*, and *Xestospongia muta*). As the phylogenetic tree of Demospongiae is not well resolved (79), I decided to proceed with the analyses discarding these two species, and so the final analyses include 152 animal species, instead of 154.

Comparative phylogenetic analyses

To examine the correlates of sociality with the demographic traits examined from the MPMs in COMADRE, I first run a phyloANOVA via the *phytools* R package (80). For each demographic trait, the response variable was its residual from the pgls regression over adult body mass, and the explanatory variable was the factor variable sociality continuum, with its five different levels, ordered as "solitary", "gregarious", "communal", "colonial", and "social". To correct for type II errors, after the tests were performed, I also implemented a Bonferroni correction with the function *p.adjust* from the *Stats* R package. The phylogenetically-corrected, body mass-corrected post-hoc Tukey test that I implemented next explicitly allowed me to test the hypothesis that the sociality continuum ranks, which ranks species along increasing degree of social interactions, associates with different life history traits, vital rates and their elasticities of population growth rate, as well as metrics of asymptotic and transient dynamics.

The phylogenetic ANOVAs performed here allowed me to examine the correlates of

sociality with each life history trait separately. However, life history traits are often

correlated into life history strategies (38). Indeed, previous works have demonstrated that

animals show two primary axes of variation: one where individuals either develop fast or

live long, the so-called fast-slow continuum (38,53,56,81), and a second one along which

species differ in their reproductive tactics, the so-called reproductive strategies

continuum (10,54,81).

To quantify the life history strategy space of the 152 study animal species, I used a

phylogenetic principal component analysis (pPCA). Briefly, PCA is a family of

multivariate statistical techniques used to examine complex data by reducing

dimensionality of the data to highlight the main factors that explain observed variation

(82). The pPCA takes into consideration the non-independence of life history traits due to

species relatedness (83,84). To implement the pPCA, I first log-transformed the life

history traits to fulfil assumptions of error normal distribution necessary in linear models.

Next, I calculated the residuals of each life history trait as a function of species' adult

body mass via a pgls with the R package caper (85) and the phylogeny. Next, I rescaled

the residuals to mean (μ) = 0 and variance (σ) = 1. This approach allowed to establish

primary axes of variation of, in this case, life history traits, as well as to estimate the

overall strength of the phylogenetic relationships among species. For the latter, I also

estimated Pagel's λ, a parameter that ranges between 0 (the pattern of association is not

explained by the phylogenetic hypothesis), and 1 (the pattern is entirely explained by

phylogenetic inertia).

Because PCA approaches require a dataset with no missing values, and my data were

somewhat gappy, I carried out imputations. The gappiness of the eight life history traits

was 19.62% (Fig. S2). I used a well-established protocol (53,54,63,81) to recover the

missing information using an iterative PCA approach. Briefly, I employed the

"*regularised*" method in the function *mice* in the R package *mice (86)* with 20 sets of

imputation. To draw strength across the full dataset, the imputation took into account the

vital rate values, elasticities, transient dynamics metrics (described below), and body

mass (above) of each species. Here, an iterative algorithm first assigns the mean of each

life history trait to the missing values of that life history trait. Next, a regular PCA is

performed, and the scores of the species with missing data are re-assigned using the

orthogonal relationships of the PCA. This iteration is repeated until convergence. As a

check, I implemented the same protocol using only the 85 species for which no life

history trait data were missing, and found that the results with imputed data are

insensitive to the imputation (not shown). The insensitivity of the results reported here to

the imputation approach is in agreement with previous comparative demographic

studies that also used COMADRE (10,54,87).

Finally, to examine how the sociality continuum governs the life history strategy space of

animals, I used the scores of the pPCA (above) to run further phylogenetically-corrected

post-hoc Tukey tests to examine if the different levels of the sociality continuum are

significantly clustered along the different PC axes.

Results

295 The phylogenetic signal of my sociality continuum, from solitary species, like the giant

296 panda (*Ailuropoda melanoleuca*; Table S1), to social species, like the red howler

297 (*Alouatta seniculus*), across the 152 examined animal species, was high. Indeed the

298 estimates of Pagel's λ for the sociality continuum across these species is 0.907 (95%

299 C.I.: 0.865-0.945; $P < 0.001$). This strong signal is also visually apparent upon examining

the phylogenetic tree shown in Fig. 1B. The phylogenetic signal of the associations between the sociality continuum and the examined demographic traits, in contrast, was variable (Table 1). The estimates of Pagel's λ from the performed pgls models ranged from virtually absent (*e.g.*, vital rates of growth, shrinkage, and reproduction (0.000), population growth rate (also called λ *(44)*; 0.001), period of oscillation (0.011)), to high (mean life expectancy (0.739), generation time (0.665), age at maturity (0.649), variance in life expectancy (0.668), survival (0.616)).

Across the five different types of demographic traits examined here (Table 1: life history traits, vital rates, vital rate elasticity, asymptotics, and transients), the only category where the sociality continuum did not raise some significant results was the metrics of asymptotic dynamics: population growth rate and deviance growth rate equilibrium. Moreover, in these two metrics, the 95% confidence intervals Pagel' λ for the respective pgls models overlapped with 0, indicating no phylogenetic effect in the relationship between sociality and long-term population performance. I found that, after correcting for adult body mass and phylogenetic relationships, more social species attain longer generation times, longer mean life expectancy, greater variance in life expectancy, and are more likely to reproduce before dying (Table 1A). Interestingly, the relationships between the sociality continuum and species' vital rates were opposed to their elasticities: more social species have an increased ability to shrink and reproduce (Table 1B), but their population growth rates would increase the most if their vital rates of survival and growth would increase (Table 1C). Finally, for the transient metrics, I found that more social species tended to display greater periods of oscillation and higher first step attenuation (*i.e.*, inverse of resistance to disturbances, *sensu* (68)).

The phylogenetic ANOVA post-hoc Tukey tests revealed important differences in the examined demographic traits as a function of the sociality continuum. For instance,

increases in sociality (solitary → gregarious → communal → colonial → social) were

associated with prolonged generation time, mean and variance of life expectancy,

probability to attain reproduction before dying, but not with age at maturity, reproductive

senescence, or Demetrius' entropy (Fig. 2). This pattern was not always gradual and

monotonic. For instance, the rate of senescence was highest (implying a lower rate of

senescence) for communal species, and lowest for gregarious species, though not

significantly so (Fig. 2). Social species also had higher survival as well as lower

shrinkage and reproduction compared to all other categories (Fig. 3). These patterns

remained largely consistent when evaluating the elasticities of population growth rate to

said vital rates. Finally, for the transient metrics, as the degree of sociality increased, so

did the period of oscillation and first step attenuation, and decreased species' reactivity

(Fig. 4).

The life history strategy space defined by the eight examined life history traits (Table 1A)

identified two principal components whose associated eigenvalues > 1. Following

Kaiser's criterion (88), I retained these two axes for the next part of the analysis. PC1

explains 27.76% of the variation in life history traits across the 152 animal species, and it

separates fast-living (left of Fig. 5) from slow-living species. Indeed, species to the right

in the PCA live long (η_e) but with high variance in life expectancy ($\Delta\eta_e$), postpone the first

reproductive event (L_α), and have a high generation time (T) (right of PCA space). PC2

explains 26.94% of the variation, and separates species that senesce fast for

survivorship (s_{lx}) and reproduction (s_{mx}) (bottom), from those that do not, and that in

addition have a high probability of reproducing before dying (p_R), and are highly

iteroparous (S). The emergent phylogenetic signal, after adult body mass correction, in

this phylogenetic PCA was relatively low, with an estimate of Pagel's λ of 0.220 (0.034-

0.289 95% C.I.).

The ranking of species along the sociality continuum raised significant differences along

PC1 (the fast-slow continuum; *P* = 0.021; Fig. 5). Indeed, social species like the Northern

muriqui (*Brachyteles hypoxanthus*) or the White-faced capuchin monkey (*Cebus*

capucinus) species, and colonial species like the Southern fulmar (*Fulmarus glacialis*) or

the black vulture (*Coragys atratus*), are associated with slower life histories, compared to

solitary species like the lake chubsucker (*Erimyzon sucetta*), gregarious species like the

European hare (*Lepus europaeus*), or communal species like the American crocodile

(*Crocodylus acutus*). I found no association between the sociality continuum and the

reproductive strategies continuum, PC2 (*P* = 0.738).

Discussion

Sociality, the degree to which animals engage in interactions and form cohesive

structures, can shape their survival (89) and reproduction (16), the building blocks of

fitness. Animals show a wide range of social behaviours, including cooperative breeding

(13), group living (3), and hierarchical structures to better protect their young (32), their

territory (90), *etc*. Though much research has examined the demographic correlates of

animal sociality, it is constrained by taxonomic siloes. Indeed, much animal sociality

research focuses on single species, or on mammals (4) and birds (27). Here, by

proposing a sociality continuum that ranks species according to the type and duration of

their interactions, from solitary, to gregarious, to communal, to colonial, to social, I show

that, across 152 animal species from 13 taxonomic classes, as the intensity of these

interactions increases, species life histories become *slower* (38), even after accounting

for the allometric effects of body mass (54,66) and potential constraints of phylogenetic

inertia (83,84). My comparative phylogenetic analyses also show that the intensity of

these social interactions does not shape long-term metrics of performance, such as the

population growth rate, contrary to the 'social buffer' hypothesis (42), but that the

sociality continuum is associated with some key attributes of demographic resilience

(65).

A wealth of scientific evidence highlights the profound impacts of sociality on life history

traits in animals. Indeed, social structures often influence reproductive strategies, with

cooperative breeding systems, for example, promoting delayed sexual maturity and

extended parental care (91). In species exhibiting eusociality, such as certain bees and

ants, caste systems shape life history trajectories for individuals within the colony

(92,93). Sociality can also affect reproductive success, as observed in the intricate

mating systems of various primates (94) and birds (13,95). Moreover, the presence of

social alliances may impact longevity and survivorship, providing advantages in terms of

predation avoidance and access to resources (32,96). Here, I show that the sociality

continuum is associated with key life history traits such as prolonged generation time,

postponed age at maturity, and increased mean and variance of life expectancy. It is

worth noting, however, that the interplay between sociality and life history traits extends

beyond reproduction and survival, influencing factors like dispersal patterns (97) not

taken into account in the present study. To understand this complexity, future work must

consider species-specific adaptations and ecological contexts and their

dispersal/migration abilities, employing a combination of field observations, experimental

manipulations, and modelling approaches based on big data (98).

In the present study, the sociality continuum does not impact the rate of actuarial nor

reproductive senescence across the examined 152 animal species. The impacts of

sociality on senescence in animals has been a subject of extensive research, with

findings offering mixed perspectives (3). Studies on various species indicate that social

interactions can either accelerate (99) or decelerate senescence processes (100). For

instance, in certain cooperative breeding systems, where individuals collectively care for

offspring, sociality may delay reproductive senescence due to shared parental

responsibilities and increased chances of survival for offspring (Kokko et al., 2002;

Clutton-Brock, 2002). Conversely, in some species with intense intrasexual competition,

the pressure to secure mates and resources may accelerate reproductive senescence

(101). Actuarial senescence, reflected in declining survival rates with age, can also be

influenced by social factors. For example, the presence of social allies may enhance

protection against predation, potentially extending lifespan (96). However, the stress

associated with social hierarchies and conflicts can have the opposite effect, accelerating

actuarial senescence (102). The complex interplay between social dynamics and

senescence, and the multiple ways through which senescence can manifest itself (103),

underscores the need for a nuanced understanding that considers species-specific traits

and environmental contexts. Ultimately, a combination of the comparative approach used

here (to identify patterns and plausible proximal mechanisms), coupled with experimental

manipulations (to identify and understand distal mechanisms), will provide the ultimate

answer of the conditions under which sociality accelerates or slows down senescence in

animals.

The relationship between sociality and the ability of animals to withstand environmental

stochasticity has garnered substantial attention. Numerous studies suggest that sociality

can serve as a buffering mechanism (42), providing a range of benefits that enhance

fitness in stochastic environments. While the social buffer hypothesis predicts that

sociality should confer adaptive advantages in mitigating the impact of environmental

stochasticity, I found that sociality is not associated with any metric of long-term

population performance that are often used as population-level proxies for fitness

(44,104). For instance, when evaluating how far from a population growth rate at

equilibrium the 152 species' population were, there was no significant correlation with

sociality. I did, however, find that, while more social species do not take a significantly

different amount of time to recover after a disturbance compared to less social species,

the period of oscillation back to stationary dynamics for the former is greater than in less

social species, implying a potential greater risk of extinction for social species (105,106).

Likewise, the reactivity of social species was lower than in gregarious species,

suggesting that, when a population is hit by a disturbance, it is less likely to take

advantage by increasing its population size (68). The past decades have witnessed the

development of indexes of demographic buffering based on structured population models

(107–110). Future research should evaluate whether the sociality continuum is

intrinsically linked to a potential demographic lability-buffering continuum (111).

Species are more than just social or not social. Thus, a continuum to classify animal

sociality is of high appeal. A previous attempt has, however, highlighted some key

challenges (24). Indeed, sociality is shaped by multiple axes, including group size, the

kind, timing, and hierarchical nature of interactions, or the partition of labour, to mention

a few. The inspiration for my proposed sociality continuum comes from work my

laboratory and I carried out to examine the anatomical correlates of actuarial senescence

(37). Finch (112) hypothesised that the ability to compartmentalise risk may allow

species to postpone the onset of senescence. Classifying the compartmentalisation of

138 plants (proxied by the distance between their xylem vessels) and 151 animals

(proxied by the degree of open-closed circulatory and respiratory systems, and degree of

organ redundancy), we found that modularity does predict actuarial senescence in plants

but not in animals. Our work there was only focused on vascular plants, for which a

standard measure of xylem distance can be obtained (113). The classification of a

continuum of modularity-integration in animals proved difficult then because of the wide

range of variation in animal anatomy. Another way to think about animal anatomy is to

examine the 'anatomy' of their groups and populations: how close are their individuals,

how often do they come in contact, and what kinds of interactions do they have. In this regard, recently (8) that older, more senescent (*i.e.*, likely to die) individuals of red deer (*Cervus elaphus*) become more isolated in the group, perhaps as a way to contain risk too, as plants do (37). My proposed sociality continuum correlates with key aspects of the demography of animals. However, further work should examine orthogonal ways in which animal societies evolve, as the predictive power of the correlations examined here remains low (Table 1).

The future of research on animal sociality and demography should prioritise interdisciplinary approaches to analyse a large number of longitudinal studies. This approach will help us unravel the nuanced complexities of social structures and their demographic implications. Integrating advanced technologies, such as GPS tracking (114), social network analysis (8), and comparative genomics (115), can provide a more comprehensive understanding of the dynamic interactions within animal societies. Embracing comparative analyses across a diverse array of taxa can uncover general principles while acknowledging species-specific nuances. As conservation issues become increasingly urgent, linking social and demographic research to conservation biology can inform strategies for preserving biodiversity in the face of environmental challenges (116). This multidimensional approach holds the key to advancing our understanding of the intricate interplay between sociality and demography in the animal kingdom.

Acknowledgments

I thank J. Firth and K. Davis for conversations regarding the interface of animal sociality in demography; J. Cant and B. Sheldon for input into the classification of sociality of

476   fishes and birds, respectively; the thousands of researchers who have contributed open-

477   access data to the COMADRE Animal Matrix Database (www.compadre-db.org); and M.

478   Paniw and E. Conquet for providing access to the demographic models of *Suricata*

479   *suricatta*. I was supported by a NERC Pushing the Frontiers grant (NE/X013766/1).

References

1. Oro D. Perturbation, Behavioural Feedbacks, and Population Dynamics in Social Animals:
When to Leave and Where to Go [Internet]. Oxford University Press; 2020. 176 p. Available
from: https://play.google.com/store/books/details?id=f4_UDwAAQBAJ

2. Polgar S. Population, Ecology, and Social Evolution [Internet]. Walter de Gruyter; 2011. 364 p.
Available from: https://play.google.com/store/books/details?id=R1S1BwAAQBAJ

3. Lucas ER, Keller L. The co□evolution of longevity and social life. Funct Ecol [Internet]. 2020
Jan;34(1):76–87. Available from: https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-
2435.13445

4. Blumstein DT, Williams DM, Lim AN, Kroeger S, Martin JGA. Strong social relationships are
associated with decreased longevity in a facultatively social mammal. Proc Biol Sci [Internet].
2018 Jan 31;285(1871). Available from: http://dx.doi.org/10.1098/rspb.2017.1934

5. Sakata JT, Gupta A, Chuang CP, Crews D. Social experience affects territorial and
reproductive behaviours in male leopard geckos, Eublepharis macularius. Anim Behav
[Internet].   2002   Mar   1;63(3):487–93.   Available   from:
https://www.sciencedirect.com/science/article/pii/S0003347201919529

6. Amdam GV, Csondes A, Fondrk MK, Page RE Jr. Complex social behaviour derived from
maternal reproductive traits. Nature [Internet]. 2006 Jan 5;439(7072):76–8. Available from:
http://dx.doi.org/10.1038/nature04340

7. Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, et al. Multiple
spatial behaviours govern social network positions in a wild ungulate. Ecol Lett [Internet]. 2021
Apr;24(4):676–86. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13684

8. Albery GF, Clutton-Brock TH, Morris A, Morris S, Pemberton JM, Nussey DH, et al. Ageing
red deer alter their spatial behaviour and become less social. Nature Ecology & Evolution
[Internet]. 2022;6(8):1231–8. Available from: https://www.nature.com/articles/s41559-022-
01817-9

9. Bateman AW, Ozgul A, Krkošek M, Clutton-Brock TH. Matrix Models of Hierarchical
Demography: Linking Group- and Population-Level Dynamics in Cooperative Breeders. Am
Nat [Internet]. 2018 Aug;192(2):188–203. Available from: http://dx.doi.org/10.1086/698217

10.   Paniw M, Maag N, Cozzi G, Clutton-Brock T, Ozgul A. Life history responses of meerkats
to seasonal changes in extreme environments. Science [Internet]. 2019 Feb 8;363(6427):631–
5. Available from: http://dx.doi.org/10.1126/science.aau5905

11.   Paniw M, Duncan C, Groenewoud F, Drewe JA, Manser M, Ozgul A, et al. Higher
temperature extremes exacerbate negative disease effects in a social mammal. Nat Clim
Chang [Internet]. 2022 Feb 7 [cited 2024 Jan 12];12(3):284–90. Available from:
https://www.nature.com/articles/s41558-022-01284-x

12.   Groenewoud F, Clutton-Brock T. Meerkat helpers buffer the detrimental effects of adverse
environmental conditions on fecundity, growth and survival. J Anim Ecol [Internet]. 2021
Mar;90(3):641–52. Available from: http://dx.doi.org/10.1111/1365-2656.13396

13.   Arnold KE, Owens IPF. Cooperative breeding in birds: a comparative test of the life
history hypothesis. Proceedings of the Royal Society of London Series B: Biological Sciences
[Internet].   1998   May   7;265(1398):739–45.   Available   from:
https://doi.org/10.1098/rspb.1998.0355

14.   Koykka C, Wild G. The association between the emergence of cooperative breeding and
clutch size. J Evol Biol [Internet]. 2016 Jan;29(1):58–76. Available from:
http://dx.doi.org/10.1111/jeb.12762

15.   Poiani A, Pagel M. EVOLUTION OF AVIAN COOPERATIVE BREEDING:
COMPARATIVE TESTS OF THE NEST PREDATION HYPOTHESIS. Evolution [Internet].
1997 Feb;51(1):226–40. Available from: http://dx.doi.org/10.1111/j.1558-5646.1997.tb02404.x

16.   Salomon M, Lubin Y. Cooperative breeding increases reproductive success in the social
spider Stegodyphus dumicola (Araneae, Eresidae). Behav Ecol Sociobiol [Internet]. 2007 Sep
1;61(11):1743–50. Available from: https://doi.org/10.1007/s00265-007-0406-2

17.   Poethke HJ, Liebig J. Risk-sensitive foraging and the evolution of cooperative breeding
and reproductive skew. BMC Ecol [Internet]. 2008 Mar 18;8:2. Available from:
http://dx.doi.org/10.1186/1472-6785-8-2

18.   Groenewoud F, Frommen JG, Josi D, Tanaka H, Jungwirth A, Taborsky M. Predation risk
drives social complexity in cooperative breeders. Proc Natl Acad Sci U S A [Internet]. 2016
Apr 12;113(15):4104–9. Available from: http://dx.doi.org/10.1073/pnas.1524178113

19.   Hellmann JK, Hamilton IM. The presence of neighbors influences defense against
predators in a cooperatively breeding cichlid. Behav Ecol [Internet]. 2014;25(2):386–91.
Available from: https://academic.oup.com/beheco/article-abstract/25/2/386/226002

20.   Schoepf I, Schradin C. Better off alone! Reproductive competition and ecological
constraints determine sociality in the African striped mouse (Rhabdomys pumilio). J Anim Ecol
[Internet]. 2012 May;81(3):649–56. Available from: http://dx.doi.org/10.1111/j.1365-
2656.2011.01939.x

21.   Kappeler PM, Cremer S, Nunn CL. Sociality and health: impacts of sociality on disease
susceptibility and transmission in animal and human societies. Philos Trans R Soc Lond B Biol
Sci [Internet]. 2015 May 26;370(1669). Available from:
http://dx.doi.org/10.1098/rstb.2014.0116

22.   Rehan SM, Richards MH, Adams M, Schwarz MP. The costs and benefits of sociality in a
facultatively social bee. Anim Behav [Internet]. 2014 Nov 1;97:77–85. Available from:
https://www.sciencedirect.com/science/article/pii/S0003347214003352

23.   Wong MYL. Abiotic stressors and the conservation of social species. Biol Conserv
[Internet]. 2012 Oct 1;155:77–84. Available from:
https://www.sciencedirect.com/science/article/pii/S0006320712002686

24.   Rubenstein DR, Botero CA, Lacey EA. Discrete but variable structure of animal societies
leads to the false perception of a social continuum. R Soc Open Sci [Internet]. 2016
May;3(5):160147. Available from: http://dx.doi.org/10.1098/rsos.160147

25.   Perony N. Comparative Analysis of Social Interactions in Animal Groups [Internet]. ETH;
2012. Available from: https://play.google.com/store/books/details?id=gedXMwEACAAJ

26.   Beauchamp G. A comparative analysis of vigilance in birds. Evol Ecol [Internet]. 2010
Sep;24(5):1267–76. Available from:
https://idp.springer.com/authorize/casa?redirect_uri=https://link.springer.com/article/10.1007/s
10682-010-9358-5&casa_token=bVvEtk1E5-8AAAAA:XHgDX_1s3QmYP-
UVl8WVvwiHy7SrEcZalln_9tgnJ248QLXiGOLTV8xIGGg1X8tcs1Rk3Yo5EgssJNNc

27.   Vágási CI, Vincze O, Lemaître JF, Pap PL, Ronget V, Gaillard JM. Is degree of sociality
associated with reproductive senescence? A comparative analysis across birds and
mammals. Philos Trans R Soc Lond B Biol Sci [Internet]. 2021 Apr 26;376(1823):20190744.

Available from: http://dx.doi.org/10.1098/rstb.2019.0744

28.   Weinstein M, Kaplan H, Lane MA. Sociality, Hierarchy, Health: Comparative
Biodemography [Internet]. National Academies Press (US); 2014 [cited 2024 Jan 16].
Available from: https://www.ncbi.nlm.nih.gov/books/NBK242461/

29.   Woodroffe R. Demography of a recovering African wild dog (Lycaon pictus) population. J
Mammal [Internet]. 2011 Apr 15 [cited 2024 Jan 12];92(2):305–15. Available from:
https://academic.oup.com/jmammal/article-abstract/92/2/305/863612

30.   Schülke O, Ostner J. Ecological and social influences on sociality. The evolution of
primate   societies   [Internet].   2012;195–219.   Available   from:
https://books.google.com/books?hl=en&lr=&id=gvwy--
ul3CQC&oi=fnd&pg=PA195&dq=sociality+increases+resource+acquisition&ots=N84vQvPVXn
&sig=_Z0We-RlqHraMvpElDrefJMNNSw

31.   Ostner J, Schülke O. Linking sociality to fitness in primates: A call for mechanisms. In:
Naguib M, Barrett L, Healy SD, Podos J, Simmons LW, Zuk M, editors. Advances in the Study
of Behavior [Internet]. San Diego, CA: Elsevier; 2018. p. 127–75. (Advances in the study of
behavior;   vol.   50).   Available   from:
https://www.sciencedirect.com/science/article/pii/S0065345417300220

32.   Silk JB. Social components of fitness in primate groups. Science [Internet]. 2007 Sep
7;317(5843):1347–51. Available from: http://dx.doi.org/10.1126/science.1140734

33.   Bekoff M, Daniels TJ, Gittleman JL. Life history patterns and the comparative social
ecology of carnivores. Annu Rev Ecol Syst [Internet]. 1984 Nov;15(1):191–232. Available
from:
https://www.annualreviews.org/doi/pdf/10.1146/annurev.es.15.110184.001203?casa_token=N
Nv7OYmhpPUAAAAA:mIZTlqV9gWQaDVYI3FyvNGh57NkFvGYghSS_H6M_VzSYofKH5F8q
SQXFRYdiOwRFBi2VkjbUw3VnyQ

34.   Kramer BH, Schrempf A, Scheuerlein A, Heinze J. Ant Colonies Do Not Trade-Off
Reproduction against Maintenance. PLoS One [Internet]. 2015 Sep 18;10(9):e0137969.
Available from: http://dx.doi.org/10.1371/journal.pone.0137969

35.   Chouvenc T, Ban PM, Su NY. Life and Death of Termite Colonies, a Decades-Long Age
Demography Perspective. Frontiers in Ecology and Evolution [Internet]. 2022;10. Available
from: https://www.frontiersin.org/articles/10.3389/fevo.2022.911042

36.   Buston P. Forcible eviction and prevention of recruitment in the clown anemonefish.
Behav Ecol [Internet]. 2003 Jul 1 [cited 2024 Jan 12];14(4):576–82. Available from:
https://academic.oup.com/beheco/article-abstract/14/4/576/211446

37.   Bernard C, Compagnoni A, Salguero-Gómez R. Testing Finch's hypothesis: The role of
organismal modularity on the escape from actuarial senescence. Funct Ecol [Internet]. 2020
Jan;34(1):88–106.   Available   from:
https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-2435.13486

38.   Stearns SC. The Evolution of Life Histories. New York, USA: Oxford University Press;
1999. 249 p.

39.   de Kroon H, van Groenendael J, Ehrlén J. Elasticities: a review of methods and model
limitations. Ecology. 2000 Mar;81(3):607–18.

40.   Stott I, Townley S, Hodgson DJ. A framework for studying transient dynamics of
population projection matrix models. Ecol Lett [Internet]. 2011 Sep;14(9):959–70. Available
from: http://dx.doi.org/10.1111/j.1461-0248.2011.01659.x

41.   Roper M, Capdevila P, Salguero-Gómez R. Senescence: why and where selection
gradients might not decline with age. Proc Biol Sci [Internet]. 2021 Jul
28;288(1955):20210851.   Available   from:
https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2021.0851

42.   Hennessy MB, Kaiser S, Sachser N. Social buffering of the stress response: diversity,
mechanisms, and functions. Front Neuroendocrinol [Internet]. 2009 Oct;30(4):470–82.
Available from: http://dx.doi.org/10.1016/j.yfrne.2009.06.001

43.   Salguero-Gomez R, Jones OR, Archer CR, Bein C, de Buhr H, Farack C, et al.
COMADRE: a global data base of animal demography. J Anim Ecol [Internet]. 2016
Mar;85(2):371–84. Available from: http://dx.doi.org/10.1111/1365-2656.12482

44.   Caswell H. Matrix Population Models: Construction, Analysis, and Interpretation. 2nd
edition. Sinauer Associates, Inc.; 2001.

45.   Jones OR, Barks P, Stott I, James TD, Levin S, Petry WK, et al. Rcompadre and Rage-
Two R packages to facilitate the use of the COMPADRE and COMADRE databases and
calculation of life-history traits from matrix population models. Methods Ecol Evol [Internet].
2022; Available from: http://dx.doi.org/10.1111/2041-210x.13792

46.   Salguero-Gómez R, Plotkin JB. Matrix dimensions bias demographic inferences:
implications for comparative plant demography. Am Nat [Internet]. 2010 Dec;176(6):710–22.
Available from: http://dx.doi.org/10.1086/657044

47.   Jones OJ, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB, et
al. Diversity of ageing across the tree of life. Nature [Internet]. 2014;505:169–73. Available
from: http://dx.doi.org/10.1038/nature12789

48.   Horvitz CC, Tuljapurkar S. Stage dynamics, period survival, and mortality plateaus. Am
Nat [Internet]. 2008 Aug;172(2):203–15. Available from: http://dx.doi.org/10.1086/589453

49.   Monchot H, Fernandez P, Gaillard JM. Paleodemographic analysis of a fossil porcupine
(Hystrix refossa Gervais, 1852) population from the Upper Pleistocene site of Geula Cave
(Mount Carmel, Israel). J Archaeol Sci [Internet]. 2012 Sep 1;39(9):3027–38. Available from:
https://www.sciencedirect.com/science/article/pii/S0305440312001707

50.   Conquet E, Ozgul A, Blumstein DT, Armitage KB, Oli MK, Martin JGA, et al. Demographic
consequences of changes in environmental periodicity. Ecology [Internet]. 2023
Mar;104(3):e3894. Available from: http://dx.doi.org/10.1002/ecy.3894

51.   Stott I, Hodgson DJ, Townley S. popdemo: an R package for population demography
using projection matrix analysis. Methods Ecol Evol [Internet]. 2012 Oct;3(5):797–802.
Available from: http://dx.doi.org/10.1111/j.2041-210X.2012.00222.x

52.   Stubben C, Milligan B. Estimating and analyzing demographic models using the popbio
package in R. J Stat Softw. 2007 Sep;22(11):1–23.

53.   Paniw M, Ozgul A, Salguero-Gomez R. Interactive life-history traits predict sensitivity of
plants and animals to temporal autocorrelation. Ecol Lett [Internet]. 2018 Feb;21(2):275–86.
Available from: http://dx.doi.org/10.1111/ele.12892

54.   Healy K, Ezard THG, Jones OR, Salguero-Gomez R, Buckley YM. Animal life history is
shaped by the pace of life and the distribution of age-specific mortality and reproduction.
Nature Ecology & Evolution [Internet]. 2019 Aug;3(8):1217 – +. Available from:
http://dx.doi.org/10.1038/s41559-019-0938-7

55.   Gaillard JM, Yoccoz NG, Lebreton JD, Bonenfant C, Devillard S, Loison A, et al.

Generation time: a reliable metric to measure life-history variation among mammalian
populations. Am Nat [Internet]. 2005 Jul;166(1):119–23; discussion 124–8. Available from:
http://dx.doi.org/10.1086/430330

56. Gaillard JM, Pontier D, Allaine D, Lebreton JD, Trouvilliez J, Clobert J. An analysis of
demographic tactics in birds and mammals. Oikos [Internet]. 1989 Sep;56(1):59–76. Available
from: http://dx.doi.org/10.2307/3566088

57. Dobson FS, Oli MK. The life histories of orders of mammals: Fast and slow breeding. Curr
Sci [Internet]. 2008;95(7):862–5. Available from: http://www.jstor.org/stable/24103180

58. Cochran ME, Ellner S. Simple methods for calculating age-based life-history parameters
for stage-structured populations. Ecol Monogr. 1992 Sep;62(3):345–64.

59. Keyfitz N, Caswell H. Applied Mathematical Demography. New York; 2005. (Springer
Science & Business Media).

60. Baudisch A. The pace and shape of ageing. Methods Ecol Evol [Internet]. Available from:
https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/j.2041-210X.2010.00087.x

61. Baudisch A, Stott I. A pace and shape perspective on fertility. Methods Ecol Evol
[Internet]. 2019 Nov;10(11):1941–51. Available from:
https://besjournals.onlinelibrary.wiley.com/doi/10.1111/2041-210X.13289

62. Demetrius L. Adaptive value, entropy and survivorship curves. Nature.
1978;275(5677):213–4.

63. Salguero-Gómez R, Jones OR, Jongejans E, Blomberg SP, Hodgson DJ, Mbeau-Ache C,
et al. Fast-slow continuum and reproductive strategies structure plant life-history variation
worldwide. Proc Natl Acad Sci U S A. 2016;113(1):230–5.

64. de Kroon H, Plaisier A, van Groenendael J, Caswell H. Elasticity: the relative contribution
of demographic parameters to population growth rate. Ecology. 1986 Oct;67(5):1427–31.

65. Neubert MG, Caswell H, Solow AR. Detecting reactivity. Ecology [Internet]. 2009
Oct;90(10):2683–8. Available from: http://dx.doi.org/10.1890/08-2014.1

66. Gamelon M, Gimenez O, Baubet E, Coulson T, Tuljapurkar S, Gaillard JM. Influence of
life-history tactics on transient dynamics: a comparative analysis across mammalian
populations. Am Nat [Internet]. 2014 Nov;184(5):673–83. Available from:
http://dx.doi.org/10.1086/677929

67. Capdevila P, Stott I, Cant J, Beger M, Rowlands G, Grace M, et al. Life history mediates
the trade-offs among different components of demographic resilience. Ecol Lett.
:2021.06.30.450480.

68. Capdevila P, Stott I, Beger M, Salguero-Gomez R. Towards a Comparative Framework of
Demographic Resilience. Trends Ecol Evol [Internet]. 2020 Sep;35(9):776–86. Available from:
http://dx.doi.org/10.1016/j.tree.2020.05.001

69. Greenspoon L, Krieger E, Sender R, Rosenberg Y, Bar-On YM, Moran U, et al. The
global biomass of wild mammals. Proc Natl Acad Sci U S A [Internet]. 2023 Mar
7;120(10):e2204892120. Available from: http://dx.doi.org/10.1073/pnas.2204892120

70. Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, et al. AVONET:
morphological, ecological and geographical data for all birds. Ecol Lett [Internet]. 2022
Mar;25(3):581–97. Available from: http://dx.doi.org/10.1111/ele.13898

71.   Froese R, Pauly D. FishBase: A Global Information System on Fishes. World Wide Web
electronic publication: www.fishbase.org [Internet]. 2002 [cited 2024 Jan 13];000–000.
Available from: https://oceanrep.geomar.de/id/eprint/2964

72.   Boettiger C, Lang DT, Wainwright PC. rfishbase: exploring, manipulating and visualizing
FishBase data from R. J Fish Biol [Internet]. 2012 Nov;81(6):2030–9. Available from:
http://dx.doi.org/10.1111/j.1095-8649.2012.03464.x

73.   Myhrvold NP, Baldridge E, Chan B, Sivam D, Freeman DL, Ernest SKM. An amniote life-
history database to perform comparative analyses with birds, mammals, and reptiles. Ecology
[Internet].   2015   Nov;96(11):3109–3000.   Available   from:
https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/15-0846R.1

74.   Bernard B, Santos G, Deere J, Capdevila P, Kusch E, Gascoigne S, et al. MOSAIC: A
Unified Trait Database to Complement Structured Population Models. bioRxiv [Internet].
2022;2022. Available from: https://mosaicdatabase.web.ox.ac.uk/

75.   Dewey T, Shefferly N, Havens A. Animal Diversity Web. University of [Internet]. 2010;
Available   from:
https://animaldiversity.ummz.umich.edu/site/accounts/information/Felis_silvestris.html

76.   Baillie J, Hilton-Taylor C, Stuart SN, IUCN Species Survival Commission. 2004 IUCN Red
List of Threatened Species: A Global Species Assessment [Internet]. IUCN; 2004. 191 p.
Available from: https://play.google.com/store/books/details?id=Djr8v_-mFzYC

77.   McTavish EJ, Hinchliff CE, Allman JF, Brown JW, Cranston KA, Holder MT, et al.
Phylesystem: a git-based data store for community-curated phylogenetic estimates.
Bioinformatics   [Internet].   2015   Sep   1;31(17):2794–800.   Available   from:
http://dx.doi.org/10.1093/bioinformatics/btv276

78.   Michonneau F, Brown JW, Winter DJ. rotl: an R package to interact with the Open Tree of
Life data. Methods Ecol Evol [Internet]. 2016 Dec;7(12):1476–81. Available from:
https://besjournals.onlinelibrary.wiley.com/doi/10.1111/2041-210X.12593

79.   Plese B, Kenny NJ, Rossi ME, Cárdenas P, Schuster A, Taboada S, et al. Mitochondrial
evolution in the Demospongiae (Porifera): Phylogeny, divergence time, and genome biology.
Mol   Phylogenet   Evol   [Internet].   2021   Feb;155:107011.   Available   from:
http://dx.doi.org/10.1016/j.ympev.2020.107011

80.   Revell LJ. R package phytools. 2013.

81.   Salguero-Gomez R. The fast-slow continuum is not the end-game of life history evolution,
human or otherwise. Evol Hum Behav. 2021;

82.   Revell LJ. Size-correction and principal components for interspecific comparative studies.
Evolution   [Internet].   2009   Dec;63(12):3258–68.   Available   from:
http://dx.doi.org/10.1111/j.1558-5646.2009.00804.x

83.   Blomberg SP, Garland T, Ives AR. Testing for phylogenetic signal in comparative data:
Behavioral traits are more labile. Evolution [Internet]. 2003 Apr;57(4):717–45. Available from:
http://dx.doi.org/10.1111/j.0014-3820.2003.tb00285.x

84.   Freckleton RP. Phylogenetic tests of ecological and evolutionary hypotheses: checking
for phylogenetic independence. Funct Ecol [Internet]. 2000 Feb;14(1):129–34. Available from:
http://doi.wiley.com/10.1046/j.1365-2435.2000.00400.x

85.   Orme D, Freckleton R, Thomson G, Petzoldt T, Fritz S, Isaac N, et al. The caper
package: comparative analysis of phylogenetics and evolution in R [Internet]. 2013. Available

from: http://cran.r-project.org/web/packages/caper/

86.   van Buuren S, Groothuis-Oudshoorn K. MICE: multivariate imputation by chained
equations   [Internet].   2013.   Available   from: http://cran.r-
project.org/web/packages/mice/index.html

87.   James TD, Salguero-Gómez R, Jones OR, Childs DZ, Beckerman AP. Bridging gaps in
demographic analysis with phylogenetic imputation. Conserv Biol [Internet]. 2021
Aug;35(4):1210–21. Available from: http://dx.doi.org/10.1111/cobi.13658

88.   Legendre P, Legendre L. Numerical Ecology. 3rd ed. London: Elsevier; 2012. 1006 p.

89.   Montero AP, Williams DM, Martin JGA, Blumstein DT. More social female yellow-bellied
marmots, Marmota flaviventer, have enhanced summer survival. Anim Behav [Internet]. 2020
Feb   1;160:113–9.   Available   from:
https://www.sciencedirect.com/science/article/pii/S0003347219304154

90.   Höner OP, Wachter B, East ML, Runyoro VA, Hofer H. The effect of prey abundance and
foraging tactics on the population dynamics of a social, territorial carnivore, the spotted hyena.
Oikos [Internet]. 2005; Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.0030-
1299.2005.13533.x?casa_token=5gDMug6OJuYAAAAA:ji20KaEeVK3_l6tbw2jnMVHKZBAJD
31Qw6-zd9rqMojV0GAuGt1YALOjCtiigwuCMrND6f0kM-ndYJNVvQ

91.   Kreider JJ, Kramer BH, Komdeur J, Pen I. The evolution of ageing in cooperative
breeders.   Evol   Lett   [Internet].   2022   Dec;6(6):450–9.   Available   from:
http://dx.doi.org/10.1002/evl3.307

92.   da Silva J. Life History and the Transitions to Eusociality in the Hymenoptera. Frontiers in
Ecology   and   Evolution   [Internet].   2021;9.   Available   from:
https://www.frontiersin.org/articles/10.3389/fevo.2021.727124

93.   Wilson EO, Hölldobler B. Eusociality: origin and consequences. Proc Natl Acad Sci U S A
[Internet].   2005   Sep   20;102(38):13367–71.   Available   from:
http://dx.doi.org/10.1073/pnas.0505858102

94.   Kappeler PM, van Schaik CP. Evolution of primate social systems. Int J Primatol
[Internet].   2002;23(4):707–40.   Available   from:
https://idp.springer.com/authorize/casa?redirect_uri=https://link.springer.com/article/10.1023/A
:1015520830318&casa_token=D4H0iLbXhxgAAAAA:Xni5gWzoFf3r1uZ6lKPj1B-
UraLsoEdAbt1xqfx_-zx9_UHjlnDYNvRu8Ell4dWPf5HlZgXXWPkcyCqkQQ

95.   Hatchwell BJ. The evolution of cooperative breeding in birds: kinship, dispersal and life
history. Philos Trans R Soc Lond B Biol Sci [Internet]. 2009 Nov 12;364(1533):3217–27.
Available from: http://dx.doi.org/10.1098/rstb.2009.0109

96.   Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, et al. The
benefits of social capital: close social bonds among female baboons enhance offspring
survival. Proc Biol Sci [Internet]. 2009 Sep 7;276(1670):3099–104. Available from:
http://dx.doi.org/10.1098/rspb.2009.0681

97.   Corcobado G, Rodríguez-Gironés MA, Moya-Laraño J, Avilés L. Sociality level correlates
with dispersal ability in spiders. Funct Ecol [Internet]. 2012 Aug;26(4):794–803. Available from:
https://besjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2435.2012.01996.x

98.   Rubenstein DR, Abbot P. Comparative Social Evolution [Internet]. Cambridge University
Press;   2017.   479   p.   Available   from:
https://play.google.com/store/books/details?id=VnSJDgAAQBAJ

99. Anderson JA, Johnston RA, Lea AJ, Campos FA, Voyles TN, Akinyi MY, et al. High social status males experience accelerated epigenetic aging in wild baboons. Elife [Internet]. 2021 Apr 6;10. Available from: http://dx.doi.org/10.7554/eLife.66128

100. Berger V, Lemaître JF, Allainé D, Gaillard JM, Cohas A. Early and Adult Social Environments Shape Sex-Specific Actuarial Senescence Patterns in a Cooperative Breeder. Am Nat [Internet]. 2018 Oct;192(4):525–36. Available from: http://dx.doi.org/10.1086/699513

101. Reznick DN, Bryant MJ, Roff D, Ghalambor CK, Ghalambor DE. Effect of extrinsic mortality on the evolution of senescence in guppies. Nature [Internet]. 2004 Oct 28;431(7012):1095–9. Available from: http://dx.doi.org/10.1038/nature02936

102. Verhulst S, Geerdink M, Salomons HM, Boonekamp JJ. Social life histories: jackdaw dominance increases with age, terminally declines and shortens lifespan. Proc Biol Sci [Internet]. 2014 Sep 22;281(1791):20141045. Available from: http://dx.doi.org/10.1098/rspb.2014.1045

103. Shefferson R, Jones OR, Salguero-Gómez R. The Evolution of Senescence in the Tree of Life. 1st ed. Press, Cambridge University, editor. London: Cambridge University Press; 2017.

104. McGraw JB, Caswell H. Estimation of Individual Fitness from Life-History Data. Am Nat [Internet]. 1996 Jan 1;147(1):47–64. Available from: https://doi.org/10.1086/285839

105. Iles D, Koons D, Adler P, Salguero-Gomez R. Transient Dynamics, Life History, and Biological Invasions. In: Student Research Symposium [Internet]. 2015 [cited 2024 Jan 15]. Available from: https://digitalcommons.usu.edu/researchweek/ResearchWeek2015/All2015/279/

106. Hodgson D, McDonald JL, Hosken DJ. What do you mean, "resilient"? Trends Ecol Evol [Internet]. 2015 Sep;30(9):503–6. Available from: http://dx.doi.org/10.1016/j.tree.2015.06.010

107. Pfister CA. Patterns of variance in stage-structured populations: evolutionary predictions and ecological implications. Proc Natl Acad Sci U S A. 1998 Jan;95(1):213–8.

108. Tuljapurkar S. An uncertain life: demography in random environments. Theor Popul Biol. 1989 Jun;35(3):227–94.

109. Le Coeur C, Yoccoz NG, Salguero-Gómez R, Vindenes Y. Life history adaptations to fluctuating environments: Combined effects of demographic buffering and lability. Ecol Lett [Internet]. 2022 Oct;25(10):2107–19. Available from: http://dx.doi.org/10.1111/ele.14071

110. Gascoigne SJL, Kajin M, Tuljapurkar S, Santos GS, Compagnoni A, Steiner UK, et al. Structured demographic buffering: A framework to explore the environment drivers and demographic mechanisms underlying demographic buffering [Internet]. bioRxiv. 2023 [cited 2024 Jan 15]. p. 2023.07.20.549848. Available from: https://www.biorxiv.org/content/biorxiv/early/2023/07/21/2023.07.20.549848

111. Santos GS, Gascoigne SJL, Dias ATC, Kajin M, Salguero-Gómez R. A unified framework to identify demographic buffering in natural populations [Internet]. bioRxiv. 2023 [cited 2024 Jan 15]. p. 2023.07.03.547528. Available from: https://www.biorxiv.org/content/10.1101/2023.07.03.547528

112. Finch CE. Longevity, senescence and the genome. Chicago, USA: University of Chicago Press; 1990.

113. Schenk HJ, Espino S, Goedhart CM, Nordenstahl M, Cabrera HIM, Jones CS. Hydraulic integration and shrub growth form linked across continental aridity gradients. Proc Natl Acad Sci U S A [Internet]. 2008 Aug;105(32):11248–53. Available from:

http://dx.doi.org/10.1073/pnas.0804294105

114.   Cagnacci F. GPS-telemetry Data: Challenges and Opportunities for Behavioural Ecology
Studies [Internet]. Royal society; 2010. 148 p. Available from:
https://play.google.com/store/books/details?id=sZqYzwEACAAJ

115.   Tong C, Najm GM, Pinter-Wollman N, Pruitt JN, Linksvayer TA. Comparative Genomics
Identifies Putative Signatures of Sociality in Spiders. Genome Biol Evol [Internet]. 2020 Mar
1;12(3):122–33. Available from: http://dx.doi.org/10.1093/gbe/evaa007

116.   Coulson T, MacNulty DR, Stahler DR, Vonholdt B, Wayne RK, Smith DW. Modeling
Effects of Environmental Change on Wolf Population Dynamics, Trait Evolution, and Life
History. Science [Internet]. 2011 Dec;334(6060):1275–8. Available from:
http://dx.doi.org/10.1126/science.1209441

 # Tables

 Table 1

Type	Demographic trait	R^2	F	d.f.	P	Pagel's λ	95% CI
A. Life history traits	Generation time (T)	3.618	1.380	5, 147	**0.024**	0.665	0.368 - 0.851
	Age at maturity (L_a)	0.661	0.244	5, 147	0.913	0.649	0.288 - 0.858
	Mean life expectancy (η_e)	1.079	1.401	5, 147	**0.008**	0.739	0.485 - 0.879
	Variance in life expectancy ($\Delta\eta_e$)	2.078	1.678	5, 147	**0.010**	0.668	0.568- 0.710
	Actuarial senescence (s_{lx})	3.420	1.301	5, 147	0.272	0.158	0.009 - 0.483
	Probability of reproduction (p_R)	0.584	1.216	5, 147	**0.009**	0.519	0.205 - 0.783
	Reproductive senescence (s_{mx})	1.374	0.512	5, 147	0.727	0.023	0.000 - 0.425
	Demetrius' entropy (S)	2.150	1.208	5, 147	0.062	0.019	0.000 - 0.297
B. Vital rates	Survival (σ)	1.372	0.511	5, 147	0.728	0.616	0.314 - 0.823
	Growth (γ)	3.477	1.324	5, 147	0.264	0.000	0.000 - 0.493
	Shrinkage (ρ)	2.970	1.125	5, 147	**0.045**	0.000	0.000 - 0.195
	Reproduction (φ)	6.637	2.612	5, 147	**0.038**	0.000	0.000 - 0.082
C. Elasticities	Survival elasticity ($E\sigma$)	6.196	2.428	5, 120	**0.040**	0.229	0.038 - 0.576
	Growth elasticity ($E\gamma$)	14.171	6.068	5, 120	**<0.001**	0.000	0.000 - 0.101
	Shrinkage elasticity ($E\rho$)	3.090	1.172	5, 120	0.326	0.405	0.216 - 0.607
	Reproduction elasticity ($E\varphi$)	1.984	1.365	5, 120	**0.033**	0.640	0.281 - 0.852
D. Asymptotics	Population growth rate (λ)	2.789	1.054	5, 120	0.381	0.001	0.000 - 0.160
	Deviance growth rate equilibrium ($|1-\lambda|$)	1.357	0.505	5, 131	0.732	0.107	0.000 - 0.443

	Damping ratio (ζ)	1.049	0.390	5, 124	0.816	0.178	0.000 - 0.587
	Period of oscillation (P_i)	9.734	3.963	5, 86	**0.004**	0.011	0.000 - 0.053
	Reactivity ($\overline{\rho}_1$)	1.861	1.319	5, 85	**0.036**	0.475	0.213 - 0.747
E. Transients	First step attenuation ($\underline{\rho}_1$)	1.507	1.874	5, 107	0.094	0.526	0.207 - 0.788

Figure and table captions

Table 1. The degree of sociality predicts key demographic traits across 152 animal species. Phylogenetic generalised least square (pgls) models of sociality and different demographic traits. The demographic traits are classified into five general types: **A**. Life history traits; **B**. Vital rates; **C**. Elasticities of population growth rate (λ) to vital rates; **D**. Metrics of asymptotic (*i.e.,* long-term) performance; and **E**. Metrics transient (*i.e.,* short-term) performance. **Bold**: $P < 0.05$. Pagel's λ quantifies the phylogenetic signal in the model, and it ranges from 0 (*i.e.,*no phylogenetic signal) to 1 (*i.e.,* pattern completely explained by phylogenetic relationships).

Figure 1. Geographic and phylogenetic representation of the 152 animal species in this study to examine the correlates of sociality and demographic traits. Species sociality is represented along a discrete continuum, from solitary (blue), gregarious (green), communal (yellow), colonial (orange), to social (red). **A**. World map depicting the geographic location of the examined studies from the COMADRE Animal Matrix Database, when GPS coordinates were available (n = 98 species). **B**. Phylogenetic representation of the 152 study species. Numbered arcs represent the taxonomic classes (from left to right): 1: Scyphozoa (n = 1 species); 2: Anthozoa (n = 4); 3: Malacostraca (n = 1); 4: Insecta (n = 4); 5: Gastropoda (n = 4); 6: Bivalvia (n = 5); 7: Thaliacea (n = 1); 8: Elasmobranchii (n = 1); 9: Actinopterygii (n = 28); 10: Amphibia (n = 1); 11: Reptilia (n = 18); 12: Aves (n = 26); and 13: Mammalia (n = 58). Silhouette source: http://phylopic.org.

Figure 2. The continuum of sociality predicts key life history traits in animals. Boxplots of

the residuals of life history traits after correction for adult body mass and phylogenetic

relationships as a function of the discrete continuum of sociality for 152 animal species.

The letters above each group represent post-hoc Tukey test significance levels from the

phylogenetic ANOVA. Groups with different letters are significantly different at $P < 0.05$.

Figure 3. The continuum of sociality is correlated with key vital rates and their elasticities

to population growth rate. Boxplots of the residuals of vital rates (left: survival σ, growth

γ, shrinkage ρ, and reproduction φ), and their elasticities to population growth rate λ

(right) after correcting for adult body mass and phylogenetic relationships, as a function

of the discrete continuum of sociality for 152 animal species. The letters above each

group represent post-hoc Tukey test significance levels from the phylogenetic ANOVA.

Groups with different letters are significantly different at $P < 0.05$.

Figure 4. The continuum of sociality is correlated with key attributes of transient (short-

term) dynamics that predict the inherent ability of natural populations to respond to

disturbances. Boxplots of the residuals of four transient dynamics metrics after correction

for adult body mass and phylogenetic relationships, as a function of the discrete

continuum of sociality for 152 animal species. Damping ratio (ζ) quantifies the speed of

recovery post disturbance; Period of oscillation (P_i) quantifies the number of cycles a

population goes through before achieving asymptotic dynamics; Reactivity ($\overline{\rho}_1$) quantifies

how much a population can grow after one time step following a disturbance, while first

step attenuation ($\underline{\rho}_1$) quantifies the loss of resistance after one time step. The letters

above each group represent post-hoc Tukey test significance levels from the

phylogenetic ANOVA. Groups with different letters are significantly different at $P < 0.05$.

Figure 5. The life history strategy space of animals predicts the placing of sociality along

the fast-slow continuum, but not the reproductive strategies continuum. Adult body-mass,

phylogenetically corrected principal component analyses (pPCA) of the eight life history

traits shown in Table 1A and Figure 2. These traits are colour-coded by investments on

survival and reproduction (black - generation time, T), survival only (grey - age at

maturity, L_α; mean life expectancy, η_e; variance in life expectancy, $\Delta\eta_e$; actuarial

senescence, s_{lx}), or reproduction only (red - probability of achieving reproduction before

dying, p_R; reproductive senescence, s_{mx}; Demetrius' entropy, S, which quantifies the

degree of semel/itero-parity). Each point in the PCA space corresponds to one animal

species, and some are represented by a silhouette (source: http://phylopic.org; refer to

Fig. 1B). The loadings of the life history traits clearly identifies: PC1 (explaining 27.76%

of the variance) as the fast-slow continuum, with species living long, having high

variance in life expectancy, and postponing their first reproductive event to the right; and

PC2 (explaining 26.94% of the variance) as the reproductive strategies continuum, with

species that senesce less for survival (s_{lx}) and reproduction (s_{mx}), have a higher

probability of achieving reproduction before death (p_R), and are highly iteroparous (S) at

the top. The side boxplots correspond to the groups along PC1 (top) and PC2 (right) as a

function of the discrete continuum of sociality.

Social
Colonial
Communal
Gregarious
Solitary

Reproductive strategies – PC2 (26.94%)

Fast–slow continuum – PC1 (27.76%)

S
p_R
S_{mx}
s_{lx}
η_e
$\Delta\eta_e$
T
L_α