More social species live longer, have higher generation times, and longer reproductive windows

Roberto Salguero-Gómez[1,*]

1 Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RB, United Kingdom, ORCID ID 0000-0002-6085-4433

* Author for correspondence: rob.salguero@biology.ox.ac.uk

Total word count: 6,181

Total figures: 5

Total tables: 1

Total references: 129

SOMs: 3 tables, 5 figures

Abstract

The role of sociality in the demography of animals has become an intense focus of

research in recent decades. However, efforts to understand the sociality-demography

nexus have focused on single species or isolated taxonomic groups. Consequently, we

lack generality regarding how sociality associates with demographic traits within the

Animal Kingdom. Here, I propose a continuum of sociality, from solitary to tightly social,

and test whether this continuum correlates with the key demographic properties of 152

species, from jellyfish to humans. After correction for body mass and phylogenetic

relationships, I show that the sociality continuum is associated with key life history traits:

more social species live longer, postpone maturity, have greater generation time, and

greater probability of achieving reproduction than solitary, gregarious, communal, or

colonial species. Contrary to the social buffering hypothesis, sociality does not result in

more buffered populations. While more social species have a lower ability to benefit from

disturbances, they display greater resistance than more solitary species. Finally, I also

show that sociality does not shape reproductive or actuarial senescence rates. This

cross-taxonomic examination of sociality across the demography of 13 taxonomic

classes highlights keyways in which individual interactions shape most aspects of animal

demography.

Keywords: eusocial, buffering, generation time, gregarious, solitary, colonial, communal,

resilience, senescence.

Introduction

Quantifying the links between sociality and demography is crucial to understand and forecast the dynamics of animal wildlife populations [1,2]. Indeed, social structures can influence various aspects of an organism's life, including its survival [3,4] and reproduction [5,6]. As such, the links between sociality and the demography of animal species has gained increasing attention in recent years [7–9], as we strive to unravel the mechanisms underlying population fluctuations and viability [10,11].

Social interactions within a group can significantly influence the key rates that shape organismal fitness: the survival and reproduction of individuals in the group. For instance, cooperative breeding, where individuals assist in the care and rearing of offspring, occurs in many animal species, from meerkats (*Suricata suricatta*; [12] to fairy-wrens (*Malurus cyaneus;* [13]. This behaviour may enhance reproductive success by optimising resource allocation [14] and providing protection against predators [15]. Conversely, in some species, competition for mates within social groups can result in skewed reproductive success [16,17]. Survival patterns within a population can also be intricately linked to sociality. Indeed, social animals often experience reduced predation risk through group vigilance [18] and cooperative defence mechanisms [19]. However, social hierarchies can also increase competition [20], and lower survival [21,22]. As such, sociality has the potential to drastically affect the demography of animal species. Understanding how these contrasting factors interact is essential for predicting and managing population dynamics in socially structured species [23].

Our understanding of how sociality shapes the demography of animals is lacking generality. Specifically, we lack a comprehensive overview of how sociality associates with various demographic attributes of animal populations across a wide range of

taxonomic classes - but see [24] for a cross-taxonomic examination of reproductive monopolisation across *ca.* 300 species of wasps, ants, birds, and mammals. Most social comparative studies have focused on birds and mammals [25–28]. In these studies, sociality has been linked to enhanced reproductive success [29], and only weakly with reproductive senescence [27]. Within primates, the impacts of sociality are modulated via complex group hierarchies, where dominance influences access to resources, mating opportunities, and reproductive success [30–32]. Among the carnivora, social species like lions (*Panthera leo*) and spotted hyenas (*Crocuta crocuta*) display complex group structures that influence demographic patterns. In them, cooperative hunting and communal care of young enhance the survival and overall fitness of individuals within the social group [33]. In eusocial insects (*e.g.*, ants, bees, and termites), the within-colony division of labour affects demographic traits via resource allocation, colony defence, and reproduction [34,35]. In certain fish species, such as the clownfish (*Amphiprion* spp.) and cleaner wrasses (*Labroides* spp.), sociality is linked to improved foraging efficiency, reduced predation risk, and increased reproductive success [36].

Sociality is multifaceted. As such, different authors define the degree of sociality depending on the taxonomic group under examination. I argue that the lack of cross-taxonomic categorisation of sociality, together with the historical scarcity of demographic data across the Tree of Life explains why a broader examination of sociality across the demographies of the Animal Kingdom does not exist. However, taking inspiration from the Plant Kingdom, one could think about the individuals in a group of animals as an analogous organisation to the xylem vessels that connect the roots and shoots in an individual vascular plant. The degree of integration-modularity of these individuals (vessels) within the animal group (individual plant) provides a framework to categorise sociality across the Animal Kingdom. Indeed, the degree of integration-modularity across

138 plant species strongly correlates with their longevity and rate of actuarial

senescence [37].

Here, I propose a continuum of animal sociality to examine whether sociality, the way

individual organisms organise themselves within a population and interact, shapes their

demography. My sociality continuum contains five categories: (1) solitary: individuals

spend their time alone, except to breed (*e.g.*, tigers (*Panthera tigris*), cheetahs (*Acinonyx*

jubatus), some wasps); (2) gregarious: individuals spend time in groups but social

interactions are loose (*e.g.*, wildebeests, zebras, flock-forming birds); (3) communal:

individuals live together in close proximity and often share a common nesting or dwelling

area, but do not engage in cooperative breeding (*e.g.*, purple martin (*Progne subis*)); (4)

colonial: individuals live in close proximity and always share a common nesting or living

area (*e.g.*, nesting birds, some wasps, coral polyps); and (5) social: individuals live in

close proximity and form stable, organised groups, engaging in social behaviours such

as cooperative breeding and hierarchical structures (*e.g.*, African elephant (*Loxodonta*

africana), most primates, dolphins, meerkat, honeybee (*Apis mellifera*)).

I apply this sociality continuum to 152 species across 13 taxonomic classes, from

cnidarians to mammals (Fig. 1), to carry out phylogenetic comparative analyses to

examine the demographic correlates of sociality across their life history traits [38], vital

rates and impacts on population growth rate [39], and metrics of long-term and short-

term performance [40]. My multi-level examination of the demographic traits of these

species allows me to test expectations regarding the expected impacts of sociality on

longevity and senescence [41] and the 'social buffer' hypothesis [42], whereby more

social organisms should be able to better mitigate environmental stochasticity while also

displaying lower temporal oscillations in population growth rate.

Methods

To test the aforementioned hypotheses, I ran a series of comparative phylogenetic analyses based on high resolution demographic, body mass, sociality, and phylogenetic data of animal species across multiple taxonomic classes.

Demographic data

To obtain demographic data across a wide range of animal species, I used the COMADRE Animal Matrix Database [43]. COMADRE is an open-access database that contains high-resolution demographic information of animal species, mostly from peer-reviewed publications. In its current version, COMADRE (v. 4.32.1) contains 3,488 matrix population models (MPMs, hereafter [44]) from 429 animal studies across 415 studies. An MPM is a numerical array that describes the demography of a species at a given location, time period, and environmental conditions, in discrete time and along discrete life cycle stages [44]. Specifically, in an MPM, the life cycle of a given population is discretised into stages (*e.g.*, age, discrete size ranges, developmental stages), such that the rates that are then stored in the MPM describe individual-level, stage-specific survival (σ), growth/development (γ), shrinkage/de-development (ρ), and reproduction (φ) from time t to t+1.

To ensure comparability across the thousands of MPMs available in COMADRE, I imposed the following series of selection criteria using the *Rcompadre* R package [45]: (1) Unmanipulated MPMs from wild populations, so that the derived demographic traits reflect the dynamics of the population under natural conditions, rather than under experimental treatments or in captivity; (2) MPMs with dimension equal to or greater than 4 × 4 (*i.e.*, 4 or more stages in the life cycle), so as to allow for enough resolution in the life cycle to successfully discern age-specific patterns of survivorship and reproduction

[46,47] and avoid mortality and fertility plateaus [48]; (3) MPMs where the rates of

survival and reproduction had been separated into sub-matrices (such that the overall

MPM $A = U + F$, where U and F represent the sub-matrices of survival and reproduction,

respectively), to facilitate the calculation of life history traits and vital rates and their

elasticities to population growth rate; (4) MPMs whose stage-specific survival values ≤ 1,

to prevent erroneous calculations of inflated longevity and actuarial senescence [47]; and

(5) MPMs corresponding to extant species (*e.g.*, a study successfully passed through

selection criteria 1-4, above, but in fact corresponds to an extinct porcupine, *Hystrix*

refossa [49]). These criteria resulted in 154 animal species, for which 23 species had at

least two available studies in COMADRE.

Next, for those 23 species with multiple studies available, I manually chose the single

study per species that provides higher representativity of the demography of the species

of interest by retaining the longer-term study, the MPM with greater dimensionality,

and/or the study where the rates of survival and/or reproduction in adults were not

assumed constant. For each of the resulting 154 animal species, I retained the 'grand

mean' MPM, which describes the dynamics of the population across the length of the

study by calculating the element-by-element arithmetic mean MPM, thus resulting in one

MPM per species. In addition, I supplemented my dataset with a recent study on meerkat

demography [50], not yet incorporated in COMADRE. For the 10 years of seasonal

MPMs available in this study, I retained the MPMs corresponding to 1998-2002,

corresponding to normal environmental conditions (M. Paniw, pers. comm.), and then

again estimated the element-by-element arithmetic grand mean MPM. This addition

increased the count to 155 animal species. The vast majority of MPMs were

parameterised with data collected once per year (n = 130, 84%; Fig. S1). However, to

make sure that all demographic traits derived from the MPMs (below) were on the same time units, MPMs not on an annual periodicity were rescaled accordingly by elevating each MPM element $a_{i,j}$ in **A** to the power of $1/P$, where P is the projection interval of the MPM, which is also archived in COMADRE [43]. The original sources of the individual demographic studies used in this paper are shown in Table S1.

For each species' grand MPM, I derived a series of demographic traits that inform the life history traits, vital rates and their relative impacts on population growth rate, as well as metrics of asymptotic (*i.e.*, long-term) and transient (*i.e.*, short-term) performance. To do so, I used the R packages *Rage* [45], *popdemo* [51] and *popbio* [52].

Life history traits are key metrics that define specific moments and periods of time along the life cycle of a species [38]. Previous examinations have demonstrated that the chosen life history traits describe ~60-80% of the variation in life history strategies of animals [53–57]. The 11 chosen life history traits are:

(1) Generation time (T) the average age of reproductive individuals in a population [55];

(2) Net reproductive output (R_0): the total number of offspring produced by the average individual in the population during its lifespan [44];

(3) Mean life expectancy (η_e): mean lifespan of a cohort [44];

(4) Variance in life expectancy ($\Delta\eta_e$): variance of lifespan in a cohort [44];

(5) Maximum longevity (L_{max}): time elapsed until 99% of a cohort has died [58];

(6) Age at maturity (L_α): age at which the first individual in a cohort reproduces [59];

(7) Reproductive window ($L_{\alpha-\omega}$): the average duration of reproduction of individuals in the population [60];

(8) Maturity probability (p_R): the probability of achieving reproduction before dying [44];

(9) Actuarial senescence (s_{lx}): the shape (*s hereafter*) of age-specific survivorship l_x

[61,62]; The value of this shape ranges between -0.5 and +0.5: $s_{lx} = 0$ indicates

negligible actuarial senescence (*i.e.,* constant survival with age), $s_{lx} < 0$ indicates

negative actuarial senescence (generally increasing survival with age), and $s_{lx} > 0$

indicates (positive) actuarial senescence (generally decreasing survival with age);

(10) Reproductive senescence (s_{mx}): the shape of age-specific reproduction m_x [63]

describes the symmetry of reproduction over age by comparing the area under a

cumulative reproduction curve over age with the area under constant reproduction,

and it ranges between -0.5 and +0.5: $s_{mx} = 0$ indicates negligible reproductive

senescence (*i.e.,* constant reproduction with age), $s_{mx} > 0$ indicates (positive)

reproductive senescence (decreasing reproduction with age), and $s_{mx} < 0$ indicates

negative reproductive senescence (increasing reproduction with age);

(11) Degree of parity (*S*): quantifies the degree of semelparity (individuals in the

population reproduce once; S = 0) *vs.* iteroparity (reproduce multiple times; S>>0) [64]

(Fig. S2).

To further characterise the potential associations between sociality and the demography

of different animal species, I examined vital rates and their relative impacts on population

growth rate. Using the Rage R package [45], I estimated an average value of survival

(σ), growth (γ), shrinkage (ρ), and reproduction (φ), weighted by the stationary

distribution of individuals in the population [60]. Next, I calculated the elasticities of

population growth rate (λ) to these vital rates. These elasticities quantify the effect that a

relative, infinitesimally small change in a vital rate would have on the long-term,

asymptotics of the population [39,65], quantified as the dominant eigenvector of the MPM

A, λ [44] (not to be confused with Pagel's λ *[66]*, an estimate of phylogenetic signal -

below). Next, I also quantified the population growth rate (λ), and the deviance of

population growth rate from $\lambda=1$ as $|1-\lambda|$, at which point the population is neither

increasing ($\lambda>1$) nor declining ($\lambda>1$). Deviance from $\lambda=1$ was of particular interest here,

as the social buffer hypothesis [42] predicts the canalisation of λ towards demographic

stability ($\lambda \approx 1$) with increased sociality.

Finally, I also calculated metrics of transient dynamics. Transient dynamics in an

structured population model, such as an MPM, quantify the short-term responses of a

population to potential disturbances across the structure of said population [40,67]. The

full range of potential responses to disturbances creates a so-called transient envelope

[40], from which one can calculate the rate of recovery to the stationary equilibrium

(damping ratio, ζ), the number of oscillations that the population undergoes before

achieving said equilibrium (period of oscillation, P_i), the maximum increase in size that

the population can undertake following a disturbance (reactivity, $\overline{\rho}_1$), and the maximum

decrease in size that the population can display after a disturbance (first step

attenuation, $\underline{\rho}_1$). I note that many other transient metrics can be used to define the

transient envelope of an MPM [40], but the four I chose here tend to be uncorrelated

[68,69], and together they inform the inherent ability of a population to respond to

disturbances, that is, their demographic resilience [70].

The vital rate elasticities, and metrics of asymptotic and transient performance were only

calculated for MPMs that are reducible, primitive, and ergodic, thus guaranteeing the

existence of a single dominant eigenvalue, λ_1 [40,44].

Adult body mass data

Most life history traits scale with adult body mass [53–56]. As such, to explore the

relationships between sociality and life history traits beyond the confounding effect of

body mass, I first obtained adult body mass data for each species. For mammals, I used

information archived in AnimalTraits [71]; for birds, AVONET [72]; for fish, FishBase [73]

via rFishBase [74]; for mammals, birds, and reptiles and amphibians, AMNIOTE [75]; and

for remaining groups, MOSAIC [76], and data from [54]. For 18 remaining species for

which adult body mass data were not available from these online databases (mostly

insects, bivalves, and corals), I obtained body mass information from peer-review

publications using the keywords "body mass" OR "weight" and the name of the species in

ISI WoS. Data were transformed to grams across all sources. The final variable was log-

transformed to fulfil assumptions of the statistical analyses employed below.

<u>Sociality data</u>

I classified sociality following my proposed sociality continuum. In the

development of this classification, my goal was to explicitly acknowledge how sociality is

not a binary category, and that it indeed has a multi-faceted nature, as animals can vary

in their social interactions according to temporal and spatial dimensions [77], as well as a

function of group size [78], group hierarchy [79], and type of interactions [80]. My

continuum of sociality takes five ordered levels, from less to more social, as follows: (1)

Solitary: individuals spend most of their life cycles alone, except to breed; (2) Gregarious:

individuals spend some time in groups, but their social interactions are loose, and these

interactions can frequently disaggregate; (3) Communal: individuals live in close

proximity, often sharing a common nesting or dwelling area, but do not engage in

cooperative breeding; (4) Colonial: individuals live in close proximity and always share a

common nesting or living area; and (5) Social: individuals live in close proximity and form

stable, organised groups, engaging in social behaviours such as cooperative breeding

and hierarchical structures. From solitary to social, the degree of spatial and temporal

interactions increases, as does the quality of the interaction.

No centralised, open-access database exists to my knowledge offering a

comprehensive categorisation of the sociality of species across kingdom Animalia. As

such, I classified species according to my continuum of sociality drawing from multiple

sources, which allowed me to carefully cross-validate the classification for each species.

I first contrasted information from the following online resources for each species in my

dataset: Animal Diversity Web [81], Smithsonian's National Zoo and Conservation

Biology   Institute   (https://nationalzoo.si.edu/),   FishBase

(https://www.fishbase.se/search.php), IUCN Red List of Threatened Species [82], and

chatGPT (https://chat.openai.com/) - for the latter, suggested references were cross-

checked independently and validated via ISI WoK. Next, I double-checked existing

classifications in my data and complemented missing information from peer-review

publications for each of the 154 animal species. Once the list was completed, I discussed

it with experts in each of the taxonomic classes, and made the appropriate corrections

(Table S1).

Phylogenetic data

The analyses below explicitly consider how the examined species are related

phylogenetically. To that end, I obtained a phylogeny from the Open Tree of Life [83] via

the R package *ROTL* [84]. I first matched the scientific names of my subset of animal

species from COMADRE to the Taxonomic Name Resolution Service via the function

tnrs_match_names. All species matched successfully, except the two species from class

Demospongiae (*Spongia graminea*, and *Xestospongia muta*). As the phylogenetic tree of

Demospongiae is not well resolved [85], I proceeded with the analyses discarding these

two species, and so the final analyses include 152 animal species, instead of 154.

Comparative phylogenetic analyses

To examine the correlates of sociality with the demographic traits examined from

the MPMs in COMADRE, I first ran a phyloANOVA via the *phytools* R package [86]. For

each demographic trait, the response variable was its residual from the pgls regression

[66] over adult body mass, and the explanatory variable was the factor variable sociality

continuum, with its five different levels, ordered as "solitary", "gregarious", "communal",

"colonial", and "social". To correct for type II errors, after the tests were performed, I also

implemented a Bonferroni correction with the function *p.adjust* from the *Stats* R package.

The phylogenetically-corrected, body mass-corrected post-hoc Tukey test that I

implemented next explicitly allowed me to test the hypothesis that the sociality

continuum, which ranks species along increasing degree of social interactions,

associates with different life history traits, vital rates and their elasticities of population

growth rate, as well as metrics of asymptotic and transient dynamics.

The phylogenetic ANOVAs performed here allowed me to examine the correlates of

sociality with each life history trait separately. However, life history traits are often

correlated with life history strategies [38]. Indeed, previous works have demonstrated

that animals show two primary axes of variation: one where individuals either develop

fast or live long, the so-called fast-slow continuum [38,53,56,87], and another one along

which species differ in their reproductive strategies, the so-called reproductive strategies

continuum [10,54,87].

To quantify the life history strategy space of the 152 study animal species, I used a

phylogenetic principal component analysis (pPCA). Briefly, PCA is a family of

multivariate statistical techniques used to examine complex data by reducing their

dimensionality to highlight the main factors that explain observed variation [88]. The

pPCA takes into consideration the non-independence of life history traits due to species

relatedness [89,90]. To avoid co-linearities in the pPCA, I first examined the pairwise

correlations of the 11 life history traits defined above, and chose only one life history trait

when the spearman correlation coefficient of a pairwise life history trait correlation was >

0.70. This step led me to exclude from pPCA analyses the variance in life expectancy

($\Delta\eta_e$) and maximum longevity (L_{max}), which were highly correlated with mean life

expectancy (η_e) (0.83 and 0.90, respectively; Fig. S3). Next, because PCA approaches

require a dataset with no missing values, and my data contained missing values (Fig.

S4), I carried out imputations only on life history traits for which the degree of missing

data was <40%. This step further excluded the rates of actuarial (s_{lx}) and reproductive

senescence (m_{lx}) from the pPCA analyses. These steps identified seven life history traits

for the pPCA: T, R_0, η_e, L_α, $L_{\alpha-\omega}$, p_R, and S.

To implement the pPCA, I first log-transformed the aforementioned seven life history

traits to fulfil assumptions of error normal distribution necessary in linear models. Next, I

calculated the residuals of each life history trait as a function of species' adult body mass

via a pgls with the R package caper [66] and the phylogeny. Next, I rescaled the

residuals to mean (μ) = 0 and variance (σ) = 1. This approach allowed to establish

primary axes of variation of, in this case, life history traits, as well as to estimate the

overall strength of the phylogenetic relationships among species. For the latter, I also

estimated Pagel's λ, a parameter that ranges between 0 (the pattern of association is not

explained by the phylogenetic hypothesis), and 1 (the pattern is entirely explained by

phylogenetic inertia).

I used a well-established protocol [53,54,60,87] to recover the missing information from

the seven life history traits using an iterative PCA approach. Briefly, I employed the

"*regularised*" method in the function *mice* in the R package *mice [91]* with 20 sets of

imputation. To draw strength across the full dataset, the imputation took into account the

vital rate values, elasticities, transient dynamics metrics (described below), and body

mass (above) of each species. Here, an iterative algorithm first assigns the mean of each

life history trait to the missing values of that life history trait. Next, a regular PCA is

performed, and the scores of the species with missing data are re-assigned using the

orthogonal relationships of the PCA. This iteration is repeated until convergence. As a

check, I (1) re-examined the pairwise correlations of the life history traits pre- (Fig. S3)

and post-imputation (Fig. S5), which did not qualitatively change in their covariation

structure and (2) implemented the same protocol using only the 85 species for which no

life history trait data were missing, and found that the results with imputed data are

insensitive to the imputation (not shown). The negligible sensitivity of the results reported

here to the imputation approach is in agreement with previous comparative demographic

studies that also used COMADRE [10,54,92].

Finally, to examine how the sociality continuum governs the life history strategy space of

animals, I used the scores of the pPCA (above) to run further phylogenetically-corrected

post-hoc Tukey tests to examine if the different levels of the sociality continuum are

significantly clustered along the different PC axes.

Results

The phylogenetic signal of my sociality continuum, from solitary species, like the giant

panda (*Ailuropoda melanoleuca*; Table S1), to social species, like the red howler

(*Alouatta seniculus*), across the 152 examined animal species, was high. Indeed the

estimates of Pagel's λ for the sociality continuum across these species is 0.907 (95%

C.I.: 0.865-0.945; $P < 0.001$). This strong signal is also visually apparent upon examining

the phylogenetic tree shown in Fig. 1B. The phylogenetic signal of the associations

between the sociality continuum and the examined demographic traits, in contrast, was

variable (Table 1). The estimates of Pagel's *λ* from the performed pgls models ranged

from virtually absent across some demographic rates (*e.g.*, degree of parity, vital rates of

individual growth, individual shrinkage, and reproduction, population growth rate (also

called *λ [44]*), period of oscillation), to high in some life history traits (age at maturity

(0.759), reproductive window (0.759), and mean life expectancy (0.745)).

Across the five different types of demographic traits examined here (Table 1: life history

traits, vital rates, vital rate elasticity, asymptotics, and transients), the only category

where the sociality continuum did not raise some statistically significant results was the

metrics of asymptotic dynamics: population growth rate and deviance growth rate

equilibrium. Moreover, in these two metrics, the 95% confidence intervals of Pagel's *λ* for

the respective pgls models overlapped with 0, indicating no phylogenetic effect in the

relationship between sociality and long-term population performance. After correcting for

adult body mass and phylogenetic relationships, I found that more social species attain

longer generation times, longer mean life expectancy, greater variance in life

expectancy, higher maximum longevities, and are more likely to reproduce before dying

(Table 1A). Interestingly, the relationships between the sociality continuum and species'

vital rates were opposed to their elasticities: more social species have an increased

ability to decrease in size and to reproduce (Table 1B), but their population growth rates

would increase the most if their vital rates of survival and growth would increase (Table

1C). Finally, for the transient metrics, I found that more social species tend to display

greater periods of oscillation, lower reactivity (*i.e.,* propensity to benefit from

disturbances, *sensu*   [70]), and higher first step attenuation (*i.e.*, resistance to

disturbances, *sensu* [70]).

The phylogenetic ANOVA post-hoc Tukey tests revealed important differences in the

examined demographic traits as a function of the sociality continuum. For instance,

increases in sociality (solitary → gregarious → communal → colonial → social) were

associated with prolonged generation time, mean and variance of life expectancy,

maximum longevity, longer reproductive windows, and higher probability of reproducing

before dying, but not with net reproductive output, age at maturity, actuarial or

reproductive senescence, nor degree of parity (Fig. 2). It is worth highlighting that, when

the pattern was significant, it did not always change monotonically. Examples include

generation time and maximum longevity, where solitary species attained the same range

of values as social species, whereas the next level on sociality, gregarious species, had

significantly lower values than those of more social species (Fig. 2). Social species also

had higher survival as well as lower individual-level shrinkage and reproduction

compared to most other categories (Fig. 3). These patterns remained largely consistent

when evaluating the elasticities of population growth rate to said vital rates. Finally, for

the transient metrics, overall, as the degree of sociality increased, so did the period of

oscillation and first step attenuation, but species' reactivity decreased (Fig. 4). Here,

again, we note that the range of values of solitary species do not differ from social

species for the period of oscillation and reactivity, but do differ between the next

category, gregarious, and the next three categories: communal, colonial, and social.

The life history strategy space defined by the seven examined life history traits (Table

1A) identified two principal components whose associated eigenvalues > 1. Following

Kaiser's criterion [93], I retained these two axes for the next part of the analysis. PC1

explains 33.72% of the variation in life history traits across the 152 animal species, and it

separates species with low (left of Fig. 5) *vs.* high reproductive schedules. Indeed,

species to the right in the pPCA attain higher net reproductive outputs (R_0), have longer

reproductive windows ($L_{α-ω}$), higher probability of reproducing before dying (p_R), and a

higher degree of iteroparity (S). PC2 explains 26.20% of the variation, and separates

species that live long *vs.* short. Along PC2, species at the top postpone reproduction

($L_α$), have greater generation times (T), and live longer ($η_e$) than those at the bottom. The

emergent phylogenetic signal, after adult body mass correction, in this phylogenetic PCA

was relatively low, with an estimated Pagel's $λ$ of 0.420 (0.341-0.689 95% C.I.).

The ranking of species along the sociality continuum raised detectable differences along

PC1 (the reproductive strategies continuum; P = 0.001; Fig. 5). Indeed, social species

like the macaque (*Macaca mulatta*) or the Asian elephant (*Elephas maximus*) or humans

(*Homo sapiens*), and colonial species like the common tern (*Sterna hirundo*) are

associated with longer reproductive life histories, compared to solitary species like the

flathead catfish (*Pylodictis olivaris*) or the common snapping turtle (*Chelydra serpentina*).

I found no association between the sociality continuum and the fast-slow continuum, PC2

(P = 0.107).

Discussion

Sociality, the degree to which animals engage in interactions and form cohesive

structures, can shape their demography (*i.e.* dynamics of population size and structure)

via impacts on their rates of survival [94] and reproduction [16], the building blocks of

fitness. Animals show a wide range of social behaviours, including cooperative breeding

[13], group living [3], and hierarchical structures that may better protect their young [32],

their territory [95], *etc.* Though much research has examined the demographic correlates

of animal sociality, these investigations are typically constrained by taxonomic siloes

[96]. Indeed, much animal sociality research focuses on single species, or on mammals

[4] or birds [27]. Here, by proposing a sociality continuum that ranks species according to

the type, spatial, and temporal extent of their interactions, from solitary, to gregarious, to

communal, to colonial, to social, I show that, across 152 animal species from 13

taxonomic classes, as the degree of these interactions increases, species life histories

attain longer, more iteroparous windows of reproduction[38], even after accounting for

the allometric effects of body mass [54,68] and potential constraints of phylogenetic

inertia [89,90]. My comparative phylogenetic analyses also show that these social

interactions do not shape long-term metrics of performance, such as population growth

rate, contrary to the 'social buffer' hypothesis [42], but that the sociality continuum is

associated with some key attributes of demographic resilience [67].

A wealth of scientific evidence highlights the profound impacts of sociality on life history

traits in animals. Indeed, social structures often influence reproductive strategies, with

cooperative breeding systems, for example, promoting delayed sexual maturity and

extended parental care [97]. In species exhibiting eusociality, such as certain bees and

ants, caste systems shape life history trajectories for individuals within the colony [98,99].

Sociality can also affect reproductive success, as observed in the intricate mating

systems of various primates [100] and birds [13,101]. Moreover, the presence of social

alliances may impact longevity and survivorship, providing advantages in terms of

predation avoidance and access to resources [32,80]. Here, I show that the sociality

continuum is associated with key life history traits such as prolonged generation time,

postponed age at maturity, and increased mean, maximum, and variance of longevity,

longer reproductive windows, and greater probability of achieving reproduction before

dying. It is worth noting, however, that the interplay between sociality and life history

traits extends beyond reproduction and survival, influencing factors like dispersal

patterns [102] not taken into account in the present study. To understand this complexity,

future work must consider species-specific adaptations and ecological contexts and their

dispersal/migration abilities, employing a combination of field observations, experimental

manipulations [103], and modelling approaches based on big data [104].

In the present study, the sociality continuum does not correlate with the rate of actuarial

nor reproductive senescence across the examined animal species. The impacts of

sociality on senescence in animals has been a subject of extensive research, with

findings offering mixed perspectives [3]. Studies on various species indicate that social

interactions can either accelerate [105] or decelerate senescence processes [106]. For

instance, in certain cooperative breeding systems, where individuals collectively care for

offspring, sociality may delay reproductive senescence due to shared parental

responsibilities and increased chances of offspring survival [27,107,108]. Conversely, in

some species with intense intrasexual competition, the pressure to secure mates and

resources may accelerate reproductive senescence [109]. Actuarial senescence,

reflected in declining survival rates with age, can also be influenced by social factors. For

example, the presence of social allies may enhance protection against predation,

potentially extending lifespan [80]. However, the stress associated with social hierarchies

and conflicts can have the opposite effect, accelerating actuarial senescence [110]. The

complex interplay between social dynamics and senescence, and the multiple ways

through which senescence can manifest itself [111], underscore the need for a nuanced

understanding that considers species-specific traits and environmental contexts.

Ultimately, a combination of the comparative approach used here (to identify patterns

and plausible proximal mechanisms), coupled with experimental manipulations (to

identify and understand distal mechanisms), will provide the ultimate understanding of

the conditions under which sociality accelerates or slows down senescence in animals.

Similar approaches will be necessary to disentangle the direction of causality between

sociality and demography. The macroecological approach presented here identifies

patterns and likely candidate species for further study (*e.g.*, species around the periphery

of the life history strategy space presented in Fig. 5) but, as any macroecological

exercise [112], mine cannot pinpoint the exact mechanism nor imply causality in the

relationships I report. Nonetheless, some comparative work in birds has shown that

families with high proportions of cooperation have high survival values, even in non-

cooperative breeders in the family [13], suggesting a demography → sociality

directionality. Indeed, phylogenetic approaches exist to reconstruct ancestral trait values

along phylogenetic hypotheses [113], which would allow one to explore whether shifts in

life history traits cause or are followed by changes in sociality along species lineages

[114]. However, I argue that these approaches bring too much uncertainty, as both life

history traits and sociality are too labile for phylogenetic reconstructions to produce

accurate estimates [89]. Instead, I propose that the best way to prove causality between

degree of sociality and species' demographies requires the collective effort of

behavioural biologists, examining populations under the same set of control and

treatment conditions, and to follow multiple generations.

The relationship between sociality and the ability of animals to withstand environmental

stochasticity has garnered substantial attention. Numerous studies suggest that sociality

can serve as a buffering mechanism [42], providing a range of benefits that enhance

fitness in stochastic environments. While the social buffer hypothesis predicts that

sociality should confer adaptive advantages in mitigating the impact of environmental

stochasticity, I found that sociality is not associated with any of the metrics of long-term

population performance that are often used as population-level proxies for fitness

[44,115]. For instance, when evaluating how far from a population growth rate at

equilibrium the 152 species' population were, there was no detectable correlation with

sociality. I did, however, find that while more social species do not take a different

amount of time to recover after a disturbance compared to less social species, the period

of oscillation back to stationary dynamics for the former is greater than in less social

species, implying a potentially greater risk of extinction for social species [116,117]. This

finding is at odds with the suggestion that sociality increases plasticity against climatic

oscillations [118]. Further work integrating phenotypic plasticity and demographic

schedules should be key in this context [119]. In this context too, future work should

examine how the degree of sociality may vary across populations within the same

species. Likewise, the reactivity of social species was lower than in gregarious species,

suggesting that, when a population is hit by a disturbance, it is less likely to take

advantage by increasing its population size [70]. The past decades have witnessed the

development of indexes of demographic buffering based on structured population models

[120–123]. Future research should evaluate whether the sociality continuum is

intrinsically linked to a potential demographic lability-buffering continuum [124].

Species are more than just social or not social. Thus, a continuum to classify animal

sociality is of high appeal. A previous attempt has, however, highlighted some key

challenges [24]. Indeed, sociality is shaped by multiple axes, including group size, the

kind, timing, and hierarchical nature of interactions, or the partition of labour, to mention

a few. The inspiration for my proposed sociality continuum comes from work my

laboratory and I carried out to examine the anatomical correlates of actuarial senescence

[37]. Finch [125] hypothesised that the ability to compartmentalise risk may allow species

to postpone the onset of senescence. Classifying the compartmentalisation of 138 plants

(proxied by the distance between their xylem vessels) and 151 animals (proxied by the

degree of open-closed circulatory and respiratory systems, and degree of organ

redundancy), we found that modularity does predict actuarial senescence in plants but

not in animals. Our work there was only focused on vascular plants, for which a standard

measure of xylem distance can be obtained [126]. The classification of a continuum of

modularity-integration in animals proved difficult then because of the wide range of

variation in animal anatomy. Another way to think about animal anatomy is to examine

the 'anatomy' of their groups and populations: how close are their individuals, how often

do they come in contact, and what kinds of interactions do they have. In this regard,

Albery and colleagues recently found that older, more senescent (*i.e.*, likely to die)

individuals of red deer (*Cervus elaphus*) become more isolated in the group [8], perhaps

as a way to contain risk too, as plants do [37]. My proposed sociality continuum

correlates with key aspects of the demography of animals. However, further work should

examine orthogonal ways in which animal societies evolve, as the predictive power of the

correlations examined here remains low (Table 1).

The future of research on animal sociality and demography should prioritise

interdisciplinary approaches to analyse a large number of longitudinal studies. This

approach will help us unravel the nuanced complexities of social structures and their

demographic implications. Integrating advanced technologies, such as GPS tracking

[127], social network analysis [8], and comparative genomics [128], can provide a more

comprehensive understanding of the dynamic interactions within animal societies.

Embracing comparative analyses across a diverse array of taxa can uncover general

principles while acknowledging species-specific nuances. As conservation issues

become increasingly urgent, linking social and demographic research to conservation

biology can inform strategies for preserving biodiversity in the face of environmental

challenges [129]. This multidimensional approach holds the key to advancing our

understanding of the intricate interplay between sociality and demography in the animal

kingdom.

Acknowledgments

I thank J. Firth and K. Davis for conversations regarding the interface of animal sociality

in demography; J. Cant and B. Sheldon for input into the classification of sociality of

fishes and birds, respectively; the thousands of researchers who have contributed open-

access data to the COMADRE Animal Matrix Database (www.compadre-db.org); M.

Paniw and E. Conquet for providing access to the demographic models of *Suricata*

suricatta; K. Davis for orthographic edits; and G. Albert, J-M. Gaillard, and two

anonymous reviewers for constructive feedback on an earlier version of this work I was

supported by a NERC Pushing the Frontiers grant (NE/X013766/1).

References

1. Oro D. 2020 *Perturbation, Behavioural Feedbacks, and Population Dynamics in Social Animals: When to Leave and Where to Go*. Oxford University Press.

2. Polgar S. 2011 *Population, Ecology, and Social Evolution*. Walter de Gruyter.

3. Lucas ER, Keller L. 2020 The co□evolution of longevity and social life. *Funct. Ecol.* **34**, 76–87.

4. Blumstein DT, Williams DM, Lim AN, Kroeger S, Martin JGA. 2018 Strong social relationships are associated with decreased longevity in a facultatively social mammal. *Proc. Biol. Sci.* **285**. (doi:10.1098/rspb.2017.1934)

5. Sakata JT, Gupta A, Chuang C-P, Crews D. 2002 Social experience affects territorial and reproductive behaviours in male leopard geckos, Eublepharis macularius. *Anim. Behav.* **63**, 487–493.

6. Amdam GV, Csondes A, Fondrk MK, Page RE Jr. 2006 Complex social behaviour derived from maternal reproductive traits. *Nature* **439**, 76–78.

7. Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, Firth JA. 2021 Multiple spatial behaviours govern social network positions in a wild ungulate. *Ecol. Lett.* **24**, 676–686.

8. Albery GF, Clutton-Brock TH, Morris A, Morris S, Pemberton JM, Nussey DH, Firth JA. 2022 Ageing red deer alter their spatial behaviour and become less social. *Nature Ecology & Evolution* **6**, 1231–1238.

9. Bateman AW, Ozgul A, Krkošek M, Clutton-Brock TH. 2018 Matrix Models of Hierarchical Demography: Linking Group- and Population-Level Dynamics in Cooperative Breeders. *Am. Nat.* **192**, 188–203.

10. Paniw M, Maag N, Cozzi G, Clutton-Brock T, Ozgul A. 2019 Life history responses of meerkats to seasonal changes in extreme environments. *Science* **363**, 631–635.

11. Paniw M, Duncan C, Groenewoud F, Drewe JA, Manser M, Ozgul A, Clutton-Brock T. 2022 Higher temperature extremes exacerbate negative disease effects in a social mammal. *Nat. Clim. Chang.* **12**, 284–290.

12. Groenewoud F, Clutton-Brock T. 2021 Meerkat helpers buffer the detrimental effects of adverse environmental conditions on fecundity, growth and survival. *J. Anim. Ecol.* **90**, 641–652.

13. Arnold KE, Owens IPF. 1998 Cooperative breeding in birds: a comparative test of the life history hypothesis. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **265**, 739–745.

14. Koykka C, Wild G. 2016 The association between the emergence of cooperative breeding and clutch size. *J. Evol. Biol.* **29**, 58–76.

15. Poiani A, Pagel M. 1997 EVOLUTION OF AVIAN COOPERATIVE BREEDING: COMPARATIVE TESTS OF THE NEST PREDATION HYPOTHESIS. *Evolution* **51**, 226–240.

16. Salomon M, Lubin Y. 2007 Cooperative breeding increases reproductive success in the

social spider Stegodyphus dumicola (Araneae, Eresidae). *Behav. Ecol. Sociobiol.* **61**, 1743–
1750.

17. Poethke HJ, Liebig J. 2008 Risk-sensitive foraging and the evolution of cooperative breeding
and reproductive skew. *BMC Ecol.* **8**, 2.

18. Groenewoud F, Frommen JG, Josi D, Tanaka H, Jungwirth A, Taborsky M. 2016 Predation
risk drives social complexity in cooperative breeders. *Proc. Natl. Acad. Sci. U. S. A.* **113**,
4104–4109.

19. Hellmann JK, Hamilton IM. 2014 The presence of neighbors influences defense against
predators in a cooperatively breeding cichlid. *Behav. Ecol.* **25**, 386–391.

20. Schoepf I, Schradin C. 2012 Better off alone! Reproductive competition and ecological
constraints determine sociality in the African striped mouse (Rhabdomys pumilio). *J. Anim.*
Ecol. **81**, 649–656.

21. Kappeler PM, Cremer S, Nunn CL. 2015 Sociality and health: impacts of sociality on disease
susceptibility and transmission in animal and human societies. *Philos. Trans. R. Soc. Lond.*
B Biol. Sci. **370**. (doi:10.1098/rstb.2014.0116)

22. Rehan SM, Richards MH, Adams M, Schwarz MP. 2014 The costs and benefits of sociality
in a facultatively social bee. *Anim. Behav.* **97**, 77–85.

23. Wong MYL. 2012 Abiotic stressors and the conservation of social species. *Biol. Conserv.*
155, 77–84.

24. Rubenstein DR, Botero CA, Lacey EA. 2016 Discrete but variable structure of animal
societies leads to the false perception of a social continuum. *R Soc Open Sci* **3**, 160147.

25. Perony N. 2012 *Comparative Analysis of Social Interactions in Animal Groups*. ETH.

26. Beauchamp G. 2010 A comparative analysis of vigilance in birds. *Evol. Ecol.* **24**, 1267–1276.

27. Vágási CI, Vincze O, Lemaître J-F, Pap PL, Ronget V, Gaillard J-M. 2021 Is degree of
sociality associated with reproductive senescence? A comparative analysis across birds and
mammals. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **376**, 20190744.

28. Weinstein M, Kaplan H, Lane MA. 2014 *Sociality, Hierarchy, Health: Comparative*
Biodemography. National Academies Press (US).

29. Woodroffe R. 2011 Demography of a recovering African wild dog (Lycaon pictus) population.
J. Mammal. **92**, 305–315.

30. Schülke O, Ostner J. 2012 Ecological and social influences on sociality. *The evolution of*
primate societies , 195–219.

31. Ostner J, Schülke O. 2018 Linking sociality to fitness in primates: A call for mechanisms. In
Advances in the Study of Behavior (eds M Naguib, L Barrett, SD Healy, J Podos, LW
Simmons, M Zuk), pp. 127–175. San Diego, CA: Elsevier.

32. Silk JB. 2007 Social components of fitness in primate groups. *Science* **317**, 1347–1351.

33. Bekoff M, Daniels TJ, Gittleman JL. 1984 Life history patterns and the comparative social
ecology of carnivores. *Annu. Rev. Ecol. Syst.* **15**, 191–232.

34. Kramer BH, Schrempf A, Scheuerlein A, Heinze J. 2015 Ant Colonies Do Not Trade-Off
Reproduction against Maintenance. *PLoS One* **10**, e0137969.

35. Chouvenc T, Ban PM, Su N-Y. 2022 Life and Death of Termite Colonies, a Decades-Long
Age Demography Perspective. *Frontiers in Ecology and Evolution* **10**.
(doi:10.3389/fevo.2022.911042)

36. Buston P. 2003 Forcible eviction and prevention of recruitment in the clown anemonefish.
Behav. Ecol. **14**, 576–582.

37. Bernard C, Compagnoni A, Salguero-Gómez R. 2020 Testing Finch's hypothesis: The role of
organismal modularity on the escape from actuarial senescence. *Funct. Ecol.* **34**, 88–106.

38. Stearns SC. 1999 *The Evolution of Life Histories*. New York, USA: Oxford University Press.

39. de Kroon H, van Groenendael J, Ehrlén J. 2000 Elasticities: a review of methods and model
limitations. *Ecology* **81**, 607–618.

40. Stott I, Townley S, Hodgson DJ. 2011 A framework for studying transient dynamics of
population projection matrix models. *Ecol. Lett.* **14**, 959–970.

41. Roper M, Capdevila P, Salguero-Gómez R. 2021 Senescence: why and where selection
gradients might not decline with age. *Proc. Biol. Sci.* **288**, 20210851.

42. Hennessy MB, Kaiser S, Sachser N. 2009 Social buffering of the stress response: diversity,
mechanisms, and functions. *Front. Neuroendocrinol.* **30**, 470–482.

43. Salguero-Gomez R *et al.* 2016 COMADRE: a global data base of animal demography. *J.*
Anim. Ecol. **85**, 371–384.

44. Caswell H. 2001 *Matrix Population Models: Construction, Analysis, and Interpretation*. 2nd
edition. Sinauer Associates, Inc.

45. Jones OR *et al.* 2022 Rcompadre and Rage-Two R packages to facilitate the use of the
COMPADRE and COMADRE databases and calculation of life-history traits from matrix
population models. *Methods Ecol. Evol.* (doi:10.1111/2041-210x.13792)

46. Salguero-Gómez R, Plotkin JB. 2010 Matrix dimensions bias demographic inferences:
implications for comparative plant demography. *Am. Nat.* **176**, 710–722.

47. Jones OJ *et al.* 2014 Diversity of ageing across the tree of life. *Nature* **505**, 169–173.

48. Horvitz CC, Tuljapurkar S. 2008 Stage dynamics, period survival, and mortality plateaus.
Am. Nat. **172**, 203–215.

49. Monchot H, Fernandez P, Gaillard J-M. 2012 Paleodemographic analysis of a fossil
porcupine (Hystrix refossa Gervais, 1852) population from the Upper Pleistocene site of
Geula Cave (Mount Carmel, Israel). *J. Archaeol. Sci.* **39**, 3027–3038.

50. Conquet E, Ozgul A, Blumstein DT, Armitage KB, Oli MK, Martin JGA, Clutton-Brock TH,
Paniw M. 2023 Demographic consequences of changes in environmental periodicity.
Ecology **104**, e3894.

51. Stott I, Hodgson DJ, Townley S. 2012 popdemo: an R package for population demography
using projection matrix analysis. *Methods Ecol. Evol.* **3**, 797–802.

52. Stubben C, Milligan B. 2007 Estimating and analyzing demographic models using the popbio
package in R. *J. Stat. Softw.* **22**, 1–23.

53. Paniw M, Ozgul A, Salguero-Gomez R. 2018 Interactive life-history traits predict sensitivity of
plants and animals to temporal autocorrelation. *Ecol. Lett.* **21**, 275–286.

54. Healy K, Ezard THG, Jones OR, Salguero-Gomez R, Buckley YM. 2019 Animal life history is
shaped by the pace of life and the distribution of age-specific mortality and reproduction.
Nature Ecology & Evolution **3**, 1217–+.

55. Gaillard J-M, Yoccoz NG, Lebreton J-D, Bonenfant C, Devillard S, Loison A, Pontier D,
Allaine D. 2005 Generation time: a reliable metric to measure life-history variation among
mammalian populations. *Am. Nat.* **166**, 119–23; discussion 124–8.

56. Gaillard JM, Pontier D, Allaine D, Lebreton JD, Trouvilliez J, Clobert J. 1989 An analysis of
demographic tactics in birds and mammals. *Oikos* **56**, 59–76.

57. Dobson FS, Oli MK. 2008 The life histories of orders of mammals: Fast and slow breeding.
Curr. Sci. **95**, 862–865.

58. Morris WF *et al.* 2008 Longevity can buffer plant and animal populations against changing
climatic variability. *Ecology* **89**, 19–25.

59. Cochran ME, Ellner S. 1992 Simple methods for calculating age-based life-history
parameters for stage-structured populations. *Ecol. Monogr.* **62**, 345–364.

60. Salguero-Gómez R, Jones OR, Jongejans E, Blomberg SP, Hodgson DJ, Mbeau-Ache C,
Zuidema PA, de Kroon H, Buckley YM. 2016 Fast-slow continuum and reproductive
strategies structure plant life-history variation worldwide. *Proc. Natl. Acad. Sci. U. S. A.* **113**,
230–235.

61. Keyfitz N, Caswell H. 2005 *Applied Mathematical Demography*. New York.

62. Baudisch A. In press. The pace and shape of ageing. *Methods Ecol. Evol.*
(doi:10.1111/j.2041-210X.2010.00087.x)

63. Baudisch A, Stott I. 2019 A pace and shape perspective on fertility. *Methods Ecol. Evol.* **10**,
1941–1951.

64. Demetrius L. 1978 Adaptive value, entropy and survivorship curves. *Nature* **275**, 213–214.

65. de Kroon H, Plaisier A, van Groenendael J, Caswell H. 1986 Elasticity: the relative
contribution of demographic parameters to population growth rate. *Ecology* **67**, 1427–1431.

66. Orme D, Freckleton R, Thomson G, Petzoldt T, Fritz S, Isaac N, Pearse W. 2013 The caper
package: comparative analysis of phylogenetics and evolution in R. See http://cran.r-
project.org/web/packages/caper/.

67. Neubert MG, Caswell H, Solow AR. 2009 Detecting reactivity. *Ecology* **90**, 2683–2688.

68. Gamelon M, Gimenez O, Baubet E, Coulson T, Tuljapurkar S, Gaillard J-M. 2014 Influence
of life-history tactics on transient dynamics: a comparative analysis across mammalian
populations. *Am. Nat.* **184**, 673–683.

69. Capdevila P, Stott I, Cant J, Beger M, Rowlands G, Grace M, Salguero-Gomez R. In press.
Life history mediates the trade-offs among different components of demographic resilience.
Ecol. Lett. , 2021.06.30.450480.

70. Capdevila P, Stott I, Beger M, Salguero-Gomez R. 2020 Towards a Comparative Framework
of Demographic Resilience. *Trends Ecol. Evol.* **35**, 776–786.

71. Greenspoon L *et al.* 2023 The global biomass of wild mammals. *Proc. Natl. Acad. Sci. U. S.*
A. **120**, e2204892120.

72. Tobias JA *et al.* 2022 AVONET: morphological, ecological and geographical data for all

birds. *Ecol. Lett.* **25**, 581–597.

73. Froese R, Pauly D. 2002 FishBase: A Global Information System on Fishes. *World Wide*
Web electronic publication: www.fishbase.org. , 000–000.

74. Boettiger C, Lang DT, Wainwright PC. 2012 rfishbase: exploring, manipulating and
visualizing FishBase data from R. *J. Fish Biol.* **81**, 2030–2039.

75. Myhrvold NP, Baldridge E, Chan B, Sivam D, Freeman DL, Ernest SKM. 2015 An amniote
life-history database to perform comparative analyses with birds, mammals, and reptiles.
Ecology **96**, 3109–3000.

76. Bernard B, Santos G, Deere J, Capdevila P, Kusch E, Gascoigne S, Salguero-Gomez R.
2022 MOSAIC: A Unified Trait Database to Complement Structured Population Models.
bioRxiv **2022**. (doi:10.1101/2022.03.09.483599)

77. Silk JB. 2007 The adaptive value of sociality in mammalian groups. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **362**, 539–559.

78. Dunbar RIM. 1992 Neocortex size as a constraint on group size in primates. *J. Hum. Evol.*
22, 469–493.

79. Flack JC, Girvan M, de Waal FBM, Krakauer DC. 2006 Policing stabilizes construction of
social niches in primates. *Nature* **439**, 426–429.

80. Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM,
Seyfarth RM, Cheney DL. 2009 The benefits of social capital: close social bonds among
female baboons enhance offspring survival. *Proc. Biol. Sci.* **276**, 3099–3104.

81. Dewey T, Shefferly N, Havens A. 2010 Animal Diversity Web. *University of*

82. Baillie J, Hilton-Taylor C, Stuart SN, IUCN Species Survival Commission. 2004 *2004 IUCN*
Red List of Threatened Species: A Global Species Assessment. IUCN.

83. McTavish EJ, Hinchliff CE, Allman JF, Brown JW, Cranston KA, Holder MT, Rees JA, Smith
SA. 2015 Phylesystem: a git-based data store for community-curated phylogenetic
estimates. *Bioinformatics* **31**, 2794–2800.

84. Michonneau F, Brown JW, Winter DJ. 2016 rotl: an R package to interact with the Open Tree
of Life data. *Methods Ecol. Evol.* **7**, 1476–1481.

85. Plese B, Kenny NJ, Rossi ME, Cárdenas P, Schuster A, Taboada S, Koutsouveli V, Riesgo
A. 2021 Mitochondrial evolution in the Demospongiae (Porifera): Phylogeny, divergence
time, and genome biology. *Mol. Phylogenet. Evol.* **155**, 107011.

86. Revell LJ. 2013 R package phytools.

87. Salguero-Gomez R. 2021 The fast-slow continuum is not the end-game of life history
evolution, human or otherwise. *Evol. Hum. Behav.*

88. Revell LJ. 2009 Size-correction and principal components for interspecific comparative
studies. *Evolution* **63**, 3258–3268.

89. Blomberg SP, Garland T, Ives AR. 2003 Testing for phylogenetic signal in comparative data:
Behavioral traits are more labile. *Evolution* **57**, 717–745.

90. Freckleton RP. 2000 Phylogenetic tests of ecological and evolutionary hypotheses: checking
for phylogenetic independence. *Funct. Ecol.* **14**, 129–134.

91. van Buuren S, Groothuis-Oudshoorn K. 2013 MICE: multivariate imputation by chained
equations. See http://cran.r-project.org/web/packages/mice/index.html.

92. James TD, Salguero-Gómez R, Jones OR, Childs DZ, Beckerman AP. 2021 Bridging gaps
in demographic analysis with phylogenetic imputation. *Conserv. Biol.* **35**, 1210–1221.

93. Legendre P, Legendre L. 2012 *Numerical Ecology*. 3rd edn. London: Elsevier.

94. Montero AP, Williams DM, Martin JGA, Blumstein DT. 2020 More social female yellow-
bellied marmots, Marmota flaviventer, have enhanced summer survival. *Anim. Behav.* **160**,
113–119.

95. Höner OP, Wachter B, East ML, Runyoro VA, Hofer H. 2005 The effect of prey abundance
and foraging tactics on the population dynamics of a social, territorial carnivore, the spotted
hyena. *Oikos*

96. Blumstein DT, Møller AP. 2008 Is sociality associated with high longevity in North American
birds? *Biol. Lett.* **4**, 146–148.

97. Kreider JJ, Kramer BH, Komdeur J, Pen I. 2022 The evolution of ageing in cooperative
breeders. *Evol Lett* **6**, 450–459.

98. da Silva J. 2021 Life History and the Transitions to Eusociality in the Hymenoptera. *Frontiers
in Ecology and Evolution* **9**. (doi:10.3389/fevo.2021.727124)

99. Wilson EO, Hölldobler B. 2005 Eusociality: origin and consequences. *Proc. Natl. Acad. Sci.
U. S. A.* **102**, 13367–13371.

100.Kappeler PM, van Schaik CP. 2002 Evolution of primate social systems. *Int. J. Primatol.* **23**,
707–740.

101.Hatchwell BJ. 2009 The evolution of cooperative breeding in birds: kinship, dispersal and life
history. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **364**, 3217–3227.

102.Corcobado G, Rodríguez-Gironés MA, Moya-Laraño J, Avilés L. 2012 Sociality level
correlates with dispersal ability in spiders. *Funct. Ecol.* **26**, 794–803.

103.Beltran R *et al.* 2024 Integrating animal tracking and trait data to facilitate global ecological
discoveries.

104.Rubenstein DR, Abbot P. 2017 *Comparative Social Evolution*. Cambridge University Press.

105.Anderson JA, Johnston RA, Lea AJ, Campos FA, Voyles TN, Akinyi MY, Alberts SC, Archie
EA, Tung J. 2021 High social status males experience accelerated epigenetic aging in wild
baboons. *Elife* **10**. (doi:10.7554/eLife.66128)

106.Berger V, Lemaître J-F, Allainé D, Gaillard J-M, Cohas A. 2018 Early and Adult Social
Environments Shape Sex-Specific Actuarial Senescence Patterns in a Cooperative Breeder.
Am. Nat. **192**, 525–536.

107.Clutton-Brock T. 2002 Breeding together: kin selection and mutualism in cooperative
vertebrates. *Science* **296**, 69–72.

108.Kokko H, Johnstone RA, Wright J. 2002 The evolution of parental and alloparental effort in
cooperatively breeding groups: when should helpers pay to stay? *Behav. Ecol.* **13**, 291–300.

109.Reznick DN, Bryant MJ, Roff D, Ghalambor CK, Ghalambor DE. 2004 Effect of extrinsic
mortality on the evolution of senescence in guppies. *Nature* **431**, 1095–1099.

110. Verhulst S, Geerdink M, Salomons HM, Boonekamp JJ. 2014 Social life histories: jackdaw
dominance increases with age, terminally declines and shortens lifespan. *Proc. Biol. Sci.*
281, 20141045.

111. Shefferson R, Jones OR, Salguero-Gómez R. 2017 *The Evolution of Senescence in the*
Tree of Life. 1st edn. London: Cambridge University Press.

112. Dolby GA. 2021 Towards a unified framework to study causality in Earth-life systems. *Mol.*
Ecol. **30**, 5628–5642.

113. Goolsby EW. 2015 Phylogenetic Comparative Methods for Evaluating the Evolutionary
History of Function-Valued Traits. *Syst. Biol.* **64**, 568–578.

114. Downing PA, Cornwallis CK, Griffin AS. 2015 Sex, long life and the evolutionary transition to
cooperative breeding in birds. *Proc. Biol. Sci.* **282**, 20151663.

115. McGraw JB, Caswell H. 1996 Estimation of Individual Fitness from Life-History Data. *Am.*
Nat. **147**, 47–64.

116. Iles D, Koons D, Adler P, Salguero-Gomez R. 2015 Transient Dynamics, Life History, and
Biological Invasions. In *Student Research Symposium*,

117. Hodgson D, McDonald JL, Hosken DJ. 2015 What do you mean, 'resilient'? *Trends Ecol.*
Evol. **30**, 503–506.

118. Komdeur J, Ma L. 2021 Keeping up with environmental change: The importance of sociality.
Ethology **127**, 790–807.

119. Vinton AC, Gascoigne SJL, Sepil I, Salguero-Gómez R. 2022 Plasticity's role in adaptive
evolution depends on environmental change components. *Trends Ecol. Evol.* **37**, 1067–
1078.

120. Pfister CA. 1998 Patterns of variance in stage-structured populations: evolutionary
predictions and ecological implications. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 213–218.

121. Tuljapurkar S. 1989 An uncertain life: demography in random environments. *Theor. Popul.*
Biol. **35**, 227–294.

122. Le Coeur C, Yoccoz NG, Salguero-Gómez R, Vindenes Y. 2022 Life history adaptations to
fluctuating environments: Combined effects of demographic buffering and lability. *Ecol. Lett.*
25, 2107–2119.

123. Gascoigne SJL *et al.* 2023 Structured demographic buffering: A framework to explore the
environment drivers and demographic mechanisms underlying demographic buffering.
bioRxiv. , 2023.07.20.549848. (doi:10.1101/2023.07.20.549848)

124. Santos GS, Gascoigne SJL, Dias ATC, Kajin M, Salguero-Gómez R. 2023 A unified
framework to identify demographic buffering in natural populations. *bioRxiv.* ,
2023.07.03.547528. (doi:10.1101/2023.07.03.547528)

125. Finch CE. 1990 *Longevity, senescence and the genome*. Chicago, USA: University of
Chicago Press.

126. Schenk HJ, Espino S, Goedhart CM, Nordenstahl M, Cabrera HIM, Jones CS. 2008
Hydraulic integration and shrub growth form linked across continental aridity gradients. *Proc.*
Natl. Acad. Sci. U. S. A. **105**, 11248–11253.

127. Cagnacci F. 2010 *GPS-telemetry Data: Challenges and Opportunities for Behavioural*
Ecology Studies. Royal society.

128. Tong C, Najm GM, Pinter-Wollman N, Pruitt JN, Linksvayer TA. 2020 Comparative
Genomics Identifies Putative Signatures of Sociality in Spiders. *Genome Biol. Evol.* **12**, 122–
133.

129. Coulson T, MacNulty DR, Stahler DR, Vonholdt B, Wayne RK, Smith DW. 2011 Modeling
Effects of Environmental Change on Wolf Population Dynamics, Trait Evolution, and Life
History. *Science* **334**, 1275–1278.

 Tables

Table 1. The degree of sociality correlates with key demographic traits across 152

animal species. Phylogenetic generalised least square (pgls) models of sociality and

different demographic traits. The demographic traits are classified into five general types:

A. Life history traits; **B**. Vital rates; **C**. Elasticities of population growth rate (λ) to vital

rates; **D**. Metrics of asymptotic (*i.e.,* long-term) performance; and **E**. Metrics transient

(*i.e.,* short-term) performance. **Bold**: $P < 0.05$. Pagel's λ quantifies the phylogenetic

signal in the model, and it ranges from 0 (*i.e.,*no phylogenetic signal) to 1 (*i.e.,* pattern

completely explained by phylogenetic relationships).

Type	Demographic trait	R^2	F	d.f.	P	Pagel's λ	95% CI
	Generation time (T)	0.077	3.080	5, 147	**0.018**	0.469	0.126 - 0.753
	Net reproductive output (R_0)	0.019	0.694	5, 147	0.597	0.000	0.000 - 0.197
	Mean life expectancy (η_e)	0.022	1.820	5, 147	**0.014**	0.745	0.496 - 0.882
	Variance in life expectancy ($\Delta\eta_e$)	0.065	2.563	5, 147	**0.041**	0.481	0.076 - 0.765
	Maximum longevity (L_{max})	0.009	1.511	5, 147	**0.044**	0.559	0.259 - 0.802
	Age at maturity (L_α)	0.004	0.146	5, 147	0.965	0.759	0.404 - 0.912
	Reproductive window ($L_{\alpha-\omega}$)	0.047	1.808	5, 147	**0.013**	0.759	0.507 - 0.895
	Actuarial senescence (s_{lx})	0.036	1.359	5, 147	0.251	0.167	0.000 - 0.541
	Probability of reproduction (p_R)	0.006	2.237	5, 147	**0.017**	0.515	0.199 - 0.781
A. Life history traits	Reproductive senescence (s_{mx})	0.042	1.615	5, 147	0.173	0.210	0.035 - 0.537
	Degree of parity (S)	0.033	1.270	5, 147	0.285	0.000	0.000 - 0.078

	Survival (σ)	0.016	0.613	5, 147	0.654	0.607	0.302 - 0.819
	Growth (γ)	0.046	1.760	5, 147	0.140	0.000	0.000 - 0.458
	Shrinkage (ρ)	0.031	2.162	5, 147	**0.033**	0.000	0.000 - 0.193
B. Vital rates	Reproduction (φ)	0.173	7.673	5, 147	**<0.001**	0.000	0.000 - 0.188
	Survival elasticity ($E\sigma$)	0.062	2.426	5, 147	**0.031**	0.321	0.064 - 0.686
	Growth elasticity ($E\gamma$)	0.142	6.066	5, 147	**<0.001**	0.000	0.000 - 0.101
	Shrinkage elasticity ($E\rho$)	0.031	1.172	5, 147	0.326	0.405	0.216 - 0.607
C. Elasticities	Reproduction elasticity ($E\varphi$)	0.014	2.523	5, 147	**0.032**	0.621	0.248 - 0.845
	Population growth rate (λ)	0.021	0.792	5, 147	0.532	0.000	0.000 - 0.277
D. Asymptotics	Deviance growth rate equilibrium ($|1-\lambda|$)	0.009	0.342	5, 147	0.849	0.089	0.000 - 0.437
	Damping ratio (ζ)	0.013	0.496	5, 147	0.739	0.156	0.000 - 0.543
	Period of oscillation (P_i)	0.067	2.652	5, 147	**0.036**	0.000	0.000 - 0.225
	Reactivity ($\underline{\rho}_1$)	0.025	2.940	5, 147	**0.024**	0.228	0.071 - 0.500
E. Transients	First step attenuation ($\underline{\rho}_1$)	0.010	3.381	5, 147	**0.012**	0.592	0.256 - 0.833

Figure captions

Figure 1. Geographic and phylogenetic representation of the 152 animal species used in this study to examine the correlates of sociality and demographic traits. Species sociality is represented along a discrete continuum, from solitary (blue), gregarious (green), communal (yellow), colonial (orange), to social (red). **A**. World map depicting the geographic location of the examined studies from the COMADRE Animal Matrix Database, when GPS coordinates were available (n = 98 species). **B**. Phylogenetic representation of the 152 study species. Numbered arcs represent the taxonomic classes (from left to right): 1: Scyphozoa (n = 1 species); 2: Anthozoa (n = 4); 3: Malacostraca (n = 1); 4: Insecta (n = 4); 5: Gastropoda (n = 4); 6: Bivalvia (n = 5); 7: Thaliacea (n = 1); 8: Elasmobranchii (n = 1); 9: Actinopterygii (n = 28); 10: Amphibia (n = 1); 11: Reptilia (n = 18); 12: Aves (n = 26); and 13: Mammalia (n = 58). Silhouette source: http://phylopic.org.

Figure 2. The continuum of sociality correlates with key life history traits. Boxplots of the residuals of life history traits after correction for adult body mass and phylogenetic relationships as a function of the discrete continuum of sociality for 152 animal species. The letters above each group represent post-hoc Tukey test significance levels from the phylogenetic ANOVA. Groups with different letters are significantly different at $P < 0.05$. The horizontal grey dashed line indicates no effect. See Table 1 for full statistics.

Figure 3. The continuum of sociality correlates with key vital rates and their elasticities to population growth rate. Boxplots of the residuals of vital rates (left: survival σ, growth γ, shrinkage ρ, and reproduction φ), and their elasticities to population growth rate λ (right)

after correcting for adult body mass and phylogenetic relationships, as a function of the

discrete continuum of sociality for 152 animal species. The letters above each group

represent post-hoc Tukey test significance levels from the phylogenetic ANOVA. Groups

with different letters are significantly different at $P < 0.05$. The horizontal grey dashed

line indicates no effect. See Table 1 for full statistics.

Figure 4. The continuum of sociality correlates with key attributes of transient (short-

term) dynamics that predict the inherent ability of natural populations to respond to

disturbances. Boxplots of the residuals of four transient dynamics metrics after correction

for adult body mass and phylogenetic relationships, as a function of the discrete

continuum of sociality for 152 animal species. Damping ratio (ζ) quantifies the speed of

recovery post disturbance; Period of oscillation (P_i) quantifies the number of cycles a

population goes through before achieving asymptotic dynamics; Reactivity ($\underline{\rho}_1$) quantifies

how much a population can grow after one time step following a disturbance, while first

step attenuation ($\underline{\rho}_1$) quantifies the loss of resistance after one time step. The letters

above each group represent post-hoc Tukey test significance levels from the

phylogenetic ANOVA. Groups with different letters are significantly different at $P < 0.05$.

The horizontal grey dashed line indicates no effect. See Table 1 for full statistics.

Figure 5. The life history strategy space of animals predicts the placing of sociality along

the reproductive strategies continuum, but not the fast-slow continuum. Adult body-mass,

phylogenetically corrected principal component analyses (pPCA) of seven life history

traits shown in Table 1A and Figure 2. These traits are colour-coded by investments in

survival and reproduction (black - generation time, T), survival only (grey - age at

maturity, L_α; mean life expectancy, η_e, or reproduction only (red - net reproductive output,

R_0, reproductive window, $L_{\alpha\text{-}\omega}$; probability of achieving reproduction before dying, p_R;

degree of iteroparity, S). Each point in the PCA space corresponds to an animal species,

with some represented by a silhouette (source: http://phylopic.org; refer to Fig. 1B). The

loadings of the life history traits identifies: PC1 (explaining 33.72% of the variance) as the

reproductive strategies continuum, with species with grater net reproductive output (R_0),

longer reproductive windows ($L_{\alpha\text{-}\omega}$), higher probability to reproducing before dying (p_R),

and higher degree of iteroparity (S) to the right; and PC2 (explaining 26.20% of the

variance) as the fast-slow continuum, with species living longer (η_e), postponing their first

reproductive event (L_α), and having greater generation times (T) towards the top. The

side boxplots correspond to the groups along PC1 (top) and PC2 (right) as a function of

the discrete continuum of sociality.

Figures

Figure 1

 Figure 2

 Figure 3

Figure 4

934   **Figure 5**

938