Single-Cell Local Stress Analysis in Tumoroids.

Rick Rodrigues de Mercado[1], Klara Beslmüller[2], Daan Vorselen[3],
Erik H.J. Danen[2], Thomas Schmidt[1,*]

[1]Physics of Life Processes, Leiden University, The Netherlands
[2]Leiden Academic Centre for Drug Research, Leiden University, The Netherlands
[3]Department of Cell Biology & Immunology, Wageningen University & Research, The Netherlands
*corresponding author: schmidt@physics.leidenuniv.nl

Abstract

The reciprocal interplay between cancer cells and their local environment, mediated by mechanical forces, necessitates a deeper experimental understanding. This requires precise quantitative measurements of cellular forces within the intricate three-dimensional context of the extracellular matrix. While methods such as traction-force microscopy and micropillar-array technology have effectively reported on cellular forces in two-dimensional cell culture, extending these techniques to three dimensions has proven exceedingly challenging. In the current study, we introduced a novel approach utilizing soft, elastic hydrogel microparticles, resembling the size of cells, to serve as specific and sensitive traction probes in three-dimensional cell culture of collagen-embedded tumoroids. Our methodology relies on high-resolution detection of microparticle deformations. These deformations are translated into spatially resolved traction fields, reaching a spatial resolution down to $1\,\mu m$ and thereby detecting traction forces as low as $30\,Pa$. By integrating this high-resolution traction analysis with three-dimensional cell segmentation, we reconstructed the traction fields originating from individual cells. Our methodology enables us to explore the relationships between cellular characteristics, extracellular traction fields, and cellular responses. We observed that cellular stresses ranged from 10 to 100 Pa, integrating to cellular forces from 0.1 to 100 nN, which correlated with the localization of the cells actin skeleton, and the interaction area that cells developed towards the microparticles. Interestingly, the interaction of cells with inert microparticles appeared to be governed by contact mechanics resembling that two soft particles. The methodology presented here not only addresses the challenges of extending traditional stress-probe techniques to three dimensions, but also opens a strategy for the study of specific interactions between cells and the local tumoroid environment in a strive to fully understand cell-matrix reciprocity in tissue.

1 Introduction

Metastasis is the primary cause of cancer mortality [1]. During cancer metastasis, cells dissociate from the primary tumor and spread to distant sites in the body. The metastatic potential of cancer cells often go hand in hand with therapy resistance [2]. All pathological signatures of metastasis, tumor growth, local tissue invasion, dissemination and therapy response are thereby controlled by biochemical as well as mechanical crosstalk between the tumor and its tissue microenvironment. An understanding of the mechanisms underlying such crosstalk in metastasis is hence crucial, and might lead to the potential development of novel lines of therapies. Here, we zoom in on the reciprocal mechanical interaction between tumor cells and their microenvironment that contributes to cell movement. Reciprocal interaction relates to the relation between active forces of cells, and the mechanical characteristics of the extracellular matrix (ECM) [3]. Essential questions to be addressed for understanding the mechanoreciprocity in cancer are, to what extent do cell-generated forces enhance their metastatic potential, in how far are the active mechanical properties of malignant cells different to those of non-malignant or healthy cells, and in how far are we able to modulate mechanical phenotypes of cells.

The role of mechanical properties of cancer cells [4] as well as that of the microenvironment, including cell or matrix stiffness, has only been studied in recent years. Most of the studies investigated the mechanical properties of tumor cells growing in cell cultures in a two-dimensional substrate dish. Methods, like traction-force microscopy either using flat soft gels [5], micropillar arrays [6], atomic-force microscopy [7, 8], and optical stretcher devices [9], are exquisitely able to report on the mechanical properties of individual cells and on cells growing in 2D. Extensions of these established methods to

three dimensions rely on measuring deformations of the complex extracellular network surrounding the cells, the mechanical properties of which being highly non-linear and inhomogeneous, which makes the extraction of cellular forces in tissue extremely difficult [10]. Yet, metastasis formation indisputably is a process that occurs in the more complex environment of a three-dimensional tissue. It has been shown that force generation and migratory behavior of cancer cells in 2D differ significantly to those in which cancer cells are embedded in a 3D ECM [11, 12].

The ECM thereby plays an important role in promoting or inhibiting cell migration during cancer metastasis. The ECM is composed of a large ensemble of proteins, including fibrillar network-forming proteins like collagen or fibronectin. Tumors produce ECM components, proteolytically degrade ECM components, and bind and exert forces onto ECM components through adhesion receptors such as integrins [3, 13]. In turn, the chemical, mechanical, and structural features of the ECM fibrous network influence tumor growth and therapy response as well as tumour cell motility and invasion [4, 14–16]. For example, cancer cells often show increased collagen degradation and remodeling compared to normal cells, which in this way contribute to the formation of invasive tumors [17]. Conversely, it has been shown that the mechanical properties of collagen, such as the stiffness, may influence the migration of cancer cells.

In order to further understand the interaction between cells and their environment, 3D *in vitro* assays have been established in which tumor and ECM interact much alike to the situation in the corresponding tissue [18, 19]. These 3D culture systems closely mimic the *in vivo* pathophysiological situation in human tumor tissue. Spherical aggregates of tumour cells, so called multicellular tumor spheroids (tumoroids), are thereby embedded in reconstituted ECM environments [20]. Such reconstituted ECM environment give structural support for cell attachment and can be made from different types of elastic hydrogels [21], ideally with equivalent mechanical properties to the natural ECM.

In earlier studies we have established automated printing of cell clusters into 3D fibrillar ECM networks and used this approach to show how tumoroids exert forces onto the surrounding ECM to locally reshape their environment [20, 22]. Forces exerted by single cells embedded in 3D ECM have been studied by analyzing the displacement of small fiducial markers that were randomly embedded in the matrix before cross-linking the gel [23]. Such an approach had some success, yet needed a detailed knowledge of the local mechanical behavior of the fibrous network, being difficult to model. Therefore it is advantageous to have methodology at hand that permits determination of small cellular forces in three dimensions directly. One approach was the development of cell-sized "pressure sensors", gel-beads with randomly embedded fiducial markers. [24–27]. Displacement of the markers thereby report on the deformation of the sensor which was subsequently translated into local traction at the resolution of the pressure-sensor.

Here we present an advanced methodology that reports on local stress at the sub-cellular, micrometer, length scale inside of ECM-embedded tumoroid cultures. The methodology does not require prior knowledge of the mechanical characteristics of the fibrous environment, and still has a resolution that permits to assign local stresses to the action of individual cells. The methodology uses soft hydrogel microparticles, and is applied in a study of the mechanical properties of cancer cells as they migrate through a collagen matrix. The cell-sized (20-50 µm) microparticles produced have mechanical homogeneous and isotropic properties that can be easily tuned in a wide range of stiffness between 100 Pa and 10 kPa, spanning the stiffness range of most cells and tissues.

In our study we aimed to characterize the interactions of individual cells with their neighbouring cells inside a tumoroid, and of individual cells with the collagen fibers of the ECM network. Microparticles were coated with bovine-serum albumin which prevented specific interactions of cells with the particles, and rendering the particles as passive probes for local stresses. Forces generated by cells resulted in a local stress-field as observed by local deformations of the microparticles. The homogeneous mechanical properties of the microparticles, subsequently allowed us to quantitatively translate the microparticle deformations into traction-fields at the particles surface to a resolution of $\approx 1\,\mu$m and traction fields as low as $\approx 30\,$Pa. Our analysis was combined with a 3D cell-segmentation. The accurate three-dimensional shape of individual cells in an image stack was determined. Together with the segmentation we obtained a detailed, cell-centered traction-field, which permits us to associate extracellular traction fields, and cellular response to a quantitative phenomological description of cell behavior.

2 Materials and methods

2.1 Microparticle synthesis

Microparticle synthesis and functionalization was performed similar to previously published [28]. Deformable acrylamide-co-acrylic-acid microparticles were synthesized using Shirasu Porous Glass membranes (SPG Thechnology, Japan). These tubular membranes with tunable pore size enabled us to produce spherical particles in large quantities and of uniform size in a range of 5-50 μm. Before use, the SPG-membrane was submerged in HPLC-grade N-heptane (Merck, 1043792511) and placed in a vacuum chamber to remove air bubbles. After sonicating the membrane for 5 min in N-heptane, the membrane was mounted into the internal pressure micro kit extruder (SPG Technology). A solution of 200 mL hexane (Fisher Science P216885) with 3 % (w/v) surfactant span80 (Fluka, 85548) was prepared and poured into a 3-neck water-jacketed flask together with a magnetic stirrer bar. This mixture defines the continuous phase. The flask was connected to a water bath, a temperature sensor was inserted, put under a nitrogen sparge, and placed on a hotplate. The solution was stirred at 350 RPM.

A solution of 150 mM NaOH, 0.3 % (v/v) tetramethylethelenediamine (TEMED; Thermo Fisher, 17919) and 150 mM MOPS sodium salt (Sigma -Aldrich, M9381) was supplemented with acrylamide (AAm), acrylic acid (AAc) and cross-linker N,N'-methylenebisacrylamide (BIS) (all Sigma-Aldrich, A9099, 147230, 146072, respectively) with a final pH of 7.4. Two acrylamide mixtures were used for particle synthesis. The total mass concentration of acrylic components $c_T = c_{AAm} + c_{AAc} + c_{BIS} = 100$ mg/mL for both mixtures. The relative concentration of acrylic acid was set to 10 % for the first mixture. The second mixture had a relative concentration of acrylamide of 10 %. A cross-linker concentration $c_c = m_{BIS}/(m_{AAm} + m_{AAc} + m_{BIS})$ of 0.65 % and 1.5 % was used for the first and second gel mixture, respectively. The Young's modulus of the particles was characterized using atomic force microscopy. The first set of particles had a Young's modulus of $E \approx 1$ kPa, the second set of $E \approx 0.6$ kPa.

The acrylamide solution (dispersed phase) was degassed for 15 min. 10 mL of the dispersed phase was transferred to the reservoir of the SPG module. The reservoir was subsequently flushed with nitrogen. The critical pressure for particle generation was estimated from the difference in surface tensions of the two liquid-mixtures [28]. The starting pressure was set just below this value. The pressure was then slowly increased, until a flow was observed. The acrylamide mixture was allowed to be pushed through the membrane for 3 h. Subsequently, the SPG module was retracted, the flask was plugged and the temperature increased to 60 °C. Polymerization was initiated by adding 1.5 mg/mL 2,2'-Azobisisobutyronitrile (AIBN; Sigma-Aldrich, 441090) to the solution. After 3 h at 60 °C, the temperature was reduced to 40 °C. After 16 h, the particles were finally collected in round-bottom tubes and washed thrice in hexane, followed by a wash in ethanol. All ethanol was evaporated and particles were stored in PBS containing 5 mM sodium azide (Sigma-Aldrich, S20002).

2.2 Microparticle functionalization

First, particles were washed in activation buffer (100 mM MES (Sigma-Aldrich, M5057) and 200 mM NaCl), and subsequently incubated for 15 min in reaction buffer of 0.1 % tween20 (Sigma-Aldrich, P7949), 4 % (w/v) EDC (Sigma-Aldrich, E7750) and 2 % (w/v) NHS (Thermo Fisher, 22500). Then, the microparticles were washed and incubated for 1 h with 5 μg/mL of BSA in PBS at pH 8. To verify the efficiency of protein binding, Alexafluor-488 conjugated BSA was used. Fluorescent images confirmed the protein binding (Fig. 1 A). Subsequently, TexasRed-Cadaverine (Thermo Fisher, T2425) was added for 30 min. Unreacted NHS groups were blocked using 300 mM Tris and 100 mM ethanolamine (Sigma-Aldrich, 398136) (pH 9). Finally, particles were washed thrice in PBS with 0.1 % tween20 and stored for use in PBS (pH 7.4) with 5 mM sodium azide.

2.3 Atomic-force microscopy.

The Young's modulus of the particles was determined by atomic-force-spectroscopy on a JPK NanoWizard 4 atomic-force microscope (AFM; Bruker, Berlin). A plastic dish was activated using UV-ozone for 10 min. Particles were incubated for 15 min in activation buffer (see section "Microparticle functionalization") with 0.1 % tween20, 4 % (w/v) EDC and 2 % (w/v) NHS. Unbound particles were removed by washing with PBS. Measurements were performed in PBS at about 25 °C. For the measurements, spherical-shaped probes (Nanoandmore GmbH, Germany) were used (CP-qp-CONT-BSG-B-5) with a nominal spring constant of 0.1 N/m, and nominal tip diameter of 10 μm. The cantilevers were calibrated before each measurement with the thermal tuning method [29, 30], and also inspected with scanning

electron-microscopy (SEM) for accurate determination of cantilever dimensions and sphere radii. For determination of the elastic modulus we employed the shallow indentation method. Using an optical microscope, the spherical probe was centered on the particles. The indentation curves where performed to a force-maximum of 1 nN. To reduce noise, 4-5 nano-indentations were performed on each bead. Force-distance curves were corrected for cantilever bending. Each FD curve was transformed into a force-separation (FS) curve. This curve reflects only the sample indentation, excluding thereby the effect of the cantilever bending. The Young's modulus was obtained by fitting the FS-curves to the Hertz model with spherical indenter [31].

2.4 Cell culture

The human melanoma cell line MV3 [32] was cultured in high-glucose Dulbecco's Modified Eagle's Medium (DMEM, Gibco, 11504496) containing L-Glutamine and Sodium Pyruvate, supplemented with 10% fetal calf serum (Thermo Scientific) and 100 µg/mL penicillin/streptomycin in a humidified incubator at 37 °C with 5% CO_2.

2.5 3D collagen matrix

1 mg/mL collagen gels were created by mixing rat tail collagen type I (5 mg/mL stock, Biotechne, 3447-020-01) with buffers. Two different buffer types were used and compared: for HEPES buffering, we mixed DMEM (Gibco, ThermoFisher Scientific, Waltham, MA, USA), HEPES 0.1 M (1M stock, Biosolve, Valkenswaard, The Netherlands) and NaHCO3 44 mM (440 mM stock, Merck, Darmstadt, Germany) as previously described [33]. For phosphate buffering we mixed disodium hydrogen phosphate (Sigma Aldrich, Cat. No. 71636) and sodium dihydrogen phosphate (Sigma Aldrich, Cat. No. 71507) to pH value 7.4 and final phosphate concentration of 200 mM as described before [34]. Fluorescent particles were added to the collagen mixture to a concentration of 80p/uL. For the HEPES buffered collagen gel, fluorescent particles with a Young's modulus of 1 kPa were used, and for phosphate buffered collagen gel fluorescent particles of 0.6 kPa were used. The collagen-particle mixture was polymerized in a 96-well Sensoplate (Greiner Bio-one, 655892) at 37 °C for 1 h.

2.6 Tumoroids formation

Tumoroid formation into 3D collagen matrix was performed as described previously [20]. Cells were mixed with fluorescent microparticles in a ratio of approximately 1000:1 and resuspended in 2 % polyvinylpyrrolidone (Sigma-Aldrich, P5288) to reach a cell concentration of 25000 cells per µL shortly before cells were injected into collagen. Tumoroids, 200 µm in diameter, were created by automated injection of cell-particle mixture into the collagen gels using the injection robotics from Life Science Methods (Leiden, The Netherlands). After injection, tumoroids were incubated with the appropriate medium at 37 °C. After 24 h, tumoroids started to invade into the 3D collagen matrix. Samples were subsequently incubated in a fix-stain solution containing final concentrations of 2 % formaldehyde (Sigma-Aldrich, 252549), 0.1 % TritonX-100 (Sigma-Aldrich, T8787), 0.4 µg/mL Hoechst (Fisher Biotech, 33258) and 0.05 µM AlexaFluor-488 Phalloidin (Invitrogen, A12379) in PBS for 3 h at room temperature. The samples were washed thoroughly in PBS and measurements were performed shortly after.

2.7 Microscopy

2.7.1 Scanning confocal microscopy

Images of tumoroids were acquired on a Nikon Eclipse Ti inverted scanning confocal microscope equipped with four laser lines 405 nm, 488 nm, 561 nm and 640 nm, and with an A1R MP scanner. The microscope has a Nikon encoded and automated stage. Its camera is controlled through NIS Element Software (Nikon Instruments Inc., Melville, NY, USA). Tumoroids were imaged with 20 µm distance between Z-slices. ECM collagen-fibers were detected by confocal reflection microscopy on the same scanning confocal microscope. The collagen fibers were scanned at 561 nm excitation with a 561 nm blocking dichroic mirror and all the light reflected back passed a through filter of bandwidth 400 nm - 750 nm. The reflected light from the collagen fiber network was collected on GaAsP-photomultiplier. Scanning confocal fluorescent microscopy of tumoroids, and scanning confocal reflection microscopy of the collagen network were performed using Plan Apo ×20/0.75 NA, and Apo LWD ×40/1.15 objectives (Nikon Instruments Inc., Melville, NY, USA), respectively.

2.7.2 Spinning-disc confocal microscopy

High-resolution imaging was performed on an inverted microscope (Zeiss Axiovert 200) equipped with a 63X, 1.15 NA water LD C-Apochromat W korr objective (Zeiss) of 600 μm working distance. The setup was expanded with a confocal spinning-disk unit (Yokogawa CSU-X1), and an emCCD camera (Andor iXon DU897). The hydrogel microparticles were imaged with 561 nm DPSS-laser illumination (Cobolt). A 488 nm DPSL-laser (Coherent) was used to image the AlexaFluor-488 Phalloidin. A 405 nm diode laser (CrystaLaser) was used to illuminate the Hoechst nuclear-stain. Z-stacks were used to obtain a 3D image-cube of the microparticles and the surrounding cells. The mismatch between the refractive indices of the immersion-oil of the objective and the culture medium caused an apparent elongation of the image cube, which was corrected by experimentally-determined correction factor: the apparent elongation of > 10 stiff (> 5 kPa) microparticles was determined by calculating the radius in the equatorial plane compared to the lateral radius of the particles. The ratio of both was used as a correction factor for the z-positions, rendering the effect of the refractive index mismatch negligible.

2.8 Local traction analysis.

Fast traction analysis was performed in spherical harmonics space [35] implemented in the Python's SHTools library [36]. Spherical harmonics are solutions to the generalized linear elasticity continuity equation (Hooke's law) for the mechanical equilibrium condition of a spherical boundary in an isotropic medium. Hooke's law in terms of the displacement field $\mathbf{u}$ reads:

$$\mu\nabla^2\mathbf{u} + \left(\frac{\mu}{1-2\nu}\right)\nabla\left(\nabla\cdot\mathbf{u}\right) = 0, \tag{1}$$

where μ is the shear modulus of the spherical object (microparticle), and ν is the Poisson's ratio of the elastic medium. The displacement field in turn is expressed in complex coefficients, $\hat{u}_{jlm}$, of the spherical harmonics:

$$u_j(\theta,\phi) = \sum_{l=0}^{l_{max}}\sum_{m=-l}^{l}\hat{u}_{jlm}Y_l^m(\theta,\phi), \tag{2}$$

where $Y_l^m(\theta,\phi)$ are the set of complex spherical harmonics functions given by the harmonic-degree l and order m. The displacement field results in a traction field, $T_j(\theta,\phi)$:

$$T_j(\theta,\phi) = \sum_{l=0}^{l_{max}}\sum_{m=-l}^{l}\hat{T}_{jlm}Y_l^m(\theta,\phi), \tag{3}$$

where the components $\hat{T}_{jlm}$ are uniquely defined by $\hat{u}_{jlm}$.

Any local stress on the microparticles led to a local deformation of the particle's surface at the location of the stress, and to a small global deformation of the particle due to the counterbalancing action of the embedding ECM. In some cases this counterbalancing action was directly observed as local deformation along a thick ECM-fiber close to the particle's surface (Supplemental Fig. S1).

The spatial resolution of the traction field is limited by the maximum harmonic degree, l_{max}, in the spherical harmonics expansion. l_{max} was chosen based on the trade-off between computational power and desired spatial resolution. The spatial resolution, $\Delta r/R$, scales almost linearly with l_{max} (see supplemental Fig. S4)

$$\frac{\Delta r}{R} = 2.4\ \frac{1}{l_{max}^{0.93}}\ , \tag{4}$$

where R is the radius of the spherical object. Here we used $l_{max} = 25$, since this gives a good balance between spatial resolution (≈ 1 μm), and computational time (≈ 5 h per particle on an office PC).

2.9 Statistics

The Mann Whitney U-test was used to determine statistical significance between populations. Data sets were significantly different with probabilities of $p < 0.001$ (***), and not significantly different for probabilities of $p > 0.05$ (ns).

3 Results

3.1 High-resolution microparticle shape reconstruction.

The equilibrium shape of the elastic microparticles is that of a sphere. Application of a force to the particle will result in its deformation. Hence, the higher the resolution that the particle's surface is reconstructed, the smaller the minimum forces are that we are able to detect. Here we developed a high-resolution 3D particle-surface reconstruction methodology that allowed us to reconstruct the particle shape to the statistical limit set by the signal-to-noise ratio of detection.

All particles were homogeneously labeled by fluorophores during the preparation step of the particles (see M&M). Z-stacks of particles were taken on a spinning-disk confocal fluorescence microscope. The z-slices were separated by at a distance of $\Delta z = 370\,\text{nm}$ which is half the depth-of-focus of the objective, close enough to comply to Nyquist's theorem. Care has been taken to adjust the excitation intensity and illumination time such, to keep the emCCD-detector out of saturation. Typically image-cubes of $112 \times 112 \times 80\,\mu\text{m}^3$ were acquired. Individual particles in the image-cube were subsequently identified, and the data cubes cropped to contain solely individual particles (Fig. 1 A,B). These sub-cubes were further analyzed following an analysis pipeline similar to that described by Vorselen *et al.* [28].

In short, the particle's edge was highlighted by applying a 3D Sobel-operator to the image cube (Fig. 1 C). Subsequently, radial line profiles of the filtered edge-data and radiating from the particles' center-of-mass were calculated using cubic interpolation (Fig. 1 D,E). Line profiles were calculated on a regular grid of a perfect sphere at a distance equivalent to half the point-spread function of the microscope. Each of the radial profiles were fit to a Gaussian, returning the position of the edge, its width, and its integrated signal (Fig. 1 E). The local radius of the particle at the position shown in the line profile was 10.3 µm. The width returned from the edge-fit was, as predicted, equal to twice the width of the point-spread-function of the microscope ($\approx 350\,nm$). The high-resolution edge positions, all spaced on a regular grid on a sphere, were ultimately used for the high-resolution reconstruction of the particles' shape using triangulation (Fig. 1 F). The particle shown in (Fig. 1) had a mean radius of $R_0 = 10.1\,\mu\text{m}$. Determination of the high-resolution particle shape was exquisite. Analysis of ideal spherical particles on a coverslip and embedded in a buffer solution showed a root-mean-squared surface roughness of 50 nm, equivalent to the predicted edge localization error set by the photoncount level, and the width of the point-spread function (Fig. S2) [37].

In Fig. 1 F we clearly identified particle deformations of up to 1 µm, way beyond the optical resolution and the potential surface roughness. The particle shown in the figure had a nominal radius of 10.1 µm. The nominal radius was determined from the particle's total volume, as we assumed that the microparticle was incompressible and that volume was conserved when the sphere was deformed. The colors in Fig. 1 F indicate the sign of the deformation: red indicates indentations, blue protrusions.

Simultaneous with the edge detection, cells in close neighborhood to the microparticle were detected. Cellular signals where identified through labeling cells with an fluorescence stain of actin (Fig. 1 G). The actin signal (green in Fig. 1 E) located within 1 µm of the particles edge was summed (Fig. 1 E, dark green data points) to obtain an indicator for cellular vincinity, and actin concentration. The resulting stain signal-profile was then projected onto the edge-coordinate of the particle (Fig. 1 H). Clearly visible in Fig. 1 is that regions of high stain signal (Fig. 1 H) were colocalized with the indentations of the particle (Fig. 1 F). Cells obviously pushed on the particles which lead to the local particle deformations observed.

A global goodness-of-fit estimator, χ^2, for each particle surface reconstruction was calculated:

$$\chi^2 = \frac{1}{N_{DOF}} \sum_i \frac{(R_i^{SH} - R_i^{edge})^2}{R_i^{edge}}, \tag{5}$$

where, $N_{DOF} = d \cdot (l_{max} + 1)^2$, are the number of degrees of freedom in the fitting routine, with d the number of dimensions, R_i^{SH}, the radius at position i of the spherical harmonics displacement field, and, R_i^{edge}, the radial edge coordinate from the Gaussian fitting routine. Only particles of $\chi^2 < 0.5$ were considered for further analysis.

3.2 Microparticles in ECM-embedded tumoroids.

Fluorescent microparticles were used in tumoroid assemblies, in which tumoroids together with particles were embedded within a collagen matrix, resembling a native ECM environment. The ECM surrounding tumoroids consists of a network of collagen fibers that each were approximately 50-100 µm in length

(Fig. 2 C). The assembly was established by microinjection of a cell / microparticle suspension into a preformed gel. 24 h after injection MV3 cells had migrated into the ECM and away from the injection site (Fig. 2 A). Z-stacks of tumoroids were taken on a confocal microscope. Multiple particles were located within the ECM-embedded tumoroid. The microparticles were coated with bovine serum albumine, which rendered them inert with respect to cell interactions. The particles localized at random into the tumoroid (Fig. 2 B).

3.3   Single-cell traction decomposition.

In Fig. 1 F-H it appeared that indentations were most prominent where cells were localized close to the particle. For quantitation of this observation we developed methodology that automatically identified cells, and allowed us to correlate particle deformations with various cellular properties. Our methodology identified regions on the particles surface that were in proximity to individual cells, from which we deduced from each cell the amount of the force it applied to the particle.

3.3.1   Cell segmentation.

Cell segmentation inside a tumoroid within the 3D image-cube was developed based on an analysis pipeline described by Grosser *et al.* [38] (supplemental Fig. S3). The method required both the nucleus and the cell-membrane to be stained. In brief, an adaptive threshold was used to generate a 3D-mask of all nuclei in the image cube. The mask was dilated, holes were filled and subsequently eroded. Neighbouring nuclei were separated using a watershed algorithm. The nuclear centers were dilated to a size smaller than a typical nucleus (≈ 2 µm), serving as seed in the thresholded channel of the cytoskeletal actin defining the cell's cytosol, and resulting in the final segmentation of the image cube. Potential lower fluorescence levels at the cell edges were cut out by subtracting a morphologically opened image. Finally, salt-and-pepper noise was reduced by applying an adaptive median filter.

In addition, the boundary of the tumoroid was identified by thresholding a blurred, entropy-filtered image of the cell-edges. A Sobel filter was used to find the the cell edges, which were subsequently smoothed with a narrow kernel to merge peaks in the gradients of the same line segment. Finally, the image was superimposed with the nuclear seeds and watershed segmentation was performed. Such watershed segments are displayed in white in Fig. 3 A.

The quality of the fluorescent signal of both the nuclei and cell membranes determine the segmentation accuracy. Simulated data at equivalent signal-to-noise values to our measured image stacks showed, that > 80 % of the line segments were correctly detected.

3.3.2   Single-cell traction analysis.

In Fig. 3 A three melanoma MV3 cells (indexed 1-3), each at different z-position with respect to a central microparticle, were identified to interact with the single microparticle shown. The microparticle was localized at the border of a collagen gel-embedded tumoroid. To characterize the force each of the cells applied on the microparticle, their nuclei (blue) and cytosol (actin signal, green) were segmented as described above. The cell-segments are indicated in white in Fig. 3 A.

From the particle deformations (Fig. 3 B top) the traction field normal to the surface of the particle (Fig. 3 B 2^{nd}) was determined as further described in the M&M section. The normal traction ranged from -200 Pa (inward, red) to +200 Pa (outward, blue). Further fig. 3 B shows the locations of the nearby actin-skeleton of cells (3^{rd} row), and the respective cell indicators (1-3, bottom). In the bottom panel, regions on the particle unexposed to cells were identified as background (indicator "Bkg", grey).

As already noted in Fig. 1, indentations of the microparticles were visible where cells touched. Cells obviously applied pushing forces (negative normal traction) to the soft microparticles. As the particle shown was located near the periphery of the tumoroid, the largest part of the surface (861 µm², 66%) was not covered by cells. The contact areas of cells varied between 26 µm² (2%, blue cell "1") to 348 µm² (26%, dark green cell "3"). At areas where cells were in proximity to the particle, the traction normal to the surface was significantly different to those areas not covered by cells ("Bkg") ($p < 0.001$). The mean of the absolute normal traction varied between 33 Pa (blue, cell "1"), to 115 Pa (light green, cell "2"), and 77 Pa (dark green, cell "3") (Fig. 3 C). By combining cell segmentation with traction analysis our method thus allowed us to quantitatively describe the mechanical interactions that cells within tumoroids develop towards their direct tissue-like microenvironment.

Combining the values for the mean cell traction with those we obtained for the respective cell area touching the microparticle, we calculated the total force that each cell exerted on the particle. Those

forces varied between 0.85 nN (blue, cell "1"), to 8.0 nN (light green, cell "2"), and 27 nN (dark green, cell "3"). It should be noted, that clearly the overall force-balance holds. The microparticles were about 10-fold bigger than the pores in the collagen gel, hence force balance is faithfully guaranteed by a counter force of the gel being spread over the full surface of the particle. These counterbalancing forces are potentially the reason for the exceeding spread of the traction in the background region in Fig. 3 C (grey), as compared to our control measurements of microparticles in buffer solution.

We further analyzed the F-actin signal for each of the cells as a means to quantify the potential activity of the cells. The mean actin signal of cell "1" was significantly smaller than that of the two other cells touching the particle (Fig. 3 D). By comparing Fig. 3 C with Fig. 3 D it appeared that the mean traction correlated with the amount of actin in the contact area of cells with the particle. Such a correlation is to be predicted, since F-actin is the main force generating machinery cells have to develop active forces on their environment.

3.4 Cellular forces increase with F-actin density and interaction area.

We wanted to verify the correlations we observed for the individual particle shown in Fig. 3, and test the hypotheses that cells individually develop forces inside the tumoroid. The predictions include that cells apply larger forces when they have a more pronounced actin skeleton, and that cells apply more forces when they are able to make more contact with their environment. Both dependencies have been observed for cells grown on a 2D substrate [39–44], yet are still to be shown for the complex environment of a tumoroid.

For quantification of the level and density of the F-actin network, the F-actin signal, S_{act} needed to be quantified. Yet, the F-actin signal and the unavoidable background signal of each cell varied considerably depending on the location of the particle relative to the core of the tumoroid. Such variation is inherent to the measurements, and relies among others on the local labeling efficiency, the local intensity of the excitation light (shadowing), and on the local efficiency of the fluorescence detection. These parameters not only vary depending on the location within the tumoroid, but also from sample to sample. Yet all the above mentioned unknowns influence the signal inside the actin-free microparticle, S_{bkg}, in an identical manner. Thus, we here define the reduced actin signal S^*:

$$S^* = \frac{S_{act} - S_{bkg}}{\sigma_{bkg}}, \tag{6}$$

which corrects for experimental uncertainties and faithfully reports on the F-actin signal for all particles. The standard deviation of the background signal σ_{bkg} is taken as reference. This rescaling to S^* equalizes the background for all images, and allows for quantitative comparison of the actin signal in different experiments.

In total, $48 + 27$ cells interacting with $16 + 6$ microparticles in 2 independent experimental runs were used for further analysis. The ensemble data verified the correlations observed for the individual particle presented in Fig. 3. Cells expressing more F-actin (increase in S^*), applied more pushing on the particle (decrease in normal traction, Fig. 4 A). Cells for which the actin signal was similar to that of the background ($S^* \approx 0$) did not develop local traction. In summary, traction significantly increased above background for cells in which F-actin was prominently detected.

Further, we investigated how the forces that a individual cells exerted varied with the interaction area the cell had with the particle. The net force an individual cell applied on a microparticle was calculated by integration of the normal traction over the interaction area. Results are shown in Fig. 4 B. With increasing interaction area the magnitude of the net force increased. The dependence appeared to follow a power-law as judged from the linear scaling in a double-logarithmic representation. From the slope, m, in double-logarithmic representation ($m = 1.0$, cq $m = 1.3$) it appeared that the net force scaled proportional ($m = 1$), or slightly super-proportional ($m > 1$) to the interaction area. A power-law relation was likewise observed between the average traction and the net force (Fig. 4 C). From the fit to a linear dependence in the double-logarithmic representation ($m = 0.54$, cq $m = 0.65$), we found that the normal traction increased with about the square root of the force. Given the experimental uncertainties in those experiments, both observation comply with a model that describes the contact mechanics of two elastic spheres, as further detailed in the discussion section.

Cell-segmentation allowed us further to determine the cell morphology and cellular orientation. Given that morphology and orientation of cells was shown to predict cellular motility [45–48], we hypothesized that cells which were oriented head-on to the particle's surface would develop a stronger normal traction on the particle, as compared to cells passing by, and likewise that cells passing by would develop an

increased shear-traction on the bead. Overall cell morphology is typically expressed in terms of the cell shape-index, s, which is the ratio of the cell's radius determined from it's volume, $r_V = 3/4\pi \, V^{1/3}$, to that determined from it's surface, $r_S = 1/4\pi \, \sqrt{S}$ [38]

$$s = \left\{ \frac{S^3}{36 \, \pi \, V^2} \right\}^{1/3}. \qquad (7)$$

A perfect spherical cell would have $s = 1$, larger values indicated more elongated cells. Here we take the cell shape-index as an indicator of the elongation of a cell. Fig. 4 D shows the relation of normal traction as it varies with cell shape-index. Unlike our prediction, we did not find a correlation between shape index and normal traction (Fig. 4 D). Neither we find a correlation with the tangential traction of cells (see supplemental Fig. S5).

4 Discussion

In the current study, we present a novel approach for quantitative analysis of local cellular forces in complex three-dimensional tissue-mimicking environments. Our method entails the combination of bulk synthesis of elastic microparticles, with high-resolution microscopy, and traction analysis inside a three-dimensional *in vitro* tumoroid experiment. By incorporating a cell segmentation pipeline into our analysis, we were able to address questions regarding cell-based inhomogeneities in stress generation inside of a tumoroid.

Cells developed stresses between 0.1 and 100 Pa and local forces between 0.1 and 100 nN, both compatible with the compliance of the gels used ($\approx$500 Pa), and typical for values reported for various tissue [10, 23, 49, 50]. We predicted that cells apply larger forces when they produced more actin, and when they were able to make larger contact with their environment. Our experiments confirmed those predictions. Cells that expressed more F-actin applied more pushing force on the microparticles, while cells with lower F-actin levels showed no local traction force. Traction increased about linearly with the local actin concentration.

The net force exerted by cells on the microparticles increased with increasing interaction area, where the net force scaled about linear with the interaction area. These findings are not too surprising when compared to the well-known problem of two soft spherical bodies in mechanical contact [31, 51]. From Hertzian contact mechanics, a similar trend is predicted in which the force F of elastic spheres of average radius R and effective modulus E increases with final contact area A as, $F = \frac{4}{3\pi^{3/2}} \cdot E/R \cdot A^{3/2}$, hence scales with the contact area to the power 1.5 [31, 51]. Likewise our finding that the normal traction scales with the square root of the net force come close to that predicted by the Hertzian model which reads that the pressure at contact P_0, which we take as equivalent to the normal traction in our experiments, is given by, $P_0 = 1/\pi \left(6 \, E^2 \, F/R^2 \right)^{1/3}$. The deviations of our results from those ideal mechanical model predictions might be due to the crude assumption of a spherical cell for which the Hertzian models hold. It is worth to mention as conclusion that the magnitude of forces, in the nN-range, exerted by individual cells in the complex environment of a 3D matrix are consistent with those reported for cells plated on 2D surfaces obtained in traction force microscopy [39, 44, 52–54] and micropillar array techniques [43].

Surprisingly though, our study did not reveal a relation between cellular traction and cell morphology. Our data suggest that traction is independent of cell shape and orientation, in contrary to what we predicted from experiments we performed for cells in two-dimensions in a culture dish [48]. This negative result indicates that in complex three-dimensions environments further factors will determine cellular forces. Those potentially include specific cell-ECM interaction sites which could be realized by covalently attaching ECM components to the microparticles in future experiments.

Overall, the results suggest that cells within a tumoroid individually push on their environment, and that the degree of contact with other cells in their environment as well as expression level of cytoskeleton proteins affect the magnitude of the force exerted by the cells. Our study describes a new approach to characterize cellular forces in the context of their complex three-dimensional environment of the ECM. Here we provided first insights into the mechanics of cellular forces and and how those relate to tumor tissue mimicry. Our findings may represent a solid steppingstone for the use of the presented methodology in our strive to contribute to a better understanding of the cellular mechanics involved in tumor growth and metastasis.

Funding

This work is part of the research program "The Active Matter Physics of Collective Metastasis" with project number Science-XL 2019.022, which is financed by the Dutch Research Council (NWO).

Acknowledgments

We thank Aditya Aniruddha Sane MSc. for assistance with setting up the microparticle fabrication.

Author Contributions

RRM, KB, ED and TS designed the experiments. RRM and KB conducted the experiments. RRM performed data analysis. RRM, KB, DV, ED and TS wrote the manuscript. All authors reviewed and edited the manuscript.

Disclosures

The authors declare no conflicts of interest.

References

[1] Danny R. Welch and Douglas R. Hurst. Defining the Hallmarks of Metastasis. *Cancer Research*, 79(12):3011–3027, 6 2019.

[2] Felix Weiss, Douglas Lauffenburger, and Peter Friedl. Towards targeting of shared mechanisms of cancer metastasis and therapy resistance. *Nature Reviews Cancer*, 22(3):157–173, 2022.

[3] Sjoerd van Helvert, Cornelis Storm, and Peter Friedl. Mechanoreciprocity in cell migration. *Nature Cell Biology*, 20:8–20, 1 2018.

[4] Antonios N Gargalionis, Kostas A Papavassiliou, and Athanasios G Papavassiliou. Mechanobiology of solid tumors. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, page 166555, 2022.

[5] Tim Oliver, Ken Jacobson, and Micah Dembo. Traction forces in locomoting cells. *Cell Motility*, 31(3):225–240, 1995.

[6] Stefano Coppola, Ilaria Carnevale, Erik HJ Danen, Godefridus J Peters, Thomas Schmidt, Yehuda G Assaraf, and Elisa Giovannetti. A mechanopharmacology approach to overcome chemoresistance in pancreatic cancer. *Drug Resistance Updates*, 31:43–51, 2017.

[7] M. Lekka, P. Laidler, D. Gil, J. Lekki, Z. Stachura, and A. Z. Hrynkiewicz. Elasticity of normal and cancerous human bladder cells studied by scanning force microscopy. *European Biophysics Journal*, 28:312–316, 5 1999.

[8] Martin Benoit, Daniela Gabriel, Günther Gerisch, and Hermann E Gaub. Discrete interactions in cell adhesion measured by single-molecule force spectroscopy. *Nature cell biology*, 2(6):313–317, 2000.

[9] Falk Wottawah, Stefan Schinkinger, Bryan Lincoln, Susanne Ebert, Karla Müller, Frank Sauer, Kort Travis, and Jochen Guck. Characterizing single suspended cells by optorheology. *Acta biomaterialia*, 1(3):263–271, 2005.

[10] Julian Steinwachs, Claus Metzner, Kai Skodzek, Nadine Lang, Ingo Thievessen, Christoph Mark, Stefan Münster, Katerina E Aifantis, and Ben Fabry. Three-dimensional force microscopy of cells in biopolymer networks. *Nature Methods*, 13:171–176, 2 2016.

[11] Claudia Tanja Mierke. The fundamental role of mechanical properties in the progression of cancer disease and inflammation. *Reports on Progress in Physics*, 77(7):076602, 2014.

[12] Edna Cukierman, Roumen Pankov, Daron R Stevens, and Kenneth M Yamada. Taking cell-matrix adhesions to the third dimension. *Science*, 294(5547):1708–1712, 2001.

[13] Cecilia Bergonzini, Kim Kroese, Annelien J. M. Zweemer, and Erik H. J. Danen. Targeting integrins for cancer therapy - disappointments and opportunities. *Frontiers in Cell and Developmental Biology*, 10, 2022.

[14] Claudia Tanja Mierke. The matrix environmental and cell mechanical properties regulate cell migration and contribute to the invasive phenotype of cancer cells. *Reports on Progress in Physics*, 82(6):064602, 2019.

[15] Bircan Coban, Cecilia Bergonzini, Annelien J. M. Zweemer, and Erik H. J. Danen. Metastasis: crosstalk between tissue mechanics and tumour cell plasticity. *British Journal of Cancer*, 124:49–57, 1 2021.

[16] Karin A. Jansen, Dominique M. Donato, Hayri E. Balcioglu, Thomas Schmidt, Erik H.J. Danen, and Gijsje H. Koenderink. A guide to mechanobiology: Where biology and physics meet. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1853(11, Part B):3043–3052, 2015. Mechanobiology.

[17] Masoud Najafi, Bagher Farhood, and Keywan Mortezaee. Extracellular matrix (ecm) stiffness and degradation as cancer drivers. *Journal of cellular biochemistry*, 120(3):2782–2790, 2019.

[18] Paraic A Kenny, Genee Y Lee, Connie A Myers, Richard M Neve, Jeremy R Semeiks, Paul T Spellman, Katrin Lorenz, Eva H Lee, Mary Helen Barcellos-Hoff, Ole W Petersen, et al. The morphologies of breast cancer cell lines in three-dimensional assays correlate with their profiles of gene expression. *Molecular oncology*, 1(1):84–96, 2007.

[19] Claudia Fischbach, Ruth Chen, Takuya Matsumoto, Tobias Schmelzle, Joan S Brugge, Peter J Polverini, and David J Mooney. Engineering tumors with 3d scaffolds. *Nature methods*, 4(10):855–860, 2007.

[20] Hoa H. Truong, Jan de Sonneville, Veerander P.S. Ghotra, Jiangling Xiong, Leo Price, Pancras C.W. Hogendoorn, Herman H. Spaink, Bob van de Water, and Erik H.J. Danen. Automated microinjection of cell-polymer suspensions in 3d ecm scaffolds for high-throughput quantitative cancer invasion screens. *Biomaterials*, 33(1):181–188, 2012.

[21] Camille Jubelin, Javier Muñoz-Garcia, Laurent Griscom, Denis Cochonneau, Emilie Ollivier, Marie-Françoise Heymann, François M Vallette, Lisa Oliver, and Dominique Heymann. Three-dimensional in vitro culture models in oncology research. *Cell & Bioscience*, 12(1):155, 2022.

[22] Hayri E. Balcioglu, Bob van de Water, and Erik H. J. Danen. Tumor-induced remote ecm network orientation steers angiogenesis. *Scientific Reports*, 6:22580, 3 2016.

[23] Wesley R Legant, Jordan S Miller, Brandon L Blakely, Daniel M Cohen, Guy M Genin, and Christopher S Chen. Measurement of mechanical tractions exerted by cells in three-dimensional matrices. *Nature Methods*, 7:969–971, 12 2010.

[24] M. E. Dolega, M. Delarue, F. Ingremeau, J. Prost, A. Delon, and G. Cappello. Cell-like pressure sensors reveal increase of mechanical stress towards the core of multicellular spheroids under compression. *Nature Communications*, 8:14056, 4 2017.

[25] Erfan Mohagheghian, Junyu Luo, Junjian Chen, Gaurav Chaudhary, Junwei Chen, Jian Sun, Randy H. Ewoldt, and Ning Wang. Quantifying compressive forces between living cell layers and within tissues using elastic round microgels. *Nature Communications*, 9:1878, 12 2018.

[26] S. Girardo, N. Träber, K. Wagner, G. Cojoc, C. Herold, R. Goswami, R. Schlüßler, S. Abuhattum, A. Taubenberger, F. Reichel, D. Mokbel, M. Herbig, M. Schürmann, P. Müller, T. Heida, A. Jacobi, E. Ulbricht, J. Thiele, C. Werner, and J. Guck. Standardized microgel beads as elastic cell mechanical probes. *Journal of Materials Chemistry B*, 6:6245–6261, 2018.

[27] N. Träber, K. Uhlmann, S. Girardo, G. Kesavan, K. Wagner, J. Friedrichs, R. Goswami, K. Bai, M. Brand, C. Werner, D. Balzani, and J. Guck. Polyacrylamide bead sensors for in vivo quantification of cell-scale stress in zebrafish development. *Scientific Reports*, 9:17031, 12 2019.

[28] Daan Vorselen, Yifan Wang, Miguel M. de Jesus, Pavak K. Shah, Matthew J. Footer, Morgan Huse, Wei Cai, and Julie A. Theriot. Microparticle traction force microscopy reveals subcellular force exertion patterns in immune cell–target interactions. *Nature Communications*, 11:20, 12 2020.

[29] John E. Sader, Ian Larson, Paul Mulvaney, and Lee R. White. Method for the calibration of atomic force microscope cantilevers. *Review of Scientific Instruments*, 66(7):3789–3798, 1995.

[30] Jeffrey L. Hutter and John Bechhoefer. Calibration of atomic-force microscope tips. *Review of Scientific Instruments*, 64(7):1868–1873, 1993.

[31] Heinrich Hertz. Ueber die berührung fester elastischer körper. *crll*, 1882:156–171, 1882.

[32] Goos N. P. van Muijen, Kees F. J. Jansen, Ine M. H. A. Cornelissen, Dominique F. C. M. Smeets, Johan L. M. Beck, and Dirk J. Ruiter. Establishment and characterization of a human melanoma cell line (mv3) which is highly metastatic in nude mice. *International Journal of Cancer*, 48(1):85–91, 1991.

[33] Hoa H. Truong, Jiangling Xiong, Veerander P. S. Ghotra, Ella Nirmala, Lizette Haazen, Sylvia E. Le Dévédec, Hayri E. Balcioğlu, Shuning He, B. Ewa Snaar-Jagalska, Erno Vreugdenhil, John H. N. Meerman, Bob van de Water, and Erik H. J. Danen. β_1 integrin inhibition elicits a prometastatic switch through the tgfβ-mir-200-zeb network in e-cadherin-positive triple-negative breast cancer. *Science Signaling*, 7(312):ra15–ra15, 2014.

[34] Alexander Hayn, Tony Fischer, and Claudia Tanja Mierke. Inhomogeneities in 3d collagen matrices impact matrix mechanics and cancer cell migration. *Frontiers in Cell and Developmental Biology*, 8, 2020.

[35] Yifan Wang, Xiaohan Zhang, and Wei Cai. Spherical harmonics method for computing the image stress due to a spherical void. *Journal of the Mechanics and Physics of Solids*, 126:151–167, 5 2019.

[36] Mark A. Wieczorek and Matthias Meschede. Shtools: Tools for working with spherical harmonics. *Geochemistry, Geophysics, Geosystems*, 19:2574–2592, 2018.

[37] Norman Bobroff. Position measurement with a resolution and noise-limited instrument. *Review of Scientific Instruments*, 57(6):1152–1157, 1986.

[38] Steffen Grosser, Jürgen Lippoldt, Linda Oswald, Matthias Merkel, Daniel M. Sussman, Frédéric Renner, Pablo Gottheil, Erik W. Morawetz, Thomas Fuhs, Xiaofan Xie, Steve Pawlizak, Anatol W. Fritsch, Benjamin Wolf, Lars-Christian Horn, Susanne Briest, Bahriye Aktas, M. Lisa Manning, and Josef A. Käs. Cell and nucleus shape as an indicator of tissue fluidity in carcinoma. *Physical Review X*, 11:011033, 2 2021.

[39] Nathalie Q. Balaban, Ulrich S. Schwarz, Daniel Riveline, Polina Goichberg, Gila Tzur, Ilana Sabanay, Diana Mahalu, Sam Safran, Alexander Bershadsky, Lia Addadi, and Benjamin Geiger. Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates. *Nature Cell Biology*, 3:466–472, 5 2001.

[40] Pascal Vallotton, Stephanie L. Gupton, Clare M. Waterman-Storer, and Gaudenz Danuser. Simultaneous mapping of filamentous actin flow and turnover in migrating cells by quantitative fluorescent speckle microscopy. *Proceedings of the National Academy of Sciences*, 101:9660–9665, 6 2004.

[41] Viola Vogel and Michael Sheetz. Local force and geometry sensing regulate cell functions. *Nature Reviews Molecular Cell Biology*, 7:265–275, 4 2006.

[42] Saba Ghassemi, Giovanni Meacci, Shuaimin Liu, Alexander A Gondarenko, Anurag Mathur, Pere Roca-Cusachs, Michael P Sheetz, and James Hone. Cells test substrate rigidity by local contractions on submicrometer pillars. *Proceedings of the National Academy of Sciences of the United States of America*, 109(14):5328—5333, April 2012.

[43] Hedde van Hoorn, Rolf Harkes, Ewa M. Spiesz, Cornelis Storm, Danny van Noort, Benoit Ladoux, and Thomas Schmidt. The nanoscale architecture of force-bearing focal adhesions. *Nano Letters*, 14(8):4257–4262, 2014. PMID: 24998447.

[44] Alberto Elosegui-Artola, Roger Oria, Yunfeng Chen, Anita Kosmalska, Carlos Pérez-González, Natalia Castro, Cheng Zhu, Xavier Trepat, and Pere Roca-Cusachs. Mechanical regulation of a molecular clutch defines force transmission and transduction in response to matrix rigidity. *Nature Cell Biology*, 18:540–548, 5 2016.

[45] Ilka B. Bischofs, Sebastian S. Schmidt, and Ulrich S. Schwarz. Effect of adhesion geometry and rigidity on cellular force distributions. *Phys. Rev. Lett.*, 103:048101, Jul 2009.

[46] Alex Mogilner and Kinneret Keren. The shape of motile cells. *Current Biology*, 19:R762–R771, 9 2009.

[47] Xingbo Yang, Dapeng Bi, Michael Czajkowski, Matthias Merkel, M. Lisa Manning, and M. Cristina Marchetti. Correlating cell shape and cellular stress in motile confluent tissues. *Proceedings of the National Academy of Sciences*, 114:12663–12668, 11 2017.

[48] Wim Pomp, Koen Schakenraad, Hayri E. Balcıoğlu, Hedde van Hoorn, Erik H. J. Danen, Roeland M. H. Merks, Thomas Schmidt, and Luca Giomi. Cytoskeletal anisotropy controls geometry and forces of adherent cells. *Phys. Rev. Lett.*, 121:178101, Oct 2018.

[49] Stacey A. Maskarinec, Christian Franck, David A. Tirrell, and Guruswami Ravichandran. Quantifying cellular traction forces in three dimensions. *Proceedings of the National Academy of Sciences*, 106:22108–22113, 12 2009.

[50] Sung Sik Hur, Yihua Zhao, Yi-Shuan Li, Elliot Botvinick, and Shu Chien. Live cells exert 3-dimensional traction forces on their substrata. *Cellular and Molecular Bioengineering*, 2:425–436, 9 2009.

[51] Valentin L. Popov, Markus Heß, and Emanuel Willert. *Handbook of Contact Mechanics*. Springer Berlin Heidelberg, 2019.

[52] J Lee, M Leonard, T Oliver, A Ishihara, and K Jacobson. Traction forces generated by locomoting keratocytes. *Journal of Cell Biology*, 127(6):1957–1964, 12 1994.

[53] Marion Ghibaudo, Alexandre Saez, Léa Trichet, Alain Xayaphoummine, Julien Browaeys, Pascal Silberzan, Axel Buguin, and Benoît Ladoux. Traction forces and rigidity sensing regulate cell functions. *Soft Matter*, 4:1836–1843, 2008.

[54] Benedikt Sabass, Margaret L. Gardel, Clare M. Waterman, and Ulrich S. Schwarz. High resolution traction force microscopy based on experimental and computational advances. *Biophysical Journal*, 94:207–220, 1 2008.

Figure 1: **Particle shape and signal reconstruction. A.** BSA conjugated with AlexaFluor-488 confirmed adhesion of BSA to the particle's surface. The particle was stained with TexasRed-Cadaverine. **B.** A z-stack of the particle was taken. **C.** Edges were found using the Sobel operator. **D.** Lines were drawn from the center of the particle radially outward. **E.** A Gaussian fit (red) was used to find the peak of the edge line-profile (blue). The same lines were used to find the line-profile of the stain signal (green). The stain signal on the particle's surface was calculated by averaging the signal within 1 μm of the edge (dark green). **F.** Triangulation was used to reconstruct the shape. **G.** The stain (AlexaFluor-488 Phalloidin, green) intensity of MV3 cells was found by drawing lines (blue) from the center of the particle (red). **H.** The signal on the particles' surface was averaged and projected onto the sphere-like shape.

Figure 2: **Tumoroid with particles in collagen gel, 24 h after tumoroid injection. A.** MV3 tumoroid with cells migrating from the injection site (marked with 'x'). The dashed line indicates the tumoroid core. F-actin was labelled with AlexaFluor-488 Phalloidin and nuclei with Hoechst. **B.** Close-up of a particle at the edge of a tumoroid. All scale bars are 20 μm. **C.** Reflection microscopy image of the collagen network.

Figure 3: **Cell segmentation in tumoroids allows for single-cell traction-analysis. A.** Slices of a z-stack of a MV3 tumoroid in a collagen gel. The particle (red) was located at the edge of the tumoroid. The cell shape was obtained by staining F-actin (green). Cells were identified by their nuclei (blue). Here, three cells were found touching the particle. The obtained cell-segments are displayed in white. **B.** From top to bottom: the deformation, normal traction, stain signal and cell index map. The cell indices correspond to the labels in (A). Unexposed regions of the particle were identified as background (bkg). **C.** Cells 2 and 3 significantly indent the particle, hence applying a negative normal traction beyond our traction resolution (grey area) ($p < 0.001$). **D.** The particle surface that was exposed to cells had an increased amount of stain signal ($p < 0.001$).

Figure 4: **Contact area and the amount of actin correlate with the force cells apply. A.** Weak correlation between mean normal traction in the cell-particle contact area and F-actin signal, which is displayed as a signal-to-noise value, for two independent experimental conditions (blue and red). **B.** The net force and contact area are strongly correlated through a power law relation. **C.** A power law relation was found between normal traction and net force. **D.** The applied normal traction appeared to be independent of the cell shape. Correlation coefficients are displayed by ρ. Power law fits are displayed by solid lines in (B-C). The power law exponent of $y = kx^m$ including the standard error of the fit is displayed in the corner.

Figure S1: **Collagen fibers apply stresses on the microparticles. A.** TexasRed-Cadaverine labelled microparticle (left) and reflection microscopy of the surrounding collagen fibers (right). The collagen fibers significantly indent the particle (yellow arrows). These deformations have the consequence that a stress-free boundary condition can not be applied for calculation of the traction field. **B.** Quantification of the microparticle shape (left) and collagen signal on the microparticle surface (right). The indentations (yellow outline) coincide with regions of high collagen signal.

Figure S2: **Edge localization accuracy of a particle suspended in a collagen gel. A.** Deformation field. Root-mean-square surface deformation is 140 nm, its standard deviation is 138 nm. **B.** Normal traction field. Root-mean-square normal traction is 17.6 Pa, its standard deviation is 17.5 Pa. **C.** Shear traction field. Root-mean-square shear traction is 4.6 Pa, its standard deviation is 2.3 Pa.

Figure S3: **Segmentation method. A.** Cell were identified based on their nuclei. **B.** A Gaussian filter was applied and nuclei were detected using an adaptive threshold. **C.** The nuclei centers function as seeds for the segmentation. Note that this is a single slice of a 3d image cube. Hence, not all seeds are visible in this slice. Additionally, the particle volume was excluded. **D.** Cell boundaries were identified by labelling of F-actin. **E.** Salt-and-pepper noise was reduced by applying an adaptive median filter. **F.** Using the Sobel operator, edges were found. A Gaussian filter was applied to merge edges originated from the same line segment. Subsequently, the image was superimposed with the seeds (yellow) and the particle (red). **G.** Overlay of the detected cell edges (white) and the composited image. All scale bars are 20 µm.

Figure S4: **Trade-off between spatial resolution and computational time.** With increased harmonics degree l, the relative error becomes smaller (**A**) while the time required per computation goes up (**B**). Power law fit is displayed in red.

Figure S5: **The applied shear traction appeared to be independent of the cell shape.** Correlation coefficients are displayed by ρ.