Title: Memory consolidation during rest forms shortcuts in a cognitive map

Authors: Cal M. Shearer[1,2,†], Annalise B. Rawson[2,3], Helen C. Barron*[2,3], Jill X. O'Reilly*[1]

1. Department of Experimental Psychology, University of Oxford, Oxford, United Kingdom; **2.** Medical Research Council
Brain Network Dynamics Unit, Nuffield Department for Clinical Neurosciences, University of Oxford, Oxford, United
Kingdom. **3.** Wellcome Centre of Integrative Neuroimaging, University of Oxford, FMRIB, John Radcliffe Hospital, Oxford,
United Kingdom

† Corresponding author: Cal Shearer (cal.shearer@psy.ox.ac.uk)

*These authors contributed equally

Abstract

Rest and sleep not only strengthen existing memories but also reorganise memories to generate new
knowledge that extends beyond direct experience. However, it remains unclear both *how* memories
are reorganised and the effect of this reorganisation on behaviour. Here, we designed a novel protocol
to casually manipulate memory consolidation during rest using awake, contextual targeted memory
reactivation (TMR). We found that promoting memory consolidation during rest qualitatively
reorganises memories by *forming 'shortcuts' between memories* which have not been experienced
together. These shortcuts in memory extend beyond direct experience to facilitate our ability to make
novel inferences. A series of control tests indicate that inference performance cannot be explained by
quantitative strengthening of the experienced component links but are rather explained by qualitative
changes in the cognitive map which involve formation of new shortcuts. Interestingly, we show that
representing a shortcut may come with limitations, as shortcuts cannot be readily updated in response
to rapid changes in the environment. Together, these findings reveal how memories are reorganised
during awake rest to construct a cognitive map of our environment, while highlighting the constraints
set by a trade-off between efficient and flexible behaviour.

Introduction

To understand our environment, we construct a cognitive map[1] forged from associations we have experienced. However, the cognitive map is more than the sum of its parts. As a simple example, if an observer has seen the relationship between A->B and B->C, they can infer the unobserved relationship A->C[2]. It remains unclear when and how these inferential relationships are built to allow us to make predictions that go beyond direct experience. For example, inference could be supported by a chain of learned associations (A->B, B->C), or by formation of a new 'shortcut' (A->C). The formation of a new shortcut (A->C) would represent a *qualitative* change in the organisation of memories, which is an essential pre-requisite for the organization of knowledge into a structured cognitive map.

The process of constructing a cognitive map is thought to be facilitated by offline periods of rest/sleep[3–10]. During these periods, memories for previous experiences are reactivated or 'replayed'[11–13]. Replay involves temporally structured spiking activity that recapitulates previous waking experience[14–16] to facilitate memory consolidation[11–13,17] and subsequent decision-making[18]. In addition, growing evidence suggests replay may play a particular role in extending the cognitive map beyond previous waking experience to anticipate upcoming events[19,20], restructure knowledge[21], or even "join-the-dots" between spatial trajectories or events that were not directly experienced together[22–24]. However, the way in which memories are reorganised and the consequences of this reorganisation for behaviour remain unknown. Here, we test the hypothesis that offline periods of rest *qualitatively* reorganise memories, rather than *quantitatively* changing the strength of associations that were actually experienced. Further, we predict that this qualitative re-organisation allows for the creation of new inferred links (or shortcuts) that are independent of experienced associations, thus forming a cognitive map that is more than the sum of its parts.

In order to causally manipulate which memories were replayed during offline periods of rest, we developed a novel protocol using awake contextual targeted memory reactivation (TMR). TMR involves playing sensory cues during offline periods of rest or sleep that have previously been linked to a specific memory during prior learning. For directly experienced associations, TMR has been shown to enhance behavioural readouts of memory for cued compared to uncued associations[25–29]. Crucially, TMR has also been shown to bias the content of replay towards cued associations[27]. Given the proposed role of replay in reorganising memory to extend the cognitive map beyond direct experience, and evidence to suggest that TMR can bias the content of replay, we predicted that TMR might provide a tool to investigate how memories are reorganised during periods of rest and the consequences for behaviour. Thus, we sought to test the hypotheses that TMR improves inferential choice and that this improvement is driven by the formation of new inferred links (or 'shortcuts') between cues that have not been experienced together[22,30].

To test our hypothesis, we designed a multi-stage inference task adapted from a protocol previously implemented in humans and mice[22]. In addition, we implemented a novel TMR protocol involving awake, contextual auditory TMR. Consistent with our predictions, we demonstrate that biasing memory reactivation during rest improves the ability to infer novel relationships between sensory cues. Furthermore, using behavioural and physiological (eye-tracking) data, we show that this improvement in inference is driven by the formation of novel shortcuts between indirectly linked cues. As a result of these shortcuts, participants are less reliant on memory of intermediary associations (A->B and B->C) since they can instead use the direct, yet unobserved, shortcut formed during rest (A->C) to inform their decisions. However, although beneficial for inference, we show that building cognitive shortcuts may come with limitations: when the component associations are

modified (e.g. B->C changes to B->C*), this change is not immediately transferred to the shortcut,
leaving the old shortcut (A->C) to compete with the updated chain of associations (A->B, B->C*)

Together our findings demonstrate the effectiveness of using awake contextual TMR as a tool for
examining the role of offline memory consolidation in shaping behaviour. We show that memory
consolidation enhances inferential choice through the formation of shortcuts between indirect
associations. However, these shortcuts cannot be rapidly updated. We thus demonstrate that
memory consolidation during rest constructs a cognitive map that is more than the sum of its parts,
allowing for efficient behaviour that is limited in flexibility.

Results

Task design and learning performance
To investigate the relationship between periods of rest and inference using TMR, we designed an
inference task (Figure 1A-B). The inference task was adapted from a previous inference protocol
implemented in humans and mice[22] that leveraged a sensory preconditioning paradigm[2]. The
inference task included three stages.

In the first stage, participants learnt to associate pairs of auditory and visual cues ('associative
learning'), with a many-to-one mapping from auditory to visual cues. The presentation of task cues
was blocked into two groups, distinguished by different contextual background music that played
throughout learning (either café or jungle music). One of these contextual background tracks was
later played during rest to bias memory consolidation to this group of cues (TMR manipulation). After
training, participants' mean accuracy on the associative learning trials was 96.67% (SD=7.41%), with
no significant difference in the maximum accuracy reached between sensory cues in the two
contexts (café vs jungle; p=0.385) or between the group of cues that were subsequently subject to
TMR and the group of cues that were not (TMR group vs no-TMR group; p=0.185) (Figure 1E).

In the second stage of the inference task, participants learnt to associate the visual cues with either a
rewarding outcome (represented by a pound coin) or a neutral outcome (represented by a wooden
coin) ('conditioning'). As in the associative learning phase, contextual background music was played
throughout learning. After training, participants achieved a mean accuracy of 99.50% on the
conditioning trials (SD=2.50%) (Figure 1F).

Importantly, auditory cues were never paired with outcome cues, providing an opportunity to assess
evidence for an inferred relationship between these indirectly related cues. Accordingly, in the final
stage participants performed an 'inference test'. In the inference test, we presented auditory cues in
isolation, without visual cues or outcome cues, and without the contextual background music. In
response to these isolated auditory cues, we measured evidence for inference by asking participants
which outcome was associated with each auditory cue, despite the fact that they had never directly
experienced this pairing.

TMR manipulation
In addition to the inference task, we included a TMR manipulation to study the consequences of
biasing memory consolidation. Participants underwent the TMR manipulation during a period of rest
(60 minutes), after completing the associative learning and conditioning phases of the task, but
before performing the inference test. Throughout this rest period, we played the contextual
background track associated with cues in one of the two groups (i.e., either café music or jungle
music, fully counterbalanced across participants) whilst participants completed a relaxing task
unrelated to the learning task (namely, they completed a jigsaw puzzle). If TMR facilitates the
construction of a cognitive map that is more than the sum of its parts, then this TMR manipulation
should facilitate inference between cues in the TMR group.

TMR improves inference performance
After the period of rest, participants performed the inference test (Figure 2A-B). Here, we presented
the auditory cues in isolation and tested participants' ability to infer a relationship with the
corresponding outcome cue (rewarding or neutral). To assess the effect of TMR on inference, we
compared participants' accuracy in response to auditory cues in the TMR and no TMR groups.

Figure 1. Inference Task Design and Learning Performance.

(A) Three-stage inference task adapted from a previous protocol[22]. In the learning phase, participants learnt to associate auditory cues with visual cues ('Associative Learning'), with a many-to-one mapping. Participants then learnt to associate visual cues with an outcome (rewarded – pound coin; neutral – wood coin) ('Conditioning'). During the 'Inference Test', the

Participants were better at inferring the correct outcome for auditory cues in the TMR group
(p=0.011) (Figure 2C). The inference task was carried out both in isolation, and in the presence of

two distractor tasks (a one-back task and a semantic judgement task, see Methods). The distractor
tasks were designed to increase the load on working memory or general cognitive resources
respectively, to modulate task difficulty and avoid ceiling effects when inference was performed in
isolation. In fact, participants were better at inferring the correct outcome for auditory cues in the
TMR group under all three conditions (p=0.015, no distractor task; p=0.031, memory task; p= 0.030,
non-memory task), with no significant difference in the TMR effect between conditions (p=0.130, no
distractor task vs memory task; p=0.372, no distractor task vs non-memory task) (Fig. S1). Hence,
hereafter results are pooled across the three distractor conditions. Overall, participants were better
at inferring the correct outcome for auditory cues in the TMR group (p=0.011) (Figure 2C). This result
demonstrates that biasing memory consolidation during periods of rest with awake, contextual TMR
facilitates inferential choice.

TMR creates a shortcut in memory between indirectly linked cues
Next, we asked *how* TMR influences the organisation of learned information to facilitate inferential
choice. Specifically, we asked: do participants build a shortcut (A->C) that is separate from the
original chain of learned associations (A->B, B->C), or do they instead strengthen the components of
the learned associations (A->B, B->C)? We reasoned that if TMR facilitates inference by building a
new short-cut between the auditory and outcome cues (A->C), the resulting cognitive map may be
less amenable to flexible updating when associations between cues change.

Specifically, if the visual-outcome association B->C changes to B->C*, the correct chain of
associations to support inference is now A->B and B->C*. If participants perform inference using the
chain of learned associations (A->B, B->C), and each of the learned associations are merely
strengthened during TMR, then it should be possible to rapidly update behaviour to account for the
changes from B->C to B->C*. However, if an old shortcut (A->C) is represented that is not rapidly
updated (to A->C*), behaviour may not account for the changes from B->C to B->C*.

To test this hypothesis, after the first inference test, we manipulated the association B->C (visual-
outcome), such that the outcome associated with a subset of visual cues was reversed ('value flip'
manipulation, B->C*). Participants then performed the inference test once more, to test their ability
to flexibly update the inferred value of auditory cues (from A->C to A->C*) (Figure 3A). Critically, if
TMR creates a shortcut (A->C) between auditory and outcome cues that is separate to the chain of
learned auditory-visual-outcome associations (A->B, B->C), then an update in the visual-outcome

leg (B->C*) should not transfer to the auditory-outcome (A->C) short-cut. Therefore, to make correct
inferences, changes to the visual-outcome (B->C*) mapping would require participants to draw on
the complete chain of learned associations (A->B, B->C*) for cues in both the TMR and no TMR
groups, negating the benefit of TMR. In addition, representation of an auditory-outcome (A->C)
shortcut should compete with flexible updating in response to changes to the visual-outcome (B->C*)
mapping.

We observed evidence for both of these predicted effects. First, for visual-outcome mappings (B->C)
where there was no change in the association ('non-flipped' associations), participants were better at
inferring the correct outcomes for auditory cues in the TMR compared to the no TMR group
(p=0.001). However, for flipped associations (B->C*), there was no significant difference between

stimuli in the TMR and no-TMR groups (p=0.276) (Figure 3B-C). These results suggest that the benefit
of TMR is negated by the value-flip manipulation. This was indeed the case, as the effect of TMR was
significantly greater for non-flipped stimuli (p=0.024) (Figure 3D). This is consistent with the proposal
that, after rapid updating in the visual-outcome (B->C*) mapping, participants had to perform
inferential choice by drawing on the complete chain of learned associations (A->B, B->C*), rather
than the shortcut (A->C), thus negating the benefit of TMR.

Next, we asked whether representing a shortcut between the auditory-outcome (A->C) cues acts in
competition with the modified chain of learned associations (A->B, B->C*). Specifically, during the
inference test performed after the value flip manipulation, we asked whether forming a shortcut (A-
>C) biases participants to look at the previously associated (but now incorrect) outcome (C) in the
inference test. Thus, we used eye-tracking data acquired during the second inference test to ask
whether, after the value flip manipulation, the shortcut (A->C) competes with the modified chain of
associations (A->B, B->C*) (Figure 3E). To do this, we calculated the percentage of trials where
participants looked at the incorrect outcome. For 'flipped' cues, the incorrect outcome corresponded
to the outcome cue (C) that was previously correct prior to the value flip manipulation.

We hypothesised that for auditory cues where the visual-outcome association was flipped (B->C*),
participants may still retain a shortcut (A->C) from the auditory cue to the (now incorrect) outcome
for cues in the TMR group (Figure 3E) and would therefore be more likely to look at C during the
decision process. This effect should be lower, or absent, for cues in the no-TMR group, as for these
cues the shortcut (A->C) was weaker or not formed, relative to the TMR cues. This was indeed the
case. For flipped cues, participants were significantly more likely to look at the incorrect outcome cue
(C) in the TMR group compared to the no TMR group (p<0.001) (Figure 3D). This suggests that when
a shortcut (A->C) forms due to TMR, knowledge of this shortcut competes with knowledge of
relevant updated associations. This competitive effect supports the view that the shortcut (A->C) is
separate from the chain of learned associations (A->B, B->C). Consequently, TMR may be considered
to support efficient behaviour by facilitating inference through the formation of a shortcut between
auditory and outcome cues even when these cues had not been directly experienced together.
However, building a shortcut in the underlying cognitive map appears to have limitations – the
shortcut cannot be flexibly updated in response to environmental change.

In a final set of analyses, we again used eye-tracking data to ask whether TMR facilitates inference by
creating a shortcut in memory between the auditory and outcome cues. If a shortcut (A->C) is
represented, we would expect participants to be less likely to use the intermediary visual cues (B) to
help make inferential choices for cues in the TMR group. To test this, we analysed eye-tracking data
acquired during a third inference test ('indirect') and during a memory test for learned associations
('direct') ('indirect/direct' test, Figure 2A, 4A-B). During the first half of this testing phase,
participants again had to decide whether the auditory cues were associated with the rewarding or
neutral outcome (inference test, 'indirect'). Replicating our previous findings, we found that
participants were again significantly more accurate at inferring the correct relationships for auditory
cues in the TMR group compared to the no TMR group (p=0.013) (Figure 4E). However, unlike in the
previous inference tests, all of the intermediary visual cues were now presented on the screen
(Figure 4B). We predicted that if participants had formed a shortcut between auditory and outcome
cues, they would be less likely to look at the visual cues to help them infer a value for each auditory
cue. On correct trials, participants spent less time looking at the visual cues for auditory cues in the
TMR group compared to the no TMR group (p=0.012) (Figure 4C). Therefore, for cues in the TMR
group, there was a reduced need to look at the intermediary visual cue to help make a

Figure 3. TMR Creates a Shortcut Which Cannot Be Readily Updated.

(A) Schematic illustrating the value flipping manipulation. After the learning and rest phases, the outcome associated with half of the visual cues was flipped ('Value Flip'). During Inference Test 2 (highlighted in red), we tested to see if participants could correctly infer the new outcome that was indirectly associated with the auditory cue.

(B) Inference test performance after the value flip. Bars: mean percentage accuracy for No TMR (red; left two bars) and TMR (blue; right two bars) groups split by whether the visual cue-outcome association was not flipped (plain; left bar of each group) or flipped (hatched; right bar of each group); black ticks: ± SEM; each data point: mean accuracy for one participant.

(C) Difference in TMR effect on inference test accuracy between auditory cues associated with not flipped and flipped. Left: raw data points for Not Flipped cues (left) and Flipped cues (right); each data point: mean effect of TMR (TMR accuracy – No TMR accuracy) for a given participant; black dots, mean; black ticks ± SEM. Right: difference in means between Not Flipped cues and Flipped cues shown using bootstrap-coupled estimation (DABEST) plots as in Figure 2C: black dot, mean; black

261 correct inference. Consistent with our previous findings, this result suggests that TMR facilitates the
262 formation of a memory shortcut between the indirectly linked auditory and outcome cues.

The effect of TMR on inference is not explained by improved memory for directly learnt associations

Finally, we asked whether awake contextual TMR also facilitates memory accuracy for directly
associated cues. Previous work has shown that TMR protocols that use punctate auditory cues
improve memory for directly learnt associations[31]. However, since our TMR protocol uses contextual
cues, we wanted to investigate whether this protocol would have similar or different effects on
memory. During the indirect/direct test (Figure 4A-B), participants were also given a memory test for
learned associations between auditory and visual cues ('direct associations'). By contrast to
performance on the inferred relationships between auditory and outcome cues ('indirect
associations') reported above (Figure 4E), when participants were tested on direct associations, they
showed no significant difference in accuracy for auditory-visual (A->B) cue mappings in the TMR
group compared to the no TMR group (p=0.478) (Figure 4F).

Since participants were trained to ceiling on the direct associations, it is possible that we did not see
an effect of TMR due to ceiling performance. To investigate this potential confound, we performed a
median split based on combined performance for TMR and no TMR groups, and applied this to both
the indirect association (inference) test (in which we saw an effect of TMR) and direct association
test (in which we did not see an effect of TMR). For direct associations, even for participants who
were not at ceiling, we continued to observe no significant effect of TMR (p=0.451, top 50%;
p=0.442, bottom 50%), suggesting that the lack of improvement under TMR was probably not due to
a ceiling effect. In contrast, for indirect associations (where we had observed an overall effect of
TMR), the TMR effect remained significant for both the bottom half (p=0.029) and top half (p=0.036)
of participants. Thus, even for participants who performed the inference task close to ceiling
performance, we continued to observe a significant effect of TMR. Together these findings suggest
that the reported difference in effect of TMR on indirect and directly learned associations is likely not
due to difference in task difficulty or due to a ceiling effect, but instead indicates that contextual TMR
applied during rest prioritises formation of shortcuts that support inference, rather than
strengthening learned memory content.

Together, these results suggest that while contextual TMR improves inferential choice across
indirectly associated cues, this improvement cannot be explained by improved performance on
directly learned associations. Therefore, the benefits of contextual TMR on inferential choice must be
explained by a mechanism that goes beyond merely strengthening learned associations. This
provides further evidence to suggest that periods of rest facilitate inference by restructuring
knowledge and creating shortcuts between items in memory that extend beyond direct experience.

Figure 4. TMR Creates a Shortcut Which Does Not Rely on Directly Learnt Associations

(A) Structure of the testing phase of the task. Data presented in (C-F) is from the 'Indirect/Direct Test' highlighted in red. In the indirect/direct test, participants were tested on both the indirect (auditory – outcome) and direct (auditory – visual) associations for each auditory cue.

(B) Schematic: example indirect trial (schematic) where auditory cues were now presented while the visual cues were on screen. Participants were required to infer the outcome cue in response to the auditory cue. Direct trials followed the same structure except, after hearing the auditory cue, participants had to choose the associated visual cue rather than the associated outcome.

(C) Gaze location from eye tracking data acquired during indirect trials in the indirect/direct test. Top: example gaze trajectories from a single indirect trial for one participant in response to an auditory cue from the No TMR group (red; left) and an auditory cue from the TMR group (blue; right). Darker colours indicate later time in the trial. Bottom left: raw data

points for No TMR group (red; left) and TMR group (blue; right); each data point: mean z-scored time spent looking at visual
cues during inference (correct trials only); black dots, mean; black ticks ± SEM. Bottom right: difference in means between
No TMR and TMR groups shown using bootstrap-coupled estimation (DABEST) plots as in Figure 2C: black dots, mean; black
ticks, 90% confidence interval; filled-curve, sampling-error distribution. Participants spent significantly less time looking at
the visual cues in response to auditory cues in the TMR group (p=0.012).
(D-E) The effect of TMR on indirect associations (D) and direct associations (E). Left of each panel: raw data points for No
TMR group (red; left) and TMR group (blue; right); each data point: mean accuracy for a given participant; black dots,
mean; black ticks ± SEM. Right of each panel: difference in means between No TMR and TMR groups shown using
bootstrap-coupled estimation (DABEST) plots as in Figure 2C: black dot, mean; black ticks, 90% confidence interval; filled-
curve, sampling-error distribution. There was a significant effect of TMR on accuracy for indirect associations (p=0.013) but
not for direct associations (p=0.478).

Discussion

Previous studies have shown the benefit of rest/sleep for inference[5–9,32,33]. However, the nature of the changes that occur during rest/sleep to facilitate inference remain unclear. In this study, the central question we ask is: _How does memory reactivation during rest reorganise memory to guide adaptive behaviour?_ We sought to tease apart two ways in which rest/sleep could affect memories: an improvement in behavioural performance on an inference task could be supported by either a quantitative strengthening of learned, component associations (A->B and B->C), or a qualitative change, involving the formation of a new, 'shortcut' link (A->C). The latter, qualitative change would reflect an essential pre-requisite for the organization of knowledge into a structured cognitive map that is more than the sum of its parts.

Using a multi-stage inference task with awake auditory contextual TMR, we show that memory reactivation during rest benefits inference by building a new shortcut (e.g. A->C), with no effect on directly trained associations (e.g. A->B, B->C). Thus, promoting memory consolidation during rest facilitates the formation of novel shortcuts in the cognitive map. This may explain why periods of rest and sleep facilitate behavioural readouts of inference, but also other cognitive processes such as insight, generalisation and abstraction[5–10]. Interestingly, both our behavioural and eye-tracking data show that these shortcuts cannot be readily updated in response to subsequent changes in the environment. Rather, shortcuts (e.g. A->C) may limit behavioural flexibility when optimal behaviour requires participants to revert to a chain of learned associations (i.e. A->B, B->C). Together, our results reveal how memory reactivation during rest plays a causal role in _restructuring_ memories, while highlighting how the organisation of knowledge sets an important trade-off between efficient and flexible behaviour.

A candidate neural mechanism that may explain our results is replay. During an inferential learning task in mice, similar to the one employed here, replay was shown to go beyond direct experience by "joining-the-dots" between indirectly linked cues[22]. During awake sharp-wave ripples (SWRs), hippocampal spiking activity represented the inferred relationship (A->C) in the absence of the intermediary cue (B). However, the behavioural consequences of this form of replay were not investigated. Here in humans, we use a TMR manipulation to bias the content of replay[14]. Unlike previous TMR protocols, we used awake, contextual TMR to demonstrate that biasing memory consolidation during rest leads participants to form a shortcut between indirectly related cues. Furthermore, this shortcut reduces the need to rely on memory for the intermediary cue. Together these findings suggest that our TMR manipulation facilitates replay of inferred relationships between cues in the TMR group, promoting the formation of a shortcut in the underlying cognitive map. However, since this study did not directly measure replay, future investigations will need to validate this interpretation using neuronal recordings.

While our ability to make novel inferences can clearly support efficient behaviour, representing inferred relationships as a shortcut (A->C) rather than a chain of learned associations (A->B, B->C) may limit behavioural flexibility. Our data demonstrates the limitation of building a short-cut in two key analyses. These two analyses are applied to data acquired after the value flip manipulation, where half of the visual-outcome mappings are 'flipped' (B->C becomes B->C*). First, we show that the TMR benefit on performing inference is no longer observed for the flipped cues. Second, for the flipped cues, eye-tracking data shows that participants spend more trials looking at the previously correct, but now incorrect outcome (C*) for TMR compared to no-TMR cues. This suggests that the outdated shortcut (A->C) may compete with the updated chain of associations (A->B, B->C*). As well as supporting the proposal that the shortcut (A->C) is indeed separate from the chain of associations (A->B->C/C*). These data demonstrate that shortcuts in memory (e.g. A->C) cannot be readily

updated in response to rapid changes in the environment and may limit behavioural flexibility when
optimal behaviour requires participants to revert to a chain of learned associations (i.e. A->B, B->C*).
We speculate that if a further rest period was included after the change in value of outcome (C), a
new updated shortcut (A->C*) would have been formed and used. Indeed, there is evidence to
suggest that rest and sleep may play a role in integrating new memories into existing knowledge[34–37].
Our findings complement this work by suggesting that adaptive re-organisation of the cognitive map
cannot always occur 'on the fly' and may instead rely on offline periods of rest or sleep. Our study
therefore provides insight into how new information is incorporated into a cognitive map, to guide
adaptive behaviour.

The above findings were made possible by our novel contextual auditory TMR protocol. The majority
of previous TMR studies have used either punctate auditory cues[31] or contextual odour cues[38]. Using
punctate auditory cues may have several drawbacks, including the need to precisely time the cue to
specific phases of sleep oscillations to see the best effects[39]. Furthermore, since punctate cues are
associated with specific stimuli, they provide a limited tool to investigate how links across multiple
stimuli are consolidated. The contextual TMR protocol presented here provides a unique opportunity
to investigate the effect of memory reactivation across an entire map of associations. We
demonstrate how memory reactivation during periods of rest influences the organisation of
memories within the cognitive map, where memory reactivation plays a causal role in forming novel
links that go beyond direct experience.

An additional benefit of the TMR protocol used here is that it is very simple to implement. Using a
contextual background track removes the need to present multiple different cues across the rest
period. Furthermore, auditory cues are simpler to deliver than the contextual odour cues that have
been used in previous contextual TMR paradigms. Finally, delivering the TMR during awake rest, as
opposed to sleep, removes a significant time investment and the need to ensure for accurate sleep
staging. Given the increasing interest in using TMR for clinical benefit in a number of
neuropsychiatric conditions[40], the simplicity of the paradigm carries important translational
potential.

Previous work, which largely focus on the effect of punctate TMR cues applied to directly
experienced associations, show variable results when using TMR during periods of awake
rest[26,31,38,41,42], which may in part be explained by variation in task difficulty and performance
accuracy. Indeed, non-invasive measures of memory reactivation suggest that replay of directly
learned information prioritises weakly learned information[43]. Consistent with this view, TMR
facilitates direct associations by reducing forgetting[31,44,45]. Therefore, by prioritising reactivation of
weaker memories, TMR may help protect memories that are most vulnerable to forgetting[43]. In our
data we report a specific benefit of TMR for indirect associations, with no effect found on direct
associations. In our protocol we included only a short time interval (1 hour) between learning and
test, which may have reduced the scope for forgetting. It is possible that if we had used a longer time
interval, we would have seen a benefit of TMR for directly learned associations. Notwithstanding this
point, the lack of an effect on direct associations implies that the observed restructuring of
knowledge does not rely on strengthening of directly learned associations, or on rescuing the direct
associations from forgetting.

We also note that the differential effect of TMR on indirect and direct associations cannot easily be
explained by differences in performance accuracy or ceiling effects. Unsurprisingly, participants
showed higher performance accuracy on direct compared to indirect associations. However, when
we performed a median split analysis using the combined performance for TMR and no TMR cues,
we show that both high and low performing participants continue to show a significant TMR effect

for indirect but not direct associations. Therefore, participants who showed performance accuracy
close to ceiling continued to show a significant TMR effect on indirect associations, while participants
where performance was not at ceiling continued to show no significant effect of TMR on direct
associations. Together these findings suggest that the differential effect of TMR on direct and
indirect associations cannot be explained by differences in performance accuracy. Nevertheless,
indirectly trained associations may share physiological properties with weaker memories, rendering
them more susceptible to TMR.

Overall, our results demonstrate that knowledge is subject to qualitative restructuring during offline
periods of rest/sleep. During rest, stimuli that are reactivated (due to contextual TMR) are organized
into cognitive maps through the creation of novel shortcuts between indirectly linked cues. These
shortcuts are distinct from the original chain of learned associations and the behavioural benefit of
forming a shortcut cannot be explained by quantitatively strengthening the component learned
associations (since there was no benefit of TMR on these learned associations even in participants
whose performance was well below ceiling). Furthermore, when the component learned
associations are rapidly modified in response to changes in the environment, shortcuts are not
updated. Indeed, an outdated shortcut can conflict or compete with an updated chain of associations
to control decision making, illustrating that the two sets of associations are separate. Together our
findings provide new insight into how integrated cognitive maps are formed during periods of rest.
Moreover, we demonstrate the effectiveness of using awake contextual TMR as a tool to reveal how
the resting brain re-organises memory to shape adaptive behaviour.

Methods

Participants

A total of 32 healthy participants were recruited for this study. 7 participants were excluded from analysis as they failed to reach the performance criterion during the initial associative learning phase (criterion: >85% performance accuracy for each contextual group of cues). The remaining 25 participants were included in the analysis (mean age of 24.72 ± 6.35 years, range 19-40, 4 males). All participants had normal or corrected-to-normal vision. All experiments were approved by the University of Oxford ethics committee (reference number: R43593/RE013).

Experimental set-up

Cues were presented on a 24-inch screen with a spatial resolution of 1920 x 1080 pixels (width x height), a background luminance of 0.5 (grey), and a refresh rate of 100 Hz. The approximate distance of participants from the screen was 64cm, which meant that one degree of visual angle corresponded to 40 pixels on the screen. Stimulus presentation was controlled using Psychophysics Toolbox-3[46] in MATLAB (version R2015b). Eye movements were recorded in the testing phase of the experiment (after the rest period) using an eye-tracking camera (EyeLink®, SR Research) tracking both eyes at a rate of 1kHz.

Inference task - overview

Participants performed an inference task (Figure 1A-B). In the first stage of the task ('associative learning'), participants learnt to associate punctate auditory cues with visual cues. The punctate auditory cues were naturalistic sounds from the BBC sound library (https://sound-effects.bbcrewind.co.uk/) and the visual cues were unsymbolic shapes[47]. Each visual cue was paired with three auditory cues. In the second stage of the task ('conditioning'), these visual cues were then associated with either a pound coin (rewarded outcome) or a wood coin (neutral outcome, of no value). The cues were split into two sets, grouped according to the contextual background music that played throughout learning. For one of the two sets of cues, participants heard jungle music in the background and the punctate auditory cues were sounds you would hear in a jungle (e.g., animal noises). For the other group, participants heard café music in the background and the punctate auditory cues were sounds you would hear in a café (e.g., a cash register). In total, 12 auditory cues and 4 visual cues were included in each group and the relationship between the conditions (i.e., which context was learnt first, and which context was subject to TMR) was fully counterbalanced across participants. Since the visual cues had no direct meaning, they were randomly assigned to each group across participants.

Behavioural protocol

The associative learning stage was designed to allow participants to learn associations between auditory and visual cues. This stage of the task included passive exposure and testing trials (Figure 1C). In the exposure phase, on each trial a punctate auditory cue was presented for 1.5s, immediately followed by presentation of a visual cue for 1.5s, followed by an inter-trial interval (ITI) of 2s. Each mini-block of exposure consisted of 12 trials (one presentation of each auditory-visual pair). Each mini-block of exposure was followed by a testing mini-block. On each trial in the testing mini-block participants heard an auditory cue for 1.5s before being presented with 3 possible visual cues. Participants were then required to select the visual cue that was paired with the auditory cue they just heard, responding within 5s using the keys 'j', 'k', and 'l'. Once a choice had been selected, a black rectangle would highlight this choice for 0.5s, followed by an ITI of 2s. At the end of each testing mini-block, participants were given feedback on their average performance. Each block in the associative learning phase was made up of one exposure mini-block followed by one testing mini-block and consisted only of cues from one of the two groups of cues, with the relevant contextual

background music playing throughout the block. Blocks alternated between the two contexts with a total of 8 blocks per context. The relationship between the conditions (i.e., which context was learnt first, and which context was subject to TMR) was fully counterbalanced across participants.

The conditioning stage was designed to allow participants to learn associations between visual and outcome cues. The blocks followed the same structure as the associative learning stage, including both passive exposure and testing trials (Figure 1D). In each trial of the exposure phase, a visual cue was first presented for 1.5s, immediately followed by the associated outcome for 1.5s, followed by an ITI of 2s. Each exposure mini-block consisted of 4 trials (one presentation of each visual cue), which was followed by a testing mini-block. In each trial of the testing mini-block, a visual cue was presented together with the two outcome cues for 1s. After this 1s, the visual cue would disappear but the outcome cues would remain on the screen and the participants had a further 3s to make their choice (total of 4s response period). Participants had to select which outcome they thought was associated with the visual cue using the keys 'j' and 'l'. Once a choice had been selected, a black rectangle would highlight their choice for 0.5s, followed by an ITI of 2s. Participants were given feedback on their average performance at the end of each mini-block of 4 trials. Each block in the conditioning phase was made up of one exposure mini-block followed by one testing mini-block and consisted only of cues from one of the two groups of cues, with the relevant contextual background music playing throughout the block. Blocks alternated between each context with a total of 2 blocks per context. The relationship between the conditions (i.e., which context was learnt first, and which context was subject to TMR) was fully counterbalanced across participants.

The TMR manipulation was performed after the conditioning stage. During TMR, participants were instructed to rest for up to 60 minutes, during which participants were asked to alternate between resting with their eyes closed for 10 minutes and doing a jigsaw puzzle for 20 minutes. In total, they spent 20mins with their eyes closed and 40mins doing a jigsaw puzzle. During this rest period, the contextual soundtrack associated with the TMR group was played throughout. The contextual soundtrack used for TMR was fully counterbalanced across participants. To encourage participants to attend to the soundtrack, the soundtrack was turned on and off for periods of between 9 and 16s, alternating with periods of silence. The length of each on and off period was selected from a random distribution.

After the TMR manipulation, participants were required to perform an inference test. During the inference test, participants were asked to infer the relevant outcome in response to each auditory cue.

Before beginning the first inference test, participants practised performing two tasks together using only two auditory cues, not taken from the task, and a 1-back task using digits rather than shapes. The practice consisted of up to 4 blocks with 24 trials per block. The timings of these practice trials matched the timings of the main inference test described below.

We designed three different versions of the inference test which included distractor tasks designed to increase the load on visual working memory or general cognitive resources (Figure S1A). For all versions, participants were presented with an auditory cue and had to choose which of the two outcomes they thought was associated with the auditory cue (Figure 2B). The auditory cue was presented together with the two outcome cues for 1.5s. After 1.5s, the auditory cue would stop but the outcome cues would remain on the screen and the participants had a further 3s to make their choice (total of 4.5s response period). Participants had to select which outcome they thought was associated with the auditory cue using the keys 'j' and 'l'. Once a choice had been selected, a black rectangle would highlight this choice for 0.5s, followed by an ITI of 1s. Participants were told they

would receive a bonus reward of up to £5 based on how many cues they correctly inferred. Participants were given feedback on their average performance at the end of each block. Auditory cues from both contextual groups were randomly interleaved in each block and there was no background music during the inference tests.

For 2 of the 6 blocks, the participants also performed a visual memory distractor task (Figure S1A). The purpose of this distractor task was to modulate task difficulty: a visual memory distractor task could impair recall of the intermediate visual cue when participants perform inference. The visual memory distractor task involved using a 1-back task with shapes similar to those seen in the task. On each trial, between presentations of each auditory cue, a shape would appear for 2s, and the participants had to respond based on whether it was the same or different from the previous shape using the keys 'a' and 'd'. The shape would become darker (participant chose 'same') or lighter (participant chose 'different') after the participant had made their choice. The shape remained on screen until the 2s was complete. The shapes would appear in a random orientation which participants were instructed to ignore. Following a pause of 1s, the auditory cue would begin, and the rest of the inference trial would commence as described above.

For another 2 of the 6 blocks, participants performed a non-memory-based distractor task (Figure S1A). The purpose of this distractor task was again to modulate task difficulty, but without the interfering effect of a visual memory load. On each trial, between presentations of each auditory cue, an image would appear for 2s, and the participants had to respond based on whether the image depicted something living (e.g., a hedgehog) or something non-living (e.g., balloons). The image would become darker (participant chose 'living') or lighter (participant chose 'non-living') after the participant had made their choice. The image remained on screen until the 2s was complete. Following a pause of 1s, the auditory cue would begin, and the rest of the inference trial would commence as described above.

For the remaining 2 blocks, there was no distractor task. After the normal ITI of 1s, there would be an additional pause of 3s to ensure trials across all versions of the inference task were the same length. Following this pause, the auditory cue would begin, and the rest of the inference trial commenced as described above. These 2 blocks which did not include distractor tasks always occurred at the end to minimise order effects of participants practising the inference task on its own.

After all 6 blocks of the first inference test, participants redid the conditioning stage of the experiment but with half of the associations between visual and outcome cues flipped ('value flip') (Figure 3A). The protocol for the value flip stage was otherwise the same as for the conditioning stage (Figure 1D) except participants completed 1 block per context (rather than the 2 blocks per context done in the initial conditioning stage). After the value flip, a second inference test ('inference test 2') was conducted using the same procedure as previously described and also included 6 blocks (Figure 2B; Figure S1A).

In the indirect/direct test, we tested both the indirect (auditory–outcome) and direct (auditory-visual) associations paired with each auditory cue (Figure 4B). All visual cues and outcomes were present on the screen. By having all cues on the screen, we could test whether there were differences in how long participants spent looking at the visual cues during inference in the TMR and no TMR groups. For the first 2 blocks, the auditory cue was presented, and participants had to select one of the two outcomes using the mouse. For the final 2 blocks, the auditory cue was presented, and participants had to select one of the 8 possible visual cues which was paired with the sound. This allowed us to assess performance on the directly learnt associations as well as the indirect auditory-outcome association. For all blocks, the auditory cue was presented for 1.5s. Following this,

567 participants had 30s to select either the outcome cue (first 2 blocks) or the visual cue (last 2 blocks)
568 associated with the auditory cue. Participants made their selection by moving the cursor over the
569 chosen cue and pressing 'spacebar' to confirm. Once a choice had been selected, a black rectangle
570 would highlight this choice for 0.5s, followed by an ITI of 1s. Participants were given feedback on
571 their performance at the end of each block.

Eye-tracking Data Preprocessing

For 2 participants, no usable eye-tracking data was collected. The remaining 23 participants were all
included in eye-tracking analyses. The conversion of the EyeLink® 1000 Edf files was done with the
Edf2Mat Matlab Toolbox designed and developed by Adrian Etter and Marc Biedermann at the
University of Zurich. Gaze position and pupillometry data were smoothed using a 50ms Gaussian
kernel. Blinks were detected from the smoothed pupillometry data and removed from the gaze
position data. For each participant, the eye with the fewest missing samples was selected and used
for all further analyses. The gaze position was epoched around the auditory cue onset for trials in the
second inference test and for indirect trials in the indirect/direct test. For the comparison of flipped
cues, we analysed the percentage of trials where participants looked at the incorrect outcome cue.
To investigate whether participants relied on knowledge of the intermediary visual cue, we analysed
the time spent looking at the visual cues in response to auditory cues in the TMR and no TMR groups
on correct trials of the indirect test. Due to the large variation in the average time spent looking at
the visual cues, we z-scored the fixation lengths across all trials for each participant.

Analysis

Analysis was conducted in MATLAB (version R2022a) and in Python v.3.6
(https://www.python.org/downloads/release/python-363/), using the Python packages DABEST[48],
scipy[49], numpy[50], matplotlib[51], seaborn[52], pandas[53]. Throughout this study, we used a bootstrap-
coupled estimation of effect sizes[54]. For each estimation plot showing a difference between TMR and
No TMR groups: the left panel shows the distribution of raw data points for the entire dataset,
superimposed on plots reporting group means ± SEM; and the right panel displays the difference
between the TMR group and the No TMR group, computed from 10,000 bootstrapped re-samples.
For each estimation plot: black dot, mean; black ticks, 90% or 95% confidence interval; and filled
curve: bootstrapped sampling error distribution. We used 90% confidence intervals (one-sided) for
tests of difference between TMR and No TMR groups. For any other tests, we used 95% confidence
intervals (two-sided).

Data and Code Availability Statement

599 The data and code used in this study will be made available via the MRC BNDU Data Sharing Platform
600 (https://data.mrc.ox.ac.uk) upon publication.

References

1. Tolman, E. C. Cognitive maps in rats and men. *Psychol. Rev.* **55**, 189–208 (1948).

2. Brogden, W. J. Sensory pre-conditioning. *J. Exp. Psychol.* **25**, 323 (1939).

3. Wamsley, E. J. Offline memory consolidation during waking rest. *Nat. Rev. Psychol.* **1**, 441–453 (2022).

4. Diekelmann, S. & Born, J. The memory function of sleep. *Nat. Rev. Neurosci.* **11**, 114–126 (2010).

5. Coutanche, M. N., Gianessi, C. A., Chanales, A. J. H., Willison, K. W. & Thompson-Schill, S. L. The role of sleep in forming a memory representation of a two-dimensional space. *Hippocampus* **23**, 1189–1197 (2013).

6. Werchan, D. M. & Gómez, R. L. Generalizing memories over time: Sleep and reinforcement facilitate transitive inference. *Neurobiol. Learn. Mem.* **100**, 70–76 (2013).

7. van Kesteren, M. T. R., Fernández, G., Norris, D. G. & Hermans, E. J. Persistent schema-dependent hippocampal-neocortical connectivity during memory encoding and postencoding rest in humans. *Proc. Natl. Acad. Sci.* **107**, 7550–7555 (2010).

8. Ellenbogen, J. M., Hu, P. T., Payne, J. D., Titone, D. & Walker, M. P. Human relational memory requires time and sleep. *Proc. Natl. Acad. Sci.* **104**, 7723–7728 (2007).

9. Lau, H., Alger, S. E. & Fishbein, W. Relational Memory: A Daytime Nap Facilitates the Abstraction of General Concepts. *PLOS ONE* **6**, e27139 (2011).

10. Craig, M., Dewar, M., Harris, M. A., Della Sala, S. & Wolbers, T. Wakeful rest promotes the integration of spatial memories into accurate cognitive maps. *Hippocampus* **26**, 185–193 (2016).

11. Foster, D. J. Replay Comes of Age. *Annu. Rev. Neurosci.* **40**, 581–602 (2017).

12. Joo, H. R. & Frank, L. M. The hippocampal sharp wave–ripple in memory retrieval for immediate use and consolidation. *Nat. Rev. Neurosci.* **19**, 744–757 (2018).

13. Buzsáki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25**, 1073–1188 (2015).

14. Bendor, D. & Wilson, M. A. Biasing the content of hippocampal replay during sleep. *Nat.*

Neurosci. **15**, 1439–1444 (2012).

15. Kudrimoti, H. S., Barnes, C. A. & McNaughton, B. L. Reactivation of Hippocampal Cell Assemblies:

Effects of Behavioral State, Experience, and EEG Dynamics. *J. Neurosci.* **19**, 4090–4101 (1999).

16. O'Neill, J., Senior, T. J., Allen, K., Huxter, J. R. & Csicsvari, J. Reactivation of experience-dependent

cell assembly patterns in the hippocampus. *Nat. Neurosci.* **11**, 209–215 (2008).

17. Ólafsdóttir, H. F., Bush, D. & Barry, C. The Role of Hippocampal Replay in Memory and Planning.

Curr. Biol. **28**, R37–R50 (2018).

18. Schuck, N. W. & Niv, Y. Sequential replay of nonspatial task states in the human hippocampus.

Science **364**, eaaw5181 (2019).

19. Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal cellular

assemblies. *Nature* **469**, 397–401 (2011).

20. Ólafsdóttir, H. F., Barry, C., Saleem, A. B., Hassabis, D. & Spiers, H. J. Hippocampal place cells

construct reward related sequences through unexplored space. *eLife* **4**, e06063 (2015).

21. Liu, Y., Dolan, R. J., Kurth-Nelson, Z. & Behrens, T. E. J. Human Replay Spontaneously Reorganizes

Experience. *Cell* **178**, 640-652.e14 (2019).

22. Barron, H. C. *et al.* Neuronal Computation Underlying Inferential Reasoning in Humans and Mice.

Cell **183**, 228-243.e21 (2020).

23. Wu, X. & Foster, D. J. Hippocampal replay captures the unique topological structure of a novel

environment. *J. Neurosci. Off. J. Soc. Neurosci.* **34**, 6459–6469 (2014).

24. Gupta, A. S., van der Meer, M. A. A., Touretzky, D. S. & Redish, A. D. Hippocampal Replay Is Not a

Simple Function of Experience. *Neuron* **65**, 695–705 (2010).

25. Schouten, D. I., Pereira, S. I. R., Tops, M. & Louzada, F. M. State of the art on targeted memory

reactivation: Sleep your way to enhanced cognition. *Sleep Med. Rev.* **32**, 123–131 (2017).

26. Oudiette, D., Antony, J. W., Creery, J. D. & Paller, K. A. The Role of Memory Reactivation during

Wakefulness and Sleep in Determining Which Memories Endure. *J. Neurosci.* **33**, 6672–6678

(2013).

27. Lewis, P. A. & Bendor, D. How Targeted Memory Reactivation Promotes the Selective

Strengthening of Memories in Sleep. *Curr. Biol.* **29**, R906–R912 (2019).

28. Hu, X., Cheng, L. Y., Chiu, M. H. & Paller, K. A. Promoting memory consolidation during sleep: A

meta-analysis of targeted memory reactivation. *Psychol. Bull.* **146**, 218–244 (2020).

29. Cellini, N. & Capuozzo, A. Shaping memory consolidation via targeted memory reactivation

during sleep. *Ann. N. Y. Acad. Sci.* **1426**, 52–71 (2018).

30. Zhou, Z., Singh, D., Tandoc, M. C. & Schapiro, A. C. Building integrated representations through

interleaved learning. *J. Exp. Psychol. Gen.* **152**, 2666–2684 (2023).

31. Rudoy, J. D., Voss, J. L., Westerberg, C. E. & Paller, K. A. Strengthening Individual Memories by

Reactivating Them During Sleep. *Science* **326**, 1079–1079 (2009).

32. Tadros, T. & Bazhenov, M. Role of Sleep in Formation of Relational Associative Memory. *J.

Neurosci.* **42**, 5330–5345 (2022).

33. Abdou, K. *et al.* Prefrontal coding of learned and inferred knowledge during REM and NREM

sleep. *Nat. Commun.* **15**, 4566 (2024).

34. Tamminen, J., Payne, J. D., Stickgold, R., Wamsley, E. J. & Gaskell, M. G. Sleep Spindle Activity is

Associated with the Integration of New Memories and Existing Knowledge. *J. Neurosci.* **30**,

14356–14360 (2010).

35. Tamminen, J., Ralph, M. A. L. & Lewis, P. A. The Role of Sleep Spindles and Slow-Wave Activity in

Integrating New Information in Semantic Memory. *J. Neurosci.* **33**, 15376–15381 (2013).

36. Tamminen, J., Lambon Ralph, M. A. & Lewis, P. A. Targeted memory reactivation of newly learned

words during sleep triggers REM-mediated integration of new memories and existing knowledge.

Neurobiol. Learn. Mem. **137**, 77–82 (2017).

37. Landmann, N. *et al.* The reorganisation of memory during sleep. *Sleep Med. Rev.* **18**, 531–541

(2014).

38. Rasch, B., Büchel, C., Gais, S. & Born, J. Odor Cues During Slow-Wave Sleep Prompt Declarative

Memory Consolidation. *Science* **315**, 1426–1429 (2007).

39. Ngo, H.-V. V. & Staresina, B. P. Shaping overnight consolidation via slow-oscillation closed-loop

targeted memory reactivation. *Proc. Natl. Acad. Sci.* **119**, e2123428119 (2022).

40. Paller, K. A. Sleeping in a Brave New World: Opportunities for Improving Learning and Clinical

Outcomes Through Targeted Memory Reactivation. *Curr. Dir. Psychol. Sci.* **26**, 532–537 (2017).

41. Diekelmann, S., Büchel, C., Born, J. & Rasch, B. Labile or stable: opposing consequences for

memory when reactivated during waking and sleep. *Nat. Neurosci.* **14**, 381–386 (2011).

42. Tambini, A., Berners-Lee, A. & Davachi, L. Brief targeted memory reactivation during the awake

state enhances memory stability and benefits the weakest memories. *Sci. Rep.* **7**, 15325 (2017).

43. Schapiro, A. C., McDevitt, E. A., Rogers, T. T., Mednick, S. C. & Norman, K. A. Human hippocampal

replay during rest prioritizes weakly learned information and predicts memory performance.

Nat. Commun. **9**, 3920 (2018).

44. Creery, J. D., Oudiette, D., Antony, J. W. & Paller, K. A. Targeted Memory Reactivation during

Sleep Depends on Prior Learning. *Sleep* **38**, 755–763 (2015).

45. Cairney, S. A., Lindsay, S., Sobczak, J. M., Paller, K. A. & Gaskell, M. G. The Benefits of Targeted

Memory Reactivation for Consolidation in Sleep are Contingent on Memory Accuracy and Direct

Cue-Memory Associations. *Sleep* **39**, 1139–1150 (2016).

46. Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3? (2007).

47. Endo, N., Saiki, J., Nakao, Y. & Saito, H. Perceptual judgments of novel contour shapes and

hierarchical descriptions of geometrical properties. *Jpn. J. Psychol.* **74**, 346–353 (2003).

48. Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P values: data analysis

with estimation graphics. *Nat. Methods* **16**, 565–566 (2019).

49. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*

Methods **17**, 261–272 (2020).

50. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).

51. Matplotlib: A 2D Graphics Environment. https://ieeexplore.ieee.org/document/4160265/.

52. Waskom, M. L. seaborn: statistical data visualization. *J. Open Source Softw.* **6**, 3021 (2021).

53. McKinney, W. Data structures for statistical computing in python. in *Proceedings of the 9th*

Python in Science Conference vol. 445 51–56 (Austin, TX, 2010).

54. Efron, B. The Bootstrap and Modern Statistics. *J. Am. Stat. Assoc.* **95**, 1293–1296 (2000).

Acknowledgements

This research was supported by funding from the Medical Research Council (UKRI-MRC)
(MR/W01971X/1, MR/L019639/1 to J.X.R.), UKRI (MR/W008939/1 to H.C.B.), and the Biotechnology
and Biological Sciences Research Council (UKRI-BBSRC) (BB/M011224/1 to A.B.R.). The Wellcome
Centre for Integrative Neuroimaging is supported by core funding from the Wellcome Trust
(203139/Z/16/Z). The funders had no role in study design, data collection and analysis, decision to
publish or preparation of the manuscript.

Author Information

Authors and Affiliations

Department of Experimental Psychology, University of Oxford, Oxford, United Kingdom

Cal M. Shearer, Jill X. O'Reilly

Medical Research Council Brain Network Dynamics Unit, Nuffield Department for Clinical
Neurosciences, University of Oxford, Oxford, United Kingdom

Cal M. Shearer, Annalise B. Rawson, Helen C. Barron

Wellcome Centre of Integrative Neuroimaging, University of Oxford, FMRIB, John Radcliffe
Hospital, Oxford, United Kingdom

Helen C. Barron, Annalise B. Rawson

Contributions

All authors contributed to the experimental design and editing the paper. C.M.S, H.C.B, and J.X.R
contributed to data analysis and writing the paper. C.M.S. collected the data.

Corresponding author

Cal Shearer (cal.shearer@psy.ox.ac.uk)

Ethics declarations

Competing interests

The authors declare no competing interests.

Supplemental Information

Figure S1. Distractor Task Does Not Change the Effect of TMR On Inferential Choice.
(A) Schematic: Example trials for different distractor task types during the inference test. Trials start with either no distractor
task (left), a memory distractor task (participants are asked if a shape is the same or different to the one seen on the

previous trial (middle)), or a non-memory secondary task (participants are asked to identify if a photo is of something living
or non-living (right)). On all trial types, participants then hear an auditory cue and are asked to infer the outcome associated
with that auditory cue. Full details of the inference part of each trial are shown in Figure 2B.
(B) Inference test performance broken down by distractor task type and TMR group. Bars: mean percentage inference
accuracy for no distractor task (left group), memory task (middle group), and non-memory task (right group) groups split by
TMR (blue; left bar of each group) and No TMR (red; right bar of each group); black ticks: ± SEM; each data point: mean
accuracy for one participant.
(C) Distractor task performance. Bars: mean percentage accuracy on the distractor task for memory task (orange; left) and
non-memory task (green; right); black ticks: ± SEM; each data point: mean accuracy for one participant.
(D) Difference in TMR effect on inference between different distractor tasks. Right: raw data points for no distractor task
(blue; right), memory task (orange; middle), non-memory task (green, right); each data point: mean effect of TMR (TMR
group accuracy – No TMR group accuracy) for a given participant; black dots, mean; black ticks ± SEM. Right: difference in
means between no distractor task (dashed black line) and other distractor task conditions shown using bootstrap-coupled
estimation (DABEST) plots as in Figure 2C: black dots, mean; black ticks, 95% confidence interval; filled-curve, sampling-
error distribution. There was no significant difference in the effect of TMR on accuracy between distractor task types
(p=0.130, no distractor task vs memory task; p=0.372, no distractor task vs non-memory task).

Figure S2. Differences in Overall Accuracy Cannot Explain the Effect of TMR on Direct and Indirect Associations.
(A-D) The effect of TMR on indirect associations (A; C) and direct associations (B; D) for participants split by overall accuracy in the top 50% (A-B) and bottom 50% (C-D). Left of each panel: raw data points for No TMR group (red; left) and TMR group (blue; right); each data point: mean accuracy for a given participant; black dots, mean; black ticks ± SEM. Right of each panel: difference in means between No TMR and TMR groups shown using bootstrap-coupled estimation (DABEST) plots as in Figure 2C: black dot, mean; black ticks, 90% confidence interval; filled-curve, sampling-error distribution. There was a significant effect of TMR on accuracy for indirect associations (i.e. inference) for both the bottom (p=0.029) and top (p=0.036) 50% of participants. There was no significant effect of TMR on accuracy of direct associations, for either the bottom of top 50% of participants (p=0.451, top 50%; p=0.442, bottom 50%)