MTMR regulates KRAS function by controlling plasma membrane levels of phospholipids

Taylor E. Lange[1], Ali Naji[2], Ransome van der Hoeven[2], Hong Liang[3], Yong Zhou[3], Gerald R.V.

Hammond[4], John F. Hancock[3] and Kwang-jin Cho[1, 5]

1. Department of Biochemistry and Molecular Biology, Boonshoft School of Medicine, Wright

State University, Dayton, Ohio 45435, USA

2. Department of Diagnostic and Biomedical Sciences, School of Dentistry, The University of

Texas Health Science Center at Houston, Houston, Texas 77030, USA

3. Department of Integrative Biology and Pharmacology, McGovern Medical School, University

of Texas Health Science Center, Houston, Texas 77030, USA

4. Department of Cell Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA

15213, USA

5. To whom correspondence should be addressed: Kwang-jin Cho, Department of Biochemistry

and Molecular Biology, School of Boonshoft Medical School, Wright State University, Dayton,

OH 45435; kwang-jin.cho@wright.edu; Tel: (937) 775-2670

27 **eTOC summary:** We discovered that silencing the phosphatidylinositol (PI) 3-phosphatase,

28 *MTMR*, disrupts the PM localization of PtdSer and KRAS. We propose a model, where *MTMR*

29 loss depletes PM PI needed for PM PI4P synthesis, an essential phospholipid for PM PtdSer

30 enrichment, thereby impairing KRAS PM localization. (44 words)

Abstract (160 words, 950 characters)

KRAS, a small GTPase involved in cell proliferation and differentiation, frequently gains

activating mutations in human cancers. For KRAS to function, it must bind the plasma

membrane (PM) via interactions between its membrane anchor and phosphatidylserine (PtdSer).

Therefore, depleting PM PtdSer abrogates KRAS PM binding and activity. From a genome-wide

siRNA screen to identify genes regulating KRAS PM localization, we identified a set of

phosphatidylinositol (PI) 3-phosphatases: myotubularin-related proteins (MTMR) 2, 3, 4, and 7.

Here, we show that silencing *MTMR 2/3/4/7* disrupts KRAS PM interactions by reducing PM PI

4-phosphate (PI4P) levels, thereby disrupting the localization and operation of ORP5, a lipid

transfer protein maintaining PM PtdSer enrichment. Concomitantly, silencing *MTMR 2/3/4/7*

elevates PM PI3P levels while reducing PM and total PtdSer levels. We also observed MTMR

2/3/4/7 expression is interdependent. We propose that the PI 3-phosphatase activity of MTMR is

required for generating PM PI, necessary for PM PI4P synthesis, promoting the PM localization

of PtdSer and KRAS.

Introduction (692 words, 3,692 characters)

H-, N- and KRAS proteins are small GTPases that oscillate between inactive GDP-bound and active GTP-bound conformational states and operate in signaling cascades that control cell growth and proliferation (Gorfe and Cho, 2019; Henkels et al., 2021). Consistent with this key regulatory role, activating mutations of RAS are present in ~20% of human cancers, with the majority occurring in the KRAS isoform (Prior et al., 2020). All RAS isoforms must be localized to the inner leaflet of the plasma membrane (PM) and be spatially organized into nanoclusters for biological activity (Gorfe and Cho, 2019; Henkels et al., 2021). Membrane interactions are mediated by the RAS C-terminal membrane anchor (Zhou et al., 2017). In the case of KRAS4B, the major expressed splice variant of KRAS (Hood et al., 2023) (hereafter KRAS), this anchor comprises a C-terminal farnesyl cysteine methyl ester, generated by posttranslational modification, and an adjacent polybasic domain of six continuous lysines (Gutierrez et al., 1989; Hancock et al., 1991; Hancock et al., 1989; Hancock et al., 1990). Recent work has shown that the KRAS membrane anchor encodes exquisite binding specificity for phosphatidylserine (PtdSer) lipids with one saturated and one desaturated acyl chain (Zhou et al., 2017). This lipid binding specificity is effectively hard wired into the anchor structure and renders KRAS PM binding, nanoclustering and hence biological function absolutely dependent on PM PtdSer content (Cho et al., 2012b; Cho et al., 2013; van der Hoeven et al., 2013; van der Hoeven et al., 2018).

PtdSer is synthesized in the ER and delivered to the PM against a concentration gradient by the lipid transfer proteins (LTPs), oxysterol-binding protein-related protein (ORP) 5 and 8, operating at ER-PM membrane contact sites. One molecule of phosphatidylinositol (PI) 4-phosphate (P), generated by PM localized PI 4-kinase IIIα (PI4KA) from PI, is exchanged by ORP5/8 for one molecule of ER localized PtdSer (Chung et al., 2015; Moser von Filseck et al., 2015). PI4P delivered to the ER is converted to PI by the SAC1 phosphatase, which operating together with PI4KA, maintains the PI4P gradient that in turn, drives PtdSer transport to the PM (Chung et al., 2015; Moser von Filseck et al., 2015). Concordantly genetic and pharmacological studies have shown that inhibition or loss of any component of the PtdSer transport machinery, including PI4KA, EFR3A (the PM anchor protein for the kinase), ORP5, ORP8 and SAC1 results in a reduction in PM PtdSer content, displacement of KRAS from the PM and abrogation of KRAS signaling (Gulyas et al., 2017; Kattan et al., 2019; Kattan et al., 2021; Sohn et al., 2018). Other pharmacological approaches to deplete PM PtdSer also dissociate KRAS from the PM (Cho et al., 2012b; Garrido et al., 2020; Miller et al., 2019; Tan et al., 2019; Tan et al., 2018; van der Hoeven et al., 2018). The PM is a dynamic organelle, thus KRAS on endomembranes is

sequestered by a chaperone protein, phosphodiesterase 6δ, and released to perinuclear
membranes in the ARL2/3-dependent manner. The negative charge on the recycling endosome
membranes then electrostatically traps KRAS, from where vesicular transport returns KRAS to
the PM (Chandra et al., 2012; Schmick et al., 2014). KRAS phosphorylation at the Ser181
residue can disrupt this electrostatic interaction, depleting KRAS from the PM (Cho et al., 2016a;
Kovar et al., 2020).

In this context we carried out a cell-based high content screen using a human genomic
short interfering (siRNA) library to identify additional potential regulators of KRAS PM
localization. Among the hits from the assay were multiple genes that encode proteins with
phosphatase activity. Four of these genes encode myotubularin-related proteins (MTMR) 2, 3, 4
and 7, which have 3-phosphatase activity towards phosphatidylinositol (PI) 3-phosphate (P) and
PI(3,5)-bisphosphate (PI(3,5)P_2) (Robinson and Dixon, 2006). In this study, we examined the
role of MTMR in the PM localization of KRAS. Our data demonstrate that knockdown (KD) of
MTMR 2, 3, 4 or 7 dissociates KRAS, but not HRAS from the PM. We show *inter alia* that the
molecular mechanism involves depletion of the PI4P and PtdSer content of the PM, extending
the repertoire of proteins required to support the operation of the ORP5/8 LTPs and hence
sustain KRAS PM localization.

Method and Material (1,617 words, 9.511 characters)
Plasmids and reagents. shRNA glycerol sets for MTMR2 (RHS4533-EG8898), MTMR3
(RHS4533-EG8897), MTMR4 (RHS4533-EG9110), MTMR7 (RHS4533-EG9108), and human
cDNA of MTMR2 (MHS6278-202760229), MTMR3 (MHS6278-202800229), MTMR4
(MHS6278-202801286) and MTMR7 (MHS6278-211688199) were purchased from GE
Healthcare Dharmacon. Antibodies were purchased for detecting MTMR2 (PA5-22748; 1:1,000)
and c-MYC (13-2500, 1:1,000) from Invitrogen, MTMR3 (sc-393779; 1:1,000) from SCBT, and
MTMR7 (AB150458; 1:1,000) from Abcam. Antibodies for detecting c-MYC (10828-1-AP,
1:3,000), β-actin (60008-1-Ig; 1:5,000), MTMR3 (21336-1-AP; 1:1,000) and GFP (66002-1-Ig;
1:4,000) were purchased from Proteintech. Anti-ppERK (4370; 1:3,000), anti-pAkt (S473) (4060;
1:1,000), anti-total ERK (4696; 1:1,000), anti-total Akt (2920; 1:1,000) antibodies for
immunoblotting were from Cell Signaling Technology. Anti-RFP antibody (sc-390909, 1:1,000)
was from Santa Cruz. Goat anti-mouse IgG (G21040; 1:2,000), anti-rabbit IgG secondary
antibodies (G21234; 1:5,000) and CellMask Deep Red plasma membrane stain (C10046;
1:5,000) were purchased from Invitrogen. Staurosporine (BIA-S1086) was purchased from
BioAustralis.

Human whole-genome siRNA screen. A whole-genome siRNA screen was performed in Caco-2 cells stably expressing GFP-KRASG12V using a pre-aliquoted Silencer siRNA library (Ambion) at 30 nM final concentration. A total of 21,687 genes were represented in the library and were arrayed into glass-bottomed sixty-nine 384-well plates in triplicate. Gene silencing was induced for 4 days via reverse transfection using Dharmafect 1 (GE Healthcare). For controls, each plate included non-targeting control siRNA and siRNA targeting *FNTA* (farnesyltransferase type1 subunit alpha) as negative and positive controls, respectively. Also, each plate included the *Kif11* siRNA that targets a gene essential for cell survival served as control for transfection and knockdown efficiency. After the gene silencing, cells were fixed with 4% paraformaldehyde (PFA) and stored in 1x phosphate-buffered saline (PBS) at 4^{o}C until imaged. Plates were imaged using Nikon A1R confocal microscope. Images were acquired as a 2 by 1 montage with a 20x objective lens using the GFP-confocal mode with NIS-Elements automated imaging software. Images were scored for KRAS PM dissociation/intracellular accumulation of GFP-KRASG12V and cytotoxicity. 253 genes were selected from the primary screen based on KRAS PM dissociation, and after a counter screen with GFP-HRASG12V to identify KRAS-specific mediators, we refined the candidate list to 157. The 157 genes were further validated with an independent secondary screen involving silencing each gene with Dharmacon SmartPool ON-TARGET plus siRNA in Caco-2 cells expressing GFP-KRASG12V and scoring for KRAS PM dissociation. A further bioinformatics analysis revealed that 8 of these genes encode proteins with phosphatase activity (Table 1).

Cell lines. T47D (HTB-13, ATCC) and BxPC3 (CRL-1687, ATCC) cells were maintained in RPMI-1640 medium (30-2001, ATCC) supplemented with 2 mM L-glutamine (CA009-010, GenDepot) and 10% fetal bovine serum (FBS) (16000044, Gibco). For T47D cells, 10 μg/mL human recombinant insulin (12585-014, Gibco) was also supplemented. MIA PaCa-2 (CRL-1420, ATCC), PANC-1 (CRL-1469, ATCC) and Caco-2 (HTB-37, ATCC) cells were maintained in DMEM medium with high glucose (CM002, Gibco) supplemented with 2 mM L-glutamine and 10% FBS. All cells were grown at 37^{o}C with 5% CO_2, and frequently tested for mycoplasma using MycoAlert Mycoplasma Detection Kit (LT07-318, Lonza).

Generating T47D stable cell lines. T47D cells were transfected with GFP-KRASG12V, -HRASG12V or -LactC2 in pEF6 vector (Invitrogen) using *Trans*IT-BrCa Transfection Reagent (MIR 5500, Mirus), and selected with 10 μg/mL blasticidin for 3 weeks. After the selection, cells were maintained in 5 μg/mL blasticidin. For generating MTMR knockdown cell lines, wild-type

(WT) T47D cells were infected with lentivirus expressing shRNA targeting *MTMR 2/3/4/7.* Cells

were selected by puromycin (1 μg/mL) for 1 week and maintained in 0.5 μg/mL puromycin.

Preparing T47D cells for imaging and calculating Manders' coefficient. 5×10^5 T47D cells

were seeded on a coverslip on day 1 in RPMI-1640 medium containing 10% FBS, 2 mM L-

glutamine and 10 μg/mL insulin (complete growth medium). On day 2, fresh complete growth

medium was supplemented. On day 3, cells were fixed with 4% PFA for 30 min, quenched with

50 mM NH_4Cl for 10 min and mounted with Vectashield (H-1000, Vector Laboratories). For

calculating Manders' coefficient, cells were incubated with CellMask stain (1:5,000) for 1 h in

complete growth medium, fixed with 4% PFA, and imaged by confocal microscopy. Using

ImageJ software 1.53k, images were converted to 8-bit, and a threshold to a control pixel of

each image was set. The fraction of red-fluorophore-conjugated CellMask co-localizing with

GFP-tagged proteins was calculated using Manders' coefficient plugin downloaded from Wright

Cell Image Facility.

Lipidomic analysis. T47D cells stably expressing GFP-KRASG12V with *MTMR 2/3/4/7*

knockdown were cultured in complete growth medium containing 0.5 μg/ml puromycin. Triplicate

samples of 1×10^6 cells were prepared in 333 μL Dulbecco's PBS (DPBS without Ca^{2+} and Mg^{2+},

14190144, Invitrogen). Lipid extraction and analysis using electron spray ionization and MS/ MS

were performed at Lipotype GmbH (Dresden, Germany), as described previously (Gerl et al.,

2012; Sampaio et al., 2011). Automated processing of acquired mass spectra, identification,

and quantification of detected lipid species were done by LipidXplorer software. Only lipid

identifications with a signal-to-noise ratio >5, an absolute abundance of at least 1 pmol, and a

signal intensity 5-fold higher than in corresponding blank samples were considered for further

data analysis. The abundance of lipids is represented as a heat map with log_2 scale relative to

control (scrambled shRNA) cells.

Measuring total PI3P level. The total amount of PI(3)P was examined in a quantitative and

competitive ELISA format assay according to the manufacturer's instructions (K-3300, Echelon

Biosciences). Briefly, *MTMR 2/3/4/ KD* T47D cells grown on T75 flasks were collected in ice-

cold 0.5 M trichloroacetic acid (TCA) and centrifuged at 1,000 x g for 10 min at 4^oC. The pellet

was washed with 3 mL 5% TCA/1 mM EDTA and neutral lipids were removed using

MeOH:$CHCl_3$ (2:1), followed by acidic lipids extraction using MeOH:$CHCl_3$:12N HCl (80:40:1).

The acidic lipid extracts were further separated into organic and aqueous phases, and the

organic phase was dried in a vacuum dryer. The dried lipid was rehydrated in PBS-T 3% protein

Western blotting. Preparation of cell lysates and immunoblotting were performed as described previously (Cho et al., 2011). Briefly, cells were washed twice with ice-cold 1x PBS. Cells were harvested in lysis buffer B containing 50 mM Tris-Cl (pH 7.5), 75 mM NaCl, 25 mM NaF, 5 mM $MgCl_2$, 5 mM EGTA, 1 mM DTT, 100 µM $NaVO_4$, 1% NP-40 plus protease and phosphatase inhibitors. SDS-PAGE and immunoblotting were generally performed using 20 – 40 µg of lysate from each sample group. Signals were detected by enhanced chemiluminescence (34578 and 34075, Thermo Fisher Scientific) and imaged using an Amersham Imager 600 (GE Healthcare). ImageJ software (v1.53k) was used to quantify band intensity.

Validating MTMR4 knockdown. Messenger RNA (mRNA) was extracted from T47D cells infected with lentivirus expressing shRNA targeting MTMR4 using the RNeasy kit (74104, Qiagen) according to the manufacturer's instructions. 2 µg of total RNA was converted to cDNA using Superscript II reverse transcriptase (18064-014, Thermo Fisher Scientific). To verify knockdown, forward and reverse primers specific for human *MTMR4* exons 1 and 5, and *GAPDH* (glyceraldehyde-3-phosphate dehydrogenase) exons 1 and 3 were designed: for *MTMR4*, 5'-CCAAGCCAAGGATCTGTTCCC-3' and 5'-TGTGTGAGACTCTCCAGACGT-'3', and for *GAPDH*, 5'- GGAGCGAGATCCCTCCAAAAT-3' and 5'-GGCTGTTGTCATACTTCTCATGG-3', respectively. cDNA was amplified by PCR (P2311, GenDepot) and the products were resolved by 1.5% agarose gel electrophoresis and visualized by ethidium bromide staining.

***C. elegans* vulva quantification assay.** *let-60* (n1046) worms were kindly provided by Swathi Arur (MD Anderson Cancer Center, Houston, TX). RNAi was induced by feeding *let-60* worms through adult stage with *E. coli* HT115, producing double-stranded RNA to target genes. The presence of the multivulva phenotype was scored using a DIC/Nomarski microscope. All RNAi clones were from the *C. elegans* RNAi (Ahringer) collection (Source BioScience) and were sequenced.

Proliferation assay. *MTMR 2/3/4/7* knockdown T47D cells stably expressing GFP-KRASG12V or -HRASG12V (1 x 10^4 per well), or human pancreatic ductal adenocarcinoma cell lines, BxPC3, MIA PaCa-2 and PANC-1 (3 x 10^4 per well) were seeded in triplicate onto a 96-well plate in complete growth medium containing 1.0 µg/mL puromycin. Cells were cultured for 4 days, and fresh complete growth medium containing 1.0 µg/mL puromycin was supplemented every 24 h. On day 5, cell proliferation was assayed by counting cells using Countess II cell counter (Invitrogen).

Annexin V binding assay. *MTMR 2/3/4/7* knockdown T47D cells stably expressing GFP-KRASG12V were incubated with Cy5-conjugated annexin V (559934, BD Pharmingen) according to the manufacturer's instructions. Annexin V-positive cells were counted using BD AccuriC6 Analyzer with a 670LP nm filter. GFP-KRASG12V T47D cells expressing scrambled shRNA were treated with 1 μM staurosporine for 6 h to induce apoptosis.

Electron microscopy. Plasma membrane (PM) sheets were prepared from Caco-2 or T47D cells expressing GFP-tagged proteins of interest and fixed as previously described (Hancock and Prior, 2005; Prior et al., 2003a; Prior et al., 2003b). For univariate analysis, PM sheets were labeled with anti-GFP antibody conjugated to 4.5-nm gold particles. Digital images of the immunogold-labeled PM sheets were taken in a transmission electron microscope. Intact 1 μm^2 areas of the PM sheet were identified using ImageJ software (v1.53k), and the (x, y) coordinates of the gold particles were determined (Hancock and Prior, 2005; Prior et al., 2003a; Prior et al., 2003b). K-functions (Ripley, 1977) were calculated and standardized on the 99% confidence interval (CI) for univariate functions (Diggle, 2000; Hancock and Prior, 2005; Prior et al., 2003a; Prior et al., 2003b). In the case of univariate functions, a value of $L(r) – r$ greater than the CI indicates significant clustering, and the maximum value of the function (L_{max}) estimates the extent of clustering. Differences between replicated point patterns were analyzed by constructing bootstrap tests as described previously (Diggle, 2000; Plowman and Hancock, 2005), and the statistical significance against the results obtained with 1,000 bootstrap samples was evaluated.

Confocal microscope imaging acquisition. *A) Make and Model of microscope:* Olympus Fluoview F1000. *B) Type, magnification and numerical aperture of the objective lenses:* Olympus UPlanSApo 60x/1.35 oil, ∞/0.17/FN26.5 UIS2 BFP1. *C) Temperature:* room temperature. *D) Imaging medium:* Vectashield (H-1000, Vector Laboratories). *E) Fluorochromes:* GFP and mCherry. *F) Camera make and model:* N/A. *G) Acquisition software:* FV10-ASW ver.4.2c. H) Any software used for imaging processing subsequent to data acquisition:* None.

A summary of Supplement Figures

Supplement Figure 1. MTMR 2/3/4/7 regulate the PM localization of KRASG12V.

Supplement Figure 2. *MTMR 2/3/4/7* KD disrupts the PM localization and nanoclustering of CTK.

Supplement Figure 3. MTMR 2/3/4/7 regulate cellular PI3P contents.

Supplement Figure 4. Ectopic expression of WT MTMR4 restores the effects of *MTMR4* loss.

Results

Genome-wide siRNA screening identifies novel genes regulating KRAS PM interaction.

To identify novel regulators of KRAS PM interaction, we performed an image-based high

content screen of a human genomic siRNA library. Caco-2 (human colorectal adenocarcinoma)

cell line stably expressing monomeric green fluorescent protein (GFP)-tagged oncogenic mutant

KRAS (KRASG12V) was generated and transfected with siRNA pools comprising four different

oligonucleotides against each gene. Four days after transfection, cells were imaged using an

automated confocal microscope to analyze the extent of KRAS mislocalization from the PM. We

then narrowed the list of hits using a counter screen against HRASG12V to identify KRASG12V-

specific mediators. Bioinformatics analysis revealed that 8 of these putative KRASG12V-specific

mediators encode proteins with phosphatase activity (Table 1), 4 of which are members of the

myotubularin-related protein (MTMR) family. There are 16 human myotubularins, which exhibit

3-phosphatase activity towards PI3P and PI(3,5)P_2, generating PI and PI5P, respectively

(Robinson and Dixon, 2006; Wang et al., 2024). MTMR family has previously been shown to

regulate many cellular processes including endocytosis, membrane trafficking, cell proliferation,

autophagy, cytokinesis and cytoskeletal dynamics (Hnia et al., 2012). We selected this *MTMR*

gene set for further analysis.

MTMR 2/3/4/7 regulate the PM localization of KRASG12V, but not HRASG12V. To validate

the siRNA screen, we generated short-hairpin RNA (shRNA)-mediated stable KD cell lines.

T47D (human mammary gland ductal carcinoma) epithelial cells stably expressing GFP-

KRASG12V were infected with lentiviruses expressing shRNA targeting *MTMR2, 3, 4* and *7*,

followed by puromycin selection. For each target gene, we tested 4 different shRNAs and

selected the two most effective KD for further experiments. Immunoblotting verified that

endogenous expression levels of MTMR2, 3 and 7 were reduced by the cognate shRNAs (Fig.

1A). MTMR4 mRNA levels were also reduced by shRNA expression; we evaluated mRNA

because the anti-MTMR4 antibodies we tested were not able to detect endogenous MTMR4

(Fig. 1A). To study KRAS cellular localization, cells were incubated with CellMask, a dye

staining cellular membranes (Hu et al., 2013; Maechler et al., 2019; Monkemoller et al., 2015;

Park et al., 2019), and imaged by confocal microscopy. To quantify the extent of KRAS

distribution to endomembranes, we calculated Manders' coefficient, which measures the fraction

of CellMask co-localized with GFP-KRASG12V (Cho et al., 2012b; Manders et al., 1993; Rehl et

al., 2023). Our data show that while GFP-KRASG12V was predominantly localized to the PM in

control cells, it was distributed intracellularly in the KD cell lines (Fig. 1B and S1A). Manders'

coefficients for CellMask are correspondingly higher in the KD cell lines, indicating that

KRASG12V is redistributed to endomembranes after silencing *MTMR 2/3/4/7*. To directly

quantify the extent of KRAS dissociation from the PM, intact basal PM sheets from T47D stably

co-expressing GFP-KRASG12V and shRNA for *MTMR 2/3/4/7* were prepared and labeled with

gold-conjugated anti-GFP antibodies and analyzed by electron microscopy (EM). We observed

significant reductions in anti-GFP immunogold labeling after the KD, indicating loss of

KRASG12V from the inner PM leaflet (Fig. 1C and S1B). Spatial mapping of GFP-KRASG12V

on the PM in *MTMR 2/3/4/7* KD cells revealed significant decreases in L_{max}, the peak value of

the *L(r)-r* clustering statistic that measures the extent of KRAS nanoclustering, which is

essential for high-fidelity RAS signal transduction (Cho and Hancock, 2013; Cho et al., 2012a;

Tian et al., 2007) (Fig. 1E). Confocal and electron microscopy further demonstrated that while

MTMR 2/3/4/7 KD did not disrupt the PM localization and nanoclustering of HRASG12V, it

disrupts that of the C-terminal truncated mutant KRAS (CTK), which contains the membrane

anchor without the G-domain, from the PM (Fig. 1B, D, F and S2). Together, our data show that

MTMR 2/3/4/7 regulate the PM localization and nanoclustering of KRAS, but not HRAS,

independent of the G-domain.

MTMR 2/3/4/7 regulate the cellular level and distribution of PtdSer. The PM interaction of

KRAS, but not HRAS, is dependent on PM PtdSer content (Cho et al., 2012b; Cho et al., 2016b;

Garrido et al., 2020; Miller et al., 2019; Tan et al., 2018; van der Hoeven et al., 2018; Zhou et al.,

2017). To quantify PM PtdSer content, T47D cells stably co-expressing GFP-LactC2, a well-

characterized PtdSer probe (Yeung et al., 2008), and shRNA targeting *MTMR 2/3/4/7* were

incubated with CellMask and imaged by confocal microscopy. GFP-LactC2 was primarily

localized to the PM in control cells but was redistributed in *MTMR 2/3/4/7* KD cells as reflected

by higher Manders' coefficient values (Fig. 2A). EM analysis of isolated intact PM sheets

prepared from T47D cells expressing GFP-LactC2 and incubated with anti-GFP antibody-

conjugated gold shows that gold particle labeling density of the inner basal PM was significantly

reduced when cells co-express shRNAs for *MTMR 2/3/4/7,* indicating depletion of inner PM

PtdSer (Fig. 2B). The loss of inner leaflet PtdSer was not due to flipping to the outer leaflet since

Annexin V binding to the outer PM leaflet, as measured by flow cytometry, did not increase in

MTMR 2/3/4/7 KD cells (Fig. 2C). Finally, whole cell lipidomics revealed that *MTMR 2/3/4/7* KD

resulted in a significant reduction in total cellular PtdSer content (Figs. 2D and E). Together,

these results show that MTMR 2/3/4/7 potently regulates the PM and total levels of PtdSer;

consistent with a mechanistic role for MTMR 2/3/4/7 in controlling PtdSer cellular distribution.

MTMR 2/3/4/7 regulate PI4P content at the PM. In mammalian cells, inner PM leaflet PI4P and PI(4,5)P_2 recruit the LTPs, ORP5 and ORP8, to ER-PM membrane contact sites (MCSs) to exchange newly synthesized PtdSer in the ER for PI4P from the PM (Chung et al., 2015; Ghai et al., 2017; Moser von Filseck et al., 2015; Sohn et al., 2018). Depleting PM PI4P or PI(4,5)P_2, concordantly depletes PtdSer and KRAS from the PM (Gulyas et al., 2017; Kattan et al., 2019). Depletion of PI4P at the Golgi complex also mislocalizes PtdSer and KRAS from the PM via an as yet unknown mechanism (Miller et al., 2019). To examine whether silencing *MTMR 2/3/4/7* perturbs cellular localization of PI4P and PI(4,5)P_2, *MTMR 2/3/4/7* KD T47D cells expressing GFP-P4M-SidM or -PH-PLCδ1, probes for PI4P (Hammond et al., 2014) and PI(4,5)P_2 (Varnai and Balla, 1998), respectively, were imaged by confocal microscopy. GFP-P4M-SidM decorated the PM and Golgi complex in control cells (closed and open arrowheads, respectively in Fig. 3A), consistent with previous studies (Hammond et al., 2014; Miller et al., 2019), whereas *MTMR 2/3/4/7* KD perturbed GFP-P4M-SidM staining of the PM, but not the Golgi complex (Fig. 3A). By contrast, the PM localization of GFP-PH-PLCδ1 was unchanged in *MTMR 2/3/4/7* KD cells (Fig. 3A). EM analysis of immunogold labelled PM sheets also shows significant reduction in PI4P PM content, but no change in PI(4,5)P_2 PM content after *MTMR 2/3/4/7* KD (Fig. 3B and C). Given the reduction in PM PI4P content, we next examined ORP5 cellular localization. Confocal imaging showed that GFP-ORP5 predominantly localized to the ER-PM MCS in control cells, but substantially redistributed, presumably, to the ER, in *MTMR 2/3/4/7* KD cells (Fig. 3A) (Du et al., 2020; Monteiro-Cardoso et al., 2022). We conclude that MTMR 2/3/4/7 regulate ORP5 localization to the ER-PM MCS by maintaining the PM level of PI4P.

MTMR 2/3/4/7 **KD elevates PM PI3P content.** PI3P is synthesized from PI by class II and III PI3Ks, whence it can be converted to PI(3,5)P_2 or PI by PI 5-kinase, or myotubularin family phosphatases, respectively (Jean and Kiger, 2014; Pulido et al., 2013). To examine if depleting *MTMR 2/3/4/7* alters cellular PI3P levels or distribution, *MTMR 2/3/4/7* KD T47D cells expressing the well-characterized PI3P probe, GFP-2xFYVE, the tandem FYVE domain of Hrs (hepatocyte growth factor-regulated tyrosine kinase substrate) (Gillooly et al., 2000) were stained with CellMask and imaged by confocal microscopy. Manders' coefficient values estimating fractions of endomembranes co-localized with GFP-2xFYVE-decorated endosomes were elevated in *MTMR 3/4/7*, but not *MTMR2*, KD cells, suggesting increased PI3P content in 2xFYVE-decorated endosomes (Figs. 4A and B). Also, individual KD of *MTMR 2/3/4/7* gene elevated total cellular PI3P level (Fig. 4C), consistent with previous studies (Cao et al., 2008;

Lahiri et al., 2015; Ma et al., 2008; Zhao et al., 2019). Our data further show some PM
decoration with GFP-2xFYVE in control and *MTMR* KD cells with no noticeable difference
between cell lines, indicative of PI3P at the PM in these cell lines (arrowheads in Fig. 4A). To
formally quantify PM PI3P content, we performed EM analysis on intact basal PM sheets
prepared from *MTMR 2/3/4/7* KD T47D cells expressing GFP-2xFYVE. The data reveal that
silencing *MTMR 2/3/4/7* significantly elevated PI3P levels in the PM compared to the control
(Fig. 4D). Intriguingly, confocal and EM imaging further show GFP-MTMR 2/3/4/7 localization to
the PM (Fig. 4E and F), supported by previous studies reporting the PM localization of other
MTMR members, MTMR5 and 9 (Kim et al., 2003; Zou et al., 2009). We showed that while
MTMR 2/3/4/7 KD modestly increases PM PI3P level by less than 2-folds, the KD reduces PM
PI4P, PtdSer and KRAS levels to greater extents by 2 to 3-folds (Figs. 1 – 3). Together, these
data suggest that MTMR 2/3/4/7 localize to the PM, such that genetic depletion disrupts PI3P
conversion to PI at the PM. This in turn, modestly increases total PI3P content including at the
PM and 2xFYVE-decorated endosomes, which leads to substantial inhibition of the PM
localization of KRAS, PtdSer and PI4P.
365
The enzymatic activity of MTMR regulates the PM localization of KRAS, PtdSer and PI4P.
To examine if MTMR requires its enzymatic activity for maintaining the PM localization of KRAS,
PtdSer and PI4P, we overexpressed mCherry-tagged MTMR3 WT or C413S, the phosphatase-
inactive mutant (Walker et al., 2001), in *MTMR3* KD T47D cells stably expressing GFP-
KRASG12V or -LactC2, and analyzed their cellular localization by confocal and EM. Our data
show that while overexpression of MTMR3 restored the PM binding of GFP-KRASG12V and -
LactC2, the C413S mutant did not show any changes (Figs. 5A – D). We further assessed the
PM PI4P and PI3P contents by measuring the PM binding of GFP-P4M-SidM and -2xFYVE,
respectively, by EM. Like KRASG12V and PtdSer, MTMR3 WT overexpression promoted the
PM binding of GFP-P4M-SidM and -2xFYVE, but not the C413S mutant (Figs. 5E – F). We
repeated the experiments with MTMR4 WT and the C407S, the phosphatase-inactive mutant
(Lorenzo et al., 2006), and observed a similar trend. Overexpression of WT MTMR4, but not the
inactive mutant, reduced PM PI3P levels, and promoted the PM binding of GFP-KRASG12V, -
LactC2 and -P4M-SidM (Fig. S4). Moreover, despite the similar expression level, ectopic
expression of WT MTMR 2/4/7 did not restore the effects of *MTMR3* KD (Figs. 5A – G). We
made a similar observation with rescuing *MTMR4* KD, in which overexpression of WT MTMR
2/3/7 did not revert the effects of *MTMR4* KD (Fig. S4). Together, these data suggest that

MTMR members require their enzymatic activity for maintaining the PM localization of KRAS,

PtdSer and PI4P, and that the enzymatic activity of each MTMR is not functionally redundant.

Protein expression of MTMR 2/3/4/7 is interdependent. While there are 9 catalytically active

MTMR members (Wang et al., 2024), single KD of *MTMR 2/3/4/7* is sufficient to elevate PI3P

contents and dissociates KRAS, PtdSer and PI4P from the PM (Figs. 1 – 4). One plausible

explanation is that depleting single MTMR member may perturb the enzymatic activities of other

MTMR members, resulting in significant disruption of PI3P metabolism. To test this notion, we

first examined if the loss of multiple MTMR members can induce additive effects in these

phenotypes. *MTMR3* KD T47D cells were infected with lentivirus expressing shRNA targeting

MTMR2, 4, or *7*, but the double KD resulted in cytotoxicity (not shown). We then measured

endogenous protein levels of MTMR 2/3/7 after KD of single MTMR member. Our immunoblot

data show that the protein expression of MTMR 2/3/7 were concomitantly reduced upon

MTMR2, *3*, *4* or *7* KD (Fig. 6A), suggesting the protein expression of MTMR 2/3/7 are

interdependent. Moreover, while ectopic expression of WT MTMR3 in *MTMR3* KD cells restored

the protein levels of MTMR2 and 7, overexpression of WT MTMR2, 4 or 7 in *MTMR3* KD cells

did not restore the endogenous MTMR3 protein level (Fig. 6B and C). These data suggest that

the protein expression, and thereby the enzymatic activity, of MTMR 2/3/4/7 are interdependent.

MTMR 2/3/4/7 regulate KRAS signaling. Since *MTMR 2/3/4/7* KD dissociates KRASG12V

from the PM, we used two biological systems to evaluate effects on KRAS signal transmission.

First, we analyzed signaling by the KRAS ortholog, let-60 in the invertebrate model system

Caenorhabditis elegans (*C. elegans*), where constitutively active mutation of *let-60* induces a

quantifiable multivulva phenotype (Reiner et al., 2008). We used RNAi designed to target *C.*

elegans MTMR orthologs and examined whether *MTMR* KD could suppress the multivulva

phenotype associated with expression of LET-60 G13D (n1046). Our data show that silencing

mtm-3 (ortholog of *MTMR3*), or *mtm-6* (ortholog of *MTMR7*) abrogated multivulva formation,

and indeed were equipotent with two previously reported potent suppressors of LET-60 G13D

signaling (*heo-1* and *riok-1*) (Smith and Levitan, 2004; Weinberg et al., 2014) (Fig. 7A).

Secondly, we examined it in human cancer model cell lines. The proliferation of *MTMR 2/3/4/7*

KD T47D cells stably expressing GFP-KRASG12V, but not -HRASG12V, was significantly

inhibited compared to parental cells replete for each *MTMR* gene (Fig. 7B and C), suggesting

the growth inhibitory effects of *MTMR 2/3/4/7* KD is KRAS-specific. We also examined the

proliferation and RAS signaling in human pancreatic ductal adenocarcinoma (PDAC) cell lines,

MIA PaCa-2 and PANC-1, which harbor KRAS with activating point mutation, G12C and G12D,
respectively, and their growth depends on the oncogenic KRAS signaling (Hayes et al., 2016;
Kovar et al., 2020; Rehl et al., 2023). For a control PDAC cell line, we included BxPC-3, which
harbors WT KRAS, and its growth is independent of the KRAS signaling. We showed that
MTMR 2/3/4/7 KD inhibited the proliferation of MIA PaCa-2 and PANC-1, but not BxPC-3 cells
(Fig. 7D). For assessing RAS signal output, we measured active phosphorylation sites of
ERK1/2 and Akt for the RAS/MAPK and RAS/PI3P/Akt pathways, respectively. Our
immunoblotting data show a tendency of decreased phospho-ERK after *MTMR 2/3/4/7* KD in
KRAS-dependent PDAC (Fig. 7E). To further examine the ERK activity, we measured the
expression level of the transcriptional factor, c-MYC, since active ERK enhances its stability
(Sears et al., 1999). Our data show that c-MYC protein level was significantly decreased upon
MTMR 2/3/4/7 KD in KRAS-dependent, but not -independent PDAC cell lines, indicative of
blocked MAPK signaling in KRAS-dependent PDAC (Fig. 7E). Akt phosphorylation was elevated
in *MTMR2* and *7* KD cells, while no significant changes were observed upon *MTMR3* and *4* KD
in all three cell lines, suggesting that MTMR 2/3/4/7 differentially regulate the PI3K/Akt signaling
in an oncogenic KRAS-independent manner. We further measured endogenous KRAS
expression level since KRAS protein stability is dysregulated after the PM dissociation (Cho et
al., 2012b; Garrido et al., 2020; Kovar et al., 2020). Our immunoblot data show that endogenous
KRAS expression was significantly reduced upon *MTMR 2/3/4/7* KD in KRAS-dependent, but
not -independent PDAC cell lines (Fig. 7E). Together, our study provides strong evidence that
the MTMR phosphatases regulate KRAS signal transmission by maintaining PtdSer and KRAS
PM localization and their genetical loss dissociates the PM localization of PtdSer and KRAS, but
not HRAS, resulting in the inhibition of KRAS/MAPK/c-MYC signal output and instability of
KRAS protein expression in KRAS-dependent PDAC.
441
Discussion
We have identified new roles for the phosphatases, MTMR 2/3/4/7, in maintaining the PM
localization of PI4P, PtdSer and KRAS. We show that KD of *MTMR 2/3/4/7* dissociates PtdSer
and KRAS from the PM, and also reduces total cellular PtdSer level. Concomitantly, *MTMR*
2/3/4/7 KD elevates PM PI3P content, reduces PM PI4P content and dissociates ORP5 from the
ER-PM MCS. Collecting these observations together, we propose that the PM-localized MTMR
phosphatases are required to help maintain PM levels of PI for generating PM PI4P by PI4KA,
which in turn maintains total PtdSer levels and drives ER to PM transport of PtdSer. PI is
synthesized from phosphatidic acid at the cytosolic face of ER, where a fraction of PI is used for

the synthesis of glycosylphosphatidylinositol-anchored proteins (Blunsom and Cockcroft, 2020),

with the remaining PI being widely distributed across intracellular cytosolic membranes (Ashlin

et al., 2021; Pemberton et al., 2020; Zewe et al., 2020). PI delivered from the ER directly to the

PM by the LTPs, Nir2/3 and TMEM24 (Posor et al., 2022), is a crucial substrate for PM

polyphosphoinositides (PPIn). Simple overexpression of Nir2/3 and TMEM24 elevates the

synthesis and total amount of PM PPIn (Chang et al., 2013; Kim et al., 2015; Lees et al., 2017),

suggesting that PM PI is constantly converted to PPIn and the PI supply to the PM is rate-

limiting. In this context, the pathway to PI generation from PI3P by MTMR 2/3/4/7 at the PM,

emerges as a critical alternative supplier of PI to allow sufficient PM PI4P generation hence

PtdSer transport by the ORP5/8 machinery (Fig. 8). In addition to the PM, MTMR 2/3/4/7

localize to endosomes and promotes their trafficking by regulating endosomal PI3P and

PI(3,5)P$_2$ levels (Bonneick et al., 2005; Franklin et al., 2011; Mochizuki and Majerus, 2003;

Naughtin et al., 2010; Previtali et al., 2007; Walker et al., 2001). Thus, perturbing endosomal

trafficking by *MTMR 2/3/4/7* KD will block the delivery of endosomal PI to the PM, further

contributing to the reduced PM PI content. Therefore, when *MTMR 2/3/4/7* is depleted, PM

PtdSer level falls and KRAS dissociates from the PM (Fig. 8). We did not investigate the

mechanism whereby *MTMR 2/3/4/7* KD reduced total cellular PtdSer content, but previous work

has shown that PM PI4P depletion indirectly blocks PtdSer synthase 1 and 2 activities (Sohn et

al., 2016). Since PM PI4P levels were reduced in *MTMR 2/3/4/7* KD cells, the same mechanism

is likely responsible. Furthermore, inactivating mutations in *MTMR2* are found in Charcot-Marie-

Tooth disease type 4B1, a severe autosomal recessive neuropathy with demyelination and

myelin outfoldings of the nerve. Studies have proposed that disrupted endosomal trafficking by

elevating endosomal PI3P and PI(3,5)P$_2$ due to inactive MTMR2 likely contributes to the

demyelination of nerve cells (Bonneick et al., 2005; Previtali et al., 2007). Intriguingly, PtdSer is

the main component of myelin sheath (Ma et al., 2022), and inhibition of PI4KA in Schwann

cells reduces PM and total levels of PtdSer and PI4P, and induces aberrant myelination

(Alvarez-Prats et al., 2018). Thus, in addition to disrupted endosomal trafficking by *MTMR2*

inactivation, it is plausible that the reduced PM levels of PtdSer and PI4P further contributes to

the demyelination observed in Charcot-Marie-Tooth disease type 4B1.

We have demonstrated that the protein expression of MTMR 2/3/4/7 are interdependent,

whence single KD of *MTMR2, 3, 4* or *7* concomitantly reduces the protein levels of MTMR

2/3/4/7 (Fig. 6A), which likely accounts for the observation that a loss of *MTMR2, 3, 4 or 7* is

sufficient to elevate cellular PI3P level despite the presence of 9 catalytically active MTMR

members (Cao et al., 2008; Lahiri et al., 2015; Ma et al., 2008; Zhao et al., 2019). Also, while

overexpression of WT MTMR3 in *MTMR3* KD cells restores endogenous protein levels of
MTMR 2/4/7, overexpression of WT MTMR 2/4/7 does not restore endogenous MTMR3 protein
level (Fig. 6B and C). These data are consistent with our rescue experiments, in which
overexpression of WT MTMR3, but not the other MTMR members, in *MTMR3* KD cells restores
the KD effects of *MTMR3*. For the interdependence of MTMR 2/3/4/7 protein expression, one
plausible model is that the catalytically active and inactive MTMR members form one large,
interconnected machinery, which may regulate the subcellular localization of MTMR members
for maintaining cellular PI3P metabolism. The 9 enzymatically active MTMR members need to
form homo- or heterodimers with the 7 inactive members via its coiled-coil domain for their
enzymatic activity and subcellular localization (Hnia et al., 2012; Wang et al., 2024). MTMR3
and 4 form homodimers and heterodimers with each other (Lorenzo et al., 2006). MTMR2 form
homodimers and heterodimers with inactive members, MTMR5, 12 and 13 (Kim et al., 2003;
Nandurkar et al., 2003; Robinson and Dixon, 2005), while MTMR7 heterodimerizes with the
inactive MTMR9 (Mochizuki and Majerus, 2003). Also, MTMR2 promotes MTMR13 membrane
localization and they depend on each other for maintaining their protein expression (Ng et al.,
2013; Robinson and Dixon, 2005). Thus, the loss of a single MTMR member could perturb the
subcellular localization of its partner MTMR members, resulting in their protein degradation. This
in turn, disrupts the interconnected network of MTMR members, resulting in the subcellular
mislocalization and degradation of multiple MTMR members, and thereby perturbed PI3P
metabolism. A systematic analysis of subcellular localization and protein expression of
catalytically active and inactive MTMR members after genetical depletion of single MTMR
member will provide deeper insights into the interdependence of MTMR members.
We have observed that *MTMR 2/3/4/7* KD reduces oncogenic KRAS protein expression
(Fig. 7E). Previous studies have reported that *MTMR* KD stimulates autophagy via elevating
PI3P content in the phagophore, a double-membrane structure that matures into
autophagosome, which recruits autophagy-related proteins for autophagosome biogenesis
(Nishimura and Tooze, 2020; Wang et al., 2024). Also, KRAS degradation occurs via the
autophagy/lysosome-mediated mechanism, in which *ADAM9* loss or inhibition enhances the
interaction between KRAS and plasminogen activator inhibitor 1 (PAI-1), which acts as a
specific autophagy receptor binding to light chain 3 (LC3) for stimulating lysosomal degradation
of KRAS (Huang et al., 2024). Together, our data suggest that the PM dissociated KRAS
undergoes its degradation by enhanced autophagy induced by *MTMR* KD. However, this may
only partially account for our observation. Since *MTMR 2/3/4/7* KD induces PM dissociation of
KRAS without the G-domain (Fig. S2), it is likely that the KD induces the PM dissociation of WT

KRAS expressed in BxPC-3, which was not degraded upon *MTMR 2/3/4/7* KD (Fig. 7E). Thus,

the KRAS degradation upon *MTMR 2/3/4/7* KD is unlikely induced by KRAS PM dissociation.

Also, while oncogenic KRAS/MAPK signaling promotes autophagy in PDAC (Bryant et al., 2019),

we observed *MTMR 2/3/4/7* KD reduces the MAPK/c-MYC signaling in PDAC harboring

oncogenic KRAS. Thus, the inhibition of MAPK/c-MYC signaling by *MTMR* KD would block

autophagy, but promote it. Deeper insights for KRAS degradation in terms of oncogenic KRAS

signaling and cellular PI3P content need to be elucidated.

In sum, our study proposes a new role for the MTMR phosphatases in maintaining PM

PI4P and PtdSer contents by maintaining an adequate supply of PM PI from PI3P. Depleting

MTMR 2/3/4/7 in turn, reduces PM and total cellular PtdSer levels, blocking KRAS PM

localization and KRAS signaling. Thus, the mechanisms that maintain PM PI content may

contain useful targets to abrogate oncogenic KRAS signaling. Our study further proposes an

interdependent network of MTMR members for maintaining their protein expression and their

phosphatase activity.

Conflict of interest statement

The authors declare no competing financial interests.

Acknowledgements

This work was supported by Wright State University Seed Grant and the National Cancer Institute [R00-CA188593 to K.-J.C.], National Institute of General Medical Sciences [2R35GM119412 to G.R.V. Hammond] and Cancer Prevention and Research Institute of Texas [CPRIT RP200047 to JFH]. We would like to thank Karen Henkels for her contribution to this work.

Authors contribution and ORCID IDs

- Taylor E. Lange (0000-0002-5172-3220) performed investigation and validation.
- Ali Naji (   ) performed investigation and validation.
- Ransome van der Hoeven (   ) performed investigation data analysis.
- Hong liang (0009-0004-1274-4661) performed investigation.
- Yong Zhou (0000-0002-0214-8151) performed investigation, validation and data analysis.
- Gerald R.V. Hammond (0000-0002-6660-3272) performed editing and review the manuscript.
- John F. Hancock (0000-0003-0542-4710) performed funding acquisition and edit/review the manuscript.
- Kwang-jin Cho (0000-0002-2234-9292) performed conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, validation, visualization, and writing the draft and editing.

	Figure legend

	Table 1. 8 novel genes, which encode proteins with phosphatase activity are identified to
	regulate KRAS/PM localization.
PI – phosphatidylinositol, Ins – inositol, MTMR – myotubularin-related protein, MINPP1 –
multiple inositol-polyphosphate phosphatase 1, TMEM55A – transmembrane protein 55A,
INPPL1 - inositol polyphosphate phosphatase Like 1, SHIP2 – SH2-containing
phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase, PPP2R5B - protein phosphatase 2
regulatory subunit β

Figure 1. MTMR 2/3/4/7 regulate the PM localization of KRASG12V. (**A**) T47D cells stably
expressing GFP-KRASG12V were infected with lentivirus expressing scrambled shRNA (SCR)
or shRNA targeting *MTMR2, 3, 4* or *7*, followed by 1 ug/mL puromycin selection. Cell lysates
were prepared and immunoblotted with anti-MTMR2, 3 and 7 antibodies. Actin blots are shown
as loading controls. For MTMR4, cDNA was amplified with primers specific for *MTMR4* exons 1
and 5, or *GAPDH* exons 2 and 3 as a loading control. (**B**) T47D cells co-expressing shRNA
targeting *MTMR 2/3/4/7* with GFP-KRASG12V or -HRASG12V were incubated with CellMask
for 1 h at 37 °C incubator, fixed with 4% PFA and imaged by confocal microscopy.
Representative GFP-KRASG12V and -HRASG12V images are shown. The corresponding
CellMask and merged images for GFP-KRASG12V are shown in Fig. S1A. The inserted values
represent a mean fraction ± S.E.M. of CellMask colocalizing with GFP-KRASG12V calculated
by Manders' coefficient from three independent experiments. Scale bar – 10 μm. (**C** and **D**)
Intact basal PM sheets prepared from T47D cells co-expressing GFP-KRASG12V or -
HRASG12V and shRNA targeting *MTMR2, 3, 4* or *7* were labeled with anti-GFP-conjugated
gold particles and visualized by EM. Representative EM images are shown in Fig. S1B. The
graphs show a mean number of gold particles ± SEM (n ≥ 15). (**D** and **E**) Spatial mapping of the
same gold-labeled PM sheets was performed. The peak values, L_{max}, representing the extent of
KRAS and HRAS spatial organization, are shown as bar graphs (n ≥ 15). Significant differences
between control (SCR-expressing) and *MTMR*-silenced cells were assessed by one-way
ANOVA tests for (B – D) and bootstrap tests for (E and F) (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,
**** $p < 0.0001$).

Figure 2. MTMR 2/3/4/7 regulate the cellular level and distribution of PtdSer. (**A**) T47D cells
stably expressing GFP-LactC2 were infected with lentivirus expressing scrambled shRNA (SCR)
or shRNA targeting *MTMR2, 3, 4* or *7*, followed by 1 ug/mL puromycin selection. Cells were

incubated with CellMask for 1 h at 37 $^{\circ}$C incubator, fixed with 4% PFA and imaged by confocal
microscopy. Representative GFP-LactC2 images are shown. The inserted values represent a
mean fraction $\pm$ S.E.M. of CellMask colocalizing with GFP-LactC2 calculated by Manders'
coefficient from three independent experiments. Scale bar – 10 μm. (**B**) Intact basal PM sheets
prepared from T47D cells co-expressing GFP-LactC2 and shRNA targeting *MTMR2, 3, 4* or *7*
were labeled with anti-GFP-conjugated gold particles and visualized by EM. The graphs show a
mean number of gold particles $\pm$ SEM (n $\geq$ 15). (**C**) *MTMR 2/3/4/7* KD and stably expressing
GFP-KRASG12V T47D cells were incubated with Cy5-conjugated annexin V and annexin V-
positive cells were counted by flow cytometry. Scrambled (SCR) shRNA-expressing cells were
treated with 1 μM staurosporine (STS) for 6 h to induce apoptosis. The graph represents a
mean (%) $\pm$ S.E.M. of annexin V-positive cells from three independent experiments. (**D**) Whole
cell PtdSer levels were measured in these cells via electron spray ionization and MS/MS from
three independent experiments. A heat map was constructed to quantify the changes of
different lipid species after *MTMR* knockdown. Each row represents a different species of
PtdSer, while each column represents a single sample. The scaled expression values of each
lipid measured is plotted in red-blue color $\log_2$ scale. In relation to control (SCR) cells, red and
blue colored tiles indicate higher and lower lipid contents, respectively. (**E**) The graph shows the
mean of total moles $\pm$ S.E.M. of PtdSer. Significant differences between control (SCR-
expressing) and *MTMR*-silenced cells were assessed using one-way ANOVA tests (* $p < 0.05$,
** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, # – not significant).

Figure 3. MTMR 2/3/4/7 regulate the localization of PM PI4P and ORP5. (**A**) WT T47D cells
were infected with lentivirus expressing scrambled shRNA (SCR) or shRNA targeting *MTMR2, 3,*
4 or *7*, followed by 1 ug/mL puromycin selection. These cells were transfected with GFP-P4M-
SidM, -PH-PLCδ1, or -ORP5, and fixed with 4% PFA and imaged by confocal microscopy.
Closed and open arrowheads for GFP-P4M-SidM indicate the staining of PM and Golgi complex,
respectively. Closed arrowheads for GFP-ORP5 indicate the ER localization of ORP5. Scale bar
– 10 μm. Intact basal PM sheet of Caco-2 cells co-expressing GFP-P4M-SidM (**B**) or -PH-
PLCδ1 (**C**) with shRNA targeting *MTMR2, 3, 4,* or *7* were labeled with anti-GFP-conjugated gold
particles and visualized by EM. The graphs show a mean number of gold particles $\pm$ SEM (n $\geq$
15). Significant differences between control (SCR-expressing) and *MTMR*-silenced cells were
assessed by one-way ANOVA tests (* $p < 0.05$, **** $p < 0.0001$, # – not significant).

Figure 4. MTMR 2/3/4/7 regulate PM PI3P contents. (**A**) T47D cells expressing scrambled shRNA (SCR) or shRNA targeting *MTMR2, 3, 4* or *7*, followed by 1 ug/mL puromycin selection, were overexpressed with GFP-2xFYVE. Cells were incubated with CellMask for 1 h at 37 ^{o}C incubator, fixed with 4% PFA and imaged by confocal microscopy. Representative GFP-2xFYVE images are shown. Their corresponding CellMask and merged images are shown in Fig. S3. A selected region (the white square) is shown at a higher magnification. Arrowheads indicate the PM-staining of GFP-2xFYVE. (**B**) The graph represents a mean fraction ± S.E.M (n = 13) of CellMask colocalizing with GFP-2xFYVE calculated by Manders' coefficient. (**C**) Acidic lipids were extracted from *MTMR 2/3/4/7* KD T47D cells, and total PI3P was measured by a quantitative ELISA. The graph shows a relative mean total PI3P level ± S.E.M from three independent experiments. (**D**) Intact basal PM sheets prepared from these T47D cells from (A), or WT T47D cells expressing GFP-MTMR members (**E**) were labeled with anti-GFP-conjugated gold particles and visualized by EM. The graphs show a mean number of gold particles ± SEM (n ≥ 15). (**F**) Cells from (E) were fixed with 4% PFA and imaged by confocal microscopy. Arrowheads indicate the PM localization of GFP-MTMR members. Scale bar – 10 μm. Significant differences between control (SCR-expressing) and *MTMR*-silenced cells were assessed by one-way ANOVA tests for (B – D) (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, # - not significant). Significant differences between untagged GFP (control) and GFP-MTMR-expressing cells were assessed by one-way ANOVA test for (E) (** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

Figure 5. Ectopic expression of WT MTMR3 restores the effects of *MTMR3* loss. T47D cells co-expressing *MTMR3* shRNA17 and GFP-KRASG12V (**A**) or -LactC2 (**B**) were infected with lentivirus expressing indicated mCherry-MTMR proteins, fixed with 4% PFA and imaged by confocal microscopy. Scale bar – 10 μm. Asterisks indicate the restored PM localization of GFP-KRASG12V or -LactC2. Intact basal PM sheets were prepared from T47D cells co-expressing *MTMR3* shRNA17 and GFP-KRASG12V (**C**) or -LactC2 (**D**), -P4M-SidM (**E**), or -2xFYVE (**F**) were infected with lentivirus expressing indicated mCherry-MTMR members, and labeled with anti-GFP-conjugated gold particles and visualized by EM. The graphs show a mean number of gold particles ± S.E.M. (n = 10). (**G**) Cell lysates from T47D cells co-expressing GFP-KRASG12V, *MTMR3* shRNA17 and mCherry-MTMR members were immunoblotted with anti-RFP antibody. Arrowheads represents the expression of indicated mCherry-MTMR members. An actin blot is shown as a loading control. Representative blots are shown from three independent experiments. Significant differences between control (*MTMR3* KD and untagged

mCherry-expressing) and other cells were assessed by one-way ANOVA tests for (C – F) (* p <
0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001, # - not significant).

Figure 6. Protein expression of MTMR 2/3/4/7 is interdependent. (**A**) Cell lysates from T47D
cells co-expressing GFP-KRASG12V and indicated *MTMR*-targeting shRNA were
immunoblotted for endogenous MTMR2, 3 and 7. (**B**) Cell lysates from T47D cells co-
expressing GFP-KRASG12V and *MTMR3* shRNA17, and infected with lentivirus expressing
indicated mCherry-MTMR members were prepared and immunoblotted for endogenous MTMR3.
The membrane was then stripped and re-blotted with anti-RFP antibody. Arrowheads represent
the expression of indicated mCherry-tagged MTMR members. (**C**) Cell lysates from (B) were
blotted for endogenous MTMR2 and 7. Actin blots are shown as loading controls.
Representative blots are shown. Values indicate mean MTMR member ± S.E.M relative to SCR-
expressing cells from three independent experiments. Significant differences between control
(*MTMR3* KD and untagged mCherry-expressing) and other cells were assessed by one-way
ANOVA tests (* p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001, # - not significant).

Figure 7. MTMR 2/3/4/7 regulate the oncogenic KRAS signal output. (**A**) RNAi was induced
by feeding *let-60* (n1046) L1 worms through adult stage with *E. coli* strain HT115, producing
dsRNA to target genes. The presence of the multi-vulva phenotype was scored using
DIC/Nomarski microscopy. 100 – 200 worms were assayed per RNAi knockdown. T47D cells
stably expressing (**B**) GFP-KRASG12V or (**C**) -HRASG12V, or (**D**) WT BxPC3, MIA PaCa-2 and
PANC-1 cells were infected with lentivirus expressing scrambled shRNA (SCR) or shRNA
targeting *MTMR 2/3/4/7*, followed by 1 µg/mL puromycin selection. Cells were seeded on a 96-
well plate and cultured for 4 days. Complete growth medium was replaced every 24 h, and cell
numbers were counted to measure cell proliferation. Graphs show the mean cell proliferation ±
S.E.M. relative to that for the control cells (SCR-expressing) from three independent
experiments. (**E**) Cell lysates prepared from (D) were immunoblotted for ppERK, pAkt (S473), c-
MYC and endogenous KRAS. Values indicate the mean ppERK, pAkt, c-MYC or endogenous
KRAS ± S.E.M relative to the control cells (SCR-expressing) from three independent
experiments. Total ERK, Akt and actin blots were used as loading controls. Representative blots
are shown. Significant differences between control (SCR-expressing) and *MTMR* KD cells were
assessed using one-way ANOVA tests (* p<0.05, ** p<0.01, *** p<0.001, # - not significant).

Figure 8. A working model of our study. PITPs deliver newly synthesized PI from the ER to PM, and MTMR at the PM concomitantly converts PI3P to PI, which are which are crucial substrate for PM PPIn like PI4P. ORP5 and 8, the lipid transfer proteins for PtdSer, exchanges ER PtdSer with PM PI4P at ER-PM membrane-contacting sites, which enriches PM PtdSer contents. This in turn, allows stable PM binding of KRAS, and thereby KRAS signal output. When MTMR is depleted, PM PI3P conversion to PI is blocked, providing insufficient PM PI contents for PI4P synthesis. This dissociates ORP5/8 from ER-PM membrane-contacting sites, resulting in reduced PM localization of PtdSer and thereby KRAS. PITPs – PI transfer proteins, PI4KA – phosphatidylinositol 4-kinase IIIα, ORP – oxysterol-binding protein related protein, PM – plasma membrane, ER – endoplasmic reticulum.

Supplement Figure 1. MTMR 2/3/4/7 regulate the PM localization of KRASG12V. (**A**) T47D cells stably expressing GFP-KRASG12V were infected with lentivirus expressing scrambled shRNA (SCR) or shRNA targeting *MTMR2, 3, 4* or *7*, followed by 1 ug/mL puromycin selection. Cells were incubated with CellMask for 1 h at 37 ^{o}C incubator, fixed with 4% PFA and imaged by confocal microscopy. The inserted values represent a mean fraction ± S.E.M. of CellMask colocalizing with GFP-KRASG12V calculated by Manders' coefficient from three independent experiments. Scale bar – 10 μm. (**B**) Intact basal PM sheets prepared from T47D cells co-expressing GFP-KRASG12V and shRNA targeting *MTMR2, 3, 4* or *7* were labeled with anti-GFP-conjugated gold particles and visualized by EM. Representative EM images are shown. Scale bar – 0.1 μm. Significant differences between control (SCR-expressing) and *MTMR*-silenced cells were assessed by one-way ANOVA tests (* $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$).

Supplement Figure 2. MT*MR 2/3/4/7* KD disrupts the PM localization and nanoclustering of CTK. (**A**) MCF-7 cells stably co-expressing GFP-CTK (the C-terminal membrane anchor of KRAS without the G-domain) and shRNA targeting MT*MR2, 3, 4 or 7* were fixed with 4% PFA and imaged by confocal microscopy. (**B**) Intact basal PM sheets of cells from (A) were prepared, labeled with anti-GFP-conjugated gold particles and visualized by EM. Scale bar – 10 μm. Significant differences between control (SCR-expressing) and *MTMR*-silenced cells were assessed by one-way ANOVA tests (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

Supplement Figure 3. MTMR 2/3/4/7 regulate cellular PI3P contents. T47D cells expressing scrambled shRNA (SCR) or shRNA targeting *MTMR2, 3, 4* or *7*, followed by 1 ug/mL puromycin

selection, were overexpressed with GFP-2xFYVE. Cells were incubated with CellMask for 1 h at
37°C incubator, fixed with 4% PFA and imaged by confocal microscopy. A selected region (the
white square) is shown at a higher magnification. Arrowheads indicate the PM-staining of GFP-
2xFYVE. Scale bar – 10 μm.

Supplement Figure 4. Ectopic expression of WT MTMR4 restores the effects of *MTMR4*
loss. T47D cells co-expressing *MTMR4* shRNA66 and GFP-KRASG12V (**A**) or -LactC2 (**B**)
were infected with lentivirus expressing indicated mCherry-MTMR proteins, fixed with 4% PFA
and imaged by confocal microscopy. Scale bar – 10 μm. Asterisks indicate the restored PM
localization of GFP-KRASG12V or -LactC2. Intact basal PM sheets were prepared from T47D
cells co-expressing *MTMR4* shRNA66 and GFP-KRASG12V (**C**) or -LactC2 (**D**), -P4M-SidM (**E**),
or -2xFYVE (**F**) were infected with lentivirus expressing indicated mCherry-MTMR members,
and labeled with anti-GFP-conjugated gold particles and visualized by EM. The graphs show a
mean number of gold particles ± S.E.M. (n = 10). Significant differences between control
(*MTMR4* KD and untagged mCherry-expressing) cells and other cells were assessed by one-
way ANOVA tests for (C – F) (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, # - not significant).

Reference

Alvarez-Prats, A., I. Bjelobaba, Z. Aldworth, T. Baba, D. Abebe, Y.J. Kim, S.S. Stojilkovic, M. Stopfer, and T. Balla. 2018. Schwann-Cell-Specific Deletion of Phosphatidylinositol 4-Kinase Alpha Causes Aberrant Myelination. *Cell Rep.* 23:2881-2890.

Ashlin, T.G., N.J. Blunsom, and S. Cockcroft. 2021. Courier service for phosphatidylinositol: PITPs deliver on demand. *Biochim Biophys Acta Mol Cell Biol Lipids.* 1866:158985.

Blunsom, N.J., and S. Cockcroft. 2020. Phosphatidylinositol synthesis at the endoplasmic reticulum. *Biochim Biophys Acta Mol Cell Biol Lipids.* 1865:158471.

Bonneick, S., M. Boentert, P. Berger, S. Atanasoski, N. Mantei, C. Wessig, K.V. Toyka, P. Young, and U. Suter. 2005. An animal model for Charcot-Marie-Tooth disease type 4B1. *Hum Mol Genet.* 14:3685-3695.

Bryant, K.L., C.A. Stalnecker, D. Zeitouni, J.E. Klomp, S. Peng, A.P. Tikunov, V. Gunda, M. Pierobon, A.M. Waters, S.D. George, G. Tomar, B. Papke, G.A. Hobbs, L. Yan, T.K. Hayes, J.N. Diehl, G.D. Goode, N.V. Chaika, Y. Wang, G.F. Zhang, A.K. Witkiewicz, E.S. Knudsen, E.F. Petricoin, 3rd, P.K. Singh, J.M. Macdonald, N.L. Tran, C.A. Lyssiotis, H. Ying, A.C. Kimmelman, A.D. Cox, and C.J. Der. 2019. Combination of ERK and autophagy inhibition as a treatment approach for pancreatic cancer. *Nat Med.* 25:628-640.

Cao, C., J.M. Backer, J. Laporte, E.J. Bedrick, and A. Wandinger-Ness. 2008. Sequential actions of myotubularin lipid phosphatases regulate endosomal PI(3)P and growth factor receptor trafficking. *Mol Biol Cell.* 19:3334-3346.

Chandra, A., H.E. Grecco, V. Pisupati, D. Perera, L. Cassidy, F. Skoulidis, S.A. Ismail, C. Hedberg, M. Hanzal-Bayer, A.R. Venkitaraman, A. Wittinghofer, and P.I. Bastiaens. 2012. The GDI-like solubilizing factor PDEdelta sustains the spatial organization and signalling of Ras family proteins. *Nat Cell Biol.* 14:148-158.

Chang, C.L., T.S. Hsieh, T.T. Yang, K.G. Rothberg, D.B. Azizoglu, E. Volk, J.C. Liao, and J. Liou. 2013. Feedback regulation of receptor-induced Ca2+ signaling mediated by E-Syt1 and Nir2 at endoplasmic reticulum-plasma membrane junctions. *Cell Rep.* 5:813-825.

Chi, H., X. Yang, P.D. Kingsley, R.J. O'Keefe, J.E. Puzas, R.N. Rosier, S.B. Shears, and P.R. Reynolds. 2000. Targeted deletion of Minpp1 provides new insight into the activity of multiple inositol polyphosphate phosphatase in vivo. *Mol Cell Biol.* 20:6496-6507.

Cho, J., J.S. King, X. Qian, A.J. Harwood, and S.B. Shears. 2008. Dephosphorylation of 2,3-bisphosphoglycerate by MIPP expands the regulatory capacity of the Rapoport-Luebering glycolytic shunt. *Proc Natl Acad Sci U S A.* 105:5998-6003.

Cho, K.J., D.E. Casteel, P. Prakash, L. Tan, D. van der Hoeven, A.A. Salim, C. Kim, R.J. Capon, E. Lacey, S.R. Cunha, A.A. Gorfe, and J.F. Hancock. 2016a. AMPK and Endothelial Nitric Oxide Synthase Signaling Regulates K-Ras Plasma Membrane Interactions via Cyclic GMP-Dependent Protein Kinase 2. *Mol Cell Biol.* 36:3086-3099.

Cho, K.J., and J.F. Hancock. 2013. Ras nanoclusters: a new drug target? *Small GTPases.* 4:57-60.

783   Cho, K.J., M.M. Hill, S. Chigurupati, G. Du, R.G. Parton, and J.F. Hancock. 2011. Therapeutic
784   levels of the hydroxmethylglutaryl-coenzyme A reductase inhibitor lovastatin activate ras
785   signaling via phospholipase D2. *Mol Cell Biol*. 31:1110-1120.

786   Cho, K.J., R.S. Kasai, J.H. Park, S. Chigurupati, S.J. Heidorn, D. van der Hoeven, S.J.
787   Plowman, A. Kusumi, R. Marais, and J.F. Hancock. 2012a. Raf inhibitors target ras
788   spatiotemporal dynamics. *Curr Biol*. 22:945-955.

789   Cho, K.J., J.H. Park, A.M. Piggott, A.A. Salim, A.A. Gorfe, R.G. Parton, R.J. Capon, E. Lacey,
790   and J.F. Hancock. 2012b. Staurosporines disrupt phosphatidylserine trafficking and
791   mislocalize Ras proteins. *J Biol Chem*. 287:43573-43584.

792   Cho, K.J., D. van der Hoeven, and J.F. Hancock. 2013. Inhibitors of K-Ras plasma membrane
793   localization. *Enzymes*. 33 Pt A:249-265.

794   Cho, K.J., D. van der Hoeven, Y. Zhou, M. Maekawa, X. Ma, W. Chen, G.D. Fairn, and J.F.
795   Hancock. 2016b. Inhibition of Acid Sphingomyelinase Depletes Cellular
796   Phosphatidylserine and Mislocalizes K-Ras from the Plasma Membrane. *Mol Cell Biol*.
797   36:363-374.

798   Chung, J., F. Torta, K. Masai, L. Lucast, H. Czapla, L.B. Tanner, P. Narayanaswamy, M.R.
799   Wenk, F. Nakatsu, and P. De Camilli. 2015. INTRACELLULAR TRANSPORT.
800   PI4P/phosphatidylserine countertransport at ORP5- and ORP8-mediated ER-plasma
801   membrane contacts. *Science*. 349:428-432.

802   Diggle, P.J., J. Mateu, and H. E. Clough. 2000. A comparison between parametric and non-
803   parametric approaches to the analysis of replicated spatial point patterns. *Adv. Appl.*
804   *Probab.* 32:331-343.

805   Du, X., L. Zhou, Y.C. Aw, H.Y. Mak, Y. Xu, J. Rae, W. Wang, A. Zadoorian, S.E. Hancock, B.
806   Osborne, X. Chen, J.W. Wu, N. Turner, R.G. Parton, P. Li, and H. Yang. 2020. ORP5
807   localizes to ER-lipid droplet contacts and regulates the level of PI(4)P on lipid droplets. *J*
808   *Cell Biol*. 219.

809   Dyson, J.M., C.J. O'Malley, J. Becanovic, A.D. Munday, M.C. Berndt, I.D. Coghill, H.H.
810   Nandurkar, L.M. Ooms, and C.A. Mitchell. 2001. The SH2-containing inositol
811   polyphosphate 5-phosphatase, SHIP-2, binds filamin and regulates submembraneous
812   actin. *J Cell Biol*. 155:1065-1079.

813   Franklin, N.E., G.S. Taylor, and P.O. Vacratsis. 2011. Endosomal targeting of the
814   phosphoinositide 3-phosphatase MTMR2 is regulated by an N-terminal phosphorylation
815   site. *J Biol Chem*. 286:15841-15853.

816   Garrido, C.M., K.M. Henkels, K.M. Rehl, H. Liang, Y. Zhou, J.U. Gutterman, and K.J. Cho. 2020.
817   Avicin G is a potent sphingomyelinase inhibitor and blocks oncogenic K- and H-Ras
818   signaling. *Sci Rep*. 10:9120.

819   Gerl, M.J., J.L. Sampaio, S. Urban, L. Kalvodova, J.M. Verbavatz, B. Binnington, D. Lindemann,
820   C.A. Lingwood, A. Shevchenko, C. Schroeder, and K. Simons. 2012. Quantitative
821   analysis of the lipidomes of the influenza virus envelope and MDCK cell apical
822   membrane. *J Cell Biol*. 196:213-221.

Ghai, R., X. Du, H. Wang, J. Dong, C. Ferguson, A.J. Brown, R.G. Parton, J.W. Wu, and H. Yang. 2017. ORP5 and ORP8 bind phosphatidylinositol-4, 5-biphosphate (PtdIns(4,5)P 2) and regulate its level at the plasma membrane. *Nat Commun*. 8:757.

Gillooly, D.J., I.C. Morrow, M. Lindsay, R. Gould, N.J. Bryant, J.M. Gaullier, R.G. Parton, and H. Stenmark. 2000. Localization of phosphatidylinositol 3-phosphate in yeast and mammalian cells. *EMBO J*. 19:4577-4588.

Gorfe, A.A., and K.J. Cho. 2019. Approaches to inhibiting oncogenic K-Ras. *Small GTPases*:1-10.

Gulyas, G., G. Radvanszki, R. Matuska, A. Balla, L. Hunyady, T. Balla, and P. Varnai. 2017. Plasma membrane phosphatidylinositol 4-phosphate and 4,5-bisphosphate determine the distribution and function of K-Ras4B but not H-Ras proteins. *J Biol Chem*. 292:18862-18877.

Gutierrez, L., A.I. Magee, C.J. Marshall, and J.F. Hancock. 1989. Post-translational processing of p21ras is two-step and involves carboxyl-methylation and carboxy-terminal proteolysis. *EMBO J*. 8:1093-1098.

Hammond, G.R., M.P. Machner, and T. Balla. 2014. A novel probe for phosphatidylinositol 4-phosphate reveals multiple pools beyond the Golgi. *J Cell Biol*. 205:113-126.

Hancock, J.F., K. Cadwallader, H. Paterson, and C.J. Marshall. 1991. A CAAX or a CAAL motif and a second signal are sufficient for plasma membrane targeting of ras proteins. *Embo J*. 10:4033-4039.

Hancock, J.F., A.I. Magee, J.E. Childs, and C.J. Marshall. 1989. All ras proteins are polyisoprenylated but only some are palmitoylated. *Cell*. 57:1167-1177.

Hancock, J.F., H. Paterson, and C.J. Marshall. 1990. A polybasic domain or palmitoylation is required in addition to the CAAX motif to localize p21ras to the plasma membrane. *Cell*. 63:133-139.

Hancock, J.F., and I.A. Prior. 2005. Electron microscopic imaging of Ras signaling domains. *Methods*. 37:165-172.

Hayes, T.K., N.F. Neel, C. Hu, P. Gautam, M. Chenard, B. Long, M. Aziz, M. Kassner, K.L. Bryant, M. Pierobon, R. Marayati, S. Kher, S.D. George, M. Xu, A. Wang-Gillam, A.A. Samatar, A. Maitra, K. Wennerberg, E.F. Petricoin, 3rd, H.H. Yin, B. Nelkin, A.D. Cox, J.J. Yeh, and C.J. Der. 2016. Long-Term ERK Inhibition in KRAS-Mutant Pancreatic Cancer Is Associated with MYC Degradation and Senescence-like Growth Suppression. *Cancer Cell*. 29:75-89.

Henkels, K.M., K.M. Rehl, and K.J. Cho. 2021. Blocking K-Ras Interaction With the Plasma Membrane Is a Tractable Therapeutic Approach to Inhibit Oncogenic K-Ras Activity. *Front Mol Biosci*. 8:673096.

Hnia, K., I. Vaccari, A. Bolino, and J. Laporte. 2012. Myotubularin phosphoinositide phosphatases: cellular functions and disease pathophysiology. *Trends Mol Med*. 18:317-327.

Hood, F.E., Y.M. Sahraoui, R.E. Jenkins, and I.A. Prior. 2023. Ras protein abundance
correlates with Ras isoform mutation patterns in cancer. *Oncogene*. 42:1224-1232.

Hu, Y., J.M. Wan, and A.C. Yu. 2013. Membrane perforation and recovery dynamics in
microbubble-mediated sonoporation. *Ultrasound Med Biol*. 39:2393-2405.

Huang, Y.K., W.C. Cheng, T.T. Kuo, J.C. Yang, Y.C. Wu, H.H. Wu, C.C. Lo, C.Y. Hsieh, S.C.
Wong, C.H. Lu, W.L. Wu, S.J. Liu, Y.C. Li, C.C. Lin, C.N. Shen, M.C. Hung, J.T. Lin,
C.C. Yeh, and Y.P. Sher. 2024. Inhibition of ADAM9 promotes the selective degradation
of KRAS and sensitizes pancreatic cancers to chemotherapy. *Nat Cancer*. 5:400-419.

Jean, S., and A.A. Kiger. 2014. Classes of phosphoinositide 3-kinases at a glance. *J Cell Sci*.
127:923-928.

Kattan, W.E., W. Chen, X. Ma, T.H. Lan, D. van der Hoeven, R. van der Hoeven, and J.F.
Hancock. 2019. Targeting plasma membrane phosphatidylserine content to inhibit
oncogenic KRAS function. *Life Sci Alliance*. 2.

Kattan, W.E., J. Liu, D. Montufar-Solis, H. Liang, B. Brahmendra Barathi, R. van der Hoeven, Y.
Zhou, and J.F. Hancock. 2021. Components of the phosphatidylserine endoplasmic
reticulum to plasma membrane transport mechanism as targets for KRAS inhibition in
pancreatic cancer. *Proc Natl Acad Sci U S A*. 118.

Kim, S.A., P.O. Vacratsis, R. Firestein, M.L. Cleary, and J.E. Dixon. 2003. Regulation of
myotubularin-related (MTMR)2 phosphatidylinositol phosphatase by MTMR5, a
catalytically inactive phosphatase. *Proc Natl Acad Sci U S A*. 100:4492-4497.

Kim, Y.J., M.L. Guzman-Hernandez, E. Wisniewski, and T. Balla. 2015. Phosphatidylinositol-
Phosphatidic Acid Exchange by Nir2 at ER-PM Contact Sites Maintains
Phosphoinositide Signaling Competence. *Dev Cell*. 33:549-561.

Kovar, S.E., C. Fourman, C. Kinstedt, B. Williams, C. Morris, K.J. Cho, and D.M. Ketcha. 2020.
Chalcones bearing a 3,4,5-trimethoxyphenyl motif are capable of selectively inhibiting
oncogenic K-Ras signaling. *Bioorg Med Chem Lett*. 30:127144.

Lahiri, A., M. Hedl, and C. Abraham. 2015. MTMR3 risk allele enhances innate receptor-induced
signaling and cytokines by decreasing autophagy and increasing caspase-1 activation.
Proc Natl Acad Sci U S A. 112:10461-10466.

Lees, J.A., M. Messa, E.W. Sun, H. Wheeler, F. Torta, M.R. Wenk, P. De Camilli, and K.M.
Reinisch. 2017. Lipid transport by TMEM24 at ER-plasma membrane contacts regulates
pulsatile insulin secretion. *Science*. 355.

Lorenzo, O., S. Urbe, and M.J. Clague. 2005. Analysis of phosphoinositide binding domain
properties within the myotubularin-related protein MTMR3. *J Cell Sci*. 118:2005-2012.

Lorenzo, O., S. Urbe, and M.J. Clague. 2006. Systematic analysis of myotubularins:
heteromeric interactions, subcellular localisation and endosome related functions. *J Cell*
Sci. 119:2953-2959.

Ma, J., F. Zeng, W.T. Ho, L. Teng, Q. Li, X. Fu, and Z.J. Zhao. 2008. Characterization and
functional studies of a FYVE domain-containing phosphatase in C. elegans. *J Cell*
Biochem. 104:1843-1852.

Ma, X., X. Li, W. Wang, M. Zhang, B. Yang, and Z. Miao. 2022. Phosphatidylserine,
inflammation, and central nervous system diseases. *Front Aging Neurosci*. 14:975176.

Maechler, F.A., C. Allier, A. Roux, and C. Tomba. 2019. Curvature-dependent constraints drive
remodeling of epithelia. *J Cell Sci*. 132.

Manders, E.M.M., F.J. Verbeek, and J.A. Aten. 1993. Measurement of co-localization of object
in dual-colour confocal images. *Journal of Microscopy*. 169:375-382.

McCright, B., A.M. Rivers, S. Audlin, and D.M. Virshup. 1996. The B56 family of protein
phosphatase 2A (PP2A) regulatory subunits encodes differentiation-induced
phosphoproteins that target PP2A to both nucleus and cytoplasm. *J Biol Chem*.
271:22081-22089.

Miller, T.E., K.M. Henkels, M. Huddleston, R. Salisbury, S.M. Hussain, A.T. Sasaki, and K.J.
Cho. 2019. Depletion of phosphatidylinositol 4-phosphate at the Golgi translocates K-
Ras to mitochondria. *J Cell Sci*. 132.

Mochizuki, Y., and P.W. Majerus. 2003. Characterization of myotubularin-related protein 7 and
its binding partner, myotubularin-related protein 9. *Proc Natl Acad Sci U S A*. 100:9768-
9773.

Monkemoller, V., C. Oie, W. Hubner, T. Huser, and P. McCourt. 2015. Multimodal super-
resolution optical microscopy visualizes the close connection between membrane and
the cytoskeleton in liver sinusoidal endothelial cell fenestrations. *Sci Rep*. 5:16279.

Monteiro-Cardoso, V.F., L. Rochin, A. Arora, A. Houcine, E. Jaaskelainen, A.M. Kivela, C.
Sauvanet, R. Le Bars, E. Marien, J. Dehairs, J. Neveu, N. El Khallouki, E. Santonico,
J.V. Swinnen, D. Tareste, V.M. Olkkonen, and F. Giordano. 2022. ORP5/8 and
MIB/MICOS link ER-mitochondria and intra-mitochondrial contacts for non-vesicular
transport of phosphatidylserine. *Cell Rep*. 40:111364.

Moser von Filseck, J., A. Copic, V. Delfosse, S. Vanni, C.L. Jackson, W. Bourguet, and G. Drin.
2015. INTRACELLULAR TRANSPORT. Phosphatidylserine transport by ORP/Osh
proteins is driven by phosphatidylinositol 4-phosphate. *Science*. 349:432-436.

Nandurkar, H.H., M. Layton, J. Laporte, C. Selan, L. Corcoran, K.K. Caldwell, Y. Mochizuki,
P.W. Majerus, and C.A. Mitchell. 2003. Identification of myotubularin as the lipid
phosphatase catalytic subunit associated with the 3-phosphatase adapter protein, 3-
PAP. *Proc Natl Acad Sci U S A*. 100:8660-8665.

Naughtin, M.J., D.A. Sheffield, P. Rahman, W.E. Hughes, R. Gurung, J.L. Stow, H.H.
Nandurkar, J.M. Dyson, and C.A. Mitchell. 2010. The myotubularin phosphatase MTMR4
regulates sorting from early endosomes. *J Cell Sci*. 123:3071-3083.

Ng, A.A., A.M. Logan, E.J. Schmidt, and F.L. Robinson. 2013. The CMT4B disease-causing
phosphatases Mtmr2 and Mtmr13 localize to the Schwann cell cytoplasm and

endomembrane compartments, where they depend upon each other to achieve wild-type
levels of protein expression. *Hum Mol Genet.* 22:1493-1506.

Nishimura, T., and S.A. Tooze. 2020. Emerging roles of ATG proteins and membrane lipids in
autophagosome formation. *Cell Discov.* 6:32.

Park, S., Y.K. Joo, and Y. Chen. 2019. Versatile and High-throughput Force Measurement
Platform for Dorsal Cell Mechanics. *Sci Rep.* 9:13286.

Pemberton, J.G., Y.J. Kim, J. Humpolickova, A. Eisenreichova, N. Sengupta, D.J. Toth, E.
Boura, and T. Balla. 2020. Defining the subcellular distribution and metabolic channeling
of phosphatidylinositol. *J Cell Biol.* 219.

Pham, H.Q., K. Yoshioka, H. Mohri, H. Nakata, S. Aki, K. Ishimaru, N. Takuwa, and Y. Takuwa.
2018. MTMR4, a phosphoinositide-specific 3'-phosphatase, regulates TFEB activity and
the endocytic and autophagic pathways. *Genes Cells.*

Plowman, S.J., and J.F. Hancock. 2005. Ras signaling from plasma membrane and
endomembrane microdomains. *Biochim Biophys Acta.* 1746:274-283.

Posor, Y., W. Jang, and V. Haucke. 2022. Phosphoinositides as membrane organizers. *Nat Rev*
Mol Cell Biol. 23:797-816.

Previtali, S.C., A. Quattrini, and A. Bolino. 2007. Charcot-Marie-Tooth type 4B demyelinating
neuropathy: deciphering the role of MTMR phosphatases. *Expert Rev Mol Med.* 9:1-16.

Prior, I.A., F.E. Hood, and J.L. Hartley. 2020. The frequency of Ras mutations in cancer. *Cancer*
Res.

Prior, I.A., C. Muncke, R.G. Parton, and J.F. Hancock. 2003a. Direct visualization of Ras
proteins in spatially distinct cell surface microdomains. *J Cell Biol.* 160:165-170.

Prior, I.A., R.G. Parton, and J.F. Hancock. 2003b. Observing cell surface signaling domains
using electron microscopy. *Sci STKE.* 2003:PL9.

Pulido, R., A.W. Stoker, and W.J. Hendriks. 2013. PTPs emerge as PIPs: protein tyrosine
phosphatases with lipid-phosphatase activities in human disease. *Hum Mol Genet.*
22:R66-76.

Rehl, K.M., J. Selvakumar, R.L. Pitsch, D. Hoang, K. Arumugam, S.W. Harshman, A.A. Gorfe,
and K.J. Cho. 2023. A new ferrocene derivative blocks K-Ras localization and function
by oxidative modification at His95. *Life Sci Alliance.* 6.

Reiner, D.J., V. Gonzalez-Perez, C.J. Der, and A.D. Cox. 2008. Use of Caenorhabditis elegans
to evaluate inhibitors of Ras function in vivo. *Methods Enzymol.* 439:425-449.

Ripley, B.D. 1977. Modelling spatial patterns. *J. R. Statist. Soc. B.* 39:172-192.

Robinson, F.L., and J.E. Dixon. 2005. The phosphoinositide-3-phosphatase MTMR2 associates
with MTMR13, a membrane-associated pseudophosphatase also mutated in type 4B
Charcot-Marie-Tooth disease. *J Biol Chem.* 280:31699-31707.

Robinson, F.L., and J.E. Dixon. 2006. Myotubularin phosphatases: policing 3-phosphoinositides.
Trends Cell Biol. 16:403-412.

Rodgers, J.T., R.O. Vogel, and P. Puigserver. 2011. Clk2 and B56beta mediate insulin-
regulated assembly of the PP2A phosphatase holoenzyme complex on Akt. *Mol Cell*.
41:471-479.

Sampaio, J.L., M.J. Gerl, C. Klose, C.S. Ejsing, H. Beug, K. Simons, and A. Shevchenko. 2011.
Membrane lipidome of an epithelial cell line. *Proc Natl Acad Sci U S A*. 108:1903-1907.

Schmick, M., N. Vartak, B. Papke, M. Kovacevic, D.C. Truxius, L. Rossmannek, and P.I.
Bastiaens. 2014. KRas localizes to the plasma membrane by spatial cycles of
solubilization, trapping and vesicular transport. *Cell*. 157:459-471.

Sears, R., G. Leone, J. DeGregori, and J.R. Nevins. 1999. Ras enhances Myc protein stability.
Mol Cell. 3:169-179.

Smith, M.M., and D.J. Levitan. 2004. The Caenorhabditis elegans homolog of the putative
prostate cancer susceptibility gene ELAC2, hoe-1, plays a role in germline proliferation.
Dev Biol. 266:151-160.

Sohn, M., P. Ivanova, H.A. Brown, D.J. Toth, P. Varnai, Y.J. Kim, and T. Balla. 2016. Lenz-
Majewski mutations in PTDSS1 affect phosphatidylinositol 4-phosphate metabolism at
ER-PM and ER-Golgi junctions. *Proc Natl Acad Sci U S A*. 113:4314-4319.

Sohn, M., M. Korzeniowski, J.P. Zewe, R.C. Wills, G.R.V. Hammond, J. Humpolickova, L. Vrzal,
D. Chalupska, V. Veverka, G.D. Fairn, E. Boura, and T. Balla. 2018. PI(4,5)P2 controls
plasma membrane PI4P and PS levels via ORP5/8 recruitment to ER-PM contact sites.
J Cell Biol. 217:1797-1813.

Tan, L., K.J. Cho, W.E. Kattan, C.M. Garrido, Y. Zhou, P. Neupane, R.J. Capon, and J.F.
Hancock. 2019. Acylpeptide hydrolase is a novel regulator of KRAS plasma membrane
localization and function. *J Cell Sci*. 132.

Tan, L., K.J. Cho, P. Neupane, R.J. Capon, and J.F. Hancock. 2018. An oxanthroquinone
derivative that disrupts RAS plasma membrane localization inhibits cancer cell growth. *J
Biol Chem*. 293:13696-13706.

Taylor, V., M. Wong, C. Brandts, L. Reilly, N.M. Dean, L.M. Cowsert, S. Moodie, and D. Stokoe.
2000. 5' phospholipid phosphatase SHIP-2 causes protein kinase B inactivation and cell
cycle arrest in glioblastoma cells. *Mol Cell Biol*. 20:6860-6871.

Tian, T., A. Harding, K. Inder, S. Plowman, R.G. Parton, and J.F. Hancock. 2007. Plasma
membrane nanoswitches generate high-fidelity Ras signal transduction. *Nat Cell Biol*.
9:905-914.

Ungewickell, A., C. Hugge, M. Kisseleva, S.C. Chang, J. Zou, Y. Feng, E.E. Galyov, M. Wilson,
and P.W. Majerus. 2005. The identification and characterization of two
phosphatidylinositol-4,5-bisphosphate 4-phosphatases. *Proc Natl Acad Sci U S A*.
102:18854-18859.

van der Hoeven, D., K.J. Cho, X. Ma, S. Chigurupati, R.G. Parton, and J.F. Hancock. 2013.
Fendiline inhibits K-Ras plasma membrane localization and blocks K-Ras signal
transmission. *Mol Cell Biol*. 33:237-251.

van der Hoeven, D., K.J. Cho, Y. Zhou, X. Ma, W. Chen, A. Naji, D. Montufar-Solis, Y. Zuo, S.E.
Kovar, K.R. Levental, J.A. Frost, R. van der Hoeven, and J.F. Hancock. 2018.
Sphingomyelin Metabolism Is a Regulator of K-Ras Function. *Mol Cell Biol*. 38.

Varnai, P., and T. Balla. 1998. Visualization of phosphoinositides that bind pleckstrin homology
domains: calcium- and agonist-induced dynamic changes and relationship to myo-
[3H]inositol-labeled phosphoinositide pools. *J Cell Biol*. 143:501-510.

Walker, D.M., S. Urbe, S.K. Dove, D. Tenza, G. Raposo, and M.J. Clague. 2001.
Characterization of MTMR3: an inositol lipid 3-phosphatase with novel substrate
specificity. *Curr Biol*. 11:1600-1605.

Wang, J., W. Guo, Q. Wang, Y. Yang, and X. Sun. 2024. Recent advances of myotubularin-
related (MTMR) protein family in cardiovascular diseases. *Front Cardiovasc Med*.
11:1364604.

Weinberg, F., E. Schulze, C. Fatouros, E. Schmidt, R. Baumeister, and T. Brummer. 2014.
Expression pattern and first functional characterization of riok-1 in Caenorhabditis
elegans. *Gene Expr Patterns*. 15:124-134.

Yeung, T., G.E. Gilbert, J. Shi, J. Silvius, A. Kapus, and S. Grinstein. 2008. Membrane
phosphatidylserine regulates surface charge and protein localization. *Science*. 319:210-
213.

Zewe, J.P., A.M. Miller, S. Sangappa, R.C. Wills, B.D. Goulden, and G.R.V. Hammond. 2020.
Probing the subcellular distribution of phosphatidylinositol reveals a surprising lack at the
plasma membrane. *J Cell Biol*. 219.

Zhao, D., C. Shen, T. Gao, H. Li, Y. Guo, F. Li, C. Liu, Y. Liu, X. Chen, X. Zhang, Y. Wu, Y. Yu,
M. Lin, Y. Yuan, X. Chen, X. Huang, S. Yang, J. Yu, J. Zhang, and B. Zheng. 2019.
Myotubularin related protein 7 is essential for the spermatogonial stem cell homeostasis
via PI3K/AKT signaling. *Cell Cycle*. 18:2800-2813.

Zhou, Y., P. Prakash, H. Liang, K.J. Cho, A.A. Gorfe, and J.F. Hancock. 2017. Lipid-Sorting
Specificity Encoded in K-Ras Membrane Anchor Regulates Signal Output. *Cell*. 168:239-
251 e216.

Zou, J., S.C. Chang, J. Marjanovic, and P.W. Majerus. 2009. MTMR9 increases MTMR6
enzyme activity, stability, and role in apoptosis. *J Biol Chem*. 284:2064-2071.

Table 1

Protein Name	Localization	Target PIs	Ref.
MTMR2	Perinuclear, cytoplasm	PI 3-phosphatase for PI3P and PI(3,5)P$_2$	(Kim et al., 2003), (Cao et al., 2008)
MTMR3	ER, cytoplasm	PI 3-phosphatase for PI3P and PI(3,5)P$_2$	(Walker et al., 2001), (Lorenzo et al., 2005)
MTMR4	Early and recycling endosomes	PI 3-phosphatase for PI3P	(Naughtin et al., 2010), (Pham et al., 2018)
MTMR7	Golgi complex, endosomes, cytoplasm	PI 3-phosphatase for PI3P	(Mochizuki and Majerus, 2003)
MINPP1	ER	3-phosphatase for Ins(1,3,4,5,6)pentakisphosphate (InsP5), inositol hexakisphosphate (InsP6) and 2,3bisphosphoglycerate	(Chi et al., 2000), (Cho et al., 2008)
TMEM55A	Lysosome, late endosome	PI 4-phosphatase for PI(4,5)P$_2$	(Ungewickell et al., 2005)
INPPL1/SHIP2	Cytoplasm, plasma membrane, cytoskeleton, lamellipodia	PI 5-phosphatase for PI(3,4,5)P$_3$	(Taylor et al., 2000), (Dyson et al., 2001)
PPP2R5B	Cytoplasm	A component of the serine/threonine-protein phosphatase 2A complex (PP2A). Mediates PP2A dephosphorylation of Akt.	(McCright et al., 1996), (Rodgers et al., 2011)

Figure 1

A

MTMR2
SCR sh84 sh85

MTMR3
SCR sh15 sh17

MTMR7
SCR sh15 sh62

MTMR4
SCR sh66 sh68

Actin

B

SCR
GFP-KRASG12V
0.43 ± 0.02

MTMR2 sh84
0.55 ± 0.02 *

MTMR2 sh85
0.56 ± 0.03 *

MTMR3 sh15
0.73 ± 0.02 ****

sMTMR3 sh17
0.54 ± 0.02 *

GFP-HRASG12V
0.41 ± 0.02

0.45 ± 0.02 #

0.41 ± 0.02 #

0.41 ± 0.02 #

0.36 ± 0.04 #

MTMR4 sh66
0.67 ± 0.01 ****

MTMR4 sh68
0.62 ± 0.04 ***

MTMR7 sh15
0.78 ± 0.03 ****

MTMR7 sh62
0.74 ± 0.03 ****

0.40 ± 0.03 #

0.36 ± 0.01 #

0.39 ± 0.02 #

0.39 ± 0.01 #

C. GFP-KRASG12V

D. GFP-HRASG12V

E. GFP-KRASG12V

F. GFP-HRASG12V

Figure 2

A

	SCR	MTMR2 sh84	MTMR2 sh85	MTMR3 sh15	MTMR3 sh17
GFP-LactC2	0.46 ± 0.02	0.73 ± 0.05 ***	0.64 ± 0.04 *	0.66 ± 0.05 *	0.63 ± 0.02 *

MTMR4 sh66	MTMR4 sh68	MTMR7 sh15	MTMR7 sh62
0.79 ± 0.03 ****	0.80 ± 0.03 ****	0.67 ± 0.03 **	0.70 ± 0.06 **

Figure 3

A

| | SCR | MTMR2 sh85 | MTMR3 sh17 | MTMR4 sh66 | MTMR7 sh62 |

GFP-P4M-SidM

GFP-PH-PLCδ1

GFP-ORP5

B. GFP-P4M-SidM   C. GFP-PH-PLCδ1

Figure 4

A

GFP-2xFYVE

| SCR | MTMR2 sh85 | MTMR3 sh17 | MTMR4 sh66 | MTMR7 sh62 |

B. Endosomal PI3P

C. Total PI3P

D. GFP-2xFYVE

E. GFP-MTMR

F

MTMR2   MTMR3   MTMR4   MTMR7

Figure 5

Figure 6

A

	SCR	MTMR2 sh85	MTMR3 sh17	MTMR4 sh66	MTMR7 sh62

MTMR2 — 75 kDa
1.00 0.46** 0.88# 0.63* 0.64*
±0.02 ±0.07 ±0.17 ±0.03 ±0.04

MTMR3 — 150 kDa
1.00 0.77# 0.21*** 0.46** 0.49**
±0.10 ±0.05 ±0.03 ±0.10 ±0.09

MTMR7 — 75 kDa
1.00 0.44** 0.54* 0.34** 0.18***
±0.16 ±0.03 ±0.08 ±0.10 ±0.08

Actin — 50 kDa

B

MTMR3 KD + mCherry-MTMR

	SCR	mCherry only	MTMR2 WT	MTMR3 WT	MTMR4 WT	MTMR7 WT

MTMR3 — 250 kDa / 150
1.00 0.12 0.14# 0.58*** 0.20# 0.13#
±0.05 ±0.03 ±0.04 ±0.11 ±0.04 ±0.03
(**** over SCR)

mCherry-MTMR — 250 kDa / 150 / 100

Actin — 50 kDa

C

MTMR3 KD

	SCR	mCherry only	MTMR3 WT

MTMR2 — 75 kDa
1.00** 0.75 0.90*
±0.01 ±0.03 ±0.02

MTMR7 — 75 kDa
1.00** 0.29 0.64*
±0.05 ±0.10 ±0.09

Actin — 50 kDa

Figure 7

A. C. elegans

B. KRASG12V

C. HRASG12V

D. PDAC

E.

Figure 8

Supplement Figure 1

A

	SCR	MTMR2 sh84	MTMR2 sh85	MTMR3 sh15	sMTMR3 sh17
mGFP-K-RasG12V					
Cell Mask					
Merged	0.43 ± 0.02	0.55 ± 0.02*	0.56 ± 0.03*	0.73 ± 0.02****	0.54 ± 0.02*

	MTMR4 sh66	MTMR4 sh68	MTMR7 sh15	MTMR7 sh62
mGFP-K-RasG12V				
Cell Mask				
Merged	0.67 ± 0.01****	0.62 ± 0.04***	0.78 ± 0.03****	0.74 ± 0.03****

B

SCR MTMR2 sh85 MTMR3 sh17 MTMR4 sh66 MTMR7 sh62

Supplement Figure 2

A

SCR   MTMR2 sh85   MTMR3 sh17   MTMR4 sh66   MTMR7 sh62

B

No. of gold particles (µm⁻²)

SCR
MTMR2 sh85 *
MTMR3 sh17 ***
MTMR4 sh66 ***
MTMR7 sh76 **

C

Lmax

SCR
MTMR2 sh85 ***
MTMR3 sh17 ****
MTMR4 sh66 **
MTMR7 sh62 ****

Supplement Figure 3

Supplement Figure 4

A GFP-KRASG12V　mCherry-MTMR　Merged

B GFP-LactC2　mCherry-MTMR　Merged

untagged mCherry

MTMR2 WT

MTMR3 WT

MTMR4 WT

MTMR4 C407S

MTMR7 WT

B. GFP-2xFYVE

C. GFP-KRASG12V

D. GFP-LactC2

E. GFP-P4M-SidM