Title: Identification of a Binding Site for Small Molecule Inhibitors Targeting Human

TRPM4

Authors

Babatunde Ekundayo[1], Prakash Arullampalam[2], Christian E. Gerber[2], Anne-Flore

Hämmerli[2], Sabrina Guichard[2], Mey Boukenna[2], Daniela Ross-Kaschitza[2], Martin Lochner[2],

Jean-Sebastien Rougier[2], Henning Stahlberg[1]*, Hugues Abriel[2]*, and Dongchun Ni[1]

Affiliations

[1] Laboratory of Biological Electron Microscopy, IPHYS, SB, EPFL, and Dept. Fundamental

Microbiology, Faculty of Biology and Medicine, UNIL, Cubotron, Rt. de la Sorge, 1015

Lausanne, Switzerland.

[2] Institute of Biochemistry and Molecular Medicine and Swiss National Centre of

Competence in Research TransCure, University of Bern, Bern, Switzerland

* Corresponding authors: henning.stahlberg@epfl.ch, hugues.abriel@unibe.ch

Abstract:

Transient receptor potential (TRP) melastatin 4 (TRPM4) protein is a calcium-activated

monovalent cation channel associated with various genetic and cardiovascular disorders.

The anthranilic acid derivative NBA is a potent and specific TRPM4 inhibitor, but its binding

site in TRPM4 has been unknown, although this information is crucial for drug development

targeting TRPM4. We determined three cryo-EM structures of full-length human TRPM4

embedded in native lipid nanodiscs without inhibitor, bound to NBA, and a new anthranilic

acid derivative, IBA. We found that the small molecules NBA and IBA were bound in a

pocket formed between the S3, S4, and TRP helices and the S4-S5 linker of TRPM4. Our

structural data and results from patch clamp experiments enable validation of a binding site

for small molecule inhibitors, paving the way for further drug development targeting TRPM4.

26 **Main text**

27 **Introduction**

Transient receptor potential (TRP) ion channels, a superfamily of cation channels, play a

crucial role in various physiological functions, including sensory perception, cellular

homeostasis, and ion transport [1,2]. The potential of TRP channels as therapeutic targets for

drug development is immense, given that mutations in the genes encoding these channels

can lead to their dysfunction or dysregulation, implicating them in numerous diseases and

genetic disorders [3,4]. This underscores the importance of our research in understanding the

role of TRP ion channels, particularly TRPM4, and their potential as therapeutic targets for

drug development.

The TRPM (melastatin-like transient receptor potential) sub-family member, TRPM4, is a

calcium-activated non-selective monovalent cation channel. The voltage-dependent

activation of TRPM4 by intracellular calcium drives a current due to monovalent cations flux

such as Na^+ and K^+ through the channel, leading to plasma membrane depolarization

facilitating, in many cell types, calcium uptake via other calcium-permeable channels [5,6].

TRPM4 activity regulates physiological processes such as cardiac conduction, smooth

muscle contraction, insulin secretion, and immune responses [7–9]. Genetic defects in *TRPM4*

are found in patients with inherited cardiac conduction disorders. Furthermore, alterations of

TRPM4 function are linked to diabetes, hypertension, and cancer [10–16]. Thus far, several

small molecules identified and developed as potent and selective TRPM4 inhibitors are used

as research tools to investigate the potential of TRPM4 as a therapeutic target and are

promising candidates for drug development [17]. These molecules include 9-phenanthrol,

flufenamic acid and the anthranilic acid derivatives CBA (4-chloro-2-[2-(2-chloro-phenoxy)-

acetylamino]-benzoic acid) and NBA (4-chloro-2-(2-(naphthalene-1-yloxy) acetamido)

benzoic acid [18–21]. NBA and CBA are thus far the most selective and potent TRPM4

inhibitors reported; however, it remains unknown how these drugs bind and inhibit TRPM4

activity [19]. To facilitate drug development targeting TRPM4, detailed structural information

revealing the mode of binding these molecules to TRPM4 is essential.

Cryo-electron microscopy (cryo-EM) enabled the structure determination of several TRP

channels. It revealed the druggable sites in these channels by structure determination in the

presence of antagonists and drug molecules [3,17]. These structures have revealed different

drug-binding pockets around the TRP channels' transmembrane domains (TMD). For

example, the synthetic molecule icilin binds into a hydrophobic pocket of the Voltage Sensor

Like Domain (VSLD) in TRPM8 and the binding of the small molecule inhibitor NDNA (*N'*-

(3,4-dimethoxybenzylidene)-2-(naphthalen-1-yl)acetohydrazide) into the cavity between the

S1-S4 domain and pore domain of TRPM5 [22,23]. These two examples revealed different

drug-binding sites within the same subfamily of TRP channels. Interestingly, lipid molecules

can share binding sites with drug molecules and inhibit drug binding and activity, as

demonstrated in the binding of 2-ABP (2-aminoethyl diphenylborinate) into the VBP

(Vallanoid Binding Pocket) of TRPV2, which could be inhibited by binding an endogenous

cholesterol molecule [24].

Structures of detergent-isolated TRPM4 studied by cryo-EM have revealed the binding sites

of the cholesterol homolog CHS (Cholesteryl Hemisuccinate) used to purify TRPM4,

suggesting potential endogenous cholesterol binding sites [25–28]. However, to date, no

structures of TRPM4 have been reported to be bound to specific inhibitors. The addition of

CHS required for stabilizing TRPM4 during detergent isolation could potentially occlude

drug-binding sites, posing a challenge in understanding TRPM4 inhibition. To address this

problem, we have isolated TRPM4 in its native lipid environment using SMA (Styrene Maleic

Anhydride) nanodiscs and determined its high-resolution structure in the presence and

absence of the small molecules NBA and a new derivative IBA (4-chloro-2-[2-(3-

iodophenoxy)-acetylamino]-benzoic acid) which has a similar potency to NBA (**Figure 1;**

Figure 2a-c and Supplementary Fig. 1-8; see method section for IBA synthesis) [29].

These structures reveal the mode of binding and inhibition of two potent and specific TRPM4

inhibitors, paving the way for future drug development and facilitating further investigations

of TRPM4 as a therapeutic target for treating related maladies.

Results

Structures of HsTRPM4 in its native lipid environment

We determined cryo-EM structures of full-length human TRPM4 in endogenous lipid

nanodiscs in apo (HsTRPM4$_{apo}$), IBA (HsTRPM4$_{IBA}$), and NBA (HsTRPM4$_{NBA}$) bound states

at overall resolutions of 3.7, 3.6, and 4.5 Å respectively (**Figure 1; Figure 2;**

Supplementary Fig. 4-8; Supplementary Table 1). All structures obtained were identical to

previously determined cryo-EM structures of TRPM4 in detergent micelles and reconstituted

nanodiscs [25–28]. However, we found that in native nanodiscs, densities for the cytosolic

regions of TRPM4 are poorly resolved compared to previously determined structures,

resulting in lower-resolution 3D reconstructions in these regions, which could be a result of

TRPM4 solubilization using SMA. This effect was more pronounced in the cytosolic N-

terminal TRPM homology regions 1 and 2 (MHR1/ 2), with the density in these regions much

less defined (**Supplementary Fig. 1c; Supplementary Fig. 2a and b**). Therefore, 3D

reconstructions could not be obtained for these regions (**Supplementary Fig. 2; 4- 6**).

These results suggest that SMA solubilization results in a less stable TRPM4 structure,

which could result from the differences in lipid composition or binding compared with

structures of TRPM4 in detergent micelles and reconstituted nanodiscs. Indeed, examination

of the TMD revealed that although the structural arrangement of the transmembrane helices

was identical, the lipid arrangement was strikingly different from previously determined

TRPM4 structures [25–28]. Notably, the published structure of HsTRPM4 in a reconstituted lipid

nanodisc, as well as cryo-EM density maps of detergent-solubilized human and mouse

Trpm4 from this study, revealed ordered densities for annular phospholipids and three CHS

molecules, one of which (CHS-2) occludes the drug binding site (**Supplementary Fig. 2 and**

3; Supplementary Table 2) [26].

On the other hand, in our structures, ordered densities for annular lipids are markedly

reduced, with primarily three densities for cholesterol molecules bound to TRPM4 observed

(**Figure 1a-d; Supplementary Fig. 2a and b**). Of the three, two cholesterol molecules

(CHR-1 and CHR-2) bind in similar positions to CHS molecules in a previous structure in a

reconstituted nanodisc (**Figure 1e-f; Supplementary Fig. 2a and b**). One of these

molecules (CHR-1) interacts with the S6 helix close to the channel pore, and another (CHR-

2) in a pocket formed between S3 and S4 helices and the S4-S5 linker equivalent to the VBP

of TRPV2; however, the density for this cholesterol molecule is weaker compared to CHR-1,

suggesting lower occupancy (**Figure 1e-f**). The third cholesterol molecule (CHR-3) in our

structures binds between the S5 and S2 helix from an adjacent protomer differing from the

previously reported position of CHS at the pre-S1 elbow (**Supplementary Fig. 2a-c**). The

presence of less density for annular lipids in the native nanodiscs was an essential factor

that enabled the assignment of density for the drugs in the HsTRPM4$_{IBA}$ and HsTRPM4$_{NBA}$

structures (**Fig. 2d-i**; **Supplementary Fig. 8**).

The binding site of the small molecule inhibitors NBA and IBA in HsTRPM4

Analysis of the HsTRPM4$_{IBA}$ and HsTRPM4$_{NBA}$ structures revealed discrete densities for the

IBA and NBA molecules located at the periphery of the inner membrane leaflet in the pocket

formed between S3 and S4 helices, S4-S5 linker, and TRP helix (**Figure 2d-f**). The

presence of the ligands in this pocket suggests that these ligands can displace the

endogenous cholesterol molecule (CHR-2) present in the HsTRPM4apo structure, which is

possibly not tightly bound as judged by the weaker density map for this molecule compared

to other bound cholesterol molecules (**Figure 1e and f**). Ligand binding induces only very

subtle conformational changes in the structure of HsTRPM4, with the overall structures

remaining broadly similar (**Figure 2g and h**). The HsTRPM4 ion conduction pore is also in

the closed state in all three structures; however, subtle changes in pore diameter are

observed in the IBA-bound structure, although it remained closed (**Figure 2i**). In this study,

exogenous calcium was not added to the SMA-extracted TRPM4 samples used for cryo-EM

because adding divalent ions destabilizes SMA nanodiscs [29]. However, our cryo-EM density

maps of TRPM4 in native nanodiscs revealed a strikingly apparent density in the calcium-

binding site as observed in structures of detergent-extracted human and mouse TRPM4

supplemented with Ca^{2+}. Indeed, upon fitting a previously determined structure of calcium-

bound TRPM4 into our cryo-EM map of SMA-extracted TRPM4, we observe that the calcium

ion together with its interacting residues Glu828, Gln831, Asp 868, and Arg905 fit very well

into our cryoEM density of SMA extracted TRPM4 similar to cryoEM density maps of calcium

supplemented detergent isolated TRPM4 which were reproduced in this study

(**Supplementary Fig. 9**). Indeed, this suggests the binding of endogenous Ca^{2+} ions in the

SMA extracted TRPM4. Still, we cannot rule out the possibility of other endogenously

derived ions occupying the binding site. Nonetheless, the occupation of the calcium-binding

site in our structures did not result in an open state of the channel like previously described

in calcium-bound TRPM4 structures.

Both IBA and NBA share a common anthranilic acid moiety but differ in that NBA possesses

an additional ring of its naphthalene substituent, making it bulkier and more hydrophobic

than the smaller 3-iodophenyl ring of IBA (**Figure 2e and f, Supplementary Fig. 8**). The

additional ring of the naphthalene substituent of NBA was an important fiducial marker for

the overall placement of the ligands in the density maps. In both HsTRPM4$_{IBA}$ and

HsTRPM4$_{NBA}$ structures, the hydrophilic anthranilic acid moiety containing acidic, amide, and

chloride groups faces towards the cytosol interacting with charged amino acid side chains

from TRPM4 (**Figure 3a and b**). Notably, the anthranilic acid moiety interacts with His908

from the S4 helix, Tyr1057, Gln1061, and Arg1064 from the TRP helix, as well as with

Ser924 from the S4-S5 linker, and Ser863 from the S3 helix in both structures (**Figure 3a**

and b). These interactions between the anthranilic acid moiety and residues of TRPM4

reveal the chemical basis behind the specific binding of the anthranilic acid derivatives to

TRPM4.

NBA and IBA have a similar potency for TRPM4 inhibition, possibly by making additional

unknown interactions in the binding pocket (**Figure 2a-c**). Our structures show that the

naphthalene substituent and the 3-iodophenyl ring of NBA and IBA bind into a hydrophobic

pocket between the S3 and S4 helices of TRPM4 (**Figure 3c and d; Supplementary Fig.**

7d and e), making hydrophobic interactions with the residues that decorate this pocket,

including Val901, Val904 and Leu907 (**Figure 3c and d**). Therefore, these structures reveal

a rationale for the potency and specificity of NBA and IBA and pave the way for designing

chemical modifications of these molecules to enable increased targeting of the binding

pocket to develop more potent and specific inhibitors.

Validation of the NBA and IBA binding site in HsTRPM4

To disrupt the drug-protein interactions in the hydrophilic region of the drug binding site, we

introduced the HsTRPM4 mutations Ser863Ala (S863A), His908Ala (H908A), Ser924Ala

(S924A), Lys925Ala (K925A), Tyr1057Ala (Y1057A), Gln1061Ala (Q1061A), Arg1064Ala

(R1064A), Arg1064Gly (R1064G), Arg1064Ser (R1064S), and the double variant

Ser863Ala/Ser924Ala (S863A/S924A). Additional Val901Trp (V901W) and Val904Trp

(V904W) are space-filling mutations that disrupt the hydrophobic pocket. These mutations

were introduced to reduce the drug's inhibitory effect compared to wildtype HsTRPM4

(**Figure 3a-d**). First, the expression of the different variants was investigated using the

western blot approach (**Supplementary Fig. 10**). Only Ser863Ala and Gln1061Ala variants

showed a significantly reduced expression compared to the wildtype channel

(**Supplementary Fig. 10b**). In parallel, functional experiments revealed that after activation

of the HsTRPM4 function by adding a solution with a concentration of 300 µM of free

calcium, almost all variants displayed a significant reduction in the sodium current of the

calcium-activated HsTRPM4 channel except Ser863Ala, Ser924Ala, Arg1064Ala, and

Arg1064Gly quantify at a membrane voltage of +100 mV (**Figure 3e-f, Supplementary Fig.**

12). Due to the reduction of activation in these mutants and the proximity of these residues

in the NBA binding site to the calcium-binding site (**Figure 4a**), we wondered if this decrease

in function observed may be due to an alteration of the calcium sensitivity of those variants**.**

The calcium sensitivity curves performed on some loss-of-function HsTRPM4 variants

(Val904Trp, Tyr1057Ala, Gln1061Ala, Arg1064Ser, and the double variant Ser863Ala/

Ser924Ala) show that the different EC_{50s} for calcium are not higher compared to wildtype

HsTRPM4 suggesting that this decrease of function is not due to a reduction of calcium

sensitivity **(Figure 4b; Supplementary Fig. 13a)**. On the contrary, the different EC_{50s} of the

variants are smaller than the wildtype EC_{50}, suggesting an increase in calcium sensitivity

(Figure 4b; Supplementary Fig. 13a). Following those experiments and knowing that the

inhibitory effect mediated by NBA will be investigated on these variants in the presence of

calcium (to activate them), control experiments were performed using HsTRPM4 wildtype

channels to explore the potential influence of NBA on calcium sensitivity. As for the variants,

the results suggest that NBA does not decrease the calcium sensitivity of the wildtype

channels (**Figure 4b; Supplementary Fig. 13b**). The absence of a decrease of the calcium

sensitivity in the presence of NBA and amino acid modifications led us to perform the final

experiment investigating the consequences of modifying those amino acids on the inhibitory

effect mediated by NBA. As a proof of concept, the mutations Ser863Ala, Val901Trp,

His908Ala, Ser924Ala, Arg1064Ala, Arg1064Gly, and the double variant Ser863Ala/

Ser924Ala were investigated. Compared to wildtype HsTRPM4, NBA dose-effect curves on

single variants of the HsTRPM4 channel show a drastic increase (up to one log of

difference) of the IC_{50s} for the NBA, suggesting a decreased efficiency of inhibition by NBA

(**Figure 5a**). Compared to the wildtype construct, but also to single amino acid variant

Ser863Ala and Ser924Ala, a more pronounced decrease in the efficiency of NBA of up to

two logs of difference is observed in the presence of the double mutation Ser863Ala/

Ser924Ala (**Fig. 5a,c, and d; Supplementary Figure 14**). In addition, a control experiment

was performed using a mutant of HsTRPM4 linked to cardiac dysfunction, which has already

been reported in the literature: the mutant HsTRPM4 T677I [30]. This variant was selected

based on its presence outside the putative NBA binding pocket and because this point

mutation did not alter the biophysical properties of channels [30]. This mutation did not alter

the IC_{50} of the NBA, as observed with the NBA binding pocket variants (**Figure 5e and f**).

Overall, these data strongly support the notion that those amino acids of HsTRPM4 are part

of the binding pocket for NBA or IBA, as determined by our cryo-EM structures.

Discussion

We have utilized cryo-EM to determine the high-resolution structures of HsTRPM4 bound to specific inhibitors. Importantly, we find that using SMA to isolate HsTRPM4 in its endogenous lipid nanodisc was necessary to assign densities for the inhibitor molecules, which electrophysiology experiments could further validate. This finding supports the relevance of previous attempts to isolate TRPM4 in its native lipid environment but, unfortunately, yielded lower-resolution structures [31]. High-resolution structures of TRPM4 have previously been determined in detergent micelles and in reconstituted lipid nanodiscs, both of which require detergent isolation of the protein from the cell and, by so doing, introduce artifacts to the lipid environment of the protein. An essential factor is that the added CHS, to maintain protein stability in detergent extraction, binds in specific sites in the protein, which may or may not represent actual cholesterol-binding sites. We observed apparent differences in the binding arrangement of annular lipids to TRPM4 from the structural data when isolated with SMA. Most striking is that a CHS molecule completely occupies the drug binding site, suggesting that obtaining a structure or performing biochemical experiments with detergent-CHS isolated protein and these inhibitors does not represent the endogenous scenario. Indeed, identifying the inhibitor binding site was ambiguous in our previous attempts to determine the structure of detergent-isolated TRPM4 in a complex with NBA or IBA. This finding also shows the importance of the composition and binding arrangement of the lipid annulus to the function and inhibition of the target membrane protein. This study shows that using SMA would be advantageous for the structure determination of drugs with membrane proteins, particularly in cases where the drug binds in the TMD exposed to lipid molecules.

During the revision of this work, another paper was published, which revealed the importance of temperature in modulating the structure of detergent-extracted TRPM4. More precisely, at the warm temperature of 37°C, TRPM4 was shown to undergo structural rearrangements, particularly in MHR1/2 domains of the ICD; interestingly, this region is also

flexible and poorly resolved in our structures, which could suggest an interplay of the lipid

environment with the temperature-induced structural movement [32]. Furthermore, the binding

site of the allosteric modulator DVT was also shown to change in a temperature-dependent

manner, leading to channel opening via the relative movement of the TRP helix to S1-S4

helices. This movement of the TRP helix is required for the activation of the TRPM2 channel

and is also required for desensitization of the TRPM8 channel [32,33,34]. These findings reveal

the important role of TRP helix movement in regulating the channel pore of the TRPM

subfamily. Based on our results and these previous studies, we propose that the binding of

NBA or IBA to TRPM4 may restrict the movement of the TRP helix, maintaining the TRPM4

channel pore in a closed state even in the presence of activation stimuli (**Figure 6**). The role

of the bound cholesterol molecule in the same pocket remains unclear. This cholesterol

could also regulate channel function in response to changes in the membrane environment.

A major limitation in our study is that we could not determine structures of SMA-extracted

TRPM4 in the presence of exogenous Ca^{2+} ions, as the addition of calcium destabilizes SMA

nanodiscs. Although we did observe density in the calcium-binding site, we cannot rule out

an effect or interrelation between the binding of Ca^{2+} ions and NBA or IBA. Moreover, we did

not observe decreases in calcium sensitivity in the presence of NBA, but the implications of

which remain unclear (**Figure 4**). However, other published structures of TRPM4 in the

presence and absence of Ca^{2+} ions reveal that the NBA or IBA binding site and residues

involved in the interaction between TRPM4 and NBA or IBA remain accessible for binding of

a CHS molecule in both states suggesting this pocket is accessible for binding of the

inhibitors in both states [26].

Our study has identified a binding pocket for inhibiting TRPM4 by the anthranilic acid

derivatives (**Figure 2d -f**). These drugs bind in a pocket between S3 and S4 helices, S4-S5

linker, and TRP helix. However, detailed insights into the mechanism of inhibition will require

structure determination of TRPM4 in its open state, which has been elusive. Adding Ca^{2+}

ions is insufficient to obtain the open state with purified TRPM4 in lipid nanodiscs. Still, the

presence of a membrane potential may be necessary. In the future, we will seek to address

this by studying TRPM4 in lipid vesicles. Nonetheless, identifying a binding pocket for

TRPM4 inhibitors presents a significant milestone towards further developing new potent

and even more specific drugs targeting the TRPM4 ion channel for therapeutic intervention.

Acknowledgments

This work was supported by the Swiss National Science Foundation (SNF Grants

CRSII5_177195 and the NCCR TransCure (185544)). We thank Emiko Uchikawa, Sergey

Nazarov, and members of the DCI, Lausanne from EPFL, UNIL, and UNIGE for their

support. We also thank Kelvin Lau, Florence Pojer, Laurence Durrer, and Soraya Quinch,

the PTPSP at EPFL, for their support with protein expression.

Author Contributions Statement

HS and HA conceptualized the project. BE and DN performed cryo-EM sample preparation

and structure determination with support from the DCI, Lausanne. PA, AH, SG, DRK, and MB

performed biochemistry and patch clamp experiments. CEG. and ML designed the target

molecule IBA and   synthesis. CEG. performed the synthesis of   NBA   and   IBA.   Project

administration and supervision was done by BE and JSR. BE wrote the original draft.

Manuscript review & editing were done by DN, BE, JSR, ML, HA, and HS. HS and HA acquired

funding. All authors read and approved the manuscript.

Competing Interests Statement

The authors declare no competing interests.

Supplementary Table 1

Data collection and processing	HsTRPM4apo (EMD-19057) (PDB 8RCR)	HsTRPM4$_{IBA}$ (EMD-19061) (8RCU)	HsTRPM4$_{NBA}$ (EMD-19069) (PDB 8RD9) (EMD-19060, local)
Nominal Magnification	165kx	165kx	165kx
Voltage (kV)	300	300	300
Recorded Micrographs	6 429	7 500	11 851
Electron exposure (e–/Å2)	50	50	50
Defocus range (μm)	0.8-2.5	0.8-2.5	0.8-2.5
Pixel size (Å)	0.726	0.726	0.726
Symmetry imposed	C4	C1/C4	C4
Initial particle images (no.)	1 719 323	1 149 090	1 805 182
Final particle images (no.)	16 308	20 345	15 954
Map resolution (Å)	3.67	3.59	4.50
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)	30-3.3	30-3.2	30-3.6
Map sharpening B factor (Å2)	-69.7	-52.2	-108.1
Refinement			
Initial model used (PDB code)	6BQV	6BQV	6BQV

Model composition
  Non-hydrogen atoms
  Protein residues
  Ligands
	8
	2420
	CHR: 12
		9
		2420
		IBA: 4
		CHR: 8
			8
			2422
			NBA: 4
			CHR: 8

B factors (Å^2)
  Protein
  Ligand
	41.23/251.46/135.21
	63.19/103.12/79.00
		0.00/93.44/38.86
		15.44/136.15/47.75
			142.10/427.50/239.20
			20.00/33.04/22.07

R.m.s. deviations
  Bond lengths (Å)
  Bond angles (°)
	0.012 (146)
	0.801 (7)
		0.003 (0)
		0.721 (0)
			0.004 (0)
			0.925 (10)

Validation
  MolProbity score
  Clashscore
  Poor rotamers (%)
	1.93
	14.67
	0.00
		2.04
		16.76
		0.19
			2.23
			24.33
			0.05

Ramachandran plot
  Favored (%)
  Allowed (%)
  Disallowed (%)
	96.16
	3.71
	0.13
		95.49
		4.34
		0.17
			94.82
			4.76
			0.42

294   **Supplementary Table 1 | Cryo-EM data collection, refinement, and validation statistics**

295   **for SMA solubilized HsTRPM4 samples.**

297 **Supplementary Table 2**

Data collection and processing	HsTRPM4+Ca^{2+}	HsTRPM4+NBA+Ca^{2+}	MmTRPM4+Ca^{2+}
Nominal Magnification	96kx	96kx	165kx
Voltage (kV)	300	300	300
Recorded Micrographs	11 877	10 219	8 313
Electron exposure (e–/Å2)	40	40	60
Defocus range (μm)	0.8-2.5	0.8-2.5	0.8-2.5
Pixel size (Å)	0.83	0.83	0.726
Symmetry imposed	C4	C4	C4
Initial particle images (no.)	4 430 540	2 774 028	4 340 237
Final particle images (no.)	128 316	40 061	100 227
Map resolution (Å)	3.04	2.98	2.87
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)	30-2.6	30-2.5	30-2.5

299 **Supplementary Table 2 | Cryo-EM data collection for DDM/CHS solubilized HsTRPM4**

300 **and MmTRPM4 samples.**

Figure 1: Overall structure of HsTRPM4$_{apo}$ in native lipid nanodiscs. (a and b) Two views

of the cryo-EM densities of HsTRPM4$_{apo}$ in surface representation. Alternating

protomers of TRPM4 are shown in blue and white. (c and d) Cartoon representation

of the HsTRPM4$_{apo}$ corresponding to the cryo-EM densities in a and b. The positions

308   of bound cholesterol molecules (CHR) are indicated. The black box indicates the

309   positions of CHR-1 and CHR-3, as shown in yellow. In contrast, the pink box

310   indicates the position of CHR-2. (e) Cryo-EM density in mesh representation for

311   CHR-2 (f) Cryo-EM density in mesh representation for CHR-1. In e and f, the density

312   threshold level is indicated by σ.

Figure 2: Structures of HsTRPM4$_{NBA}$ and HsTRPM4$_{IBA}$. (a) Representative sodium current of the calcium-activated HsTRPM4 wildtype (WT) channel in the presence of a solution with 300 µM of free calcium (grey line) and in the presence of a solution with 300 µM of free calcium with 0.5 µM of NBA (black line). (b) in the presence of a solution with 300 µM of free calcium (grey line) and in the presence of a solution with

300 µM of free calcium with 0.5 µM of IBA (black line). (c) NBA and IBA dose-

response curves of wildtype (WT). (n): number of cells. (d) The structure of inhibitor-

bound HsTRPM4 with box and dashed lines indicating the inhibitor binding site. (e

and f) Zoom-in of IBA and NBA with the cryo-EM density in mesh representation at

two different density threshold levels as indicated by σ. (g and h) Superimposition of

HsTRPM4$_{apo}$ in light grey, HsTRPM4$_{NBA}$ in purple and HsTRPM4$_{IBA}$ in cyan. Two

views of the structures are shown. (i) Pore radius for HsTRPM4$_{apo}$, HsTRPM4$_{NBA}$ and

HsTRPM4$_{IBA}$, all calculated using MOLE. In the top right corner are the pore-forming

helices in HsTRPM4$_{apo}$. The pore profile is shown as a space-filling model (grey).

Figure 3: Amino acid interactions in the drug binding pocket. (a) Amino acid residues of

HsTRPM4 involved in interactions with the anthranilic acid moiety of NBA (in purple) and (b)

IBA (cyan) are shown. (c) Amino acid residues of HsTRPM4 involved in hydrophobic

interactions with NBA (in purple) and (d) IBA (in cyan) are shown. (e) Graphical plot of

normalized recorded sodium current of the calcium-activated HsTRPM4 channel for the

wildtype (WT) and HsTRPM4 variants. (*) represents the p-value < 0.05 compared to the

wildtype condition. At least 5 cells per condition have been recorded. (f and g)

Representative traces of wildtype (WT) and variants sodium currents of the calcium-

activated HsTRPM4 channels: S924A and Q1061A. Sodium currents of the calcium-

activated HsTRPM4 variant channels are generally smaller than WT currents, as depicted in

Figure 3e, due to the amino acid alteration.

Figure 4: Calcium sensitivity of TRPM4 variants. (a) Zoom-in of the calcium-binding site with

the cryo-EM density in mesh representation. The inhibitor binding site is also indicated with a

red arrow. (b) Normalized calcium sensitivity curves for wildtype (WT) HsTRPM4 and loss-

of-function HsTRPM4 variant S863A/S924A, and the effect of a solution containing the

compounds NBA (200 nM) on this calcium sensitivity. (n): number of cells.

Figure 5: Effect of HsTRPM4 variants on NBA inhibition. (a) NBA dose-response curves of wildtype (WT) and variants of the predicted binding pocket of the HsTRPM4 channel, as a function of the decadic logarithm of the NBA concentration. (n): number of cells. (b) Locations of S863 and S924 in the drug binding pocket (c and d) Representative traces of wildtype (WT) and variant HsTRPM4 sodium currents: S863A/S924A in the absence of calcium (dotted black line), in the presence of a solution with a concentration of 300 µM of free calcium (grey line) and in the presence of a solution with a concentration of 300 µM of free calcium and 1 µM of NBA (black line). (e) Representation of the T677I variant located outside of the drug binding pocket of the TRPM4 channel. (f) NBA dose-response curves of wildtype (WT) and a T677I variant. (n): number of cells.

free TRP helix in closed state   locked TRP helix

Ca^{2+}

NBA/IBA

TRP helix

Ca^{2+} Na$^+$

free TRP helix in open state

Figure 6: Proposed inhibition mechanism. The binding of anthranilic acid drugs such as
NBA, CBA, and IBA in the TRPPM4 drug binding site leads to channel blocking, possibly by
restricting the movement of the TRP helix, thereby preventing channel opening upon calcium
activation.

Supplementary Fig. 1: Purification of TRPM4 in SMA-extracted native lipid nanodiscs. (a)

Biochemical workflow for isolating and purifying HsTRPM4 in native lipid nanodiscs using

SMA (SMALP-200). (b) SDS-PAGE gel of purified HsTRPM4 following affinity purification.

(c) Cryo-EM 2D class averages of TRPM4 in SMA-extracted native lipid nanodiscs and

LMNG detergent micelles. White arrows indicate the positions of MHR1/2 in the TRPM4 2D

classes.

Supplementary Fig. 2: Comparison of the annular lipid environment between the cryo-EM

maps of HsTRPM4 in reconstituted lipid nanodiscs [26] (a) and SMA-extracted native lipid

nanodiscs (b). CHS represents cholesteryl hemisuccinate in (a), and CHR represents

cholesterol in (b). (c) The location of CHR-1 and CHR-3 is shown from the cryo-EM structure

of HsTRPM4 in native lipid nanodiscs.

Supplementary Fig. 3: Cryo-EM maps of detergent-solubilized TRPM4 generated in this

study highlight the presence of CHS in the drug binding site when fitted with the published

model (pdb-id; 6BQV). The dashed line indicates the location of the drug-binding site

determined in this study.

Supplementary Fig. 4: Cryo-EM data processing workflow and map resolution (a) The

391   image processing workflow of HsTRPM4$_{apo}$. (b) micrograph and 2D classes (c) Local

392   resolution (d) Model vs map FSC curves. (e) particle direction distribution. (f) FSC curves.

Supplementary Fig. 5: Cryo-EM data processing workflow and 3D reconstructions (a) The

395   image processing workflow of HsTRPM4$_{IBA}$. (b) Micrograph and 2D classes. (c) Local

396   resolution. (d) Model vs map FSC curves. (e) Particle direction distribution. (f) FSC curves.

398 **Supplementary Fig. 6**: Cryo-EM data processing workflow and 3D reconstructions. (a) The

399 image processing workflow of HsTRPM4_NBA. (b) Micrograph and 2D classes. (c) Local

400 resolution. (d) Model vs. map FSC curves. (e) Particle direction distribution. (f) FSC curves.

Supplementary Fig. 7: Zoom-in view of the cryo-EM maps of structural elements HsTRPM4$_{apo}$, HsTRPM4$_{IBA}$ and HsTRPM4$_{NBA}$.

404
Supplementary Fig. 8: Local resolution maps of SMA solubilized HsTRPM4 generated in this

study. (a) Map for the HsTRPM4$_{apo}$ with the density contained in the drug binding site

highlighted. (b) Map for HsTRPM4$_{IBA}$ with the density contained in the drug binding site

highlighted. The chemical structure of IBA is shown on the right. (c) Map for HsTRPM4$_{NBA}$ with

the density contained in the drug binding site highlighted. The chemical structure of the NBA

is shown on the right. (d and e) Zoom-in of NBA and IBA with the cryo-EM density in mesh

representation in different views compared to Figure 2 e and f.

SMA_{solubilized}-human sample without extra Ca²⁺(CHS free)

DDM:CHS_{solubilized}-mouse sample + Ca²⁺

DDM:CHS_{solubilized}-human sample + Ca²⁺

413 **Supplementary Fig. 9**: Identification of cryo-EM densities for Ca^{2+} ions in the maps generated

414 in this study. The published HsTRPM4 model (pdb-id: 6BQV) is fitted into each map. Clear

415 density is observed for Ca^{2+} in the map for the SMA solubilized human TRPM4 sample without

416 the addition of extra calcium, which is comparable to the density shown for detergent-

417 solubilized mouse and human TRPM4 with calcium added to the sample.

Supplementary Fig. 10: Expression of HsTRPM4 variants. (a) Western blot showing the

expression of wildtype (WT) and HsTRPM4 variants expressed in HEK293 cells. (b)

Graphical plot of relative western blot intensities between wildtype (WT) and HsTRPM4

variants. (*) represents the p-value <0.05 compared to the wildtype condition.

a

b

Supplementary Fig. 11: Time course of wildtype calcium-activated sodium current for

HsTRPM4. (a) Example of raw traces (currents) from three different time points 1, 2, and 3

from the time course shown in (b) recorded from one cell expressing HsTRPM4 channels.

Without free calcium (time point 1 in (b)), no calcium-activated sodium current is recordable

at -100 mV or +100 mV (trace n°1 in (a)). Perfusion of 300 µM of free calcium concentration

leads to the rapid activation of the channel (time point 2 in (b)), and a calcium-activated

sodium current is recordable (trace n°2 in (a)) at both voltages (-100 mV and +100 mV).

Following this quick activation, a rapid run-down of the calcium-activated sodium current

mediated by the HsTRPM4 channel, due to the depletion of the PIP2 element, is observable

until the current reaches stability (time point 3 in (b)). After the run-down, the calcium-

activated sodium current amplitude is quantified by calculating the amplitude of this current

during the last 100 ms at +100 mV (grey zone in (a)). As depicted in trace number 3 in (a),

the run-down is generally more pronounced at -100 mV, leading to almost a complete

inhibition of the calcium-activated sodium current at this voltage.

Supplementary Fig. 12: Calcium-activated sodium currents for HsTRPM4. (a to l)

Representative traces of wildtype (WT, gray traces) and variants HsTRPM4 currents (black

traces): S863A, V901W, V904W, H908A, S924A, K925A, Y1057A, Q1061A, R1064A,

R1064G, and R1064S. Calcium-activated sodium currents for HsTRPM4 variant channels

are generally smaller than WT currents, as depicted in Figure 3e, due to the amino acid

alteration.

a

■ HsTRPM4 V904W (n=6) (EC$_{50}$: n.d)
▲ HsTRPM4 Y1057A (n=7) (EC$_{50}$: 246 ± 97 µM)
○ HsTRPM4 Q1061A (n=6) (EC$_{50}$: 645 ± 42 µM)
□ HsTRPM4 R1064S (n=6) (EC$_{50}$: 507 ± 57 µM)
△ HsTRPM4 S863A/S924A (n=6) (EC$_{50}$: 462 ± 11 µM)

b

● HsTRPM4 WT + vehicle (n=7) (EC$_{50}$: 966 ± 119 µM)
◆ HsTRPM4 WT + NBA 200 nM (n=6) (EC$_{50}$: 528 ± 62 µM)

Supplementary Fig. 13: Calcium sensitivity curves of wildtype and variant HsTRPM4. (a) Calcium sensitivity curves of a few loss-of-function HsTRPM4 variants. (n): number of cells. n.d: not determinable. (b) Effect of NBA (200 nM) on the calcium sensitivity on wildtype (WT) HsTRPM4 channels.

451 **Supplementary Fig. 14**: Calcium-activated sodium currents from HsTRPM4 channels. (a

452 and b) Representative traces of variants HsTRPM4 current: S863A and S924A in the

453 absence of free calcium in the solution (dotted black line), in the presence of 300 µM of free

454 calcium concentration (grey line), and in the presence of 300 µM of free calcium

455 concentration and 1 µM NBA (black line).

References

1. Clapham, D. E. TRP channels as cellular sensors. *Nature* **426**, 517–524 (2003).

2. Song, M. Y. & Yuan, J. X.-J. Introduction to TRP channels: structure, function, and

regulation. *Adv. Exp. Med. Biol.* **661**, 99–108 (2010).

3. Koivisto, A.-P., Belvisi, M. G., Gaudet, R. & Szallasi, A. Advances in TRP channel drug

discovery: from target validation to clinical studies. *Nat. Rev. Drug Discov.* **21**, 41–59

(2022).

4. Fallah, H. P. *et al.* A Review on the Role of TRP Channels and Their Potential as Drug

Targets_An Insight Into the TRP Channel Drug Discovery Methodologies. *Front.*

Pharmacol. **13**, 914499 (2022).

5. Launay, P. *et al.* TRPM4 is a Ca2+-activated nonselective cation channel mediating cell

membrane depolarization. *Cell* **109**, 397–407 (2002).

6. Nilius, B. *et al.* Voltage dependence of the Ca2+-activated cation channel TRPM4. *J.*

Biol. Chem. **278**, 30813–30820 (2003).

7. Wang, C., Naruse, K. & Takahashi, K. Role of the TRPM4 Channel in Cardiovascular

Physiology and Pathophysiology. *Cells* **7**, (2018).

8. Guinamard, R., Demion, M. & Launay, P. Physiological roles of the TRPM4 channel

extracted from background currents. *Physiology* **25**, 155–164 (2010).

9. Cheng, H. *et al.* TRPM4 controls insulin secretion in pancreatic beta-cells. *Cell Calcium*

41, 51–61 (2007).

10. Bianchi, B., Ozhathil, L. C., Medeiros-Domingo, A., Gollob, M. H. & Abriel, H. Four

TRPM4 Cation Channel Mutations Found in Cardiac Conduction Diseases Lead to

Altered Protein Stability. *Front. Physiol.* **9**, 177 (2018).

11. Liu, H. *et al.* Gain-of-function mutations in TRPM4 cause autosomal dominant isolated

cardiac conduction disease. *Circ. Cardiovasc. Genet.* **3**, 374–385 (2010).

12. Liu, H. *et al.* Molecular genetics and functional anomalies in a series of 248 Brugada

cases with 11 mutations in the TRPM4 channel. *PLoS One* **8**, e54131 (2013).

13. Janin, A. *et al.* TRPM4 mutations to cause autosomal recessive and not autosomal

dominant Brugada type 1 syndrome. *Eur. J. Med. Genet.* **62**, 103527 (2019).

14. Stallmeyer, B. *et al.* Mutational spectrum in the Ca(2+)--activated cation channel gene

TRPM4 in patients with cardiac conductance disturbances. *Hum. Mutat.* **33**, 109–117

(2012).

15. Hof, T. *et al.* TRPM4 non-selective cation channel variants in long QT syndrome. *BMC*

Med. Genet. **18**, 31 (2017).

16. Borgström, A., Peinelt, C. & Stokłosa, P. TRPM4 in Cancer—A New Potential Drug

Target. *Biomolecules* **11**, 229 (2021).

17. Zhao, Y., McVeigh, B. M. & Moiseenkova-Bell, V. Y. Structural Pharmacology of TRP

Channels. *J. Mol. Biol.* **433**, 166914 (2021).

18. Guinamard, R., Hof, T. & Del Negro, C. A. The TRPM4 channel inhibitor 9-phenanthrol.

Br. J. Pharmacol. **171**, 1600–1613 (2014).

19. Arullampalam, P. *et al.* Species-Specific Effects of Cation Channel TRPM4 Small-

Molecule Inhibitors. *Front. Pharmacol.* **12**, 712354 (2021).

20. Ozhathil, L. C. *et al.* Identification of potent and selective small molecule inhibitors of the

cation channel TRPM4. *Br. J. Pharmacol.* **175**, 2504–2519 (2018).

21. Guinamard, R., Simard, C. & Del Negro, C. Flufenamic acid as an ion channel

modulator. *Pharmacol. Ther.* **138**, 272–284 (2013).

22. Ruan, Z. *et al.* Structures of the TRPM5 channel elucidate mechanisms of activation

and inhibition. *Nat. Struct. Mol. Biol.* **28**, 604–613 (2021).

23. Zhao, C. *et al.* Structures of a mammalian TRPM8 in closed state. *Nat. Commun.* **13**,

3113 (2022).

24. Su, N. *et al.* Structural mechanisms of TRPV2 modulation by endogenous and

exogenous ligands. *Nat. Chem. Biol.* **19**, 72–80 (2023).

25. Guo, J. *et al.* Structures of the calcium-activated, non-selective cation channel TRPM4.

Nature **552**, 205–209 (2017).

26. Autzen, H. E. *et al.* Structure of the human TRPM4 ion channel in a lipid nanodisc.

Science **359**, 228–232 (2018).

27. Winkler, P. A., Huang, Y., Sun, W., Du, J. & Lü, W. Electron cryo-microscopy structure

of a human TRPM4 channel. *Nature* **552**, 200–204 (2017).

28. Duan, J. *et al.* Structure of full-length human TRPM4. *Proc. Natl. Acad. Sci. U. S. A.*

115, 2377–2382 (2018).

29. Dörr, J. M. *et al.* The styrene-maleic acid copolymer: a versatile tool in membrane

research. *Eur. Biophys. J.* **45**, 3–21 (2016).

30. Syam, N. *et al.* Variants of Transient Receptor Potential Melastatin Member 4 in

Childhood Atrioventricular Block. *J. Am. Heart Assoc.* **5**, (2016).

31. Smith, A. A. A. *et al.* Lipid Nanodiscs via Ordered Copolymers. *Chem* **6**, 2782–2795

(2020).

32. Hu, J. *et al.* Physiological temperature drives TRPM4 ligand recognition and gating.

Nature **630**, 509–515 (2024).

33. Yin, Y. *et al.* Structural basis of cooling agent and lipid sensing by the cold-activated

TRPM8 channel. *Science* **363**, (2019).

34. Wang, L. *et al.* Structures and gating mechanism of human TRPM2. *Science* **362**,

(2018).

Methods

Synthesis of 4-Chloro-2-(2-(3-iodophenoxy)acetamido)benzoic acid (IBA)

General Remarks

Reagents and organic solvents were purchased from commercial suppliers and used without further purification. Deionized water produced in-house or commercially available Milli-Q® water was used depending on the application. Aqueous solutions of sodium hydroxide, hydrogen chloride, saturated ammonium chloride, and saturated sodium chloride (brine) were prepared with deionized water. Thin layer chromatography (TLC) was performed using Macherey-Nagel ALUGRAM® Xtra SIL G/UV$_{254}$ plates coated with 0.20 mm silica gel 60 containing fluorescent indicator. High-pressure liquid chromatography (HPLC) was performed using a Thermo Fisher Scientific UltiMate 3000 RSLCnano System composed of a DIONEX UltiMate 3000 Pump, a DIONEX UltiMate 3000 Sampler, a DIONEX UltiMate 3000 Column Compartment, and a DIONEX UltiMate 3000 Diode Array Detector. HPLC measurements were conducted using Milli-Q® water (+ 0.1 % TFA) and acetonitrile (+ 0.1 % TFA) as eluents and an Acclaim™ 120 C18 column (Thermo Scientific™). Flash column chromatography (LC) was performed using the Teledyne Isco Combi*Flash©Rf*+ system. Teledyne Isco Redi*Sep©Rf* dry load cartridges were used to prepare dry loads. If not stated otherwise, dry loads were prepared on silica gel. Teledyne Isco Silica Redi*Sep©Rf* prepacked silica flash columns of two sizes (24g and 80g) were used. Nuclear magnetic resonance spectroscopy (NMR) was performed at the Department of Chemistry, Biochemistry, and Pharmaceutical Sciences, Universität Bern (Furrer Group) using a Bruker AVANCE III HD 300 GA spectrometer with a magnetic field of 7.05 Tesla and operating frequencies of 300.13 MHz for [1]H measurements and 75.48 MHz for [13]C measurements. High-resolution mass spectrometry (HRMS) was performed by the mass spectrometry service (Schürch group) at the Department of Chemistry, Biochemistry and Pharmaceutical Sciences, Universität Bern. The measurements were performed using electrospray ionization (ESI) and a ThermoScientific LTQ Orbitrap XL mass spectrometer with high mass resolution (m/Δm > 100'000) and accuracy (Δm < 3ppm). The previously published synthetic

procedure (Ozhathil *et al.*, 2018) to generate similar anthranilic anilide compounds was

slightly adapted.

Methyl 4-chloro-2-(2-chloroacetamido) benzoate

Methyl 2-amino-4-chlorobenzoate (1.7876 g, 9.6308 mmol) and potassium carbonate

(2.6620 g, 19.2616 mmol, 2 eq.) were dissolved in tetrahydrofuran (150 mL) and stirred for

10 minutes at room temperature. Not all potassium carbonate dissolved completely. The

mixture was then cooled in an ice bath, and chloroacetyl chloride (1.0919 g, 9.6676 mmol,

0.77 mL, 1 eq.) was added dropwise via a syringe. This mixture was stirred at 0 °C for 10

minutes and further stirred at room temperature for 16 hours. The reaction mixture showed a

pale pink color and was monitored by thin layer chromatography (eluent: cyclohexane /ethyl

acetate, 4:1). After full conversion of the starting material, water was added to the reaction

mixture and the product was extracted with ethyl acetate. The combined organic phases

were then washed with brine, dried over magnesium sulfate, filtered through celite, and the

volatiles were evaporated under reduced pressure. The crude pale-yellow and solid product

was purified by flash column chromatography (eluent: cyclohexane /ethyl acetate, gradient

from 0 % to 20 % ethyl acetate). The product (white powder) was dried *in vacuo*. Yield

Methyl 4-chloro-2-(2-chloroacetamido) benzoate quant., 2.5 g, 9.5387 mmol. ^{1}H NMR (300

MHz, DMSO-d_6) δ 11.44 (s, 1H), 8.52 (d, *J* = 2.2 Hz, 1H), 8.00 (d, *J* = 8.6 Hz, 1H), 7.33 (dd,

J = 8.6, 2.2 Hz, 1H), 4.48 (s, 2H), 3.89 (s, 3H). ^{13}C NMR (75 MHz, DMSO-d_6) δ 166.75,

165.67, 140.29, 138.75, 132.51, 123.77, 119.84, 115.57, 52.82, 43.35. HRMS (ESI) *m/z*

[M+H]$^+$ calculated for $C_{10}H_9Cl_2NO_3$ 262.0032, found 262.0037.

Methyl 4-chloro-2-(2-(3-iodophenoxy)acetamido)benzoate

Methyl 4-chloro-2-(2-chloroacetamido) benzoate (0.3270 g, 1.2475 mmol) was dissolved in

dimethylformamide (6 mL), and potassium carbonate (0.3465 g, 2.5073 mmol, 2 eq.) was

added. This mixture was stirred for 10 minutes at room temperature. Not all potassium

carbonate dissolved completely. 3-Iodophenol (0.3019 g, 1.3723 mmol, 1.1 eq.) was then

dissolved in dimethylformamide (6 mL) and added via syringe to the reaction mixture, which

was stirred at 80 °C. The color of the reaction mixture changed to deep brown. The reaction

was monitored by thin layer chromatography (eluent: cyclohexane /ethyl acetate, 4:1). Upon

full conversion of the starting material after 60 minutes, water was added to the mixture,

resulting in a brown and cloudy suspension. The product was extracted with ethyl acetate,

and the combined organic phases were then washed with brine, dried over magnesium

sulfate, filtered through celite, and the volatiles were evaporated under reduced pressure.

The crude product was purified by flash column chromatography (eluent: cyclohexane /ethyl

acetate, gradient from 0 % to 20 % ethyl acetate), and the purified product (white powder)

was dried *in vacuo*. Yield Methyl 4-chloro-2-(2-(3-iodophenoxy)acetamido)benzoate 84 %,

0.4653 g, 1.0442 mmol. ^{1}H NMR (300 MHz, DMSO-d_6) δ 11.76 (s, 1H), 8.70 (d, J = 2.1 Hz,

1H), 8.01 (d, J = 8.6 Hz, 1H), 7.48 (t, J = 1.9 Hz, 1H), 7.40 (ddd, J = 6.4, 2.5, 1.6 Hz, 1H),

7.30 (dd, J = 8.6, 2.2 Hz, 1H), 7.15 – 7.09 (m, 2H), 4.79 (s, 2H), 3.91 (s, 3H). ^{13}C NMR (75

MHz, DMSO-d_6) δ 167.24, 166.87, 157.61, 140.57, 138.94, 132.56, 131.50, 130.66, 123.67,

123.34, 119.33, 114.75, 114.64, 95.04, 67.35, 52.76. HRMS (ESI) *m/z* [M+H]$^+$ calculated for

$C_{16}H_{13}ClINO_4$ 445.9651, found 445.9647.

4-Chloro-2-(2-(3-iodophenoxy)acetamido)benzoic acid (IBA)

Methyl 4-chloro-2-(2-(3-iodophenoxy)acetamido)benzoate (0.4531 g, 1.0168 mmol) was

dissolved in methanol (100 mL) and potassium hydroxide (0.1744 g, 3.1075 mmol, 3 eq.)

was dissolved in Milli-Q® water (20 mL). The aqueous potassium hydroxide solution was

added to the reaction mixture, which was stirred at 65° C, and progress was monitored by

thin layer chromatography (eluent: dichloromethane /methanol, 9:1). Upon full conversion of

the starting material after 30 minutes, the reaction mixture was cooled down to room

temperate. Aqueous hydrochloric acid solution (1 M, ca. 100 mL) was then added, resulting

in immediate precipitation of the product, which was filtered off using a glass filter frit (Por.

4), and the solid white product was dried *in vacuo*. Yield 4-Chloro-2-(2-(3-

iodophenoxy)acetamido)benzoic acid 79 %, 0.3466 g, 0.8030 mmol. Purity (HPLC): 98.52 %

(at 254 nm). ^{1}H NMR (300 MHz, DMSO-d_6) δ 14.08 (s, 1H), 12.25 (s, 1H), 8.77 (d, J = 2.1

Hz, 1H), 8.03 (d, J = 8.6 Hz, 1H), 7.46 (m, 1H), 7.38 (dt, J = 7.0, 1.7 Hz, 1H), 7.27 (dd, J =

8.6, 2.2 Hz, 1H), 7.19 – 7.08 (m, 2H), 4.77 (s, 2H). ^{13}C NMR (75 MHz, DMSO-d_6) δ 168.78,

167.24, 157.58, 141.12, 138.69, 133.05, 131.48, 130.55, 123.74, 123.06, 118.78, 114.94,

114.29, 95.06, 67.21. HRMS (ESI) *m/z* [M-H]$^-$ calculated for $C_{15}H_{11}ClINO_4$ 429.9349, found

429.9348.

Description of drugs

Stock solutions of 4-chloro-2-(2-(naphthalene-1-yloxy)acetamido)benzoic acid (NBA) and the

iodo-modified congener; 4-chloro-2-(2-(3-iodo-phenoxy)acetamino)benzoic acid (IBA) were

dissolved in 100% DMSO to a final concentration of 10 mM. IBA was synthesized to

increase the electron densities when using the cryo-EM approaches. Although, as proof of

concept, only NBA has been used in the functional experiments presented in this study, the

inhibitory efficacity of IBA has been evaluated and is the same as of NBA. NBA and IBA

solutions were freshly made before each experiment. The lipophilicity of these drugs (cLog

P) has been calculated using the algorithms on Swiss ADME from the Swiss Institute of

Bioinformatics (http://www.swissadme.ch/). The Log *P* corresponds to the compound's

concentration ratio at equilibrium between organic (octanol) and aqueous phases. A

negative Log *P* means the compound is hydrophilic, and a positive value for Log *P* denotes a

more lipophilic compound. At pH = 7.4, the NBA and IBA compounds (both protonated

(carboxylic acid) and deprotonated (carboxylate) forms) tend to be lipophilic. NBA is slightly

more lipophilic than IBA (consensus cLog *P* protonated/deprotonated: NBA: 3.59/3.32; IBA:

3.25/3.05). The decadic $\log_{10}$ is used here.

639
Plasmid constructs design and cloning
641
642   For this study, the wild-type codon-optimized *H. sapiens* (Hs) and *M. musculus* (Mm)

643   TRPM4 genes coding full-length TRPM4 were synthesized, fused to consecutive C-terminal

644   HA (hemagglutinin)—and FLAG-tags, and cloned into a pCDNA-3.1 plasmid for expression

645   in HEK293 cells by the CMV (cytomegalovirus) promoter (GenScript Biotech). Mutant

646   variants were generated by site-directed mutagenesis (GenScript Biotech).

647
Protein expression and purification

The full-length *H. sapiens* TRPM4 and *M. musculus* TRPM4 were expressed and purified

from HEK293F cells grown in suspension. For expression, HEK293F cells were transfected

with 1 mg of a plasmid containing the TRPM4 gene per liter of cells using PEI

(polyethylenimine). The cultures were grown at 37°C and 5% CO_2 for 48 hours. The cultures

were harvested by centrifugation at 3,000 xg for 30 mins at 4°C and washed in 1x PBS,

followed by another round of centrifugation. The pellets were carefully resuspended in lysis

buffer containing HEPES-NaOH, pH 7.5, 200 mM NaCl and supplemented with cOmplete™

EDTA-free Protease Inhibitor Cocktail (Roche). 4 tablets of protease inhibitor cocktail were

added per 100ml of buffer. Following resuspension, the cells were lysed by sonication for a

total of 1 min in 10 sec On and OFF cycles and an amplitude of 35%. After sonication, the

membrane fraction was harvested by centrifugation using an Optima XPN-100

ultracentrifuge (Beckman Coulter) with the Ti45 rotor and spun at 70,560 × g for 30 minutes

at 4°C. The resulting pellets were stored at -80°C. For solubilization of the membrane

fraction with SMALP200, 12 g of pellet was resuspended in 30 ml of solubilization buffer

containing HEPES-NaOH, pH 7.5, 200 mM NaCl and 0.5% SMALP200 (CubeBiotech

GmbH) supplemented with 2 tablets of cOmplete™ EDTA-free Protease Inhibitor Cocktail

(Roche). The resuspended pellet was homogenized manually in a 40 ml Kimble glass

homogenizer (Sigma). The tube containing the mixture was then placed in a bottle with a stir

bar and left to incubate at 4°C for 1 hour 30 mins. The homogenate was clarified by

centrifugation for 30 min at 70,560 × g at 4°C in an Optima XPN Ultracentrifuge (Beckman

Coulter) using a Ti-45 rotor. The supernatant, which contained soluble FLAG-tagged

HsTRPM4 mixed with 1ml of Anti-FLAG® M2 affinity gel (Millipore, Billerica, MA) pre-

equilibrated with wash buffer containing 25 mM HEPES-NaOH, pH 7.5, 200 mM NaCl. The

beads were washed with 100 mL of wash buffer containing 25 mM HEPES-NaOH, pH 7.5,

200 mM NaCl and eluted with 4 mL elution buffer containing 25 mM HEPES-NaOH, pH 7.5,

200 mM NaCl, 120 µg/ml of 3xFLAG peptide, followed by concentration on 100 K Amicon

Ultra-15 concentrators (Millipore, Billerica, MA) to an absorbance at 280 nm of 1.0 to prepare

cryo-EM grids.

For detergent solubilization, 12 g of pellet was resuspended in 30 ml of solubilization buffer

containing HEPES-NaOH, pH 7.5, 200 mM NaCl, 1% n-Dodecyl-beta-Maltoside (DDM) and

0.1% Cholesteryl Hemissucinate (CHS)

(Avanti Pola lipids) supplemented with 2 tablets of cOmplete™ EDTA-free Protease Inhibitor

Cocktail (Roche). The resuspended pellet was homogenized manually in a 40 ml Kimble

glass homogenizer (Sigma). The tube containing the mixture was then placed in a bottle with

a stir bar and left to incubate at 4°C for 2 hours. The homogenate was clarified by

centrifugation for 30 min at 70,560 × g at 4°C in an Optima XPN Ultracentrifuge (Beckman

Coulter) using a Ti-45 rotor. The supernatant, which contains soluble FLAG-tagged

HsTRPM4 mixed with 1ml of Anti-FLAG® M2 affinity gel (Millipore, Billerica, MA) pre-

equilibrated with wash buffer containing 25 mM HEPES-NaOH, pH 7.5, 200 mM NaCl and

1% DDM/ 0.1% CHS. The beads were washed with 100 mL of wash buffer containing 25 mM

HEPES-NaOH, pH 7.5, 200 mM NaCl, and 1% DDM/ 0.1% CHS and eluted with 4 mL

elution buffer containing 25 mM HEPES-NaOH, pH 7.5, 200 mM NaCl, 120 µg/ml of 3xFLAG

peptide and 1% DDM/ 0.1% CHS. The purified protein was run on a Superose 6 gel filtration

column pre-equilibrated with 0.005% Lauryl Maltose Neopentyl Glycol (LMNG)/ 0.0005%

CHS and the peak fraction was concentrated on 100 K Amicon Ultra-15 concentrators

(Millipore, Billerica, MA) to an absorbance at 280 nm of 1.0 to prepare cryo-EM grids.

SDS-PAGE Analysis

An SDS-PAGE analysis was performed to assess the purity of purified proteins. 15 µL of

protein sample was supplemented with 5 µL of 4X NuPAGE LDS Sample Buffer (Thermo

Scientific). Samples were incubated at 95°C for 10 minutes before loading on 4-12%

SurePAGE™ Bis-Tris precast gels (Witec AG). Spectra™ Prestained Protein Ladder

(Thermo Scientific) (10 to 180 kDa) was also loaded on the gel to run as a size marker. Gels

were run in 1X Tris-MOPS SDS running buffer (Witec AG) at 200 V for 30 minutes, washed

briefly in MilliQ water, and stained for 2 hours with QuickBlue Protein Stain (LuBioScience

GmbH) with shaking. Gels were washed in MilliQ water before imaging on an iBright FL1500

Imaging System (Thermo Scientific).

Cryo-EM sample preparation and data collection

Purified TRPM4 was incubated with a final concentration of 0.2mM IBA for $TRPM4_{IBA}$ and

NBA for $TRPM4_{NBA}$ for 30 mins at room temperature before freezing cryo-EM grids. Cryo-EM

grids were prepared by applying 3 µl of concentrated sample onto 400-mesh R1.2/1.3

UltrAuFoil grids (Quantifoil Micro Tools GmbH), which had rendered hydrophilic by glow

discharging at 15mA for 60seconds with a PELCO EasyGlow device (TED PELLA, INC).

The samples were immediately blotted and plunged frozen into liquid ethane using a Vitrobot

Mark IV plunge freezer (Thermo Fisher Scientific). Cryo-EM data were collected using the

automated data acquisition software EPU (Thermo Fisher Scientific) on a Titan Krios G4

transmission electron microscope (Thermo Fisher Scientific), operating at 300kV and

equipped with a cold-FEG electron source, a SelectrisX energy filter and a Falcon4 direct

detection camera. Images were recorded in counting mode at a nominal magnification of

165kx, corresponding to a physical pixel size of 0.726 Å at the sample level. Datasets were

collected at a defocus range of 0.8 to 2.5 µm with a total electron dose of 60 $e^-/Å^2$. Image

data were saved as Electron Event Recordings (EER).

Cryo-EM image processing, model building, and refinement

The cryo-EM image processing was performed using cryoSPARC v3.4 [34].

Patch-based motion correction (cryoSPARC implementation) was used to align the EM

movie stacks and apply dose-dependent resolution weighting to recorded movies. CTF

estimation was performed using the patch-based option as well. For the data of the

HsTRPM4$_{apo}$, a total of 6,429 movies at 0.726 Å per pixel were collected, and 1000 particles

were manually picked and used for one round of 2D classification for template creation.

Template-based automated particle picking was then used on the recorded image data,

which resulted in a set of 1'719'323 particles at a size of 450 pixels. Two rounds of 2D

classification were performed for the initial step of particle cleaning, resulting in 255'053

particles in the first round and 16'308 in the second round. Ab initio and non-uniform

refinement yielded one 3D reconstruction with a map at 3.67 Å overall resolution in C4

symmetry (**Supplementary Fig. 4**).

For the data of the HsTRPM4$_{IBA}$, a total of 7,500 movies at 0.726 Å per pixel were collected.

2D classes from the HsTRPM$_{apo}$ dataset were used for template-based picking. Template-

based automated particle picking resulted in a set of 1'149'090 particles at a size of 450

pixels. Two rounds of 2D classification were performed for the initial step of particle cleaning,

resulting in 185'741 particles at 400 pixels in the first round and 35'399 particles in the

second round. Ab initio and hetero-refinement refinement yielded two 3D reconstructions.

One reconstruction representing 57.5% of particles was selected for further non-uniform

refinement, resulting in a map at 3.62 Å overall resolution in C4 symmetry. Following

symmetry expansion, the overall resolution of the map could be improved to an overall

resolution of 3.62 Å (**Supplementary Fig. 5**).

For the data of the HsTRPM4$_{NBA}$, a total of 11,851 movies at 0.726 Å per pixel were

collected. 2D classes from the HsTRPM$_{apo}$ dataset were used for template-based picking.

Template-based automated particle picking resulted in a set of 1'805'182 particles at 400

pixels. Four rounds of 2D classification were performed for the initial step of particle

cleaning, resulting in 40'633 particles. Ab initio and hetero-refinement refinement yielded

four 3D reconstructions. One reconstruction representing 15'954 particles was selected for

further non-uniform refinement, resulting in a map at 4.50 Å overall resolution in C4

symmetry from 15,954 particles (**Supplementary Fig. 6**).

Cryo-EM data was also collected for DDM/CHS solubilized mouse and human TRPM4.

MmTRPM4 data was collected with 5mM Calcium chloride present in the sample. Two

datasets were collected for DDM/CHS solubilized HsTRPM4 with 5mM Calcium chloride

present in one sample and both 5mM calcium chloride and 0.2mM NBA present in the other.

Details of the data collection are included (**Supplementary Table 2**).

Atomic models for HsTRPM$_{apo}$, HsTRPM4$_{IBA}$, and HsTRPM4$_{NBA}$ structures were mainly built

in Coot 0.9.4 [35], using a model PDB id: 6BQV as an initial model. Real-space refinement for

all built models was performed using Phenix, version 1.19.2-4158, by applying a general

restraints setup [33].

Cell Culture

TsA-201 cells were cultured with Dulbecco's Modified Eagle's culture Medium DMEM

(Gibco, Basel, Switzerland) supplemented with 10% FBS, 0.5% penicillin, and streptomycin

(10,000 U/mL) at 37°C in a 5% CO_2 incubator.

Transfections

Sixty mm dishes (BD Falcon, Durham, North Carolina, USA) at 80% of confluence were

transiently transfected using Xtreme Gene 9™ transfection reagent (Sigma Aldrich Merck,

Switzerland) and following the manufacturer's instructions. In brief, 1,000 ng of either empty

vector or HsTRPM4 WT or variants of HsTRPM4 (TRPM4 S863A, TRPM4 V901W, TRPM4

V904W, TRPM4 H908A, TRPM4 S924A, TRPM4 K925A, TRPM4 Y1057A, TRPM4 Q1061A,

TRPM4 R1064A, TRPM4 R1064G, TRPM4 R1064S or TRPM4 S863A/S924A) was mixed

with a solution containing 100 ng of a reporter gene coding for GFP and 1,000 ng of empty

vector. The expression of GFP was used to evaluate the transfection efficiency procedure,

and the empty vector was added to reach a certain amount of total cDNA to ensure an

efficient transfection. These final cDNA solutions (2,100 ng) were mixed with 210 µL of opti-

MEM (Gibco, Basel, Switzerland) and 6.3 µL of Xtreme Gen9™ reagent (ration 1/3). After 30

minutes at room temperature, the cDNA solutions were applied to the cells. Forty-eight hours

post-transfection, the cells were harvested for the western blot experiments.

Western-blots

The expression of the human TRPM4 channel was assessed in whole-cell lysates. First,

cells were washed with PBS 1X and then lysed for 1 hour at 4 °C in lysis buffer (50 mM

HEPES pH 7.4, 1.5 mM $MgCl_2$, 150 mM NaCl, 1 mM EGTA pH 8, 10% glycerol, 1% Triton X-

100, and Complete® protease inhibitor cocktail (Roche Diagnostics, Mannheim, Germany)).

The pellet was discarded after centrifugation at 4 °C, 16,000 g for 15 minutes. Protein

concentrations of each lysate sample were measured in triplicate by Bradford assay and

interpolated by a bovine serum albumin (BSA) standard curve. Samples were denatured at

30 °C for 37 minutes before loading on a gel. Twenty µg of protein for each sample was run

at 150 V for 1 hour on 9% polyacrylamide gels. The Turbo Blot dry blot system (Biorad,

Hercules, CA, USA) was used to transfer the samples to a nitrocellulose membrane. All

membranes were stained with Ponceau as a qualitative check for equivalent total protein

loading. Membranes were then rinsed twice with TBS 1X and blocked with 5% BSA in TBS

1X for 1 hour. After this blocking step, the membranes were incubated for 2 hours with rabbit

anti-human TRPM4 antibody (epitope: $_{1137}$CRDKRESDSERLKRTSQKV$_{1155}$, Pineda, Berlin,

Germany) diluted 1:1,000 in TBS 1X + 0.1% tween and mouse anti-Na$^+$/K$^+$ ATPase antibody

(Abcam ab 7671) diluted 1:1,000 in TBS 1X + 0.1% tween. The membranes were washed 4

times in TBS 1X + 0.1% tween before incubating with fluorescent secondary antibodies.

Secondary antibodies   IR Dye 800 CW, anti-rabbit diluted (1:20,000) in TBS 1X + 0.1%

tween and IR Dye 700 CW, anti-mouse diluted (1:20,000) in TBS 1X + 0.1% tween (LI-COR

Biosciences, Lincoln, NE, USA) were added for 1 hour. After 4 washes with TBS 1X + 0.1%

tween and 3 washes in TBS 1X, membranes were scanned with the FUSION FX Spectra®

Infrared Imaging System (VILBER smart imaging, Marne-la-Vallée, France) to detect

fluorescent protein. Subsequent quantitative analysis of protein content was achieved by

measuring and comparing band densities (equivalent to fluorescence intensities of the

bands) using the *Evolution-Capt software* (VILBER smart imaging, Marne-la-Vallée, France)*.*

The background was first subtracted for each band (human TRPM4 and Na$^+$/K$^+$ ATPase),

then TRPM4 intensity was divided by the intensity of the Na$^+$/K$^+$ ATPase band (for a given

sample) and normalized for comparison.

Electrophysiology

Transfections

Thirty-five mm dishes (BD Falcon, Durham, North Carolina, USA) at 80% of confluence were

transiently transfected using JetPEI^{TM} transfection reagent (Polyplus transfection, Illkirch,

France) and following the instructions of the manufacturer. In brief, 500 ng of either empty

vector or HsTRPM4 WT or variants of HsTRPM4 (HsTRPM4 T677I, HsTRPM4 S863A,

HsTRPM4 V901W, HsTRPM4 V904W, HsTRPM4 H908A, HsTRPM4 S924A, HsTRPM4

K925A, HsTRPM4 Y1057A, HsTRPM4 Q1061A, HsTRPM4 R1064A, HsTRPM4 R1064G,

HsTRPM4 R1064S or HsTRPM4 S863A/S924A) was mixed with 200 ng of a reporter gene

coding for GFP. Expression of GFP was used to identify transfected cells during patch clamp

experiments. Coding DNAs (cDNAs from HsTRPM4 and GFP) were mixed with 4 µl of

JetPEI^{TM} and 46 µl of 150 mM NaCl. After 15 minutes at room temperature, the cDNA

solutions were applied to the cells. Twenty-four hours post-transfection, the cells were plated

at low density in a new 35 mm dish coated with poly-l-lysine. The cells were patched 24

hours post-platting (48 hours post-transfection).

<u>Inside-out patch clamp</u>

Electrophysiological recordings were performed in the inside-out patch-clamp configuration

with patch pipettes (1 and 2 µm tip opening) pulled from 1.5 mm borosilicate glass capillaries

(Zeitz-Instruments GmbH, München, Germany) using micropipette puller P 97 (Sutter

Instruments, Novato, CA, United States). The tips were polished for 2–4 MΩ pipette

resistance in the bath solution. The pipette solution contained 150 mM NaCl, 10 mM

HEPES, and 2 mM CaCl$_2$ (pH 7.4 with NaOH). The initial bath solution with a concentration

of 0 µM of free calcium contained 150 mM NaCl, 10 mM HEPES, and 2 mM HEDTA (pH 7.4

with NaOH). After reaching the inside-out configuration, different solutions were perfused at

the intracellular side of the membrane patch using a modified rapid solution exchanger

(Perfusion Fast-Step SF-77B; Warner Instruments Corp. CT, United States).

<u>For calcium sensitivity experiments</u>. The first solution, applied for 3 to 5 sweeps to reach a

stable current, is the bath solution (150 mM NaCl, 10 mM HEPES, and 2 mM HEDTA; pH

7.4 with NaOH) with a concentration of 0 µM of free calcium. Then, to activate TRPM4

channels, a solution (150 mM NaCl, 10 mM HEPES; pH 7.4 with NaOH) containing different

concentrations of free calcium (0 µM, 50 µM, 100 µM, 300 µM, 500 µM, 1000 µM, 5000 µM)

was applied. The effect of the calcium was quantified when the sodium current of the

calcium-activated HsTRPM4 reached stability. Finally, the bath solution with a concentration

of 0 µM of free calcium was used to quantify the potential leak current that can occur during

such recordings.

For drug dose-response experiments. The first solution, applied for 3 to 5 sweeps to reach

the stable current, is the bath solution (150 mM NaCl, 10 mM HEPES, and 2 mM HEDTA;

pH 7.4 with NaOH) with a concentration of 0 µM of free calcium. Then, after stabilization of

the current, a solution (150 mM NaCl, 10 mM HEPES; pH 7.4 with NaOH) containing a

concentration of 300 µM free calcium without compounds was applied to activated

HsTRPM4 channels and recorded the sodium current of the calcium-activated HsTRPM4

until its stabilization. Then, a solution (150 mM NaCl, 10 mM HEPES; pH 7.4 with NaOH)

containing a concentration of 300 µM of free calcium and the compound at different

concentrations (10 nM, 50 nM, 100 nM, 500 nM, 1000 nM, 5000 nM, 10'000 nM, 50'000 nM)

were applied. The drug's effect was quantified when the sodium current of the calcium-

activated HsTRPM4 reached stability. Finally, the bath solution with a concentration of 0 µM

of free calcium was used to quantify the potential leak current that may occur during such

recordings. No investigation and comparison have been made concerning the kinetics of

block and washout, and only ''stable'' currents have been used to calculate the percentage

of inhibition.

The sodium current of the calcium-activated HsTRPM4 was recorded with a Multiclamp

700B amplifier (Molecular Devices, Sunnyvale, CA, United States) controlled by Clampex 10

via Digidata 1332A (Molecular Devices, Sunnyvale, CA, United States). Data were low-pass

filtered at 5 kHz and sampled at 10 kHz. Experiments were performed at room temperature

(20–25°C). The holding potential was 0 mV. The stimulation protocol consisted of two pulses

totaling 1'000 ms for measuring steady-state currents repeated at 0.2Hz (1 sweep every 5

sec.). The first pulse was at −100 mV for 500 ms, and the second was at +100 mV for

500 ms. For analysis, the effect of the compounds on calcium-activated sodium current

HsTRPM4 has been calculated by averaging the values of this current during the last

100 ms of the second sweep (at +100 mV) on the trace recorded during the stable phase.

Due to the stronger run-down observable at -100 mV (first pulse of the stimulation protocol)

for the HsTRPM4 channels leading to, after stabilization of the current, at sometimes almost

no sodium current quantifiable at -100 mV of the calcium-activated HsTRPM4 channel

(supplementary Figure 11b, trace number 3 at -100 mV), the effects of the compounds at -

100 mV have not been quantified. Electrophysiology data were exported and analyzed using

Prism7.05 GraphPad™ software (GraphPad by Dotmatics, San Diego, CA, USA).

Concentration-response curves were fitted using the log(inhibitor) vs. response-variable

slope (four parameters) equation (Y=Bottom + (Top-Bottom)/(1+10^((LogIC50-X)

*HillSlope))) where X corresponds to the log of concentration and Y current recorded in pA.

The decadic $\log_{10}$ is used here.

Data analyses and statistics

Data are represented as means ± SEM. Statistical analyses were performed using

Prism7.05 GraphPad™ software (GraphPad by Dotmatics, San Diego, CA, USA). An

unpaired Mann-Whitney U test was used to compare two unpaired groups. $p < 0.05$ was

considered significant. No muti-group comparison has been performed in this study. Western

blots were done in triplicate, and at least 6 cells were used for different doses in

electrophysiology experiments. Important information. Due to the limited channels available

in our perfusion system (7 channels available), only 5 different concentrations for

compounds (NBA or IBA) could be applied to the same cells (the 2 remaining channels were

used for the solution with a concentration of 0 μM of free calcium without no compounds to

record the background current not related to the activation of HsTRPM4 and the solution

with a concentration of 300 μM of free calcium with no compounds to activate the HsTRPM4

channel and record the calcium-activated sodium current). Not all the concentrations (8 in

total) for compounds presented in the different figures have been applied to the same cell.

This limitation did prohibit the determination of the IC_{50s} per cell; otherwise, the fundamental

statistical rule of independence would be violated. Consequently, IC_{50s} were calculated

based on the average data point collected for x cells per concentration, as presented in the

different figures, and no statistical analysis between the different conditions was performed

on those variables. However, a two-way ANOVA with two-factor repeated measures was

used to investigate any statistical difference of inhibition, mediated by the compounds,

between WT and variant conditions at 1µM of a compound, a concentration close to the

general IC_{50s} of the variants investigated. The percentages of inhibition for all variants,

except the TRPM4 T677I, at the concentration of 1µM of a compound, significantly differed

($p<0.05$) from the WT and are smaller, suggesting a decrease of inhibition, mediated by the

compounds, on these variants of HsTRPM4 channel.

Data visualization

Gel images were processed and prepared on ImageJ (Version 1.53k). Figures were

rendered using PyMOL, (The PyMOL Molecular Graphics System, version 1.8.2.0,

Schrödinger, LLC), UCSF Chimera, UCSF ChimeraX [36], and Adobe Illustrator

(https://adobe.com/products/illustrator).

Data and Code Availability

The reconstructed maps are available from the EMDB database under access codes

HsTRPM4apo (EMD-19057), HsTRPM4$_{IBA}$ (EMD-19061), HsTRPM4$_{NBA}$ (EMD-19069) (EMD-

19060, local). The atomic models are available in the PDB database, access codes

HsTRPM4apo (PDB 8RCR), HsTRPM4$_{IBA}$ (8RCU) and HsTRPM4$_{NBA}$ (PDB 8RD9).

Methods-only references

35. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms

for rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296

(2017).

36. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of

Coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).

37. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons

and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**,

861–877 (2019).

38. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization

and analysis. *Protein Sci.* **27**, 14–25 (2018).

937