Brain resident macrophages regulate sleep, with repopulated ones being unable to reestablish the original sleep circuits

Ali Seifinejad[1*], Gianni Monaco[1], Mojtaba Bandarabadi[2], Meriem Haddar[2],
Saskia Wundt[1], Victoria Wingert[3], Luciana Hannibal[3], Thomas Blank[1], Mehdi Tafti[2], and
Marco Prinz[1, 4*],

[1] *Institute of Neuropathology, Faculty of Medicine, University of Freiburg, Freiburg, Germany*
[2] *Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland*
[3] *Department of General Paediatrics, Adolescent Medicine and Neonatology, Laboratory of Clinical Biochemistry and Metabolism, Medical Center, University of Freiburg, Faculty of Medicine, Freiburg, Germany*
[4] *Signalling Research Centres BIOSS and CIBSS, University of Freiburg, Freiburg, Germany*

[*]*Correspondence to:*
Marco Prinz: marco.prinz@uniklinik-freiburg.de or
Ali Seifinejad: ali.seifinejad@mail.ch

Summary

Brain resident macrophages such as microglia and CNS-associated macrophages (CAMs) are already established before birth and play therefore a crucial role for normal brain functioning during development. However, their involvement in fine-tuning complex physiological functions such as vigilance states (VS) after birth remains poorly understood. Here, we investigated the reciprocal interaction of microglia and VS using multimodal high throughput transcriptional, electrophysiological and metabolomic profiling in mice. We found that sleep deprivation caused severe transcriptional changes in microglia and CAMs and absence of the wake-promoting neuropeptides hypocretin/orexin intensified these effects. Depletion of embryonic microglia robustly increased sleep quantity during the active period, while decreased sleep quality that was reflected in reduced power of brain sleep oscillations. Unexpectedly, subsequent repopulation by postnatal microglia failed to reestablish normal sleep-wake patterns, and even induced additional abnormalities such as sleep fragmentation. Moreover, we found a substantial excitatory-inhibitory synaptic imbalance following microglia depletion, which was not normalized after microglial repopulation and even lead to an increase of inhibitory synapses in the brain. At the metabolite level, we observed a distinct metabolite pattern after microglia depletion, which largely returned to normal levels after repopulation. Our findings suggest a so far largely unknown interaction between microglia and brain VS and emphasizes striking functional differences between embryonic and postnatal-derived microglia, thereby potentially paving the way for the further exploration of microglia of different origin and their roles in sleep disorders and synaptic connectivity.

Keywords: Microglia; CAM, vigilance states, sleep; HCRT

Introduction

Tissue-resident myeloid cells in the CNS comprising parenchymal microglia and CAMs [1-4], play significant roles in pathology and protecting the CNS from damage [5]. Although the function of CAMs in regulating brain function is not well understood, microglia participate in a range of physiological functions including synaptic pruning [6], remodeling and displacement [7], pain control [8,9], learning and memory [10,11], neurogenesis [11], and sleep-wake regulation [12].

Microglial cells and leptomeningeal macrophages (as part of the CAMs) are derived from prenatal sources known as erythormyeloid precursors that engraft into the developing CNS [13,14] and facilitate the establishment of neuronal networks especially until the first postnatal weeks [6]. If embryonic microglial cells are depleted during adulthood, they repopulate from surviving endogenous cells [15]. The *de novo* establishment of the micoglial networks occurs by surviving cells through clonal expansion [16] and these postnatal-derived microglial cells exhibit a different transcriptinal signature [17]. However, whether these adultood-derived microglia exhibit any functional differences to orginal embryo-derived ones is currently largely unknown.

VS are divided into wakefulness, non-rapid eye movement sleep (NREMS), and rapid eye movement sleep (REMS), which are dynamic complex states and vital for the normal brain functioning. They are measured using electroencephalogram (EEG) and characterized by different brain oscillations, e.g. delta waves (0.5-4Hz) for NREMS and theta waves (6-9Hz) for REMS in rodents. Dysregulation of VS is associated with a variety of diseases such as sleep disorders, cardiovascular and neurodegenerative disorders such as Alzheimer's disease (AD) [18,19]. VS regulation and sleep oscillatory patterns rely on a complex CNS connectivity, and different cell types, mainly neuronal populations interact with each other (via synaptic connections and neurotransmitter release) to fine-tune the regulation of VS and brain oscillations. Among the major neuropeptides and neurotransmitters involved in VS regulation are hypocretin/orexin (HCRT) and melanin-concentrating hormone neurons (MCH) in the lateral hypothalamus (LH), serotonin (5HT)

and noradrenaline (NA) in the brainstem (BS), and acetylcholine in the BS and basal forebrain

(BF), and midbrain dopaminergic neurons [20,21].

While a few studies have demonstrated a bidirectional relationship between embryo-derived

microglia and the VS [12,22-24], a comprehensive understanding of how these two compartments

modulate each other is still lacking. Specifically, we do not yet know the transcriptional signature

of microglia and CAMs in response to sleep loss, both in healthy individuals and in disease

conditions, such as narcolepsy with cataplexy, where patients have undetectable HCRT

neuropeptide levels. This deficiency in HCRT neuropeptide is largely thought to be immune-

mediated, and there have been reports of increased MHC-II molecule expression in microglia at

the onset of canine narcolepsy [25]. Additionally, it remains unclear whether adulthood-derived

microglia are capable of re-establishing neural circuitries in the same manner as prenatal

microglial cells, including synaptic plasticity and neurochemical balance, which are crucial for the

normal VS regulation.

Here, we first studied the impact of sleep deprivation on the transcriptional signature of microglia

and CAMs in both wild-type (WT) and HCRT cell-ablated narcolepsy mouse model (*Hcrt-ataxin-*

3) mice. Next, through in-depth analysis, we explored whether the absence of embryo-derived

microglia and CAMs influences normal VS patterns and whether repopulated postnatal microglia

and CAMs can normalize these patterns. Finally, we investigated how the absence of microglia

and CAMs affects brain network connectivity and neurochemistry.

Results

HCRT neuron ablation significantly alters transcriptional profile of microglia and CAMs in response to sleep loss

Sleep deprivation (SD) induces functional alterations in microglia [26,27], however its impact on the transcriptional profile of microglia and CAMs has not yet been documented. Furthermore, it is uncertain whether any of the substances that regulate VS mentioned earlier could worsen or alleviate the impacts of sleep deficiency on microglia and CAMs. To better understand this, we isolated microglia and CAMs from the cortex and hypothalamus, which are the two main sleep-regulating centers, of both wild-type (WT) and *Hcrt-ataxin-3* mice (hereafter referred to as Ataxin mice) [28]. HCRT neurons play a central role in regulating wakefulness [29] and waking behaviors that drive sleep need [30], and HCRT deficiency is a hallmark biomarker of human narcolepsy with cataplexy. By studying these mice, we aimed at determining whether the lack of HCRT neuropeptide has any influence on the transcriptional profile of microglia and CAMs in response to SD possibly contributing to narcolepsy with cataplexy phenotype.

We deprived mice of sleep for 6 h (starting from light onset) and immediately afterward, isolated microglia (CD11b$^+$, CD45intermediate) and CAMs (CD11b$^+$, CD206$^+$) from the cortex and hypothalamus of both WT and Ataxin mice (Figure 1A, B). The control group was not sleep deprived. We extracted RNA and performed bulk RNA sequencing (RNA-seq) to compare their transcriptome across different conditions. After 6 hours SD, no changes were observed in expression of microglial core genes (*Hexb*, *P2ry12*, *Trem2*, and *TMEM119*) across the cortex and hypothalamus (Supplementary Fig. 1A). Furthermore, we did not observe any transcriptional changes when comparing microglia from wild-type (WT) and Ataxin mice under normal conditions, and only two *Lyz2* and *Slc31a2* genes were differentially regulated after SD in these two mouse lines (Supplementary Fig. 1B). When comparing the microglial transcriptome between SD and control conditions across the cortex and hypothalamus of WT mice, we observed many differentially expressed genes (DEGs) (110 genes) (Figure 1C, E). However, the same

comparison in Ataxin mice resulted in a larger number of DEGs compared to WT mice (270 genes)

(Figure 1D, F). This finding is intriguing because it suggests that the HCRT neuropeptide directly

or indirectly interacts with and modulates microglia during enforced wakefulness (SD). While we

did not detect the expression of HCRT receptors in microglia, we observed the expression of

downstream circuit receptors, including noradrenaline and serotonin receptors, in microglia. One

such gene is the beta-2 adrenergic receptor (*Adrb2*), which is predominantly expressed in

hypothalamic microglia compared to cortical microglia (Supplementary Fig. 1C). We also

observed that SD primarily affects cortical, rather than hypothalamic microglia in Ataxin mice. In

WT mice, however, the hypothalamic microglia are more impacted by SD (Supplementary Fig. 2),

indicating that the cortical microglia are more susceptible to the lack of HCRT.

Several genes were primarily regulated under the SD condition, such as *Cirbp*, *Ighd*, and *Npnt*

(Fig. 1G). Additionally, we found genes that were mainly expressed in the hypothalamus

compared to the cortex (Fig. 1H). Furthermore, we observed that genes such as *Sgk1* and *Pecr*

were largely regulated after SD in the hypothalamus but not in the cortex, suggesting structure-

specific microglia changes after SD (Fig. 1I). Lastly, we found genes that were exclusively

regulated in Ataxin mice like *Slc31a2*, *Lyz2*, and *Ccl4* (Fig. 1J). Gene set enrichment analysis of

the DEGs in WT mice revealed an enrichment of pathways related to protein folding, such as

"Positive regulation of protein folding" (Supplementary Fig. 3A). In contrast, the analysis of DEGs

in Ataxin mice indicated an enrichment of pathways like "Virion attachment to host cell"

(Supplementary Fig. 3B). This finding is intriguing, as the 2009 H1N1 vaccination has been linked

to a high incidence of narcolepsy disease [31], and it remains to be carefully investigated whether

microglia play a role in this process.

We next investigated how SD affects the transcriptome of CAMs in WT and Ataxin mice. As CAMs

are not abundant in the brain, we sorted them from the entire brain. Our results revealed that 6

hours of SD induced numerous DEGs in CAMs sorted from WT mice compared to the control

condition (Fig. 1K). Among these DEGs were the *App* and *Lyve1* genes (Fig. 1L), both of which

have been linked to Alzheimer's disease (AD) pathogenesis [32,33]. It is intriguing to note that *Lyve1*

gene expression almost tripled after 6 hours of SD. Gene set enrichment analysis of DEGs of

CAMs in WT mice revealed enrichment for pathways related to AD, including positive regulation

of tau-protein kinase activity (Supplementary Fig. 3C). Nevertheless, we were astonished to

discover an opposing result of only a small number of DEGs while contrasting SD versus control

in CAMs of Ataxin mice (Fig. 1M), as we witnessed numerous DEGs in microglia of Ataxin mice

after SD. As the DEGs found in CAMs of WT mice have been implicated in AD, a thorough

literature search uncovered a study by Roh et al. [34] indicating that a null mutation in the *Hcrt* gene

in an AD mouse model leads to a significant reduction in Aβ pathology in the brain. It can be

speculated that this effect is mediated by CAMs and not microglia.

Overall, these data indicate that both microglia and CAMs transcriptional networks are modulated

by VS and HCRT neuropeptide that regulates VS, suggesting a reciprocal interaction between

both systems.

Repopulated adulthood-derived microglia fall short in rebuilding normal NREMS circuitry

We demonstrated that there is an interaction between brain-resident macrophages and VS, which

leads us to expect that brain-resident macrophages are also involved in regulating sleep. While

there is a limited number of studies examining the regulatory role of microglia in VS [12,23,35], a

comprehensive understanding of this regulation, as well as the specific parameters of the VS

affected by brain-resident macrophages, remains unclear. To address this, we depleted microglia

from the brains of mice for two weeks using colony-stimulating factor 1 receptor (CSF1R) inhibition

via PLX5622 [36] and allowed them to repopulate for four weeks (Fig. 2A). CSF1 receptor is vital

for microglia survival. Antagonizing it severely depletes microglia within three days, and upon the

termination of treatment, a fast repopulation can be observed within a few days [17]. We observed

substantial microglia depletion (>86%) after two weeks of PLX5622 treatment and complete

repopulation four weeks after withdrawal (Fig. 2B, C).

We recorded EEG signals before depletion (control), after depletion and after repopulation (Fig.

2A). We included the repopulation phase to determine if it restores any potential changes in the

VS induced after microglia depletion. Since CSF1 inhibition also depletes CAMs [37], the effects we

report here are due to the depletion of both cell types, which we refer to collectively as MG. Our

experimental conditions therefore included control, MG-Dep (microglia/CAM depleted), and MG-

Repop (microglia/CAM repopulation). To evaluate the distribution of vigilance states, we

employed our standard sleep/wake phenotyping protocol [38], with two days of baseline (which were

averaged and presented as one day in the figures) and in the third day we had 6 hours of SD

(ZT0-ZT6) and 18 hours of recovery (Fig. 1D). In baseline conditions, all experimental mice

demonstrated the characteristic rhythmic diurnal behavior, with higher levels of wakefulness

during the dark period and higher levels of NREMS during the light period and exhibited the

increase in NREMS amounts in response to SD (Fig. 1E, F). The distribution of wakefulness and

NREMS in all experimental groups was typical, and the appearance of one state occurred at the

cost of the other. However, significant differences (NREMS increase) were observed in the

baseline dark period when comparing the MG-Dep to the control condition. Interestingly, the

repopulation of MG did not normalize this effect (Control: 12.75±0.72 min, MG-Dep: 16.78±0.77

min, MG-Repop: 16.54±0.70 min, mean ± SEM) (Fig. 2E, F, bottom panels). Additionally, during

the dark period, there was a notable reduction in wakefulness (Fig. 2E, F, top panels).

To gain insight into the homeostatic regulation of sleep in different mice groups and to determine

whether increased sleep is due to an increased need for sleep, we analyzed the time course of

NREMS EEG delta power (1-4 Hz, mainly referred to as sleep need) across our 3-day recordings.

All mice groups showed a typical decline in delta power during the light period and a strong

rebound after SD, indicating the homeostatic regulation of sleep need in all groups (Fig. 2G).

However, the buildup in delta power during the baseline dark period was significantly lower in

MG-Dep and MG-Repop mice compared to WT mice (Fig. 2G). This attenuation was more

pronounced in MG-Repop mice and may be partly related to increased sleep during the baseline

dark period (Fig. 2E, lower panel). However, when mice were exposed to enforced wakefulness

(SD), the EEG delta power significantly increased and this effect was more noticeable in the MG-

Repop condition (Fig. 2G). Altogether, these data indicate a deficient buildup in homeostatic sleep

need during the waking period.

Spectral analysis of NREMS showed that MG-Dep mice exhibited higher slow-delta (1-2 Hz)

activity compared to control mice during the baseline light period (Fig. 2H, left). This frequency

band appears to be more affected, even in MG-Repop mice, as it was significantly increased

during the recovery light period after the mice were exposed to SD (Fig. 2I, left). The brain circuits

that are more vulnerable to microglia depletion and repopulation, resulting in those changes in

EEG power, are still unknown. However, our previous study has shown an increase in slow delta

activity during the light period in 5HT transporter knockout (5HTT KO), 5HTT; Hcrt double KO,

and DR_Hcrt1&2cKO mice [38]. In contrast, the fast delta (3-4 Hz) was blunted during both baseline

and recovery light periods and only in the MG-Dep condition (Fig. 2H, I, left). During the baseline

dark period, large blunting was observed in both MG-Repop (1.5-3.5Hz) and MG-Dep (3-5.5Hz)

conditions. These deficits remained nearly the same during the recovery dark period, indicating

that NREMS of both experimental mice appears to be largely shallower (Fig. 2H, I, right).

Collectively, it can be concluded that the presence of MG is necessary for the precise regulation

of fast and slow-delta power as a proxy for the sleep homeostatic process. MG repopulation not

only fails to restore NREMS regulation but worsens it.

To further determine whether the increase in NREMS is due to consolidation of this state or more

NREMS episodes, we conducted the sleep fragmentation analysis. We found that while MG-Dep

mice have similar sleep bout duration patterns as the control condition, MG-Repop mice showed

a large number of shorter NREMS bouts (16 seconds [Control: 6.7±0.63, MG-Dep: 6.8±0.7, MG-

Repop: 9.84±1.12]) and 64 seconds [Control: 12.12±1.15, MG-Dep: 11.02±0.81, MG-Repop:

17.63±1.2]) and fewer longer NREMS bouts (>256 seconds [Control: 1.96±0.28, MG-Dep:

1.74±0.33, MG-Repop: 0.62±0.17]), suggesting a strong destabilization of NREMS in the MG-

Repop brain (Fig. 2J).

During NREMS, brief bursts of brain activity occur at the frequency range of 9-16 Hz, called

spindles, which are generated by thalamo-cortical interactions, and play roles in processes such

as learning, memory, and cognition [39]. To determine if the lack of MG and its repopulation

influence spindle generation, we quantified spindle rate during NREMS episodes and found a

significant reduction in MG-Dep mice compared to the control condition (Fig. 2K, L; see

"Methods"). However, in contrast to other features of NREMS, MG repopulation normalized the

occurrence rate of spindle events, indicating that MG repopulation partially rescued the circuit

deficits seen in MG depletion.

Altogether, these data suggest that MG is integral part of NREMS establishment and maintenance

that might operate in multiple ways like keeping the homeostasis of different neurotransmitters

and neuromodulatory connections. Moreover, repopulated postnatal MG fail to settle the normal

NREMS regulatory functions of embryo-derived MG.

Depletion of microglia prolongs REMS and degrades its quality.

Different neural pathways and neurotransmitters regulate REMS compared to NREMS. However,

the role of glial cells and MG in the regulation of REMS is not well understood. We quantified

REMS amount across the dark/light periods and found that MG-Dep mice display more REMS

compared to the control and MG-Repop conditions during the dark period (Control: 0.67±0.1, MG-

Dep: 1.17±0.21 min/h, mean ± SEM). However, in contrast to NREMS, MG repopulation corrected

the abnormal expression of REMS (MG-Repop: 0.61±0.13 min/h) (Fig. 3A, B). To investigate if

the increase in REMS is due to the consolidation of this state, we performed fragmentation

analysis and found that the number of shorter REMS bouts (8 s) is largely decreased (Control:

8.7±2.05, MG-Dep: 2.8±1.2, MG-Repop: 8.8±3.3) and longer bouts (>2 min) are increased

(Control: 1.5±0.8, MG-Dep: 6.1±1.1, MG-Repop: 4.71±1.4, mean ± SEM) (Fig. 3C), suggesting

further stabilization of REMS in the absence of MG.

Theta oscillations dominate EEG signal during REMS, where theta phase dynamically modulates

gamma amplitude in hippocampal and cortical networks [40]. We next investigated the contribution

of MG in theta oscillations and theta-gamma interactions. Spectral analysis of REMS revealed a

significant dampening of theta (6-8Hz) power in MG-Dep mice during the baseline dark period

(Fig. 3D), which was restored after MG repopulation. We have previously observed REMS

stabilization and reduced theta power in mice lacking *Hcrtr*1&2 genes in dorsal raphe serotonergic

neurons [38]. We also found that the theta-gamma coupling is significantly reduced in the MG-Dep

condition compared to the control. Although MG repopulation attempts to bring it back to the

normal level, it fails significantly (Fig. 3E, F). Altogether, these findings suggest that the presence

of MG is necessary for the normal expression of REMS, and the local MG that contribute in the

generation of REMS and theta oscillations can re-establish large part of their normal function in

REMS circuit formations after repopulation.

Microglia contribute to the maintenance of active wakefulness

The quality of prior wakefulness contributes to the expression of both NREMS and REMS. Along

these lines, a specific sub-state of prior wakefulness was recently proposed to drive NREMS need

[30], and some aspects of wakefulness may functionally substitute for REMS [41]. To investigate how

wakefulness is expressed in our experimental conditions, we performed a dynamic analysis of

the full waking EEG spectrum across three recording days. This global analysis revealed

differential expression of mainly higher waking frequency bands during light-dark cycles across

three days, and most notably during enforced wakefulness (SD). As shown in Figure 4a, both

MG-Dep and MG-Repop mice exhibit a large dampening of the lower gamma (≈45-55Hz) band

during the baseline dark period, which is severely exacerbated during the recovery dark period.

Additionally, the MG-Dep mice exhibit a large blunting of the higher gamma band (≈70-90Hz)

during SD, which partially normalizes after MG repopulation. However, the lower frequency bands

are not easily distinguishable on the scale of this analysis. We, therefore, performed PSD analysis

and found that MG-Dep mice exhibit a large increase in slow delta (1.35-1.75Hz) activity

compared to the control mice (Fig. 4B). The increase in this waking frequency band has been

previously reported in mice lacking the serotonin transporter gene [38].

Mice spend some portion of their waking time engaged in exploratory and motivated behaviors,

such as running, nest building, or drinking, often referred to in rodents as "active wakefulness."

These behaviors are associated with the EEG signature called "theta-dominated wakefulness"

(TDW 6.0-9.5 Hz), which is highly dependent on the expression of the HCRT neuropeptide, and

has been suggested to be the principal driver of the sleep homeostat (Vassalli and Franken,

2017). We quantified TDW in our mice and found a large decrease in its amount in both MG-Dep

and MG-Repop mice during the baseline dark period (Fig. 4C, D). This reduction was more

pronounced in MG-Repop mice, which was also seen during SD (Fig. 3C, E). TDW is associated

with heightened gamma activity. We analyzed the time course of gamma activity during

wakefulness and found an increase in slow (32-45Hz) frequency bands in the MG-Repop

condition compared to the controls (Fig. 4F). The increase in these frequency bands in MG-Repop

mice is quite clear on time-frequency heat maps in Figure 4a.

Overall, these data suggest a significant imbalance in the theta/gamma-rich waking state in both

the MG-Dep and MG-Repop conditions, and that there is a heightened gamma activity after MG

repopulation.

Disrupted balance of excitatory and inhibitory synaptic signaling and neurotransmitters

upon microglial depletion and repopulation

Neurons communicate with each other at synapses, and synaptic plasticity is associated with

changes in VS [42]. The expression of synaptic proteins increases during wakefulness and sleep

loss and decreases during sleep [43]. Homer1a, a postsynaptic protein and core brain molecular

correlate of sleep loss [44], accumulates in neurons during waking period but is excluded from

synapses due to high expression of noradrenaline [45]. Microglia play a substantial role in synaptic

function, homeostasis, and plasticity in the adult brain. They enhance synaptic activity, prevent

hyperexcitability, and maintain brain network synchronization within a physiological range [46-48].

These functions are dependent on neuronal activity and VS [49].

To investigate whether MG depletion and repopulation induce changes in brain network

connectivity at synaptic and neurochemical levels, we analyzed the excitatory and inhibitory

synaptic density and metabolite content of the brains of our experimental mice. Using an ImageJ

macro, we counted the number of synapses on cortical and hypothalamic neurons (Fig. 5A, B).

We found that the number of excitatory Vglut1-Homer1 synapses on NeuN-positive cortical

somatosensory neurons was significantly reduced in both MG-Dep and MG-Repop conditions.

However, this reduction was more noticeable in MG-Repop mice (Fig. 5C). Interestingly, this

effect was regulated at the postsynaptic level rather than the presynaptic level (Fig. 5D, E). We

furthermore found that the number of parvalbumin (PV)-originated [50] Synaptotagmin-2-Gephyrin

(Syt2-Geph) inhibitory synapses on the cortical somatosensory NeuN neurons was significantly

increased under the MG-Repop condition (Fig. 5F, G). This increase could possibly be due to an

increase in the activity of PV neurons, as we found that the number of Vglut1-Homer1 excitatory

synapses on PV neurons was significantly increased under both MG-Dep and MG-Repop

conditions (Fig. 5H, I). Recently, cortical PV neurons have been proposed to be responsible for

homeostatic sleep regulation [51]. As the quality of VS largely depends on HCRT signaling [30], we

next quantified Vglut1 presynaptic protein synapses that interact with HCRT neurons, based on

their colocalization with Homer1 molecules (Fig. 5J). Our findings indicate a significant decrease

in the number of Vglut1-Homer1 synapses on hypothalamic HCRT neurons, both in MG-Dep and

MG-Repop conditions (Fig. 5K). These results suggest that the function of HCRT neurons may

be impaired when MG are depleted from the brain, and that repopulation fails to restore it.

To further investigate changes in network connectivity in the brains of MG-Dep and MG-Repop

mice, we analyzed additional synaptic markers and specific neurotransmitter receptors on RNA

extracted from cortical samples of these mice. Our bulk analysis revealed that some general

synaptic markers, such as *Synaptophysin I* (*syp*), are largely downregulated in both experimental

mice, while *Slc17a7* and *Homer1* genes are significantly downregulated in the cortex of the MG-

Repop condition. Among the specific neurotransmitter markers analyzed, the adrenergic receptor

family (e.g. *Adrb2*, *adra2a*, *adrb3*, and *Adra1a*) genes showed a tendency to be less expressed

in both MG-Dep and MG-Repop conditions, with a marked decrease in the expression of the

Adra2a gene. Additionally, we observed a reduction in expression of *cholinergic receptor*

muscarinic 2 (*Chrm2*) and serotonergic receptor *5-Hydroxytryptamine Receptor 1A* (*Htr1a*) genes,

which mainly occur in the brain of MG-Repop mice (Fig. 6A).

Synapses serve as the junction between pre- and postsynaptic elements in the brain. However,

their significance goes far beyond just structural interaction, as they play a crucial role in

transmitting neurotransmitters and conveying information that is essential for brain activity. We

therefore conducted targeted metabolomics to analyze the neurotransmitter content and

metabolite reservoir of the cortex, which is the final functional unit and the region where EEG

signals are acquired. This analysis was performed using LC-MS/MS on control, as well as MG-

Dep and MG-Repop mice. The purpose of this analysis was to gain a deeper understanding of

the changes that might occur in the brain of these mice and how these changes could affect the

expression of VS.

Our results showed that pharmacological depletion of MG led to a distinct clustering of

metabolites, as indicated by the principal component analysis (PCA) (Fig. 6B). This suggests that

MG plays an important role in maintaining the overall metabolic composition of the cortex.

However, MG repopulation restored the cortical submetabolome to a pattern similar to that

observed under control experimental conditions (Figure 6B). Univariate analysis revealed a

reduction in norepinephrine upon MG depletion, which was restored after repopulation (Figure

6C). The concentrations of acetylcholine, choline, dopamine, and serotonin remained unchanged

between the experimental conditions, with a slight tendency to decrease in MG-Dep condition.

Furthermore, we observed that MG depletion led to a marked reduction of L-2-amino adipic acid

(L-2-AAA), which was only partially recuperated after repopulation (Figure 6C). We also analyzed

the precursors of neurotransmitters and found a significant decrease in the amino acids L-serine

and asparagine, which could only be partially restored upon MG repopulation (Figure 6D).

Similarly, there was a trend towards a decrease in glycine and L-glutamic acid, although not

significantly (Figure 6D). These findings indicate substantial alterations in metabolic pathways

that play an important role in the regulation of VS.

We showed above that the excitatory synaptic input to HCRT neurons was significantly decreased

in both MG-Dep and MG-Rep conditions. We therefore quantified the *Hcrt* transcript expression

in the hypothalamus of our experimental mice and found a decrease in *Hcrt* expression which

was only partially rescued after MG repopulation (Figure 6E). Taken together, our data suggest

that while network connectivity is significantly impaired in MG-Dep condition with no restoration

after MG repopulation, the underlying substrates (neurotransmitters and metabolites) are largely

restored after MG repopulation.

Discussion

In this study, we uncovered the reciprocal interaction between VS and brain resident macrophages. While the effects of vigilance states on oligodendrocytes [52] and astrocytes [53] have been documented, their effects on brain resident macrophages are less explored. We demonstrated that lack of sleep induces significant transcriptional changes in both MG and CAMs, and these changes are further amplified when hypothalamic HCRT neurons are ablated. Furthermore, we observed regional differential gene expression in cortical and hypothalamic microglia, which is also impacted by SD. These findings suggest that even a few hours of SD can affect the immune compartments of the brain, potentially influencing their function, such as synaptic elimination [22]. While the transcriptional changes in microglia of HCRT neuron-ablated mice is intriguing, it is crucial to explore the mechanisms through which this neuropeptide influences the biology of these cell types.

We also demonstrated that MG depletion has a significant impact on the homeostatic regulation of sleep, and this effect is not fully reversed after MG repopulation. Both NREMS and wakefulness are greatly affected by MG depletion and repopulation, whereas REMS is primarily influenced by MG deficiency, but with significant recovery after MG repopulation. Previous studies have shown that the brain global transcriptomes are little influenced by repopulated microglia [17]. Furthermore, it has been suggested that after MG repopulation, neuroprotection is achieved through IL-6 induction in neurons [54]. Other studies demonstrated full functionality of retinal microglia after repopulation, including their role in the continuous surveillance of the retina, behavioral and physiological responses to retinal injury, and maintenance of synaptic structure [55]. However, our data reveal that MG repopulation not only fails to completely restore various aspects of sleep regulation, but in some cases, it exacerbates certain parameters, such as the fragmentation of NREMS and the increase in sleep need. This demomstrates striking funtional differences between microlgial cells derived from pre-or postantal sources. We speculate that not all microglia-neuronal interactions are properly reestablished by adulthood-dervied microglia after

repopulation. We observed a significant imbalance between excitatory and inhibitory synapses

after depletion of embryo-derived microglia and after repopulation from adult microglia, with a

notable increase in inhibitory synapses during repopulation and a significant rise in excitatory

synapses interacting with PV inhibitory neurons. Previous studies have suggested that microglia

play a role in restraining excessive neuronal activation that cannot be sufficiently suppressed by

inhibitory neurons alone [48]. Our results suggest that prenatal microglia also control excessive

excitation of inhibitory neurons, but the reasons why this is not corrected with adulthood-derived

microglia remain to be investigated.

Deficient synaptic neurotransmission was proposed as a potential cause for abnormal sleep

regulation [56]. We found here an imbalanced neurotransmitter and metabolite content of the brain

in mice devoid of prenatal microglia. It appears that the norepinephrine neurotransmission is

particularly sensitive to MG deficiency, as both the levels of norepinephrine metabolites and the

corresponding receptors are significantly affected by MG depletion. Norepinephrine is implicated

in both wakefulness [21] and sleep regulation [57,58], and it has been suggested that microglial

surveillance process relies heavily on norepinephrine signaling [59,60]. Additionally, we observed a

decrease in the expression of neurotransmitter precursors such as L-serine, which was mirrored

by changes in the concentrations of glycine, asparagine, and glutamic acid. Previous studies have

proposed serine as a link between metabolism and neurotransmission [61], and it serves as a

precursor for synthesizing ceramide [62], which has been suggested to be essential for maintaining

stable wakefulness [35]. Our findings indicate that microglia play a more substantial role in

maintaining the physical connectivity of brain networks rather than the neurochemical

connectivity, as MG repopulation largely restores metabolites but not synapses. To gain a detailed

understanding of microglial regulation of sleep, it is necessary to explore local and specific

microglia depletion or disruption of microglia-neuron interactions using inducible systems.

Additionally, investigating the separate contributions of microglia and CAMs in sleep regulation

using Cre lines under the promoter of *Hexb* [63] and *Mrc1* [64] genes, respectively, would be a starting

point. The observed VS phenotypes may also be influenced by disruptions in other central factors (such as protein phosphorylation orchestrated by microglial TNF-α) [12] and peripheral players (CSF1R inhibition also effects circulating and tissue macrophages) [65] warranting further investigations. Overall, our results shed new light on the complex interaction between the neuroimmune system and sleep regulation, and it paves the way for future investigations in this field.

Data availability

All data are available from the corresponding author upon reasonable request. Bulk RNA sequencing datasets were deposited into the Gene Expression Omnibus database under accession number XXXXX and are available at the following URL: XXXXXXXX.

Acknowledgements

MP was supported by the German Research Foundation (DFG) (TRR167 Project-ID: 259373024, CRC 1479 Project-ID: 441891347, CRC 1160 Project-ID: 256073931, TRR 359 Project-ID: 491676693). MP was further supported by the Novo Nordisk Foundation, the Faber foundation, the Ernst Jung Foundation, the DFG (CRC 992 Project-ID: 192904750, a Reinhart Koselleck grant, the Gottfried Wilhelm Leibniz prize), and the Alzheimer Forschung Initiative e.V. (AFI). This study was further supported by the DFG under Germany's Excellence Strategy (CIBSS, EXC-2189, Project-ID 390939984). MT was supported by Swiss NSF (grant 173126), and the State of Vaud (Faculty of Biology and Medicine, University of Lausanne). AS was supported by Novartis Foundation for medical-biological Research (#23B079).

Author contributions

AS & MP conceptualized and designed the study. AS, MH, SW, VW and LH performed experiments. AS, GM, MB and LH analyzed data. AS, MP, MT, MB, LH and TB wrote the manuscript.

Declaration of interests

446   The authors declare no competing interests.

Methods

Animals. Wilde type (C57BL/6) and Hcrt-ataxin-3 transgenic mice were used in this study. All animals were
between 12 and 14 weeks old during the time of experiments and animal experiments were approved by
the Ministry for Nature, Environment and Consumers` Protection of the state of Baden-Württemberg and
were performed in accordance to the respective national, federal, and institutional regulations. At all times,
care was taken to minimize animal discomfort and avoid pain.

Microglia and CAM isolation and flow cytometry. Mice brains were collected and microglia were
enriched using density gradient separation and were prepared as described previously [66,67]. The cell
suspension was then incubated with Fixable Viability Dye eFluor® 780 (1:1000, 65-0865-14, Invitrogen) for
10 min at 4 °C. Subsequently, the following antibodies were used: antiCD11b (1:200, clone M1/70, 101235
Biolegend), anti-CD45 (1:200, clone 30-F11, 103108 BioLegend), anti-CD11c (1:100, 12-0114-81,
Invitrogen), anti-Gr1 (1:200, 108408, Biolegend) and anti-CD206 (1:200, 141708, Biolegend). Cell sorting
for RNA-seq was performed on a MoFlo Astrios (Beckman Coulter).

Total RNA extraction. Total RNA was extracted from FACS sorted mouse brain cells stabilized in
RNAprotect buffer according to the "Purification of total RNA from animal and human cells" protocol of the
RNeasy   Plus   Micro   Kit   (QIAGEN,   Hilden,   Germany).
In brief, cells were stored and shipped in buffer RNAprotect at 2-8 °C. After pelleting by centrifugation for 5
minutes at 5,000 x g, the RNAprotect was replaced by 350 µl buffer RLT Plus and the samples were
homogenized by vortexing for 30 seconds. Genomic DNA contamination was removed by using gDNA
Eliminator spin columns. Next one volume of 70 % ethanol was added and the samples were applied to
RNeasy MinElute spin columns followed by several wash steps. Finally the total RNA was eluted in 12 µl
of nuclease free water. Purity and integrity of the RNA was assessed on the Agilent 2100 Bioanalyzer with
the RNA 6000 Pico LabChip reagent set (Agilent, Palo Alto, CA, USA).

RNA sequencing. The SMARTer Ultra Low Input RNA Kit for Sequencing v4 (Clontech Laboratories, Inc.,
Mountain View, CA, USA) was used to generate first strand cDNA from approximately 1 ng total-RNA.
Double stranded cDNA was amplified by LD PCR (13 cycles) and purified via magnetic bead clean-up.

Library preparation was carried out as described in the Illumina Nextera XT Sample Preparation Guide (Illumina, Inc., San Diego, CA, USA). Thereby 150 pg of input cDNA were tagmented (tagged and fragmented) by the Nextera XT transposome. The products were purified and amplified via a limited-cycle PCR program to generate multiplexed sequencing libraries. For the PCR step 1:5 dilutions of the unique dual indexing (i7 and I5) adapters were used. The libraries were quantified using the KAPA Library Quantification Kit - Illumina/ABI Prism User Guide (Roche Sequencing Solutions, Inc., Pleasanton, CA, USA). Equimolar amounts of each library were sequenced on an Illumina NextSeq 2000 instrument controlled by the NextSeq 2000 Control Software (NCS) v1.4.0.39521, using one 50 cycles P3 Flow Cell with the dual index, single-read (SR) run parameters. Image analysis and base calling were done by the Real Time Analysis Software (RTA) v3.9.2. The resulting .cbcl files were converted into .fastq files with the bcl2fastq v2.20 software. RNA extraction, library preparation and RNAseq were performed at the Genomics Core Facility "KFB - Center of Excellence for Fluorescent Bioanalytics" (University of Regensburg, Regensburg, Germany; www.kfb-regensburg.de). Bulk RNA-seq data was aligned with STAR v2.7.10a [68] to the mouse reference genome Gencode M26 [69]. Counts were obtained from the bam files with the tool featureCounts v2.0.3. Genes with count expression higher than 5 in at least 3 samples were kept for the differential expression analysis. Differential expression was performed using the limma trend method [70] on the log2 counts per million (CPM) values. Venn diagrams and Volcano plots were generated with the R packages limma and EnhancedVolcano, respectively. Enrichment analysis was performed using ShinyGO [71].

Gene expression analysis. For qPCR analysis, RNA extracted from the cortical samples of control, MG-Dep and MG-Repop mice. RNA was extracted using RNeasy micro kit (74004) and cDNA synthesis was performed using random hexamer primers and M-MLV Reverse Transcriptase Promega kit (M3682). SYBR green-based and Taqman probes were used to detect the gene expression. The assay was performed using a LightCycler480 (Roche).

Immunofluorescence and synapse counting. Mice were anesthetized with ketamine (100mg per kg body weight) and xylazine (10mg per kg body weight) followed by perfusion with 1× PBS. For histology, brains were kept overnight in 4% paraformaldehyde (PFA). Brains were then dehydrated in 30% sucrose for about 48 hours and then snap-frozen on dry ice. Cryosections (14µm for synapse counting and 20µm for microglia and CAM staining) from brain tissue were prepared as free floating in PBS. The following antibodies were used to stain microglia, CAM and synapses: Rabbit-anti-Iba1 (ab178846, abcam), Rat-anti-CD206 (Biorad MCA2235), Goat-anti-Collagene type 4 (Millipore AB769), anti-NeuN (ab177487, ab104224, abcam), anti-Vglut1 (135304, synaptic system), anti-Homer1 (ABN37, Millipore), anti-Gephyrin (147318, synaptic system), anti-Synaptotagmin-2 (ZDB-ATB-081002-25, ZFIN), anti-Parvalbumin (MAB1572, Merck) and anti-Hcrt-A (SC-8070). The DAPI was used to stain the nucleus. Rat-anti-CD206 and Goat-anti-Collagene type 4 antibodies were incubated for 48 hours at 4 °C to have high quality staining. Iba1 antibody was

incubated overnight at 4 °C and the primary antibodies to detect synapses were incubated for 1.45hr at room temperature. The brain cuts were incubated with secondary antibodies for 1-2 hours at room temperature. Fluorescence imaging was performed with TCS SP8 X (Leica Microsystems). Imaging of synapses was performed using 63X objective with zoom factor 8. Iba1+ cells were counted using Fiji. Synapse analysis and quantification were conducted using Fiji macro developed previously [72] with little modifications.

Quantitative profiling of metabolites by liquid chromatography and mass spectrometry (LC-MS/MS). Mice brain tissue biopsies were thawed and homogenised with ice-cold PBS supplemented with 1% protease inhibitor cocktail (Sigma Nr. P8340-5ML), using a cordless pestle motor and disposable pellet mixers (VWR Nr. 47747-366). Buffer volume was adjusted to reach a target of 0.1 to 0.3 mg wet-tissue/mL lysis buffer. Whole tissue homogenates were aliquoted for downstream measurement of total protein concentration, flash-frozen and stored at -80 °C. Sulfur-containing metabolites as well as creatinine, S-adenosylmethionine and S-adenosylhomocysteine were determined according to a previously published procedure [73,74]. Lactate, TCA and glycolysis intermediates and other organic acids, and folates were determined as described in previous work [75]. Amino acids and neurotransmitters were determined using a previously described protocol [76,77]. A commercially available standardized amino acid mixture was utilized to generate a calibration curve for amino acids (Amino acid standards, physiological, Sigma, Nr. A9906-10ML). Calibration curves for all other metabolites were prepared from individual stock solutions prepared in house. Quantitation accuracy was examined by monitoring homocysteine and methylmalonic acid concentrations in an external quality control, namely, the Control Special Assays in Serum, European Research Network for the evaluation and improvement of screening, diagnosis, and treatment of Inherited disorders of Metabolism (ERNDIM) IQCS, SAS-02.1 and SAS-02.2 from MCA Laboratories, Winterswijk, Netherlands. For all other metabolites, quantitation trueness was tested by examining metabolite concentrations in plasma from a previously validated sample isolated from a healthy control individual with respect to standard reference ranges, using the same calibration curves and LC-MS/MS running conditions. Quantification of metabolites was carried out with Analyst® 1.7.2 software, 2022 AB Sciex.

Treatments. In order to eliminate cells dependent on CSF1R (MG) mice were administered a CSF1R inhibitor called PLX5622 (Plexxikon). The inhibitor was combined with AIN-76A standard chow at a concentration of 1,200ppm (Research Diets) and the mice were given unrestricted access to it for two weeks. The control group received AIN-76A standard chow without any modifications. After two weeks the PLX5622 food was replaced with control food to start repopulation of MG.

Mice surgery, data acquisition and vigilance state analysis. Mice surgery and data acquisition were performed as previously described [38]. MATLAB scripts were created to quantify various parameters related to wakefulness, NREMS and REMS. These parameters include time distribution and amount of episodes,

as well as the fragmentation of vigilance states, all of which were defined based on previously established

criteria [38]. TDW analysis was conducted following the method described by Vassalli and Franken [30].

Power spectral analysis. Using Somnologica-3TM (Medcare) software, each 4 s epoch of the EEG signal

was subjected to discrete Fourier transform to determine EEG power density (spectra 0–90 Hz at a

frequency resolution of 0.25 Hz). We used artifact-free, same-state–flanked 4 s epochs to calculate the

mean EEG spectral profiles for each behavioral states and time intervals. To account for differences among

animals in absolute EEG power, the mean values of spectral profiles were converted into percentages

relative to a baseline EEG power reference value of 100%. This reference value was determined for each

mouse by summing the power within the frequency range of 0.75–47.5 Hz across all states and throughout

the first day of baseline recordings. To control for differences in the amount of wakefulness, NREMS and

REMS, this reference was weighted so that the relative contribution of each state was identical for all mice

[78]. Time-frequency heatmaps of EEG power during wakefulness were generated using methods described

previously [79].

Time-course analysis. In order to examine the dynamic changes of EEG power within specific frequency

ranges throughout the day and night, the time-course of the activity of that frequency band was computed

for 4 s epochs scored as state of interest (NREMS, REMS and wake). For this purpose, the epochs for

each state were divided into percentiles, ensuring that each percentile contained a roughly equal number

of epochs. The number of percentiles for NREMS and REMS were: 12 for light periods, 6 for dark periods,

and 8 for the 6-h light period after SD; for waking state: 6 for light periods, 12 for dark periods, 8 for the 6-

h light SD, and 4 for the 6-h light after SD. The average EEG power within the specified frequency range

was computed for each percentile, and subsequently normalized based on the type of analysis. This

normalization was either done relative to the total power of the baseline state or with respect to the power

achieved within the delta frequency band during slow-wave sleep at ZT8-12. Like power density analysis,

single epochs were excluded and only power values of the epochs that themselves, as well as the two

adjacent ones, were scored as artifact-free same-state were included in the analysis.

Theta-gamma cross-frequency coupling. We used the modulation index (MI) to measure the theta-

gamma phase-amplitude coupling [40,80]. Using finite impulse response filters with an order equal to three

cycles of the low cutoff frequency, we first bandpass-filtered EEG signals into theta (6-11 Hz) and fast-

gamma (54-90 Hz) in both forward and reverse directions to eliminate phase distortion. We then estimated

instantaneous phase of theta and the envelope of fast-gamma using the Hilbert transform. Theta phase

was discretized into 18 equal bins (N=18, each 20°) and the average value of fast-gamma envelope within

each bin was calculated. The resulting phase-amplitude histogram (P) was compared with a uniform

distribution (U) using the Kullback-Leibler distance, $D_{KL}(P,U) = \sum_{j=1}^{N} P(j) * log[P(j)/U(j)]$, and normalized

by $log(N)$ to obtain the modulation index, $MI = D_{KL} / log(N)$. To explore possible coupling patterns between

different pairs of low and high frequency bands, we used the comodulogram analysis [80]. We considered 16

frequency bands for phase (1-18 Hz, 1-Hz increments, 2-Hz bandwidth), and 14 frequency bands for

amplitude (15-90 Hz, 5-Hz increments, 10-Hz bandwidth). MI values were then calculated for all these pairs

to obtain the comodulogram graph.

Detection of sleep spindles. NREMS spindles were detected automatically using an optimized wavelet-

based method as previously described [39]. Briefly, the power of EEG signals within 9-16Hz was estimated

using the complex B-spline wavelet function, and smoothed using a 200 ms Hanning window, and then a

threshold equal to 3 SD (SD: standard deviation) above the mean was applied to detect the potential spindle

events. Events shorter than 400 ms or longer than 2 s were discarded. Using band pass-filtered EEG signals

in the spindle range (9–16 Hz), we automatically counted the number of cycles of each detected event and

excluded those with <5 cycles or more than 30 cycles. To discard artefacts, events with a power in the

spindle band lower than 6–8.5 Hz or 16.5–20 Hz power bands were not included.

Statistical analysis. The specific number (n) of replicates employed in each experiment is provided in the

corresponding figure legends. The individuals responsible for conducting SD and scoring the sleep

recordings were unaware of the genotype of the animals, ensuring blinding during these procedures. The

results are presented as mean ± SEM or mean ± SD. Statistical analyses were conducted using GraphPad

Prism. The significance of comparisons was determined using one- or two-way ANOVA or mix effect model

followed by relevant post hoc tests and are reported in the figure legends.

    Figure legends

Figure 1. Large transcriptional changes in the hypothalamic and cortical microglia and
CAMs of WT and Ataxin mice after six hours of SD. (**A**) (left) parenchymal Iba1$^+$ cells
(microglia) in green and perivascular space (yellow arrowheads) and meningeal (white
arrowheads) Iba1$^+$ cells (CAMs) in red. CAMs are positive for CD206 protein expression and are
found in the collagen (Col) IV$^+$ perivascular compartment (right). (**B**) CD11b$^+$; CD45Intermediate
microglia were sorted separately from the cortex and hypothalamus of WT and Ataxin mice.
CD11b and CD206 positive CAMs were sorted from the whole brain of the both genotypes. (**C**)
heatmap showing differentially expressed genes in microglia from hypothalamus and cortex of
WT mice under SD and control conditions. Each column represents microglia gene expression
from one individual mouse, with 4-5 mice per group. Color code represents linear scale. (**D**)
heatmap showing differentially expressed genes in microglia from hypothalamus and cortex of
Ataxin mice under SD and control conditions. (**E**) Volcano plots depicting differentially regulated
cortical and hypothalamic microglial genes between the SD and control groups in WT mice.
NS = not significant. (**F**) Venn diagram depicting the number of differentially regulated and
overlapping genes between microglia of WT and Ataxin mice. Red and dark blue represents up
and downregulated genes respectively. (**G**, **H**, **I** and **J**) Radar plots representing DEGs in microglia
of cortex (Cx) and hypothalamus (Hy) in WT (green dots) and Ataxin (blue dots) mice under SD
(white, upper part) and control (noSD) condition (yellow, bottom). (**K**) Volcano plots depicting
differentially regulated CAMs genes between the SD and control groups in WT mice. NS = not
significant. (**L**) Bar plots showing the expression of *App* and *Lyve-1* genes under the SD and
control conditions in CAMs of WT mice. Values are presented as mean ± SEM, n =4-5 (t-test, *P
< 0.05; **P < 0.01; ***P < 0.001). (**M**) Volcano plots depicting differentially regulated CAMs genes
between the SD and control groups in Ataxin mice.

Figure 2. MG are essential for the proper regulation of NREMS. (**A**) Schematic figure
illustrating the experimental procedure conducted on adult mice to assess the brain's electrical
activity using EEG. Prior to drug administration, the first recordings were obtained. These initial
recordings served as controls. Subsequently, the mice were fed food containing PLX5622 for a
duration of two weeks. The second recording was conducted during the final three days of this
period. Following the drug treatment, the mice were once again provided with control food. Upon
completion of this phase, the third recording was performed. (**B, C**) PLX5622 substantially
depletes the microglia from the brain, and removing it brings the microglia back to the brain. (**D**)

Recording paradigm. The first two days of recordings served as baseline, and were averaged and represented as one day. The second day consisted of 6 hours of SD and 18 hours of recovery. (**E**) Time-course of vigilance states; Wakefulness (top) and NREMS (bottom). MG-Dep and – Repop mice show a large increase in NREMS amount during the dark period both at baseline and during recovery (n = 9 for control and MG-Depleted conditions and n = 6 for MG-Repopulated condition, two-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars are significant differences between control mice and MG-Dep and MG-Repop mice respectively). Data points are shown in minutes per hour (min/h) and represent the average of 2 h. (**F**) From left to right: the total amount of wakefulness (top) and NREMS (bottom) during baseline light, dark and recovery light and dark periods (Mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**G**) Time-course of the EEG delta power (1–4 Hz) in the course of the 3-day recording. EEG delta power denotes sleep need. As shown, both MG-Repop and MG-Dep animals have deficiency to build up sleep need during dark period (two-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars are significant differences between control mice and MG-Dep and MG-Repop mice respectively). (**H**) EEG power spectra of all experimental mice in NREMS during baseline light (left) and dark (right) periods. Insets indicate amplification for specific EEG frequencies (two-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars are significant differences between control mice and MG-Depleted and MG-Repopulated mice respectively). (**I**) EEG power spectra of all experimental mice in NREMS during recovery light (left) and dark (right) periods. (**J**) Distribution of NREMS bout durations during baseline dark period across three experimental mice. Sleep bout length are presented on the x-axis (one-way ANOVA, followed by Tukey test). (**K**) (Top) Heatmap showing the average time–frequency representations of spindles. (Bottom) Representative EEG/EMG signals of a detected spindle. The dashed horizontal lines indicate upper and lower thresholds to detect the spindle using the wavelet energy within 9-16 Hz, and the dashed vertical lines indicate the start and the end of the detected spindle. (**L**) Spindle rate during NREMS episodes of different conditions. Spindle rate significantly decreased in MG-depleted mice (Mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001)).

Figure 3. REMS regulation largely depends on the presence of the microglia. (**A**) Time-course of distribution of REMS amount across three conditions. MG-Dep mice exhibit a large increase in REMs amount during the baseline dark period (n = 9 for control and MG-Dep conditions and n = 6 for MG-Repop condition, two-way ANOVA, followed by Dunnett's test, *P <

0.05; **P < 0.01; ***P < 0.001, orange and blue stars are significant differences between control

mice and MG-Dep and MG-Repop mice respectively). Data points are shown in minutes per hour

(min/h) and represent the average of 2 h. (**B**) Total amount of REMS during baseline dark period

(Mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**C**)

Distribution of REMs bout durations during baseline dark period across three experimental mice.

Sleep bout length are presented on the x-axis (one-way ANOVA, followed by Tukey test). (**D**)

EEG power spectra of all experimental mice in REMS during baseline dark period. Insets indicate

amplification for specific EEG frequencies (two-way ANOVA, followed by Dunnett's test, *P <

0.05; **P < 0.01; ***P < 0.001, orange and blue stars are significant differences between control

mice and MG-Dep and MG-Repop mice respectively). (**E**) Comodulogram graphs show the

modulation index for a wide range of frequency pairs, obtained from 12-h recordings of one animal

in different conditions. (**F**) Normalized theta-gamma coupling across three experimental

conditions. Theta-gamma coupling is significantly decreased in MG-depleted and -repopulated

conditions (Mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P <

0.001)).

.

Figure 4. MG Repopulation fails to normalize the waking state deficits observed under the

MG Depleted condition. (**A**) Heatmaps of time x frequency x power representations of the waking

EEG of control, MG-Dep and MG-Repop mice across the 3 recorded days (48h baseline, a 6h

sleep deprivation and 18h recovery) show large changes in higher frequency bands. (**B**) EEG

power spectra of all experimental mice in wakefulness during baseline light and dark periods (two-

way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange star are

significant differences between control mice and MG-Dep mice). (**C**) Time-course distribution of

TDW amounts indicates decreased amount during the dark period in MG-Dep and MG-Rep. Data

points are shown in minutes/hour and represent the average of 2 hours (baselines are averaged),

(two-way ANOVA followed by Dunnett's test, orange and blue stars are significant differences

between control mice and MG-Dep and MG-Repop mice respectively). (**D**) Quantification of total

amount of TDW during baseline light and dark period (Mixed-effects analysis, followed by

Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**E**) Total amount of TDW during SD. (**F**)

Waking EEG slow-gamma (32-45 Hz) power dynamics in the course of the 3-day recording

(baseline one and two are averaged) (two-way ANOVA, followed by Dunnett's test).

Figure 5. Large synaptic imbalance after MG depletion and Repopulation in the cortex and hypothalamus. (**A**) Schematic figure depicting the cell types on which the number of synapses was counted. NeuN (blue) and PV (orange) positive neurons in the cortex and Hcrt neurons in the hypothalamus. The number of the co-localized excitatory Vglut1 – Homer1 synaptic proteins on NeuN, PV and Hcrt positive neurons were counted. Additionally, the number of the inhibitory Syt2-Gephyrin synaptic proteins on NeuN neurons were counted. (**B**) Representative image depicting pre, postsynaptic proteins and their colocalization with NeuN positive neurons. The synapses are shown with arrowheads and marked with pink color by ImageJ macro (right). Scale bar: 10μm. (**C**) number of vGlut1-Homer1 synapses/μm2 of NeuN+ soma. N=342 cell from 12 mice for control, 324 cell from 11 mice for MG-Dep and 357 cell from 12 mice for MG-Repop conditions. The results show reduced number of these synapses in both MG-Dep and MG-Repop conditions (**D**) Number of Homer1 puncta/μm2 of NeuN+ soma. (**E**) Number of Vglut1 puncta/μm2 of NeuN+ soma. (**F**) Number of Syt2-Gephyrin synapses/μm2 of NeuN+ soma. N=299 cell from 12 mice for control, 270 cell from 11 mice for MG-Dep and 306 cell from 12 mice for MG-Repop conditions. (**G**) Syt2-Gephyrin synapses on NeuN positive cortical neuron. Large increase in this inhibitory is observed under MG-Repop condition. Scale bar: 10μm. (**H**) distribution of the PV positive neurons in the cortex. Scale bar: 100 and 10 μm. (**I**) Number of vGlut1-Homer1 synapses/μm2 of PV+ soma. N=29 cell from 5 mice for control, 29 cell from 5 mice for MG-Dep and 23 cell from 4 mice for MG-Repop conditions. (**J**) Hcrt neurons distribution in the lateral hypothalamus. Red: Hcrt neuron, green (Vglut1) and Cyan (Homer1). Scale bar: 100 and 10 μm. (**K**) number of vGlut1-Homer1 synapses/μm2 of Hcrt+ soma. N=70 cell from 3 mice for control, 73 cell from 3 mice for MG-Dep and 77 cell from 4 mice for MG-Repop conditions. One-way ANOVA, followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

Figure 6. Significant changes occur in neurotransmitter receptors and metabolite content following MG depletion and Repopulation. (**A**) Gene expression level of synaptic proteins and neurotransmitter receptors at RNA level following MG depletion and Repopulation. N=3 for control and MG-Dep and n=4 for MG-Repop conditions. One-way ANOVA, followed by Tukey test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$. (**B**) PCA on targeted metabolites of cortex isolated from control (n=3), MG depleted (n=3) and Repopulated (n=4) animals. (**C**) LC-MS/MS measured neurotransmitter content in the brain of control, MG-Dep and MG-Repop mice. (**D**) Down regulation of amino acids in the brain of experimental mice. (**E**) QPCR analysis of the *Hcrt* gene expression in the hypothalamus of the control, MG-Dep and MG-Repop mice.

Figure S1: Transcriptional Changes in Microglial core genes after SD. (**A**) Bar plots showing
the expression of microglial core genes in the cortex and hypothalamus under SD and NoSD
(control) conditions. The color codes are provided. (**B**) Bar plots showing the expression of *Lyz2*
and *Slc31a2* genes. (**C**) Bar plots showing the expression of the *Adrb2* gene. *P < 0.05; **P <
0.01; ***P < 0.001.

Figure S2: Region and genotype specific transcriptional changes in microglial genes after
SD.

Heatmap showing differentially expressed genes in microglia of the cortex and hypothalamus in
WT and Ataxin mice under SD and control conditions. Each column represents microglia data
from one individual mouse, with 4-5 mice per group. The color code represents a linear scale.

Figure S3: Gene set enrichment analysis for DEGs of microglia and CAM in WT and Ataxin
mice.

(**A**) Gene set enrichment analysis for DEGs in microglia of WT and Ataxin mice (**B**), as well as
CAMs of WT mice (**C**).

References

Amann, L., Masuda, T. & Prinz, M. Mechanisms of myeloid cell entry to the healthy and diseased central nervous system. *Nature immunology* **24**, 393-407 (2023). https://doi.org:10.1038/s41590-022-01415-8

Kierdorf, K., Masuda, T., Jordão, M. J. C. & Prinz, M. Macrophages at CNS interfaces: ontogeny and function in health and disease. *Nature reviews. Neuroscience* **20**, 547-562 (2019). https://doi.org:10.1038/s41583-019-0201-x

Prinz, M., Masuda, T., Wheeler, M. A. & Quintana, F. J. Microglia and Central Nervous System-Associated Macrophages-From Origin to Disease Modulation. *Annual review of immunology* **39**, 251-277 (2021). https://doi.org:10.1146/annurev-immunol-093019-110159

Sankowski, R. *et al.* Multiomic spatial landscape of innate immune cells at human central nervous system borders. *Nat Med* (2023). https://doi.org:10.1038/s41591-023-02673-1

Prinz, M., Jung, S. & Priller, J. Microglia Biology: One Century of Evolving Concepts. *Cell* **179**, 292-311 (2019). https://doi.org:10.1016/j.cell.2019.08.053

Paolicelli, R. C. *et al.* Synaptic pruning by microglia is necessary for normal brain development. *Science (New York, N.Y.)* **333**, 1456-1458 (2011). https://doi.org:10.1126/science.1202529

Chen, Z. *et al.* Microglial displacement of inhibitory synapses provides neuroprotection in the adult brain. *Nature communications* **5**, 4486 (2014). https://doi.org:10.1038/ncomms5486

Tansley, S. *et al.* Microglia-mediated degradation of perineuronal nets promotes pain. *Science (New York, N.Y.)* **377**, 80-86 (2022). https://doi.org:10.1126/science.abl6773

Kohno, K. *et al.* A spinal microglia population involved in remitting and relapsing neuropathic pain. *Science (New York, N.Y.)* **376**, 86-90 (2022). https://doi.org:10.1126/science.abf6805

Cornell, J., Salinas, S., Huang, H. Y. & Zhou, M. Microglia regulation of synaptic plasticity and learning and memory. *Neural regeneration research* **17**, 705-716 (2022). https://doi.org:10.4103/1673-5374.322423

Shigemoto-Mogami, Y., Hoshikawa, K., Goldman, J. E., Sekino, Y. & Sato, K. Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal subventricular zone. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**, 2231-2243 (2014). https://doi.org:10.1523/jneurosci.1619-13.2014

Pinto, M. J. *et al.* Microglial TNFα orchestrates protein phosphorylation in the cortex during the sleep period and controls homeostatic sleep. *The EMBO journal* **42**, e111485 (2023). https://doi.org:10.15252/embj.2022111485

Kierdorf, K. *et al.* Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nature neuroscience* **16**, 273-280 (2013). https://doi.org:10.1038/nn.3318

Schulz, C. *et al.* A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science (New York, N.Y.)* **336**, 86-90 (2012). https://doi.org:10.1126/science.1219179

Bruttger, J. *et al.* Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity* **43**, 92-106 (2015). https://doi.org:10.1016/j.immuni.2015.06.012

Tay, T. L. *et al.* A new fate mapping system reveals context-dependent random or clonal expansion of microglia. *Nature neuroscience* **20**, 793-803 (2017). https://doi.org:10.1038/nn.4547

Huang, Y. *et al.* Repopulated microglia are solely derived from the proliferation of residual microglia after acute depletion. *Nature neuroscience* **21**, 530-540 (2018). https://doi.org:10.1038/s41593-018-0090-8

18   Ramar, K. *et al.* Sleep is essential to health: an American Academy of Sleep Medicine position
statement. *Journal of clinical sleep medicine : JCSM : official publication of the American*
Academy of Sleep Medicine **17**, 2115-2119 (2021). https://doi.org:10.5664/jcsm.9476

19   Irwin, M. R. & Vitiello, M. V. Implications of sleep disturbance and inflammation for Alzheimer's
disease dementia. *The Lancet. Neurology* **18**, 296-306 (2019). https://doi.org:10.1016/s1474-
4422(18)30450-2

20   Saper, C. B., Chou, T. C. & Scammell, T. E. The sleep switch: hypothalamic control of sleep and
wakefulness. *Trends Neurosci* **24**, 726-731 (2001). https://doi.org:10.1016/s0166-
2236(00)02002-6

21   Seifinejad, A., Vassalli, A. & Tafti, M. Neurobiology of cataplexy. *Sleep Med Rev* **60**, 101546
(2021). https://doi.org:10.1016/j.smrv.2021.101546

22   Choudhury, M. E. *et al.* Phagocytic elimination of synapses by microglia during sleep. *Glia* **68**, 44-
59 (2020). https://doi.org:10.1002/glia.23698

23   Corsi, G. *et al.* Microglia modulate hippocampal synaptic transmission and sleep duration along
the light/dark cycle. *Glia* **70**, 89-105 (2022). https://doi.org:10.1002/glia.24090

24   Ingiosi, A. M., Opp, M. R. & Krueger, J. M. Sleep and immune function: glial contributions and
consequences of aging. *Current opinion in neurobiology* **23**, 806-811 (2013).
https://doi.org:10.1016/j.conb.2013.02.003

25   Tafti, M. *et al.* Major histocompatibility class II molecules in the CNS: increased microglial
expression at the onset of narcolepsy in canine model. *The Journal of neuroscience : the official*
journal of the Society for Neuroscience **16**, 4588-4595 (1996).
https://doi.org:10.1523/jneurosci.16-15-04588.1996

26   Tuan, L. H. & Lee, L. J. Microglia-mediated synaptic pruning is impaired in sleep-deprived
adolescent mice. *Neurobiology of disease* **130**, 104517 (2019).
https://doi.org:10.1016/j.nbd.2019.104517

27   Bellesi, M. *et al.* Sleep Loss Promotes Astrocytic Phagocytosis and Microglial Activation in Mouse
Cerebral Cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*
37, 5263-5273 (2017). https://doi.org:10.1523/jneurosci.3981-16.2017

28   Hara, J. *et al.* Genetic ablation of orexin neurons in mice results in narcolepsy, hypophagia, and
obesity. *Neuron* **30**, 345-354 (2001). https://doi.org:10.1016/s0896-6273(01)00293-8

29   Adamantidis, A. R., Zhang, F., Aravanis, A. M., Deisseroth, K. & de Lecea, L. Neural substrates of
awakening probed with optogenetic control of hypocretin neurons. *Nature* **450**, 420-424 (2007).
https://doi.org:10.1038/nature06310

30   Vassalli, A. & Franken, P. Hypocretin (orexin) is critical in sustaining theta/gamma-rich waking
behaviors that drive sleep need. *Proceedings of the National Academy of Sciences of the United*
States of America **114**, E5464-e5473 (2017). https://doi.org:10.1073/pnas.1700983114

31   Dauvilliers, Y. *et al.* Post-H1N1 narcolepsy-cataplexy. *Sleep* **33**, 1428-1430 (2010).
https://doi.org:10.1093/sleep/33.11.1428

32   Drieu, A. *et al.* Parenchymal border macrophages regulate the flow dynamics of the
cerebrospinal fluid. *Nature* **611**, 585-593 (2022). https://doi.org:10.1038/s41586-022-05397-3

33   Zhang, Y. W., Thompson, R., Zhang, H. & Xu, H. APP processing in Alzheimer's disease. *Molecular*
brain **4**, 3 (2011). https://doi.org:10.1186/1756-6606-4-3

34   Roh, J. H. *et al.* Potential role of orexin and sleep modulation in the pathogenesis of Alzheimer's
disease. *The Journal of experimental medicine* **211**, 2487-2496 (2014).
https://doi.org:10.1084/jem.20141788

35   Liu, H. *et al.* Microglia modulate stable wakefulness via the thalamic reticular nucleus in mice.
Nature communications **12**, 4646 (2021). https://doi.org:10.1038/s41467-021-24915-x

36 Dagher, N. N. *et al.* Colony-stimulating factor 1 receptor inhibition prevents microglial plaque
association and improves cognition in 3xTg-AD mice. *Journal of neuroinflammation* **12**, 139
(2015). https://doi.org:10.1186/s12974-015-0366-9
37 Van Hove, H. *et al.* A single-cell atlas of mouse brain macrophages reveals unique transcriptional
identities shaped by ontogeny and tissue environment. *Nature neuroscience* **22**, 1021-1035
(2019). https://doi.org:10.1038/s41593-019-0393-4
38 Seifinejad, A., Li, S., Possovre, M. L., Vassalli, A. & Tafti, M. Hypocretinergic interactions with the
serotonergic system regulate REM sleep and cataplexy. *Nat Commun* **11**, 6034 (2020).
https://doi.org:10.1038/s41467-020-19862-y
39 Bandarabadi, M. *et al.* A role for spindles in the onset of rapid eye movement sleep. *Nature*
communications **11**, 5247 (2020). https://doi.org:10.1038/s41467-020-19076-2
40 Bandarabadi, M. *et al.* Dynamic modulation of theta-gamma coupling during rapid eye
movement sleep. *Sleep* **42**, 1-11 (2019). https://doi.org:10.1093/sleep/zsz182
41 Endo, T. *et al.* Selective REM sleep deprivation in humans: effects on sleep and sleep EEG. *The*
American journal of physiology **274**, R1186-1194 (1998).
https://doi.org:10.1152/ajpregu.1998.274.4.R1186
42 Tononi, G. & Cirelli, C. Sleep and synaptic homeostasis: a hypothesis. *Brain research bulletin* **62**,
143-150 (2003). https://doi.org:10.1016/j.brainresbull.2003.09.004
43 Gilestro, G. F., Tononi, G. & Cirelli, C. Widespread changes in synaptic markers as a function of
sleep and wakefulness in Drosophila. *Science (New York, N.Y.)* **324**, 109-112 (2009).
https://doi.org:10.1126/science.1166673
44 Maret, S. *et al.* Homer1a is a core brain molecular correlate of sleep loss. *Proceedings of the*
National Academy of Sciences of the United States of America **104**, 20090-20095 (2007).
https://doi.org:10.1073/pnas.0710131104
45 Diering, G. H. *et al.* Homer1a drives homeostatic scaling-down of excitatory synapses during
sleep. *Science (New York, N.Y.)* **355**, 511-515 (2017). https://doi.org:10.1126/science.aai8355
46 Merlini, M. *et al.* Microglial G(i)-dependent dynamics regulate brain network hyperexcitability.
Nature neuroscience **24**, 19-23 (2021). https://doi.org:10.1038/s41593-020-00756-7
47 Akiyoshi, R. *et al.* Microglia Enhance Synapse Activity to Promote Local Network
Synchronization. *eNeuro* **5** (2018). https://doi.org:10.1523/eneuro.0088-18.2018
48 Badimon, A. *et al.* Negative feedback control of neuronal activity by microglia. *Nature* **586**, 417-
423 (2020). https://doi.org:10.1038/s41586-020-2777-8
49 Hristovska, I. *et al.* Sleep decreases neuronal activity control of microglial dynamics in mice.
Nature communications **13**, 6273 (2022). https://doi.org:10.1038/s41467-022-34035-9
50 Bouhours, B., Gjoni, E., Kochubey, O. & Schneggenburger, R. Synaptotagmin2 (Syt2) Drives Fast
Release Redundantly with Syt1 at the Output Synapses of Parvalbumin-Expressing Inhibitory
Neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **37**,
4604-4617 (2017). https://doi.org:10.1523/jneurosci.3736-16.2017
51 Kon, K. *et al.* Cortical parvalbumin neurons are responsible for homeostatic sleep rebound
through CaMKII activation. *bioRxiv*, 2023.2004.2029.537929 (2023).
https://doi.org:10.1101/2023.04.29.537929
52 Bellesi, M. *et al.* Effects of sleep and wake on oligodendrocytes and their precursors. *The Journal*
of neuroscience : the official journal of the Society for Neuroscience **33**, 14288-14300 (2013).
https://doi.org:10.1523/jneurosci.5102-12.2013
53 Bellesi, M., de Vivo, L., Tononi, G. & Cirelli, C. Effects of sleep and wake on astrocytes: clues from
molecular and ultrastructural studies. *BMC biology* **13**, 66 (2015).
https://doi.org:10.1186/s12915-015-0176-7

54   Willis, E. F. *et al.* Repopulating Microglia Promote Brain Repair in an IL-6-Dependent Manner.
904    *Cell* **180**, 833-846.e816 (2020). https://doi.org/10.1016/j.cell.2020.02.013
905 55   Zhang, Y. *et al.* Repopulating retinal microglia restore endogenous organization and function
906    under CX3CL1-CX3CR1 regulation. *Science advances* **4**, eaap8492 (2018).
907    https://doi.org/10.1126/sciadv.aap8492
908 56   Guillaumin, M. C. C. *et al.* Deficient synaptic neurotransmission results in a persistent sleep-like
909    cortical activity across vigilance states in mice. *bioRxiv*, 2023.2005.2011.540034 (2023).
910    https://doi.org/10.1101/2023.05.11.540034
911 57   Osorio-Forero, A. *et al.* Noradrenergic circuit control of non-REM sleep substates. *Current*
912    *biology : CB* **31**, 5009-5023.e5007 (2021). https://doi.org/10.1016/j.cub.2021.09.041
913 58   Ma, C. *et al.* Microglia Regulate Sleep via Calcium-Dependent Modulation of Norepinephrine
914    Transmission. *bioRxiv*, 2023.2007.2024.550176 (2023).
915    https://doi.org/10.1101/2023.07.24.550176
916 59   Stowell, R. D. *et al.* Noradrenergic signaling in the wakeful state inhibits microglial surveillance
917    and synaptic plasticity in the mouse visual cortex. *Nature neuroscience* **22**, 1782-1792 (2019).
918    https://doi.org/10.1038/s41593-019-0514-0
919 60   Liu, Y. U. *et al.* Neuronal network activity controls microglial process surveillance in awake mice
920    via norepinephrine signaling. *Nature neuroscience* **22**, 1771-1781 (2019).
921    https://doi.org/10.1038/s41593-019-0511-3
922 61   Maugard, M., Vigneron, P. A., Bolanos, J. P. & Bonvento, G. l-Serine links metabolism with
923    neurotransmission. *Prog Neurobiol* **197**, 101896 (2021).
924    https://doi.org/10.1016/j.pneurobio.2020.101896
925 62   Vávrová, K., Hrabálek, A., Dolezal, P., Holas, T. & Zbytovská, J. L-Serine and glycine based
926    ceramide analogues as transdermal permeation enhancers: polar head size and hydrogen
927    bonding. *Bioorganic & medicinal chemistry letters* **13**, 2351-2353 (2003).
928    https://doi.org/10.1016/s0960-894x(03)00409-8
929 63   Masuda, T. *et al.* Novel Hexb-based tools for studying microglia in the CNS. *Nature immunology*
930    **21**, 802-815 (2020). https://doi.org/10.1038/s41590-020-0707-4
931 64   Masuda, T. *et al.* Specification of CNS macrophage subsets occurs postnatally in defined niches.
932    *Nature* **604**, 740-748 (2022). https://doi.org/10.1038/s41586-022-04596-2
933 65   Lei, F. *et al.* CSF1R inhibition by a small-molecule inhibitor is not microglia specific; affecting
934    hematopoiesis and the function of macrophages. *Proceedings of the National Academy of*
935    *Sciences of the United States of America* **117**, 23336-23338 (2020).
936    https://doi.org/10.1073/pnas.1922788117
937 66   Mezö, C. *et al.* Different effects of constitutive and induced microbiota modulation on microglia
938    in a mouse model of Alzheimer's disease. *Acta neuropathologica communications* **8**, 119 (2020).
939    https://doi.org/10.1186/s40478-020-00988-5
940 67   Mossad, O. *et al.* Gut microbiota drives age-related oxidative stress and mitochondrial damage
941    in microglia via the metabolite N(6)-carboxymethyllysine. *Nature neuroscience* **25**, 295-305
942    (2022). https://doi.org/10.1038/s41593-022-01027-3
943 68   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)* **29**,
944    15-21 (2013). https://doi.org/10.1093/bioinformatics/bts635
945 69   Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE Project.
946    *Genome research* **22**, 1760-1774 (2012). https://doi.org/10.1101/gr.135350.111
947 70   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
948    microarray studies. *Nucleic acids research* **43**, e47 (2015). https://doi.org/10.1093/nar/gkv007

71   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and
plants. *Bioinformatics (Oxford, England)* **36**, 2628-2629 (2020).
https://doi.org/10.1093/bioinformatics/btz931
72   Favuzzi, E. *et al.* GABA-receptive microglia selectively sculpt developing inhibitory circuits. *Cell*
184, 5686 (2021). https://doi.org/10.1016/j.cell.2021.10.009
73   Behringer, S. *et al.* Targeted Metabolic Profiling of Methionine Cycle Metabolites and Redox
Thiol Pools in Mammalian Plasma, Cells and Urine. *Metabolites* **9** (2019).
https://doi.org/10.3390/metabo9100235
74   Bravo, A. C. *et al.* Analysis of S-Adenosylmethionine and S-Adenosylhomocysteine: Method
Optimisation and Profiling in Healthy Adults upon Short-Term Dietary Intervention. *Metabolites*
12, 373 (2022).
75   Hannibal, L. *et al.* Metabolic Profiling in Human Fibroblasts Enables Subtype Clustering in
Glycogen Storage Disease. *Front Endocrinol (Lausanne)* **11**, 579981 (2020).
https://doi.org/10.3389/fendo.2020.579981
76   Maier, J. P. *et al.* Inhibition of metabotropic glutamate receptor III facilitates sensitization to
alkylating chemotherapeutics in glioblastoma. *Cell Death Dis* **12**, 723 (2021).
https://doi.org/10.1038/s41419-021-03937-9
77   Moritz, L., Klotz, K., Grunert, S. C., Hannibal, L. & Spiekerkoetter, U. Metabolic phenotyping in
phenylketonuria reveals disease clustering independently of metabolic control. *Mol Genet*
Metab **138**, 107509 (2023). https://doi.org/10.1016/j.ymgme.2023.107509
78   Franken, P., Malafosse, A. & Tafti, M. Genetic variation in EEG activity during sleep in inbred
mice. *The American journal of physiology* **275**, R1127-1137 (1998).
https://doi.org/10.1152/ajpregu.1998.275.4.R1127
79   Li, S., Franken, P. & Vassalli, A. Bidirectional and context-dependent changes in theta and
gamma oscillatory brain activity in noradrenergic cell-specific Hypocretin/Orexin receptor 1-KO
mice. *Scientific reports* **8**, 15474 (2018). https://doi.org/10.1038/s41598-018-33069-8
80   Tort, A. B. *et al.* Dynamic cross-frequency couplings of local field potential oscillations in rat
striatum and hippocampus during performance of a T-maze task. *Proc Natl Acad Sci U S A* **105**,
20517-20522 (2008). https://doi.org/10.1073/pnas.0810524105

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

C

Figure S1

Figure S2

A

Microglia | Wildtype

- Pos. reg. of protein folding
- Reg. of protein folding
- Chaperone−mediated protein folding
- Protein folding
- Cellular response to unfolded protein
- Cellular response to topologically incorrect protein
- Response to endoplasmic reticulum stress
- Organonitrogen compound catabolic proc.
- Organic substance catabolic proc.

Fold Enrichment

Fold Enrichment
- 25
- 50
- 75
- 100

−log10(FDR)
3
4
5

B

- Virion attachment to host cell
- Adhesion of symbiont to host cell
- Reg. of translational fidelity
- Cytoplasmic translation
- Translation
- Peptide biosynthetic proc.
- Amide biosynthetic proc.
- Peptide metabolic proc.
- Cellular amide metabolic proc.
- Organonitrogen compound biosynthetic proc.

Fold Enrichment

Hcrt-ataxin

Fold Enrichment
- 10
- 20
- 30
- 40

−log10(FDR)
5.0
7.5
10.0
12.5

C

CAMs | Wildtype

- Pos. reg. of tau−protein kinase activity
- Reg. of tau−protein kinase activity
- Reg. of mononuclear cell migration
- Mononuclear cell migration
- Leukocyte chemotaxis
- Cell chemotaxis
- Leukocyte migration
- Chemotaxis
- Taxis
- Defense response

Fold Enrichment

Fold Enrichment
- 50
- 100
- 150

−log10(FDR)
2.2
2.3
2.4
2.5
2.6
2.7

Figure S3