Embryo-derived brain-resident macrophages sustain sleep-wake circuits

Ali Seifinejad[1#*], Mojtaba Bandarabadi[2], Meriem Haddar[2],
Saskia Wundt[1], Anne Vassalli[2], Mehdi Tafti[2*], Gianni Monaco[1*]

[1] *Institute of Neuropathology, Faculty of Medicine, University of Freiburg, Freiburg, Germany*

[2] *Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland*

#Present address: Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland

[*]*Correspondence to:*

Dr. Ali Seifinejad: ali.seifinejad@mail.ch or

Prof. Mehdi Tafti: Mehdi.tafti@unil.ch or

Dr. Gianni Monaco: mongianni1@gmail.com

Summary

Sleep is a complex behavior regulated by a variety of brain cell types. The roles of brain-resident macrophages, such as microglia and CNS-associated macrophages (CAMs), including those derived postnatally, are not well defined. Here, we investigated the reciprocal interaction of brain-resident macrophages and sleep using multimodal high-throughput transcriptional, electrophysiological and metabolomic profiling in mice. We found that sleep deprivation caused profound transcriptional changes in microglia and CAMs, which were intensified by impaired sleep regulation in the absence of the important sleep-regulatory neuropeptide hypocretin/orexin (HCRT). Depletion in embryonically-derived brain macrophages caused increased sleep in the active period, but reduced its quality, reflected in reduced power of brain sleep oscillations. This was observed both for the Non-REM and REM sleep stages. Subsequent repopulation by postnatal brain macrophages unexpectedly failed to reestablish normal sleep-wake patterns and additionally induced sleep fragmentation. Furthermore, brain macrophage depletion caused excitatory-inhibitory synaptic imbalance, which was resistant to repopulation, and led to increased inhibitory synapses. At the metabolite level, the distinct metabolite profile induced by brain macrophage depletion largely returned to normal after repopulation. Our findings suggest a so far largely unknown interaction between brain-resident macrophages and sleep and emphasizes striking functional differences between embryonic and postnatally-derived brain macrophages, paving the way to future exploration of the role of brain macrophages of different origin in sleep disorders and synaptic connectivity.

Keywords: Microglia; CAM, vigilance states, sleep; HCRT

Introduction

Sleep is a complex behavior essential for the normal functioning of animals, consistently linked to various diseases when compromised(1). Sleep regulation involves circadian and homeostatic processes, determining respectively sleep timing and sleep need(2).

Sleep oscillatory patterns rely on the intricate connectivity of the central nervous system (CNS) at synaptic and neurotransmitter levels. Various cell types, primarily neuronal populations located in the hypothalamus, brainstem (BS), and basal forebrain (BF), interact to finely regulate sleep. The expression of synaptic proteins increases during wakefulness and sleep loss, and decreases during sleep(3), and the postsynaptic protein HOMER1A is a well-established core brain molecular correlate of sleep loss (4). Additionally, the release of key neuropeptides and neurotransmitters such as HCRT (also called Orexin), serotonin (5HT), noradrenaline (NA), and acetylcholine across different brain regions actively modulates sleep-wake cycles(5). This intricate interplay manifests in electroencephalographically (EEG) measurable readouts distinguishing wakefulness, non-rapid eye movement sleep (NREMS), and rapid eye movement sleep (REMS), characterized by distinct brain oscillatory activities such as delta waves (0.5-4 Hz) for NREMS and theta waves (6-9 Hz) for REMS in rodents. However, the contribution of non-neuronal cell types, such as brain-resident macrophages, to brain oscillations and sleep architecture remains largely unexplored.

Brain-resident macrophages comprising parenchymal microglia and CAMs(6-8), play significant roles in pathology and protecting the CNS from damage(9). Although the function of CAMs in regulating brain function is not well understood, microglia participate in a range of physiological functions including synaptic pruning(10). Microglial cells and leptomeningeal macrophages (as part of the CAMs) are derived from prenatal sources known as erythromyeloid precursors that engraft into the developing CNS(11, 12) and facilitate the establishment of neuronal networks especially in the first postnatal weeks (10). If embryonic microglial cells are depleted in adulthood, they repopulate from surviving endogenous cells (13). D*e novo* establishment of micoglial networks occurs by surviving cells through clonal

expansion (14) and these postnatal-derived microglial cells exhibit a different transcriptional

signature (15). However, whether adulthood-derived microglia exhibit functional differences

relative to orginal embryo-derived populations is largely unknown.

While a few studies have demonstrated a bidirectional relationship between embryo-derived

microglia and sleep/wake states (16-19), a comprehensive understanding of how these two

compartments modulate each other is still lacking. Specifically, the transcriptional signature of

microglia and CAMs in response to sleep loss remains unknown. Conversely, the extent to

which brain-resident macrophages impact sleep EEG activity requires thorough investigation

through extensive EEG analysis. Moreover, the influence of brain macrophages on sleep-

regulating neurotransmitters, neurochemical content, and synaptic connectivity warrants

further exploration. Furthermore, it remains unclear whether adulthood-derived microglia are

capable of re-establishing neural circuitries in the same manner as prenatal microglial cells,

including synaptic plasticity and neurochemical balance, which are crucial for normal sleep

regulation.

Here, we first studied the impact of sleep deprivation on the transcriptional signature of

microglia and CAMs. Next, through in-depth analysis, we explored whether absence of

embryo-derived microglia and CAMs influences sleep EEG patterns and whether repopulated

postnatal microglia and CAMs can normalize these patterns. Finally, we investigated how the

absence of microglia and CAMs affects brain network connectivity and neurochemistry.

Results

Sleep loss markedly alters the transcriptional profile of microglia and CAMs

We first investigated how lack of sleep impacts brain-resident macrophages. Sleep deprivation (SD) has shown to induce functional alterations in microglia[20, 21], however, its impact on the transcriptional profile of microglia and CAMs has not yet been documented. We thus deprived wild-type (WT) mice of sleep for 6 h (starting from light onset) and immediately afterward isolated microglia (CD11b[+], CD45[intermediate]) and CAMs (CD11b[+], CD206[+], Fig. 1 A-B) from the cortex and hypothalamus, 2 major sleep-regulating centers, and performed gene expression analysis. The control group was not sleep deprived. We extracted RNA and performed bulk RNA sequencing (RNA-seq) to compare the transcriptome between SD and control conditions. After 6 hours SD, no changes were observed in expression of microglial core genes (*Hexb*, *P2ry12*, *Trem2*, and *TMEM119*) across the cortex and hypothalamus (Supplementary Fig. 1A). However, we observed many other differentially expressed genes (DEGs) (110 genes) (Figure 1C). While almost an equal number of genes were differentially regulated in cortex and hypothalamus after SD, we found genes only affected either in the cortex (e.g. *Cul1*, *Ints6l*, and *Ccnl2*) or the hypothalamus (e.g. *Slc1a3*, *Pdia3*, and *Rpn1*) (Supplementary Fig. 2) indicating region-specific microglial response to SD. Gene set enrichment analysis of the DEGs in WT mice revealed an enrichment of pathways related to protein folding, such as 'Positive regulation of protein folding' (Fig. 1D).

We next investigated how SD affects the transcriptome of CAMs. As CAMs are not abundant in the brain, we sorted them from the entire brain. Our results revealed that 6 h of SD induced numerous DEGs in CAMs (Fig. 1E). Among them, both the *App* and *Lyve1* genes (Fig. 1F) were previously linked to Alzheimer's disease (AD) pathogenesis (22, 23). Remarkably, *Lyve1* gene expression almost tripled after 6 h of SD. Gene set enrichment analysis of DEGs of CAMs in WT mice revealed enrichment for pathways related to AD, including 'Positive regulation of tau-protein kinase activity' (Fig. 1G).

To further understand the effect of sleep on brain-resident macrophages, we additionally, analyzed the transcriptome of microglia and CAM in mice with a disrupted regulation of sleep,

the *Hcrt-ataxin-3* transgenic mice (hereafter referred to as Ox-Ataxin mice), which experience

HCRT neuronal degeneration during development (24). HCRT neurons play a critical role in

waking behaviors that drive sleep need (25). Ox-Ataxin mice represent a model for narcolepsy

disease, manifesting excessive daytime sleepiness, fragmentation of wake and sleep.

Transcriptional comparison of SD and control conditions in microglia of OX-Ataxin mice

resulted in a larger number of DEGs compared to WT mice (270 vs 110 genes) (Figure 1H, I),

suggesting that sleep loss significantly increases the transcriptional change in microglia in the

mice with disrupted sleep regulatory pathways. Among the DEGs, some were shared between

the two mouse lines, however, Ox-Ataxin mice exhibited many DEGs that were not

differentially regulated in WT mice in response to sleep loss (e.g. *Naglu*, *Lyz2*, *Rsp9*, *Rps29*,

Stip1) (Fig. I). We additionally found that SD primarily affects cortical, rather than hypothalamic

microglia in Ox-Ataxin mice, indicating that the cortical microglia are more susceptible to

vigilance state instability induced by the lack of wake-promoting neuropeptide HCRT

(Supplementary Fig. 2). While we did not detect the expression of HCRT receptors in

microglia, we observed the expression of downstream circuit receptors, including

noradrenaline and serotonin receptors, in microglia. One such gene is the beta-2 adrenergic

receptor (*Adrb2*), which is predominantly expressed in hypothalamic microglia compared to

cortical microglia (Supplementary Fig. 1B). Gene set enrichment analysis of the DEGs in

microglia of Ox-Ataxin mice revealed an enrichment of pathways like "Virion attachment to

host cell" (Fig. 1J).

Overall, in the two mouse lines, we found microglial genes primarily regulated under the SD

condition, such as *Cirbp*, *Ighd*, and *Npnt* (Fig. 1K). Additionally, we found genes that were

more expressed in the hypothalamus compared to the cortex (Fig. 1L). Furthermore, we

observed that genes such as *Sgk1* and *Pecr* were largely regulated after SD in the

hypothalamus but not in the cortex, suggesting structure-specific microglia changes after SD

(Fig. 1M). Lastly, we found genes that were exclusively regulated in Ox-Ataxin mice such as

Slc31a2, *Lyz2*, and *Ccl4* (Fig. 1N).

Analysis of SD vs control conditions in CAMs of Ox-Ataxin mice surprisingly revealed a

profoundly diminished number of DEGs relative to WT mice (only 6 genes), suggesting that

CAMs show a blunted response to sleep deprivation in absence of a functional HCRT system

(Fig. 1O).

In summary, our data show that sleep significantly influences the transcriptional profiles of

brain-resident macrophages, indicating intricate interactions between these two systems. SD

induces major changes in the transcriptional profile of microglia and CAM genes involved in

disease-related pathways.

Repopulated adulthood-derived microglia are deficient in rebuilding normal NREMS

circuitry

We next examined to what extent embryo-derived brain-resident macrophages contribute to

sleep regulation and whether postnatal-derived macrophages can perform the same function.

To address this, we depleted microglia from the brains of mice for two weeks using colony-

stimulating factor 1 receptor (CSF1R) inhibition via PLX5622 (26), and then allowed them to

repopulate for four weeks (Fig. 2A). CSF1 receptor is vital for microglia survival. Antagonizing

it severely depletes microglia within three days, and upon the termination of treatment,

repopulation can be readily observed within a few days(15). We observed substantial microglia

depletion (>86%) after two weeks of PLX5622 treatment, and complete repopulation four

weeks after withdrawal (Fig. 2B, C).

We recorded EEG signals before depletion (control), after depletion and after repopulation

(Fig. 2A). Since CSF1 inhibition also depletes CAMs(27), the effects we report here are due

to the depletion of both cell types, which we refer to collectively as MG. Our experimental

conditions therefore included control, MG-Dep (microglia/CAM depleted), and MG-Repop

(microglia/CAM repopulation). To evaluate the distribution of vigilance states, we employed

our standard sleep/wake phenotyping protocol(28), with two days of baseline (which were

averaged and presented as one day in the figures) and in the third day mice underwent 6 h of

SD (ZT0-ZT6) and 18 h of recovery (Fig. 2D). In baseline conditions, all experimental mice

demonstrated the characteristic cycling behavior, with higher levels of wakefulness during the

dark period and higher levels of NREMS during the light period and exhibited the increase in

NREMS amounts in response to SD (Fig. 2E, F). However, a significant NREMS increase was

observed in the baseline dark period when comparing the MG-Dep to the control condition.

Interestingly, the repopulation of MG did not normalize this effect (Control: 12.75±0.72 min,

MG-Dep: 16.78±0.77 min, MG-Repop: 16.54±0.70 min, mean ± SEM) (Fig. 2E, F, bottom

panels). Reciprocally, there was a notable reduction in wakefulness during the dark period

(Fig. 2E, F, top panels).

To gain insight into sleep homeostatic regulation in different mouse groups and determine

whether increased sleep is due to increased need for sleep, we analyzed the time course of

NREMS EEG delta power (1-4 Hz, a recognized quantitative readout of sleep need) across

our 3-day recordings. All mice groups showed a typical decline in delta power during the light

period and a strong rebound after SD, indicating homeostatic regulation of sleep need in all

groups (Fig. 2G). However, the buildup in delta power during the baseline dark period was

significantly lower in MG-Dep and MG-Repop mice compared to WT mice (Fig. 2G). This

attenuation was more pronounced in MG-Repop mice and may be partly related to the

increased amount of sleep during the baseline dark period (Fig. 2E, lower panel). In contrast,

when mice were exposed to enforced wakefulness (SD), the EEG delta power increase was

significantly more pronounced in MG-Dep and MG-Repop than in control conditions, and again

this effect was more extensive in MG-Repop than in MG-Dep (Fig. 2G). Altogether, these data

indicate impaired buildup of homeostatic sleep need during the waking period.

Spectral analysis of NREMS revealed that MG-Dep mice exhibited higher slow-delta (1-2 Hz)

activity compared to control mice in baseline light period (Fig. 2H, left). This frequency band

appears particularly affected also in MG-Repop mice, as it was significantly increased in the

recovery light period following SD (Fig. 2I, left). The fast-delta (3-4 Hz) frequency range was

in contrast blunted during both baseline and recovery light periods only in the MG-Dep

condition (Fig. 2H, I, left). During the baseline dark period, large delta power blunting was

observed in both MG-Repop (1.5-3.5 Hz) and MG-Dep (3-5.5 Hz) conditions. These deficits

remained nearly the same during the recovery dark period, indicating that NREMS of both

experimental mice appears to be shallower (Fig. 2H, I, right). Collectively, it can be concluded

that the presence of MG is necessary for the precise regulation of fast and slow-delta power

as a proxy of the sleep homeostatic process. MG repopulation not only fails to restore NREMS

regulation but worsens it.

To determine whether increased NREMS duration is due to enhanced state initiation

(increased number of NREMS episodes) and/or to enhanced state maintenance (increased

NREMS episode duration), we conducted a sleep fragmentation analysis. We found that while

MG-Dep mice have similar sleep bout duration patterns as the control condition, MG-Repop

mice showed a larger number of shorter NREMS bouts (16 seconds [Control: 6.7±0.63, MG-

Dep: 6.8±0.7, MG-Repop: 9.84±1.12]) and 64 seconds [Control: 12.12±1.15, MG-Dep:

11.02±0.81, MG-Repop: 17.63±1.2]) and fewer longer NREMS bouts (>256 seconds [Control:

1.96±0.28, MG-Dep: 1.74±0.33, MG-Repop: 0.62±0.17]), suggesting a strong destabilization

of NREMS in the MG-Repop brain (Fig. 2J).

During NREMS, brief bursts of brain activity in the 9-16 Hz frequency range, called spindles,

are generated by thalamo-cortical interactions and play roles in processes such as learning,

memory, and cognition(29). To determine if MG depletion and repopulation influence spindle

generation, we quantified spindle rate during NREMS episodes and found a significantly

reduced spindle rate in MG-Dep mice compared to the control condition (Fig. 2K, L; see

"Methods"). In contrast to other features of NREMS, MG repopulation was found to normalize

the occurrence rate of spindle events, indicating that MG repopulation partially rescues the

circuit deficits seen after MG depletion.

Altogether, these data suggest that embryo-derived MG are an integral part of NREMS

establishment and maintenance, and repopulated postnatal MG largely fail to fulfill the normal

NREMS regulatory functions of embryo-derived MG.

Depletion of microglia prolongs REMS and degrades its quality

Different neural pathways and neurotransmitters regulate REMS compared to NREMS.

However, the role of MG in the regulation of REMS is not well understood. We quantified

REMS amount across the dark/light periods and found that MG-Dep mice display more REMS

compared to the control and MG-Repop conditions during the baseline dark period (Control:

0.67±0.1, MG-Dep: 1.17±0.21 min/h, mean ± SEM). However, in contrast to NREMS, MG

repopulation corrected the abnormal expression of REMS (MG-Repop: 0.61±0.13 min/h) (Fig.

3A, B). To investigate if the increase in REMS is due to the consolidation of this state, we

performed fragmentation analysis and found that the number of shorter REMS bouts (8 s) is

largely decreased (Control: 8.7±2.05, MG-Dep: 2.8±1.2, MG-Repop: 8.8±3.3) and longer

bouts (>2 min) are increased (Control: 1.5±0.8, MG-Dep: 6.1±1.1, MG-Repop: 4.71±1.4, mean

± SEM) (Fig. 3C) in MG-Dep mice, suggesting stabilization of REMS in the absence of MG.

Theta oscillations dominate the EEG signal during REMS, where theta phase dynamically

modulates gamma amplitude in hippocampal and cortical networks(30). We next investigated

the contribution of MG in theta oscillations and theta-gamma interactions. Spectral analysis of

REMS revealed a significant dampening of theta (6-8 Hz) power in MG-Dep mice during the

baseline dark period (Fig. 3D), which was restored after MG repopulation. We also found that

the theta-gamma coupling is significantly reduced in the MG-Dep condition compared to the

control. Although MG repopulation attempts to bring it back to the normal level, it fails

significantly (Fig. 3E, F).

Altogether, these findings suggest that the presence of MG is necessary for the normal

expression of REMS.

Microglia contribute to the maintenance of active wakefulness

The quality of prior wakefulness contributes to the expression of both NREMS and REMS.

Along these lines, a specific sub-state of prior wakefulness was recently proposed to drive

NREMS need (25), and some aspects of wakefulness may functionally substitute for REMS

(31). To investigate how wakefulness is expressed in our experimental conditions, we

performed a dynamic analysis of the full waking EEG spectrum across three recording days.

This global analysis revealed differential expression of the higher waking frequency bands

during light-dark cycles across three days, and most notably during enforced wakefulness

(SD). As suggested in Figure 4 (Fig. 4A), both MG-Dep and MG-Repop mice exhibit a large

dampening of the lower gamma (≈45-55 Hz) band during the baseline dark period, which is

exacerbated during the recovery dark period. Additionally, the MG-Dep mice exhibit a large

blunting of the higher gamma band (≈70-90 Hz) during SD, which partially normalizes after

MG repopulation. To better quantify the lower frequency bands, we performed PSD analysis

and found that MG-Dep mice exhibit a large increase in slow delta (1.35-1.75 Hz) activity

compared to the control mice (Fig. 4B).

Mice spend a portion of their waking time engaged in exploratory and motivated behaviors,

such as running, nest building, or drinking, often referred to in rodents as "active wakefulness".

These behaviors are associated with the EEG signature called "theta-dominated wakefulness"

(TDW 6.0-9.5 Hz), which has been suggested to be the principal driver of the sleep homeostat

(25). We quantified TDW in our mice and found a large decrease in its amount in both MG-

Dep and MG-Repop mice during the baseline dark period (Fig. 4C, D). This reduction was

more pronounced in MG-Repop mice, which was also seen during SD (Fig. 3C, E). TDW is

associated with heightened gamma activity. We analyzed the time course of gamma activity

during wakefulness and found an increase in slow (32-45Hz) frequency bands in the MG-

Repop condition compared to the controls (Fig. 4F). The increase in these frequency bands

in MG-Repop mice is also apparent on time-frequency heat maps in Figure 4A.

Overall, these data suggest a significant imbalance in the theta/gamma-rich waking state in

both the MG-Dep and MG-Repop conditions, and appearance of heightened gamma activity

after MG repopulation.

Disrupted balance of synaptic signaling upon microglial depletion and repopulation

Sleep is associated with changes in synaptic plasticity(32) and microglia play a substantial

role in synaptic functions, which are dependent on neuronal activity and vigilance states

(VS)(33). To investigate whether MG depletion and repopulation induce changes in brain

network connectivity at the synaptic level, we analyzed a subset of excitatory and inhibitory

synaptic density proteins that are involved in sleep regulation, (e.g., the core brain molecular

correlate of sleep loss, HOMER1) in the brains of our experimental mice.

Using an ImageJ macro, we counted the number of synapses on cortical and hypothalamic

neurons (Fig. 5A, B) and found that the number of excitatory VGLUT1-HOMER1 synapses on

NEUN$^+$ cortical neurons was significantly reduced in both MG-Dep and MG-Repop conditions.

This reduction was more noticeable in MG-Repop mice (Fig. 5C). Interestingly, this effect was

regulated at the postsynaptic level (HOMER1-associated, Fig. 5D) rather than the presynaptic

level (VGLUT1-associated, Fig. 5E). Recently, cortical PV neurons have been proposed to be

responsible for homeostatic sleep regulation(34). We found that the number of parvalbumin

(PV)-originated(35) Synaptotagmin-2-Gephyrin (Syt2-Geph) inhibitory synapses on the

cortical NEUN$^+$ neurons was significantly increased under the MG-Repop condition (Fig. 5F,

G). This increase could possibly be due to an increase in the activity of PV neurons, as we

found that the number of VGLUT1-HOMER1 excitatory synapses on PV neurons was

significantly increased under both MG-Dep and MG-Repop conditions (Fig. 5H, I). We

additionally investigated the changes in the count of VGLUT1-HOMER1 synapses on

representative hypothalamic neurons (HCRT) and found a significant decrease in the number

of VGLUT1-HOMER1 synapses on HCRT neurons in both MG-Dep and MG-Repop conditions

(Fig. 5J, K).

To further investigate changes in network connectivity in the brains of MG-Dep and MG-Repop

mice, we analyzed additional synaptic markers and sleep-specific neurotransmitter receptors

(e.g., receptors of monoaminergic systems and neuropeptides) on RNA extracted from cortical

samples of these mice. Our bulk analysis revealed that some general synaptic markers, such

as *Synaptophysin I* (*Syp*), are largely downregulated in both experimental mice, while *Slc17a7*

(encoding Vesicular glutamate transporter 1) and *Homer1* genes are significantly

downregulated in the cortex of the MG-Repop condition. Among the specific neurotransmitter

markers analyzed, the adrenergic receptor family (e.g. *Adrb2, Adra2a, Adrb3,* and *Adra1a*)

genes showed a tendency to be less expressed in both MG-Dep and MG-Repop conditions,

with a marked decrease in the expression of the *Adra2a* gene. Additionally, we observed a reduction in expression of *Cholinergic receptor muscarinic 2* (*Chrm2*) and the serotonergic receptor *5-Hydroxytryptamine receptor 1A* (*Htr1a*) genes, which mainly occur in the brain of MG-Repop mice (Fig. 5L).

Taken together, these data suggest that brain connectivity at the synaptic level is largely impaired following the depletion of brain-resident macrophages. Repopulating the brain with these macrophages not only failed to normalize connectivity but, in many cases analyzed, worsened it.

Neurotransmitter release is largely restored after the repopulation of resident macrophages

Synapses, in addition to providing structural connections in the brain, play a crucial role in transmitting neurotransmitters essential for the function of complex neural systems. We therefore questioned whether the levels of essential neurotransmitters involved in sleep regulation are affected by the depletion and repopulation of brain-resident macrophages. We conducted targeted metabolomics to analyze the neurotransmitter content and metabolite reservoir of the cortex, target of most neurotransmitters and region where EEG signals are acquired. Assessment of neurotransmitter content was performed using chromatography with tandem mass spectrometry (LC-MS/MS) in the brain of control, as well as MG-Dep and MG-Repop mice. We found that pharmacological depletion of MG led to a distinct clustering of metabolites, as indicated by principal component analysis (PCA) (Fig. 6A). This suggests that MG plays an important role in maintaining the overall metabolic composition of the cortex. MG repopulation however restored the cortical submetabolome to a pattern similar to the one observed under control experimental conditions (Figure 6A). Univariate analysis revealed a reduction in norepinephrine upon MG depletion, which was restored after repopulation (Figure 6B). The concentrations of acetylcholine, choline, dopamine, and serotonin remained unchanged between the experimental conditions, with a slight tendency to decrease in MG-Dep condition.

Furthermore, we observed that MG depletion led to a marked reduction of L-2-amino adipic

acid (L-2-AAA), which was only partially recovered after repopulation (Figure 6B). We also

analyzed the precursors of neurotransmitters and found a significant decrease in the amino

acids L-serine and asparagine under depleted condition, which could be restored upon MG

repopulation (Figure 6C). Similarly, there was a trend towards decreased glycine and L-

glutamic acid content (Figure 6C).

Taken together, our data suggest that while network connectivity is significantly impaired in

MG-Dep condition and resistant to MG repopulation, the underlying substrates

(neurotransmitters and metabolites) are largely restored after MG repopulation.

Discussion

In this study, we investigated the reciprocal interaction between sleep and brain-resident

macrophages. We first analyzed the transcriptional changes in microglia and CAMs in

response to sleep loss. While the effects of sleep on oligodendrocytes(36) and astrocytes(37)

have been documented, its effects on brain-resident macrophages are less explored. We

show that lack of sleep induces significant transcriptional changes in both MG and CAMs,

These changes are further amplified when the normal sleep-regulatory system is impaired, as

seen in the narcolepsy OX-Ataxin mouse model. We found that the altered genes are involved

in disease-related pathways including regulation of protein folding and Tau-protein kinase

activity. These pathways are involved in neurodegeneration(38, 39), and the role of microglia

in sleep loss-associated neurodegeneration is under investigation(17, 40). Our data showed

that removing the important sleep-regulating neuropeptide, HCRT, induces massive

transcriptional changes in both microglia (increased number of DEGs) and CAMs (decreased

number of DEGs). Previous studies have shown that a null mutation in the *Hcrt* gene in an AD

mouse model leads to a significant reduction in Aβ brain pathology(41), and human narcolepsy

patients who lack a functional HCRT system display reduced amyloid burden(42). It can be

speculated that this effect is mediated by either microglia or CAMs. Further investigations are

warranted to decipher these interactions. We additionally found region-specific transcriptional

response to sleep loss in microglia highlighting that local microglia differentially participate in

neural functioning of the brain as previously suggested(43, 44).

In a reciprocal analysis, we also investigated whether embryo-derived MG contribute to sleep

regulation, and whether postnatal-derived MG serve a similar role. We demonstrate that MG

depletion has a significant impact on the homeostatic regulation of sleep. The brain circuits

most vulnerable to microglia depletion and repopulation, causing the changes in sleep EEG

that we document, are still unknown. However, changes in slow-delta activity in NREM sleep,

state stabilization and reduced theta power in REMS, as well as changes in waking slow-delta

activity, were reported in mice with defects in hypothalamic HCRT and brainstem serotonergic

and noradrenergic systems(25, 28, 45, 46). This is suggestive of functionally important local

interactions between microglial populations and sleep regulating circuits, as mentioned above.

We showed that the changes in sleep patterns after MG depletion are not fully reversed after

MG repopulation. Previous studies have shown that the brain global transcriptomes are little

influenced by repopulated microglia(15). Furthermore, it has been suggested that after MG

repopulation, neuroprotection is achieved through IL-6 induction by neurons(47). Other

studies reported full functionality of retinal microglia after repopulation, including continuous

retinal surveillance, maintenance of synaptic structure, and normal behavioral and

physiological responses to retinal injury (48). However, our data reveal that MG repopulation

not only fails to completely restore various aspects of sleep regulation, but in some cases

exacerbates altered parameters, such as fragmentation of NREMS and dysregulation of sleep

need. This demonstrates striking funtional differences between microgial cells derived from

pre-or postantal sources. We speculate that not all microglia-neuronal interactions are

properly reestablished by adulthood-derived microglia after repopulation. Whether a longer

repopulation period might correct the abnormalities observed in our analysis warrants further

investigation. We observed a significant imbalance between excitatory and inhibitory

synapses after depletion of embryo-derived microglia and after repopulation with adult

microglia, with a notable increase in inhibitory synapses following repopulation. Studies in

juvenile mice prevously reported that microglia depletion in the developing brain significantly

increases the number of both excitatory (VGLUT1-HOMER1) and inhibitory (Syt2-Gephyrin)

synapses on cortical neurons, an effect that persisted after repopulation until P30. However,

in the adult (P60) brain, synapse density returned to normal control levels, suggesting that

depleting microglia during development causes long-lasting, albeit not permanent, effects(49).

Our data, however, show that depleting microglia in adulthood, and then repopulating them,

induces different results, with a reduction in excitatory synapses in both MG-Dep and MG-

Repop conditions, and an increase in inhibitory synapses in MG-Repop conditions. In the

developing brain, it has been suggested that MG-Dep impairs synaptic pruning processes and

alters related gene expression(10, 49). Whether the same process or any specific pathways

are induced by MG depletion and repopulation in the adult brain, such as aberrant reactivation

of developmental programs(50), remains to be found.

Furthermore,other studies have suggested that microglia play a role in restraining excessive

neuronal activation that cannot be sufficiently suppressed by inhibitory neurons alone(51). Our

results suggest that embryo-derived microglia also control excessive activity of inhibitory PV

neurons, but the reasons why this is not corrected with adulthood-derived microglia remain to

be investigated.

Deficient synaptic neurotransmission was proposed as a potential cause for abnormal sleep

regulation(52). We found here an imbalanced neurotransmitter and metabolite content of the

brain in mice devoid of prenatal microglia. It appears that norepinephrine neurotransmission

is particularly sensitive to MG deficiency, as both the levels of norepinephrine metabolites and

the corresponding receptors are significantly affected by MG depletion. Norepinephrine is

implicated in both wakefulness(53) and sleep regulation(54-56), and it has been suggested

that microglial surveillance process relies heavily on norepinephrine signaling(57, 58).

Additionally, we observed a decrease in the expression of neurotransmitter precursors such

as L-serine, which was mirrored by changes in the concentrations of glycine, asparagine, and

glutamic acid. Previous studies have proposed serine as a link between metabolism and

neurotransmission (59), and serine serves as a precursor for synthesizing ceramide (60),

which has been suggested to be essential for maintaining stable wakefulness (61).

Our findings indicate that MG affect vigilane state regulation via multiple pathways and play

a more substantial role in maintaining the physical connectivity of brain networks rather than

neurochemical connectivity, as MG repopulation largely restores metabolites but not

synapses. To gain a detailed understanding of MG regulation of sleep, it will be necessary to

explore local specific microglial depletion, and disruption of microglia-neuron interactions

using inducible systems. Additionally, investigating the separate contributions of microglia and

CAMs in sleep regulation using Cre lines under the promoter of *Hexb* (62) and *Mrc1* (63)

genes, respectively, would be a starting point. The observed vigilance state phenotypic

alterations may also be influenced by disruptions in central factors (such as protein

phosphorylation orchestrated by microglial TNF-α)(16), and peripheral players (since CSF1R

inhibition also effects circulating and tissue macrophages)(64) warranting further

investigations. Overall, our results shed new light on the complex interaction between the

neuroimmune system and sleep regulation, and it paves the way for future investigations in

this important physiopathological field.

Data availability

445 All data are available from the corresponding author upon reasonable request. Bulk RNA

446 sequencing datasets were deposited into the Gene Expression Omnibus database under

447 accession number XXXXX and are available at the following URL: XXXXXXXX.

Acknowledgements

We extend our gratitude to the Metabolomics Core Facility at the Department of General

Pediatrics, University of Freiburg, with special thanks to Dr. Luciana Hannibal and Victoria

Wingert for their invaluable support and expertise. AS was supported by Novartis Foundation

for medical-biological Research (#23B079). MB was supported by the Swiss National Science

Foundation (grant 190605). MT was supported by Swiss NSF (grant 201235), and the State

of Vaud (Faculty of Biology and Medicine, University of Lausanne). AV was supported by the

Swiss National Science Foundation (grant 31003A_182613).

Author contributions

AS conceptualized and designed the study. AS, MH, and SW performed experiments. AS,

GM, and MB analyzed data. AS, GM, AV, MT, and MB wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

Methods

Animals. Wild type (C57BL/6J) and Ox-ataxin-3 transgenic mice were used in this study. All animals

were between 12 and 14 weeks old during the time of experiments and animal experiments were

approved by the Ministry for Nature, Environment and Consumers` Protection of the state of Baden-

Württemberg and were performed in accordance to the respective national, federal, and institutional

regulations. At all times, care was taken to minimize animal discomfort and avoid pain.

Microglia and CAM isolation and flow cytometry. Mice brains were collected and microglia were

enriched using density gradient separation and were prepared as described previously (65, 66). The

cell suspension was then incubated with Fixable Viability Dye eFluor® 780 (1:1000, 65-0865-14,

Invitrogen) for 10 min at 4 °C. Subsequently, the following antibodies were used: antiCD11b (1:200,

clone M1/70, 101235 Biolegend), anti-CD45 (1:200, clone 30-F11, 103108 BioLegend), anti-CD11c

(1:100, 12-0114-81, Invitrogen), anti-Gr1 (1:200, 108408, Biolegend) and anti-CD206 (1:200, 141708,

Biolegend). Cell sorting for RNA-seq was performed on a MoFlo Astrios (Beckman Coulter).

Total RNA extraction. Total RNA was extracted from FACS sorted mouse brain cells stabilized in

RNAprotect buffer according to the "Purification of total RNA from animal and human cells" protocol of

the RNeasy Plus Micro Kit (QIAGEN, Hilden, Germany).

In brief, cells were stored and shipped in buffer RNAprotect at 2-8 °C. After pelleting by centrifugation

for 5 minutes at 5,000 x g, the RNAprotect was replaced by 350 µl buffer RLT Plus and the samples

were homogenized by vortexing for 30 seconds. Genomic DNA contamination was removed by using

gDNA Eliminator spin columns. Next one volume of 70 % ethanol was added and the samples were

applied to RNeasy MinElute spin columns followed by several wash steps. Finally the total RNA was

eluted in 12 µl of nuclease free water. Purity and integrity of the RNA was assessed on the Agilent 2100

Bioanalyzer with the RNA 6000 Pico LabChip reagent set (Agilent, Palo Alto, CA, USA).

RNA sequencing. The SMARTer Ultra Low Input RNA Kit for Sequencing v4 (Clontech Laboratories,

Inc., Mountain View, CA, USA) was used to generate first strand cDNA from approximately 1 ng total-

RNA. Double stranded cDNA was amplified by LD PCR (13 cycles) and purified via magnetic bead

clean-up. Library preparation was carried out as described in the Illumina Nextera XT Sample

Preparation Guide (Illumina, Inc., San Diego, CA, USA). Thereby 150 pg of input cDNA were tagmented

(tagged and fragmented) by the Nextera XT transposome. The products were purified and amplified via

a limited-cycle PCR program to generate multiplexed sequencing libraries. For the PCR step 1:5

dilutions of the unique dual indexing (i7 and I5) adapters were used. The libraries were quantified using

the KAPA Library Quantification Kit - Illumina/ABI Prism User Guide (Roche Sequencing Solutions, Inc.,

Pleasanton, CA, USA). Equimolar amounts of each library were sequenced on an Illumina NextSeq

2000 instrument controlled by the NextSeq 2000 Control Software (NCS) v1.4.0.39521, using one 50

cycles P3 Flow Cell with the dual index, single-read (SR) run parameters. Image analysis and base

calling were done by the Real Time Analysis Software (RTA) v3.9.2. The resulting .cbcl files were

converted into .fastq files with the bcl2fastq v2.20 software. RNA extraction, library preparation and

RNAseq were performed at the Genomics Core Facility "KFB - Center of Excellence for Fluorescent

Bioanalytics" (University of Regensburg, Regensburg, Germany; www.kfb-regensburg.de). Bulk RNA-

seq data was aligned with STAR v2.7.10a (67) to the mouse reference genome Gencode M26 (68).

Counts were obtained from the bam files with the tool featureCounts v2.0.3. Genes with count

expression higher than 5 in at least 3 samples were kept for the differential expression analysis.

Differential expression was performed using the limma trend method (69) on the log2 counts per million

(CPM) values. Venn diagrams and Volcano plots were generated with the R packages limma and

EnhancedVolcano, respectively. Enrichment analysis was performed using ShinyGO (70).

Gene expression analysis. For qPCR analysis, RNA extracted from the cortical samples of control,

MG-Dep and MG-Repop mice. RNA was extracted using RNeasy micro kit (74004) and cDNA synthesis

was performed using random hexamer primers and M-MLV Reverse Transcriptase Promega kit

(M3682). SYBR green-based and Taqman probes were used to detect the gene expression. The assay

was performed using a LightCycler480 (Roche).

Immunofluorescence and synapse counting. Mice were anesthetized with ketamine (100 mg per kg

body weight) and xylazine (10 mg per kg body weight) followed by perfusion with 1× PBS. For histology,

brains were kept overnight in 4% paraformaldehyde (PFA). Brains were then dehydrated in 30%

sucrose for about 48 hours and then snap-frozen on dry ice. Cryosections (14µm for synapse counting

and 20µm for microglia and CAM staining) from brain tissue were prepared as free floating in PBS. The

following antibodies were used to stain microglia, CAM and synapses: Rabbit-anti-IBA1 (ab178846,

abcam), Rat-anti-CD206 (Biorad MCA2235), Goat-anti-Collagen type 4 (Millipore AB769), anti-NEUN

(ab177487, ab104224, abcam), anti-VLUT1 (135304, synaptic system), anti-HOMER1 (ABN37,
Millipore), anti-Gephyrin (147318, synaptic system), anti-Synaptotagmin-2 (ZDB-ATB-081002-25,
ZFIN), anti-Parvalbumin (MAB1572, Merck) and anti-HCRT-A (SC-8070). The DAPI was used to stain
the nucleus. Rat-anti-CD206 and Goat-anti-Collagen type 4 antibodies were incubated for 48 hours at
4 °C to have high quality staining. Iba1 antibody was incubated overnight at 4 °C and the primary
antibodies to detect synapses were incubated for 1.45hr at room temperature. The brain cuts were
incubated with secondary antibodies for 1-2 hours at room temperature. Fluorescence imaging was
performed with TCS SP8 X (Leica Microsystems). Imaging of synapses was performed using 63X
objective with zoom factor 8. Iba1+ cells were counted using Fiji. Synapse analysis and quantification
were conducted using Fiji macro developed previously (71) with little modifications.
536
Quantitative profiling of metabolites by liquid chromatography and mass spectrometry (LC-
MS/MS). Mice brain tissue biopsies were thawed and homogenised with ice-cold PBS supplemented
with 1% protease inhibitor cocktail (Sigma Nr. P8340-5ML), using a cordless pestle motor and
disposable pellet mixers (VWR Nr. 47747-366). Buffer volume was adjusted to reach a target of 0.1 to
0.3 mg wet-tissue/mL lysis buffer. Whole tissue homogenates were aliquoted for downstream
measurement of total protein concentration, flash-frozen and stored at -80 °C. Sulfur-containing
metabolites as well as creatinine, S-adenosylmethionine and S-adenosylhomocysteine were
determined according to a previously published procedure (72, 73). Lactate, TCA and glycolysis
intermediates and other organic acids, and folates were determined as described in previous work (74).
Amino acids and neurotransmitters were determined using a previously described protocol (75, 76). A
commercially available standardized amino acid mixture was utilized to generate a calibration curve for
amino acids (Amino acid standards, physiological, Sigma, Nr. A9906-10ML). Calibration curves for all
other metabolites were prepared from individual stock solutions prepared in house. Quantitation
accuracy was examined by monitoring homocysteine and methylmalonic acid concentrations in an
external quality control, namely, the Control Special Assays in Serum, European Research Network for
the evaluation and improvement of screening, diagnosis, and treatment of Inherited disorders of
Metabolism (ERNDIM) IQCS, SAS-02.1 and SAS-02.2 from MCA Laboratories, Winterswijk,
Netherlands. For all other metabolites, quantitation trueness was tested by examining metabolite
concentrations in plasma from a previously validated sample isolated from a healthy control individual
with respect to standard reference ranges, using the same calibration curves and LC-MS/MS running
conditions. Quantification of metabolites was carried out with Analyst® 1.7.2 software, 2022 AB Sciex.
558
559 **Treatments.** In order to eliminate cells dependent on CSF1R (MG) mice were administered a CSF1R
560 inhibitor called PLX5622 (Plexxikon). The inhibitor was combined with AIN-76A standard chow at a
561 concentration of 1,200ppm (Research Diets) and the mice were given unrestricted access to it for two
562 weeks. The control group received AIN-76A standard chow without any modifications. After two weeks
563 the PLX5622 food was replaced with control food to start repopulation of MG.
564

Mice surgery, data acquisition and vigilance state analysis. Mice surgery and data acquisition were performed as previously described (28). MATLAB scripts were created to quantify various parameters related to wakefulness, NREMS and REMS. These parameters include time distribution and amount of episodes, as well as the fragmentation of vigilance states, all of which were defined based on previously established criteria (28). TDW analysis was conducted following the method described by Vassalli and Franken (25).

Power spectral analysis. Using Somnologica-3TM (Medcare) software, each 4 s epoch of the EEG signal was subjected to discrete Fourier transform to determine EEG power density (spectra 0–90 Hz at a frequency resolution of 0.25 Hz). We used artifact-free, same-state–flanked 4 s epochs to calculate the mean EEG spectral profiles for each behavioral states and time intervals. To account for differences among animals in absolute EEG power, the mean values of spectral profiles were converted into percentages relative to a baseline EEG power reference value of 100%. This reference value was determined for each mouse by summing the power within the frequency range of 0.75–47.5 Hz across all states and throughout the first day of baseline recordings. To control for differences in the amount of wakefulness, NREMS and REMS, this reference was weighted so that the relative contribution of each state was identical for all mice (77). Time-frequency heatmaps of EEG power during wakefulness were generated using methods described previously (78).

Time-course analysis. To examine the dynamic changes of EEG power within specific frequency ranges throughout the day and night, the time-course of the activity of that frequency band was computed for 4 s epochs scored as state of interest (NREMS, REMS and wake). For this purpose, the epochs for each state were divided into percentiles, ensuring that each percentile contained a roughly equal number of epochs. The number of percentiles for NREMS and REMS were: 12 for light periods, 6 for dark periods, and 8 for the 6-h light period after SD; for waking state: 6 for light periods, 12 for dark periods, 8 for the 6-h light SD, and 4 for the 6-h light after SD. The average EEG power within the specified frequency range was computed for each percentile, and subsequently normalized based on the type of analysis. This normalization was either done relative to the total power of the baseline state or with respect to the power achieved within the delta frequency band during slow-wave sleep at ZT8-12. Like power density analysis, single epochs were excluded and only power values of the epochs that themselves, as well as the two adjacent ones, were scored as artifact-free same-state were included in the analysis.

Theta-gamma cross-frequency coupling. We used the modulation index (MI) to measure the theta-gamma phase-amplitude coupling (30, 79). Using finite impulse response filters with an order equal to three cycles of the low cutoff frequency, we first bandpass-filtered EEG signals into theta (6-11 Hz) and fast-gamma (54-90 Hz) in both forward and reverse directions to eliminate phase distortion. We then estimated instantaneous phase of theta and the envelope of fast-gamma using the Hilbert transform. Theta phase was discretized into 18 equal bins (N=18, each 20°) and the average value of fast-gamma envelope within each bin was calculated. The resulting phase-amplitude histogram (P) was compared

with a uniform distribution (U) using the Kullback-Leibler distance, $D_{KL}(P, U) = \sum_{j=1}^{N} P(j) *$

$log[P(j)/U(j)]$, and normalized by *log(N)* to obtain the modulation index, *MI = D_{KL} / log(N)*. To explore

possible coupling patterns between different pairs of low and high frequency bands, we used the

comodulogram analysis (79). We considered 16 frequency bands for phase (1-18 Hz, 1-Hz increments,

2-Hz bandwidth), and 14 frequency bands for amplitude (15-90 Hz, 5-Hz increments, 10-Hz bandwidth).

MI values were then calculated for all these pairs to obtain the comodulogram graph.

Detection of sleep spindles. NREMS spindles were detected automatically using an optimized

wavelet-based method as previously described (29). Briefly, the power of EEG signals within 9-16Hz

was estimated using the complex B-spline wavelet function, and smoothed using a 200 ms Hanning

window, and then a threshold equal to 3 SD (SD: standard deviation) above the mean was applied to

detect the potential spindle events. Events shorter than 400 ms or longer than 2 s were discarded. Using

band pass-filtered EEG signals in the spindle range (9–16 Hz), we automatically counted the number

of cycles of each detected event and excluded those with <5 cycles or more than 30 cycles. To discard

artefacts, events with a power in the spindle band lower than 6–8.5 Hz or 16.5–20 Hz power bands

were not included.

Statistical analysis. The specific number (n) of replicates employed in each experiment is provided in

the corresponding figure legends. The individuals responsible for conducting SD and scoring the sleep

recordings were unaware of the genotype of the animals, ensuring blinding during these procedures.

The results are presented as mean ± SEM or mean ± SD. Statistical analyses were conducted using

GraphPad Prism. The significance of comparisons was determined using one- or two-way ANOVA or

mix effect model followed by relevant post hoc tests and are reported in the figure legends.

Figure legends

Figure 1. Large transcriptional changes in the microglia and CAMs of WT and Ox-Ataxin mice after SD (**A**) (Left) Representative confocal image of parenchymal microglia (IBA1$^+$, green) and perivascular (yellow arrowheads) and meningeal (white arrowheads) CAMs (IBA1$^+$ and CD206$^+$, red). (Right) CAMs are positive for CD206 protein expression and are found in the collagen (Col) IV$^+$ perivascular and meningeal compartment. (**B**) Microglia were sorted as CD11b$^+$; CD45^{Intermediate} populations, separately from the cortex and hypothalamus of WT and Ox-Ataxin-Tg mice. CAMs were sorted as CD11b$^+$ and CD206$^+$ populations from the whole brain of both genotypes. (**C**) Heatmap representation showing differentially expressed genes in microglia from hypothalamus and cortex of WT mice under SD or control conditions. Each column represents microglial gene expression from one individual mouse, with 4-5 mice per group. Linear color scale. (**D**) Gene set enrichment analysis for DEGs in microglia of WT mice. (**E**) Heatmaps depicting DEGs in CAMs of WT mice under SD or control conditions. Each column represents CAM gene expression from one individual mouse, with 4-5 mice per group. Linear color scale. (**F**) Bar plots depicting *App* and *Lyve-1* gene expression under SD or control conditions in CAMs of WT mice. Values are mean ± SEM, n=4-5 (t-test, *P < 0.05; **P < 0.01; ***P < 0.001). (**G**) Gene set enrichment analysis for DEGs in CAMs of WT mice. (**H**) Heatmaps depicting DEGs in microglia from hypothalamus and cortex of Ox-Ataxin mice under SD or control conditions. (**I**) Venn diagram depicting the numbers of differentially regulated and overlapping genes between microglia of WT and OX-Ataxin mice. Red and blue represent up- and down-regulated genes, respectively. (**J**) Gene set enrichment analysis for DEGs in microglia of Ox-Ataxin mice. (**K**, **L**, **M** and **N**) Radar plots representing DEGs in microglia of cortex (Cx) and hypothalamus (Hy) in WT (green dots) and Ox-Ataxin (blue dots) mice under SD (white, top) or control (noSD) condition (yellow, bottom). (**O**) Heatmaps depicting DEGs between the SD and control groups in CAMs of Ox-Ataxin mice. Each column represents CAM gene expression from one individual mouse, with 4-5 mice per group. Linear color scale. The number of DEGs in response to sleep loss in CAMs of Ox-Ataxin mice is markedly diminished relative to WT mice

Figure 2. Essential role of MG in regulation of NREMS and associated EEG oscillations.
(**A**) Schematic illustration of the experimental procedure conducted on adult mice to assess the brain's electrical activity using EEG. Recordings prior to PLX5622 drug administration served as controls. Mice were then fed PLX5622-containing food during 2 weeks. During the final 3 days of this 2-week period, a 2^{nd} series of recordings was conducted. Following PLX5622 treatment, mice were provided control food for 4 weeks. After 4 weeks under normal food, a 3^{rd} recording was performed. All 3 recordings were performed on the same mice. (**B, C**) PLX5622 substantially depleted the microglia from the brain within 2 weeks (compare control and MG-Dep). Return to normal food repopulated microglia into the brain after 4 weeks (compare MG-Dep and MG-Repop). (**D**) Recording paradigm. The first 2 days of recordings served as baseline, averaged as one day. The 2^{nd} day consisted of 6 h SD followed by 18 h of recovery. (**E**) Time-course of vigilance states; Wakefulness (top) and NREMS (bottom). Compared to controls (black datapoints), MG-Dep (orange datapoints) and –Repop (blue datapoints) mice show a large increase in NREMS duration during the dark period both at baseline and during recovery (n = 9 for control and MG-Depleted conditions and n = 6 for MG-Repopulated condition, 2-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars, significant differences between control and MG-Dep and MG-Repop mice, respectively). Data points are shown in minutes per hour (min/h) and represent the average of 2 h. (**F**) From left to right: Time in wakefulness (top) and NREMS (bottom) during baseline light, dark and recovery light and dark periods (Mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**G**) Time-course of EEG delta power (1–4 Hz) during the 3-day recording. EEG delta power is a proxy of sleep need. Both MG-Dep and MG-Repop mice show deficiency in building up sleep need during the dark period (2-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars, significant differences between control and MG-Dep and MG-Repop mice, respectively). (**H**) EEG power spectra of experimental mice in NREMS during baseline light (left) and dark (right) periods. EEG power is expressed as % of baseline power of control condition. Insets show amplification of specific EEG frequencies (2-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars indicate significant differences between control and MG-Depleted and MG-Repopulated mice, respectively). (**I**) EEG power spectra of experimental mice in NREMS during recovery light (left) and dark (right) periods. (**J**) Distribution of NREMS bout durations during baseline dark period across the experimental mice. Sleep bout duration is represented on the x-axis (1-way ANOVA, followed by Tukey test). (**K**) (Top) Representative time-frequency heatmap of spindles. (Bottom) Representative EEG/EMG signals of a detected spindle. Dashed horizontal lines indicate upper and lower thresholds used to detect spindles with wavelet energy within 9-16 Hz, and

dashed vertical lines indicate the start and end of the detected spindle. (**L**) Spindle rate during

NREMS episodes in different conditions. Spindle rate significantly decreased in MG-depleted

mice (mixed-effects analysis, followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$)).

Figure 3. REMS regulation largely depends on the presence of the MG. (**A**) Time-course

of REMS duration in control (black), MG-Dep (orange) and MG-Repop (blue) mouse groups.

MG-Dep mice exhibit a large increase in REMS time during the baseline dark period (n = 9 for

control and MG-Dep conditions and n = 6 for MG-Repop condition, 2-way ANOVA, followed

by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, orange and blue stars represent

significant differences between control and MG-Dep and MG-Repop mice, respectively). Data

points are shown in minutes per hour (min/h) and represent the average of 2 h. (**B**) Time in

REMS during baseline dark period (mixed-effects analysis, followed by Dunnett's test, *$P <$

0.05; **$P < 0.01$; ***$P < 0.001$). (**C**) Distribution of REMs bout durations during baseline dark

period across 3 mouse groups. Sleep bout duration is presented as x-axis (1-way ANOVA,

followed by Tukey test). (**D**) EEG power spectra of all experimental mice in REMS during

baseline dark period. Insets show amplification for specific EEG frequencies (2-way ANOVA,

followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, orange and blue stars are

significant differences between control mice and MG-Dep and MG-Repop mice respectively).

(**E**) Comodulogram graphs show the modulation index for a wide range of frequency pairs,

obtained from 12-h recordings of one animal in different conditions. (**F**) Normalized theta-

gamma coupling across 3 experimental conditions. Theta-gamma coupling is significantly

decreased in MG-depleted and -repopulated conditions (mixed-effects analysis, followed by

Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$)).

.

Figure 4. MG Repopulation fails to normalize the waking state deficits induced by MG

depletion. (**A**) Time x frequency x power heatmap representations of the waking EEG of

control, MG-Dep and MG-Repop mice across the 3 recorded days (48 h baseline, 6 h sleep

deprivation and 18 h recovery) show large changes in higher frequency bands. (**B**) EEG power

spectra of experimental mice in wakefulness during baseline light and dark periods (2-way

ANOVA, followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, orange star are

significant differences between control mice and MG-Dep mice). (**C**) Time-course of TDW

amounts indicates decreased TDW time during the dark period in MG-Dep and MG-Rep.

Datapoints are shown in min/h and represent the average of 2 h (baseline days 1 and 2 are

averaged), (2-way ANOVA followed by Dunnett's test, orange and blue stars are significant

differences between control and MG-Dep and MG-Repop mice, respectively). (**D**)

Quantification of time in TDW (min/h) during baseline light and dark period (mixed-effects

analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**E**) Time in TDW

during SD. (**F**) Waking EEG slow-gamma (32-45 Hz) power dynamics during the 3-day

recording (baseline days 1 and 2 are averaged) (2-way ANOVA, followed by Dunnett's test).

Figure 5. Large synaptic imbalance after MG depletion and repopulation. (**A**) Schematic

illustration of the cell types for which the number of synapses was counted. NEUN (blue) and

PV (orange) positive neurons in the cortex and HCRT neurons in the hypothalamus. The

number of co-localized excitatory VGLUT1-HOMER1 synaptic proteins on NEUN, PV and

HCRT positive neurons were counted. Additionally, the number of co-localized inhibitory

SYT2-Gephyrin synaptic proteins on NEUN neurons were counted. (**B**) Representative image

depicting pre, postsynaptic proteins and their colocalization in a NEUN positive neuron.

Synapses (arrowheads) are marked with pink color using ImageJ macro (right). Scale bar:

10μm. (**C**) Number of VGLUT1-HOMER1 synapses/μm^2 of NEUN$^+$ soma. N=342 cells from

12 mice for control, 324 cells from 11 mice for MG-Dep, and 357 cells from 12 mice for MG-

Repop conditions. The data show a reduced number of these synapses in both MG-Dep and

MG-Repop conditions (**D**) Number of HOMER1 puncta/μm^2 of NEUN$^+$ soma. (**E**) Number of

VGLUT1 puncta/μm^2 of NEUN+ soma. (**F**) Number of SYT2-Gephyrin synapses/μm^2 of

NEUN$^+$ soma. N=299 cells from 12 mice for control, 270 cells from 11 mice for MG-Dep, and

306 cells from 12 mice for MG-Repop conditions. (**G**) Representative images depicting SYT2-

Gephyrin synapses in NEUN$^+$ cortical neuron. Large increase in these inhibitory synapses is

observed in the MG-Repop condition. Scale bar: 10 μm. (**H**) Distribution of PV$^+$ neurons (red)

in the cortex. Scale bar: 100 μm (left), 10 μm (right). (**I**) Number of vGLUT1-HOMER1

synapses/μm^2 of PV$^+$ soma. N=29 cells from 5 mice for control, 29 cells from 5 mice for MG-

Dep, and 23 cells from 4 mice for MG-Repop conditions. (**J**) Left, distribution of HCRT$^+$

neurons (red) in the lateral hypothalamus. Right, a HCRT$^+$ neuron stained with VGLUT1

(green) and HOMER1 (Cyan). Scale bar: 100 and 10 μm. (**K**) Number of VGLUT1-HOMER1

synapses/μm^2 of HCRT+ soma. N=70 cells from 3 mice for control, 73 cells from 3 mice for

MG-Dep, and 77 cells from 4 mice for MG-Repop conditions. (**L**) Gene expression of synaptic

proteins and neurotransmitter receptors at RNA level following MG depletion and repopulation.

N=3 for control and MG-Dep and n=4 for MG-Repop conditions. 1-way ANOVA, followed by

Tukey test, *P < 0.05; **P < 0.01; ***P < 0.001.

Figure 6. MG repopulation largely restores the metabolite changes induced by

microglial depletion. (**A**) PCA of targeted metabolites in cortex isolated from control (n=3),

MG depleted (n=3) and repopulated (n=4) mice. (**B**) LC-MS/MS assessment of the

neurotransmitter content in the brain of control, MG-Dep and MG-Repop mice. (**C**)

Downregulation of amino acids in the brain of experimental mice after MG depletion. 1-way

ANOVA, followed by Tukey test, *P < 0.05; **P < 0.01; ***P < 0.001.

Figure S1: Transcriptional changes in microglial core genes following sleep

deprivation. (**A**) Bar plots showing the expression of microglial core genes in the cortex and

hypothalamus under SD and NoSD (control) conditions. The color codes are provided. (**B**) Bar

plots showing the expression of the *Adrb2* gene. *P < 0.05; **P < 0.01; ***P < 0.001.

Figure S2: Region and genotype specific transcriptional changes in microglial genes

following sleep deprivation.

Heatmap showing differentially expressed genes in microglia of the cortex and hypothalamus

in WT and Ox-Ataxin mice under SD and control conditions. Each column represents microglia

data from one individual mouse, with 4-5 mice per group. The color code represents a linear

scale.

References

1. K. Ramar *et al.*, Sleep is essential to health: an American Academy of Sleep Medicine position
statement. *J Clin Sleep Med* 10.5664/jcsm.9476 (2021).

2. A. A. Borbély, A two process model of sleep regulation. *Hum Neurobiol* **1**, 195-204 (1982).

3. G. F. Gilestro, G. Tononi, C. Cirelli, Widespread changes in synaptic markers as a function of
sleep and wakefulness in Drosophila. *Science* **324**, 109-112 (2009).

4. S. Maret *et al.*, Homer1a is a core brain molecular correlate of sleep loss. *Proc Natl Acad Sci*
U S A **104**, 20090-20095 (2007).

5. C. B. Saper, T. C. Chou, T. E. Scammell, The sleep switch: hypothalamic control of sleep and
wakefulness. *Trends Neurosci* **24**, 726-731 (2001).

6. L. Amann, T. Masuda, M. Prinz, Mechanisms of myeloid cell entry to the healthy and
diseased central nervous system. *Nat Immunol* **24**, 393-407 (2023).

7. K. Kierdorf, T. Masuda, M. J. C. Jordão, M. Prinz, Macrophages at CNS interfaces: ontogeny
and function in health and disease. *Nat Rev Neurosci* **20**, 547-562 (2019).

8. M. Prinz, T. Masuda, M. A. Wheeler, F. J. Quintana, Microglia and Central Nervous System-
Associated Macrophages-From Origin to Disease Modulation. *Annual review of immunology*
39, 251-277 (2021).

9. M. Prinz, S. Jung, J. Priller, Microglia Biology: One Century of Evolving Concepts. *Cell* **179**,
292-311 (2019).

10. R. C. Paolicelli *et al.*, Synaptic pruning by microglia is necessary for normal brain
development. *Science (New York, N.Y.)* **333**, 1456-1458 (2011).

11. K. Kierdorf *et al.*, Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-
dependent pathways. *Nature neuroscience* **16**, 273-280 (2013).

12. C. Schulz *et al.*, A lineage of myeloid cells independent of Myb and hematopoietic stem cells.
Science (New York, N.Y.) **336**, 86-90 (2012).

13. J. Bruttger *et al.*, Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in
the Mammalian Central Nervous System. *Immunity* **43**, 92-106 (2015).

14. T. L. Tay *et al.*, A new fate mapping system reveals context-dependent random or clonal
expansion of microglia. *Nature neuroscience* **20**, 793-803 (2017).

15. Y. Huang *et al.*, Repopulated microglia are solely derived from the proliferation of residual
microglia after acute depletion. *Nat Neurosci* **21**, 530-540 (2018).

16. M. J. Pinto *et al.*, Microglial TNFα orchestrates protein phosphorylation in the cortex during
the sleep period and controls homeostatic sleep. *Embo j* **42**, e111485 (2023).

17. M. E. Choudhury *et al.*, Phagocytic elimination of synapses by microglia during sleep. *Glia* **68**,
44-59 (2020).

18. G. Corsi *et al.*, Microglia modulate hippocampal synaptic transmission and sleep duration
along the light/dark cycle. *Glia* **70**, 89-105 (2022).

19. A. M. Ingiosi, M. R. Opp, J. M. Krueger, Sleep and immune function: glial contributions and
consequences of aging. *Current opinion in neurobiology* **23**, 806-811 (2013).

20. L. H. Tuan, L. J. Lee, Microglia-mediated synaptic pruning is impaired in sleep-deprived
adolescent mice. *Neurobiol Dis* **130**, 104517 (2019).

21. M. Bellesi *et al.*, Sleep Loss Promotes Astrocytic Phagocytosis and Microglial Activation in
Mouse Cerebral Cortex. *J Neurosci* **37**, 5263-5273 (2017).

22. A. Drieu *et al.*, Parenchymal border macrophages regulate the flow dynamics of the
cerebrospinal fluid. *Nature* **611**, 585-593 (2022).

23. Y. W. Zhang, R. Thompson, H. Zhang, H. Xu, APP processing in Alzheimer's disease. *Mol Brain*
4, 3 (2011).

24. J. Hara *et al.*, Genetic ablation of orexin neurons in mice results in narcolepsy, hypophagia,
and obesity. *Neuron* **30**, 345-354 (2001).

25. A. Vassalli, P. Franken, Hypocretin (orexin) is critical in sustaining theta/gamma-rich waking
behaviors that drive sleep need. *Proc Natl Acad Sci U S A* **114**, E5464-e5473 (2017).

26. N. N. Dagher *et al.*, Colony-stimulating factor 1 receptor inhibition prevents microglial
plaque association and improves cognition in 3xTg-AD mice. *J Neuroinflammation* **12**, 139
(2015).
27. H. Van Hove *et al.*, A single-cell atlas of mouse brain macrophages reveals unique
transcriptional identities shaped by ontogeny and tissue environment. *Nat Neurosci* **22**,
1021-1035 (2019).
28. A. Seifinejad, S. Li, M. L. Possovre, A. Vassalli, M. Tafti, Hypocretinergic interactions with the
serotonergic system regulate REM sleep and cataplexy. *Nat Commun* **11**, 6034 (2020).
29. M. Bandarabadi *et al.*, A role for spindles in the onset of rapid eye movement sleep. *Nat*
Commun **11**, 5247 (2020).
30. M. Bandarabadi *et al.*, Dynamic modulation of theta-gamma coupling during rapid eye
movement sleep. *Sleep* **42**, 1-11 (2019).
31. T. Endo *et al.*, Selective REM sleep deprivation in humans: effects on sleep and sleep EEG.
Am J Physiol **274**, R1186-1194 (1998).
32. G. Tononi, C. Cirelli, Sleep and synaptic homeostasis: a hypothesis. *Brain Res Bull* **62**, 143-150
(2003).
33. I. Hristovska *et al.*, Sleep decreases neuronal activity control of microglial dynamics in mice.
Nat Commun **13**, 6273 (2022).
34. K. Kon *et al.*, Cortical parvalbumin neurons are responsible for homeostatic sleep rebound
through CaMKII activation. *bioRxiv* 10.1101/2023.04.29.537929, 2023.2004.2029.537929
(2023).
35. B. Bouhours, E. Gjoni, O. Kochubey, R. Schneggenburger, Synaptotagmin2 (Syt2) Drives Fast
Release Redundantly with Syt1 at the Output Synapses of Parvalbumin-Expressing Inhibitory
Neurons. *J Neurosci* **37**, 4604-4617 (2017).
36. M. Bellesi *et al.*, Effects of sleep and wake on oligodendrocytes and their precursors. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **33**, 14288-14300
(2013).
37. M. Bellesi, L. de Vivo, G. Tononi, C. Cirelli, Effects of sleep and wake on astrocytes: clues
from molecular and ultrastructural studies. *BMC biology* **13**, 66 (2015).
38. S. Dujardin *et al.*, Tau molecular diversity contributes to clinical heterogeneity in Alzheimer's
disease. *Nat Med* **26**, 1256-1263 (2020).
39. N. Naidoo, Cellular stress/the unfolded protein response: relevance to sleep and sleep
disorders. *Sleep Med Rev* **13**, 195-204 (2009).
40. L. Rajendran, R. C. Paolicelli, Microglia-Mediated Synapse Loss in Alzheimer's Disease. *J*
Neurosci **38**, 2911-2919 (2018).
41. J. H. Roh *et al.*, Potential role of orexin and sleep modulation in the pathogenesis of
Alzheimer's disease. *The Journal of experimental medicine* **211**, 2487-2496 (2014).
42. A. Gabelle *et al.*, Reduced brain amyloid burden in elderly patients with narcolepsy type 1.
Ann Neurol **85**, 74-83 (2019).
43. Y. L. Tan, Y. Yuan, L. Tian, Microglial regional heterogeneity and its role in the brain. *Mol*
Psychiatry **25**, 351-367 (2020).
44. P. Ayata *et al.*, Epigenetic regulation of brain region-specific microglia clearance activity. *Nat*
Neurosci **21**, 1049-1060 (2018).
45. S. R. Morairty *et al.*, A role for cortical nNOS/NK1 neurons in coupling homeostatic sleep
drive to EEG slow wave activity. *Proc Natl Acad Sci U S A* **110**, 20272-20277 (2013).
46. C. Cirelli, R. Huber, A. Gopalakrishnan, T. L. Southard, G. Tononi, Locus ceruleus control of
slow-wave homeostasis. *J Neurosci* **25**, 4503-4511 (2005).
47. E. F. Willis *et al.*, Repopulating Microglia Promote Brain Repair in an IL-6-Dependent Manner.
Cell **180**, 833-846.e816 (2020).
48. Y. Zhang *et al.*, Repopulating retinal microglia restore endogenous organization and function
under CX3CL1-CX3CR1 regulation. *Sci Adv* **4**, eaap8492 (2018).

49. E. Favuzzi *et al.*, GABA-receptive microglia selectively sculpt developing inhibitory circuits.
Cell **184**, 4048-4063.e4032 (2021).
50. D. K. Wilton, L. Dissing-Olesen, B. Stevens, Neuron-Glia Signaling in Synapse Elimination.
Annu Rev Neurosci **42**, 107-127 (2019).
51. A. Badimon *et al.*, Negative feedback control of neuronal activity by microglia. *Nature* **586**,
417-423 (2020).
52. M. C. C. Guillaumin *et al.*, Deficient synaptic neurotransmission results in a persistent sleep-
like cortical activity across vigilance states in mice. *bioRxiv* 10.1101/2023.05.11.540034,
2023.2005.2011.540034 (2023).
53. A. Seifinejad, A. Vassalli, M. Tafti, Neurobiology of cataplexy. *Sleep Med Rev* **60**, 101546
(2021).
54. A. Osorio-Forero *et al.*, Noradrenergic circuit control of non-REM sleep substates. *Current*
biology : CB **31**, 5009-5023.e5007 (2021).
55. C. Ma *et al.*, Microglia Regulate Sleep via Calcium-Dependent Modulation of Norepinephrine
Transmission. *bioRxiv* 10.1101/2023.07.24.550176, 2023.2007.2024.550176 (2023).
56. C. Ma *et al.*, Microglia regulate sleep through calcium-dependent modulation of
norepinephrine transmission. *Nat Neurosci* **27**, 249-258 (2024).
57. R. D. Stowell *et al.*, Noradrenergic signaling in the wakeful state inhibits microglial
surveillance and synaptic plasticity in the mouse visual cortex. *Nat Neurosci* **22**, 1782-1792
(2019).
58. Y. U. Liu *et al.*, Neuronal network activity controls microglial process surveillance in awake
mice via norepinephrine signaling. *Nat Neurosci* **22**, 1771-1781 (2019).
59. M. Maugard, P. A. Vigneron, J. P. Bolanos, G. Bonvento, l-Serine links metabolism with
neurotransmission. *Prog Neurobiol* **197**, 101896 (2021).
60. K. Vávrová, A. Hrabálek, P. Dolezal, T. Holas, J. Zbytovská, L-Serine and glycine based
ceramide analogues as transdermal permeation enhancers: polar head size and hydrogen
bonding. *Bioorg Med Chem Lett* **13**, 2351-2353 (2003).
61. H. Liu *et al.*, Microglia modulate stable wakefulness via the thalamic reticular nucleus in
mice. *Nat Commun* **12**, 4646 (2021).
62. T. Masuda *et al.*, Novel Hexb-based tools for studying microglia in the CNS. *Nat Immunol* **21**,
802-815 (2020).
63. T. Masuda *et al.*, Specification of CNS macrophage subsets occurs postnatally in defined
niches. *Nature* **604**, 740-748 (2022).
64. F. Lei *et al.*, CSF1R inhibition by a small-molecule inhibitor is not microglia specific; affecting
hematopoiesis and the function of macrophages. *Proc Natl Acad Sci U S A* **117**, 23336-23338
(2020).
65. C. Mezö *et al.*, Different effects of constitutive and induced microbiota modulation on
microglia in a mouse model of Alzheimer's disease. *Acta neuropathologica communications*
8, 119 (2020).
66. O. Mossad *et al.*, Gut microbiota drives age-related oxidative stress and mitochondrial
damage in microglia via the metabolite N(6)-carboxymethyllysine. *Nature neuroscience* **25**,
295-305 (2022).
67. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
68. J. Harrow *et al.*, GENCODE: the reference human genome annotation for The ENCODE
Project. *Genome Res* **22**, 1760-1774 (2012).
69. M. E. Ritchie *et al.*, limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
70. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and
plants. *Bioinformatics* **36**, 2628-2629 (2020).
71. E. Favuzzi *et al.*, GABA-receptive microglia selectively sculpt developing inhibitory circuits.
Cell **184**, 5686 (2021).

72. S. Behringer *et al.*, Targeted Metabolic Profiling of Methionine Cycle Metabolites and Redox
Thiol Pools in Mammalian Plasma, Cells and Urine. *Metabolites* **9** (2019).
73. A. C. Bravo *et al.*, Analysis of S-Adenosylmethionine and S-Adenosylhomocysteine: Method
Optimisation and Profiling in Healthy Adults upon Short-Term Dietary Intervention.
Metabolites **12**, 373 (2022).
74. L. Hannibal *et al.*, Metabolic Profiling in Human Fibroblasts Enables Subtype Clustering in
Glycogen Storage Disease. *Front Endocrinol (Lausanne)* **11**, 579981 (2020).
75. J. P. Maier *et al.*, Inhibition of metabotropic glutamate receptor III facilitates sensitization to
alkylating chemotherapeutics in glioblastoma. *Cell Death Dis* **12**, 723 (2021).
76. L. Moritz, K. Klotz, S. C. Grunert, L. Hannibal, U. Spiekerkoetter, Metabolic phenotyping in
phenylketonuria reveals disease clustering independently of metabolic control. *Mol Genet
Metab* **138**, 107509 (2023).
77. P. Franken, A. Malafosse, M. Tafti, Genetic variation in EEG activity during sleep in inbred
mice. *Am J Physiol* **275**, R1127-1137 (1998).
78. S. Li, P. Franken, A. Vassalli, Bidirectional and context-dependent changes in theta and
gamma oscillatory brain activity in noradrenergic cell-specific Hypocretin/Orexin receptor 1-
KO mice. *Sci Rep* **8**, 15474 (2018).
79. A. B. Tort *et al.*, Dynamic cross-frequency couplings of local field potential oscillations in rat
striatum and hippocampus during performance of a T-maze task. *Proc Natl Acad Sci U S A*
105, 20517-20522 (2008).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

Figure S1

Figure S2