Contrasting contribution of embryo- and postnatally-derived brain-resident macrophages in sustaining sleep-wake circuitry

Ali Seifinejad[1#*], Mojtaba Bandarabadi[2], Meriem Haddar[2],

Saskia Wundt[1], Mehdi Tafti[2], Anne Vassalli[2], Abbas Khani[3], Gianni Monaco[1*]

1 Institute of Neuropathology, Faculty of Medicine, University of Freiburg, Freiburg, Germany

2 Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland

3Institute of Neuroinformatics, Federal Institute of Technology (ETH) Zurich, Switzerland

#Present address: Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland

*Correspondence to:

Dr. Ali Seifinejad: ali.seifinejad@mail.ch or

Dr. Gianni Monaco: mongianni1@gmail.com

Summary

Sleep is a complex behavior regulated by various brain cell types. However, the roles of brain-

resident macrophages, including microglia and CNS-associated macrophages (CAMs),

particularly those derived postnatally, in sleep regulation remain poorly understood. Here, we

investigated the effects of natural (embryo-derived) and repopulated (postnatally derived)

brain-resident macrophages on the regulation of vigilance states. We found that depletion in

embryonically-derived brain macrophages caused increased sleep in the active period, but

reduced its quality, reflected in reduced power of brain sleep oscillations. This was observed

both for the Non-REM and REM sleep stages. Subsequent repopulation by postnatal brain

macrophages unexpectedly failed to reestablish normal sleep-wake patterns and additionally

induced sleep fragmentation. Furthermore, brain macrophage depletion caused excitatory-

inhibitory synaptic imbalance, which was resistant to repopulation, and led to increased

inhibitory synapses. At the metabolite level, the distinct metabolite profile induced by brain

macrophage depletion largely returned to normal after repopulation. Our findings suggest a

so far largely unknown interaction between brain-resident macrophages and sleep and

emphasizes striking functional differences between embryonic and postnatally-derived brain

macrophages, paving the way to future exploration of the role of brain macrophages of

different origin in sleep disorders and synaptic connectivity.

Keywords: Microglia, CAMs, EEG, vigilance states, sleep, synapse

Introduction

Sleep is a complex behavior essential for the normal functioning of animals, consistently linked to various diseases when compromised[1]. Sleep regulation involves circadian and homeostatic processes, determining respectively sleep timing and sleep need [2].

Sleep oscillatory patterns rely on the intricate connectivity of the central nervous system (CNS) at synaptic and neurotransmitter levels. Various cell types, primarily neuronal populations located in the hypothalamus, brainstem (BS), and basal forebrain (BF), interact to finely regulate sleep. The expression of synaptic proteins increases during wakefulness and sleep loss, and decreases during sleep[3], and the postsynaptic protein HOMER1A is a well-established core brain molecular correlate of sleep loss [4]. Additionally, the release of key neuropeptides and neurotransmitters such as HCRT (also called Orexin), serotonin (5HT), noradrenaline (NA), and acetylcholine across different brain regions actively modulates sleep-wake cycles[5]. This intricate interplay manifests in electroencephalographically (EEG) measurable readouts distinguishing wakefulness, non-rapid eye movement sleep (NREMS), and rapid eye movement sleep (REMS), characterized by distinct brain oscillatory activities such as delta waves (0.5-4 Hz) for NREMS and theta waves (6-9 Hz) for REMS in rodents. However, the contribution of non-neuronal cell types, such as brain-resident macrophages, to brain oscillations and sleep architecture remains largely unexplored.

Brain-resident macrophages comprising parenchymal microglia and CAMs [6-8], play significant roles in pathology and protecting the CNS from damage [9]. They participate in a range of physiological functions including synaptic pruning, neurogenesis, immune surveillance [10-12], and are derived from prenatal sources known as erythromyeloid precursors that engraft into the developing CNS [13,14], facilitating the establishment of neuronal networks especially in the first postnatal weeks [10]. If embryo-derived brain-resident macrophage cells are depleted in adulthood, they repopulate from surviving endogenous cells [15]. De novo establishment of micoglial networks occurs by surviving cells through clonal expansion [16] and these postnatal-derived microglial cells exhibit a different transcriptional signature [17].

However, whether adulthood-derived microglia exhibit functional differences relative to orginal

embryo-derived populations is hotly disputed.

There is evidence that immune system activation, such as during illness, results in increased

sleepiness, suggesting an interaction between the immune system and the sleep-wake cycle

[18,19]. Brain-resident macrophages, being the closest immune components to sleep-regulating

centers in the brain, might play a crucial role in this interaction. While some studies have

demonstrated a clear influence of sleep loss on embryo-derived microglia [20-22], the opposite

relationship and how the brain-resident macrophages modulate sleep/wake cycle and

vigilance states remain largely unexplored. Therefore, it is essential to thoroughly investigate

the extent to which brain-resident macrophages influence sleep states and brain activity during

these states. EEG-recorded brain activity will therefore be an important indicator of the effects

of the immune system on brain oscillations. Additionally, the influence of brain macrophages

on sleep-regulating neurotransmitters, and synaptic connectivity warrants further exploration.

Finally, it remains unclear whether adulthood-derived brain-resident macrophages are

capable of re-establishing neural circuitries in the same manner as prenatal cells, including

synaptic plasticity and neurochemical balance, which are crucial for normal sleep regulation.

Here, we undertook to address these questions through a sound experimental design and in-

depth analyses. Using a comprehensive characterization of EEG signals recorded from the

mice brain, we first explored whether the absence of embryo-derived brain-resident

macrophages influences sleep-related brain oscillatory patterns. Next, we studied whether

repopulated postnatal macrophages can normalize these patterns. Finally, we investigated

how the absence and re-presence of microglia and CAMs affects brain network connectivity

and neurochemistry. Our data show functional differences between embryo-derived and

postnatally adulthood-derived brain-resident macrophages.

Results

Repopulated postnatally-derived brain-resident macrophages are deficient in rebuilding normal NREMS circuitry

We examined to what extent embryo-derived brain-resident macrophages contribute to sleep regulation and whether postnatal-derived macrophages mirror the same functionality. To address this, we depleted microglia from the brains of mice for two weeks using colony-stimulating factor 1 receptor (CSF1R) inhibition via PLX5622 [23], and then allowed them to repopulate for four weeks (Fig. 1A). CSF1 receptor is vital for microglia survival. Antagonizing it severely depletes microglia within three days, and upon the termination of treatment, repopulation can be readily observed within a few days[17]. We observed substantial microglia depletion (>86%) after two weeks of PLX5622 treatment, and complete repopulation four weeks after withdrawal (Fig. 1B, C).

We recorded EEG signals before depletion (control), after depletion and after repopulation (Fig. 1A). Since CSF1 inhibition also depletes CAMs [24], the effects we report here are due to the depletion of both cell types, which we refer to collectively as MG. Our experimental conditions therefore included control, MG-Dep (microglia/CAM depleted), and MG-Repop (microglia/CAM repopulation). To evaluate the distribution of vigilance states, we employed our standard sleep/wake phenotyping protocol [25], with two days of baseline (which were averaged and presented as one day in the figures) and in the third day mice underwent 6 h of SD (ZT0-ZT6) and 18 h of recovery (Fig. 1D). In baseline conditions, all experimental mice demonstrated the characteristic cycling behavior, with higher levels of wakefulness during the dark period and higher levels of NREMS during the light period and exhibited the increase in NREMS amounts in response to SD (Fig. 1E, F). However, a significant NREMS increase was observed in the baseline dark period when comparing the MG-Dep to the control condition. Interestingly, the repopulation of MG did not normalize this effect (Control: 12.75±0.72 min, MG-Dep: 16.78±0.77 min, MG-Repop: 16.54±0.70 min, mean ± SEM) (Fig. 1E, F, bottom panels). Reciprocally, there was a notable reduction in wakefulness during the dark period (Fig. 1E, F, top panels).

To gain insight into sleep homeostatic regulation in different mouse groups and determine

whether increased sleep is due to increased need for sleep, we analyzed the time course of

NREMS EEG delta power (1-4 Hz, a recognized quantitative readout of sleep need) across

our 3-day recordings. All mice groups showed a typical decline in delta power during the light

period and a strong rebound after SD, indicating homeostatic regulation of sleep need in all

groups (Fig. 1G). However, the buildup in delta power during the baseline dark period was

significantly lower in MG-Dep and MG-Repop mice compared to WT mice (Fig. 1G). This

attenuation was more pronounced in MG-Repop mice and may be partly related to the

increased amount of sleep during the baseline dark period (Fig. 1E, lower panel). In contrast,

when mice were exposed to enforced wakefulness (SD), the EEG delta power increase was

significantly more pronounced in MG-Dep and MG-Repop than in control conditions, and again

this effect was more extensive in MG-Repop than in MG-Dep (Fig. 1G). Altogether, these data

indicate impaired buildup of homeostatic sleep need during the waking period.

Spectral analysis of NREMS revealed that MG-Dep mice exhibited higher slow-delta (1-2 Hz)

activity compared to control mice in baseline light period (Fig. 1H, left). This frequency band

appears particularly affected also in MG-Repop mice, as it was significantly increased in the

recovery light period following SD (Fig. 1I, left). The fast-delta (3-4 Hz) frequency range was

in contrast blunted during both baseline and recovery light periods only in the MG-Dep

condition (Fig. 1H, I, left). During the baseline dark period, large delta power blunting was

observed in both MG-Repop (1.5-3.5 Hz) and MG-Dep (3-5.5 Hz) conditions. These deficits

remained nearly the same during the recovery dark period, indicating that NREMS of both

experimental mice appears to be shallower (Fig. 1H, I, right). Collectively, it can be concluded

that the presence of MG is necessary for the precise regulation of fast and slow-delta power

as a proxy of the sleep homeostatic process. MG repopulation not only fails to restore NREMS

regulation but worsens it.

To determine whether increased NREMS duration is due to enhanced state initiation

(increased number of NREMS episodes) and/or to enhanced state maintenance (increased

NREMS episode duration), we conducted a sleep fragmentation analysis. We found that while

MG-Dep mice have similar sleep bout duration patterns as the control condition, MG-Repop

mice showed a larger number of shorter NREMS bouts (16 seconds [Control: 6.7±0.63, MG-

Dep: 6.8±0.7, MG-Repop: 9.84±1.12]) and 64 seconds [Control: 12.12±1.15, MG-Dep:

11.02±0.81, MG-Repop: 17.63±1.2]) and fewer longer NREMS bouts (>256 seconds [Control:

1.96±0.28, MG-Dep: 1.74±0.33, MG-Repop: 0.62±0.17]), suggesting a strong destabilization

of NREMS in the MG-Repop brain (Fig. 1J).

During NREMS, brief bursts of brain activity in the 9-16 Hz frequency range, called spindles,

are generated by thalamo-cortical interactions and play roles in processes such as learning,

memory, and cognition [26]. To determine if MG depletion and repopulation influence spindle

generation, we quantified spindle rate during NREMS episodes and found a significantly

reduced spindle rate in MG-Dep mice compared to the control condition (Fig. 1K, L; see

"Methods"). In contrast to other features of NREMS, MG repopulation was found to normalize

the occurrence rate of spindle events, indicating that MG repopulation partially rescues the

circuit deficits seen after MG depletion.

Altogether, these data suggest that embryo-derived MG are an integral part of NREMS

establishment and maintenance, and repopulated postnatal MG largely fail to fulfill the normal

NREMS regulatory functions of embryo-derived MG.

Depletion of brain-resident macrophages prolongs REMS and degrades its quality

Different neural pathways and neurotransmitters regulate REMS compared to NREMS.

However, the role of MG in the regulation of REMS is not well understood. We quantified

REMS amount across the dark/light periods and found that MG-Dep mice display more REMS

compared to the control and MG-Repop conditions during the baseline dark period (Control:

0.67±0.1, MG-Dep: 1.17±0.21 min/h, mean ± SEM). However, in contrast to NREMS, MG

repopulation corrected the abnormal expression of REMS (MG-Repop: 0.61±0.13 min/h) (Fig.

2A, B). To investigate if the increase in REMS is due to the consolidation of this state, we

performed fragmentation analysis and found that the number of shorter REMS bouts (8 s) is

largely decreased (Control: 8.7±2.05, MG-Dep: 2.8±1.2, MG-Repop: 8.8±3.3) and longer

bouts (>2 min) are increased (Control: 1.5±0.8, MG-Dep: 6.1±1.1, MG-Repop: 4.71±1.4, mean

± SEM) (Fig. 2C) in MG-Dep mice, suggesting stabilization of REMS in the absence of MG.

Theta oscillations dominate the EEG signal during REMS, where theta phase dynamically

modulates gamma amplitude in hippocampal and cortical networks [27]. We next investigated

the contribution of MG in theta oscillations and theta-gamma interactions. Spectral analysis of

REMS revealed a significant dampening of theta (6-8 Hz) power in MG-Dep mice during the

baseline dark period (Fig. 2D), which was restored after MG repopulation. We also found that

the theta-gamma coupling is significantly reduced in the MG-Dep condition compared to the

control. Although MG repopulation attempts to bring it back to the normal level, it fails

significantly (Fig. 2E, F).

Altogether, these findings suggest that the presence of MG is necessary for the normal

expression of REMS.

Brain-resident macrophages contribute to the maintenance of active wakefulness

The quality of prior wakefulness contributes to the expression of both NREMS and REMS.

Along these lines, a specific sub-state of prior wakefulness was recently proposed to drive

NREMS need [28], and some aspects of wakefulness may functionally substitute for REMS [29].

To investigate how wakefulness is expressed in our experimental conditions, we performed a

dynamic analysis of the full waking EEG spectrum across three recording days. This global

analysis revealed differential expression of the higher waking frequency bands during light-

dark cycles across three days, and most notably during enforced wakefulness (SD). As

suggested in Figure 3 (Fig. 3A), both MG-Dep and MG-Repop mice exhibit a large dampening

of the lower gamma (≈45-55 Hz) band during the baseline dark period, which is exacerbated

during the recovery dark period. Additionally, the MG-Dep mice exhibit a large blunting of the

higher gamma band (≈70-90 Hz) during SD, which partially normalizes after MG repopulation.

To better quantify the lower frequency bands, we performed PSD analysis and found that MG-

Dep mice exhibit a large increase in slow delta (1.35-1.75 Hz) activity compared to the control

mice (Fig. 3B).

Mice spend a portion of their waking time engaged in exploratory and motivated behaviors,

such as running, nest building, or drinking, often referred to in rodents as "active wakefulness".

These behaviors are associated with the EEG signature called "theta-dominated wakefulness"

(TDW 6.0-9.5 Hz), which has been suggested to be the principal driver of the sleep homeostat

[28]. We quantified TDW in our mice and found a large decrease in its amount in both MG-Dep

and MG-Repop mice during the baseline dark period (Fig. 3C, D). This reduction was more

pronounced in MG-Repop mice, which was also seen during SD (Fig. 3C, E). TDW is

associated with heightened gamma activity. We analyzed the time course of gamma activity

during wakefulness and found an increase in slow (32-45Hz) frequency bands in the MG-

Repop condition compared to the controls (Fig. 3F). The increase in these frequency bands

in MG-Repop mice is also apparent on time-frequency heat maps in Figure 3A.

Overall, these data suggest a significant imbalance in the theta/gamma-rich waking state in

both the MG-Dep and MG-Repop conditions, and appearance of heightened gamma activity

after MG repopulation.

Disrupted balance of synaptic signaling upon MG depletion and repopulation

Sleep is associated with changes in synaptic plasticity [30] and microglia play a substantial role

in synaptic functions, which are dependent on neuronal activity and vigilance states (VS)[31]. To

investigate whether MG depletion and repopulation induce changes in brain network

connectivity at the synaptic level, we analyzed a subset of excitatory and inhibitory synaptic

density proteins that are involved in sleep regulation, (e.g., the core brain molecular correlate

of sleep loss, HOMER1) in the brains of our experimental mice.

Using an ImageJ macro, we counted the number of synapses on cortical and hypothalamic

neurons (Fig. 4A, B) and found that the number of excitatory VGLUT1-HOMER1 synapses on

NEUN$^+$ cortical neurons was significantly reduced in both MG-Dep and MG-Repop conditions.

This reduction was more noticeable in MG-Repop mice (Fig. 4C). Interestingly, this effect was

regulated at the postsynaptic level (HOMER1-associated, Fig. 4D) rather than the presynaptic

level (VGLUT1-associated, Fig. 4E). Recently, cortical PV neurons have been proposed to be

responsible for homeostatic sleep regulation[32]. We found that the number of parvalbumin (PV)-

originated[33] Synaptotagmin-2-Gephyrin (Syt2-Geph) inhibitory synapses on the cortical

NEUN[+] neurons was significantly increased under the MG-Repop condition (Fig. 4F, G). This

increase could possibly be due to an increase in the activity of PV neurons, as we found that

the number of VGLUT1-HOMER1 excitatory synapses on PV neurons was significantly

increased under both MG-Dep and MG-Repop conditions (Fig. 4H, I). We additionally

investigated the changes in the count of VGLUT1-HOMER1 synapses on representative

hypothalamic neurons (HCRT) and found a significant decrease in the number of VGLUT1-

HOMER1 synapses on HCRT neurons in both MG-Dep and MG-Repop conditions (Fig. 4J,

K).

To further investigate changes in network connectivity in the brains of MG-Dep and MG-Repop

mice, we analyzed additional synaptic markers and sleep-specific neurotransmitter receptors

(e.g., receptors of monoaminergic systems and neuropeptides) on RNA extracted from cortical

samples of these mice. Our bulk analysis revealed that some general synaptic markers, such

as *Synaptophysin I* (*Syp*), are largely downregulated in both experimental mice, while *Slc17a7*

(encoding Vesicular glutamate transporter 1) and *Homer1* genes are significantly

downregulated in the cortex of the MG-Repop condition. Among the specific neurotransmitter

markers analyzed, the adrenergic receptor family (e.g. *Adrb2*, *Adra2a*, *Adrb3*, and *Adra1a*)

genes showed a tendency to be less expressed in both MG-Dep and MG-Repop conditions,

with a marked decrease in the expression of the *Adra2a* gene. Additionally, we observed a

reduction in expression of *Cholinergic receptor muscarinic 2* (*Chrm2*) and the serotonergic

receptor *5-Hydroxytryptamine receptor 1A* (*Htr1a*) genes, which mainly occur in the brain of

MG-Repop mice (Fig. 4L).

Taken together, these data suggest that brain connectivity at the synaptic level is largely

impaired following the depletion of brain-resident macrophages. Repopulating the brain with

these macrophages not only failed to normalize connectivity but, in many cases analyzed,

worsened it.

Neurotransmitter release is largely restored after the repopulation of resident macrophages

Synapses, in addition to providing structural connections in the brain, play a crucial role in transmitting neurotransmitters essential for the function of complex neural systems. We therefore questioned whether the levels of essential neurotransmitters involved in sleep regulation are affected by the depletion and repopulation of brain-resident macrophages. We conducted targeted metabolomics to analyze the neurotransmitter content and metabolite reservoir of the cortex, target of most neurotransmitters and region where EEG signals are acquired. Assessment of neurotransmitter content was performed using chromatography with tandem mass spectrometry (LC-MS/MS) in the brain of control, as well as MG-Dep and MG-Repop mice. We found that pharmacological depletion of MG led to a distinct clustering of metabolites, as indicated by principal component analysis (PCA) (Fig. 5A). This suggests that MG plays an important role in maintaining the overall metabolic composition of the cortex. MG repopulation however restored the cortical submetabolome to a pattern similar to the one observed under control experimental conditions (Fig. 5A). Univariate analysis revealed a reduction in norepinephrine upon MG depletion, which was restored after repopulation (Fig. 5B). The concentrations of acetylcholine, choline, dopamine, and serotonin remained unchanged between the experimental conditions, with a slight tendency to decrease in MG-Dep condition.

Furthermore, we observed that MG depletion led to a marked reduction of L-2-amino adipic acid (L-2-AAA), which was only partially recovered after repopulation (Fig. 5B). We also analyzed the precursors of neurotransmitters and found a significant decrease in the amino acids L-serine and asparagine under depleted condition, which could be restored upon MG repopulation (Fig. 5C). Similarly, there was a trend towards decreased glycine and L-glutamic acid content (Fig. 5C).

Taken together, our data suggest that while network connectivity is significantly impaired in MG-Dep condition and resistant to MG repopulation, the underlying substrates (neurotransmitters and metabolites) are largely restored after MG repopulation.

Discussion

In this study, we investigated the extent to which embryo-derived brain-resident macrophages contribute to sleep regulation and whether postnatally derived ones serve a similar role. We demonstrate that MG depletion has a significant impact on the homeostatic regulation of sleep. The brain circuits most vulnerable to microglia depletion and repopulation, causing the changes in sleep EEG that we document, are still unknown. However, changes in slow-delta activity in NREM sleep, state stabilization and reduced theta power in REMS, as well as changes in waking slow-delta activity, were reported in mice with defects in hypothalamic HCRT and brainstem serotonergic and noradrenergic systems[25,28,34,35]. This is suggestive of functionally important local interactions between microglial populations and sleep regulating circuits, as mentioned above.

We showed that the changes in sleep patterns after MG depletion are not fully reversed after MG repopulation. Previous studies have shown that the brain global transcriptomes are little influenced by repopulated microglia [17]. Furthermore, it has been suggested that after MG repopulation, neuroprotection is achieved through IL-6 induction by neurons[36]. Other studies reported full functionality of retinal microglia after repopulation, including continuous retinal surveillance, maintenance of synaptic structure, and normal behavioral and physiological responses to retinal injury [37]. However, our data reveal that MG repopulation not only fails to completely restore various aspects of sleep regulation, but in some cases exacerbates altered parameters, such as fragmentation of NREMS and dysregulation of sleep need. This demonstrates striking funtional differences between microgial cells derived from pre-or postantal sources. We speculate that not all microglia-neuronal interactions are properly reestablished by adulthood-derived microglia after repopulation. Whether a longer repopulation period might correct the abnormalities observed in our analysis warrants further investigation. We observed a significant imbalance between excitatory and inhibitory synapses after depletion of embryo-derived microglia and after repopulation with adult

microglia, with a notable increase in inhibitory synapses following repopulation. Studies in

juvenile mice prevously reported that microglia depletion in the developing brain significantly

increases the number of both excitatory (VGLUT1-HOMER1) and inhibitory (Syt2-Gephyrin)

synapses on cortical neurons, an effect that persisted after repopulation until P30. However,

in the adult (P60) brain, synapse density returned to normal control levels, suggesting that

depleting microglia during development causes long-lasting, albeit not permanent, effects[38].

Our data, however, show that depleting microglia in adulthood, and then repopulating them,

induces different results, with a reduction in excitatory synapses in both MG-Dep and MG-

Repop conditions, and an increase in inhibitory synapses in MG-Repop conditions. In the

developing brain, it has been suggested that MG-Dep impairs synaptic pruning processes and

alters related gene expression[10,38]. Whether the same process or any specific pathways are

induced by MG depletion and repopulation in the adult brain, such as aberrant reactivation of

developmental programs[39], remains to be found.

Furthermore,other studies have suggested that microglia play a role in restraining excessive

neuronal activation that cannot be sufficiently suppressed by inhibitory neurons alone[40]. Our

results suggest that embryo-derived microglia also control excessive activity of inhibitory PV

neurons, but the reasons why this is not corrected with adulthood-derived microglia needs

further investigation.

Deficient synaptic neurotransmission was proposed as a potential cause for abnormal sleep

regulation[41]. We found here an imbalanced neurotransmitter and metabolite content of the

brain in mice devoid of prenatal microglia. It appears that norepinephrine neurotransmission

is particularly sensitive to MG deficiency, as both the levels of norepinephrine metabolites and

the corresponding receptors are significantly affected by MG depletion. Norepinephrine is

implicated in both wakefulness[42] and sleep regulation[43-45], and it has been suggested that

microglial surveillance process relies heavily on norepinephrine signaling[46,47]. Additionally, we

observed a decrease in the expression of neurotransmitter precursors such as L-serine, which

was mirrored by changes in the concentrations of glycine, asparagine, and glutamic acid.

Previous studies have proposed serine as a link between metabolism and neurotransmission

[48], and serine serves as a precursor for synthesizing ceramide [49], which has been suggested

to be essential for maintaining stable wakefulness [50].

Our findings indicate that MG affect vigilane state regulation via multiple pathways and play

a more substantial role in maintaining the physical connectivity of brain networks rather than

neurochemical connectivity, as MG repopulation largely restores metabolites but not

synapses. To gain a detailed understanding of MG regulation of sleep, it will be necessary to

explore local specific microglial depletion, and disruption of microglia-neuron interactions

using inducible systems. Additionally, investigating the separate contributions of microglia and

CAMs in sleep regulation using Cre lines under the promoter of *Hexb* [51] and *Mrc1* [52] genes,

respectively, would be a starting point. The observed vigilance state phenotypic alterations

may also be influenced by disruptions in central factors (such as protein phosphorylation

orchestrated by microglial TNF-α)[53], and peripheral players (since CSF1R inhibition also

effects circulating and tissue macrophages)[54] warranting further investigations. Overall, our

results shed new light on the complex interaction between the neuroimmune system and sleep

regulation, and it paves the way for future investigations in this important physiopathological

field.

Data availability

All data are available from the corresponding author upon reasonable request.

Acknowledgements

We extend our gratitude to the Metabolomics Core Facility at the Department of General

Pediatrics, University of Freiburg, with special thanks to Dr. Luciana Hannibal and Victoria

Wingert for their invaluable support and expertise. AS was supported by Novartis Foundation

for medical-biological Research (#23B079). MB was supported by the Swiss National Science

Foundation (grant 190605). MT was supported by Swiss NSF (grant 201235), and the State

of Vaud (Faculty of Biology and Medicine, University of Lausanne). AV was supported by the

Swiss National Science Foundation (grant 31003A_182613). AK was supported by institute of

neuroinformatics ETH-Zurich. GM was supported by H2020-MSCA-IF grant, n. 101025176.

Author contributions

AS conceptualized and designed the study. AS, MH, and SW performed experiments. AS,

GM, and MB analyzed data. AS, GM, AK, AV, MT, and MB wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

Methods

Animals. Wild type (C57BL/6J) mice were used in this study. All animals were between 12 and 14

weeks old during the time of experiments and animal experiments were approved by the Ministry for

Nature, Environment and Consumers` Protection of the state of Baden-Württemberg and were

performed in accordance to the respective national, federal, and institutional regulations. At all times,

care was taken to minimize animal discomfort and avoid pain.

Gene expression analysis. For qPCR analysis, RNA extracted from the cortical samples of control,

MG-Dep and MG-Repop mice. RNA was extracted using RNeasy micro kit (74004) and cDNA synthesis

was performed using random hexamer primers and M-MLV Reverse Transcriptase Promega kit

(M3682). SYBR green-based and Taqman probes were used to detect the gene expression. The assay

was performed using a LightCycler480 (Roche).

Immunofluorescence and synapse counting. Mice were anesthetized with ketamine (100 mg per kg

body weight) and xylazine (10 mg per kg body weight) followed by perfusion with 1× PBS. For histology,

brains were kept overnight in 4% paraformaldehyde (PFA). Brains were then dehydrated in 30%

sucrose for about 48 hours and then snap-frozen on dry ice. Cryosections (14µm for synapse counting

and 20µm for microglia and CAM staining) from brain tissue were prepared as free floating in PBS. The

following antibodies were used to stain microgliaand synapses: Rabbit-anti-IBA1 (ab178846, abcam),

anti-NEUN (ab177487, ab104224, abcam), anti-VLUT1 (135304, synaptic system), anti-HOMER1

(ABN37, Millipore), anti-Gephyrin (147318, synaptic system), anti-Synaptotagmin-2 (ZDB-ATB-081002-

25, ZFIN), anti-Parvalbumin (MAB1572, Merck) and anti-HCRT-A (SC-8070). The DAPI was used to

stain the nucleus. Iba1 antibody was incubated overnight at 4 °C and the primary antibodies to detect

synapses were incubated for 1.45hr at room temperature. The brain cuts were incubated with secondary

antibodies for 1-2 hours at room temperature. Fluorescence imaging was performed with TCS SP8 X

(Leica Microsystems). Imaging of synapses was performed using 63X objective with zoom factor 8.

Iba1[+] cells were counted using Fiji. Synapse analysis and quantification were conducted using Fiji

macro developed previously [55] with little modifications.

Quantitative profiling of metabolites by liquid chromatography and mass spectrometry (LC-

MS/MS). Mice brain tissue biopsies were thawed and homogenised with ice-cold PBS supplemented

with 1% protease inhibitor cocktail (Sigma Nr. P8340-5ML), using a cordless pestle motor and

disposable pellet mixers (VWR Nr. 47747-366). Buffer volume was adjusted to reach a target of 0.1 to

0.3 mg wet-tissue/mL lysis buffer. Whole tissue homogenates were aliquoted for downstream

measurement of total protein concentration, flash-frozen and stored at -80 $^\circ$C. Sulfur-containing

metabolites as well as creatinine, S-adenosylmethionine and S-adenosylhomocysteine were

determined according to a previously published procedure [56,57]. Lactate, TCA and glycolysis

intermediates and other organic acids, and folates were determined as described in previous work [58].

Amino acids and neurotransmitters were determined using a previously described protocol [59,60]. A

commercially available standardized amino acid mixture was utilized to generate a calibration curve for

amino acids (Amino acid standards, physiological, Sigma, Nr. A9906-10ML). Calibration curves for all

other metabolites were prepared from individual stock solutions prepared in house. Quantitation

accuracy was examined by monitoring homocysteine and methylmalonic acid concentrations in an

external quality control, namely, the Control Special Assays in Serum, European Research Network for

the evaluation and improvement of screening, diagnosis, and treatment of Inherited disorders of

Metabolism (ERNDIM) IQCS, SAS-02.1 and SAS-02.2 from MCA Laboratories, Winterswijk,

Netherlands. For all other metabolites, quantitation trueness was tested by examining metabolite

concentrations in plasma from a previously validated sample isolated from a healthy control individual

with respect to standard reference ranges, using the same calibration curves and LC-MS/MS running

conditions. Quantification of metabolites was carried out with Analyst® 1.7.2 software, 2022 AB Sciex.

Treatments. In order to eliminate cells dependent on CSF1R (MG) mice were administered a CSF1R

inhibitor called PLX5622 (Plexxikon). The inhibitor was combined with AIN-76A standard chow at a

concentration of 1,200ppm (Research Diets) and the mice were given unrestricted access to it for two

weeks. The control group received AIN-76A standard chow without any modifications. After two weeks

the PLX5622 food was replaced with control food to start repopulation of MG.

Mice surgery, data acquisition and vigilance state analysis. Mice surgery and data acquisition were

performed as previously described [25]. MATLAB scripts were created to quantify various parameters

related to wakefulness, NREMS and REMS. These parameters include time distribution and amount of

episodes, as well as the fragmentation of vigilance states, all of which were defined based on previously

established criteria [25]. TDW analysis was conducted following the method described by Vassalli and

Franken [28].

Power spectral analysis. Using Somnologica-3TM (Medcare) software, each 4 s epoch of the EEG
signal was subjected to discrete Fourier transform to determine EEG power density (spectra 0–90 Hz
at a frequency resolution of 0.25 Hz). We used artifact-free, same-state–flanked 4 s epochs to calculate
the mean EEG spectral profiles for each behavioral states and time intervals. To account for differences
among animals in absolute EEG power, the mean values of spectral profiles were converted into
percentages relative to a baseline EEG power reference value of 100%. This reference value was
determined for each mouse by summing the power within the frequency range of 0.75–47.5 Hz across
all states and throughout the first day of baseline recordings. To control for differences in the amount
of wakefulness, NREMS and REMS, this reference was weighted so that the relative contribution of
each state was identical for all mice [61]. Time-frequency heatmaps of EEG power during wakefulness
were generated using methods described previously [62].
465
Time-course analysis. To examine the dynamic changes of EEG power within specific frequency
ranges throughout the day and night, the time-course of the activity of that frequency band was
computed for 4 s epochs scored as state of interest (NREMS, REMS and wake). For this purpose, the
epochs for each state were divided into percentiles, ensuring that each percentile contained a roughly
equal number of epochs. The number of percentiles for NREMS and REMS were: 12 for light periods,
6 for dark periods, and 8 for the 6-h light period after SD; for waking state: 6 for light periods, 12 for
dark periods, 8 for the 6-h light SD, and 4 for the 6-h light after SD. The average EEG power within the
specified frequency range was computed for each percentile, and subsequently normalized based on
the type of analysis. This normalization was either done relative to the total power of the baseline state
or with respect to the power achieved within the delta frequency band during slow-wave sleep at ZT8-
12. Like power density analysis, single epochs were excluded and only power values of the epochs that
themselves, as well as the two adjacent ones, were scored as artifact-free same-state were included in
the analysis.
479
Theta-gamma cross-frequency coupling. We used the modulation index (MI) to measure the theta-
gamma phase-amplitude coupling [27,63]. Using finite impulse response filters with an order equal to three
cycles of the low cutoff frequency, we first bandpass-filtered EEG signals into theta (6-11 Hz) and fast-
gamma (54-90 Hz) in both forward and reverse directions to eliminate phase distortion. We then
estimated instantaneous phase of theta and the envelope of fast-gamma using the Hilbert transform.
Theta phase was discretized into 18 equal bins (*N*=18, each 20°) and the average value of fast-gamma
envelope within each bin was calculated. The resulting phase-amplitude histogram (*P*) was compared
with a uniform distribution (*U*) using the Kullback-Leibler distance, $D_{KL}(P,U) = \sum_{j=1}^{N} P(j) *$
$log[P(j)/U(j)]$, and normalized by *log(N)* to obtain the modulation index, *MI = D_{KL} / log(N)*. To explore
possible coupling patterns between different pairs of low and high frequency bands, we used the
comodulogram analysis [63]. We considered 16 frequency bands for phase (1-18 Hz, 1-Hz increments,
2-Hz bandwidth), and 14 frequency bands for amplitude (15-90 Hz, 5-Hz increments, 10-Hz bandwidth).
MI values were then calculated for all these pairs to obtain the comodulogram graph.
493

Detection of sleep spindles. NREMS spindles were detected automatically using an optimized wavelet-based method as previously described [26]. Briefly, the power of EEG signals within 9-16Hz was estimated using the complex B-spline wavelet function, and smoothed using a 200 ms Hanning window, and then a threshold equal to 3 SD (SD: standard deviation) above the mean was applied to detect the potential spindle events. Events shorter than 400 ms or longer than 2 s were discarded. Using band pass-filtered EEG signals in the spindle range (9–16 Hz), we automatically counted the number of cycles of each detected event and excluded those with <5 cycles or more than 30 cycles. To discard artefacts, events with a power in the spindle band lower than 6–8.5 Hz or 16.5–20 Hz power bands were not included.

Statistical analysis. The specific number (n) of replicates employed in each experiment is provided in the corresponding figure legends. The individuals responsible for conducting SD and scoring the sleep recordings were unaware of the genotype of the animals, ensuring blinding during these procedures. The results are presented as mean ± SEM or mean ± SD. Statistical analyses were conducted using GraphPad Prism. The significance of comparisons was determined using one- or two-way ANOVA or mix effect model followed by relevant post hoc tests and are reported in the figure legends.

Figure legends

Figure 1. Essential role of MG in regulation of NREMS and associated EEG oscillations. (**A**) Schematic illustration of the experimental procedure conducted on adult mice to assess the brain's electrical activity using EEG. Recordings prior to PLX5622 drug administration served as controls. Mice were then fed PLX5622-containing food during 2 weeks. During the final 3 days of this 2-week period, a 2nd series of recordings was conducted. Following PLX5622 treatment, mice were provided control food for 4 weeks. After 4 weeks under normal food, a 3rd recording was performed. All 3 recordings were performed on the same mice. (**B, C**) PLX5622 substantially depleted the microglia from the brain within 2 weeks (compare control and MG-Dep). Return to normal food repopulated microglia into the brain after 4 weeks (compare MG-Dep and MG-Repop). (**D**) Recording paradigm. The first 2 days of recordings served as baseline, averaged as one day. The 2nd day consisted of 6 h SD followed by 18 h of recovery. (**E**) Time-course of vigilance states; Wakefulness (top) and NREMS (bottom). Compared to controls (black datapoints), MG-Dep (orange datapoints) and –Repop (blue datapoints) mice show a large increase in NREMS duration during the dark period both at baseline and during recovery (n = 9 for control and MG-Depleted conditions and n = 6 for MG-Repopulated condition, 2-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars, significant differences between control and MG-Dep and MG-

Repop mice, respectively). Data points are shown in minutes per hour (min/h) and represent
the average of 2 h. (**F**) From left to right: Time in wakefulness (top) and NREMS (bottom)
during baseline light, dark and recovery light and dark periods (Mixed-effects analysis,
followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001). (**G**) Time-course of EEG delta
power (1–4 Hz) during the 3-day recording. EEG delta power is a proxy of sleep need. Both
MG-Dep and MG-Repop mice show deficiency in building up sleep need during the dark period
(2-way ANOVA, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and
blue stars, significant differences between control and MG-Dep and MG-Repop mice,
respectively). (**H**) EEG power spectra of experimental mice in NREMS during baseline light
(left) and dark (right) periods. EEG power is expressed as % of baseline power of control
condition. Insets show amplification of specific EEG frequencies (2-way ANOVA, followed by
Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars indicate significant
differences between control and MG-Depleted and MG-Repopulated mice, respectively). (**I**)
EEG power spectra of experimental mice in NREMS during recovery light (left) and dark (right)
periods. (**J**) Distribution of NREMS bout durations during baseline dark period across the
experimental mice. Sleep bout duration is represented on the x-axis (1-way ANOVA, followed
by Tukey test). (**K**) (Top) Representative time-frequency heatmap of spindles. (Bottom)
Representative EEG/EMG signals of a detected spindle. Dashed horizontal lines indicate
upper and lower thresholds used to detect spindles with wavelet energy within 9-16 Hz, and
dashed vertical lines indicate the start and end of the detected spindle. (**L**) Spindle rate during
NREMS episodes in different conditions. Spindle rate significantly decreased in MG-depleted
mice (mixed-effects analysis, followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001)).
554
Figure 2. REMS regulation largely depends on the presence of the MG. (**A**) Time-course
of REMS duration in control (black), MG-Dep (orange) and MG-Repop (blue) mouse groups.
MG-Dep mice exhibit a large increase in REMS time during the baseline dark period (n = 9 for
control and MG-Dep conditions and n = 6 for MG-Repop condition, 2-way ANOVA, followed
by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars represent
significant differences between control and MG-Dep and MG-Repop mice, respectively). Data
points are shown in minutes per hour (min/h) and represent the average of 2 h. (**B**) Time in
REMS during baseline dark period (mixed-effects analysis, followed by Dunnett's test, *P <
0.05; **P < 0.01; ***P < 0.001). (**C**) Distribution of REMs bout durations during baseline dark
period across 3 mouse groups. Sleep bout duration is presented as x-axis (1-way ANOVA,
followed by Tukey test). (**D**) EEG power spectra of all experimental mice in REMS during
baseline dark period. Insets show amplification for specific EEG frequencies (2-way ANOVA,
followed by Dunnett's test, *P < 0.05; **P < 0.01; ***P < 0.001, orange and blue stars are
significant differences between control mice and MG-Dep and MG-Repop mice respectively).

(**E**) Comodulogram graphs show the modulation index for a wide range of frequency pairs,

obtained from 12-h recordings of one animal in different conditions. (**F**) Normalized theta-

gamma coupling across 3 experimental conditions. Theta-gamma coupling is significantly

decreased in MG-depleted and -repopulated conditions (mixed-effects analysis, followed by

Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$)).

Figure 3. MG Repopulation fails to normalize the waking state deficits induced by MG

depletion. (**A**) Time x frequency x power heatmap representations of the waking EEG of

control, MG-Dep and MG-Repop mice across the 3 recorded days (48 h baseline, 6 h sleep

deprivation and 18 h recovery) show large changes in higher frequency bands. (**B**) EEG power

spectra of experimental mice in wakefulness during baseline light and dark periods (2-way

ANOVA, followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, orange star are

significant differences between control mice and MG-Dep mice). (**C**) Time-course of TDW

amounts indicates decreased TDW time during the dark period in MG-Dep and MG-Rep.

Datapoints are shown in min/h and represent the average of 2 h (baseline days 1 and 2 are

averaged), (2-way ANOVA followed by Dunnett's test, orange and blue stars are significant

differences between control and MG-Dep and MG-Repop mice, respectively). (**D**)

Quantification of time in TDW (min/h) during baseline light and dark period (mixed-effects

analysis, followed by Dunnett's test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$). (**E**) Time in TDW

during SD. (**F**) Waking EEG slow-gamma (32-45 Hz) power dynamics during the 3-day

recording (baseline days 1 and 2 are averaged) (2-way ANOVA, followed by Dunnett's test).

Figure 4. Large synaptic imbalance after MG depletion and repopulation. (**A**) Schematic

illustration of the cell types for which the number of synapses was counted. NEUN (blue) and

PV (orange) positive neurons in the cortex and HCRT neurons in the hypothalamus. The

number of co-localized excitatory VGLUT1-HOMER1 synaptic proteins on NEUN, PV and

HCRT positive neurons were counted. Additionally, the number of co-localized inhibitory

SYT2-Gephyrin synaptic proteins on NEUN neurons were counted. (**B**) Representative image

depicting pre, postsynaptic proteins and their colocalization in a NEUN positive neuron.

Synapses (arrowheads) are marked with pink color using ImageJ macro (right). Scale bar:

10μm. (**C**) Number of VGLUT1-HOMER1 synapses/μm^2 of NEUN$^+$ soma. N=342 cells from

12 mice for control, 324 cells from 11 mice for MG-Dep, and 357 cells from 12 mice for MG-

Repop conditions. The data show a reduced number of these synapses in both MG-Dep and

MG-Repop conditions (**D**) Number of HOMER1 puncta/μm^2 of NEUN$^+$ soma. (**E**) Number of

VGLUT1 puncta/μm^2 of NEUN+ soma. (**F**) Number of SYT2-Gephyrin synapses/μm^2 of

NEUN$^+$ soma. N=299 cells from 12 mice for control, 270 cells from 11 mice for MG-Dep, and

306 cells from 12 mice for MG-Repop conditions. (**G**) Representative images depicting SYT2-

Gephyrin synapses in NEUN$^+$ cortical neuron. Large increase in these inhibitory synapses is
observed in the MG-Repop condition. Scale bar: 10 µm. (**H**) Distribution of PV$^+$ neurons (red)
in the cortex. Scale bar: 100 µm (left), 10 µm (right). (**I**) Number of vGLUT1-HOMER1
synapses/µm^2 of PV$^+$ soma. N=29 cells from 5 mice for control, 29 cells from 5 mice for MG-
Dep, and 23 cells from 4 mice for MG-Repop conditions. (**J**) Left, distribution of HCRT$^+$
neurons (red) in the lateral hypothalamus. Right, a HCRT$^+$ neuron stained with VGLUT1
(green) and HOMER1 (Cyan). Scale bar: 100 and 10 µm. (**K**) Number of VGLUT1-HOMER1
synapses/µm^2 of HCRT+ soma. N=70 cells from 3 mice for control, 73 cells from 3 mice for
MG-Dep, and 77 cells from 4 mice for MG-Repop conditions. (**L**) Gene expression of synaptic
proteins and neurotransmitter receptors at RNA level following MG depletion and repopulation.
N=3 for control and MG-Dep and n=4 for MG-Repop conditions. 1-way ANOVA, followed by
Tukey test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

Figure 5. MG repopulation largely restores the metabolite changes induced by
microglial depletion. (**A**) PCA of targeted metabolites in cortex isolated from control (n=3),
MG depleted (n=3) and repopulated (n=4) mice. (**B**) LC-MS/MS assessment of the
neurotransmitter content in the brain of control, MG-Dep and MG-Repop mice. (**C**)
Downregulation of amino acids in the brain of experimental mice after MG depletion. 1-way
ANOVA, followed by Tukey test, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

625
626
627

References

628
1. Ramar, K., Malhotra, R.K., Carden, K.A., Martin, J.L., Abbasi-Feinberg, F., Aurora, R.N., Kapur,
V.K., Olson, E.J., Rosen, C.L., Rowley, J.A., et al. (2021). Sleep is essential to health: an
American Academy of Sleep Medicine position statement. J Clin Sleep Med.
10.5664/jcsm.9476.
2. Borbély, A.A. (1982). A two process model of sleep regulation. Hum Neurobiol *1*, 195-204.
3. Gilestro, G.F., Tononi, G., and Cirelli, C. (2009). Widespread changes in synaptic markers as a
function of sleep and wakefulness in Drosophila. Science *324*, 109-112.
10.1126/science.1166673.
4. Maret, S., Dorsaz, S., Gurcel, L., Pradervand, S., Petit, B., Pfister, C., Hagenbuchle, O., O'Hara,
B.F., Franken, P., and Tafti, M. (2007). Homer1a is a core brain molecular correlate of sleep
loss. Proc Natl Acad Sci U S A *104*, 20090-20095. 10.1073/pnas.0710131104.
5. Saper, C.B., Chou, T.C., and Scammell, T.E. (2001). The sleep switch: hypothalamic control of
sleep and wakefulness. Trends Neurosci *24*, 726-731. 10.1016/s0166-2236(00)02002-6.
6. Amann, L., Masuda, T., and Prinz, M. (2023). Mechanisms of myeloid cell entry to the
healthy and diseased central nervous system. Nat Immunol *24*, 393-407. 10.1038/s41590-
022-01415-8.
7. Kierdorf, K., Masuda, T., Jordão, M.J.C., and Prinz, M. (2019). Macrophages at CNS interfaces:
ontogeny and function in health and disease. Nat Rev Neurosci *20*, 547-562.
10.1038/s41583-019-0201-x.
8. Prinz, M., Masuda, T., Wheeler, M.A., and Quintana, F.J. (2021). Microglia and Central
Nervous System-Associated Macrophages-From Origin to Disease Modulation. Annual
review of immunology *39*, 251-277. 10.1146/annurev-immunol-093019-110159.
9. Prinz, M., Jung, S., and Priller, J. (2019). Microglia Biology: One Century of Evolving Concepts.
Cell *179*, 292-311. 10.1016/j.cell.2019.08.053.
10. Paolicelli, R.C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M.,
Ferreira, T.A., Guiducci, E., Dumas, L., et al. (2011). Synaptic pruning by microglia is
necessary for normal brain development. Science (New York, N.Y.) *333*, 1456-1458.
10.1126/science.1202529.
11. Shigemoto-Mogami, Y., Hoshikawa, K., Goldman, J.E., Sekino, Y., and Sato, K. (2014).
Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal
subventricular zone. J Neurosci *34*, 2231-2243. 10.1523/jneurosci.1619-13.2014.
12. Nimmerjahn, A., Kirchhoff, F., and Helmchen, F. (2005). Resting microglial cells are highly
dynamic surveillants of brain parenchyma in vivo. Science *308*, 1314-1318.
10.1126/science.1110647.
13. Kierdorf, K., Erny, D., Goldmann, T., Sander, V., Schulz, C., Perdiguero, E.G., Wieghofer, P.,
Heinrich, A., Riemke, P., Hölscher, C., et al. (2013). Microglia emerge from erythromyeloid
precursors via Pu.1- and Irf8-dependent pathways. Nature neuroscience *16*, 273-280.
10.1038/nn.3318.
14. Schulz, C., Gomez Perdiguero, E., Chorro, L., Szabo-Rogers, H., Cagnard, N., Kierdorf, K.,
Prinz, M., Wu, B., Jacobsen, S.E., Pollard, J.W., et al. (2012). A lineage of myeloid cells
independent of Myb and hematopoietic stem cells. Science (New York, N.Y.) *336*, 86-90.
10.1126/science.1219179.
15. Bruttger, J., Karram, K., Wörtge, S., Regen, T., Marini, F., Hoppmann, N., Klein, M., Blank, T.,
Yona, S., Wolf, Y., et al. (2015). Genetic Cell Ablation Reveals Clusters of Local Self-Renewing
Microglia in the Mammalian Central Nervous System. Immunity *43*, 92-106.
10.1016/j.immuni.2015.06.012.
16. Tay, T.L., Mai, D., Dautzenberg, J., Fernández-Klett, F., Lin, G., Sagar, Datta, M., Drougard, A.,
Stempfl, T., Ardura-Fabregat, A., et al. (2017). A new fate mapping system reveals context-
dependent random or clonal expansion of microglia. Nature neuroscience *20*, 793-803.
10.1038/nn.4547.

17.   Huang, Y., Xu, Z., Xiong, S., Sun, F., Qin, G., Hu, G., Wang, J., Zhao, L., Liang, Y.X., Wu, T., et al.
680   (2018). Repopulated microglia are solely derived from the proliferation of residual microglia
681   after acute depletion. Nat Neurosci 21, 530-540. 10.1038/s41593-018-0090-8.
682   18.   Ingiosi, A.M., Opp, M.R., and Krueger, J.M. (2013). Sleep and immune function: glial
683   contributions and consequences of aging. Curr Opin Neurobiol 23, 806-811.
684   10.1016/j.conb.2013.02.003.
685   19.   Imeri, L., and Opp, M.R. (2009). How (and why) the immune system makes us sleep. Nat Rev
686   Neurosci 10, 199-210. 10.1038/nrn2576.
687   20.   Tuan, L.H., and Lee, L.J. (2019). Microglia-mediated synaptic pruning is impaired in sleep-
688   deprived adolescent mice. Neurobiol Dis 130, 104517. 10.1016/j.nbd.2019.104517.
689   21.   Gentry, N.W., McMahon, T., Yamazaki, M., Webb, J., Arnold, T.D., Rosi, S., Ptáček, L.J., and
690   Fu, Y.H. (2022). Microglia are involved in the protection of memories formed during sleep
691   deprivation. Neurobiol Sleep Circadian Rhythms 12, 100073. 10.1016/j.nbscr.2021.100073.
692   22.   Bellesi, M., de Vivo, L., Chini, M., Gilli, F., Tononi, G., and Cirelli, C. (2017). Sleep Loss
693   Promotes Astrocytic Phagocytosis and Microglial Activation in Mouse Cerebral Cortex. J
694   Neurosci 37, 5263-5273. 10.1523/jneurosci.3981-16.2017.
695   23.   Dagher, N.N., Najafi, A.R., Kayala, K.M., Elmore, M.R., White, T.E., Medeiros, R., West, B.L.,
696   and Green, K.N. (2015). Colony-stimulating factor 1 receptor inhibition prevents microglial
697   plaque association and improves cognition in 3xTg-AD mice. J Neuroinflammation 12, 139.
698   10.1186/s12974-015-0366-9.
699   24.   Van Hove, H., Martens, L., Scheyltjens, I., De Vlaminck, K., Pombo Antunes, A.R., De Prijck, S.,
700   Vandamme, N., De Schepper, S., Van Isterdael, G., Scott, C.L., et al. (2019). A single-cell atlas
701   of mouse brain macrophages reveals unique transcriptional identities shaped by ontogeny
702   and tissue environment. Nat Neurosci 22, 1021-1035. 10.1038/s41593-019-0393-4.
703   25.   Seifinejad, A., Li, S., Possovre, M.L., Vassalli, A., and Tafti, M. (2020). Hypocretinergic
704   interactions with the serotonergic system regulate REM sleep and cataplexy. Nat Commun
705   11, 6034. 10.1038/s41467-020-19862-y.
706   26.   Bandarabadi, M., Herrera, C.G., Gent, T.C., Bassetti, C., Schindler, K., and Adamantidis, A.R.
707   (2020). A role for spindles in the onset of rapid eye movement sleep. Nat Commun 11, 5247.
708   10.1038/s41467-020-19076-2.
709   27.   Bandarabadi, M., Boyce, R., Gutierrez Herrera, C., Bassetti, C.L., Williams, S., Schindler, K.,
710   and Adamantidis, A. (2019). Dynamic modulation of theta-gamma coupling during rapid eye
711   movement sleep. Sleep 42, 1-11. 10.1093/sleep/zsz182.
712   28.   Vassalli, A., and Franken, P. (2017). Hypocretin (orexin) is critical in sustaining theta/gamma-
713   rich waking behaviors that drive sleep need. Proc Natl Acad Sci U S A 114, E5464-e5473.
714   10.1073/pnas.1700983114.
715   29.   Endo, T., Roth, C., Landolt, H.P., Werth, E., Aeschbach, D., Achermann, P., and Borbély, A.A.
716   (1998). Selective REM sleep deprivation in humans: effects on sleep and sleep EEG. Am J
717   Physiol 274, R1186-1194. 10.1152/ajpregu.1998.274.4.R1186.
718   30.   Tononi, G., and Cirelli, C. (2003). Sleep and synaptic homeostasis: a hypothesis. Brain Res
719   Bull 62, 143-150. 10.1016/j.brainresbull.2003.09.004.
720   31.   Hristovska, I., Robert, M., Combet, K., Honnorat, J., Comte, J.C., and Pascual, O. (2022). Sleep
721   decreases neuronal activity control of microglial dynamics in mice. Nat Commun 13, 6273.
722   10.1038/s41467-022-34035-9.
723   32.   Kon, K., Ode, K.L., Mano, T., Fujishima, H., Tone, D., Shimizu, C., Shiono, S., Yada, S., Garçon,
724   J.Y., Kaneko, M., et al. (2023). Cortical parvalbumin neurons are responsible for homeostatic
725   sleep rebound through CaMKII activation. bioRxiv, 2023.2004.2029.537929.
726   10.1101/2023.04.29.537929.
727   33.   Bouhours, B., Gjoni, E., Kochubey, O., and Schneggenburger, R. (2017). Synaptotagmin2
728   (Syt2) Drives Fast Release Redundantly with Syt1 at the Output Synapses of Parvalbumin-
729   Expressing Inhibitory Neurons. J Neurosci 37, 4604-4617. 10.1523/jneurosci.3736-16.2017.

34.   Morairty, S.R., Dittrich, L., Pasumarthi, R.K., Valladao, D., Heiss, J.E., Gerashchenko, D., and
Kilduff, T.S. (2013). A role for cortical nNOS/NK1 neurons in coupling homeostatic sleep drive
to EEG slow wave activity. Proc Natl Acad Sci U S A *110*, 20272-20277.
10.1073/pnas.1314762110.

35.   Cirelli, C., Huber, R., Gopalakrishnan, A., Southard, T.L., and Tononi, G. (2005). Locus ceruleus
control of slow-wave homeostasis. J Neurosci *25*, 4503-4511. 10.1523/jneurosci.4845-
04.2005.

36.   Willis, E.F., MacDonald, K.P.A., Nguyen, Q.H., Garrido, A.L., Gillespie, E.R., Harley, S.B.R.,
Bartlett, P.F., Schroder, W.A., Yates, A.G., Anthony, D.C., et al. (2020). Repopulating
Microglia Promote Brain Repair in an IL-6-Dependent Manner. Cell *180*, 833-846.e816.
10.1016/j.cell.2020.02.013.

37.   Zhang, Y., Zhao, L., Wang, X., Ma, W., Lazere, A., Qian, H.H., Zhang, J., Abu-Asab, M., Fariss,
R.N., Roger, J.E., and Wong, W.T. (2018). Repopulating retinal microglia restore endogenous
organization and function under CX3CL1-CX3CR1 regulation. Sci Adv *4*, eaap8492.
10.1126/sciadv.aap8492.

38.   Favuzzi, E., Huang, S., Saldi, G.A., Binan, L., Ibrahim, L.A., Fernández-Otero, M., Cao, Y., Zeine,
A., Sefah, A., Zheng, K., et al. (2021). GABA-receptive microglia selectively sculpt developing
inhibitory circuits. Cell *184*, 4048-4063.e4032. 10.1016/j.cell.2021.06.018.

39.   Wilton, D.K., Dissing-Olesen, L., and Stevens, B. (2019). Neuron-Glia Signaling in Synapse
Elimination. Annu Rev Neurosci *42*, 107-127. 10.1146/annurev-neuro-070918-050306.

40.   Badimon, A., Strasburger, H.J., Ayata, P., Chen, X., Nair, A., Ikegami, A., Hwang, P., Chan, A.T.,
Graves, S.M., Uweru, J.O., et al. (2020). Negative feedback control of neuronal activity by
microglia. Nature *586*, 417-423. 10.1038/s41586-020-2777-8.

41.   Guillaumin, M.C.C., Harding, C.D., Krone, L.B., Yamagata, T., Kahn, M.C., Blanco-Duque, C.,
Banks, G.T., Nolan, P.M., Peirson, S.N., and Vyazovskiy, V.V. (2023). Deficient synaptic
neurotransmission results in a persistent sleep-like cortical activity across vigilance states in
mice. bioRxiv, 2023.2005.2011.540034. 10.1101/2023.05.11.540034.

42.   Seifinejad, A., Vassalli, A., and Tafti, M. (2021). Neurobiology of cataplexy. Sleep Med Rev *60*,
101546. 10.1016/j.smrv.2021.101546.

43.   Osorio-Forero, A., Cardis, R., Vantomme, G., Guillaume-Gentil, A., Katsioudi, G., Devenoges,
C., Fernandez, L.M.J., and Lüthi, A. (2021). Noradrenergic circuit control of non-REM sleep
substates. Current biology : CB *31*, 5009-5023.e5007. 10.1016/j.cub.2021.09.041.

44.   Ma, C., Li, B., Silverman, D., Ding, X., Li, A., Xiao, C., Huang, G., Worden, K., Muroy, S., Chen,
W., et al. (2023). Microglia Regulate Sleep via Calcium-Dependent Modulation of
Norepinephrine Transmission. bioRxiv, 2023.2007.2024.550176.
10.1101/2023.07.24.550176.

45.   Ma, C., Li, B., Silverman, D., Ding, X., Li, A., Xiao, C., Huang, G., Worden, K., Muroy, S., Chen,
W., et al. (2024). Microglia regulate sleep through calcium-dependent modulation of
norepinephrine transmission. Nat Neurosci *27*, 249-258. 10.1038/s41593-023-01548-5.

46.   Stowell, R.D., Sipe, G.O., Dawes, R.P., Batchelor, H.N., Lordy, K.A., Whitelaw, B.S., Stoessel,
M.B., Bidlack, J.M., Brown, E., Sur, M., and Majewska, A.K. (2019). Noradrenergic signaling in
the wakeful state inhibits microglial surveillance and synaptic plasticity in the mouse visual
cortex. Nat Neurosci *22*, 1782-1792. 10.1038/s41593-019-0514-0.

47.   Liu, Y.U., Ying, Y., Li, Y., Eyo, U.B., Chen, T., Zheng, J., Umpierre, A.D., Zhu, J., Bosco, D.B.,
Dong, H., and Wu, L.J. (2019). Neuronal network activity controls microglial process
surveillance in awake mice via norepinephrine signaling. Nat Neurosci *22*, 1771-1781.
10.1038/s41593-019-0511-3.

48.   Maugard, M., Vigneron, P.A., Bolanos, J.P., and Bonvento, G. (2021). l-Serine links
metabolism with neurotransmission. Prog Neurobiol *197*, 101896.
10.1016/j.pneurobio.2020.101896.

49.   Vávrová, K., Hrabálek, A., Dolezal, P., Holas, T., and Zbytovská, J. (2003). L-Serine and glycine
based ceramide analogues as transdermal permeation enhancers: polar head size and
hydrogen bonding. Bioorg Med Chem Lett *13*, 2351-2353. 10.1016/s0960-894x(03)00409-8.

50.   Liu, H., Wang, X., Chen, L., Chen, L., Tsirka, S.E., Ge, S., and Xiong, Q. (2021). Microglia
modulate stable wakefulness via the thalamic reticular nucleus in mice. Nat Commun *12*,
4646. 10.1038/s41467-021-24915-x.

51.   Masuda, T., Amann, L., Sankowski, R., Staszewski, O., Lenz, M., P, D.E., Snaidero, N., Costa
Jordão, M.J., Böttcher, C., Kierdorf, K., et al. (2020). Novel Hexb-based tools for studying
microglia in the CNS. Nat Immunol *21*, 802-815. 10.1038/s41590-020-0707-4.

52.   Masuda, T., Amann, L., Monaco, G., Sankowski, R., Staszewski, O., Krueger, M., Del Gaudio,
F., He, L., Paterson, N., Nent, E., et al. (2022). Specification of CNS macrophage subsets
occurs postnatally in defined niches. Nature *604*, 740-748. 10.1038/s41586-022-04596-2.

53.   Pinto, M.J., Cottin, L., Dingli, F., Laigle, V., Ribeiro, L.F., Triller, A., Henderson, F., Loew, D.,
Fabre, V., and Bessis, A. (2023). Microglial TNFα orchestrates protein phosphorylation in the
cortex during the sleep period and controls homeostatic sleep. Embo j *42*, e111485.
10.15252/embj.2022111485.

54.   Lei, F., Cui, N., Zhou, C., Chodosh, J., Vavvas, D.G., and Paschalis, E.I. (2020). CSF1R inhibition
by a small-molecule inhibitor is not microglia specific; affecting hematopoiesis and the
function of macrophages. Proc Natl Acad Sci U S A *117*, 23336-23338.
10.1073/pnas.1922788117.

55.   Favuzzi, E., Huang, S., Saldi, G.A., Binan, L., Ibrahim, L.A., Fernández-Otero, M., Cao, Y., Zeine,
A., Sefah, A., Zheng, K., et al. (2021). GABA-receptive microglia selectively sculpt developing
inhibitory circuits. Cell *184*, 5686. 10.1016/j.cell.2021.10.009.

56.   Behringer, S., Wingert, V., Oria, V., Schumann, A., Grunert, S., Cieslar-Pobuda, A., Kolker, S.,
Lederer, A.K., Jacobsen, D.W., Staerk, J., et al. (2019). Targeted Metabolic Profiling of
Methionine Cycle Metabolites and Redox Thiol Pools in Mammalian Plasma, Cells and Urine.
Metabolites *9*. 10.3390/metabo9100235.

57.   Bravo, A.C., Aguilera, M.N.L., Marziali, N.R., Moritz, L., Wingert, V., Klotz, K., Schumann, A.,
Grünert, S.C., Spiekerkoetter, U., Berger, U., et al. (2022). Analysis of S-Adenosylmethionine
and S-Adenosylhomocysteine: Method Optimisation and Profiling in Healthy Adults upon
Short-Term Dietary Intervention. Metabolites *12*, 373.

58.   Hannibal, L., Theimer, J., Wingert, V., Klotz, K., Bierschenk, I., Nitschke, R., Spiekerkoetter, U.,
and Grunert, S.C. (2020). Metabolic Profiling in Human Fibroblasts Enables Subtype
Clustering in Glycogen Storage Disease. Front Endocrinol (Lausanne) *11*, 579981.
10.3389/fendo.2020.579981.

59.   Maier, J.P., Ravi, V.M., Kueckelhaus, J., Behringer, S.P., Garrelfs, N., Will, P., Sun, N., von Ehr,
J., Goeldner, J.M., Pfeifer, D., et al. (2021). Inhibition of metabotropic glutamate receptor III
facilitates sensitization to alkylating chemotherapeutics in glioblastoma. Cell Death Dis *12*,
723. 10.1038/s41419-021-03937-9.

60.   Moritz, L., Klotz, K., Grunert, S.C., Hannibal, L., and Spiekerkoetter, U. (2023). Metabolic
phenotyping in phenylketonuria reveals disease clustering independently of metabolic
control. Mol Genet Metab *138*, 107509. 10.1016/j.ymgme.2023.107509.

61.   Franken, P., Malafosse, A., and Tafti, M. (1998). Genetic variation in EEG activity during sleep
in inbred mice. Am J Physiol *275*, R1127-1137. 10.1152/ajpregu.1998.275.4.R1127.

62.   Li, S., Franken, P., and Vassalli, A. (2018). Bidirectional and context-dependent changes in
theta and gamma oscillatory brain activity in noradrenergic cell-specific Hypocretin/Orexin
receptor 1-KO mice. Sci Rep *8*, 15474. 10.1038/s41598-018-33069-8.

63.   Tort, A.B., Kramer, M.A., Thorn, C., Gibson, D.J., Kubota, Y., Graybiel, A.M., and Kopell, N.J.
(2008). Dynamic cross-frequency couplings of local field potential oscillations in rat striatum
and hippocampus during performance of a T-maze task. Proc Natl Acad Sci U S A *105*, 20517-
20522. 10.1073/pnas.0810524105.

Figure 1

Figure 2

Figure 3

Figure 4

A

PCA - Cortex

B

C

Figure 5