When the tap runs dry: The multi-tissue gene expression and physiological responses of

water deprived *Peromyscus eremicus*

Danielle M. Blumstein[1], Matthew D. MacManes[1]

University of New Hampshire, Molecular, Cellular, and Biomedical Sciences Department,

Durham, NH 03824 (DMB Dani.Blumstein@unh.edu, Orcid ID: 0000-0003- 8 2129-5541;

MDM Matthew.MacManes@unh.edu, Orcid ID: 0000-0002-2368-6960)

Abstract

The harsh and dry conditions of desert environments have resulted in genomic adaptations,

allowing for desert organisms to withstand prolonged drought, extreme temperatures, and limited

food resources. Here, we present a comprehensive exploration of gene expression across five

tissues (kidney, liver, lung, gastrointestinal tract, and hypothalamus) and 19 phenotypic

measurements to explore the whole-organism physiological and genomic response to water

deprivation in the desert-adapted cactus mouse (*Peromyscus eremicus*). The findings encompass

the identification of differentially expressed genes and correlative analysis between phenotypes

and gene expression patterns across multiple tissues. Specifically, we found robust activation of

the vasopressin renin-angiotensin-aldosterone system (RAAS) pathways, whose primary

function is to manage water and solute balance. Animals reduce food intake during water

deprivation, and upregulation of *PCK1* highlights the adaptive response to reduced oral intake

via its actions aimed at maintained serum glucose levels. Even with such responses to maintain

water balance, hemoconcentration still occurred, prompting a protective downregulation of genes

responsible for the production of clotting factors while simultaneously enhancing angiogenesis

which is thought to maintains tissue perfusion. In this study, we elucidate the complex

mechanisms involved in water balance in the desert-adapted cactus mouse, *P. eremicus*. By

prioritizing a comprehensive analysis of whole-organism physiology and multi-tissue gene

expression in a simulated desert environment, we describe the complex and successful response

of regulatory processes.

Introduction

Genomic adaptations play a pivotal role in enabling life to persist in the harsh and dynamic conditions of desert environments. Evolutionary processes have shaped the genomes of these organisms to enhance their capacity to withstand prolonged drought, extreme temperatures, and limited food resources (Colella et al., 2021; Tigano et al., 2022, 2020; Wu et al., 2014; Yang et al., 2016). Understanding these genetic underpinnings of desert adaptation not only contributes to our comprehension of evolutionary biology, but it also holds promise for insights into how these adaptations may be leveraged to address challenges posed by water scarcity and climate change in other organisms, including humans, and in other ecosystems. Studies of desert mammals have provided evidence of positive selection on genes related to food storage (Jirimutu et al., 2012; Wu et al., 2014), water reabsorption (Jirimutu et al., 2012; Marra et al., 2014, 2012; Yang et al., 2016), osmoregulation (Colella et al., 2021; Kordonowy and MacManes, 2017, 2016; MacManes and Eisen, 2014), fat metabolism (Chebii et al., 2020; Colella et al., 2021; Kim et al., 2016; Sugden et al., 2018; Tigano et al., 2020), thyroid-induced metabolism (Malaspinas et al., 2016), and salt regulation (Ababaikeri et al., 2020). These genetic insights suggest the molecular basis of observed phenotypes, including enhanced metabolic water production (Frank, 1988; MacMillen and Hinds, 1983; Walsberg, 2000), reduced water loss (Blumstein and MacManes, 2023; Frank, 1988; Schmidt-Nielsen, 1975), tolerance to high-salt diets (Ali et al., 2019; Jirimutu et al., 2012), and coping with starvation and dehydration (Blumstein and MacManes, 2023, 2023; Boumansour et al., 2021; Kordonowy et al., 2017; Kordonowy and MacManes, 2017; MacManes, 2017), which are all common in desert-dwelling mammals. However, it remains unclear how water deprivation affects the activities of other organs with respect to their expression of genes in the whole-organism context.

Osmoregulation, or the process through which animals manage water and solute balance, is critical for desert animals. It involves the maintenance of internal fluid homeostasis, with water intake being dependent on factors such as drinking, dietary sources, and metabolic water, while water output is regulated through processes including waste removal (i.e., urine and feces), respiration, perspiration, and reduced food intake (Bouby and Fernandes, 2003; Popkin et al.,

2010; Watts and Boyle, 2010). Failure to maintain water and solute homeostasis can result in

impaired renal, reproductive, and cardiovascular function, affect an animal's ability to regulate

its body temperature, and ultimately lead to death (Popkin et al., 2010). During water

deprivation, a decrease in extracellular water volume results in heightened plasma osmolality due

to an elevated concentration of solutes, primarily sodium, is detected by osmoreceptors

(Greenleaf, 1992; Leib et al., 2016; Thornton, 2010). Osmotic balance is then intricately

managed through thought two distinct multisystem mechanisms, the renin-angiotensin-

aldosterone system (RAAS) and by vasopressin (Aisenbrey et al., 1981; Bouby and Fernandes,

2003; Greenleaf, 1992; Roberts et al., 2011).

In response to changes in osmotic pressure, osmoreceptors in the hypothalamus become

activated, aldosterone is released from the adrenal glands, and vasopressin is produced in the

hypothalamus and released from the posterior pituitary gland (Leib et al., 2016; Thornton, 2010;

Yoshimura et al., 2021). This promotes water reabsorption in the kidneys by enhancing the water

permeability of epithelial cells lining renal collecting ducts (Fuller et al., 2020). Additionally, the

kidneys retain sodium while also excreting solutes via urination (Qian, 2018). Proteins related to

the transport of water are translocated from within the cell to the cell surface, forming water

channels, resulting in increased water reabsorption from the tubule system of the nephrons back

into the bloodstream (Brown et al., 1995; Kortenoeven and Fenton, 2014; Verkman, 2002) and

the retention of sodium in the distal tubules (Goodfriend, 2006). Renin is released from the

kidney, and acts on angiotensinogen that is released from the liver (Blair et al., 1997; Greenleaf,

1992). This triggers the formation of angiotensin I, which is then converted into angiotensin II, a

hormone involved in regulating blood pressure and peripheral circulation, by the release of

angiotensin-converting enzyme in the lungs (Fountain et al., 2023; Greenleaf, 1992; Santos et al.,

2019). As a result, RAAS impacts water and solute retention by promoting vasoconstriction,

stimulating the release of aldosterone, and enhancing the reabsorption of sodium and water in the

kidneys. This orchestrated response helps regulate blood pressure and maintain fluid balance in

the body.

If water homeostasis is not achieved, blood volume continues to decrease, resulting in

hemoconcentration. As dehydration leads to a higher concentration of erythrocytes, its viscosity

increases, potentially enhancing the risk of spontaneous clot formation and hindering blood flow

through vessels. This elevated blood viscosity might impede the efficient delivery of oxygen and

nutrients to various tissues and organs, triggering alterations in vascular dynamics in nutrient-

deprived, hypoxic environments (Alonso et al., 2005; Cavaglia et al., 2001; Swain et al., 2003).

Studies suggest that impaired flow-dynamics typical of dehydration may induce reversible

changes in angiogenesis, altering the growth and development of blood vessels to regulate blood

flow and distribution (Alim et al., 2019; Dumas et al., 2020). The modulation of angiogenesis

during dehydration reflects the body's dynamic response in adjusting vascular networks to

manage the metabolic demands of tissues.

Vasopressin receptors expressed in the kidneys, lungs, liver, hypothalamus etc., further

affect vasoconstriction, glycogenolysis, water reabsorption, thermoregulation, and food intake

(Yoshimura et al., 2021). Water deprived animals have been shown to limit food intake

(Armstrong et al., 1980, Blumstein, MacManes, personal observation) as an adaptive mechanism

allowing for osmotically sequestered water in the GI to be reabsorbed into the systemic

vasculature (Kutscher, 1968; Lepkovsky et al., 1957; Schoorlemmer and Evered, 1993) which

thereby reduces solute load (Rowland, 2007). Decreased food intake may be driven by the

expression of vasopressin in the hypothalamus (Yoshimura et al., 2021), suggesting another link

between eating and drinking. To survive reduced food intake during water deprivation (i.e.,

dehydration anorexia) while maintaining blood glucose concentrations, rodents have been

previous studies have shown increased glycogenolysis, lipolysis, and/ or gluconeogenesis (Salter

and Watts, 2003; Schoorlemmer and Evered, 2002; Watts and Boyle, 2010).

In this this study, we have performed a comprehensive analysis of gene expression across

five tissues relevant to the response to dehydration (kidney, liver, lung, gastrointestinal tract, and

hypothalamus) and used 19 phenotypic measurements to assess the whole-organism

physiological and genomic response to water deprivation in a hot and dry environment in the

desert-adapted cactus mouse (*Peromyscus eremicus*). The results of this study, including: 1) a

robust activation of RAAS, as seen by upregulation of *AGT* across all five tissues, 2)

upregulation of *PCK1*, reflecting an adaptive response to maintain blood glucose levels during

decreased oral intake, 3) a broad decrease in genes related to coagulation, possibly in response to

hemoconcentration and 4) a clear signal of vascular remodeling. Overall, the lung experienced

the largest number of changes in gene expression, followed by tissues involved in RAAS and

then the hypothalamus. Each tissue differed in its own way with regards to the number of genes

with highly correlated expression profiles; however, this was not due to different gene expression
between the various tissues.
126
Methods
Animal Care, RNA Extraction, and Sequencing
Captive born, sexually mature, non-reproductive healthy male and female *P. eremicus* were
reared in an environmental chamber designed to simulate the Sonoran desert (Blumstein et al.,
2022; Blumstein and MacManes, 2023; Colella et al., 2021; Kordonowy et al., 2017). All mice
were subjected to standard animal care procedures before the experiment which included a health
assessment conducted by licensed veterinary staff following animal care procedures guidelines
established by the American Society of Mammalogists (Sikes et al., 2016) and approved by the
University of New Hampshire Institutional Animal Care and Use Committee under protocol
number 210602. Mice were provided a standard diet and fed *ad libitum* (LabDiet® 5015*,
26.101% fat, 19.752% protein, 54.148% carbohydrates, energy 15.02 kJ/g, food quotient [FQ]
0.89). Animals were randomly selected and assigned to the two water treatment groups (n=9 of
each group, female mice with water, female mice without water, male mice with water, and male
mice without water, total n=36). Prior to the start of the experiment, a temperature-sensing
passive integrated transponder (PIT) tag (BioThermo13, accuracy ±0.5°C, BioMark®, Boise, ID,
USA) was implanted subdermally. At the start of the experiment (day 0, time 0hr, 10:00), mice
were weighed (rounded to the nearest tenth of a gram) on a digital scale and water was removed
from chambers corresponding to those animals in the dehydration group. Mice were
metabolically phenotyped for the duration of the experiment (Blumstein and MacManes, 2023)
using a pull flow-through respirometry system from Sable Systems International (SSI). Rates of
CO_2 production, O_2 consumption, and water loss were calculated using equations 10.6, 10.5, and
10.9, respectively, from Lighton (2018). Respiratory quotient (RQ, the ratio of VCO_2 to VO_2)
and energy expenditure (EE) kJ hr^{-1} were calculated as in Lighton (2018, eq. 9.15). For
downstream analysis, we calculated the mean of the last hour of water loss, EE, and RQ for each
mouse.
At the conclusion of the experiment (day 3, time 72hr, 12:00) as described in Blumstein
and MacManes (2023), body temperature was recorded via a Biomark® HPR Plus reader, mice
were weighed, animals were euthanized with an overdose of isoflurane, and 120 µl of trunk

blood was collected for serum electrolyte measurement and analyzed with an Abaxis i-STAT®

Alinity machine using i-STAT CHEM8+ cartridges (Abbott Park, IL, USA, Abbott Point of Care

Inc). We measured the concentration of sodium (Na, mmol/L), potassium (K, mmol/L), blood

urea nitrogen (BUN, mmol/L), hematocrit (Hct, % PCV), ionized calcium (iCa, mmol/L),

glucose (Glu, mmol/L), osmolality (mmol/L), hemoglobin (Hb, g/dl), chlorine (Cl, mEq/L), total

CO_2 (TCO₂, mmol/L), and Anion gap (AnGap, mEq/L). Using Na, Glu, and BUN, we calculated

serum osmolality. To test for statistically significant ($p < 0.05$) differences, we used a student's

two-tailed t-test (stats::t.test) between the sexes for each experimental group in R v 4.0.3 (R Core

Team, 2020).

The lung, liver, kidney, a section of the large intestines (referred to as GI throughout),

and hypothalamus were collected and stored in RNAlater (Ambion) at 4°C for 12hr before being

frozen at −80°C for long-term storage. Prior to RNA extraction, the tissues were removed from

the RNAlater and a small section was dissected off. Care was taken to retain an anatomically

similar region of tissue from each animal. Tissues were mechanically lysed using a Bead Beater,

and RNA was then extracted using a standardized Trizol protocol. RNA libraries were prepared

using standard poly-A tail purification, prepared using Illumina primers, and individually dual-

barcoded using a New England Biolabs Ultra II Directional kit (NEB #E7765). Individually

barcoded samples were pooled and sequenced paired end and 150 bp in length on two lanes of a

Novaseq at the University of New Hampshire Hubbard Center of Genome Studies.

Genome Alignment and Differential Gene Expression

All the code used to analyze the data are located at the GitHub repository

(https://github.com/DaniBlumstein/dehy_rnaseq). The *P. eremicus* genome version 2.0.1 from

the DNA Zoo Consortium (dnazoo.org) was indexed, and reads from each individual were

aligned to the genome using STAR version 2.7.10b (Dobin et al., 2013), allowing a 10 base

mismatches, a maximum of 20 multiple alignments per read, and discarding reads that mapped at

<30% of the read length. Aligned reads were counted using HTSEQ-COUNT version 2.0.2

(Anders et al., 2015).

Counts from HTSEQ-COUNT were exported as csv files, and all downstream statistical

analyses were conducted in R v4.0.3 (R Core Team, 2020). Counts were merged into a gene-

level count by combining all counts that mapped to the same gene. Low expression genes

(defined as having 10 or less counts in 8 or more individuals) were removed from downstream

analyses. Differential gene expression analysis was conducted in R using DESEQ2 (Love et al.,

2014). For the dataset as a whole, we performed three models to test for the effects of sex, water

access, and tissue type. For each tissue, we performed two models, testing the effect of and

identifying genes specific to sex and water access with a Wald test. Results were visualized using

GGPLOT2 (Wickham, 2016).

Weighted Gene Correlation Network Analysis

To identify the regulation of gene expression associated with responses to water access, we

performed a weighted gene correlation network analysis (WGCNA), a network-based statistical

approach that identifies clusters of genes with highly correlated expression profiles (modules),

(Langfelder and Horvath, 2008) for each tissue independently. This approach allows us to relate

gene expression with physiological phenotypes (mean EE, water loss, RQ, total weight loss,

proportional weight loss, sex, body temperature, water access, and the panel of electrolytes).

Prior to WGCNA, read counts were normalized within tissues using DESEQ2 (Love et al.,

2014). Module detection was done using WGCNA::blockwiseModules with networkType set to

"signed" but otherwise default parameters were used. We estimated a soft threshold power (β)

for each tissue dataset by plotting this value against mean connectivity to determine the

minimum value at which mean connectivity asymptotes, which represents scale-free topology

(liver = 15, kidney = 21, GI = 14, lung = 20, hypothalamus = 14).

Canonical Correlation Analysis

We used a Canonical Correlation Analysis (CCA) implemented in the R package vegan

(Oksanen, 2010) to investigate multivariate correlation of gene expression, by tissue, water

access, and sex, with metabolic variables (mean EE, mean RQ, mean water loss, body

temperature, and proportional weight loss) and display the three levels of information in a triplot.

We used an ANOVA to identify what response variables were significant. Significant response

variables were graphed as vectors and allowed us to identify their correlative nature; vectors

pointing in the same direction are positively correlated, while vectors pointing in opposite

direction are negatively correlated. To identify genes of interest, we selected genes that graphed

two standard deviations away from the mean for CCA1 and CCA2.

Gene Ontology

To examine gene ontology of DE genes and WGCNA modules, we cross-referenced our gene

IDs with *Homo sapiens* gene IDs via Ensembl before running Gene Ontology (GO) analyses.

Each analysis above resulted in a list or lists of genes that were used as input for the GO analysis

using the R package gprofiler (Kolberg et al., 2023). From there, we identified the topmost 20

significant GO terms based on g:SCS corrected p-values (Reimand et al., 2007) for each up and

downregulated list for each tissue and for each significant module from the individual WGCNA

analysis.

Consensus gene list and KEGG pathway analysis

We generated a high confidence consensus list of genes from the results of the three orthogonal

analyses; DE, WGCNA, and genes located two standard deviations away from the origin in the

CCA. We then selected three KEGG pathways (Renin-Angiotensin system KEGG pathway

[hsa04614], vasopressin-regulated water reabsorption pathway [hsa04962], and insulin resistance

KEGG pathway [hsa04931], Kanehisa et al., 2023; Kanehisa and Goto, 2000) based on genes in

our consensus gene set and cross references the genes in those pathways with significantly

differently expressed genes in our five tissue datasets.

Results

No health issues were detected by veterinary staff, no animals were removed prior to the end of

the experiment, and all mice were active at the end of the experiment.

Genomic Data

We obtained an average of 21.44 million reads (+- 11.6 million SD) per sample

(PRJNA1048512). On average, 78.33% of reads were uniquely mapped per sample (+- 2.12%

SD). Data on the number of reads and mapping rate per sample are located in Supplemental File

1, raw read files are archived at NCBI SRA BioProject: PRJNA1048512, and all gene expression

count data and code used to analyze the data are located at the GitHub repository

(https://github.com/DaniBlumstein/dehy_rnaseq).

Electrolytes and Physiological Phenotypes

The same mice used to generate the electrolyte and physiology data more fully described in

Blumstein and MacManes (2023) are also used in the study described herein for RNAseq

analysis. When comparing males and females separately, the following electrolytes showed

significant differences with and without access to water: Na (male and female Na p = 0.0016 and

p = 0.0026 respectively), BUN (p = 0.001/0.003), Hct (p = 0.002/0.001), osmolality (p = 8.2^{-}

^{05}/0.0001), Cl (p = 0.02/0.007), Hb (p = 0.017/0.009), and TCO2 (female p = 0.017) (Table 1).

When comparing males to females within each water treatment (with or without access to water),

no significant differences were found in the electrolyte levels (Table 1). Both males and females

experienced significant weight loss (p = 0.001, 0.005) and proportional weight loss (p = 2.2e-16,

2.659e-09) at the end of the experiment (Blumstein and MacManes, 2023). Body temperature

was significantly lower for female mice without access to water, but not for males (p = 0.0003).

To relate whole-organism physiology data to gene expression data, we calculated the

means for each mouse from the last hour of data collected in Blumstein and MacManes (2023,

data located at: https://github.com/DaniBlumstein/dehy_phys) for WLR, EE, and RQ for the

same 18 adult females and 18 adult males (n=9 of each treatment, total n=36) used in this study.

Within sex, WLR significantly different between water groups (male and female, p = 0.001 and

0.002), RQ was significantly different between water groups for males (p=0.0003), and EE was

not significantly different for either males or females.

sex	all		female		male	
water access	no	yes	no	yes	no	yes
Hct	43.31	33.18	44.38	32.50	42.25	33.78
Hb	14.73	11.28	15.09	11.05	14.36	11.48
Na	155.75	144.47	159.25	144.75	152.25	144.22
K	6.58	5.88	6.19	5.85	6.98	5.91
Cl	123.63	115.59	125.88	115.38	121.38	115.78
TCO2	24.31	20.53	25.63	20.50	23.00	20.56
BUN	56.75	33.53	54.25	33.63	59.25	33.44
Crea	0.24	0.20	0.28	0.20	0.21	0.20
Glu	116.88	121.53	117.38	122.38	116.38	120.78
iCa	1.25	1.30	1.28	1.31	1.22	1.28
AnGap	15.19	15.18	14.50	15.38	15.88	15.00
osmolality	300.68	290.57	306.13	274.05	295.22	307.10

change in weight	-5.07	-0.03	-4.75	0.08	-5.38	-0.15
body temperature	35.46	36.13	34.94	36.16	35.98	36.10

Table 1

Mean measurements for serum electrolyte measurements (Na = Sodium (mmol/L), K =

Potassium (mmol/L), Cr = Creatinine (μmol/L), BUN = Blood Urea Nitrogen (mmol/L), Hct =

Hematocrit (% PCV), iCa = Ionized Calcium (mmol/L), and osmolality (mmol/L), change in

weight (g), and body temperature (°C) for female (n=18), male (n=18), and all *Peromyscus*

eremicus (n=36) with and without access to water. Data were collected and are further described

in Blumstein and MacManes (2023).

Differential Gene Expression

We cross-referenced our gene IDs with *Homo sapiens* gene IDs. Patterns of gene expression data

are largely driven by tissue type (PC1: 42% variance and PC2: 26% variance, Supplemental

Figure 1).

We then conducted all downstream analyses (except for CCA, see below) on each tissue

independently. Count and sample data were filtered to include only the tissue of interest and low

expression genes were removed. This resulted in removing 4929 genes in the kidney and left

12083, 4390 genes removed in the GI leaving 12622 genes, 3623 genes removed in the

hypothalamus leaving 13389 genes, 5887 genes were removed in the liver leaving 11125 genes,

and 4070 genes were removed in the lung leaving 12942 genes. Within each tissue, we found

many differentially expressed genes ($p < 0.05$) between water treatments (Figure 1, Table 2) and

few differentially expressed genes between sex (Table 2).

Figure 1

Log2FC of all genes across the lung (lu), liver (liv), gastrointestinal tract (gi), hypothalamus

(hyp), and kidney (kid) of *Peromyscus eremicus* with water vs without water. Blue and green

colored dots indicate p<0.05, whereas grey dots indicate p>=0.05.

Tissue	DE between water treatments with water vs without water		DE between sexes males vs females	
	Up	Down	Up	Down
kidney	793	1056	5	7
liver	933	1122	1	3
lung	2190	2183	5	5
hypothalamus	1	8	1	2
gastrointestinal tract	956	703	1	1

Table 2

The number of differentially expressed (DE) genes in the lung, liver, gastrointestinal tract,

hypothalamus, and kidney of *Peromyscus eremicus* for with water vs without water and males vs

females (adjusted p-value < 0.05).

Weighted Gene Correlation Network Analysis

A total of 12083 genes in the kidney were successfully assigned into 13 modules with the

number of genes per module ranging from 31 – 7008. A full list of gene assignments is available

in Supplemental Table 3. Of the 13 modules identified, 9 modules were significant for three or

more phenotypes (Supplemental Table 2). We identified 24 individual modules using 12622

genes for the GI. The modules contained 33-3530 genes each (Supplemental Table 2). Of these

modules, 14 were significantly correlated with three or more phenotypes (Supplemental Table 2).

In the lung, 12942 genes were assigned to 13 modules. Each module contained 23-7116 genes

(Supplemental Table 2). Nine modules were significant for three or more phenotypes

(Supplemental Table 2). A total of 13389 genes were assigned to 18 different modules in the

hypothalamus. Modules contained 30-2319 genes (Supplemental Table 2). Of the 18 modules,

three modules were significant for three or more phenotypes (Supplemental Table 2). Finally,

11125 genes were assigned to 18 modules in the liver, with the number of genes per module

ranging from 28-3116 (Supplemental Table 2). Of the 18 modules, 13 modules were significant

for three or more phenotypes (Supplemental Table 2).

Gene Ontology

After filtering to the top 20 GO terms for the upregulated and downregulated significantly

differentially expressed genes for each tissue, we identified 149 unique terms. Of the 149 GO

terms, 14 terms were identified in two different tissues (Supplemental Figure 2).

For each tissue WGCNA analysis, we identified GO terms for each significant module

phenotype combination. After filtering each combination for the top 20 GO terms, we identified

a total of 379 unique GO terms. 199 GO terms were identified in the hypothalamus with 168

terms only appearing in one module, 25 identified in two modules, five in three modules, and

one in four modules (Figure 2). Within the kidney 82 unique GO terms were identified.

Specifically, 78 terms were in one module and four terms were in two modules. There were 122

unique GO terms identified in the liver with 100 of the go terms appearing in one module, nine

terms in two modules, and three terms in three different modules (Figure 2). Lastly, the lung had

52 unique GO terms identified, the fewest number any tissue, and there was no GO term overlap

for any of the modules (Figure 2). When comparing GO terms between tissues, 53 terms were in

two different tissues, five terms were in three of the tissues, and one term were in four tissues

(Figure 2).

Figure 2

Visualization of gene ontology (GO) terms to show common WGCNA modules within and

between the lung (lu), liver (liv), gastrointestinal tract (gi), hypothalamus (hyp), and kidney (kid)

of *Peromyscus eremicus*. Visualized are selections of the top 20 significant GO terms for each

phenotype module combination. The number of genes in the GO term are indicated by size of the

dots.

Canonical Correlation Analysis

We examined the relationship between gene expression and water access, physiological variables

(EE, RQ, WLR, proportional weight loss, body temperature), and tissue type using CCA

(Oksanen, 2010). CCA suggests a significant overall association between the physiological

variables and gene expression across tissues (F = 105.45, p = 0.001; CCA1 – 38.02% and CCA2

– 21.14%, Figure 3, Table 4). We identified 1233 genes two standard deviations from the origin

and found a high degree of overlap between the genes located two standard deviations in the

CCA and genes assigned to WGCNA modules (kidney: 37/10906, GI: 60/11032, lung: 69/12527,

hypothalamus: 53/1971, liver: 40/11051). Here, the proportional weight loss and WLR are

significantly correlated (both p=0.001, Table 4) with gene expression in the hypothalamus and

GI (Figure 3), but there is no overlap between the genes and the two vectors. This suggests that

proportional weight loss and WLR exert different effects on gene expression in the

hypothalamus and the GI.

	Df	ChiSquare	F	Pr(>F)
water access	1	0.00012	6.437	0.001
proportional weight loss	1	0.00009	4.801	0.001
RQ	1	0.00002	0.990	0.407
EE	1	0.00003	1.474	0.179
WLR	1	0.00004	1.986	0.096
body temperature	1	0.00002	1.212	0.261
sex	1	0.00003	1.730	0.120
tissue	4	0.02141	285.317	0.001
model	11	0.02176	105.450	0.001
residual	147	0.00276		

359   Table 4

 ANOVA results from the Canonical Correspondence Analysis. Formula: gene expression ~

water access + proportional weight loss + RQ + EE + WLR + body temperature + sex + tissue).

Figure 3

Canonical correspondence analysis (CCA) indicates correlations between normalized

differentially expressed genes and physiological measurements for *Peromyscus eremicus* with

and without access to water. The distribution of tissue samples in Euclidian space as a function

of their gene expression values is shown (points colored by tissue type and water treatment).

Outlier genes (defined as two standard deviations or more from the mean) are colored blue. Inlier

genes (defined as less than two standard deviations from the mean) are colored grey. CCA

reveals a significant relationship between proportion weight loss (F = 4.8006, P = 0.001) and,

while not significant, a strong relationship for water loss rate (WLR) ($F = 1.9856$, $P = 0.096$).

This can be seen by a subset of genes (blue) pulled in the direction of the proportion weight loss

(pro_weight) ordination vector and a subset of genes (blue) pulled in the direction of the WLR

(mean_H_2Og) ordination vector.

Consensus gene set

We identified 41 genes that were significantly DE and assigned to at least one significant module

for all five tissues as well as were located at least two standard deviations away from the origin

in our CCA triplot (Figure 3).

general function	genes
housekeeping	ANKRD13D, GPRASP2, ACOX2, ALDOB, GSTA3, HAL, HOGA1, INMT, MTARC1, SPTBN2, DES, EPHX2, FMO5
ion homeostasis	ATP1A2, ALB, APOA4, CYP2E1, CYP4B1, RGN, SLC38A4, AGT, SLC6A13, TF, HP
central nervous system	MBP, MAG, PLP1, MAOB, SLC6A13
apoptosis	RELL2, PYCARD
coagulation	SERPINF2
blood pressure	ALB, AGT, CASZ1
angiogenesis	ANG, CASZ1
lipid	APOA2, APOA4, APOB, AZGP1, CYP2E1, CYP4B1, SLC27A2
immune response	CFI (renal failure), CTSC, PYCARD, HP
gluconeogenesis	CYP2E1 (induced by starvation), PCK1, AGT, SERPINA6
bitter taste	AZGP1

Table 5

Genes identified in all three analyses (i.e. significantly differentially expressed genes, assigned to

a significant module in WGCNA, and are outliers in CCA) in all five tissues in the study (lung

(lu), liver (liv), gastrointestinal tract (gi), hypothalamus (hyp), and kidney (kid) grouped by the

general function of the gene.

Discussion

Extensive research has been conducted on the genomic and physiological mechanisms that

control water balance in mice (Blumstein and MacManes, 2023; McCue et al., 2017; Rocha et

al., 2021). This is particularly intriguing in the context of exploring adaptations to extreme

conditions, given that the management of water has significant implications for survival. Studies
have largely focused on characterizing the response in kidneys (Rocha et al., 2023; MacManes,
2017; Peng et al., 2023; Rocha et al., 2021), which, while important, represent only a fraction of
the physiological and genomic response to dehydration. Indeed, the organismal response to an
environmental stressor (such as dehydration), is likely to involve coordination of multiple organ
systems and physiological process. However, there are key challenges to studying such responses
at the organismal level. Although we understand that organ-systems operate in tandem with other
organ systems, the identification of coordination at the level of gene expression is challenging,
even when organismal physiology is well-characterized. In addition to complications related to
biology, technical complications exist. Here, when looking at gene expression levels across
tissues, results highlight differences between the tissues rather than to elucidate the ways in
which processes in one depend on the actions of the other, thereby potentially obscuring the gene
expression signal of coordination. Further, gene interaction maps (*e.g.,* KEGG) that span organ
systems do not currently exist, mostly because the models used for their development (*e.g.,*
yeast, flies) lack such complexity present in mammalian systems.

The study of coordination at the level of physiology has been similarly difficult to
elucidate, due to limitations in our ability to collect and analyze phenotypic data at a temporal
scale that is relevant to the biological phenomenon under study (but see Blumstein et al., 2022;
Blumstein and MacManes, 2023; Colella et al., 2021; McKechnie et al., 2021; Ramirez et al.,
2022). Despite the challenges, understanding the complex interplay of multi-tissue networked
gene expression with whole-organism physiological response is critical to developing a more in-
depth understanding of how organisms respond to environmental stressors. Our focus on
physiological measurements in a simulated desert environment, as well as on collection of multi-
tissue gene expression data in the context of a whole-organism response, represents a distinctive
contribution to the field. The findings underscore the complexity of the genetic landscape
governing physiological responses to water deprivation and emphasize the need for a broader
organismal understanding of the physiological and genomic mechanisms orchestrating successful
adaptations.

Global response to acute water deprivation

The inclusion of physiology and multi-tissue gene expression data in this study provides

opportunities to understand how the tissues work together to compensate for water deprivation.

During water deprivation, blood volume decreases, which if inadequately managed, may result in

a decrease in organ perfusion and ultimately organ failure. The multisystem response to water

deprivation involves a coordinated response, including renin-angiotensin-aldosterone system

(RAAS) activation, reduced food intake, with compensatory activation of systems for preserving

levels of serum glucose, a widespread reduction of genes responsible for clotting factors, and

distinct indications of vascular restructuring to support perfusion.

The first response to water deprivation involves the upregulation of processes that are

aimed at solute management. In part regulated by the RAAS, this process begins with decreased

blood volume detected by baroreceptors found in the carotid sinuses and aortic arch. As a result,

the kidneys release the enzyme renin (Blair et al., 1997; Greenleaf, 1992). Renin then acts on a

plasma protein called angiotensinogen, produced by the liver, and converts it into angiotensin I

(Fountain et al., 2023; Santos et al., 2019). Angiotensin I is subsequently converted into

angiotensin II by the angiotensin-converting enzyme (Fountain et al., 2023; Greenleaf, 1992;

Santos et al., 2019). Angiotensin II serves as a potent vasoconstrictor, which helps increase tissue

perfusion pressure and stimulate the release of aldosterone from the adrenal glands (Fountain et

al., 2023; Santos et al., 2019). Aldosterone in turn prompts the kidneys to retain sodium while

excreting potassium, which encourages renal water retention. In further support of the activation

of the RAAS mechanism, multiple genes in the Renin-Angiotensin system KEGG pathway

(hsa04614: Kanehisa et al., 2023; Kanehisa and Goto, 2000) were found to be differentially

expressed. Among these genes were *AGT* (angiotensinogen), *ACE2* (angiotensin-converting

enzyme), *REN1* (Renin), *CMA1* (converts angiotensin I to angiotensin II), and *AGTR1*

(Angiotensin II Receptor) (supplemental table 4), which were all more highly expressed in

water-deprived mice. Further, we found two compete pathways of genes to be significantly

differentially expressed genes within hsa04614, enhanced vasoconstriction and coagulation

cascade as well as vasoconstriction, inflammation, fibrosis, antinatriuresis, reactive oxygen

species activation, and Na and water retention (Kanehisa et al., 2023). Interestingly, the

candidate gene and essential component of the RAAS mechanism, *AGT,* was significantly

upregulated in dehydrated mice, assigned to a significant WGCNA module, and defined as an

outlier in the CCA analysis (Table 5). Together with the physiological data, the consistent

genetic signature of RAAS activation, using orthogonal analytical methods, suggests a robust

whole-body RAAS response which is critical for the cactus mouse to regulate water balance.

456   At the same time, independent of the RAAS pathway but stimulated by its products,

vasopressin is released for the primary function of conserving water. Vasopressin binds to

receptors, activating a signaling cascade in the kidneys which functions to retain water by

inducing expression of water transport proteins in the late distal tubule and collecting duct,

increasing the permeability of the membrane to water. The increased water permeability allows

water to move through, from the collecting ducts back into the bloodstream, further aiding in

regulating blood volume and fluid balance (Fountain et al., 2023; Santos et al., 2019). Support

for the genetic activation of the vasopressin pathway is shown with various genes within the

vasopressin-regulated water reabsorption pathway (hsa04962, Kanehisa et al., 2023; Kanehisa

and Goto, 2000) are significantly differentially expressed. Specifically, *STX4, RAB5C, RAB11B,*

CREB3L2, *AQP3*, *VAMP2*, *DYNLL2*, *GNAS*, and *AVPI1* (supplemental table 4) were upregulated

in dehydrated mice. Interestingly, *AQP2*, a key gene in hsa04962, did not have any reads mapped

to it. It is worth noting here that the identification of differentially expressed transcripts on a

KEGG pathway can be thought of in terms of a stoichiometric problem, albeit one where neither

current tools nor current data allow us to satisfactorily solve. When attempting to use differential

expression to support the upregulation of a given pathway, it may be that only one gene, the

quantity of which is rate limiting, may be more highly expressed. Without understanding which

transcripts are rate limiting, the interpretation of KEGG pathway mapping may suffer from either

over- or under-valuation.

475   During water deprivation, animals are known to reduce the amount of solid food intake

(Armstrong et al., 1980; Hamilton and Flaherty, 1973; Salter and Watts, 2003; Schoorlemmer

and Evered, 2002; Watts and Boyle, 2010). Known as dehydration associated anorexia, this

secondary response reduces the amount of water required for digestion and facilitates water

reabsorption from the kidneys and gastrointestinal tract back to systemic circulation (Rowland,

2007; Watts and Boyle, 2010). While the magnitude of the impact in cactus mice is unknown,

diets high in fiber, like the diet consumed in this study, can result in high fecal volume, and has

been shown to account for as much as 25% of total daily water loss in rats (Radford Jr, 1959;

Rowland, 2007). Perhaps even more importantly, but unquantified, the processing of solid food

requires the production of significant quantities of digestive enzymes, all of which are water rich.

In humans, as much as 10L of fluids is excreted daily (Ma and Verkman, 1999), which

represents a significant investment of water resources. While some of these enzymes continue to

be produced during dehydration anorexia, their quantity is likely decreased. As a direct result,

the reduction of oral food intake may result in significant water savings, which at least in the

short term with the presence of sufficient glycogen stores (see below), should be related to a

reduction in water use and electrolyte derangement and therefore enhanced survival.

With limited food intake, there is no external source of glucose, nevertheless, we

observed that glucose levels were still maintained (Table 1, Further explained in Blumstein and

MacManes, 2023) which is critical for surviving dehydration. This can be achieved by enhancing

glycogenolysis and gluconeogenesis (Salter and Watts, 2003; Schoorlemmer and Evered, 2002;

Watts and Boyle, 2010), responses that are designed to maintain blood glucose levels during

fasting or starvation. In mammals, there is a vasopressin receptor in the liver that when bound

activates gluconeogenesis (Bankir et al., 2017), with secondary contributions from the kidney

(Nordlie et al., 1999). This suggests a successful two-pronged response to vasopressin secretion,

such as vasoconstriction and reabsorption of water from kidney as well as a role in feeding

behavior and energy balance. In the current study, we identified the candidate genes *PCK1,* a

main control point for the regulation of gluconeogenesis (Hatting et al., 2018) as well as

CYP2E1, a gene induced by starvation with products involved in gluconeogenesis (Harjumäki et

al., 2021; Schattenberg and Czaja, 2014), as upregulated in water-deprived animals, Further,

these genes were assigned to a significant module in all five tissues, and defined as an outlier in

the CCA analysis (Table 5). Further, the insulin resistance KEGG pathway (hsa04931, Kanehisa

et al., 2023; Kanehisa and Goto, 2000) involves various mechanisms contributing to altered

glucose metabolism and insulin responsiveness. Notably, all genes but two genes in hsa04931

are significantly up regulated in dehydrated animals (supplemental table 4)*,* all of which are

upstream of *PCK1* are in the pathway. Two of these important pathway genes (AGT and

SLC27A2, Table 5) were found in our consensus gene set. This upregulation is likely a

compensatory response to ensure the maintenance of blood glucose levels in the absence of

dietary intake. Lastly, *INMT* (Table 5), a gene in the Tryptophan metabolism pathway

(hsa00390, Kanehisa et al., 2023; Kanehisa and Goto, 2000) is upstream of the

Glycolysis/gluconeogenesis pathway (map00010, Kanehisa et al., 2023; Kanehisa and Goto,

2000), suggesting complex precursor genes and gene pathways may be contributing to glucose

homeostasis as well. The significant changes in expression observed in these genes underscore

their role in modulating glucose metabolism, shedding light on how organisms adjust to sustain

vital glucose levels during periods of reduced food intake and water deprivation.

519    During water deprivation, a key challenge is to cope with altered fluid balance and

maintain effective blood circulation and nutrient delivery under water-deficient conditions. This

process can involve the formation of new blood vessels or the remodeling of existing ones to

optimize perfusion, addressing the challenges posed by reduced fluid availability and potential

hemoconcentration. We found several overlapping GO terms related to vascular development

across all five tissues (vascular development [GO:0001944], circulatory system development

[GO:0072359], angiogenesis [GO:0001525], blood vessel development [GO:0001568], blood

vessel morphogenesis [GO:0048514], and regulation of vasculature development

[GO:1901342]). These GO terms were identified in a myriad of gene modules (water treatment,

EE, Hb, TCO2, WLR, and body temperature, Figure 2). Additionally, we identified two genes

from our consensus gene set that are mediators of new blood vessel formation and

morphogenesis (*ANG*, *CASZ*, Table 5) Prior research has indicated that persistent activation of

neuronal systems could modify local blood circulation through angiogenesis. Specifically, rats

reared in complex environments had increased capillary density in the visual cortex (Cavaglia et

al., 2001). Long-term motor activity has been documented to induce the development of new

blood vessels within the cerebellar cortex (Black et al., 1990) and primary motor cortex (Swain

et al., 2003), and hyperosmotic stimuli, similar to what experimental animals experience, has

been shown to modify the vasculature and induced reversible angiogenesis throughout the

hypothalamic nuclei (Alonso et al., 2005). Furthermore, Alim et al. (2019) observed seasonal

differences in vascularization, showing blood capillaries were thicker during the winter,

suggesting less diffusion across the membrane, compared summer in the dromedary camel.

540    To maintain blood flow and prevent the formation of clots that could impede circulation,

downregulation of coagulation factors during periods of reduced fluid intake could be a

protective mechanism against the potential risks associated with increased blood viscosity due to

hemoconcentration. As dehydration leads to reduced water content in the blood, resulting in

thicker blood consistency there is a risk for blood clot formation. We found several

downregulated GO terms related to hemoconcentration (regulation of coagulation

[GO:0050818], blood microparticle [GO:0072562], hemopoiesis [GO:0030097], serine

hydrolase activity [GO:0017171], and regulation of hemostasis [GO:1900046]) were identified

in the hypothalamus and liver (Figure 3) as well as four genes in our consensus gene list

involved in blood composition and concentration (*ALB*, *HP*, *SERPINA6*, *SERPINF2*, Table 5) .

This could indicate several things: 1) To mitigate the impact of hemoconcentration, genes

responsible for clotting factors are downregulated to reduce the risk associated with increased

blood viscosity as thicker blood is more prone to clotting. 2) to decrease clotting during body

temperature dysregulation. Blumstein and MacManes (2023) discuss heterothermy as a

mechanism for substantial energy and water savings. In addition to this, relative hypothermia can

result in inactivation of coagulation enzymes and/or platelet adhesion defect (Paal et al., 2016),

suggesting further mechanisms for maintaining perfusion and limit clotting. However, it's

important to note that the timing of expression poses a potential concern regarding correlations.

Decreased body temperature was measured in Blumstein and MacManes (2023) during the dark

phase of the experiment and tissues were collected during the light phase of the experiment.

Using a single time point snapshot for each experimental condition might overlook certain

interactions, given the lag between the expression of a gene (such as a transcription factor) and

the expression of downstream effectors.

Tissue Specific Responses

When analyzing the RNAseq data across all the tissues, we observed that, as expected, samples

of the same tissue type clustered together regardless of the water-access treatment, suggesting

that the signature of tissues-specific gene expression overpowers the signature of experimental

treatment. During individual tissue analysis, the degree of sample separation in PCA space

(Supplemental Figure 1) and number of differentially expressed genes suggests tissue specific

responses. This is in part supported by downstream gene ontology (GO) terms. Specifically, GO

terms related to metabolic processes were identified in multiple significant WGCNA modules in

the kidney, but not other tissues. Hydrolase activity and renal system development uniquely in

the GI. Lipid metabolic process and detoxification uniquely in the liver. Immune response

uniquely in the lung. Lastly, starvation response uniquely in the hypothalamus (Supplemental

Figure 2). It is well known that the hypothalamus is the central regulating unit in the brain for

maintenance of energy homeostasis (Tran et al., 2022). However, few genes were identified as

differentially expressed in the hypothalamus (Figure 1), while our other analyses, WGCNA and

CCA, uncovered genes and GO terms that responded to water deprivation. This suggests that the
hypothalamus may not be well suited for bulk RNAseq studies due to the heterogeneity of the
tissue. Future studies, particularly those using single cell methods (Kephart, 2023; Marquez-
Galera et al., 2022; Yue et al., 2023) may further clarify the role that the hypothalamus plays in
the overall response to dehydration.

Conclusion

Here, we highlight the intricate mechanisms involved in regulating water balance in the desert
cactus mouse, *P. eremicus*. Our emphasis on whole-organism physiological and multi-tissue
(kidney, GI, hypothalamus, liver, and lung) gene expression analysis within a simulated desert
environment allowed us to achieve an understanding of genomic mechanisms of water
homeostasis. Previous genome scan studies (Colella et al., 2021; Kim et al., 2016; Rocha et al.,
2021; Tigano et al., 2020; Wu et al., 2014) have identified many of the same processes we found
(i.e., metabolic processes, renal system development, immune response, and starvation response)
however, our study design allows us to further interpret function because we were able to both
described the tissue specific location and link the processes to physiological measurements.

At a whole-organismal scale, we observed a robust response of the renin-angiotensin-
aldosterone system (RAAS) in dehydrated cactus mice, with upregulation of *AGT* in all five
tissues as well as upregulation of other pathway genes. Additionally, the compensatory action of
PCK1 was activated in all tissues, further supporting reduced food intake during water
deprivation and underscores the body's adaptive response. However, despite efforts to maintain
blood volume, hemoconcentration still occurs, but in response there was a downregulation of
genes responsible for coagulation (*e.g., SERPINF2*) as a protective measure against blood
clotting in all five tissues, a gene with the major role of regulating the blood clotting pathway.
The consequential thickened blood consistency poses challenges to effective blood flow through
vessels, compelling the body to initiate the construction of additional vessels to enhance blood
movement, further supported by the upregulation of *ANG* in all five tissues, further illustrating
the complex interplay of regulatory processes in response to fluid balance disturbances.

Acknowledgments

We would like to thank members of the MacManes lab for helpful comments and support on
previous versions of the manuscript; Adam Stuckert at the University of Huston for lively
discussion, valuable insight, and code development; The Animal Resources Office and
veterinary care staff at the University of New Hampshire for colony maintenance and care. This
work was supported by the National Institute of Health National Institute of General Medical
Sciences (R35 GM128843 to M.D.M.).

Author Contributions

617 Conceptualization: M.D.M.; Methodology: D.M.B.; Formal analysis: D.M.B., Investigation:
618 D.M.B., Resources: M.D.M.; Writing - original draft: D.M.B.; Writing - review & editing:
619 D.M.B., M.D.M.; Visualization: D.M.B; Supervision: M.D.M.; Project administration: M.D.M.;
620 Funding acquisition: M.D.M.

Competing Interests

623 No competing interests declared.

Data Availability

626 https://github.com/DaniBlumstein/dehy_rnaseq

627 BioProject ID: PRJNA1048512

628 http://www.ncbi.nlm.nih.gov/bioproject/1048512

References

Ababaikeri B, Abduriyim S, Tohetahong Y, Mamat T, Ahmat A, Halik M. 2020. Whole-genome
sequencing of Tarim red deer (Cervus elaphus yarkandensis) reveals demographic history
and adaptations to an arid-desert environment. *Frontiers in zoology* **17**:1–15.
Aisenbrey GA, Handelman WA, Arnold P, Manning M, Schrier RW. 1981. Vascular effects of
arginine vasopressin during fluid deprivation in the rat. *J Clin Invest* **67**:961–968.
doi:10.1172/jci110146
Ali A, Baby B, Vijayan R. 2019. From desert to medicine: a review of camel genomics and
therapeutic products. *Frontiers in genetics* **10**:17.
Alim FZD, Romanova EV, Tay Y-L, Rahman AY bin A, Chan K-G, Hong K-W, Rogers M,
Southey BR, Greenwood MP, Mecawi AS, Mustafa MR, Mahy N, Campbell C, Antunes-
Rodrigues J, Sweedler JV, Murphy D, Hindmarch CCT. 2019. Seasonal adaptations of
the hypothalamo-neurohypophyseal system of the dromedary camel. *PLOS ONE*
14:e0216679. doi:10.1371/journal.pone.0216679

Alonso G, Galibert E, Duvoid-Guillou A, Vincent A. 2005. Hyperosmotic stimulus induces
reversible angiogenesis within the hypothalamic magnocellular nuclei of the adult rat: a
potential role for neuronal vascular endothelial growth factor. *BMC Neurosci* **6**:20.
doi:10.1186/1471-2202-6-20
Anders S, Pyl PT, Huber W. 2015. HTSeq—a Python framework to work with high-throughput
sequencing data. *bioinformatics* **31**:166–169.
Armstrong S, Coleman G, Singer G. 1980. Food and water deprivation: changes in rat feeding,
drinking, activity and body weight. *Neuroscience & Biobehavioral Reviews* **4**:377–402.
Bankir L, Bichet DG, Morgenthaler NG. 2017. Vasopressin: physiology, assessment and
osmosensation. *Journal of Internal Medicine* **282**:284–297. doi:10.1111/joim.12645
Black JE, Isaacs KR, Anderson BJ, Alcantara AA, Greenough WT. 1990. Learning causes
synaptogenesis, whereas motor activity causes angiogenesis, in cerebellar cortex of adult
rats. *Proceedings of the National Academy of Sciences* **87**:5568–5572.
Blair ML, Woolf PD, Felten SY. 1997. Sympathetic activation cannot fully account for increased
plasma renin levels during water deprivation. *Am J Physiol* **272**:R1197-1203.
doi:10.1152/ajpregu.1997.272.4.R1197
Blumstein DM, Colella JP, Linder E, MacManes MD. 2022. High total water loss driven by low-
fat diet in desert-adapted mice. *bioRxiv*.
Blumstein DM, MacManes MD. 2023. When the tap runs dry: The physiological effects of acute
experimental dehydration in Peromyscus eremicus. *Journal of Experimental Biology*
jeb.246386. doi:10.1242/jeb.246386
Bouby N, Fernandes S. 2003. Mild dehydration, vasopressin and the kidney: animal and human
studies. *Eur J Clin Nutr* **57**:S39–S46. doi:10.1038/sj.ejcn.1601900
Boumansour L, Benhafri N, Guillon G, Corbani M, Touati H, Dekar-Madoui A, Ouali-
Hassenaoui S. 2021. Vasopressin and oxytocin expression in hypothalamic supraoptic
nucleus and plasma electrolytes changes in water-deprived male Meriones libycus.
Animal Cells and Systems **0**:1–10. doi:10.1080/19768354.2021.1986130
Brown D, Katsura T, Kawashima M, Verkman AS, Sabolic I. 1995. Cellular distribution of the
aquaporins: A family of water channel proteins. *Histochem Cell Biol* **104**:1–9.
doi:10.1007/BF01464780
Cavaglia M, Dombrowski SM, Drazba J, Vasanji A, Bokesch PM, Janigro D. 2001. Regional
variation in brain capillary density and vascular response to ischemia. *Brain research*
910:81–93.
Chebii VJ, Oyola SO, Kotze A, Domelevo Entfellner J-B, Musembi Mutuku J, Agaba M. 2020.
Genome-wide analysis of Nubian ibex reveals candidate positively selected genes that
contribute to its adaptation to the desert environment. *Animals* **10**:2181.
Colella Jocelyn P., Blumstein DM, MacManes MD. 2021. Disentangling environmental drivers
of circadian metabolism in desert-adapted mice. *Journal of Experimental Biology* **224**.
doi:10.1242/jeb.242529
Colella Jocelyn P, Tigano A, Dudchenko O, Omer AD, Khan R, Bochkov ID, Aiden EL,
MacManes MD. 2021. Limited Evidence for Parallel Evolution Among Desert-Adapted
Peromyscus Deer Mice. *Journal of Heredity* **112**:286–302. doi:10.1093/jhered/esab009
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.
doi:10.1093/bioinformatics/bts635

Dumas SJ, Meta E, Borri M, Goveia J, Rohlenova K, Conchinha NV, Falkenberg K, Teuwen L-
A, de Rooij L, Kalucka J, Chen R, Khan S, Taverna F, Lu W, Parys M, De Legher C,
Vinckier S, Karakach TK, Schoonjans L, Lin L, Bolund L, Dewerchin M, Eelen G,
Rabelink TJ, Li X, Luo Y, Carmeliet P. 2020. Single-Cell RNA Sequencing Reveals
Renal Endothelium Heterogeneity and Metabolic Adaptation to Water Deprivation. *J Am*
Soc Nephrol **31**:118–138. doi:10.1681/ASN.2019080832
Fountain JH, Kaur J, Lappin SL. 2023. Physiology, Renin Angiotensin SystemStatPearls.
Treasure Island (FL): StatPearls Publishing.
Frank CL. 1988. Diet Selection by a Heteromyid Rodent: Role of Net Metabolic Water
Production. *Ecology* **69**:1943–1951. doi:10.2307/1941171
Fuller A, Maloney SK, Blache D, Cooper C. 2020. Endocrine and metabolic consequences of
climate change for terrestrial mammals. *Current Opinion in Endocrine and Metabolic*
Research **11**:9–14.
Goodfriend TL. 2006. Aldosterone—a hormone of cardiovascular adaptation and maladaptation.
The Journal of Clinical Hypertension **8**:133–139.
Greenleaf JE. 1992. Problem: thirst, drinking behavior, and involuntary dehydration. *Medicine &*
Science in Sports & Exercise **24**:645.
Hamilton LW, Flaherty CF. 1973. Interactive effects of deprivation in the albino rat. *Learning*
and Motivation **4**:148–162.
Harjumäki R, Pridgeon CS, Ingelman-Sundberg M. 2021. CYP2E1 in Alcoholic and Non-
Alcoholic Liver Injury. Roles of ROS, Reactive Intermediates and Lipid Overload.
International Journal of Molecular Sciences **22**:8221. doi:10.3390/ijms22158221
Hatting M, Tavares CDJ, Sharabi K, Rines AK, Puigserver P. 2018. Insulin regulation of
gluconeogenesis. *Ann N Y Acad Sci* **1411**:21–35. doi:10.1111/nyas.13435
Jirimutu, Wang Zhen, Ding G, Chen G, Sun Y, Sun Zhihong, Zhang H, Wang L, Hasi S, Zhang
Y, Li Jianmei, Shi Y, Xu Z, He C, Yu S, Li S, Zhang W, Batmunkh M, Ts B, Narenbatu,
Unierhu, Bat-Ireedui S, Gao H, Baysgalan B, Li Q, Jia Z, Turigenbayila, Subudenggerile,
Narenmanduhu, Wang Zhaoxia, Wang J, Pan L, Chen Yongcan, Ganerdene Y, Dabxilt,
Erdemt, Altansha, Altansukh, Liu T, Cao M, Aruuntsever, Bayart, Hosblig, He F, Zha-ti
A, Zheng G, Qiu F, Sun Zikui, Zhao L, Zhao W, Liu B, Li C, Chen Yunqin, Tang X, Guo
C, Liu W, Ming L, Temuulen, Cui A, Li Yi, Gao J, Li Jing, Wurentaodi, Niu S, Sun T,
Zhai Z, Zhang M, Chen C, Baldan T, Bayaer T, Li Yixue, Meng H, The Bactrian Camels
Genome Sequencing and Analysis Consortium. 2012. Genome sequences of wild and
domestic bactrian camels. *Nat Commun* **3**:1202. doi:10.1038/ncomms2192
Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. 2023. KEGG for
taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res* **51**:D587–D592.
doi:10.1093/nar/gkac963
Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*
Res **28**:27–30. doi:10.1093/nar/28.1.27
Kephart M. 2023. A 10x Visium Approach: A Spatial RNA-Seq Analysis of Renal Tissue in
Peromyscus eremicus. Durham: University of New Hampshire.
Kim E-S, Elbeltagy AR, Aboul-Naga AM, Rischkowsky B, Sayre B, Mwacharo JM, Rothschild
MF. 2016. Multiple genomic signatures of selection in goats and sheep indigenous to a
hot arid environment. *Heredity* **116**:255–264. doi:10.1038/hdy.2015.94

Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. 2023. g: Profiler—interoperable
web service for functional enrichment analysis and gene identifier mapping (2023
update). *Nucleic Acids Research* gkad347.
Kordonowy L, Lombardo KD, Green HL, Dawson MD, Bolton EA, LaCourse S, MacManes
MD. 2017. Physiological and biochemical changes associated with acute experimental
dehydration in the desert adapted mouse, *Peromyscus eremicus*. *Physiol Rep* **5**:e13218.
doi:10.14814/phy2.13218
Kordonowy L, MacManes M. 2017. Characterizing the reproductive transcriptomic correlates of
acute dehydration in males in the desert-adapted rodent, Peromyscus eremicus. *BMC*
Genomics **18**:473. doi:10.1186/s12864-017-3840-1
Kordonowy LL, MacManes MD. 2016. Characterization of a male reproductive transcriptome
for *Peromyscus eremicus* (Cactus mouse). *PeerJ* **4**:e2617. doi:10.7717/peerj.2617
Kortenoeven MLA, Fenton RA. 2014. Renal aquaporins and water balance disorders. *Biochimica*
et Biophysica Acta (BBA) - General Subjects, Aquaporins **1840**:1533–1549.
doi:10.1016/j.bbagen.2013.12.002
Kutscher C. 1968. Plasma volume change during water-deprivation in gerbils, hamsters, guinea
pigs and rats. *Comparative Biochemistry and Physiology* **25**:929–936. doi:10.1016/0010-
406X(68)90581-1
L. Rocha J, Silva P, Santos N, Nakamura M, Afonso S, Qninba A, Boratynski Z, Sudmant PH,
Brito JC, Nielsen R, Godinho R. 2023. North African fox genomes show signatures of
repeated introgression and adaptation to life in deserts. *Nat Ecol Evol* **7**:1267–1286.
doi:10.1038/s41559-023-02094-w
Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**:559. doi:10.1186/1471-2105-9-559
Leib DE, Zimmerman CA, Knight ZA. 2016. Thirst. *Current Biology* **26**:R1260–R1265.
doi:10.1016/j.cub.2016.11.019
Lepkovsky S, Lyman R, Fleming D, Nagumo M, Dimick MM. 1957. Gastrointestinal regulation
of water and its effect on food intake and rate of digestion. *American Journal of*
Physiology-Legacy Content **188**:327–331.
Lighton JRB. 2018. Measuring Metabolic Rates: A Manual for Scientists, 2nd ed. Oxford
University Press. doi:10.1093/oso/9780198830399.001.0001
Love M, Anders S, Huber W. 2014. Differential analysis of count data–the DESeq2 package.
Genome Biol **15**:10–1186.
Ma T, Verkman AS. 1999. Aquaporin water channels in gastrointestinal physiology. *J Physiol*
517:317–326. doi:10.1111/j.1469-7793.1999.0317t.x
MacManes MD. 2017. Severe acute dehydration in a desert rodent elicits a transcriptional
response that effectively prevents kidney injury. *Renal Physiol* 11.
MacManes MD, Eisen MB. 2014. Characterization of the transcriptome, nucleotide sequence
polymorphism, and natural selection in the desert adapted mouse Peromyscus eremicus.
PeerJ **2**:e642. doi:10.7717/peerj.642
MacMillen RE, Hinds DS. 1983. Water Regulatory Efficiency in Heteromyid Rodents: A Model
and Its Application. *Ecology* **64**:152–164. doi:https://doi.org/10.2307/1937337
Malaspinas A-S, Westaway MC, Muller C, Sousa VC, Lao O, Alves I, Bergström A,
Athanasiadis G, Cheng JY, Crawford JE. 2016. A genomic history of Aboriginal
Australia. *Nature* **538**:207–214.

Marquez-Galera A, de la Prida LM, Lopez-Atalaya JP. 2022. A protocol to extract cell-type-
specific signatures from differentially expressed genes in bulk-tissue RNA-seq. *STAR*
Protocols **3**:101121. doi:10.1016/j.xpro.2022.101121

Marra NJ, Eo SH, Hale MC, Waser PM, DeWoody JA. 2012. A priori and a posteriori
approaches for finding genes of evolutionary interest in non-model species:
Osmoregulatory genes in the kidney transcriptome of the desert rodent Dipodomys
spectabilis (banner-tailed kangaroo rat). *Comparative Biochemistry and Physiology Part*
D: Genomics and Proteomics **7**:328–339. doi:10.1016/j.cbd.2012.07.001

Marra NJ, Romero A, DeWoody JA. 2014. Natural selection and the genetic basis of
osmoregulation in heteromyid rodents as revealed by RNA-seq. *Molecular Ecology*
23:2699–2711. doi:10.1111/mec.12764

McCue MD, Sandoval J, Beltran J, Gerson AR. 2017. Dehydration Causes Increased Reliance on
Protein Oxidation in Mice: A Test of the Protein-for-Water Hypothesis in a Mammal.
Physiological and Biochemical Zoology **90**:359–369. doi:10.1086/690912

McKechnie AE, Gerson AR, Wolf BO. 2021. Thermoregulation in desert birds: scaling and
phylogenetic variation in heat tolerance and evaporative cooling. *Journal of Experimental*
Biology **224**:jeb229211. doi:10.1242/jeb.229211

Nordlie RC, Foster JD, Lange AJ. 1999. Regulation of glucose production by the liver. *Annual*
review of nutrition **19**:379–406.

Oksanen J. 2010. Vegan: community ecology package. *http://vegan r-forge r-project org/*.

Paal P, Gordon L, Strapazzon G, Brodmann Maeder M, Putzer G, Walpoth B, Wanscher M,
Brown D, Holzer M, Broessner G, Brugger H. 2016. Accidental hypothermia–an update.
Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine **24**:111.
doi:10.1186/s13049-016-0303-7

Peng X, Cheng J, Li H, Feijó A, Xia L, Ge D, Wen Z, Yang Q. 2023. Whole-genome sequencing
reveals adaptations of hairy-footed jerboas (Dipus, Dipodidae) to diverse desert
environments. *BMC Biol* **21**:182. doi:10.1186/s12915-023-01680-5

Popkin BM, D'Anci KE, Rosenberg IH. 2010. Water, Hydration and Health. *Nutr Rev* **68**:439–
458. doi:10.1111/j.1753-4887.2010.00304.x

Qian Q. 2018. Salt, water and nephron: Mechanisms of action and link to hypertension and
chronic kidney disease. *Nephrology* **23**:44–49. doi:10.1111/nep.13465

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Radford Jr EP. 1959. Factors modifying water metabolism in rats fed dry diets. *American*
Journal of Physiology-Legacy Content **196**:1098–1108.

Ramirez RW, Riddell EA, Beissinger SR, Wolf BO. 2022. Keeping your cool: thermoregulatory
performance and plasticity in desert cricetid rodents. *Journal of Experimental Biology*
225:jeb243131. doi:10.1242/jeb.243131

Reimand J, Kull M, Peterson H, Hansen J, Vilo J. 2007. g:Profiler—a web-based toolset for
functional profiling of gene lists from large-scale experiments. *Nucleic Acids Res*
35:W193–W200. doi:10.1093/nar/gkm226

Roberts EM, Pope GR, Newson MJF, Lolait SJ, O'Carroll A-M. 2011. The Vasopressin V1b
Receptor Modulates Plasma Corticosterone Responses to Dehydration-Induced Stress.
Journal of Neuroendocrinology **23**:12–19. doi:10.1111/j.1365-2826.2010.02074.x

Rocha JL, Godinho R, Brito JC, Nielsen R. 2021. Life in Deserts: The Genetic Basis of
Mammalian Desert Adaptation. *Trends in Ecology & Evolution* **36**:637–650.
doi:10.1016/j.tree.2021.03.007

Rowland NE. 2007. Food or fluid restriction in common laboratory animals: balancing welfare
considerations with scientific inquiry. *Comp Med* **57**:149–160.

Salter D, Watts AG. 2003. Differential suppression of hyperglycemic, feeding, and
neuroendocrine responses in anorexia. *American Journal of Physiology-Regulatory,*
Integrative and Comparative Physiology **284**:R174–R182.
doi:10.1152/ajpregu.00275.2002

Santos RAS, Oudit GY, Verano-Braga T, Canta G, Steckelings UM, Bader M. 2019. The renin-
angiotensin system: going beyond the classical paradigms. *Am J Physiol Heart Circ*
Physiol **316**:H958–H970. doi:10.1152/ajpheart.00723.2018

Schattenberg JM, Czaja MJ. 2014. Regulation of the effects of CYP2E1-induced oxidative stress
by JNK signaling. *Redox Biology* **3**:7–15. doi:10.1016/j.redox.2014.09.004

Schmidt-Nielsen K. 1975. Desert Rodents: Physiological Problems of Desert Life In: Prakash I,
Ghosh PK, editors. Rodents in Desert Environments, Monographiae Biologicae.
Dordrecht: Springer Netherlands. pp. 379–388. doi:10.1007/978-94-010-1944-6_18

Schoorlemmer GH, Evered MD. 1993. Water and solute balance in rats during 10 h water
deprivation and rehydration. *Canadian journal of physiology and pharmacology* **71**:379–
386.

Schoorlemmer GHM, Evered MD. 2002. Reduced feeding during water deprivation depends on
hydration of the gut. *American Journal of Physiology-Regulatory, Integrative and*
Comparative Physiology **283**:R1061–R1069. doi:10.1152/ajpregu.00236.2002

Sikes RS, the Animal Care and Use Committee of the American Society of Mammalogists. 2016.
2016 Guidelines of the American Society of Mammalogists for the use of wild mammals
in research and education. *J Mammal* **97**:663–688. doi:10.1093/jmammal/gyw078

Sugden LA, Atkinson EG, Fischer AP, Rong S, Henn BM, Ramachandran S. 2018. Localization
of adaptive variants in human genomes using averaged one-dependence estimation.
Nature communications **9**:703.

Swain RA, Harris AB, Wiener EC, Dutka MV, Morris HD, Theien BE, Konda S, Engberg K,
Lauterbur PC, Greenough WT. 2003. Prolonged exercise induces angiogenesis and
increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience*
117:1037–1046.

Thornton SN. 2010. Thirst and hydration: Physiology and consequences of dysfunction.
Physiology & Behavior, Beverages and Health **100**:15–21.
doi:10.1016/j.physbeh.2010.02.026

Tigano A, Colella JP, MacManes MD. 2020. Comparative and population genomics approaches
reveal the basis of adaptation to deserts in a small rodent. *Molecular Ecology* **29**:1300–
1314. doi:10.1111/mec.15401

Tigano A, Khan R, Omer AD, Weisz D, Dudchenko O, Multani AS, Pathak S, Behringer RR,
Aiden EL, Fisher H. 2022. Chromosome size affects sequence divergence between
species through the interplay of recombination and selection. *Evolution* **76**:782–798.

Tran LT, Park S, Kim SK, Lee JS, Kim KW, Kwon O. 2022. Hypothalamic control of energy
expenditure and thermogenesis. *Exp Mol Med* **54**:358–369. doi:10.1038/s12276-022-
00741-z

Verkman AS. 2002. Physiological importance of aquaporin water channels. *Annals of Medicine*
34:192–200. doi:10.1080/ann.34.3.192.200
Walsberg GE. 2000. Small Mammals in Hot Deserts: Some Generalizations Revisited.
BioScience **50**:109. doi:10.1641/0006-3568(2000)050[0109:SMIHDS]2.3.CO;2
Watts AG, Boyle CN. 2010. The functional architecture of dehydration-anorexia. *Physiology &*
Behavior, Proceedings from the 2009 Meeting of the Society for the Study of Ingestive
Behavior **100**:472–477. doi:10.1016/j.physbeh.2010.04.010
Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
Wu H, Guang X, Al-Fageeh MB, Cao J, Pan S, Zhou H, Zhang L, Abutarboush MH, Xing Y,
Xie Z, Alshanqeeti AS, Zhang Y, Yao Q, Al-Shomrani BM, Zhang D, Li J, Manee MM,
Yang Z, Yang L, Liu Y, Zhang J, Altammami MA, Wang Shenyuan, Yu L, Zhang W,
Liu S, Ba L, Liu C, Yang X, Meng F, Wang Shaowei, Li L, Li E, Li X, Wu K, Zhang S,
Wang Junyi, Yin Y, Yang H, Al-Swailem AM, Wang Jun. 2014. Camelid genomes
reveal evolution and adaptation to desert environments. *Nature Communications* **5**:1–10.
doi:10.1038/ncomms6188
Yang J, Li W-R, Lv F-H, He S-G, Tian S-L, Peng W-F, Sun Y-W, Zhao Y-X, Tu X-L, Zhang M,
Xie X-L, Wang Y-T, Li J-Q, Liu Y-G, Shen Z-Q, Wang F, Liu G-J, Lu H-F, Kantanen J,
Han J-L, Li M-H, Liu M-J. 2016. Whole-Genome Sequencing of Native Sheep Provides
Insights into Rapid Adaptations to Extreme Environments. *Mol Biol Evol* **33**:2576–2592.
doi:10.1093/molbev/msw129
Yoshimura M, Conway-Campbell B, Ueta Y. 2021. Arginine vasopressin: Direct and indirect
action on metabolism. *Peptides* **142**:170555. doi:10.1016/j.peptides.2021.170555
Yue L, Liu F, Hu J, Yang P, Wang Y, Dong J, Shu W, Huang X, Wang S. 2023. A guidebook of
spatial transcriptomic technologies, data resources and analysis approaches.
Computational and Structural Biotechnology Journal.
892
893

Supplemental

Supplemental Figure 1

Principal component analysis of gene expression of the lung (lu), liver (liv), gastrointestinal tract

(gi), hypothalamus (hyp), and kidney (kid) of *Peromyscus eremicus*. The axes are labelled with

the proportion of the data explained by principal components 1 and 2.

Supplemental figure 2

Visualization of gene ontology (GO) terms of up and down regulated differential gene

expression between the lung (lu), liver (liv), gastrointestinal tract (gi), hypothalamus (hyp), and

kidney (kid) of *Peromyscus eremicus*. Visualized are selections of the top 20 significant GO

terms for each phenotype module combination. The number of genes in the GO term are

indicated by size of the dots.

Supplemental table 1

The number of reads and mapping rate for each RNAseq sample.

Supplemental table 2

Gene assignments to WGCNA modules for the lung, liver, gastrointestinal tract, hypothalamus,

and kidney of *Peromyscus eremicus*.

trait	kidney	gastrointestinal tract	liver	lung	hypothalamus
sex	0	0	0	1	0
delta weight	5	13	9	7	2
proportional weight loss	7	13	9	8	2
Na	6	14	12	9	2
BUN	7	14	8	8	3
AnGap	1	0	0	0	0
K	1	0	0	0	0
Cr	2	3	2	5	4
Htc	7	12	10	8	3
Cl	6	12	12	7	2
Glu	0	0	0	0	0
Hb	7	12	10	8	3
TCO2	5	9	10	6	2
iCa	2	0	4	2	0
RQ	5	1	1	1	2
EE	0	0	2	1	1
WLR	4	3	6	6	1
body temperature	2	3	9	5	2
water access	7	12	9	7	1
Total modules	12	24	18	13	18

Supplemental table 3

The number of WGCNA modules for each phenotypic measurement for the lung, liver,

gastrointestinal tract, hypothalamus, and kidney of *Peromyscus eremicus*.

Supplemental table 4

Results, including the log fold change values, p values, and adjusted p values, from the

differential gene expression analysis for for the lung (lu), liver (liv), gastrointestinal tract (gi),

hypothalamus (hyp), and kidney (kid) of *Peromyscus eremicus.*