Interplay of *Mycobacterium abscessus* and *Pseudomonas aeruginosa* in

coinfection: Biofilm Dynamics and Host Immune Response

Víctor Campo-Pérez[1,2], Esther Julián[2,*] and Eduard Torrents[1,3,*]

[1] Bacterial Infections and Antimicrobial Therapy Group, Institute for Bioengineering of Catalonia

(IBEC), The Barcelona Institute of Science and Technology (BIST), Baldiri Reixac 15-21, 08028

Barcelona, Spain.

[2] Departament de Genètica i de Microbiologia, Facultat de Biociències, Universitat Autònoma de

Barcelona, 08193 Barcelona, Spain.

[3] Microbiology Section, Department of Genetics, Microbiology and Statistics, Faculty of Biology,

University of Barcelona, 643 Diagonal Ave., 08028, Barcelona, Spain.

* Corresponding authors:

Dr. Esther Julián   E-mail: esther.julian@uab.cat

Dr. Eduard Torrents   E-mail: etorrents@ibecbarcelona.eu/eduard.torrents@ub.edu

Abstract

The incidence of infection by nontuberculous mycobacteria, mainly *Mycobacterium abscessus*, in patients with cystic fibrosis and other chronic pulmonary illnesses is increasing, translating into an acceleration in the decline of lung function. In most cases, *M. abscessus* coinfects with *Pseudomonas aeruginosa*, the most common pathogen in these chronic diseases. However, it is unknown how these two bacterial species interact when coinfecting. This study aims to explore the behavior of both species in three relevant pathogenic settings: dual-species biofilm development using a recently developed method to monitor individual species in dual-species biofilms; coinfection in bronchial epithelial cells using *in vitro* assays; and *in vivo* coinfection using the *Galleria mellonella* model. The results demonstrate the capability of both species to form stable mixed biofilms and to reciprocally inhibit single-biofilm progression. Coinfections in bronchial epithelial cells were correlated with significantly decreased cell viability, while in *G. mellonella,* coinfections induced lower survival rates than individual infections. Outstandingly, the analysis of the immune response triggered by each bacterium in bronchial epithelial cell assays and *G. mellonella* larvae revealed that *P. aeruginosa* induces the overexpression of proinflammatory and melanization cascade responses, respectively. In contrast, *M. abscessus* and *P. aeruginosa* coinfection significantly inhibited the immune response in both models, resulting in worse consequences for the host than those generated by single *P. aeruginosa* infection. Overall, the presence of *M. abscessus* produces a decline in the immune responses that worsens the infection and compromises the host.

Importance

The appearance of bacterial infections in the respiratory tract of patients with chronic respiratory diseases suppose a serious and difficult to treat health problem. This complication is exacerbated by the increase resistance against antibiotics generated by pathogenic microorganisms. The most common and virulent pathogenic bacteria reported in the respiratory airway is *Pseudomonas aeruginosa*. It is a Gram-negative, ubiquitous, and intrinsic resistant to antibiotics bacteria. However, the incidence of a rapidly growing, multi-drug resistant mycobacteria; *Mycobacterium abscessus*, is growing worldwide. The pulmonary coinfection by both pathogens is directly related with higher rates of morbidity and mortality of patients. The significance of our research is characterizing the behavior of these two pathogens when they coinfects together, exploring the immune response triggered by the host and its impact in the survival. The purpose is enhancing the limited understanding we have of this clinically relevant coinfection to favor the development of new effective treatments.

Keywords: dual-species biofilm, *Galleria mellonella*, nontuberculous mycobacteria, pulmonary epithelia.

Introduction

Cystic fibrosis (CF) is an autosomal recessive inherited condition caused by the presence of mutations in the gene encoding the cystic fibrosis transmembrane conductance regulator (CFTR) protein (1). This protein acts as a transporter channel that conducts chloride ions (Cl^-) across epithelial cell membranes to maintain the balance of salt and water on many surfaces in the body and is especially relevant in the respiratory system airways (2). A malfunction in CFTR protein prevents proper hydration of the cellular surface, which leads to the loss of the mucus covering the cells, becoming thick and sticky (3). These characteristics of the mucus in CF patients hinder the primary innate defense mechanism of the lung, mucociliary clearance (4), leading to an increased risk of infections from different pathogens, especially *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Burkholderia spp.* and nontuberculous *Mycobacterium spp.* (NTM), which are associated with a high risk of pulmonary compromise and lead to chronic infections (5). In fact, lung tract infections in CF patients are complex and polymicrobial, capable of developing intermicrobial and host–pathogen interactions impacting clinically relevant features such as virulence, persistent colonization of the airways, and antimicrobial recalcitrance (6, 7). Although CF polymicrobial infections are highly individualized in each patient, *S. aureus* and *P. aeruginosa* dominate in children and adult microbial communities, respectively. In this sense, the coinfection by these two dominant pathogens or one of them with other species negatively affect respiratory function compared to single infections (7).

P. aeruginosa is a ubiquitous gram-negative opportunistic pathogen that is metabolically flexible and considered the most relevant bacterium associated with CF due to its high prevalence and pathogenesis. Due to its ability to adapt to the CF pulmonary environment, *P. aeruginosa* is also the primary pathogen present in advanced CF lung disease (8). *P. aeruginosa* PAO1 is the reference laboratory strain for research and PAET1 strain is a clinical isolate from a cystic fibrosis chronically infected patient. *P. aeruginosa* develops biofilms on the thick and dehydrated mucoid surface, which represents a difficult barrier to antibiotic penetration and to host innate immune effectors, in CF (9). These biofilms cause chronic infections through higher antibiotic tolerance, resistance to phagocytosis, and immune-mediated chronic inflammation (10). Furthermore, biofilms are usually polymicrobial and are related to an exacerbated inflammatory immune response that seriously compromises respiratory function, producing the most extensive lung lesions (11).

A significant increase in the incidence of infections caused by NTM in CF patients has been recently reported (12, 13), with fast-growing *Mycobacterium abscessus* being the most prominent and worrisome pathogen of this genus (14). *M. abscessus* infections are associated with an increase in morbidity and mortality due to the long duration of treatment related to significant adverse effects, its multiresistance to antimicrobials, and consequently, the low success rate in its elimination from the respiratory tract (15). Another relevant element in *M. abscessus* pathogenicity is its ability to modify its cell wall, which allows the identification of two different morphotypes in this species: smooth (S), which contains abundant glycopeptidolipids (GPLs) in the outer layer, and rough (R), which is devoid of GPLs (16). Similar to *P.*

aeruginosa, *M. abscessus* also has a biofilm-forming capacity, making treatment even more difficult.

Remarkably, 58-78% of patients with *M. abscessus* infection are also infected by *P. aeruginosa* (17). The

colonization of the lung epithelium begins with S morphotypes, which progressively switch to R mutants,

giving a more aggressive, invasive pulmonary infection (18). In this sense, biofilms developed by each

morphotype of *M. abscessus* present different mechanical properties, with R biofilms having higher rigidity

than S biofilms. Interestingly, *M. abscessus* R/S biofilms present more resistance to mechanical forms of

clearance from the lung than *P. aeruginosa* biofilms (19). Furthermore, *M. abscessus* biofilm bacilli change

the mycolic acid proportions and produce a lipid-relevant extracellular matrix with other components, such

as carbohydrates, proteins, or extracellular DNA (20).

Coinfections of *P. aeruginosa* and *M. abscessus* in CF have become clinically more relevant over the last

two decades; the incidence of NTM infections among CF patients has risen from 3.3% to 22.6%, resulting

in an increase in morbidity and mortality (16). In 16% to 68% of NTM-positive sputum cultures, the *M.*

abscessus complex is present, and its prevalence is increasing (12). Furthermore, the presence of *M.*

abscessus lung disease has been considered a strong relative contraindication to lung transplantation (21),

which can extend and improve the quality of life in properly CF patients, increasing their mean life

expectancy by 7.4 years (22).

Despite the challenges stated above, studies focused on the interaction between both species are scarce. It

is therefore crucial to further elucidate and characterize the *P. aeruginosa*-*M. abscessus* coinfections. For

this reason, the objective of the present study is to unravel the interaction mechanisms between both

bacteria, focusing on relevant aspects of pathogenicity: dual-species biofilm development using recently

developed coculture techniques, including S and R *M. abscessus* morphotypes, and the viability outcome

and the analysis of the triggered immune response by each species in individual and coinfection experiments

using *in vitro* and *in vivo* models. A better understanding of the behavior between *P. aeruginosa* and *M.*

abscessus would provide us with tools to develop efficient treatment strategies against these pathogens in

CF patients.

119
Results

M. abscessus reduces the progression of P. aeruginosa biofilms in vitro

In vitro biofilm development assays showed that the addition of *M. abscessus* R or S to mature *P.*

aeruginosa biofilms limits their growth capacity (**Figure 1A**). This inhibitory effect was observed for both

strains of *P. aeruginosa* PAO1 and PAET1 biofilms. Specifically, the presence of *M. abscessus* (R and S)

resulted in a growth reduction of approximately 20% for PAO1 and 30% for PAET1 biofilms (**Figure 1A**)

compared with the nontreated *P. aeruginosa* biofilms (represented as 100% growth). Nevertheless, the

highest rate of growth inhibition was reported with *M. abscessus* S at 10^4 cfus/well, with approximately

40% PAO1 biofilm reduction. The addition of heat-killed *M. abscessus* or bacterial culture supernatants

had no inhibitory effect on the growth of *P. aeruginosa* mature biofilms (**Figure 1A**). Similarly, the use of

E. coli (gram-negative) or *B. thuringiensis* (gram-positive) did not have a growth-limiting effect on *P.*

aeruginosa biofilms and even, in the case of heat-killed *E. coli* and supernatant from *E. coli* cultures,

induced *P. aeruginosa* PAO1 biofilm growth (**Figure 1A**).

When *P. aeruginosa* strains were cocultured simultaneously with all bacteria tested, the results were

comparable with the previous assay results but with slight inhibition percentage changes in some cases

(**Figure 1B**). *P. aeruginosa* PAO1 cocultured with the *M. abscessus* R morphotype showed a 30% reduction

in PAO1 biofilm compared to the control, while with the S morphotype, the reduction in PAO1 biofilm was

only approximately 20%. In *P. aeruginosa* PAET1 cocultures, the results using *M. abscessus* R or S were

quite similar, with an approximately 30% decrease in PAET1 biofilm compared to the control (growth of

P. aeruginosa alone) (**Figure 1B**). As previously shown in Figure 1A, the addition of *E. coli* and *B.*

thuringiensis and the use of heat-killed cultures or supernatants did not have an inhibitory effect on the *P*

aeruginosa PAO1 and PAET1 biofilms (**Figure 1B**).

After observation of a clear *P. aeruginosa* biofilm reduction with the addition of *M. abscessus*, either added

in a mature biofilm or when simultaneously grown in a dual-species biofilm, the structure and composition

of coculture biofilms were analyzed by confocal microscopy. The images showed that after 24 h of

coculture, *M. abscessus* was established in a *P. aeruginosa* mature biofilm, demonstrating the capacity to

settle and simultaneously grow under these conditions (**Figure 1C**). The presence of *P. aeruginosa* cells

(in blue) was clearly higher, but *M. abscessus* bacilli (in green) were clearly observed in both *P. aeruginosa*

PAO1- and PAET1-formed biofilms and quantitatively represented between 25 to 35% of the sample, as

shown in the pixel color quantification graph (**Figure 1C**). This result demonstrates that both species can

survive and coexist in a dual-species biofilm. Apparently, the presence of both morphotypes R and S of *M.*

abscessus is higher in the *P. aeruginosa* PAET1 biofilm than in PAO1, consistent with the results obtained

in **Figure 1A,** where growth inhibition was higher for the PAET1 strain using the same experimental

conditions than for the PAO strain.

In view of the results obtained, a possible direct inhibitory effect between *P. aeruginosa* PAO1/PAET1

growth and some molecules released by *M. abscessus* R and S, *E. coli* or *B. thuringiensis* was tested. For

this purpose, all bacteria were cultivated on agar plates in close streaks where no inhibitory effects were

observed on the growth of any bacteria by the presence of the others (**Supplementary Figure 3A**). Kinetic

curves of *P. aeruginosa* PAO1 and PAET1 planktonic growth were also generated by cocultivation with

culture supernatants of the other bacteria, and similar to the solid agar cultures, no significant inhibitory

effect on the growth of *Pseudomonas* was reported (**Supplementary Figure 3B and 3C**).

***P. aeruginosa* reduces the development of *M. abscessus* biofilms**

Although infection by *M. abscessus* in CF patients generally occurs in patients previously infected with *P.*

aeruginosa, we aimed to study the effect of *P. aeruginosa* on well-developed biofilms of *M. abscessus*.

Over a mature *M. abscessus* biofilm (120 h old), the addition of *P. aeruginosa* is directly related to a clear

reduction in the development of the *Mycobacterium* biofilm (both for R and S biofilms) (**Figure 2A**),

similar to the opposite case (**Figure 1A**). In detail, biofilm inhibition was notably higher in the *M. abscessus*

S morphotype than in the R morphotype for all the conditions tested. Additionally, *P. aeruginosa* PAO1
induced a higher growth inhibitory effect in both *M. abscessus* biofilm morphotypes than the *P. aeruginosa*
PAET1 strain. These results obtained by GFP quantification were corroborated using confocal microscopy
(**Figure 2B**). The images show that after treatment of an *M. abscessus* biofilm with *P. aeruginosa,* both
bacteria form a combined biofilm in which *P. aeruginosa* expands rapidly due to its competitive advantage
in growth rate. The detection of mycobacteria by GFP in coculture conditions was reduced compared with
that of the control biofilms (without *P. aeruginosa*), and a lower presence of *M. abscessus* was also detected
in the S morphotype compared to the R in combination with both *P. aeruginosa* PAO1 and PAET1 (**Figure
2B**).
177
Coinfection of M. abscessus and P. aeruginosa in bronchial epithelial cells reduces the viability and
immune response triggered with respect to single P. aeruginosa infection
For determination of the effect of bacterial coinfection in the CF context, CFBE41o- (homozygous for the
ΔF508 CFTR mutation) and 16HBE14o- (usually differentiated bronchial epithelial cells) bronchial
epithelial cells were infected with *P. aeruginosa*, *M. abscessus, B. thuringiensis* or combinations of
bacteria. The results showed a reduction in viability of approximately 10-30% in infected cells with respect
to uninfected cells in all cases (**Figure 3A**). Remarkably, while single infection with *M. abscessus* variants
or *B. thuringiensis* reduced viability up to 5-15% in both cell lines, when cells were coinfected with either
of the two *P. aeruginosa* strains, viability was significantly diminished with respect to single infection
(Figure 3A). In all cases, the viability levels of coinfected cells were similar to those obtained with single
infection of *P. aeruginosa* strains, with the exception of the *P. aeruginosa* strains PAO1 and PAET1, and
M. abscessus R coinfected cells, in which lower viability was obtained (Figure 3A). No differences were
obtained between cell lines, indicating that the inhibitory effect of the bacteria is not dependent on the
epithelial characteristics.
Differences were observed for each culture condition when IL-6 and IL-8 production was analyzed. When
cytokine production was analyzed in cell culture supernatants, *P. aeruginosa*-infected cell lines showed the
highest amount, indicating an elevated inflammatory response (**Figure 3B**). Single infections with *M.*
abscessus R and S and *B. thuringiensis* showed different stimulatory capabilities, while *M. abscessus* S did
not induce the production of cytokines, *B. thuringiensis* triggered only IL-8 production in both cell lines,
and *M. abscessus* R triggered IL-6 production in both cell lines and IL-8 only in the control cell line (**Figure
3B**).
The most relevant result was observed in cells coinfected with both bacteria. *M. abscessus* coinfection was
able to inhibit the higher production that was reported by *P. aeruginosa* single-infected cells, and the release
of IL-6 and IL-8 was reduced to levels similar to those triggered by *M. abscessus* single infection (**Figure
3B**). *M. abscessus* S and *P. aeruginosa* (both strains) coinfection reduced the production of cytokines to
levels similar to those in the control cells. Strikingly, this effect observed with *M. abscessus* was not
observed when *B. thuringiensis* was coinfecting with *P. aeruginosa*, triggering cytokine levels similar to

those produced by *P. aeruginosa* single infections (Figure 3B). No differences were observed between cell

lines.

M. abscessus and P. aeruginosa coinfection accelerates G. mellonella larval death

The *G. mellonella* larvae *in vivo* model presents an immune system broadly similar to mammalian innate

immunity being useful for the characterization of microbial infections and host-pathogen interactions.

Infection by both *P. aeruginosa* PAO1 and PAET1 strains is lethal for *G. mellonella* larvae. However, *M.*

abscessus infections need prolonged times (up to 144 h) and high concentrations (10^6 cfus/larva) to become

lethal (see Materials and Methods section and **Supplementary Figure 1**). Coinfection with *M. abscessus*

R and S but not with *B. thuringiensis* accelerated the survival decline in the larvae in both strains and all

concentrations tested compared to the infections of *P. aeruginosa* strains alone (**Figure 4**). However,

although this overtaking lethality was clearly observed, it was only significant for the *P. aeruginosa* PAET1

at the lowest dose (10^4 cfus/larva) coinfected with *M. abscessus* R and S. We corroborated that the causal

agent of larval death was the presence of *P. aeruginosa* PAO1 or PAET1 in the larvae by detecting the

luminescence in their interior trough *lux* detection in a luminescence analysis (**Supplementary Figure 2**).

As Figure 4 shows, *B. thuringiensis* coinfection did not modify larval survival rates.

M. abscessus and P. aeruginosa coinfection restricts the immune response of G. mellonella larvae

G. mellonella larvae produce several antimicrobial peptides (AMPs) and enzymes in response to infection.

The results of immune-relevant *G. mellonella* gene expression showed a significant decrease in the

transcriptional induction for several genes tested, comparing *P. aeruginosa* individual infections with

different coinfection conditions (**Figure 5**). For gloverin, cecropin D, transferrin, moricin, lysozyme, insect

metalloproteinase inhibitor (IMPI) and nitric oxide synthase (NOS) gene expression, a comparable trend

was observed between all the infections assayed. Briefly, high induction of gene expression in PAO1- and

PAET1-infected larvae was observed, with lower induction of gene expression in larvae coinfected with *B.*

thuringiensis and *P. aeruginosa* strains and notably lower gene expression in the case of larvae infected

with both *M. abscessus* R and S alone and coinfected with *P. aeruginosa* (**Figure 5**). These data indicated

that the presence of *M. abscessus* generated a decrease in gene expression compared with the infection of

single *P. aeruginosa* strains, leaving the induction of larvae coinfected by both bacteria at the levels

established by the mycobacteria alone.

Apolipophorin III (Apo III) and glutathione S-transferase (GST) resulted in a similar trend, with no

significant differences detected compared to the control (PBS-infected larvae) in any conditions except for

the *P. aeruginosa* PAO1-infected larvae, which showed evident repression of both genes. NADPH oxidase

(NOX4) gene expression was the only one that did not show significant differences in any of the infections

tested. In the case of hemolin, increased gene expression was observed in *M. abscessus*-infected larvae,

much higher than in the other studied genes and higher than in *B. thuringiensis* infection conditions. This

result contradicts the rest of the immuno-relevant gene tendencies. Hemolin acts as an opsonin and mediates

the interaction between microorganisms and hemocytes, stimulating phagocytosis. In this work, we have

also shown that *M. abscessus* R and S could be phagocytosed by *G. mellonella* hemocytes (**Supplementary**

Figure 4A) and progressively eliminated from the larva (**Supplementary Figure 4B**), which would explain

this contradictory result compared to the rest of the genes.

Discussion

Infections of the respiratory tract epithelium are the leading cause of morbidity and mortality in CF patients.

P. aeruginosa is the most relevant pathogen due to its robust virulence systems and biofilm-forming

capacity, which trigger increased tolerance to antibiotics and resist phagocytosis (9, 23). Although *P.*

aeruginosa is the most common and prevalent pathogen in CF patients, other microorganisms, such as *S.*

aureus, *Haemophilus influenzae*, *Stenotrophomonas maltophilia*, NTM, and *Burkholderia* species, are

presently forming a polymicrobial environment. Within the NTM group, *M. abscessus* is the most relevant

pathogen in CF patients since its multidrug-resistant and biofilm formation capabilities (20, 24) widely

increase antibiotic resistance and prevalence. In this work, the relationship between *P. aeruginosa* PAO1

(laboratory strain)/PAET1 (clinical isolate) and *M. abscessus* R/S morphotypes was studied for the first

time in different contexts of critical interest: biofilm development, bronchial epithelial cell *in vitro* cultures

and *G. mellonella* larvae *in vivo* infection.

Regarding biofilm development, the results indicate a decrease in the progression of *P. aeruginosa* biofilms

when live *M. abscessus* R and S were introduced, independent of whether the biofilm was already well

established (mature biofilm) or in coculture biofilm growth. Inhibition of *Pseudomonas* biofilm

development by cocultivation with other pathogens commonly found in the lungs of CF patients, such as

Aspergillus, or potential probiotics, such as lactobacilli, has already been reported (25, 26). Nevertheless,

no inhibitory effect has been reported previously for mycobacteria. This inhibition could be related to the

capacity of *M. abscessus* to inactivate quorum sensing quinolone signals from *P. aeruginosa* (27). In

addition, the results showed that both species coexist in dual-species biofilms, with *P. aeruginosa* being

the dominant species, most likely related to its competitive advantage in terms of growth since *M. abscessus*

has a division time of 4 to 5 h, while *P. aeruginosa* has a division time of only 25 to 35 min (28, 29).

Conversely, biofilms of *M. abscessus* are also inhibited by the presence of *P. aeruginosa,* which

outcompetes mycobacteria, as previously reported in biofilm coculture experiments (30). The mechanism

responsible of this inhibition remain still unknown and *P. aeruginosa* interbacterial competition secretion

systems and quorum sensing seem to be not involved (31). Distinctly, the growth of S morphotype biofilms,

despite forming biofilm more easily by the abundance of GPLs in its cell wall (32), is inhibited more by

the presence of *P. aeruginosa* than R morphotype biofilms, which could be related to the aggregative

phenotype that confers higher resistance to antibiotics as well as structural rigidity (33). These results

indicate a decrease in both species' biofilm progression when coculturing. It can be hypothesized that this

effect could not be due to direct inhibition but rather a consequence of the combined growth of both species

sharing the same space. This hypothesis is partly supported by the lack of direct inhibition observed in agar

plate cultures using *M. abscessus* R and S cultured close to *P. aeruginosa* PAO1 and PAET1 or in

planktonic growth kinetics curves of *P. aeruginosa* PAO1 and PAET1 in combination with culture

supernatants of the rest of bacteria, in which no significant effect on the growth of *P. aeruginosa* was

highlighted (**see Supplementary Figure 3**). It has been demonstrated that in dual-species biofilms,

antibiotic treatment to eliminate *P. aeruginosa* supposes a competitive advantage for *M. abscessus*

development (17). In any case, *M. abscessus* establishes itself effectively and, despite its difference in

growth rate, prevails in dual-species biofilms, demonstrating its resistance and survival capabilities (34).

Regarding the effect of coinfection on bronchial epithelial cell viability *in vitro*, it has been shown that

coinfection results in higher mortality compared to individual *M. abscessus* and *B. thuringiensis* infections,

as expected. However, as previously reported (35), the viability of the cell lines tested, CFBE41o- and

16HBE14o-, was not excessively compromised, being approximately 80% of the control in all tested

conditions. *M. abscessus* and *P. aeruginosa* coinfections correlate with an elevated pulmonary function

decline and worse disease severity (36, 37); therefore, lower viability of the epithelial cells could be

involved in these complications. The inflammatory response triggered by epithelial cells shows a clearly

exacerbated production of proinflammatory cytokines, IL-6 and IL-8, in response to PAO1 and PAET1

single infections (see Figure 3). This immune response against *P. aeruginosa* is harmful to patients, as

excessive inflammation is responsible for most of the morbidity and mortality in CF (38–40). Surprisingly,

coinfection of *M. abscessus* and *P. aeruginosa* with epithelial cells showed an evident inhibition of the

proinflammatory response promoted by the mycobacteria since single *M. abscessus*-infected cells presented

similar production levels as the coinfected cells, being especially prominent in the case of *M. abscessus* S.

In general, microorganisms that occupy the airways together with *P. aeruginosa* also induce a

proinflammatory response, as in the case of *B. cenocepacia, H. influenzae, S. aureus* and *Candida spp.*

among others (41, 42). A comparable inflammatory reduction effect to that observed for *M. abscessus* in

our results has only been demonstrated in *P. aeruginosa* coinfections with *Prevotella spp.* an anaerobic

gram-negative bacterium belonging to the respiratory tract microbiota (43). In the case of *M. abscessus*,

previous works demonstrated that bronchial epithelial cells generate a hyporesponsive response since *M.*

abscessus S fail to elicit a Toll-like receptor 2 (TLR2) response and only a precise cell fraction from R

morphotype exhibits TLR2 stimulating activity (44), which is necessary for the activation of the signaling

cascade that triggers the secretion of the cytokines IL-6 and IL-8 (45). This lack of immune response

activation observed is particularly high in the case of the S morphotype (as supported by our results) since

the presence of GPL in its cell wall masks other molecules that would activate TLR2 signaling (46). In

most cases, this factor also explains why initial *M. abscessus* colonization of lung airways occurs with an

S morphotype strain. In the CF context, a beneficial immune response consists of the capacity to activate

an inflammatory response to manage infection but to avoid an exuberant response that can be harmful (47).

In the present study, cocultures of *M. abscessus* and *P. aeruginosa* showed less inflammatory activation

and the highest loss of viability among infected cells, while cocultures of *B. thuringiensis* and *P. aeruginosa*

showed a high inflammatory response, leading to a high loss of viability. These results could indicate that

an excessive proinflammatory response and significant immune inhibition could be equally harmful to
infected lung epithelial cells.

The results of the *in vitro* cultures were corroborated by infections in the *G. mellonella* larva model, in
which an increased lethality was observed in coinfection conditions with respect to the single *P. aeruginosa*
infection. *G. mellonella* is highly susceptible to *P. aeruginosa* infections since 100% larval lethality is
reached in less than 24 h with few inoculated bacteria (10-30 CFU) (48, 49). However, *M. abscessus,*
despite being pathogenic for larvae, requires much higher doses and prolonged infection times than *P.*
aeruginosa to be equally lethal ((50, 51), and Supplementary Figure 1). In addition, in agreement with our
in vitro results, the immune response of *G. mellonella* larvae was clearly inhibited under coinfection
conditions (Figure 5). The gene expression of most proteins and enzymes related to larval immunity showed
a significant decrease compared with that of *P. aeruginosa* strain single infections (52). In detail, relevant
AMPs involved in antimicrobial activity, such as gloverin, cecropin D and moricin, were clearly inhibited
in coinfection conditions. All of them act against the bacterial cell wall by interacting with LPS and other
components, especially affecting the outer membrane of gram-negative bacteria (53–55); specifically,
moricin is capable of forming pores in gram-positive and gram-negative cell walls (56). Under coinfection
conditions, insect metalloproteinase inhibitor (IMPI) expression was also inhibited; this peptide has only
been described thus far in *G. mellonella* (57) and inhibits the activity of proteases secreted by invading
microorganisms (58). In this sense, it is well known that *P. aeruginosa* produces various proteases as
virulence factors (59); thus, high levels of IMPI would potentially slow the progression of *P. aeruginosa*
infection. Furthermore, IMPI inhibits the activation of phenoloxidase, the activator enzyme of the
melanization cascade (60). The produced melanin is deposited around the pathogenic microorganism,
limiting its ability to damage the larva, but in parallel to melanin production, dangerous substances and
reactive substances are also produced, resulting in tissue damage and cell death, which could be self-
defeating for the host (61). Additionally, the expression levels of immune-relevant proteins and enzymes,
transferrin, lysozyme and nitric oxide synthetase (NOS), are notably inhibited when *M. abscessus* and *P.*
aeruginosa are infecting simultaneously. Lysozyme is a cationic peptide with muramidase activity (62) that
degrades peptidoglycan, making it more effective against gram-positive bacteria and, similar to IMPI,
preventing an excessive melanization response (63). Transferrin mediates nutritional immunity by
sequestering iron from invading pathogens, limiting their ability to grow and develop (64). *P. aeruginosa*
needs substantial amounts of iron to grow and produces two siderophores to take up this metal (65), so high
levels of transferrin in the hemolymph are beneficial to host survival. The production of nitric oxide by
NOS under infection conditions is increased, although its role in immunity is not entirely clear (66),
although it is hypothesized that it is transformed into reactive nitrogen species that are highly toxic to
bacteria but equally toxic to host cells (67). Moreover, the expression levels of glutathione S-transferase
(GST) and apolipophorin III (Apo III) are clearly repressed only in *P. aeruginosa* PAO1-infected larvae,
while they remain unchanged in the other conditions, including *P. aeruginosa* PAET1, which indicates
differences in the immune response against both *P. aeruginosa* strains. GST is a detoxification enzyme that

protects the host from oxidative stress (68). To reduce pathogen progression, larvae induce a
prophenoloxidase response, which triggers the production of reactive oxygen species to damage bacteria,
although it also affects host cells (69); therefore, it would make sense to inhibit antioxidative enzymes, such
as GST, under severe infections. Apo III is a lipid transport protein with relevant immune effects since it
mediates hemocyte adhesion, phagocytosis and nodule formation (70). *P. aeruginosa* serine protease IV
was shown to degrade Apo III (71, 72); however, the results of our study show that Apo III levels are also
reduced at the transcriptional level. In addition to the protease activity of *P. aeruginosa*, low levels of Apo
III in hemolymph could be reinforced by a repressive effect on its synthesis. NADPH oxidase (NOX4) is
the only enzyme that did not present significant differences between infected and uninfected larvae. This
enzyme produces superoxide and reactive oxygen species in response to infection, suggesting that it is not
overexpressed under the infection conditions tested in this study due to the production of these toxic agents
by the prophenoloxidase pathway (73), which is highly induced, as previously mentioned.

The repressive effect of *M. abscessus* R and S, together with *P. aeruginosa* PAO1 and PAET1 coinfections,
was observed in most of the genes evaluated. However, as the only exception, an induction of transcriptional
expression of the hemolin gene in *M. abscessus*-infected larvae (especially the R variant) was shown.
Hemolin is an immunoglobulin-like protein with no direct antibacterial properties associated with
hemocytes in the pathogen recognition process. This protein functions as an opsonin that mediates between
bacteria and hemocytes, stimulating the phagocytosis process. Surprisingly, this study shows the phagocytic
capacity of *G. mellonella* hemocytes against *M. abscessus* bacilli (**Supplementary Figure 4**), which could
be directly related to this singular overexpression detected in mycobacterium-infected larvae.

Overall, the transcriptional expression levels of the different genes investigated show high induction levels
in the single infections of *P. aeruginosa*, although slight differences between PAO1 and PAET1 strains are
marked, showing different responses. In *M. abscessus*-infected larvae, expression levels are notably lower
and remain at similar levels in both single and coinfections. Notably, larvae infected with *B. thuringiensis*
follow trends similar to those of individual *P. aeruginosa* infections, although interestingly showing lower
levels than these.

The observed results in the immune response triggered by bronchial epithelial cells and *G. mellonella* larvae
are compatible with the decreased response degree under coinfection and contrary to the overexpressed
response under *P. aeruginosa* single infections. The hyperinflammatory response produced in the airways
by *P. aeruginosa* infection is well described in patients (74), producing more severe respiratory malfunction
usually treated with corticosteroids (75). In the case of *M. abscessus*, it has been shown in a murine model
of CF that the proinflammatory response produced is higher in the R morphotype than in the S morphotype
(76), in accordance with the IL-6 and IL-8 differences detected in our study. The immune repression shown
in the coinfections of *M. abscessus* and *P. aeruginosa* is described for the first time in this work. A
disproportionate immune response is dangerous for the host since excess inflammation and the production
of toxic compounds for bacteria and the host trigger a decline in survival. However, the weak immune
response observed in coinfections with *M. abscessus* is even worse since the bacteria can spread

unrestrictedly, triggering lethality at shorter times, as shown in the results of the *in vivo* survival and *in*

vitro viability assays in this work.

Materials and Methods

Bacterial strains and culture conditions.

Two different *P. aeruginosa* strains were used: PAO1 (ATCC 15692) as a reference laboratory strain and

PAET1, a clinical isolate from a chronic CF-infected patient (77). Both strains were transformed with a

plasmid (pETS130lux) expressing a constitutive bacterial luciferase gene cassette (*lux*) as a bioluminescent

bioreporter system (78). Both *P. aeruginosa* strains were grown in Luria-Bertani (LB, Sharlab, Barcelona,

Spain) agar with 50 µg/ml gentamycin (Gm) for PAO1 and 300 µg/ml carbenicillin (Cb) for PAET1 as a

selection marker. For biofilm formation assays, *P. aeruginosa* was grown in tryptic soy broth (TSB,

Sharlab) without antibiotics. *Bacillus thuringiensis* (ATCC 10792) was cultured in tryptic soy agar (TSA,

Sharlab) or TSB, and *Escherichia coli* MG1655 (ATCC 700926) was grown in LB (agar and broth). The

M. abscessus DSMZ 44196 (ATCC 19977) original smooth (S) morphotype and the rough (R) variant, a

natural mutant of the wild-type strain previously obtained (79), were grown on TSB with agitation and 0.5

mm diameter glass beads (DDBiolab, Barcelona, Spain) to prevent cell aggregation and on TSA agar for

colony forming unit (CFU) counting. As previously reported, both variants of *M. abscessus* were

transformed with the plasmid pETS218 (80). For further detail regarding the strains used, see

Supplementary Table S1.

DNA manipulation and plasmid construction

Molecular biology enzymes and kits were purchased from Thermo Fisher Scientific (Madrid, Spain) and

used according to the manufacturer's instructions. DNA amplification was performed by PCR using

DreamTaq MasterMix (2X) or High-Fidelity PCR Enzyme Mix (Thermo Fisher Scientific). The primers

used are listed in **Supplementary Table S2**. Plasmid pETS218 was constructed by cloning the promoter

region of the class Ib ribonucleotide reductase (encoded by the *nrdHIE* genes) from *M. brumae* (GenBank

assembly accession: GCA_900073015.1). All other recombinant DNA manipulations were performed

using standard procedures (81). For construction of mycobacterial *nrdH* transcriptional GFP fusions, a 537

bp long fragment encompassing the *nrdH* promoter region was amplified by PCR using the primer pair

MbruPnrdHIE-For-MbruPnrdHIE-Rev and *M. brumae* genomic DNA; the obtained DNA fragment was

cloned into pJET1.2 and transformed into *E. coli* DH5α cells. *BamHI* and *Apa*I restriction enzymes were

used for fragment digestion and cloning into pFPV27 (82) to generate pETS218. The absence of mutations

introduced during cloning was verified via DNA sequencing. *M. abscessus* S and R morphotypes were

grown in TSA plus 50 µg/ml kanamycin (Kn) when they were transformed with pETS218 as we previously

described (80).

Biofilm progression inhibition quantification

Biofilm quantifications were performed using a microtiter plate screening assay previously optimized and

validated in our group (83). Briefly, mature *P. aeruginosa* PAO1 and PAET1 biofilms (72 h growth) were

cocultured with *B. thuringiensis*, *E. coli* and *M. abscessus* R and S for 24h further. For this purpose, ON

cultures of *B. thuringiensis* and *E. coli* were adjusted to an OD550 nm of 1 and appropriately diluted to

obtain the desired concentrations. *M. abscessus* R and S were grown for 3 days on TSB cultures, and the

concentrations of both live and heat-killed (121 °C for 21 min) cell suspensions were determined by

comparison to the MacFarland 1 (McF) turbidity standard. Bacterial culture supernatants were also obtained

by centrifugation of these cultures at 12,000 g. Serial dilutions of the bacterial suspensions were plated to

verify the accurate concentration in each experiment for all bacteria tested. The level of lux expression (*P.*

aeruginosa cells) in the biofilms was quantified using a Spark® multimode microplate reader (Tecan,

Männedorf, Switzerland). For simultaneous dual-species biofilm formation experiments, *P. aeruginosa*

PAO1 and PAET1 suspensions were mixed with the rest of bacteria or supernatants and grown for 72 h.

M. abscessus biofilm formation and P. aeruginosa coexistence

Three-day-old liquid cultures of *M. abscessus* R and S transformed with pETS218 plasmid were adjusted

to a concentration of 5×10^6 cfu/ml and grown in TSB medium + 0.2% glucose in Costar® 96-Well Black

Polystyrene Plates (200 µl/well) (Corning) at 37 °C in saturation humidity conditions to allow

mycobacterial biofilm formation. After 120 h of incubation, 3 washes with PBS were carried out, and

suspensions of *P. aeruginosa* PAO1 and PAET1 cultures were added at an $OD_{550\ nm} = 0.1$ for an additional

72 h. Finally, the biofilms were homogenized in PBS and quantified by measuring the GFP and *lux*

expression associated with *M. abscessus* and PAO1/PAET1 growth, respectively, using a Spark®

multimode microplate reader.

Biofilm analysis through confocal microscopy

The *P. aeruginosa* and mycobacterial biofilms developed in 96-well plates were collected, and bacteria

were stained with 4',6-diamidino-2-phenylindole (DAPI). *M. abscessus* R and S (transformed with

pETS218) bacilli were specifically detected by measuring their GFP expression. Representative images

were obtained in a Zeiss LSM 800 confocal laser scanning microscope (CSLM, Zeiss, Oberkochen,

Germany) using the 63×/1.4 oil objective with excitation wavelengths of 405 and 488 nm for DAPI and

GFP, respectively. Five representative Z-stack images were selected and analyzed for pixel color

quantification using the color histogram tool from Fiji-ImageJ software and Zen software (Zeiss).

Galleria mellonella maintenance and infection

G. mellonella larvae were fed *at libitum* with an artificial diet as previously described (78). The larvae were

kept at 34 °C in a dark environment until they reached their optimum size (approximately 200 mg). *G.*

mellonella larvae were injected with 10 µl of bacterial suspension containing the desired concentration

through the top right proleg using a 25 µl Hamilton microsyringe (Hamilton, Reno, USA). In previous

experiments, lethal and innocuous doses of each bacterium were determined (**Supplementary Figure 1**).

Based on these results, the highest safe doses were selected: 10^4 CFU/larva for *M. abscessus* R and S and

10^5 CFU/larva for *B. thuringiensis* and *E. coli*. In the case of *P. aeruginosa*, even the lowest dose tested (10

CFU/larva) was lethal. The virulence of the *P. aeruginosa* PAO1 strain was much higher than that of

PAET1; therefore, the PAET1 strain dose was adjusted to 10^4 CFU/larva, similar to 10 CFU/larva of PAO1.

Ten larvae were infected with each condition, *E. coli*, *B. thuringiensis* and *M. abscessus* R/S, and were then

injected with the *P. aeruginosa* PAO1 or PAET1 strains. Additionally, larvae infected with only each *P.*

aeruginosa strain were used as controls for comparison purposes. Infected larvae were kept at 37 °C to

allow the development of infection for 14 h. At that time, five larvae were used to extract RNA, and ten

were monitored for survival for 24 h.

For analysis of the phagocytosis of *M. abscessus* R and S by *G. mellonella* hemocytes and the concentration

of mycobacteria in the hemolymph (**Supplementary Figure 4**), the larvae were infected with 10^4

CFU/larvae of each variant. For collection of hemolymph, larvae were placed on ice for 10 min. Once

anesthetized, the anal proleg was cut with a surgical blade, and the hemolymph was collected in a 1.5 ml

Eppendorf tube. Hemolymph from 10 larvae was pooled and appropriately diluted and cultured on TSA

plates for mycobacterial CFU counts. The hemolymph from the other 10 larvae was centrifuged and washed

3 times with PBS (5 min at 500 g at 4 °C). Precipitated hemocytes were resuspended in PBS and stained

with DAPI (blue) (4′,6-diamidino-2-phenylindole) to observe nuclei and with FM 4-64 (red) (N-3-

triethylammoniumpropyl-4-6-4-diethylamino phenyl hexatrienyl pyridinium dibromide) for dying

hemocyte vesicles and plasma membrane and observed under a confocal microscope (Zeiss LSM 800

confocal laser scanning microscope (CSLM)).

G. mellonella RNA extraction and immune-relevant gene expression analysis

At 14 h post-infection, five larvae were placed on ice for 10 min. The larvae were homogenized using a

rotor-stator homogenizer (IKA, Staufen, Germany), and 30 mg of the homogenate was used for RNA

extraction following the GeneJET RNA Purification Kit instructions (Thermo Fisher Scientific). RNA

obtained was treated with 10x TURBO DNase (Life Technologies, Carlsbad, USA) for 1 h to eliminate

possible DNA contamination. DNA absence was verified by PCR amplification of the 18S rRNA

housekeeping gene using genomic DNA of *G. mellonella* as a positive control. RNA was quantified in an

M200 PRO microplate reader (Tecan). RNA to cDNA reverse-transcription was performed using Maxima

Reverse Transcriptase (Thermo Fisher Scientific) with Oligo (dT)18 primers (Thermo Fisher Scientific) by

following the manufacturer's instructions. The cDNA obtained was stored at -20 °C until use. Quantitative

real-time PCR (qRT-PCR) was performed using PowerUp™ SYBR™ Green Master Mix (Applied

Biosystems, Foster City, CA, USA) in a StepOnePlus™ Real-Time PCR System (Applied Biosystems)

according to the manufacturer's protocol. All qRT-PCRs used specific primers of *G. mellonella* immune-

relevant genes listed in **Supplementary Table 2**.

Bronchial epithelial cell line culture and infection

Two human bronchial epithelial cell lines were used: CFBE41o-, isolated from a CF patient homozygous for the ΔF508 CFTR mutation, and 16HBE14o-, isolated from a cardiopulmonary patient with overexpression of CFTR (35). Cells were maintained in Dulbecco's modified Eagle's medium: nutrient mixture F12 (DMEM/F12; Thermo Fisher Scientific) with 10% (v/v) decomplemented fetal bovine serum (dFBS) (Gibco, Paisley, UK) and 1% (v/v) penicillin–streptomycin (Thermo Fisher Scientific). Cells were maintained in a humidified incubator at 37 °C and 5% (v/v) CO_2 (ICOmed, Memmert). For infection experiments, cells ($5x10^4$ cell/well) were incubated in 96-well plates (Corning) using antibiotic- and serum-free medium for 3 h to allow cell adhesion. Subsequently, the cells were infected at a multiplicity of infection (MOI) of 50:1 for *P. aeruginosa* PAO1 and PAET1 (50:1) and of 10:1 for *M. abscessus* R / S and *B. thuringiensis*. Infections with each bacterium as well as coinfections combining both strains of *P. aeruginosa* and *M. abscessus* were carried out. Three hours later, the wells were washed 3 times with PBS to remove extracellular bacteria, and the plates were incubated for 24 hours. The supernatant was then removed and stored at -80 °C for cytokine measurements. The effect of *P. aeruginosa*, *M. abscessus* and their combination on cell viability was analyzed by Presto Blue assay (Thermo Fisher Scientific) following the manufacturer's instructions, using uninfected cells as a control. Cell infections were performed in triplicate in three independent experiments.

IL-6 and IL-8 detection by enzyme immunoassays (ELISAs) in cell supernatants

IL-6 and IL-8 production was measured in cell culture supernatants using ELISA kits according to the manufacturer's instructions (Becton and Dickinson, BD, San Diego, USA). Briefly, wells were coated with IL-6 or IL-8 capture antibody diluted in coating buffer (0.1 M sodium carbonate, pH 9.5) overnight at 4 °C. Plates were then washed and blocked with assay diluent (PBS with 10% FBS) for 1 h at room temperature (RT). Plates were prepared with standards diluted in assay diluent, and samples were incubated for 2 h at RT. After washing, streptavidin-horseradish peroxidase conjugate mixed with biotinylated detection antibody was then added to each well for 1 h. Finally, tetramethylbenzidine (TMB) substrate solution and hydrogen peroxide (BD, San Diego, USA) were added to the plates for 30 min in the dark. Absorbance ($OD_{630 \text{ nm}}$) was measured in an Infinite 200 PRO microplate reader (Tecan) and analyzed with Magellan software (Tecan).

Author Contribution

The manuscript was written through the contributions of all authors. Víctor Campo performed the experiments. Esther Julián and Eduard Torrents directed the research, revised the experimental data, and wrote the manuscript. All authors have approved the final version of the manuscript.

Acknowledgments

This study was supported by grants from the Ministerio de Ciencia, Innovación y Universidades and Agencia Estatal de Investigación (AEI), Spain, cofunded by Fondo Europeo de Desarrollo Regional, FEDER, European Union (RTI2018-098777-B-I00, PID2021-122331OB-I00, PID2021-125801OB-100, PDC2022-133577-I00 and PLEC2022-009356), the CERCA programme and *AGAUR-Generalitat de Catalunya* (2021SGR-0092 and 2021SGR-0145), the European Regional Development Fund (FEDER), the Catalan Cystic Fibrosis association and Obra Social "La Caixa". VC-P is a recipient of a Ph.D. contract (FI) from the Generalitat de Catalunya.

Declaration of Competing Interest

The authors declare no financial or commercial conflicts of interest.

Data Availability

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Figure 1: Coculture of *P. aeruginosa* PAO1 and PAET1 biofilms with different bacteria. A) Growth of well-developed mature *P. aeruginosa* biofilm when different bacterial cultures (live and heat-killed) or bacterial culture supernatants were added. **B)** Growth of *P. aeruginosa* in coculture biofilms. The values correspond to the luciferase expression (*lux*) of the *P. aeruginosa* strains generated in each condition, normalized based on the average expression reported in control wells (marked with a dashed line as 100% *P. aeruginosa* growth). The effects of *M. abscessus* R and S morphotypes, *E. coli* MG1655 and *B. thuringiensis* on both *P. aeruginosa* strains were tested as well as the supernatant (SN) and heat-killed (HK) form of these bacterial cultures. Data are presented as the mean values ± SDs of at least three independent experiments including six replicates in each experimental condition; significant differences were determined using one-way ANOVA and Tukey's multiple comparisons tests; ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$. **C)** Representative confocal microscopy images of coculture biofilms and pixel color quantification of the images. *P. aeruginosa* PAO1 and PAET1 biofilms were grown for 72 h, and *M. abscessus* R and S were inoculated for an additional 24 h. Z-stack compositions of 96 h-old *P. aeruginosa* PAO1 and PAET1 biofilms in combination with *M. abscessus* R and S morphotypes. DAPI stained all cells blue, and green fluorescent protein (GFP) of plasmid pETS218 allowed mycobacterial cells to be detected in green. Scale bars correspond to 20 μm. Pixel quantifications are represented as the mean values ± SDs from three representative images of each condition.

Figure 2: Effect of the *P. aeruginosa* PAO1 and PAET1 strains on *M. abscessus* R and S biofilm

development. A) Graph represents the inhibition of *M. abscessus* biofilms when cocultured for 72 h with

P. aeruginosa by measuring GFP expression. The control corresponds to 100% of single *M. abscessus*

biofilm growth by analyzing GFP expression (dotted line). Data are presented as the mean ± SD of three

independent experiments including six replicates in each experimental condition, and significant

differences were reported using the Mann–Whitney *t* test, ****, $p<0.0001$. **B)** Confocal microscopy

representative coculture biofilm images and pixel color quantification. Z-stack compositions of *M.*

abscessus variant 120 h biofilms in combination with *P. aeruginosa strains*. DAPI stained all cells blue,

and green fluorescent protein (GFP) of plasmid pETS218 allowed mycobacterial cells to be detected in

green. Scale bars correspond to 20 µm. Pixel quantifications are represented as the mean values ± SDs from

three representative images of each condition.

Figure 3: Bronchial epithelial cell viability and proinflammatory cytokine production after *P.*
***aeruginosa* PAO1/PAET1, *M. abscessus* R/S and *B. thuringiensis* infections. A)** Viability of CFBE41o-
and 16HBE14o- cell lines after bacterial infections. Presto blue cell viability assay results in which data are
expressed with respect to uninfected wells (considered as 100% cell viability). Data are represented as the
mean ± SD from three independent experiments. Significance was analyzed using one-way ANOVA and
Tukey's multiple comparisons test. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$. Comparison of
cocultures with individual PAO1/PAET1 infections: ##, $p<0.01$; ###, $p<0.001$. **B)** Cytokine (IL-6 and IL-8)
production detected in culture supernatants from bronchial epithelial cell infections. Statistical differences
were determined using one-way ANOVA and Dunnett's test, which compares each column with the control.
Data are represented as the mean ± SD from three independent experiments; *, $p<0.05$; **, $p<0.01$; ***,
$p<0.001$; ****, $p<0.0001$. Comparison of results obtained in *M. abscessus* R infections and those in *M.*
abscessus S infections were determined using one-way ANOVA, Tukey's multiple comparisons test; #,
$p<0.05$; ##, $p<0.01$; ###, $p<0.001$; ####, $p<0.0001$.

606
607
608

624
625
626

References

1. O'Sullivan BP, Freedman SD. 2009. Cystic fibrosis. Lancet 373:1891–1904.

2. Linsdell P. 2018. Cystic fibrosis transmembrane conductance regulator (CFTR): Making an ion channel out of an active transporter structure. Channels (Austin) 12:284–290.

3. Kreda SM, Davis CW, Rose MC. 2012. CFTR, mucins, and mucus obstruction in cystic fibrosis. Cold Spring Harb Perspect Med 2.

4. Bustamante-Marin XM, Ostrowski LE. 2017. Cilia and mucociliary clearance. Cold Spring Harb Perspect Biol 9.

5. Coutinho H, Falcão-Silva VS, Gonçalves G. 2008. Pulmonary bacterial pathogens in cystic fibrosis patients and antibiotic therapy: a tool for the health workers. Int Arch Med 1:24.

6. Filkins LM, O'Toole GA. 2015. Cystic fibrosis lung infections: polymicrobial, complex, and hard to treat. PLoS Pathog 11:e1005258.

7. Jean-Pierre F, Vyas A, Hampton TH, Henson MA, O'toole GA. 2021. One versus many: Polymicrobial communities and the cystic fibrosis airway. mBio 12:1–7.

8. Gaspar MC, Couet W, Olivier JC, Pais AACC, Sousa JJS. 2013. *Pseudomonas aeruginosa* infection in cystic fibrosis lung disease and new perspectives of treatment: a review. Eur J Clin Microbiol Infect Dis 32:1231–1252.

9. Høiby N, Ciofu O, Bjarnsholt T. 2010. *Pseudomonas aeruginosa* biofilms in cystic fibrosis. Future Microbiol 5:1663–1674.

10. Moreau-Marquis S, Stanton BA, O'Toole GA. 2008. *Pseudomonas aeruginosa* biofilm formation in the cystic fibrosis airway. Pulm Pharmacol Ther 21:595–599.

11. Hartl D, Gaggar A, Bruscia E, Hector A, Marcos V, Jung A, Greene C, McElvaney G, Mall M, Döring G. 2012. Innate immunity in cystic fibrosis lung disease. J Cyst Fibros 11:363–382.

12. Martiniano SL, Nick JA, Daley CL. 2016. Nontuberculous mycobacterial infections in cystic fibrosis. Clin Chest Med 37:83–96.

13. Abidin NZ, Gardner AI, Robinson HL, Haq IJ, Thomas MF, Brodlie M. 2021. Trends in nontuberculous mycobacteria infection in children and young people with cystic fibrosis. J Cyst Fibros 20:737–741.

14. Esther CR, Esserman DA, Gilligan P, Kerr A, Noone PG. 2010. Chronic *Mycobacterium abscessus* infection and lung function decline in cystic fibrosis. J Cyst Fibros 9:117–123.

15. Brugha R, Spencer H. 2021. *Mycobacterium abscessus* in cystic fibrosis. Science 372:465–466.

16. Degiacomi G, Sammartino JC, Chiarelli LR, Riabova O, Makarov V, Pasca MR. 2019. *Mycobacterium abscessus*, an emerging and worrisome pathogen among cystic fibrosis patients. Int J Mol Sci 20.

17. Rodríguez-Sevilla G, Crabbé A, García-Coca M, Aguilera-Correa JJ, Esteban J, Pérez-Jorge C. 2019. Antimicrobial treatment provides a competitive advantage to *Mycobacterium abscessus* in a dual-species biofilm with *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 63.

18. Halloum I, Carrère-Kremer S, Blaise M, Viljoen A, Bernut A, le Moigne V, Vilchèze C, Guérardel Y, Lutfalla G, Herrmann JL, Jacobs WR, Kremer L. 2016. Deletion of a

dehydratase important for intracellular growth and cording renders rough *Mycobacterium*
abscessus avirulent. Proc Natl Acad Sci U S A 113:E4228–E4237.

19.   Gloag ES, Wozniak DJ, Stoodley P, Hall-Stoodley L. 2021. *Mycobacterium abscessus*
biofilms have viscoelastic properties which may contribute to their recalcitrance in chronic
pulmonary infections. Sci Rep 11:5020.

20.   Dokic A, Peterson E, Arrieta-Ortiz ML, Pan M, di Maio A, Baliga N, Bhatt A. 2021.
Mycobacterium abscessus biofilms produce an extracellular matrix and have a distinct
mycolic acid profile. Cell Surf 7.

21.   Gilljam M, Scherstén H, Silverborn M, Jönsson B, Ericsson Hollsing A. 2010. Lung
transplantation in patients with cystic fibrosis and *Mycobacterium abscessus* infection.
Journal of Cystic Fibrosis 9:272–276.

22.   Thabut G, Christie JD, Mal H, Fournier M, Brugière O, Leseche G, Castier Y, Rizopoulos
D. 2013. Survival benefit of lung transplant for cystic fibrosis since lung allocation score
implementation. Am J Respir Crit Care Med 187:1335–1340.

23.   Maurice NM, Bedi B, Sadikot RT. 2018. *Pseudomonas aeruginosa* biofilms: host response
and clinical implications in lung infections. Am J Respir Cell Mol Biol 58:428–439.

24.   Esteban J, García-Coca M. 2018. Mycobacterium Biofilms. Front Microbiol 8.

25.   Shokri D, Khorasgani MR, Mohkam M, Fatemi SM, Ghasemi Y, Taheri-Kafrani A. 2018.
The inhibition effect of *Lactobacilli* against growth and biofilm formation of *Pseudomonas*
aeruginosa. Probiotics Antimicrob Proteins 10:34–42.

26.   Reece E, Doyle S, Greally P, Renwick J, McClean S. 2018. *Aspergillus fumigatus* inhibits
Pseudomonas aeruginosa in co-culture: implications of a mutually antagonistic relationship
on virulence and inflammation in the CF airway. Front Microbiol 9.

27.   Birmes FS, Wolf T, Kohl TA, Rüger K, Bange F, Kalinowski J, Fetzner S. 2017.
Mycobacterium abscessus subsp. abscessus is capable of degrading *Pseudomonas*
aeruginosa quinolone signals. Front Microbiol 8.

28.   Cortes MAM, Nessar R, Singh AK. 2010. Laboratory maintenance of *Mycobacterium*
abscessus. Curr Protoc Microbiol Chapter 10.

29.   LaBauve AE, Wargo MJ. 2012. Growth and laboratory maintenance of *Pseudomonas*
aeruginosa. Curr Protoc Microbiol Chapter 6.

30.   Rodríguez-Sevilla G, García-Coca M, Romera-García D, Aguilera-Correa JJ, Mahíllo-
Fernández I, Esteban J, Pérez-Jorge C. 2018. Non-Tuberculous Mycobacteria multispecies
biofilms in cystic fibrosis: development of an in vitro *Mycobacterium abscessus* and
Pseudomonas aeruginosa dual species biofilm model. Int J Med Microbiol 308:413–423.

31.   Idosa AW, Wozniak DJ, Hall-Stoodley L. 2022. Surface dependent inhibition of
Mycobacterium abscessus by diverse *Pseudomonas aeruginosa* strains. Microbiol Spectr
10.

32.   Clary G, Sasindran SJ, Nesbitt N, Mason L, Cole S, Azad A, McCoy K, Schlesinger LS,
Hall-Stoodley L. 2018. *Mycobacterium abscessus* smooth and rough morphotypes form
antimicrobial-tolerant biofilm phenotypes but are killed by acetic acid. Antimicrob Agents
Chemother 62.

33.   Bernut A, Herrmann JL, Kissa K, Dubremetz JF, Gaillard JL, Lutfalla G, Kremer L. 2014.
Mycobacterium abscessus cording prevents phagocytosis and promotes abscess formation.
Proc Natl Acad Sci U S A 111.

34. de Groote MA, Huitt G. 2006. Infections due to rapidly growing mycobacteria. Clin Infect
Dis 42:1756–1763.

35. del Mar Cendra M, Torrents E. 2020. Differential adaptability between reference strains and
clinical isolates of *Pseudomonas aeruginosa* into the lung epithelium intracellular lifestyle.
Virulence 11:862–876.

36. Hsieh MH, Lin CY, Wang CY, Fang YF, Lo YL, Lin SM, Lin HC. 2018. Impact of
concomitant nontuberculous mycobacteria and *Pseudomonas aeruginosa* isolates in non-
cystic fibrosis bronchiectasis. Infect Drug Resist 11:1137–1143.

37. Brennan MJ. 2017. Biofilms and *Mycobacterium tuberculosis*. Infect Immun 85.

38. Courtney JM, Ennis M, Elborn JS. 2004. Cytokines and inflammatory mediators in cystic
fibrosis. J Cyst Fibros 3:223–231.

39. Roesch EA, Nichols DP, Chmiel JF. 2018. Inflammation in cystic fibrosis: An update.
Pediatr Pulmonol 53:S30–S50.

40. Kube D, Sontich U, Fletcher D, Davis PB. 2001. Proinflammatory cytokine responses to *P.
aeruginosa* infection in human airway epithelial cell lines. Am J Physiol Lung Cell Mol
Physiol 280.

41. Gangell C, Gard S, Douglas T, Park J, de Klerk N, Keil T, Brennan S, Ranganathan S,
Robins-Browne R, Sly PD. 2011. Inflammatory responses to individual microorganisms in
the lungs of children with cystic fibrosis. Clin Infect Dis 53:425–432.

42. Heijerman H. 2005. Infection and inflammation in cystic fibrosis: a short review. J Cyst
Fibros 4 Suppl 2:3–5.

43. Bertelsen A, Elborn JS, Schock BC. 2021. Microbial interaction: *Prevotella spp.* reduce *P.
aeruginosa* induced inflammation in cystic fibrosis bronchial epithelial cells. J Cyst Fibros
20:682–691.

44. Le Moigne V, Roux AL, Jobart-Malfait A, Blanc L, Chaoui K, Burlet-Schiltz O, Gaillard
JL, Canaan S, Nigou J, Herrmann JL. 2020. A TLR2-Activating fraction from
Mycobacterium abscessus rough variant demonstrates vaccine and diagnostic potential.
Front Cell Infect Microbiol 10.

45. Kempaiah P, Davidson LB, Perkins DJ, Byrd TF. 2013. Cystic fibrosis CFBE41o- cells
contain TLR1 SNP I602S and fail to respond to *Mycobacterium abscessus*. J Cyst Fibros
12:773.

46. Davidson LB, Nessar R, Kempaiah P, Perkins DJ, Byrd TF. 2011. *Mycobacterium
abscessus* Glycopeptidolipid prevents respiratory epithelial TLR2 signaling as measured by
HβD2 gene expression and IL-8 release. PLoS One 6:29148.

47. van Heeckeren AM, Sutton MT, Fletcher DR, Hodges CA, Caplan AI, Bonfield TL. 2021.
Enhancing cystic fibrosis immune regulation. Front Pharmacol 12.

48. Jander G, Rahme LG, Ausubel FM. 2000. Positive correlation between virulence of
Pseudomonas aeruginosa mutants in mice and insects. J Bacteriol 182:3843–3845.

49. Ramarao N, Nielsen-Leroux C, Lereclus D. 2012. The insect *Galleria mellonella* as a
powerful infection model to investigate bacterial pathogenesis. J Vis Exp
https://doi.org/10.3791/4392.

50. Meir M, Grosfeld T, Barkan D. 2018. Establishment and validation of *Galleria mellonella*
as a novel model organism to study *Mycobacterium abscessus* infection, pathogenesis, and
treatment. Antimicrob Agents Chemother 62.

51. Li Y, Spiropoulos J, Cooley W, Khara JS, Gladstone CA, Asai M, Bossé JT, Robertson BD,
Newton SM, Langford PR. 2018. *Galleria mellonella* - a novel infection model for the
Mycobacterium tuberculosis complex. Virulence 9:1126–1137.

52. Andrejko M, Mak P, Siemińska-Kuczer A, Iwański B, Wojda I, Suder P, Kuleta P, Regucka
K, Cytryńska M. 2021. A comparison of the production of antimicrobial peptides and
proteins by *Galleria mellonella* larvae in response to infection with two *Pseudomonas*
aeruginosa strains differing in the profile of secreted proteases. J Insect Physiol 131.

53. Xu XX, Zhong X, Yi HY, Yu XQ. 2012. *Manduca sexta* gloverin binds microbial
components and is active against Bacteria and Fungi. Dev Comp Immunol 38:275.

54. Yi HY, Chowdhury M, Huang YD, Yu XQ. 2014. Insect antimicrobial peptides and their
applications. Appl Microbiol Biotechnol 98:5807–5822.

55. Moore AJ, Beazley WD, Bibby MC, Devine DA. 1996. Antimicrobial activity of cecropins.
J Antimicrob Chemother 37:1077–1089.

56. Hara S, Yamakawa M. 1995. Moricin, a novel type of antibacterial peptide isolated from
the silkworm, *Bombyx moriv*. J Biol Chem 270:29923–29927.

57. Vertyporokh L, Wojda I. 2017. Expression of the insect metalloproteinase inhibitor IMPI
in the fat body of *Galleria mellonella* exposed to infection with *Beauveria bassiana*. Acta
Biochim Pol 64:273–278.

58. Wojda I. 2017. Immunity of the greater wax moth *Galleria mellonella*. Insect Sci 24:342–
357.

59. Galdino ACM, Branquinha MH, Santos ALS, Viganor L. 2017. *Pseudomonas aeruginosa*
and its arsenal of proteases: Weapons to battle the host. Pathophysiological Aspects of
Proteases 381–397.

60. Clermont A, Wedde M, Seitz V, Podsiadlowski L, Lenze D, Hummel M, Vilcinskas A.
2004. Cloning and expression of an inhibitor of microbial metalloproteinases from insects
contributing to innate immunity. Biochemical Journal 382:315.

61. Fallon JP, Troy N, Kavanagh K. 2011. Pre-exposure of *Galleria mellonella* larvae to
different doses of *Aspergillus fumigatus* conidia causes differential activation of cellular
and humoral immune responses. Virulence 2:413–421.

62. Catalão MJ, Pimentel M. 2018. Mycobacteriophage lysis enzymes: targeting the
mycobacterial cell envelope. Viruses 10.

63. Rao XJ, Ling E, Yu XQ. 2010. The role of lysozyme in the prophenoloxidase activation
system of *Manduca sexta*: an in vitro approach. Dev Comp Immunol 34:264–271.

64. Iatsenko I, Marra A, Boquete JP, Peña J, Lemaitre B. 2020. Iron sequestration by transferrin
1 mediates nutritional immunity in *Drosophila melanogaster*. Proc Natl Acad Sci U S A
117:7317–7325.

65. Minandri F, Imperi F, Frangipani E, Bonchi C, Visaggio D, Facchini M, Pasquali P,
Bragonzi A, Visca P. 2016. Role of iron uptake systems in *Pseudomonas aeruginosa*
virulence and airway infection. Infect Immun 84:2324–2335.

66. Sadekuzzaman M, Stanley D, Kim Y. 2018. Nitric oxide mediates insect cellular immunity
via phospholipase A2 activation. J Innate Immun 10:70–81.

67. Hillyer JF, Estévez-Lao TY. 2010. Nitric oxide is an essential component of the hemocyte-
mediated mosquito immune response against bacteria. Dev Comp Immunol 34:141–149.

68. Vogel H, Altincicek B, Glöckner G, Vilcinskas A. 2011. A comprehensive transcriptome
and immune-gene repertoire of the lepidopteran model host *Galleria mellonella*. BMC
Genomics 12:1–19.

69. Lu A, Zhang Q, Zhang J, Yang B, Wu K, Xie W, Luan YX, Ling E. 2014. Insect
prophenoloxidase: the view beyond immunity. Front Physiol 5.

70. Zakarian RJ, Dunphy GB, Albert PJ, Rau ME. 2002. Apolipophorin-III affects the activity
of the haemocytes of *Galleria mellonella* larvae. J Insect Physiol 48:715–723.

71. Andrejko M, Mizerska-Dudka M, Jakubowicz T. 2008. Changes in *Galleria mellonella*
apolipophorin III level during *Pseudomonas aeruginosa* infection. J Invertebr Pathol 97:14–
19.

72. Andrejko M, Cytryńska M, Jakubowicz T. 2005. Apolipophorin III is a substrate for
protease IV from *Pseudomonas aeruginosa*. FEMS Microbiol Lett 243:331–337.

73. Bergin D, Reeves EP, Renwick J, Wientjes FB, Kavanagh K. 2005. Superoxide production
in *Galleria mellonella* hemocytes: identification of proteins homologous to the NADPH
oxidase complex of human neutrophils. Infect Immun 73:4161–4170.

74. Malhotra S, Hayes D, Wozniak DJ. 2019. Cystic fibrosis and *Pseudomonas aeruginosa*: the
host-microbe interface. Clin Microbiol Rev 32.

75. Dinwiddie R. 2005. Anti-inflammatory therapy in cystic fibrosis. J Cyst Fibros 4 Suppl
2:45–48.

76. Caverly LJ, Caceres SM, Fratelli C, Happoldt C, Kidwell KM, Malcolm KC, Nick JA,
Nichols DP. 2015. *Mycobacterium abscessus* morphotype comparison in a murine model.
PLoS One 10.

77. Crespo A, Gavaldà J, Julián E, Torrents E. 2017. A single point mutation in class III
ribonucleotide reductase promoter renders *Pseudomonas aeruginosa* PAO1 inefficient for
anaerobic growth and infection. Sci Rep 7.

78. Moya-Andérico L, Admella J, Fernandes R, Torrents E. 2020. Monitoring gene expression
during a *Galleria mellonella* bacterial infection. Microorganisms 8:1–14.

79. Brambilla C, Llorens-Fons M, Julián E, Noguera-Ortega E, Tomàs-Martínez C, Pérez-
Trujillo M, Byrd TF, Alcaide F, Luquin M. 2016. Mycobacteria clumping increase their
capacity to damage macrophages. Front Microbiol 7:1562.

80. Campo-Pérez V, Cendra M del M, Julián E, Torrents E. 2021. Easily applicable
modifications to electroporation conditions improve the transformation efficiency rates for
rough morphotypes of fast-growing mycobacteria. N Biotechnol 63:10–18.

81. Falcon I. 1982. Molecular cloning: A laboratory manual third edition. Molecular Cloning:
A Laboratory Manual Third Edition.

82. Perrone F, de Siena B, Muscariello L, Kendall SL, Waddell SJ, Sacco M. 2017. A novel
TetR-like transcriptional regulator is induced in acid-nitrosative stress and controls
expression of an efflux pump in Mycobacteria. Front Microbiol 8.

83.   Campo-Pérez V, Alcàcer-Almansa J, Julián E, Torrents E. 2023. A high-throughput
839   microtiter plate screening assay to quantify and differentiate species in dual-species
840   biofilms. Microorganisms 2023, Vol 11, Page 2244 11:2244.

 Supplementary information

844
845
846 **Supplementary Table 1: Strains and plasmids.** Detail of the strains and plasmids used in this study, *P.*
847 *aeruginosa* strains PAO1 and PAET1 were transformed with the pETS130Lux plasmid to quantify the
848 expression of luciferase (*Lux*). The *M. abscessus* strains were transformed with the pETS218 plasmid for
849 constitutive expression of green fluorescent protein (GFP).
850

Strains	Observations
Pseudomonas aeruginosa PAO1	Reference strain for laboratory studies. Compatible with a cystic fibrosis acute infection.
Pseudomonas aeruginosa PAET1	Cystic fibrosis strain isolated from a chronic patient (77)
Mycobacterium abscessus Smooth morphotype	Type strain. (DSMZ 44916)
Mycobacterium abscessus Rough morphotype	Strain obtained by several passages of the original type strain (79)
Escherichia coli str. K-12 substr. MG1655	Model strain in molecular biology. Safe strain used as control.
Bacillus thuringiensis	Type strain. (CECT 197)
Escherichia coli DH5α	Versatile strain used for general cloning applications. In the study it was used to preserve the different plasmids: pFPV27, pJET 1.2 + PnrdH, pETS218. All were transformed by thermal shook.

Plasmids	Observations
pFPV27	Mycobacteria expression, it has a kanamycin resistance gene and another that codes for GFP.
pJET1.2	Positive selection cloning vector
pETS130Lux	Plasmid previously constructed to obtain *Lux* constitutive expression in *P. aeruginosa*
pETS218 (MbruPnrdHIE + pFPV27)	Plasmid constructed to obtain GFP constitutive expression in *M. abscessus*

852
853
854

Supplementary Table 2: List of primers used for pETS218 construction and the RT-PCRs experiments. The list includes the genes with relevance in the immune response of the *Galleria mellonella* larva against infection by bacteria used in this study.

Primer name	Sequence 5´-3´	Observations
MbruPnrdHIE For	AAGGATCCGCGGTTCGCGACGCCGTC	Primers used to amplify the promoter region of the class Ib ribonucleotide reductase. This sequence was then used to clone into pFPV27 to obtain a constitutive expression of GFP (pETS218)
MbruPnrdHIE Rev	AAGGGCCCAGCGCCTTGTAGGTCGCGTT	
18S rRNA For	ATGGTTGCAAAGCTGAAACT	Housekeeping gene: Used to normalize gene expression data in RT-PCRs.
18S rRNA Rev	TCCCGTGTTGAGTCAAATTA	
Apo III For	AGACTTGCACGCCATCAAGA	Apolipophorin III: activity as pathogen recognition receptor, stimulating the activity of defense peptides, and possessing antimicrobial activity itself.
Apo III Rev	TGCATGCTGTTTGTCACTGC	
Gloverin For	AGATGCACGGTCCTACAG	Interacts with lipopolysaccharides (LPS) and inhibit the formation of Gram-negative bacterial outer membrane.
Gloverin Rev	GATCGTAGGTGCCTTGTG	
Lysozyme For	TCCCAACTCTTGACCGACGA	Is a muramidase that cleaves the linkages between N-acetylomuramid acid and N-acetylglucosamine in bacterial peptidoglycan.
Lysozyme Rev	AGTGGTTGCGCCATCCATAC	
Cecropin D For	CTGCGCCATGTTCTTCA	Cationic antimicrobial peptide that disrupts microbial membranes, which eventually results in microbial cell death.
Cecropin D Rev	TCGCATCTCTGATCCTCTG	
Moricin For	GCTGTACTCGCTGCACTGAT	Antibacterial activity against both Gram-positive and Gram-negative bacteria. Forms ion channels in the bacterial membrane.
Moricin Rev	TGGCGATCATTGCCCTCTTT	
Hemolin For	CCCGAAGACGCTGGTGAATA	Functions as an opsonin that facilitates pathogen recognition and mediates hemocytic immune responses
Hemolin Rev	CGCACGTTCATTTGCTGTTC	
GST For	GACAGAAGTCCTCCGGTCAG	Glutathione S-transferase: protect cells from oxidative stress, but they also play a central role in the detoxification of both endogenous and xenobiotic compounds.
GST Rev	TCCGTCTTCAAGCAAAGGCA	
NOX-4 For	TGGCACGGCATCAGTTATCA	NADPH oxidase: is a pro-oxidative stress enzyme, induce oxidative stress humoral responses and secrete reactive oxygen species (ROS).
NOX-4 Rev	ACAGCGACTGTCATGTGGAA	
NOS For	ATGAAGGTGCTGAAGTCACAA	Nitric oxide synthase: NO active the gene encoding of some antimicrobial peptides.
NOS Rev	GCCATTTTACAATCGCCACAA	
IMPI For	ATTTGTAACGGTGGACACGA	Insect metalloproteinase inhibitor: protect AMPs against digestion by metalloproteinases secreted by the invading bacteria.
IMPI Rev	CGCAAATTGGTATGCATGG	
Transferrin For	CCCGAAGATGAACGATCAC	Mediates nutritional immunity by sequestering iron from invading pathogens. Insects trigger a hypoferremic response after infection to limit iron availability to invading microbes.
Transferrin Rev	CGAAAGGCCTAGAACGTTTG	

861 **Supplementary Figure 1: Kaplan–Meier survival curves of *Galleria mellonella* for the optimization**

862 **of bacterial innocuous and lethal doses.** The strains used in this study were tested to determine doses of

863 interest. *P. aeruginosa* was lethal at all doses tested. The doses of PAET1 strain were adjusted to obtain

864 similar results as PAO1 (more virulent). Ten larvae were used for each bacterium and condition (n=10).

Supplementary Figure 2: Images of *lux* expression in *G. mellonella* larvae to report *P. aeruginosa* infection. Images were taken every 2 h until all the larvae died (-) using an Image Quant LAS 4000 (GE Healthcare, Chicago, USA) measuring chemiluminescence at a 30 sec exposure.

	14 h	16h	18 h	20 h	22 h	24 h
PAO1 10						-
M. abscessus S + PAO1 10					-	-
PAO1 10²					-	-
M. abscessus S + PAO1 10²				-	-	-
PAO1 10³				-	-	-
M. abscessus S + PAO1 10³			-	-	-	-

	14 h	16h	18 h	20 h	22 h	24 h
PAET1 10⁴						
M. abscessus S + PAET1 10⁴						-
PAET1 10⁵					-	-
M. abscessus S + PAET1 10⁵				-	-	-
PAET1 10⁶				-	-	-
M. abscessus S + PAET1 10⁶				-	-	-

	14 h	16h	18 h	20 h	22 h	24 h
PAET1 10^4						
B. thuringiensis + PAET1 10^4						-
PAET1 10^5						-
B. thuringiensis + PAET1 10^5						-
PAET1 10^6				-	-	-
B. thuringiensis + PAET1 10^6				-	-	-

	14 h	16h	18 h	20 h	22 h	24 h
PAO1 10						
B. thuringiensis + PAO1 10						
PAO1 10^2						-
B. thuringiensis + PAO1 10^2						-
PAO1 10^3				-	-	-
B. thuringiensis + PAO1 10^3				-	-	-

Supplementary Figure 3: Direct growth inhibition assay and growth kinetics of ***P. aeruginosa***

strains culture with bacterial culture supernatants. A) Images of agar plates with *M. abscessus*

R and S cultures and perpendicular streaks of the different strains of *P. aeruginosa, E. coli* and *B.*

thuringiensis to visualize possible inhibitory growth effects. **B)** *P. aeruginosa* PAO1 and PAET1

liquid cultures growth kinetics after adding different amounts of supernatant from cultures of *M.*

abscessus R and S. **C)** *P. aeruginosa* PAO1 and PAET1 growth kinetics by adding different

amounts of supernatant from cultures of *E. coli* and *B. thuringiensis*. The absorbance (OD $_{550\,nm}$)

and bioluminescence (Relative Light Units, RLUs) of *P. aeruginosa* planktonic cultures are

quantified in 96-well plates (100 µl/well) every 15 min in contact with different volumes of other

bacterial supernatants (SN); 10, 30 and 50 µl.

A)

B)

904
905

C)

906
907

Supplementary Figure 4: Phagocytosis of *M. abscessus* **bacilli by** *G. mellonella* **hemocytes and**
prevalence inside the larvae over time. A) Confocal microscope micrographs and Imaris software
reconstruction of *M. abscessus* phagocytosed by *G. mellonella* hemocytes. Cell nuclei was dye with DAPI
(blue), vesicular network and plasma membrane were dye with FM 4-64 (red) (N-3-
Triethylammoniumpropyl-4-6-4-Diethylamino Phenyl Hexatrienyl Pyridinium Dibromide) and
mycobacteria was detected through GFP (green) expression. Scale bars correspond to 10 μm in XY and
5μm in XZ reconstructions. **B)** *M. abscessus* R and S morphotypes colony forming units (cfus) recovered
from *G. mellonella* hemolymph after infection over time.

916
917

A)

M. abscessus Biofilm Inhibition

% of growth

**** ****

+ PAO1 + PAET1   + PAO1 + PAET1
M. abscessus R   *M. abscessus* S

B)

	Control	PAO1	PAET1
M. abscessus R			
M. abscessus S			

Pixel quantification in the biofilm

% of pixels

PAO1 PAET1   PAO1 PAET1
M. abscessus R   *M. abscessus* S

■ *M. abscessus*
■ *P. aeruginosa*

M. abscessus R + PAO1 coinfection

M. abscessus R + PAET1 coinfection

M. abscessus S + PAO1 coinfection

M. abscessus S + PAET1 coinfection

B. thuringiensis + PAO1 coinfection

B. thuringiensis + PAET1 coinfection